1 **Naturally occurring variation in gene-associated transposable elements impacts gene**
2 **expression and phenotypic diversity in woodland strawberry**
3

4 Santiago Priego-Cubero[1,7], Rocio Tolley[2,6,7], Julia Llinares-Gómez[1], Camila Zlauvinen[1],
5 Tuomas Toivainen[3], Timo Hytönen[3], Carmen Martín-Pizarro[1*], Ileana Tossolini[2,4,5*], Pablo A.
6 Manavella[1*]
7

AFFILIATIONS
[1]Instituto de Hortofruticultura Subtropical y Mediterránea "La Mayora" (IHSM "La Mayora"),
Universidad de Málaga-Consejo Superior de Investigaciones Científicas (UMA-CSIC), 29010
Málaga, Spain.
[2]Facultad de Ingeniería, Universidad Nacional de Entre Ríos, 3100 Oro Verde, Argentina.
[3]Department of Agricultural Sciences, Viikki Plant Science Centre, University of Helsinki,
Finland.
[4]Instituto de Agrobiotecnología del Litoral (CONICET-UNL), 3000 Santa Fe, Argentina.
[5]Current address: Instituto de Investigación y Desarrollo en Bioingeniería y Bioinformática
(IBB, CONICET-UNER), 3100 Oro Verde, Argentina.
[6]Current address: Instituto Andino Patagónico de Tecnologias Biológicas y Geoambientales
(IPATEC, CONICET-UNComa), 8400 Rio Negro, Argentina.
[7]These authors contributed equally to this work.
21

22 *To whom correspondence should be addressed. E-mail: ileana.tossolini@uner.edu.ar;
23 pablomanavella@ihsm.uma-csic.es; cmmartin@uma.es
24

ABSTRACT

Transposable elements (TEs) are major components of plant genomes and powerful drivers
of regulatory and phenotypic diversity, yet their mechanistic contributions to gene regulation
in non-model crops remain poorly understood. Here, we generate a comprehensive and
curated TE annotation for woodland strawberry (*Fragaria vesca*), revealing that TEs comprise
~30% of the genome and uncovering an exceptional enrichment of gene-proximal miniature
inverted-repeat transposable elements (MITEs) and short inverted repeats (IRs).
Integrating small RNA, DNA methylation and transcriptomic data across tissues and fruit
developmental stages, we show that most gene-associated MITEs and IRs are epigenetically
active, producing 24-nt siRNAs and displaying dynamic CHH methylation patterns during fruit
ripening. Genome-wide analyses reveal that MITE- and IR-proximal genes occupy a distinct

regulatory landscape compared to genes associated with other TE classes, consistent with a
role in context-dependent transcriptional regulation.

As a mechanistic proof of concept, we demonstrate that a Mutator-like MITE located near
FvAAT1, encoding a key alcohol acyltransferase required for aroma ester biosynthesis,
modulates gene expression through siRNA-dependent DNA methylation and ripening-stage-
specific chromatin reorganization. Transient perturbation of this MITE epigenetic state alters
FvAAT1 expression, and chromatin conformation capture reveals MITE-associated looping at
this locus. Notably, this regulatory MITE is conserved in commercial strawberry species,
suggesting an evolutionarily preserved function in fruit quality traits. Extending these findings
to natural variation, we characterize thousands of TE insertion and deletion polymorphisms
across 210 European *F. vesca* accessions and show that TE-based variation recapitulates
population structure. A TE-informed genome-wide association analysis identifies significant
associations between MITE and IR polymorphisms and fruit volatile compounds, including key
aroma determinants.

Together, our results establish MITEs and short IRs as widespread, epigenetically active
regulatory elements that shape gene expression, chromatin architecture and phenotypic
diversity in strawberry. This work provides a conceptual and methodological framework to
integrate TE-driven regulatory variation into population genomics, breeding and genome-
editing strategies aimed at improving fruit quality.

55
56

57 **KEYWORDS**
58 Transposons, Strawberries, DNA methylation, MITE, Inverted Repeats, Fragaria, Epigenetics
59

INTRODUCTION
Transposable elements (TEs) are repetitive, mobile DNA sequences widely dispersed
across eukaryotic genomes. They constitute the most abundant genomic component in many
species, accounting for nearly 60% of the human (*Homo sapiens*) genome [1] and up to 85%
of the wheat (*Triticum aestivum*) genome [2]. Their extensive genomic presence directly
impacts genome architecture, evolution, and gene regulation, particularly in response to
environmental stressors [3-7].

TEs are classified into two major classes based on their transposition mechanisms [8].
Class I elements, or retrotransposons, transpose via an RNA intermediate through a "copy-
and-paste" mechanism, whereas Class II elements, known as DNA transposons, move via a
"cut-and-paste" mechanism [9, 10] or, in the case of Helitrons, a "peel-and-paste" replicative
mechanism that uses a circular DNA intermediate [11]. TEs are further categorized into

autonomous elements, which encode the proteins required for their own transposition, and
non-autonomous elements, which rely on autonomous TEs for mobility [8].

Within these classes, TEs are subdivided into orders, superfamilies, and families based on
structural and functional characteristics, such as replication strategies, sequence
conservation, and enzymatic activity [8]. In plants, Class I Long terminal repeat (LTR)
retrotransposons (e.g., Gypsy and Copia) are particularly abundant. Non-LTR
retrotransposons (e.g., LINEs and SINEs), though more extensively studied in animals, also
contribute significantly to plant genome evolution.

Beyond their contribution to genome size and structure, specific types of TEs are
increasingly recognized for their regulatory potential. In this context, miniature inverted-repeat
transposable elements (MITEs) represent a prominent group of short (<800 bp), non-
autonomous Class II TEs that are highly abundant in eukaryotic genomes [7, 12, 13]. While
many TEs accumulate in pericentromeric regions, MITEs and other short TE-derived inverted
repeats (IRs) are frequently located near genes, where they can function as regulatory
elements [14, 15]. A key feature that makes MITEs potent regulatory elements is their ability
to form hairpin-shaped double-stranded RNAs (dsRNAs) upon transcription. These dsRNAs
are processed by DCL3 into 24-nt small interfering RNAs (siRNAs) that trigger DNA
methylation through a RNA-dependent RNA Polymerase 2 (RDR2)-independent variant of the
RNA-directed DNA methylation (RdDM) pathway. This non-canonical form of RdDM plays a
crucial role in silencing TEs to maintain genome stability [16-18]. Different from the canonical
RdDM-initiated silencing of TEs through heterochromatin formation, RdDM-mediated DNA
methylation of MITEs does not commonly act as a silencing mark but instead could impact the
expression, both positively and negatively, of nearby genes [7].

TE research has expanded significantly with the increasing availability of genome
sequences, enabling improved annotation, classification, and functional analysis of these
elements. However, challenges remain, particularly in non-model species with incomplete
TE libraries [19]. Various computational approaches have been developed for TE
annotation [20], including *de novo* prediction [21-23], motif-based searches [24, 25], and
similarity-based methods [26, 27]. Combining multiple strategies has proven most
effective in generating accurate and comprehensive TE annotations [19, 23, 28, 29].

One of the most compelling open questions in TE research is understanding how the
large natural variation in TE content within a species contributes to phenotypic diversity.
Thanks to advancements in long-read high-throughput sequencing technologies, the study
of TE polymorphisms has gained attention due to their role in complex traits and their potential
as genetic markers for plant research and breeding [30, 31]. Integrating TE polymorphisms

with Genome-Wide Association Studies (GWAS) has provided valuable insights into the functional implications of TEs in various biological processes [32-35]. For instance, transposon insertion polymorphisms (TIPs) have been shown to improve predictions of agronomic traits beyond what is possible with single-nucleotide polymorphism (SNP)-based approaches alone [34, 36-38].

The woodland strawberry (*Fragaria vesca*) is a wild diploid species from the Rosaceae family that serves as a model for strawberry research. It is one of the four ancestral species contributing to the polyploidization of the cultivated octoploid *F. × ananassa*, the most widely consumed strawberry species. Among the four subgenomes of *F. × ananassa*, the *F. vesca* subgenome exhibits the lowest density of TEs near genes and encodes significantly more dominantly expressed homeologs than the other three subgenomes combined [39]. While previous studies have examined TE diversity in the *F. vesca* genome [40-42], their complete annotation, functional characterization, and linkage with phenotypic variation remain poorly explored.

In addition to its genomic significance, *F. vesca* fruits are distinguished by their unique volatilome, which contributes to their intense floral and fruity aroma [10, 43]. The species exhibits substantial phenotypic diversity influenced by both genetic variation and environmental factors. The contribution of TEs to this phenotypic diversity remains largely unexplored. Given its importance as a genetic resource, understanding TE-driven variation in *F. vesca* could provide valuable insights into fruit development and quality traits, with potential applications for breeding.

In this study, we systematically annotated transposable elements in the *Fragaria vesca* reference genome, generating a curated TE library. Using this resource, we analyzed TE insertion and deletion polymorphisms across 210 re-sequenced European *F. vesca* accessions [44], revealing their contribution to genetic diversity and subpopulation structure. We then integrated TE polymorphisms into a genome-wide association framework, identifying significant associations with economically important fruit traits, including volatile compound profiles.

As a proof of concept, we functionally validated the regulatory impact of a MITE located near FvH4_7g18570, which encodes the alcohol acyl transferase *FvAAT1*, a putative ortholog of *SAAT* involved in volatile ester biosynthesis in *F. × ananassa [45]*. Combining gene expression analyses with chromatin conformation profiling at this locus, we demonstrate that MITE-derived siRNAs trigger DNA methylation that reshapes local chromatin organization, thereby modulating *FvAAT1* expression in a ripening-stage-dependent manner.

While genome-wide TE distribution patterns, such as LTR enrichment in
pericentromeric regions and MITE localization near genes, have been described in other
plant species, the TE landscape of *F. vesca* has remained poorly annotated and
functionally unexplored. Our work provides a comprehensive TE resource for this species
and establishes a framework to investigate the regulatory potential of MITEs and short
inverted repeats, particularly during fruit development. By integrating TE-driven variation
into population-genomic and functional analyses, this study offers new insights into how
TEs contribute to gene regulation and phenotypic diversity in strawberry, and highlights
their potential as targets for breeding and genome-editing strategies aimed at improving
fruit quality.

RESULTS

Comprehensive identification and annotation of transposable elements in *Fragaria*
vesca

Transposable elements (TEs) do not typically experience selective pressures that preserve
their sequences over time. Consequently, they tend to become fragmented due to the
accumulation of random mutations, which drives divergence among different TEs and
complicates their identification [23, 46, 47]. To address these challenges, we combined two
recent bioinformatic pipelines, EDTA [48] and DeepTE [49], and applied TEtrimmer [50], a
consensus refinement and de-redundancy tool, to generate a semi-curated de novo TE library
in *F. vesca* (Figure S1). This allowed us to comprehensively identify and classify TEs in the *F.*
vesca v4.0 reference genome [51, 52] into classes, orders, and superfamilies (Table 1, Figure
1A).

The complete set of identified TEs accounted for approximately 30% of the *F. vesca*
genome (Figure 1B). In plants, TE content is typically proportional to genome size [53, 54].
For example, the maize genome (~2,500 Mb) contains over 80% TEs and repetitive elements,
whereas the smaller *Arabidopsis thaliana* genome (~135 Mb) contains only 21% TEs [55-57].
The proportion of TEs identified for *F. vesca* (~30%), which has a genome size of ~220 Mb
[58], aligns with these observations (Figure 1B).

In contrast to most plant species, where Class I retrotransposons typically dominate, the
genome of *F. vesca* exhibits a relatively balanced proportion between Class I and Class II
elements. Class I retrotransposons constitute approximately 15% of the total genomic DNA,
whereas Class II DNA transposons represent around 13.4% (Table 1). However, domain-
based analyses revealed that only a small fraction of Class II elements retain detectable

transposase-associated domains, indicating that most DNA transposon copies correspond to non-autonomous or fragmented elements rather than potentially autonomous transposons. Within Class I, LTR retrotransposons remain the most prevalent, comprising most of these elements, with the Gypsy superfamily being particularly abundant (Figure 1A), and non-LTR retrotransposons contribute only a minor fraction (~0.25%) of the genome. The relatively moderate proportion of Class I retrotransposons observed in *F. vesca* might reflect its smaller genome size compared to other plant genomes and could be further influenced by limitations in accurately assembling centromeric regions, which are enriched with these elements.

Interestingly, *Fragaria* species exhibit higher diversity among DNA transposons, compared to other plant genomes, highlighting the unique composition of their genomes [40]. After re-classifying DNA transposons into MITES and non-MITEs with DeepTE, almost 13% of identified TEs belong to subclass 1, including both MITEs and other DNA transposons from the CACTA, Harbinger, TcMar, hAT, and Mutator superfamilies. The remaining 0.62% corresponds to subclass 2, specifically Helitrons (Figure 1A). Despite their known difficulty in identification, the proportion of Helitrons identified in the *F. vesca* genome is consistent with reports in other plant genomes, where they typically constitute below 4% of the genome [40, 48, 59].

We then analyzed the size distribution of TEs for each class based on the annotation generated for *F. vesca*, including both fragmented and intact TEs (Figure 1C, Figure S2, Table S1). Overall, no clear differences were observed in size distributions among the results obtained from EDTA, DeepTE, and the curated library (Figure 1C). The average length of each intact and fragmented TE type is detailed in Figure S2 and Table S1. The similarity in size distributions across libraries suggests that the final curated TE library preserves the structural characteristics of both intact and fragmented elements, supporting the robustness of the library for subsequent analyses.

A high proportion of fragmented TEs was identified across the genome. This explains the relatively low mean length values observed for the complete TE library compared to the intact subset (Figure S2), a pattern consistent with the long-term accumulation of mutations and deletions in ancient TEs. While Class I elements generally have lengths below 4 kb, the longest sequences belong to the LTR order (Figure 1D, Figure S2). In contrast, Class II TEs display a broader size range strongly linked to their respective superfamily classification (Figure 1D). The distinct size patterns observed among TE classes suggest differences in replication strategies and selective pressures acting on different TE families.

To better understand the structural preservation of these elements, we partitioned the library into two mutually exclusive sets: intact and fragmented elements. A marked contrast in the distribution of TE classes between these groups was observed. While DNA transposons

dominated both fractions, they showed a significant increase of 12.3% in the intact library compared to the fragmented set. This enrichment was driven primarily by MITEs, which accounted for 36.6% of the intact elements, a substantial increase of 16.1% relative to their 20.5% representation in the fragmented library. Conversely, LTRs were more prevalent among fragmented sequences than in the intact subset, representing a decrease of 9.5%. This highlights a higher rate of structural decay for these larger elements. The abundance of intact MITEs, which are known for being relatively short, can be attributed to their unique structural nature. Unlike autonomous DNA transposons or LTR retrotransposons, MITEs are characterized by their compact sequence length (Figure 1D) and lack of internal coding regions. From a genomic perspective, a shorter sequence presents a significantly smaller "mutational target" for stochastic fragmentation, deletions, or nested TE insertions. While the integrity of an LTR element is easily compromised by a single internal deletion, MITEs are more likely to remain structurally recognizable as "intact" over evolutionary time. Furthermore, because MITEs do not rely on internal Open Reading Frames (ORFs) for their identification they are less sensitive to the sequence decay that typically fragments autonomous elements. Consequently, while larger autonomous elements are rapidly converted into genomic "noise" or fragmented relics, MITEs maintain higher structural stability.

Genome-wide distribution and abundance of Transposable Elements

The distribution, abundance, and genome coverage of TEs vary substantially across species and are strongly influenced by subfamily classification. While some TEs tend to accumulate near protein-coding genes or toward the centromeres, others are more evenly distributed across chromosomes. To assess the distribution of TE types in *F. vesca*, circos ideograms were generated for each TE class, with gene density included as a reference (Figure 2).

As shown in Figure 2A, regions with higher Class I TE content tend to have lower gene density, a pattern particularly evident for chromosomes Fvb1, Fvb3, and Fvb4. This TE distribution is predominantly driven by LTR transposons, which are the most abundant among Class I. A large fraction of LTR elements appeared located more than 3,000 bp from genes, suggesting a low likelihood of gene association (Figure S3). This distribution pattern of Class I elements has been previously reported in several plant species, including tomato (*Solanum lycopersicum*) and rice (*Oryza sativa) [56, 57, 60]*).

In contrast, Class II TEs are more concentrated near pericentromeric regions compared to chromosome arms except for Helitrons, which exhibit a more widespread distribution across the genome (Figure 2B). Helitrons positioning varies considerably between species. For example, Helitrons display a more erratic pattern of distribution in potato and rice [57, 61]

compared to Arabidopsis, where they are enriched in gene-poor pericentromeric regions, or
in maize (*Zea mays*), where they are more abundant in gene-rich regions [62].

MITEs exhibit a distinct distribution pattern characterized by a notable concentration in
gene-rich regions, particularly in the immediate vicinity of, or within, coding sequences (Figure
S3). This pattern aligns with previous reports describing the regulatory role of MITEs in gene
expression [56, 57, 63]. Notably, 46% of *F. vesca* genes harbor a MITE within 3,000 bp of their
coding sequence, and 15% of these genes contain MITEs within their coding regions. This
proximity to genes reinforces the potential role of MITEs in influencing gene expression
through epigenetic and chromatin-level mechanisms encouraging us to focus on the study of
these elements.

Characterization of MITEs and short TE-derived IRs in *F. vesca*

MITEs are frequently found in close proximity to genes, where they influence gene
expression and act as evolutionary agents contributing to adaptation and developmental
processes [64, 65]. This regulatory impact is often mediated by the production of 24-nt small
RNAs through the non-canonical RdDM pathway [64, 66, 67]. Such processes trigger localized
DNA methylation and induce structural changes in chromatin organization, leading to the
formation of short-range chromatin interactions that modulate the transcriptional activity at
neighboring genes [64-67]. A similar effect can be triggered by gene-associated TE-derived
short inverted repeats (IRs) elements, which often originate from MITEs that have lost
distinctive structural elements. The presence of IRs reflects the progressive fragmentation and
degeneration of TEs over evolutionary time [7]. In any case, and for the purpose of the
regulatory effects studied here, IRs and MITEs operate similarly.

Given the mechanistic similarities between IRs and MITEs, we employed *einverted* from
the European Molecular Biology Open Software Suite (EMBOSS) [68] to also annotate all IRs
in *F. vesca*. We then classified the putative nature of the identified IRs and compared them
with the superfamilies previously assigned to MITEs (Figure 3A). IRs can originate from both
MITE-type and DNA transposons, Class I elements, or even non-TE sequences [64].
Consistently, 84% of the identified IRs were classified as MITE-derived elements, in line with
the matching distribution patterns of MITEs and IRs (Figure S4A). Additionally, the overall
superfamily distribution of IRs was consistent with that of the annotated MITEs in our TE library
(Figure 3A). However, the proportion of IRs from the MITE Harbinger superfamily was lower
than that of the annotated MITEs. This suggests that certain IRs may have originated from
more ancient MITE elements that have since diverged from their original sequence patterns.
Lastly, 16% of the IRs were classified as elements distinct from MITEs ("Other" category in
Figure 3C), predominantly belonging to DNA Mutator/CACTA, LTR Gypsy/Copia, and

unknown origin (Figure S4B). Considering this classification, we observed that 58% of the
identified IRs overlapped by at least 20% of their length with MITEs annotated in our TE library.
Therefore, subsequent analyses were performed separately for MITEs and MITE-independent
IRs.

Most of these identified IRs (4,374, ~80%) were found near or within annotated protein-
coding regions (Figure 3B). Of these, 1,047 IRs are situated 500 bp from a gene, 734 between
501 bp and 1,000 bp, and 1,493 between 1,001 bp and 3,000 bp (Figure 3B). In total, 14% of
F. vesca genes harbor an IR within these ranges. Similarly, of the 26,060 MITEs annotated
(including fragmented elements), 18,221 MITEs were located near or within annotated protein-
coding regions. Specifically, 5,277 MITEs were located within 500 bp of a gene, 3,154 between
501 bp and 1,000 bp, and 6,127 between 1,001 and 3,000 bp (Figure 3A). In total, 51% of *F.*
vesca genes harbor at least one MITE or IR within 3,000 bp of their coding regions. Notably,
1,100 IRs (~20%) and 3,665 MITEs (~14%) are fully located within protein-coding genes,
primarily in introns (Figure 3B-C), representing approximately 8% of all coding genes in the *F.*
vesca genome. TE insertions in introns can have a significant regulatory effect. For example,
in maize, insertion of a CACTA TE derived element into the *Par1* gene creates an epiallele
whose methylation status directly affects gene expression and plant architecture [69].

The detected IRs ranged in size from 100 bp to 700 bp (the length limit of our analysis),
with an average length of 445 bp. The annotated MITEs exhibited a broader size distribution,
ranging from 51 bp to 2,120 bp, with an average length of 261 bp. An overrepresentation of
elements between 100 bp and 500 bp (73%) was observed (Figure 1D), values aligning with
the typical size range of MITEs in plants [70].

Most MITEs and IRs in *F. vesca* are epigenetically active

Upon transcription, MITEs and IRs can be processed into siRNA that trigger non-canonical
RdDM, leading to the regulation of neighboring genes and turning these TEs into autonomous
regulatory elements. To identify potentially active regulatory MITEs and IRs, we used sRNA
and bisulfite (BS) sequencing to determine whether the identified MITEs and IRs produced
24-nt siRNAs and triggered DNA methylation—clear indicators of active and potentially
regulatory TEs. Three distinct tissues and stages were examined: leaf, and fruit receptacle at
both white (immature) and red (ripe) developmental stages (Figure 3D, Figure S5).

Our analysis revealed that 66% of MITEs and 77% of IRs produced siRNAs in at least one
of the tested tissues (Table S2). Notably, approximately 80% of these elements are located
within 3 kb of a gene, which is consistent with previous reports showing that gene-hosted
MITEs and IRs are commonly cotranscribed by RNA polymerase II (Pol II) and are
epigenetically active [64]. Metagene analysis of 24-nt siRNAs mapping to the MITEs and IRs

showed a characteristic two-peak profile, indicating that the siRNAs are produced from the terminal inverted repeats (TIR) through DLC3 processing (Figure 3D). This behavior was particularly pronounced for IRs located within 1,000 bp of genes. In contrast, for MITEs, this pattern persisted even for elements located farther away from genes, although their expression levels decreased with increasing distance (Figure S5A).

BS-sequencing metagene analysis revealed that annotated TEs, MITEs, and IRs are methylated under all tested conditions (Figure 3E, Figure S5B). Gene-associated MITEs and IRs, particularly those closest to genes, showed a signature two-peak CHH methylation pattern, consistent with the siRNA production profile and RdDM-dependent methylation. Moreover, siRNA levels and CHH methylation decreased as the distance between MITEs or IRs and the associated gene increased (Figure S5B). This distance-dependent epigenetic activity of MITEs and IRs is consistent with previous reports showing greater transcriptional and regulatory activity of elements closer to a gene [64]. This observation supports the idea that 24-nt siRNAs derived from MITEs and IRs near protein-coding genes initiate a non-canonical RdDM pathway likely dependent on Pol II transcription.

In contrast, methylation in the CpG and CHG contexts remained relatively stable regardless of the distance between TEs, MITEs, or IRs and nearby genes (Figure S5B). When these elements were located farther from genes, the surrounding genomic context showed higher CpG and CHG methylation levels, likely reflecting the heterochromatic nature of these regions. Furthermore, CpG and CHG methylation levels were higher in leaves compared to ripe fruits, contrasting with the siRNA production profile and consistent with their independence from siRNA activity (Figure 3D-E, Figure S5B). This contrasts with immature and ripe fruits, which showed higher CHH methylation, suggesting that RdDM activity is required for dynamic regulation of TEs during fruit development and maturation. These differences underscore the distinct epigenetic regulation of transposons across tissues, with stable maintenance of CpG and CHG methylation in somatic tissues and active RdDM-driven CHH methylation in reproductive tissues. This tissue-specific regulatory pattern also reflects the dynamic role of TEs in gene regulation and genome plasticity during fruit ripening and development.

MITEs and IRs exhibit context-dependent regulatory signatures at the genome-wide level

To investigate the potential regulatory roles of gene-hosted MITEs and IRs in gene expression, we performed a differential accumulation analysis of 24-nt siRNAs derived from these elements as a proxy for MITE differential expression and processing. We compared siRNA profiles between leaves and fruits at two ripening stages (immature and ripe) in *F.vesca*. The analysis revealed a distinct production of 24-nt siRNAs derived from MITEs and

IRs across these conditions, considering thresholds of |log2FC| ≥1 and adjusted p-value ≤0.05

(Figure 4A). For MITEs, we identified 17 upregulated and 12 downregulated elements in ripe

fruits compared to immature fruits, 318 upregulated and 89 downregulated in ripe fruits

compared to leaves, and 89 upregulated and 349 downregulated in leaves compared to

immature fruits. Similarly, for IRs, 4 elements were upregulated and 3 downregulated in ripe

fruits compared to immature fruits, 144 upregulated and 28 downregulated in ripe fruits

compared to leaves, and 26 upregulated and 119 downregulated in leaves compared to

immature fruits. This approach allowed us to identify MITEs and IRs actively producing siRNAs

in a specific sample, which may serve as potential regulators of gene expression in a tissue

and stage-specific manner.

We then identified genes located within 3,000 bp of the differentially expressed (DE) MITEs

and IRs and evaluated whether these genes also exhibited differential expression under the

same conditions (Table S3, Table S4). More than 24% of DE MITEs or IRs were located near

DE genes (Figure 4B), with this proportion increasing to 50% during fruit ripening. We also

found a, a comparable proportion of non-differentially expressed (non-DE) MITEs and IRs

located near DE genes across all conditions analyzed. This indicate that the MITE/IR specific

regulation is not necessarily coupled to the transcriptional behavior of the neighboring gene.

These cases may parallel our previous observations at the *HaWRKY6* locus in sunflower,

where tissue-specific gene regulation is mediated by a MITE-triggered chromatin loop, but the

regulatory outcome is controlled by differential DNA methylation at a second anchoring region

rather than at the MITE itself, whose expression remains largely stable across tissues but is a

sine-qua-non requirement for the loop formation [66].

To further assess whether genes associated with MITEs/IRs display distinct transcriptional

behavior, we compared the probability of differential expression among genes located near

MITEs/IRs, genes associated with non-MITE TEs, and genes in TE-free regions. Genes

located in the vicinity (3 kb) of non-MITE TEs showed a significantly lower probability of being

differentially expressed compared to genes in TE-free regions (Fisher's exact test, OR = 0.74,

p = 3.7×10^{-15}), which is expected given that most TE classes are typically located in

repressive genomic contexts characterized by reduced transcriptional activity. In contrast,

MITE/IR-proximal genes displayed a significantly higher likelihood of differential expression

(OR = 1.13, p = 3.3×10^{-5}), indicating that MITE/IR elements are associated with a distinct

regulatory landscape. One limitation in this analysis is that the MITE effect caused by

triggering short-range chromatin loops not necessarily work as an on/off switch but affect

transcriptional dynamics, transcript structure, or regulatory responsiveness without

necessarily producing large expression fold-changes detectable at the genome-wide level
[67].

MITEs and IRs have been also reported to act as enhancers or silencers of gene
expression by generating cis-regulatory elements [71, 72] or, by modulating local chromatin
organization, altering the gene product without changing overall expression levels (e.g., by
triggering alternative splicing or termination) [67]. Their regulatory impact depends on multiple
factors, including their genomic position relative to the gene, their interaction with regulatory
elements such as transcription factors or epigenetic readers, and the way they reshape
chromatin architecture, all of which can vary between conditions.

Given the established role of 24-nt siRNAs in mediating RdDM and chromatin remodeling,
we next integrated siRNA accumulation with DNA methylation and gene expression data to
redefine the identification of MITE/IR-gene pairs with regulatory potential. As a proof-of-
concept, we focused on fruit ripening and analyzed publicly available mRNA-seq data from
strawberries treated with the DNA methylation inhibitor 5-azacytidine (5-AZA) (Figure 4B,
Table S5). Over 35% of DE MITEs/IRs were located near genes that also exhibited significant
expression changes upon 5-AZA treatment, with this proportion exceeding 50% during the
ripening stage. While this overlap alone does not demonstrate causality, the concordant
response of siRNA production, DNA methylation perturbation, and gene expression changes
supports the prioritization of specific loci for downstream analysis.

A MITE Mutator element modulates FvAAT1 expression through epigenetically driven
411
chromatin reorganization
412

Based on above criteria, we examined individual loci in detail as representative case
studies. We focused on elements located near genes with established roles in fruit ripening
and associated metabolic or developmental processes, which are highlighted in bold in
Supplementary Tables S3 and S4. These case studies are intended to illustrate plausible
regulatory configurations supported by epigenetic and transcriptional signals, and are not
intended to imply global enrichment or uniform behavior across all analyzed loci.

During fruit ripening we found 29 up- or downregulated MITEs and 7 up- or downregulated
IRs near genes associated with fruit ripening, metabolic or developmental processes (Figure
4A, Figure 5A). Among them, we identified a 416 bp element which corresponds to a MITE
Mutator, that showed a 12-fold higher siRNA production at the ripening stage (Table S3). This
MITE is located 1,915 bp upstream of the gene *FvAAT1* (FvH4_7g18570) (Figure S6), which
encodes a putative ortholog to alcohol acyl transferase SAAT, a key enzyme responsible for
the biosynthesis of aroma volatile esters in *F. × ananassa [45]*. Consistent with its functional

role, *FvAAT1* displayed substantial upregulation during fruit ripening; at the ripe stage, gene
expression was more than 1,700 times higher than at the immature stage (Table S5), aligning
with previous findings that underscore its importance in producing characteristic ester-based
aroma compounds during ripening [73, 74]. This gene is also upregulated during the turning
stage in *F. × ananassa* compared to *F. vesca*, emphasizing its importance in commercial fruit
quality traits [75]. By contrast, treatment with 5-AZA resulted in a fivefold reduction in *FvAAT1*
expression (Table S5), suggesting that the identified epigenetically active MITE can influence
its expression.

In the same genomic region, we identified another MITE located 1,064 bp downstream of
FvAAT1 and an IR element (IR_9194) located 4,761 bp upstream of *FvAAT1* and 3 bp
downstream of FvH4_7g18550, a gene encoding a papain family cysteine protease implicated
in protein degradation and defense responses during fruit ripening (Figure S6). This gene
showed a marked expression increase (~105-fold) in ripe fruit compared to immature fruit
(Table S5), mirroring patterns observed in other plant species in which cysteine proteases
have been proposed to regulate ripening processes [76-78].

We then evaluated siRNA production and CHH methylation patterns around these
elements to gain further insights into potential epigenetic regulation (Figure 5B and 5C). We
observed that IR_9194 and especially the MITE Mutator produce 24-nt siRNA with higher
levels in fruit ripening, consistent with active elements (Figure 5B, top and middle panel,
respectively). This increase in siRNA production in mature fruits coincides with a CHH
methylation signal (Figure 5C). Interestingly, the MITE located downstream of FvH4_7g18570
displayed the highest methylation levels at its TIRs during immature fruit stage, albeit we did
not observe a significant accumulation of siRNA (Figure 5B and 5C, bottom panel). This may
indicate that this MITE might be in a permanently silenced state, in contrast to IR_9194 and
the MITE Mutator, which remain epigenetically active.

Although such MITE appears to be transcriptionally inactive, the increase in DNA
methylation at this element may serve as an "anchor point" in conjunction with the more
dynamic MITE Mutator element to trigger chromatin looping and control *FvAAT1*, as observed
in other systems [66]. To test the hypothesis that the promotion of a chromatin loop between
these two elements impact the expression of FvAAT1, we designed an experiment to
transiently modulate the epigenetic status of the MITE Mutator. To do so, we infiltrated *F.*
vesca fruits with a construct either overexpressing the viral silencing suppressor P19, which
reduces siRNA activity and thus RdDM, or an additional copy of the MITE Mutator under the
35S promoter to increase the production of its specific siRNAs *in trans* and thus induce
methylation at this element. We have previously used these strategies to induce or repress
MITEs activity [65, 66]. We then measured the expression of *FvAAT1* in infiltrated fruits and

mock infiltrated controls. Consistent with the proposed regulatory effect of the MITE Mutator
over *FvAAT1,* we observed a correlated expression of the gene when RdDM at the MITE was
induced or repressed (Figure 5D). The activation *in trans* of the MITE Mutator correlated with
a significant increase in the expression of *FvAAT1* (Figure 5D). Conversely, the epigenetic
inactivation of the MITE Mutator, achieved by transiently overexpressing either a MITE Mutator
sponge sequence fused to GFP, which sequesters the MITE-derived siRNAs, or the viral
silencing suppressor P19, which nonspecifically sequester siRNAs, correlated with a
significant repression of the gene *FvAAT1* (Figure 5D)*. Furthermore, using a chromatin
conformation capture (3C) approach, we observed a ripening stage dependent change in
chromatin organization, with the MITE Mutator acting most likely as the anchor point (Figure
5E). In this sense, a chromatin loop encompassing the entire loci was increasingly detected
with the maturation of the fruits, suggesting a positive effect on gene transcription, as
previously reported for example during Pol II reclining happening on gene looping events [67].
Given the apparent regulatory importance of the MITE Mutator, we assessed its
conservation in the orthologous genomic regions of the commercial strawberry species *F.
chiloensis* and *F. × ananassa* through sequence alignment. In *F. chiloensis*, a copy of the
element*, situated 1,674 bp upstream of the *FvAAT1* homologue was detected, displaying
89.4% sequence identity (Figure S7A, left). A similar pattern was observed in *F. × ananassa*
(Camarosa genome assembly), where a copy of the MITE associated with *FaAAT1* exhibits
95.7% sequence identity (Figure S7A, right). Analysis of the predicted secondary structures
using RNAfold for the three elements revealed the formation of a long dsRNA hairpin structure,
suggesting that they are likely DCL3 substrates and will function similarly to those observed
in *F. vesca* (Figure S7B). These findings underscore the evolutionary conservation of this
MITE near key ripening-related genes across different *Fragaria* species, potentially reflecting
a conserved regulatory function in fruit quality traits.
We identified additional IR/MITE elements adjacent to genes with key roles in fruit
development and metabolism, suggesting potential regulatory contributions through small
RNA production, DNA methylation, and gene expression changes (Table S4 and Table S5).
For example, a 439 bp IR element (IR_8440) exhibiting approximately a 4.5-fold increase in
expression in red fruit compared to leaf tissue (Table S4) and pronounced CHH methylation
in fruit was identified 1,972 bp upstream of the SEPALLATA-like MADS-box gene
FvH4_6g46420 (Figure S8A). This IR is adjacent to a MITE Harbinger. A MITE hAT located
downstream of FvH4_6g46420 displays relatively constant expression across tissues,
suggesting it could serve as a potential "anchoring" region for 3D chromatin configurations.
The FvH4_6g46420 gene itself is significantly upregulated in white fruit vs. leaf tissue,
consistent with prior studies underscoring the role of its ortholog in *F. × ananassa*, *FaMADS9,*

which is involved in strawberry fruit development and ripening [79]. Silencing of *FaMADS9* in
F. × ananassa delays or inhibits normal ripening and alters both auxin and ABA metabolism,
implicating this MADS-box transcription factor in pleiotropic regulatory networks [80, 81].
Furthermore, *FaMADS9* is downregulated under AZA treatment, highlighting the broader
interplay between epigenetic modifications and fruit ripening pathways [82].

Additional cases reinforce the potential role of IRs and MITEs in regulating nearby gene
expression. A MITE hAT (334 bp) is located near FvH4_2g30230, annotated as molybdate
transporter 1 (Figure S8B). Molybdenum is an essential trace element in plant nutrition and
physiology, and its effects on strawberry flavor and fruit quality are well documented [83]. The
upregulation of this gene in ripe fruit and its downregulation in 5-AZA-treated strawberries,
along with the distinct expression profiles of the MITE in leaf tissue and in fruits at the white
and red ripening stages, together with localized methylation and siRNA mapping, further
support a role for IR-associated epigenetic control. Moreover, IR_6046 (448 bp) is located
upstream of FvH4_5g23180 (Figure S8C), encoding a probable xyloglucan
endotransglucosylase/hydrolase (XTH30), which is critical for cell wall remodeling during fruit
softening [84, 85]. Differential expression of both the element and XTH30, alongside distinct
siRNA clusters in leaf versus fruit ripening conditions, aligns with a mechanism involving
dynamic chromatin organization. Fruit softening is a critical factor affecting the marketability
and economic value of fruits; therefore, targeting cell wall-related genes, including XTHs, may
offer an effective strategy to delay this process and extend postharvest shelf life.

Taken together, these observations underscore the potential of IR and MITE elements to
modulate gene activity through epigenetic pathways, thereby influencing fruit maturation and
quality traits. An idea strongly supported by the validations made over the *FvAAT1* loci.

Discovery of TE Polymorphisms in resequenced *F. vesca* genomes
Due to their inherent mobility, TEs frequently vary between natural populations in copy
number and genomic position, producing insertion or deletion polymorphisms [18]. A TE
insertion polymorphism (TIP) occurs when a TE is present in an accession but absent from
the reference genome, whereas a TE absence polymorphism (TAP) represents a TE found in
the reference but missing in other accessions. Because many TEs harbor their own regulatory
sequences, they can strongly influence the expression of nearby genes, often more than other
types of structural variation [36, 86]. Moreover, our previous results indicate that MITEs and
IRs potentially exert a regulatory effect upon insertion into intergenic regions proximal to genes
[64]. Historically, genetic diversity analyses have primarily focused on SNPs, while TE
insertion/deletion polymorphisms remain an underexplored source of genomic variation [87-
90]. TEs activity is often influenced by stress conditions and many causative TE insertions

result from recent movements and are not effectively tagged by SNPs, meaning TIPs can be
used as direct markers to uncover these adaptive genetic factors [32, 91]. As such, they hold
the potential to affect phenotypes and key biotechnological traits in nutritionally and
commercially important plant species.

We analyzed 210 resequenced *F. vesca* genomes from a European germplasm collection,
representing the species' continental diversity [44] (PRJNA1018297), to study natural variation
in TEs, MITEs, and IRs using TEPID [90] (Figure S9). A total of 8,184 TE, MITEs and IRs
insertions (TIPs) and 11,345 deletions (TAPs) were identified across the population (Table 2).
As reported for other plant species [32, 92, 93], the TE landscape was prevalent in rare
variants; 84% of TE insertions and 88% of IR insertions exhibited a MAF below 5%. This strong
skew toward rare alleles suggests that these insertions are either evolutionarily recent or are
being actively purged by purifying selection to maintain genomic integrity. Notably, deletion
polymorphisms reached higher frequencies in the population than insertions; for instance,
28.7% of TE deletions were classified as common (MAF ≥ 5%) compared to only 15.8% of

insertions. This disparity suggests that many deletions represent the loss of ancestral
elements present in the reference genome that have not reached fixation in the European
germplasm. Among the identified TE insertions, 35% belonged to Class I and 65% to Class II,
closely matching the general TE distribution observed in the reference genome (33% and
67%, respectively). A similar trend was found for deletion polymorphisms, where Class I
elements accounted for 35% and Class II for 65% of the total events Insertion polymorphisms
were particularly enriched for LTR/Gypsy, hAT, Mutatorand MITEs elements, suggesting
recent bursts of activity that have not yet spread across the population [94] (Figure 6A).
Conversely, LTR/Gypsy deletions were notably frequent among common variants, indicating
a significant turnover of these retrotransposons in specific lineages.

A principal component analysis (PCA) based on the identified TE polymorphisms revealed
a clear subpopulation structure (Figure 6B), mirroring the geographical clustering previously
reported using SNP-based markers [44]. In the TE-based PCA, the PC1 and 2 captured ~19%
of the variance, similar to the SNP-based PCA by Toivainen et al. (2026) [44]. This provides
strong confidence in the accuracy and reliability of our detected polymorphisms, as both
genomic features capture the same fundamental evolutionary signals. Our TE-based analysis
also reflected the differentiation reported in the *F. vesca* volatilome by Urrutia et al. [95], such
as the distinct separation of the Icelandic and Baltic Sea clusters. The TE-based PCA also
showed a clear resolution for the Iberian and Scandinavian peninsulas, and a separation of
Western and Southern European accessions from Northern and Eastern populations. In
Scandinavia, this is particularly evident and may be linked to the admixture of western and

eastern post-glacial colonization routes [44]. These results suggest that TE-derived structural variation provides a highly sensitive molecular signature for uncovering fine-scale population structure in *F. vesca*.

The genome-wide distribution of insertion and deletion polymorphisms in *F. vesca* revealed a clear non-random pattern shaped by genome architecture and selective constraints (Figure 6C). Insertions and deletions exhibited partially distinct patterns. Deletions showed a stronger avoidance of exonic regions than insertions, in line with their greater potential to compromise gene integrity (Figure S10A). Across all TEs superfamilies, polymorphisms were strongly depleted within gene bodies, with the most pronounced avoidance observed in exons and untranslated regions (UTRs), consistent with strong purifying selection acting against disruptive structural variation in functional elements (Figure 6C, Figure S10B). Introns showed a comparatively milder depletion, indicating a higher tolerance for structural variation relative to coding and regulatory regions. In contrast, polymorphisms were consistently enriched in gene-flanking regions, where chromatin is typically less condensed, likely facilitating access for the TE mobilization machinery (Figure 6C, Figure S10). Similar patterns have been observed in other species, where TE insertions tend to occur in gene-rich regions, although variations can occur at both the TE family and species levels [36, 94, 96]. Overall, more than half of all identified polymorphisms were located within genes or their immediate flanking sequences (up to 1,000 bp upstream/downstream). Specifically, MITEs and IRs showed a preferential accumulation within promoter regions (Figure 6C, Figure S10B). This high frequency in upstream regions and to some extent downstream, suggests a significant potential for these elements to influence gene expression, as observed in the results described above or in other crop species, like *Brassica rapa* and *Solanum lycopersicum* [36, 37].

Taken together, these results suggest that the landscape of TE polymorphisms in *F. vesca* is characterized by a high turnover of elements within regulatory proximity to genes. The robust population structure resolved by these variants underscores the role of TEs as dynamic footprints of post-glacial colonization. The high frequency of rare variants suggests that most transposable element polymorphisms (TEPs) sare subjected to purifying selection or occurred recently in certain lineages. Moreover, these TEPs occur preferentially in promoter regions. Collectively, this represents a significant and underexplored reservoir of genetic diversity with the potential to impact complex phenotypic traits in European woodland strawberry germplasm.

Linking TE polymorphisms to phenotypic traits in woodland strawberry fruits

Several phenotypic traits, including fruit dimensions (length, width, volume) and the concentrations of 99 volatile secondary metabolites (the volatilome), were previously

measured in a collection of 116 and 154 wild *F. vesca* accessions across two independent
harvest seasons (H16 and H17) [95]. Since the genomes of these accessions were included
in our polymorphism analysis, we leveraged the detected TEPs to perform a genome-wide
association study (GWAS) to investigate the impact of TE and IR natural diversity in those
strawberry fruit traits.

Several phenotypic traits, including fruit dimensions (length, width, volume) and the
concentrations of 99 volatile secondary metabolites (the volatilome), were previously
measured in a collection of 116 and 154 wild *F. vesca* accessions across two independent
harvest seasons (H16 and H17) [95]. Since the genomes of these accessions were included
in our polymorphism analysis, we leveraged the detected TEPs to perform a genome-wide
association study (GWAS) to investigate the impact of TE and IR natural diversity in those
strawberry fruit traits.

For the volatilome dataset, 68 distinct TEPs were significantly associated (5% Bonferroni
threshold) with 40 VOCs across harvests H16 and H17 resulting in 78 significant trait–
polymorphism associations (Table S7). None of these associations corresponded to TEP
directly impacting the coding sequence of genes known to be involved in strawberry VOC
metabolism; this is consistent with our observation that only a very small fraction of these
variants occur in exonic regions and tend to occur more frequently in intergenic regions
proximal to genes where they could exert a regulatory effect (Figure 6C). This is also
suggested in the results in this work presented, in which we observed the regulatory effect of
different MITEs on the expression of *FvAAT1*.

Considering our results, we next focused on statistically significant associations that exhibit
potential biological relevance regulating the genes in their vicinity based on current knowledge,
specially focusing on MITE or IR polymorphisms. For instance, a MITE Mutator insertion was
associated with 6-methyl-5-hepten-2-one (sulcatone) on chromosome Fvb3 (P = 5.6512 x 10^{-6};
PVE = 9.20%) (Figure 7A-B, Figure S12A). Sulcatone is a common apocarotenoid that
originates from lycopene cleavage [97]. Apocarotenoids contribute to the aromas of different
flowers and fruits and are highly valued by the flavor and fragrance industry [97]. Unsaturated
methyl ketones, such as sulcatone, have been proposed to act as highly reactive VOCs with
defensive and pesticidal properties and have been associated with unpleasant taste in fruits
affected by powdery mildew disease [98-100]. Accessions carrying this insertion (n = 34)
showed lower 6-methyl-5-hepten-2-one levels in harvest H17 (Figure 7C). Interestingly, the
insertion is located in a gene-dense genomic region (FvH4_3g02700, FvH4_3g02710,
FvH4_3g02720, FvH4_3g02730 and FvH4_3g02731 genes) enriched in stress- and ROS-
related signalling components and redox enzymes, including LEA/NDR1-like (LEA-2) proteins,
NDR1 a secretory class III peroxidase, a cysteine-rich receptor-like kinase (CRK-like), and an

LpxD-like N-acyltransferase. Class III peroxidases act as major apoplastic regulators of H_2O_2

homeostasis and oxidative reactions, shaping the local redox environment that feeds into

downstream oxidative volatile formation [101]. In parallel, NDR1 is a central component of

plant defence signalling, and CRK family kinases are widely implicated in ROS-associated

signal transduction during stress responses. The presence of a MITE element in this gene

module could impact the regulation of these genes, ultimately affecting the response to

oxidative stress and indirectly influencing the oxidative flux required for carotenoid degradation

and the sulcanone levels.

A Harbinger Class II TE deletion was associated with lower levels of the 6-C aldehyde

green leaf volatile (GLV) (E)-2-hexenal in H17 (P = 9.51 x 10^{-8}; PVE = 2%) (Figure 7D-F). GLV

are VOCs emitted in response to wounding or plant damage and are responsible for the

characteristic "green" aroma of plants (reviewed in [102]). The deletion is located on

chromosome Fvb1, 13 kb upstream of a gene annotated as 2-ALKENAL REDUCTASE

(FvH4_1g00330). Alkenal reductases facilitate the reduction of double-bonds in aldehydes,

including short-chain aldehydes such as (E)-2-hexenal [103, 104]. This gene, or any of its

eight downstream paralogs, represents a potential candidate for the conversion of (E)-2-

hexenal to hexanal in strawberry. Notably, this locus has not been identified in previous QTL

analysis in *F. vesca* or cultivated strawberry [105, 106], highlighting the potential of TEPs-

GWAS to identify novel QTLs.

Other associations further suggest linkage between the TEP and the underlying candidate

gene. A primary example is methyl 2-aminobenzoate (methyl anthranilate), a precursor of

tryptophan involved in primary metabolism and associated with a grape-like aroma in fruits

[107]. We identified a significant association with the deletion of a MITE Mutator on

chromosome Fvb5 (P = 1 x 10^{-14}; PEV = 0) (Figure S13A-C). Interestingly, although this variant

appears to explain only a marginal amount of phenotypic variance, this locus has been

consistently associated with methyl anthranilate levels in octoploid strawberry [108]. Barbey

et al. described a QTL at the beginning of chr5-4, the same region where the TEP is located.

This site is 380 kb from an L-tryptophan—pyruvate aminotransferase (TAA1) gene

(FvH4_5g05880), which utilizes L-tryptophan for auxin biosynthesis; this gene may be related

to varying levels of the intermediate methyl anthranilate.

Our analysis also identified a MITE Harbinger deletion on chromosome Fvb3 that was

associated with γ-decalactone levels across both harvest's seasons (P = 4.2911 x 10^{-6} in H16

and P = 1.4523 x 10^{-6} in H17) (Figure S13D-F, Figure S12B). This variant appears to explain

only a marginal amount of phenotypic variance in both harvests. Accessions carrying this

deletion (n = 87 and 115 in H16 and H17, respectively number) showed higher levels of γ-

decalactone, a key compound previously identified as one was one of the most variable in the

F. vesca volatilome analysis by Urrutia et al. [95]. In *F. × ananassa*, γ-decalactone is the most

abundant lactone [79], reaching maximum levels in fully ripe fruits [79] and is positively

correlated with consumer perception of favorable organoleptic traits [95, 109]. The deletion

falls within an intron of the gene FvH4_3g30910, annotated as a replication factor-A C-terminal

domain protein, which shows no evident functional connection to γ-decalactone. Notably,

however, the deletion lies ~5 kb downstream of FvH4_3g30901, a gene encoding a serine

carboxypeptidase-like (SCPL) enzyme. SCPL enzymes belong to the α/β-hydrolase

superfamily, functioning as acyltransferases in plant secondary metabolism which include

esterases and acyltransferases that act on diverse lipid substrates [110, 111]. Given the

functional plasticity of these enzymes, it is likely that many SCPL members involved in fruit

specialized metabolism are yet to be characterized. In this context, structural variation

affecting chromatin state or *cis*-regulation near the SCPL gene could indirectly contribute to

quantitative differences in γ-decalactone.

Taken together, these results suggest that natural polymorphisms in transposable

elements, particularly MITEs and IRs, are linked to variation in strawberry fruit volatilome

composition, primarily through effects in gene-proximal regions. While these associations do

not imply direct causality, they reveal candidate regulatory loci whose effects are shaped by

local genomic and chromatin context.

DISCUSSION

Our study provides a comprehensive analysis of TEs in *Fragaria vesca*, highlighting their

role in genome composition, organization, and potential functional impact. By systematically

annotating TEs and analyzing their polymorphisms across 210 *F. vesca* accessions, we have

gained valuable insights into their evolutionary significance and their potential contributions to

gene regulation and phenotypic diversity.

Our TE annotation revealed that TEs occupy ~30% of the *F. vesca* genome, a proportion

consistent with expectations for plant species of similar genome size and slightly higher than

the 22% reported in the initial 2011 *F. vesca* genome assembly [41]. Unlike most plant

genomes, where LTR retrotransposons are dominant [54] *F. vesca* exhibits a comparable

proportion of Class II DNA transposons, with MITEs representing a substantial fraction of intact

elements. This enrichment is likely explained by their compact structure and lack of coding

regions, which reduce their susceptibility to fragmentation over evolutionary time. In contrast,

LTR retrotransposons, particularly the larger elements, appear more frequently fragmented,

consistent with higher rates of structural decay. The genome-wide distribution of TEs further

reveals distinct patterns among classes: LTRs are enriched in gene-poor, pericentromeric

regions, although their relatively low abundance in *F. vesca* may reflect incomplete assembly

of centromeric regions, whereas MITEs, IRs, and other DNA transposons are concentrated near or within coding sequences, suggesting a potential role in the regulation of adjacent genes. These observations underscore the unique composition of the *F. vesca* genome and provide a foundation for exploring how TE-mediated regulatory variation contributes to phenotypic diversity in strawberry. One of the most significant findings of our study is the widespread production of small RNAs (siRNAs) from MITEs and IRs in *F. vesca*, supporting their potential regulatory role in gene expression. On average, 71% of these non-autonomous elements produced 24-nt siRNAs in at least one tissue type studied, with a characteristic two-peak siRNA profile indicative of processing through the RdDM pathway. The correlation observed between siRNA production and CHH methylation, particularly in reproductive tissues, underscores the functional relevance of these elements in tissue-specific gene regulation. In contrast, CpG and CHG methylation patterns remained relatively stable, supporting the role of siRNA-driven CHH methylation as a key mechanism controlling TE activity and gene expression in a dynamic and reversible manner. These findings highlight the functional significance of MITEs and IRs as autonomous regulatory elements that modulate gene expression through small RNA-mediated epigenetic modifications. These observations position MITEs and IRs not merely as silencing targets, but as active components of regulatory networks capable of modulating local chromatin states.

While genome-wide analyses revealed that differentially expressed MITEs and IRs are not significantly more likely than non-DE elements to be located near differentially expressed genes, this finding underscores an important conceptual point: TE proximity alone is insufficient to infer regulatory function. This is not entirely surprising as we previously found that one of the most important features of MITE affecting gene expression is not their sequence but their capacity to trigger chromatin loops, which inevitably require an additional anchor point. In a previous study we showed that, if a MITE is expressed, its regulatory function can be modulated by alterations (methylation, accessibility, etc.) on the second anchor point [66]. The lack of global enrichment of differentially expressed genes near DE MITEs or IRs during ripening further indicates that these elements do not act alone. Instead, their regulatory potential appears to emerge from the convergence of multiple features, including tissue-specific siRNA production, local methylation dynamics, transcriptional responsiveness, and the genomic neighborhood.

This concept is exemplified by the detailed analysis of selected loci associated with fruit ripening. At the *FvAAT1* locus, a MITE Mutator element displays coordinated increases in siRNA accumulation and a clear CHH methylation signature during ripening, coinciding with dramatic upregulation of the adjacent gene encoding an alcohol acyltransferase, a key enzyme responsible for volatile ester biosynthesis in strawberry [45, 112, 113]. The

upregulation of *FvAAT1* during ripening, coupled with siRNA production from the adjacent
MITE, supports a model in which this TE influences gene expression through methylation-
dependent chromatin remodeling. Further validation using 5-AZA demethylation assays
confirmed that *FaAAT1* expression is methylation-dependent, highlighting the direct regulatory
effect of MITEs in *F. vesca* fruit ripening. The conservation of this particular MITE in *F.*
chiloensis and *F. × ananassa*, with high sequence identity and structural similarities, suggests
that it plays a conserved role in fruit aroma regulation across strawberry species. This
discovery adds to the growing evidence that TEs contribute to the evolution of crop traits by
introducing regulatory variation that affects key metabolic pathways [7, 36, 114, 115]. It also
positions TEs, especially gene-associated MITEs and short IRs, as promising targets for
genome editing. The CRISPR/Cas9-mediated removal of a MITE could significantly alter the
expression of adjacent genes and the associated phenotypic traits. Genome editing of MITEs
also offers the advantage that these elements are non-coding and naturally variable in
populations, potentially simplifying regulatory approval.

In addition to their regulatory potential, TE polymorphisms provide an important source of
genetic diversity that can influence trait variation [37]. Our GWAS analyses link TE
polymorphisms to volatile composition, demonstrating the phenotypic relevance of mobile
elements in *F. vesca*. Notably, some of the identified TEPs correspond to previously
unreported loci affecting key flavor compounds, such as sulcatone, γ-decalactone, and methyl
anthranilate, highlighting the potential of TEP-focused GWAS to uncover novel regulatory loci
that are missed by traditional SNP-based approaches. While these associations do not
establish direct causality, they provide candidate regulatory elements for functional validation
and suggest that integrating TE polymorphisms into breeding programs could expand the
spectrum of allelic variation available for trait improvement. Our findings demonstrate that TE-
driven variation significantly contributes to the natural diversity of fruit metabolic profiles, with
potential implications for breeding strategies aimed at improving fruit aroma and flavor.
Furthermore, understanding how TEs modulate gene expression and metabolite production
at the molecular level provides valuable insights into the complex regulatory networks
underlying crop traits. The high degree of TE polymorphism observed in *F. vesca*, combined
with the functional relevance of associated traits, suggests that TEs represent an untapped
source of genetic variation for crop improvement.
780

CONCLUSIONS
Our findings demonstrate that, as in other plants, TEs in strawberries are not merely
passive genomic elements but rather key contributors to genome function, gene regulation,
and phenotypic variation. By integrating TE annotation, small RNA profiling, methylation data,

gene expression analyses, and TEP-based GWAS, we identify specific TE–gene
configurations with plausible regulatory roles in fruit development and quality traits. Although
TE proximity alone does not predict regulatory function at the genome-wide level, selected loci
illustrate how epigenetically active MITEs and IRs can modulate gene expression in
biologically relevant contexts, such as fruit ripening.

This identification of TE-derived regulatory elements and polymorphisms associated with
fruit traits provides a valuable resource for future breeding efforts aimed at improving
strawberry quality. In particular, the regulatory roles of MITEs and IRs in fruit ripening suggest
that targeted epigenetic modifications could be leveraged to optimize fruit quality. The use of
genome-editing tools such as CRISPR/Cas9 to precisely manipulate TE-derived regulatory
regions may enable the fine-tuning of gene expression without introducing foreign DNA,
making it a promising strategy for crop improvement.

Additionally, the conservation of key regulatory MITEs in commercial strawberry species
suggests that similar TE-driven mechanisms may underlie important agronomic traits in
cultivated varieties. Further comparative genomic studies across *Fragaria* species could help
elucidate the broader evolutionary impact of TEs and their role in shaping fruit diversity.

METHODS

Transposable element identification and classification

The genome sequence of *Fragaria vesca* and annotation files (v4.0.a2) were downloaded from
the Phytozome web portal at https://phytozome-next.jgi.doe.gov/ [116]. TEs were detected
and preliminarily classified using the EDTA v2.1.0 [48], executed as follows: `EDTA.pl --`
`genome $genome --exclude $annotation_bed --anno 1`. where `$genome` is the *F.*
vesca reference genome, and `$annotation_bed` is the assembly annotation converted to
BED format from the original GFF3 file using the *gff2bed* tool from BEDOPS v2.4.41 [117].
EDTA produces a filtered non-redundant TE library for annotation of structurally intact and
fragmented elements. TEs are classified into several non-overlapping categories, including
LTR (referring to LTR retrotransposons), non-LTR (including SINEs and LINEs), TIR (referring
to DNA transposons with TIRs, including MITEs), Helitron, and non-TE repeat sequences.
TEs identified by EDTA were reclassified with DeepTE [49] that uses convolutional neural
networks and detects domains inside TEs to correct false classification. Within Class II,
DeepTE allows for the subclassification of MITEs and non-MITEs (DNA TEs). DeepTE was
run with the following command: `DeepTE.py -d /path/intermed_files -o`
`/path/results -i $TE_lib -sp P -m_dir /path/Plants_model`. Where `$TE_lib`
is the FASTA file with the TE sequences detected by EDTA, the parameter `-sp P` specifies

a plant organism, and the `-m_dir` flag indicates the path to the training model. A custom

bash script was then used to merge the GFF3 EDTA output files with the DeepTE classification

(TE order and superfamily). For TEs categorized as "unknown" by DeepTE, the original EDTA

classification was preserved, translating it into the classification used in this work (Figure 1B).

The consensus TE library was then curated using TEtrimmer [118], a tool designed to

automate the manual curation of TEs, which performs automated consensus sequence

curation, clusters related elements, and removes duplicates. This step markedly improves TE

library quality and facilitates more accurate repeat masking and classification. TEtrimmer was

run with default parameters and the `--classify_all` option. Then, the consensus library

file after de-duplication step was used as input for RepeatMasker to finally annotate the

curated TE library.

Two GFF3 files were generated with the genomic coordinates and assigned TE

class/superfamily. One includes all TEs (including fragments), while the other contains only

structurally intact TEs, that is, those containing all domains corresponding to their superfamily.

TE-derived IRs were classified by grouping sequences based on their sequence identity, using

the CD-HIT-EST module of CD-HIT v4.8.1 [119], as previously reported for *Arabidopsis*

thaliana [63], running the following command: `cd-hit-est -i $fa -o $output -c 0.8`

`-A 0.8 -T 2 -n 5`. A minimum sequence identity threshold of 80% (`-c`), a minimum

alignment coverage of 80% (`-A`), and a word size of 5 (`-n`) were set. Next, DeepTE was run

as previously described for TE annotation, using the representative sequences from each

cluster to classify the IRs. When IRs overlapped with annotated TEs but were not initially

classified as Class II, they were reclassified based on the overlapping element's category

using a custom bash script. The presence of transposase-associated domains in Class II

elements was evaluated using the DeepTE_domains.py module, which detects conserved

protein domains in TE sequences, applied to both Class II consensus sequences and genome-

wide annotated copies.

IR and MITE analysis

Inverted repeats were identified in the *F. vesca* v4 genome using *einverted* from the EMBOSS

program suite v6.6.0.0 [68]. The following parameters were used: maximum repeat of 700 bp,

a gap penalty of 8, a minimum score threshold of 150, a match score of 3, and a mismatch

score of −4. A custom Perl script was used to parse the output and generate a GFF3 file with

the IR annotation. To classify IRs as MITE-independent elements, we used bedtools intersect

v2.26.0 [59], discarding elements that exhibited more than 20% overlap between MITEs and
IRs.

To assess the conservation of the MITE Mutator, the Tripal MegaSearch tool from the Genome
Database for Rosaceae (GDR) website (https://www.rosaceae.org/) was first used to identify
orthologous genes in the strawberry species *Fragaria chiloensis* (KIB CAS genome assembly)
and *Fragaria × ananassa* (Camarosa genome assembly). A 3,000 bp sequence upstream of
each gene was retrieved from JBrowse (GDR) for the respective genomes and locally aligned
with the *F. vesca* MITE Mutator sequence using EMBOSS Water
(https://www.ebi.ac.uk/jdispatcher/psa/emboss_water), which employs the Smith-Waterman
algorithm for pairwise alignment.

The predicted secondary structures with minimum free energy for each MITE were generated
using the RNAfold web server (http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi)
with default parameters.

Plant material and growth conditions
Plants from the *F. vesca* accession 'Reine des Vallées' used in this study were grown and
maintained in a greenhouse under natural sunlight conditions (IHSM "La Mayora", Málaga,
Spain), using a mixture of universal substrate and river sand (3:1 ratio by volume). Fruits at
the immature (white) and ripe (red) stages were collected for small RNA-seq (sRNA-seq) and
RNA-seq transcriptome analyses. Strawberry fruits infiltrations were performed as previously
described [120]. MITE-Mutator siRNA sponge was created by fussing half of its sequence
(only one of the TIR to prevent dsRNA hairpin formation) with the GFP cloning sequence.

sRNA-seq
RNA was extracted as previously described [121] from fruit receptacles at two developmental
stages, immature (white) and ripe (red). In detail, for each stage, two biological replicates were
prepared; each biological replicate was composed of 30 receptacles after the removal of the
achenes. RNA quality of the samples was measured on a Bioanalyzer 2100 (Agilent
Technologies, Santa Clara, CA). RNA purity was checked using the NanoPhotometer®
spectrophotometer (IMPLEN, CA, USA). RNA integrity was analyzed by lab-on-a-chip
electrophoresis using the RNA Nano 6000 Assay Kit with the Agilent Bioanalyzer 2100 system
(Agilent Technologies, CA, USA). Quantification of samples was performed with Qubit® RNA
Assay Kit in the Qubit® 3.0 Flurometer (Life Technologies, CA, USA). Subsequently, 1 µg of
total RNA per sample was used to generate small RNA libraries using the Truseq® Small RNA
Library Prep Kit according to manufacturer's instructions (Illumina Inc., San Diego, CA, USA).
Library pool was sequenced on the Illumina Nextseq550 instrument at the University of Málaga

889 (Spain), with 1 × 75 cycles according to the manufacturer's protocol. Publicly available sRNA-

890 Seq reads from *F. vesca* leaves were retrieved from GEO under accession number

891 GSM1513956 and analyzed together with immature and ripe reads.

892 The small RNA reads were first cut to remove 3' adapters using cutadapt v4.9 and their quality

893 checked using FastQC v0.12.1 (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/)

894 and MultiQC v1.25.1 [122]. They were then mapped with ShortStack v4.1.1 [123] to *F. vesca*

895 v4 genome using the following parameters: `--locifile` (with MITE or IR annotation files), `-`

896 `-nohp` (prevents search for microRNAs) and `--make_bigwigs`. The DESeq2 Bioconductor

897 package ([124], R Core Team 2022) was used to normalize the total raw read counts from

898 ShortStack for differential expression analysis. MITEs/IRs with 10 or more 24-nucleotide reads

899 across their entire region in all samples were considered to have potential siRNA production.

900 After count normalization, the expression of IR/MITE-derived siRNA was compared between

901 leaf tissue vs. immature stage, ripe vs. immature stages, and ripe stage vs. leaf tissue, using

902 the negative binomial test. Differentially expressed MITEs/IRs were determined by adjusting

903 p-values with the Benjamini–Hochberg false discovery rate, and those with an absolute log2

904 fold change ≥1 and an adjusted p-value ≤0.05 were defined as differentially expressed

905 elements.

907 **RNA-seq**

908 Total RNA was isolated as described previously [120] using the same protocol applied to the

909 fruit receptacles used for sRNA-seq analysis. For the RNA-seq experiment, however, three

910 independent biological replicates were prepared per developmental stage. After validating the

911 quality and quantity of the total RNA using Bioanalyzer 2100 (Agilent Technologies Santa

912 Clara, CA, USA) and Qubit 2.0, libraries were prepared using Illumina's TruSeq RNA Sample

913 Prep Kit v2. Pools of libraries were sequenced by paired-end sequencing (100x2) in an

914 Illumina HiSeq 2500 sequencer by Sistemas Genómicos (Valencia, Spain). Also,

915 transcriptome data for leaf tissue (GEO: GSM3243286 and GSM3243287), fruits treated with

916 5-AZA and its control (GEO: GSE151534) produced by previous studies were used [82, 125].

917 The analysis started by quality trimming and filtering the raw reads with Trimmomatic v0.39

918 [126], the reads were then aligned to the *F. vesca* v4 reference genome using STAR v2.7.11b

[127] which was guided by the gene and exon annotation from *F. vesca* v4.0.a2. *Sort* and

bamCoverage tools from Samtools v1.21 [128] were then used to sort BAM files by genomic

coordinates and generate normalized coverage tracks for visualization and further analysis,

respectively. Read quality before and after trimming was analyzed with FastQC v0.12.1

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and, together with mapping

efficiency, they were summarized with MultiQC v1.25.1 [122]. Read counts on each gene were

then calculated with featureCounts v2.0.8 [129]. This pipeline was run with the aid of the

Snakemake workflow engine [130]. Differential expression analysis was performed by the

DESeq2 package [124], filtering out genes with counts below 10 in all samples. Genes were

then annotated using the information (BLAST, InterPro, GO term, GO accession) provided by

the GDR database (https://www.rosaceae.org/).

BS-seq

Publicly available Bisulfite-Seq data [125] from *F. vesca* leaf tissue, immature receptacle and

ripe receptacle were retrieved from GEO under accession numbers GSM3243379,

GSM3243380 and GSM3243381, respectively. Although it is not specified whether tissues

were collected from the same individual plants, the observed methylation trends, including

global hypomethylation and consistent DMRs across replicates, support their reliability for

integrative methylome analysis. The EpiDiverse WGBS v1.0 pipeline for bisulfite read

trimming,   mapping   and   methylation   calling   was   implemented

(https://github.com/EpiDiverse/wgbs), which was specifically designed for non-model plant

species [131]. For each sample, bedGraph files corresponding to each sequence context

(CpG, CHG, CHH) were obtained. Then *bedGraphToBigWig* v.2.10 from ENCONDE

(https://www.encodeproject.org/software/bedgraphtobigwig/) was used to generate coverage

tracks for visualization and plotting.

3C assay and RT–qPCR

The 3C assay was conducted following a published procedure [66]. Approximately 2 g of tissue

was collected per sample and crosslinked with 1% formaldehyde. Crosslinking was stopped

by adding glycine to a final concentration of 0.125 M, after which nuclei were isolated. To

detect FvAAT1 loops DNA was digested with XhoI (Thermo Fisher Scientific). DNA fragments

were ligated with 25 U of high concentrated T4 DNA ligase (Thermo Fisher Scientific) at 22

°C for 5 h in a 2-ml volume. Subsequent steps included reverse crosslinking, proteinase K

treatment (Qiagen) and DNA purification with the phenol–chloroform. To measure interaction

frequency, qPCR was performed, and results were analyzed with the 2−ΔΔCt formula using

ACT2 as a control. The final qPCR products were subjected to agarose gel electrophoresis

(2%) and bands corresponding to the expected size were purified and sequenced using the

Sanger method to confirm ligation products in the case of 3C experiments. For RT–qPCR, 500

ng of total RNA underwent RT using the RevertAid RT kit (Thermo Fisher Scientific) after

treatment with DNase (Thermo Fisher Scientific). Subsequently, qPCR was conducted using

SYBR green (Thermo Fisher Scientific Maxima SYBR Green qPCR Master Mix (2×)). Relative

expression levels were determined using the 2−ΔΔCt method with *FvCHP1* as the

housekeeping gene control.

Identification of TE Polymorphisms

A set of 217 resequenced woodland strawberry genomes from a European germplasm

collection (PRJNA1018297), representing the species' continental diversity, was used for the

polymorphism study [44]. Detailed information on each accession's geographic origin, read

depth, and chromosomal coverage can be found in Table S6. Since the study of TE

polymorphisms requires a coverage over 15X and no less than 10X [90], resequencing data

was analyzed using SAMtools *coverage* v1.17 [132]. The results were processed using

custom bash and R scripts. Next, the accessions were filtered based on a mean per-

chromosome coverage exceeding 10X. Consequently, seven of the 217 available accessions

(Alta, FIN55, NOR2, NOR-P1-10, RUS3, UK3, and UK9) were excluded from subsequent

analyses, resulting in a final set of 210 accessions.

TEPID v0.10 [90] was employed for the detection of polymorphisms, using the curated non-

redundant TE library, and the IR annotation. The three stages of the TEPID algorithm were

executed using the commands listed below:

Initial Mapping Stage:

`tepid-map -x $bowtie_index -y $yaha_index -p $threads -s $insert_size`

`-n $reseqid -1 $r1 -2 $r2 -z`

Where `$bowtie_index` and `$yaha_index` indicate the paths to the reference indices

generated by each program; `$insert_size` refers to the average insert size of the

sequencing library, set to 250 bp based on data obtained using the *CollectInsertSizeMetrics*

module from Picard v3.1.1 (https://broadinstitute.github.io/picard/); `$reseqid` represents the

genome under analysis; `$r1` and `$r2` denote the forward and reverse reads, respectively;

and the `-z` flag is used to handle compressed FASTQ files.

<u>Discovery Stage:</u>

`tepid-discover -p $threads --name $reseqid --conc $conc_BAM --split`

`$split_BAM --te $TE_bed`

Where `$reseqid` is the genome identifier; `$conc_BAM` and `$split_BAM` are the Bowtie2

[133] and YAHA [134] alignment files resulting from the mapping stage, respectively; and

`$TE_bed` is the BED file with TE annotations in the format specified by TEPID.

After the discovery stage, an intermediate step involved merging the TE insertions and

deletions detected in the population into a single BED file using the TEPID scripts

`merge_insertions.py` and `merge_deletions.py`. Briefly, insertions of the same TE

were merged if their coordinates were within ±100 bp, generating a list of accessions

containing that insertion. Similarly, TE deletions were merged to create a file representing all

deletions detected in the population, along with the list of accessions containing each deletion.

<u>Refinement Stage</u>

`tepid-refine -p $threads --insertions $insertions --deletions`

`$deletions --name $reseqid --conc $conc_BAM --split $split_BAM -a`

`$reseq_list --te $TE_bed`

Where `$insertions` and `$deletions` contain the identified insertion and deletion

polymorphisms for all samples in the population; `$reseqid` is the genome identifier to run

the *refine* command on; `$conc_BAM` and `$split_BAM` are the Bowtie2 and YAHA alignment

files, respectively; `$reseq_list` is the list with all the genome identifiers in the population;

and `$TE_bed` is the TE annotation BED file.

This step, designed to reduce false-negative calls, re-evaluates genomic regions with less

stringent settings where a TE variant was identified in at least one sample of the population

but not in the focal sample. It produces two types of BED files for each accession: (i)

second_pass, referring to polymorphisms not detected in the discovery step but rescued after

re-evaluating the region, and (ii) ambiguous, referring to TE presence/absence calls that could

not be reliably determined during the refinement step. For each accession, second_pass

insertions and deletions were combined with the variants discovered in step 2. The resulting

final insertion and deletion files for each accession were then merged using the scripts

`merge_insertions.py` and `merge_deletions.py`.

The resulting BED files were processed using custom R scripts.

To determine the genomic distribution of TE polymorphisms (Figure 6C and Figure S10) we

developed a custom script using R packages GenomicRanges and GenomicFeatures [135].

Insertions and deletions were mapped against the *Fragaria vesca* v4.0.a2 reference

annotation. Genomic features were defined hierarchically to resolve overlapping annotations:

regions 1 kb upstream of the transcription start site (TSS) were classified as promoters

("Upstream"), followed by 1 kb downstream of the transcription termination site (TTS). Within

gene bodies, determining priority was given to Exons, followed by UTRs (5' and 3'), and finally

Introns. Polymorphisms falling outside these regions were classified as distal intergenic (>1

kb or <-1 kb). To assess whether the observed distribution deviated from random expectation,

we calculated the "Genomic fraction" occupied by each feature type. This was calculated by

collapsing the reference genome into non-overlapping unique base pairs (unstranded) to

represent the physical target size available for mutation. Enrichment or depletion was

quantified using the Log_2 ratio of the Observed frequency to the Expected frequency

(percentage of the genome occupied by a specific feature).

1031

Genome-Wide Association Study

GWAS was conducted on the 4395 and 4340 polymorphisms present in 116 and 154

accessions for harvests H16 and H17 [91], respectively, with a MAF ≥ 5%. The association

analyses were performed by GAPIT (version 3.5) [136] using a Multi-Locus Mixed Model

(MLMM) [137]. The model included the three first principal components derived from all the

polymorphisms used for each group of accessions (H16 and H17) (Figure S11) as covariates

to reduce false positives due to population stratification. Additionally, a SNP-based kinship

matrix capturing the relationship between accessions was included as random effects. Biallelic

variants with a minor allele frequency greater than 0.01 were retained for kinship estimation

using the VanRaden method [138]. The Bonferroni correction was used to establish a

threshold of significance for all the associations.

1043

Data processing and visualization

To calculate the number of genes near each type of TE and differentially expressed

MITEs/IRs, the tool *slop* and *intersect* from bedtools v2.26.0 [59] were used for distances of

500, 1000, and 3000 bp both upstream and downstream of each element. Additionally,

intersect was used to count elements located within coding regions, utilizing different

annotation files for each feature.

To evaluate whether MITEs or IRs are preferentially associated with gene expression changes

relative to other genomic contexts, genes were classified into mutually exclusive categories

based on their proximity (≤ 3 kb) to: (i) MITEs or IRs, (ii) other TE classes, or (iii) no annotated

TE (TE-free regions). Each gene was assigned to a single category based on genomic overlap
identified using bedtools intersect. For each comparison, genes were further classified as DE
or non-DE using the list of DE genes identified from the RNA-seq analysis comparing mature
versus immature fruit stages. Contingency tables were constructed, and statistical significance
was assessed using Fisher's exact test. Odds ratios (OR) and p-values were used to estimate
the relative likelihood of differential expression across genomic contexts.
In order to evaluate the global methylation and siRNA level, metagene analysis was performed
using deepTools v3.5.5 [125]. First, *computeMatrix* was executed to generate read abundance
from all samples over gene, TE, MITE and IR regions. This matrix was then used to create,
using *plotProfile*, a metagene profile from 2 kb upstream of the Transcription Start Site (TSS)
to 2 kb downstream of the Transcription End Site (TES).
Moreover, circos ideograms were created using Circos software vv0.69-6 [126], displaying
only the assembled chromosomes due to their higher density of gene and transposon
annotations. Also, JBrowse2 desktop v2.18.0 [127] was used to visualize gene and MITE/IR
annotations, along with RNA-seq and BS-seq data, at specific *loci* in the *F. vesca* v4 genome.
Principal Component Analysis (PCA) was performed using the R package prcomp [139]. Data
obtained from the different analyses of TE annotation and from the sequencing experiments
were further processed, statistically analyzed, and plotted in R using a diversity of packages,
including dplyr (https://github.com/tidyverse/dplyr), tidyr (https://github.com/tidyverse/tidyr),
scales (https://github.com/r-lib/scales), forcats (https://github.com/tidyverse/forcats), ggplot2,
RColorBrewer (https://CRAN.R-project.org/package=RColorBrewer), MetBrewer
(https://github.com/BlakeRMills/MetBrewer), rtracklayer [140], and ggman
(https://github.com/drveera/ggman).
1076
1077 **Ethics approval and consent to participate**
1078 Not applicable.
1079
1080 **Consent for publication**
1081 Not applicable.
1082
1083 **Availability of data and materials**
1084 The datasets generated during the current study are available from European Nucleotide
1085 Archive (ENA) under the accession number PRJEB86847. The curated TE, MITE, and IR
1086 libraries, as well as the code for TE polymorphism and GWAS analyses, are available at:
1087 https://github.com/spriego/PriegoCubero_Tolley_et_al_Fvesca
1088

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by grants from the Spanish Ministry of Science and Innovation (PID2022-137037NB-I00 and CNS2023-145312 to P.A.M., and PID2021-123677OB-I00 to D.P.), from the European Research Council (ERC-2014-Stg 638134 to D.P), and from the Junta de Andalucía ("Proyecto QUAL21 012 IHSM, Consejería de Universidad, Investigación e Innovación, Junta de Andalucía". UMA20-FEDERJA-093 and Postdoctoral program, POSTDOC_21_00893 to C.M.-P.).

Authors' contributions

R.T. carried out TE identification and classification. **I.T.** conducted MITEs and IRs bioinformatic analysis. **S.P.C.** performed TE polymorphism identification and GWAS analysis. **J.L.G**. and **C.Z** conducted all the validations of the MITE-triggered effects over *FvAAT1*. **C.M.P.** collected the samples and performed sRNA-sequencing and RNA-seq experiments. **T.H.**, and **T.T.** provided genomic data of *F. vesca* accessions (bam files). **I.T.**, **I.T., C.M.P.**, and **P.A.M.** conceived the study. **P.A.M.** secured project funding. **I.T.** and **P.A.M.** supervised the work and wrote the manuscript with the help of all authors.

Acknowledgements

We would like to acknowledge the Sequencing Unit from the University of Malaga for the support they have provided for the small RNA-seq analysis. We thankfully acknowledge the computer resources (Picasso Supercomputer), technical expertise and assistance provided by the SCBI (Supercomputing and Bioinformatics) center of the University of Malaga. This work was supported by grants from the Spanish Ministry of Science and Innovation (PID2022-137037NB-I00 and CNS2023-145312) to P.A.M.

REFERENCES

1. Matsushima W, Planet E, Trono D: **Ancestral genome reconstruction enhances transposable element annotation by identifying degenerate integrants.** *Cell Genom* 2024, **4:**100497.
2. International Wheat Genome Sequencing C: **Shifting the limits in wheat research and breeding using a fully annotated reference genome.** *Science* 2018, **361**.
3. Hirsch CD, Springer NM: **Transposable element influences on gene expression in plants.** *Biochim Biophys Acta Gene Regul Mech* 2017, **1860:**157-165.
4. Hosaka A, Kakutani T: **Transposable elements, genome evolution and transgenerational epigenetic variation.** *Curr Opin Genet Dev* 2018, **49:**43-48.
5. Judd J, Feschotte C: **Transposons take remote control.** *Elife* 2018, **7**.
6. Liu P, Cuerda-Gil D, Shahid S, Slotkin RK: **The Epigenetic Control of the Transposable Element Life Cycle in Plant Genomes and Beyond.** *Annu Rev Genet* 2022, **56:**63-87.
7. Tossolini I, Mencia R, Arce AL, Manavella PA: **The genome awakens: transposon-mediated gene regulation.** *Trends in Plant Science* 2025.
8. Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, Flavell A, Leroy P, Morgante M, Panaud O, et al: **A unified classification system for eukaryotic transposable elements.** *Nat Rev Genet* 2007, **8:**973-982.
9. Kumar A, Bennetzen JL: **Plant retrotransposons.** *Annu Rev Genet* 1999, **33:**479-532.
10. Ulrich D, Komes D, Olbricht K, Hoberg E: **Diversity of aroma patterns in wild and cultivated Fragaria accessions.** *Genetic Resources and Crop Evolution* 2007, **54:**1185-1196.
11. Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, Imbeault M, Izsvak Z, Levin HL, Macfarlan TS, et al: **Ten things you should know about transposable elements.** *Genome Biol* 2018, **19:**199.
12. Hu J, Zheng Y, Shang X: **MiteFinderII: a novel tool to identify miniature inverted-repeat transposable elements hidden in eukaryotic genomes.** *BMC Med Genomics* 2018, **11:**101.
13. Nouroz F, Noreen S, Heslop-Harrison JS: **Evolutionary genomics of miniature inverted-repeat transposable elements (MITEs) in Brassica.** *Mol Genet Genomics* 2015, **290:**2297-2312.
14. Fattash I, Rooke R, Wong A, Hui C, Luu T, Bhardwaj P, Yang G: **Miniature inverted-repeat transposable elements: discovery, distribution, and activity.** *Genome* 2013, **56:**475-486.
15. Kornienko AE, Nizhynska V, Molla Morales A, Pisupati R, Nordborg M: **Population-level annotation of lncRNAs in Arabidopsis reveals extensive expression variation associated with transposable element-like silencing.** *Plant Cell* 2023, **36:**85-111.
16. Cuerda-Gil D, Slotkin RK: **Non-canonical RNA-directed DNA methylation.** *Nat Plants* 2016, **2:**16163.
17. Vaucheret H, Voinnet O: **The plant siRNA landscape.** *Plant Cell* 2024, **36:**246-275.
18. Yang DL, Huang K, Deng D, Zeng Y, Wang Z, Zhang Y: **DNA-dependent RNA polymerases in plants.** *Plant Cell* 2023, **35:**3641-3661.
19. Bell EA, Butler CL, Oliveira C, Marburger S, Yant L, Taylor MI: **Transposable element annotation in non-model species: The benefits of species-specific repeat libraries using semi-automated EDTA and DeepTE de novo pipelines.** *Mol Ecol Resour* 2022, **22:**823-833.
20. Loreto ELS, Melo ES, Wallau GL, Gomes T: **The good, the bad and the ugly of transposable elements annotation tools.** *Genet Mol Biol* 2024, **46:**e20230138.
21. Guo R, Li YR, He S, Ou-Yang L, Sun Y, Zhu Z: **RepLong: de novo repeat identification using long read sequencing data.** *Bioinformatics* 2018, **34:**1099-1107.
22. Shi J, Liang C: **Generic Repeat Finder: A High-Sensitivity Tool for Genome-Wide De Novo Repeat Detection.** *Plant Physiol* 2019, **180:**1803-1815.

23. Storer JM, Hubley R, Rosen J, Smit AFA: **Methodologies for the De novo Discovery of**
Transposable Element Families. *Genes (Basel)* 2022, **13**.
24. Mao H, Wang H: **SINE_scan: an efficient tool to discover short interspersed nuclear**
elements (SINEs) in large-scale genomic datasets. *Bioinformatics* 2017, **33:**743-745.
25. Ou S, Jiang N: **LTR_retriever: A Highly Accurate and Sensitive Program for**
Identification of Long Terminal Repeat Retrotransposons. *Plant Physiol* 2018,
176:1410-1422.
26. Huda A, Jordan IK: **Analysis of transposable element sequences using CENSOR and**
RepeatMasker. *Methods Mol Biol* 2009, **537:**323-336.
27. Kennedy RC, Unger MF, Christley S, Collins FH, Madey GR: **An automated homology-**
based approach for identifying transposable elements. *BMC Bioinformatics* 2011,
12:130.
28. Baril T, Galbraith J, Hayward A: **Earl Grey: A Fully Automated User-Friendly**
Transposable Element Annotation and Analysis Pipeline. *Mol Biol Evol* 2024, **41**.
29. Riehl K, Riccio C, Miska EA, Hemberg M: **TransposonUltimate: software for**
transposon classification, annotation and detection. *Nucleic Acids Res* 2022, **50:**e64.
30. Castanera R, Morales-Díaz N, Gupta S, Purugganan M, Casacuberta JM: **Transposons**
are important contributors to gene expression variability under selection in rice
populations. *eLife* 2023, **12:**RP86324.
31. Rohilla M, Mazumder A, Saha D, Pal T, Begam S, Mondal TK: **Genome-wide**
identification and development of miniature inverted-repeat transposable elements
and intron length polymorphic markers in tea plant (Camellia sinensis). *Sci Rep*
2022, **12:**16233.
32. Akakpo R, Carpentier MC, Ie Hsing Y, Panaud O: **The impact of transposable elements**
on the structure, evolution and function of the rice genome. *New Phytol* 2020,
226:44-49.
33. Liang Y, Liu HJ, Yan J, Tian F: **Natural Variation in Crops: Realized Understanding,**
Continuing Promise. *Annu Rev Plant Biol* 2021, **72:**357-385.
34. Yan H, Haak DC, Li S, Huang L, Bombarely A: **Exploring transposable element-based**
markers to identify allelic variations underlying agronomic traits in rice. *Plant*
Commun 2022, **3:**100270.
35. Yin Z, Yang Q, Shen D, Liu J, Huang W, Dou D: **Online data resource for exploring**
transposon insertion polymorphisms in public soybean germplasm accessions.
Plant Physiol 2023, **193:**1036-1044.
36. Cai X, Lin R, Liang J, King GJ, Wu J, Wang X: **Transposable element insertion: a**
hidden major source of domesticated phenotypic variation in Brassica rapa. *Plant*
Biotechnol J 2022, **20:**1298-1310.
37. Dominguez M, Dugas E, Benchouaia M, Leduque B, Jimenez-Gomez JM, Colot V,
Quadrana L: **The impact of transposable elements on tomato diversity.** *Nat Commun*
2020, **11:**4058.
38. Vourlaki IT, Castanera R, Ramos-Onsins SE, Casacuberta JM, Perez-Enciso M:
Transposable element polymorphisms improve prediction of complex agronomic
traits in rice. *Theor Appl Genet* 2022, **135:**3211-3222.
39. Edger PP, Poorten TJ, VanBuren R, Hardigan MA, Colle M, McKain MR, Smith RD, Teresi
SJ, Nelson ADL, Wai CM, et al: **Origin and evolution of the octoploid strawberry**
genome. *Nat Genet* 2019, **51:**541-547.
40. Lyu K, Xiao J, Lyu S, Liu R: **Comparative Analysis of Transposable Elements in**
Strawberry Genomes of Different Ploidy Levels. *Int J Mol Sci* 2023, **24**.
41. Shulaev V, Sargent DJ, Crowhurst RN, Mockler TC, Folkerts O, Delcher AL, Jaiswal P,
Mockaitis K, Liston A, Mane SP, et al: **The genome of woodland strawberry (Fragaria**
vesca). *Nat Genet* 2011, **43:**109-116.

42. Yu Z, Li J, Wang H, Ping B, Li X, Liu Z, Guo B, Yu Q, Zou Y, Sun Y, et al: **Transposable**
elements in Rosaceae: insights into genome evolution, expression dynamics, and
syntenic gene regulation. *Hortic Res* 2024, **11:**uhae118.
43. Ulrich D, Olbricht K: **Diversity of metabolite patterns and sensory characters in wild**
and cultivated strawberries 1. *Journal of Berry Research* 2014, **4:**11-17.
44. Toivainen T, Salonen JS, Kirshner J, Lembinen S, De Kort H, Lyyski A, Edger PP,
Hilmarsson HS, Hallsson JH, Sargent DJ, et al: **The Late Quaternary climate impact on**
the genome of the woodland strawberry (Fragaria vesca), a perennial herb. *Commun*
Biol 2026.
45. Aharoni A, Keizer LC, Bouwmeester HJ, Sun Z, Alvarez-Huerta M, Verhoeven HA, Blaas
J, van Houwelingen AM, De Vos RC, van der Voet H, et al: **Identification of the SAAT**
gene involved in strawberry flavor biogenesis by use of DNA microarrays. *Plant Cell*
2000, **12:**647-662.
46. Goerner-Potvin P, Bourque G: **Computational tools to unmask transposable**
elements. *Nat Rev Genet* 2018, **19:**688-704.
47. Mhiri C, Borges F, Grandbastien MA: **Specificities and Dynamics of Transposable**
Elements in Land Plants. *Biology (Basel)* 2022, **11**.
48. Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, Lugo CSB, Elliott TA, Ware D,
Peterson T, et al: **Benchmarking transposable element annotation methods for**
creation of a streamlined, comprehensive pipeline. *Genome Biol* 2019, **20:**275.
49. Yan H, Bombarely A, Li S: **DeepTE: a computational method for de novo**
classification of transposons with convolutional neural network. *Bioinformatics*
2020, **36:**4269-4275.
50. Qian J, Xue H, Ou S, Storer J, Fürtauer L, Wildermuth MC, Kusch S, Panstruga R:
TEtrimmer: a novel tool to automate the manual curation of transposable elements.
bioRxiv 2024**:**2024.2006.2027.600963.
51. Edger PP, VanBuren R, Colle M, Poorten TJ, Wai CM, Niederhuth CE, Alger EI, Ou S,
Acharya CB, Wang J, et al: **Single-molecule sequencing and optical mapping yields**
an improved genome of woodland strawberry (Fragaria vesca) with chromosome-
scale contiguity. *Gigascience* 2018, **7:**1-7.
52. Li Y, Pi M, Gao Q, Liu Z, Kang C: **Updated annotation of the wild strawberry Fragaria**
vesca V4 genome. *Hortic Res* 2019, **6:**61.
53. Beric A, Mabry ME, Harkess AE, Brose J, Schranz ME, Conant GC, Edger PP, Meyers
BC, Pires JC: **Comparative phylogenetics of repetitive elements in a diverse order**
of flowering plants (Brassicales). *G3 (Bethesda)* 2021, **11**.
54. Wang D, Zheng Z, Li Y, Hu H, Wang Z, Du X, Zhang S, Zhu M, Dong L, Ren G, Yang Y:
Which factors contribute most to genome size variation within angiosperms? *Ecol*
Evol 2021, **11:**2660-2668.
55. Buisine N, Quesneville H, Colot V: **Improved detection and annotation of transposable**
elements in sequenced genomes using multiple reference sequence sets.
Genomics 2008, **91:**467-475.
56. Noshay JM, Marand AP, Anderson SN, Zhou P, Mejia Guerra MK, Lu Z, O'Connor CH,
Crisp PA, Hirsch CN, Schmitz RJ, Springer NM: **Assessing the regulatory potential of**
transposable elements using chromatin accessibility profiles of maize
transposons. *Genetics* 2021, **217:**1-13.
57. Zavallo D, Crescente JM, Gantuz M, Leone M, Vanzetti LS, Masuelli RW, Asurmendi S:
Genomic re-assessment of the transposable element landscape of the potato
genome. *Plant Cell Rep* 2020, **39:**1161-1174.
58. Zhou Y, Xiong J, Shu Z, Dong C, Gu T, Sun P, He S, Jiang M, Xia Z, Xue J, et al: **The**
telomere-to-telomere genome of Fragaria vesca reveals the genomic evolution of
Fragaria and the origin of cultivated octoploid strawberry. *Hortic Res* 2023,
10:uhad027.

59. Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic**
features. *Bioinformatics* 2010, **26:**841-842.
60. Jouffroy O, Saha S, Mueller L, Quesneville H, Maumus F: **Comprehensive repeatome**
annotation reveals strong potential impact of repetitive elements on tomato
ripening. *BMC Genomics* 2016, **17:**624.
61. Yang L, Bennetzen JL: **Structure-based discovery and description of plant and**
animal Helitrons. *Proc Natl Acad Sci U S A* 2009, **106:**12832-12837.
62. Yang L, Bennetzen JL: **Distribution, diversity, evolution, and survival of Helitrons in**
the maize genome. *Proc Natl Acad Sci U S A* 2009, **106:**19922-19927.
63. Guo C, Spinelli M, Ye C, Li QQ, Liang C: **Genome-Wide Comparative Analysis of**
Miniature Inverted Repeat Transposable Elements in 19 Arabidopsis thaliana
Ecotype Accessions. *Sci Rep* 2017, **7:**2634.
64. Arce AL, Mencia R, Cambiagno DA, Lang PL, Liu C, Burbano HA, Weigel D, Manavella
PA: **Polymorphic inverted repeats near coding genes impact chromatin topology**
and phenotypic traits in Arabidopsis thaliana. *Cell Rep* 2023, **42:**112029.
65. Mencia R, Arce AL, Houriet C, Xian W, Contreras A, Shirsekar G, Weigel D, Manavella
PA: **Transposon-triggered epigenetic chromatin dynamics modulate EFR-related**
pathogen response. *Nat Struct Mol Biol* 2025, **32:**199-211.
66. Gagliardi D, Cambiagno DA, Arce AL, Tomassi AH, Giacomelli JI, Ariel FD, Manavella
PA: **Dynamic regulation of chromatin topology and transcription by inverted repeat-**
derived small RNAs in sunflower. *Proc Natl Acad Sci U S A* 2019, **116:**17578-17583.
67. Gagliardi D, Manavella PA: **Short-range regulatory chromatin loops in plants.** *New*
Phytol 2020, **228:**466-471.
68. Rice P, Longden I, Bleasby A: **EMBOSS: the European Molecular Biology Open**
Software Suite. *Trends Genet* 2000, **16:**276-277.
69. Li R, Yao J, Cai S, Fu Y, Lai C, Zhu X, Cui L, Li Y: **Genome-wide characterization and**
evolution analysis of miniature inverted-repeat transposable elements in Barley
(Hordeum vulgare). *Front Plant Sci* 2024, **15:**1474846.
70. Lu C, Chen J, Zhang Y, Hu Q, Su W, Kuang H: **Miniature inverted-repeat transposable**
elements (MITEs) have been accumulated through amplification bursts and play
important roles in gene expression and species diversity in Oryza sativa. *Mol Biol*
Evol 2012, **29:**1005-1017.
71. Sun Q, He Z, Wei R, Ye J, Chai L, Cheng Y, Xu Q, Deng X: **Red peel regulator 1 links**
ethylene response factor 25 and beta-citraurin biosynthetic genes to regulate
ethylene-induced peel reddening in citrus. *Plant Cell* 2024, **37**.
72. Yang X, Liu Q, Wang MM, Wang XY, Han MQ, Liu FP, Lu TF, Liu J, Wang YZ: **A single**
dominant GLOBOSA allele accounts for repeated origins of hose-in-hose flowers
in Sinningia (Gesneriaceae). *Plant Cell* 2024, **37**.
73. Defilippi BG, Kader AA, Dandekar AM: **Apple aroma: alcohol acyltransferase, a rate**
limiting step for ester biosynthesis, is regulated by ethylene. *Plant Science* 2005,
168:1199-1210.
74. Zhang Z, Yu S, Zhang Z, Zhang J, Li H: **Comparative Characterization of Fruit Volatiles**
and Volatile-Related Genes Expression of 'Benihoppe' Strawberry and Its
Somaclonal Mutant. *Plants (Basel)* 2023, **12**.
75. Li Y-p, Liu T-j, Luo H-f, Liu S-c: **The transcriptional landscape of cultivated strawberry**
(Fragaria×ananassa) and its diploid ancestor (Fragaria vesca) during fruit
development. *Journal of Integrative Agriculture* 2021, **20:**1540-1553.
76. Nieuwenhuizen NJ, Maddumage R, Tsang GK, Fraser LG, Cooney JM, De Silva HN,
Green S, Richardson KA, Atkinson RG: **Mapping, complementation, and targets of the**
cysteine protease actinidin in kiwifruit. *Plant Physiol* 2012, **158:**376-388.
77. Wang W, Wang Y, Chen T, Qin G, Tian S: **Current insights into posttranscriptional**
regulation of fleshy fruit ripening. *Plant Physiol* 2023, **192:**1785-1798.

78. Zhai Y, Cui Y, Song M, Vainstein A, Chen S, Ma H: **Papain-Like Cysteine Protease Gene Family in Fig (Ficus carica L.): Genome-Wide Analysis and Expression Patterns.** *Front Plant Sci* 2021, **12:**681801.

79. Sanchez-Sevilla JF, Cruz-Rus E, Valpuesta V, Botella MA, Amaya I: **Deciphering gamma-decalactone biosynthesis in strawberry fruit using a combination of genetic mapping, RNA-Seq and eQTL analyses.** *BMC Genomics* 2014, **15:**218.

80. Seymour GB, Ryder CD, Cevik V, Hammond JP, Popovich A, King GJ, Vrebalov J, Giovannoni JJ, Manning K: **A SEPALLATA gene is involved in the development and ripening of strawberry (Fragaria x ananassa Duch.) fruit, a non-climacteric tissue.** *J Exp Bot* 2011, **62:**1179-1188.

81. Vallarino JG, Merchante C, Sanchez-Sevilla JF, de Luis Balaguer MA, Pott DM, Ariza MT, Casanal A, Pose D, Vioque A, Amaya I, et al: **Characterizing the involvement of FaMADS9 in the regulation of strawberry fruit receptacle development.** *Plant Biotechnol J* 2020, **18:**929-943.

82. Martinez-Rivas FJ, Blanco-Portales R, Molina-Hidalgo FJ, Caballero JL, Perez de Souza L, Alseekh S, Fernie AR, Munoz-Blanco J, Rodriguez-Franco A: **Azacytidine arrests ripening in cultivated strawberry (Fragaria x ananassa) by repressing key genes and altering hormone contents.** *BMC Plant Biol* 2022, **22:**278.

83. Liu L, Xiao W, Ji M-l, Yang C, Li L, Gao D-s, Fu X-l: **Effects of molybdenum on nutrition, quality, and flavour compounds of strawberry (Fragaria×ananassa Duch. cv. Akihime) fruit.** *Journal of Integrative Agriculture* 2017, **16:**1502-1512.

84. Cosgrove DJ: **Growth of the plant cell wall.** *Nat Rev Mol Cell Biol* 2005, **6:**850-861.

85. Del Olmo I, Romero I, Alvarez MD, Tarradas R, Sanchez-Ballesta MT, Escribano MI, Merodio C: **Transcriptomic analysis of CO(2)-treated strawberries (Fragaria vesca) with enhanced resistance to softening and oxidative stress at consumption.** *Front Plant Sci* 2022, **13:**983976.

86. Ariel FD, Manavella PA: **When junk DNA turns functional: transposon-derived non-coding RNAs in plants.** *J Exp Bot* 2021, **72:**4132-4143.

87. Alonge M, Wang X, Benoit M, Soyk S, Pereira L, Zhang L, Suresh H, Ramakrishnan S, Maumus F, Ciren D, et al: **Major Impacts of Widespread Structural Variation on Gene Expression and Crop Improvement in Tomato.** *Cell* 2020, **182:**145-161 e123.

88. Quadrana L, Etcheverry M, Gilly A, Caillieux E, Madoui MA, Guy J, Bortolini Silveira A, Engelen S, Baillet V, Wincker P, et al: **Transposition favors the generation of large effect mutations that may facilitate rapid adaption.** *Nat Commun* 2019, **10:**3421.

89. Saxena RK, Edwards D, Varshney RK: **Structural variations in plant genomes.** *Brief Funct Genomics* 2014, **13:**296-307.

90. Stuart T, Eichten SR, Cahn J, Karpievitch YV, Borevitz JO, Lister R: **Population scale mapping of transposable element diversity reveals links to gene regulation and epigenomic variation.** *Elife* 2016, **5**.

91. Tossolini I, Mencia R, Arce AL, Manavella PA: **The genome awakens: transposon-mediated gene regulation.** *Trends Plant Sci* 2025, **30:**857-871.

92. Contreras-Garrido A, Galanti D, Movilli A, Becker C, Bossdorf O, Drost HG, Weigel D: **Transposon dynamics in the emerging oilseed crop Thlaspi arvense.** *PLoS Genet* 2024, **20:**e1011141.

93. Quadrana L, Bortolini Silveira A, Mayhew GF, LeBlanc C, Martienssen RA, Jeddeloh JA, Colot V: **The Arabidopsis thaliana mobilome and its impact at the species level.** *Elife* 2016, **5**.

94. Catlin NS, Josephs EB: **The important contribution of transposable elements to phenotypic variation and evolution.** *Curr Opin Plant Biol* 2022, **65:**102140.

95. Urrutia M, Meco V, Rambla JL, Martin-Pizarro C, Pillet J, Andres J, Sanchez-Sevilla JF, Granell A, Hytonen T, Pose D: **Diversity of the volatilome and the fruit size and shape in European woodland strawberry (Fragaria vesca).** *Plant J* 2023, **116:**1201-1217.

96. Baduel P, Leduque B, Ignace A, Gy I, Gil J, Jr., Loudet O, Colot V, Quadrana L: **Genetic**
and environmental modulation of transposition shapes the evolutionary potential
of Arabidopsis thaliana. *Genome Biol* 2021, **22:**138.
97. Shi J, Cao C, Xu J, Zhou C: **Research Advances on Biosynthesis, Regulation, and**
Biological Activities of Apocarotenoid Aroma in Horticultural Plants. *Journal of*
Chemistry 2020, **2020:**2526956.
98. Riikonen J, Ruhanen H, Uimari A, Poteri M, Toljamo A, Kokko H, Blande JD, Kumpula R,
Kivimäenpää M: **Impact of pre-harvest UVC treatment on powdery mildew infection**
and strawberry quality in tunnel production in Nordic conditions. *Scientia*
Horticulturae 2024, **338:**113706.
99. Achimon F, Peschiutta ML, Brito VD, Ulla SB, Pizzolitto RP: **Sulcatone as a Plant-**
Derived Volatile Organic Compound for the Control of the Maize Weevil and Its
Associated Phytopathogenic Fungi in Stored Maize. *Plants (Basel)* 2024, **13**.
100. Antonious GF, Dahlman DL, Hawkins LM: **Insecticidal and acaricidal performance**
of methyl ketones in wild tomato leaves. *Bull Environ Contam Toxicol* 2003, **71:**400-
407.
101. Shigeto J, Tsutsumi Y: **Diverse functions and reactions of class III peroxidases.**
New Phytol 2016, **209:**1395-1402.
102. Matsui K: **How and Why Plants Came to Smell Green: The Origins, Biosynthesis,**
and Roles of Green Leaf Volatiles. *J Exp Bot* 2025.
103. Youn B, Kim S-J, Moinuddin SGA, Lee C, Bedgar DL, Harper AR, Davin LB, Lewis
NG, Kang C: **Mechanistic and Structural Studies of Apoform, Binary, and Ternary**
Complexes of the Arabidopsis Alkenal Double Bond Reductase At5g16970*. *Journal*
of Biological Chemistry 2006, **281:**40076-40088.
104. Zhang Y, Teoh KH, Reed DW, Covello PS: **Molecular cloning and characterization**
of Dbr1, a 2-alkenal reductase from Artemisia annuaThe nucleotide sequence
reported in this article has been deposited in the GenBank database under
accession No. FJ750460.This paper is one of a selection of papers published in a
Special Issue from the National Research Council of Canada – Plant Biotechnology
Institute. *Botany* 2009, **87:**643-649.
105. Jiménez-Muñoz R, Urrutia M, Meco V, Rambla JL, Toivainen T, Pérez-Hedo M,
Sánchez-Sevilla JF, Urbaneja A, Salas JJ, Hytönen T, et al: **Uncovering the genetic**
basis of fruit volatiles in Fragaria vesca through GWAS reveals FvJMT2
as a methyl benzoate biosynthesis gene with insect-repellent function. *bioRxiv*
2025**:**2025.2011.2018.689018.
106. Urrutia M, Rambla JL, Alexiou KG, Granell A, Monfort A: **Genetic analysis of the wild**
strawberry (Fragaria vesca) volatile composition. *Plant Physiol Biochem* 2017,
121:99-117.
107. Holland CK, Watson AP, Chiang E: **Anthranilate at the interface of tryptophan and**
specialized metabolite biosynthesis. *Front Plant Sci* 2025, **16:**1625337.
108. Barbey CR, Hogshead MH, Harrison B, Schwartz AE, Verma S, Oh Y, Lee S, Folta
KM, Whitaker VM: **Genetic Analysis of Methyl Anthranilate, Mesifurane, Linalool, and**
Other Flavor Compounds in Cultivated Strawberry (Fragaria x ananassa). *Front*
Plant Sci 2021, **12:**615749.
109. Schwieterman ML, Colquhoun TA, Jaworski EA, Bartoshuk LM, Gilbert JL, Tieman
DM, Odabasi AZ, Moskowitz HR, Folta KM, Klee HJ, et al: **Strawberry flavor: diverse**
chemical compositions, a seasonal influence, and effects on sensory perception.
PLoS One 2014, **9:**e88446.
110. Bontpart T, Cheynier V, Ageorges A, Terrier N: **BAHD or SCPL acyltransferase?**
What a dilemma for acylation in the world of plant phenolic compounds. *New Phytol*
2015, **208:**695-707.
111. Mugford ST, Milkowski C: **Serine carboxypeptidase-like acyltransferases from**
plants. *Methods Enzymol* 2012, **516:**279-297.

112. Beekwilder J, Alvarez-Huerta M, Neef E, Verstappen FW, Bouwmeester HJ, Aharoni A: **Functional characterization of enzymes forming volatile esters from strawberry and banana.** *Plant Physiol* 2004, **135:**1865-1878.

113. Gonzalez M, Gaete-Eastman C, Valdenegro M, Figueroa CR, Fuentes L, Herrera R, Moya-Leon MA: **Aroma development during ripening of Fragaria chiloensis fruit and participation of an alcohol acyltransferase (FcAAT1) gene.** *J Agric Food Chem* 2009, **57:**9123-9132.

114. Gill RA, Scossa F, King GJ, Golicz AA, Tong C, Snowdon RJ, Fernie AR, Liu S: **On the Role of Transposable Elements in the Regulation of Gene Expression and Subgenomic Interactions in Crop Genomes.** *Critical Reviews in Plant Sciences* 2021, **40:**157-189.

115. Studer A, Zhao Q, Ross-Ibarra J, Doebley J: **Identification of a functional transposon insertion in the maize domestication gene tb1.** *Nat Genet* 2011, **43:**1160-1163.

116. Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U, Putnam N, Rokhsar DS: **Phytozome: a comparative platform for green plant genomics.** *Nucleic Acids Res* 2012, **40:**D1178-1186.

117. Neph S, Kuehn MS, Reynolds AP, Haugen E, Thurman RE, Johnson AK, Rynes E, Maurano MT, Vierstra J, Thomas S, et al: **BEDOPS: high-performance genomic feature operations.** *Bioinformatics* 2012, **28:**1919-1920.

118. Qian J, Xue H, Ou S, Mann L, Storer J, Furtauer L, Heitkam T, Wildermuth MC, Kusch S, Panstruga R: **TEtrimmer: a tool to automate the manual curation of transposable elements.** *Nat Commun* 2025, **16:**8429.

119. Fu L, Niu B, Zhu Z, Wu S, Li W: **CD-HIT: accelerated for clustering the next-generation sequencing data.** *Bioinformatics* 2012, **28:**3150-3152.

120. Li X, Martin-Pizarro C, Zhou L, Hou B, Wang Y, Shen Y, Li B, Pose D, Qin G: **Deciphering the regulatory network of the NAC transcription factor FvRIF, a key regulator of strawberry (Fragaria vesca) fruit ripening.** *Plant Cell* 2023, **35:**4020-4045.

121. Gambino G, Perrone I, Gribaudo I: **A Rapid and effective method for RNA extraction from different tissues of grapevine and other woody plants.** *Phytochem Anal* 2008, **19:**520-525.

122. Ewels P, Magnusson M, Lundin S, Kaller M: **MultiQC: summarize analysis results for multiple tools and samples in a single report.** *Bioinformatics* 2016, **32:**3047-3048.

123. Axtell MJ: **ShortStack: comprehensive annotation and quantification of small RNA genes.** *RNA* 2013, **19:**740-751.

124. Love MI, Huber W, Anders S: **Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.** *Genome Biol* 2014, **15:**550.

125. Lu P, Yu S, Zhu N, Chen YR, Zhou B, Pan Y, Tzeng D, Fabi JP, Argyris J, Garcia-Mas J, et al: **Genome encode analyses reveal the basis of convergent evolution of fleshy fruit ripening.** *Nat Plants* 2018, **4:**784-791.

126. Bolger AM, Lohse M, Usadel B: **Trimmomatic: a flexible trimmer for Illumina sequence data.** *Bioinformatics* 2014, **30:**2114-2120.

127. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR: **STAR: ultrafast universal RNA-seq aligner.** *Bioinformatics* 2013, **29:**15-21.

128. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project Data Processing S: **The Sequence Alignment/Map format and SAMtools.** *Bioinformatics* 2009, **25:**2078-2079.

129. Liao Y, Smyth GK, Shi W: **featureCounts: an efficient general purpose program for assigning sequence reads to genomic features.** *Bioinformatics* 2014, **30:**923-930.

130. Koster J, Rahmann S: **Snakemake--a scalable bioinformatics workflow engine.** *Bioinformatics* 2012, **28:**2520-2522.

131.   Nunn A, Can SN, Otto C, Fasold M, Diez Rodriguez B, Fernandez-Pozo N, Rensing
SA, Stadler PF, Langenberger D: **EpiDiverse Toolkit: a pipeline suite for the analysis**
of bisulfite sequencing data in ecological plant epigenetics. *NAR Genom Bioinform*
2021, **3:**lqab106.
132.   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane
T, McCarthy SA, Davies RM, Li H: **Twelve years of SAMtools and BCFtools.**
Gigascience 2021, **10**.
133.   Langmead B, Trapnell C, Pop M, Salzberg SL: **Ultrafast and memory-efficient**
alignment of short DNA sequences to the human genome. *Genome Biol* 2009,
10:R25.
134.   Faust GG, Hall IM: **YAHA: fast and flexible long-read alignment with optimal**
breakpoint detection. *Bioinformatics* 2012, **28:**2417-2424.
135.   Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, Morgan MT,
Carey VJ: **Software for Computing and Annotating Genomic Ranges.** *PLOS*
Computational Biology 2013, **9:**e1003118.
136.   Wang J, Zhang Z: **GAPIT Version 3: Boosting Power and Accuracy for Genomic**
Association and Prediction. *Genomics Proteomics Bioinformatics* 2021, **19:**629-640.
137.   Segura V, Vilhjalmsson BJ, Platt A, Korte A, Seren U, Long Q, Nordborg M: **An**
efficient multi-locus mixed-model approach for genome-wide association studies
in structured populations. *Nat Genet* 2012, **44:**825-830.
138.   VanRaden PM: **Efficient methods to compute genomic predictions.** *J Dairy Sci*
2008, **91:**4414-4423.
139.   Sigg CD, Buhmann JM: **Expectation-maximization for sparse and non-negative**
PCA. In *Proceedings of the 25th international conference on Machine learning*. pp. 960–
967. Helsinki, Finland: Association for Computing Machinery; 2008:960–967.
140.   Lawrence M, Gentleman R, Carey V: **rtracklayer: an R package for interfacing with**
genome browsers. *Bioinformatics* 2009, **25:**1841-1842.
1507
1508

FIGURE LEGENDS

1511 **Figure 1. Transposable Element (TE) diversity in *Fragaria vesca*. (A)** Total coverage of

1512 the *F. vesca* genome assembly by the main TE classes, shown as percentages relative to the

1513 entire genome. Representation of TE classification scheme generated in this study is

1514 displayed. **(B)** Repetitive sequence content as a proportion of total genome size in several

1515 plant species, listed in ascending order: *Arabidopsis thaliana*, woodland strawberry (*Fragaria*

1516 *vesca*), rice (*Oryza sativa*), tomato (*Solanum lycopersicum*), and maize (*Zea mays*). **(C)**

1517 Length distribution of TEs (including fragmented TEs) in the *F. vesca* v4.0.a2 annotation. The

1518 classifications are listed in descending order for EDTA, DeepTE, and the curated annotation.

1519 **(D)** Length distribution of intact TEs in the curated *F. vesca* v4 annotation.

1520

1521

Figure 2. Distribution of TEs in the *F. vesca* genome, represented as density plots within 500 kb windows. (A) Class I TEs: LTR (Track b in blue), non-LTR (Track c in green). **(B)** Class II TEs: DNA (Track "d" red), MITE (Track "e" in orange), Helitron (Track "f" in purple). The color scale indicates the number of elements found within each 500 kb window. The distribution of genes is shown in grayscale (Track "a") in both graphs. Centromeres are marked as a yellow dot in the brown outer line.

1528

Figure 3. Characterization of TE-Derived IRs and MITEs in *F. vesca*.

(A) Proportion of the identified IRs (left) and annotated MITEs (right) classified by TE superfamily in *F. vesca.* (B) Number of IRs (left) and MITEs (right) detected in the *F. vesca* v4.0.a2 reference genome, counting elements located within gene bodies ('intragenic') or within the upstream or downstream regions defined as 0–500 bp ('500'), 501–1000 bp ('1000'), and 1001–3000 bp ('3000') of annotated genes (the counts for these flanking regions also include any intragenic elements). Black numbers indicate the total counts of IRs and MITEs,

while white numbers show the non-cumulative counts. **(C)** Number of IRs and MITEs that are

entirely within a single gene feature ('exon', 'intron', '3′-UTR', or '5′-UTR') or that span more

than one gene feature as indicated. **(D)** Metagene profile of 24-nt siRNAs mapping to all IRs

and MITEs in leaf (green), and immature (gray) and ripe fruits (pink). Plots show MITEs/IRs

scaled from the start to the end plus 2,000 bp on each side. sRNA-seq replicates are plotted

together. **(E)** Metagene profile of CpG, CHG, and CHH DNA methylation of all genes, TEs,

MITEs, and IRs in leaf (green), immature (shades of gray), and ripe fruits (shades of pink).

Plots show Genes, TEs, MITEs, and IRs scaled from the start to the end plus 2,000 bp on

each side. Individual BS-seq replicates are plotted.

1545

1546

Figure 4. Regulatory role of MITEs and IRs in gene expression.

(A) Differential accumulation analysis of 24-nt siRNAs derived from IRs and MITEs across leaf

tissue, immature and ripe fruits, using thresholds of |log2-fold change| ≥ 1 and an adjusted p-

value (padj) ≤ 0.05. **(B)** Differential accumulation analysis of 24-nt siRNAs derived from IRs

(left) and MITEs (right) elements across leaf tissue, immature and ripe fruits, using thresholds

of log2-fold change (FC) ≥0.5 and adjusted p-value (padj) ≤0.05. The percentages of elements

located within 3,000 bp of either differentially expressed genes (DE) or non-DE genes are

shown.

1555

Figure 5. A *FvAAT1*-hosted MITE control the gene expression through epigenetically driven chromatin reorganization

(A) Volcano plots showing the log2FC of the differential expression analysis for IRs (left) and MITEs (right) between immature and ripe stages. Red points represent elements with an adjusted p-value ≤0.05. **(B)** 24-nt siRNAs mapping to IR_9194 and the two MITEs near FvH4_7g18550 and FvH4_7g18570 genes (highlighted by a gray box) in immature (gray) and ripe (pink) fruits. Each element is scaled from start to end plus 500 bp on each side, and all sRNA-seq replicates are plotted together. **(C)** Cytosine methylation in the CHH context for *IR_9194*, MITE Mutator, and MITE, including 500-bp flanking regions, in immature and ripe fruits. The average of individual BS-seq replicates is plotted for each condition. Average percentage of CHH methylation in each element and its 500-bp flanking regions. **(D)** *FvAAT1* expression in transiently transformed strawberry receptacles. RNA levels are expressed relative to the control samples (35S:GFP) plants. Values of n = 6 biologically independent samples. Data are presented as the mean values ± s.d. P values were calculated with a two-tailed unpaired t-test with Welch's correction. **(E)** Chromatin conformation capture (3C)

analysis showing the interaction frequency of MITE Mutator with the FvAAT1 locus in immature and ripe fruits as measured by 3C–qPCR. Values of n = 4 biologically independent samples are expressed relative to green immature stage. Data are presented as the mean values ± s.d. P values were calculated with a two-tailed unpaired t-test with Welch's correction.

Figure 6. Distribution of insertion and deletion polymorphisms in the *F. vesca* v4.0.a2 annotation.

(A) Proportion (%) of identified polymorphisms by TE superfamily. Rare TE polymorphisms are indicated by minor allele frequency (MAF) < 5% and frequent TE polymorphisms by MAF ≥ 5%. (B) Principal Component Analysis (PCA) based on TE insertion and deletion polymorphisms (MAF ≥ 5%). The geographic origin of the analyzed F. vesca accessions is indicated according to Urrutia et al [95]. **(C)** Distribution of polymorphic elements over genic features, including 5' UTR, Exons, Introns, and 3' UTR (UTR: unstranslated transcribed region; Multi-feature: polymorphisms overlpaping multiple annotations). Stacked bar plots represent the relative proportion of Insertions, Deletions, and specific TE superfamilies. Polymorphisms located within 1000 bp of a gene are shown in shades of blue, representing Upstream and Downstream regions. Intergenic polymorphisms located more than 1000 bp away from those regions are indicated in grey (<-1000 bp and >1000 bp). The 'Genomic fraction' bar represents the percentage of base pairs occupied by each feature in the reference genome.

Figure 7. TE polymorphisms linked to volatile compound variation in a collection of *F. vesca* accessions. (A) Manhattan plot of TEP-GWAS for 6-methyl-5-hepten-2-one in harvest season H17. The dotted gray line marks the significance threshold determined by the Bonferroni correction, and the significantly associated polymorphism is highlighted. **(B)** Observed and expected distribution of *p* values for TEP-GWAS for 6-methyl-5-hepten-2-one. **(C)** Violin plot showing the relative abundance of 6-methyl-5-hepten-2-one in accessions carrying the MITE insertion (+, coral) versus non-carriers (-, steel blue). Horizontal lines in (-) and (+) indicate the mean. **(D)** Manhattan plot of TEP-GWAS for (E)-2-hexenal in harvest season H17. The dotted gray line marks the significance threshold determined by the Bonferroni correction, and the significantly associated polymorphism is highlighted. **(E)** Observed and expected distribution of *p* values for TEP-GWAS for (E)-2-hexenal. **(F)** Violin plot showing the relative abundance of (E)-2-hexenal in accessions carrying the MITE insertion (+, coral) versus non-carriers (-, steel blue). Horizontal lines in (-) and (+) indicate the mean.

TABLES

Table 1. Number of unique TEs, number of copies, and total TE coverage of the *F. vesca* genome assembly v4.0, according to their classification.

Classification	N° of TEs	N° of Copies	Genome Coverage (%)
Class I			
LTR			
Copia	131	8,647	3.46
Gypsy	262	25,977	10.43
Unclassified	35	3,536	0.91
Total LTRs	428	38,160	14.80
non-LTR			
LINE (L1)	8	292	0.07
PLE	7	548	0.18
Total non-LTRs	15	840	0.25
Total Class I	443	39,000	15.05
Class II			
Subclass 1			
MITE			
CACTA	7	1,084	0.08
Harbinger	25	4,243	0.46
Mutator	56	7,495	0.77
TcMar	23	2,721	0.28
hAT	53	5,780	0.68
Unclassified	28	4,737	0.51
Total MITEs	192	26,060	2.78
DNA (non-MITE)			
CACTA	50	8,495	2.97
Harbinger	20	2,376	0.49
Mutator	166	13,946	1.95
TcMar	63	7,391	1.66
hAT	134	12,237	2.00
Unclassified	63	6,184	0.96
Total DNA	496	50,629	10.03
Subclass 2			
Helitron	13	2,291	0.62
Total Class II	701	78,981	13.43

Table 2.

1622   Polymorphisms detected by TEPID for each element annotation. MAF ≥ 5% refers to

1623   polymorphisms with a minor allele frequency above 5%. The percentage shown represents

1624   the proportion of each value relative to the total number of polymorphic elements detected for

1625   each element and each type (insertion/deletion).

1626

Polymorphisms		TEs Amount	%	MITEs Amount	%	IRs Amount	%
Insertions	Total	5974	100.0	1859	100.0	351	100.0
	MAF ≥ 5%	946	15.8	336	18.1	42	12.0
	MAF < 5%	5028	84.2	1523	81.9	309	88.0
Deletions	Total	7855	100.0	2507	100.0	983	100.0
	MAF ≥ 5%	2256	28.7	601	24.0	197	20.0
	MAF < 5%	5599	71.3	1906	76.0	786	80.0

SUPPLEMENTARY MATERIAL

Table S1. Size distribution of Transposable Elements by Class annotated in *Fragaria vesca*, including both fragmented and intact TEs.

Table S2. ShortStack analysis of sRNA-seq data from *Fragaria vesca* leaves, immature (white stage), and ripe (red stage) fruits using annotated loci files for MITEs and IRs.

Table S3. Differential expression analysis of 24-nt siRNA-producing MITEs and adjacent genes (< 3 kb) in *Fragaria vesca*. Genomic coordinates, functional annotation and expression changes between leaf tissue, and immature and ripe stages are shown. Gene IDs associated with key roles in fruit development and metabolism are indicated in bold.

Table S4. Differential expression analysis of 24-nt siRNA-producing IRs and adjacent genes (< 3 kb) in *Fragaria vesca*. Genomic coordinates, functional annotation and expression changes between leaf tissue, and immature and ripe stages are shown. Gene IDs associated with key roles in fruit development and metabolism are indicated in bold.

Table S5. Differential gene expression analysis of all *F. vesca* genes across conditions: leaf tissue vs. immature stage, ripe stage vs. leaf tissue, ripe vs. immature stages, and AZA-treated vs. control strawberries.

Table S6. Resequenced genomes of *Fragaria vesca* analyzed in the present study, including accession identifiers, country and origin coordinates, read counts, and average chromosome coverage (excluding unassembled contigs). Accessions in italics were excluded from analysis due to low coverage; accessions in bold were included in the GWAS study.

Table S7. Summary of significant genome-wide associations between Transposable Element Polymorphisms (TEPs) and agronomic and volatilome traits across two harvest seasons (H16 and H17) in *Fragaria vesca*. Associations described in the Results section are highlighted in bold.

Figure S1. Annotation and classification of transposable elements (TEs) in the *Fragaria vesca* v4.0.a2 reference genome. (A) Representation of the pipeline used in this work for transposon annotation. Software employed are highlighted in blue boxes, while input and output files are shown in pink boxes, with their formats in parentheses. **(B)** Distribution of TE

superfamilies as annotated by EDTA (left), by DeepTE (center), and by TEtrimmer (right) for the *F. vesca* v4.0 genome.

Figure S2. The average length of intact TEs and all annotated TEs in *F. vesca* genome (including fragmented TEs).

Figure S3. Distribution of intact TE superfamilies in the *Fragaria vesca* v4.0.a2 annotation based on their distance from a gene. Categories include: within a gene ('intragenic'), within 1 kb ('1000'), between 1 and 3 kb ('3000'), and more than 3 kb ('>3000'). Left: Total counts by superfamily. Right: Proportion by superfamily.

Figure S4. (A) Genome distribution of IRs and MITEs. **(B)** Proportion of IRs classified as 'Other', excluding MITE-type elements. Values are shown as percentages relative to the total number of IRs in the 'Other' category (16.1%).

Figure S5. (A) Metagene profile of 24-nt siRNAs mapping to MITEs and IRs located at different ranges of distance to the closest protein-coding gene, in leaf (green), and immature (gray) and ripe fruits (pink). Plots show MITEs/IRs scaled from the start to the end plus 2,000 bp to each side. sRNA-seq replicates are plotted together. **(B)** Metagene profile of CpG, CHG, and CHH DNA methylation at MITEs/IRs located at different ranges of distance to the closest protein-coding gene, in leaf (green), and immature (gray) and ripe fruits (pink). Plots show MITEs/IRs scaled from the start to the end plus 2,000 bp to each side. Individual BS-seq replicates are plotted.

Figure S6. Genomic region of the *F. vesca* v4 assembly containing the MITE Mutator locus near FvH4_7g18550 and FvH4_7g18570 genes. IRs and MITEs identified in this region are also displayed.

Figure S7. Conservation of the MITE Mutator from *Fragaria vesca* in the orthologous genomic regions of commercial strawberry species *Fragaria chiloensis* and *Fragaria × ananassa*. (A) Local alignment of *F. vesca* MITE Mutator with the MITE located near the FvH4_7g18570 ortholog in *F. chiloensis* (left) and *F. × ananassa* (right). **(B)** Minimum free energy (MFE) RNA secondary structure predicted by RNAfold server of each MITE. The color scale shows base-pairing probability 0 (purple) to 1 (red).

Figure S8. Region of the *F. vesca* v4.0.a2 reference genome containing the FvH4_6g46420 **(A)**, FvH4_2g30230 **(B),** and FvH4_5g23180 **(C)** loci displaying the IR and MITE elements annotated in the region and expression and epigenetic profiles. (I) 24-nt siRNAs mapping to the genomic regions as determined by sRNA sequencing of leaf (green), immature (gray) and ripe (pink) tissues. Replicates are plotted together. (II) Expression of each gene in leaf (green), immature (gray) and ripe (pink) tissues as well as strawberries treated with 5-azacytidine (5-AZA) and its control, measured by RNA-seq. (III) Cytosine DNA methylation in CG, CHG, and CHH contexts in leaf (green), immature (gray) and ripe (pink) fruits. The average of individual BS-seq replicates is plotted for each condition.

Figure S9. (A) Representation of the pipeline used in this study. The main inputs are shown at the top, while the main outputs are placed at the bottom. File formats are indicated in parentheses. **(B)** Representation of the TEPID pipeline used for identifying transposable element insertion polymorphisms (TIPs) and absence polymorphisms (TAPs).

Figure S10. Enrichment analysis of TE insertion and deletion polymorphisms across genomic features in *Fragaria vesca* v4.0.a2. (A) Genome-wide enrichment of all polymorphisms, insertions, and deletions across annotated genomic features. **(B)** Enrichment profiles of TE-associated polymorphisms across major TE superfamilies. Values are shown as log2(observed/expected), where 0 indicates a random distribution relative to genomic fraction. Positive values (red) indicate enrichment, whereas negative values (blue) indicate depletion. Bubble size reflects the absolute magnitude of enrichment or depletion.

Figure S11. Principal Component Analysis (PCA) of the subset of *Fragaria vesca* accessions used for GWAS analysis. Each point represents a single accession, with colors indicating their respective country of origin.

Figure S12. Genomic region of the *F. vesca* v4.0.a2 reference genome harboring selected TEP polymorphisms on chromosome Fvb3 associated with variation in: (A) 6-methyl-5-hepten-2-one levels. (I) Coverage of mapped reads (light green) and split reads (dark green) in some accessions carrying the polimorphism (from top to bottom: ES14, ES-13, UK11, and IT15). (II) Coverage of mapped reads (light blue) and split reads (dark blue) in non-carriers accesions (from top to bottom: FIN10, FIN12, FIN6, and FIN51). **(B)** γ-decalactone levels. (I) Coverage of mapped reads (light green) and split reads (dark green) in some accessions carrying the polimorphism (from top to bottom: DK1, ES14, IT6, and IT15). (II)

1732 Coverage of mapped reads (light blue) and split reads (dark blue) in non-carriers accesions
1733 (from top to bottom: FR1, FIN25, FIN51, and NOR131-2).

1734

Figure S13. Addition TE polymorphisms linked to volatile compound variation in a
collection of *F. vesca* accessions. **(A)** Manhattan plot of TEP-GWAS for methyl 2-
aminobenzoate in harvest season H16. The dotted gray line marks the significance threshold
determined by the Bonferroni correction, and the significantly associated polymorphism is
highlighted. **(E)** Observed and expected distribution of p values for TEP-GWAS for methyl 2-
aminobenzoate. **(F)** Violin plot showing the relative abundance of methyl 2-aminobenzoate in
accessions carrying the MITE Mutator deletion (+, coral) versus non-carriers (-, steel blue).
Horizontal lines in (-) and (+) indicate the mean. **(D)** Manhattan plot of TEP-GWAS for γ-
decalactone across both harvest seasons (H16 and H17, represented by circles and triangles,
respectively). The dotted gray line marks the significance threshold determined by the
Bonferroni correction, and the significantly associated polymorphisms are highlighted. **(E)**
Observed and expected distribution of p values for TEP-GWAS for γ-decalactone. **(F)** Violin
plot showing the relative abundance of γ-decalactone in accessions carrying the MITE deletion
(+, coral) versus non-carriers (-, steel blue). Horizontal lines in (-) and (+) indicate the mean.

1749

1750