A Prelude to Conservation Genomics: First Chromosome-Level Genome Assembly of a Flying Squirrel (Pteromyini: *Pteromys volans*)

Authors:

Gerrit Wehrenberg[1] https://orcid.org/0000-0003-4828-3057,
Angelika Kiebler[1] https://orcid.org/0009-0001-2762-5057,
Carola Greve[2, 3] https://orcid.org/0000-0003-4993-1378,
Núria Beltrán-Sanz[4] https://orcid.org/0000-0001-8081-6234,
Alexander Ben Hamadou[2, 3] https://orcid.org/0009-0008-9826-902X,
René Meißner[5] https://orcid.org/0000-0003-2829-1607,
Sven Winter[5, 6, 7] https://orcid.org/0000-0002-1890-0977,
and Stefan Prost[1,8] https://orcid.org/0000-0002-6229-3596

Affiliations:

[1] Ecology and Genetics Research Unit, University of Oulu, P.O. Box 8000, 90014, Oulu, Finland

[2] Centre for Translational Biodiversity Genomics (LOEWE-TBG), Frankfurt am Main, Germany

[3] Senckenberg Forschungsinstitut und Naturmuseum, Senckenberganlage 25, 60325 Frankfurt am Main, Germany

[4] Centro di Ateneo Orto Botanico, University of Padua, Via Orto Botanico 15, 35123 Padua, Italy

[5] Research Institute of Wildlife Ecology, University of Veterinary Medicine, Vienna, Austria

[6] Faculty of Science and Technology, University of the Faroe Islands, Tórshavn, The Faroe Islands

[7] Senckenberg Biodiversity and Climate Research Centre, Frankfurt am Main, Germany

[8] Central Research Laboratories, Natural History Museum Vienna, Vienna, Austria

Correspondence to Gerrit Wehrenberg: gerrit.wehrenberg@oulu.fi

Keywords:

Pteromys volans, Siberian flying squirrel, Omni-C, PacBio HiFi, reference genome, Sciuridae, phylogeny

38 Abstract [English]

The Siberian flying squirrel (*Pteromys volans*) represents the only European
Pteromyini species. Thus, it is biogeographically unique due to its specialised
anatomy and biology as a volant rodent. As a result of habitat fragmentation and
destruction, Siberian flying squirrels experience severe and ongoing population
declines throughout most of their distribution. While considered *Least Concern*
throughout their immense Eurasian distribution, this species is red-listed as
Vulnerable and even *Critically Endangered* in parts of its range. More knowledge
about the population structure and overall biology is needed to improve conservation
efforts for this umbrella and flagship species of old-growth boreal forests. Here, we
present the first chromosome-level genome assembly of any Pteromyini, represented
by *P. volans* (Uoulu_pteVol_1.0). The final assembly has a total length of 2.85 Gbp in
19 chromosome-scale scaffolds with only minor differences in the chromosomal
structure compared to other Sciuridae. All chromosome-scale scaffolds show
indications for telomeres at both ends; the N50 value and busco as well as *k*-mer
completeness scores are high with 157.39 Mbp and 97 – 99 %, respectively,
indicating chromosome-level quality of the assembly. Based on whole-genome data
from 17 rodent species, *P. volans* clusters according to known evolutionary
relationships. Additionally, we present a new 16,511 bp long mitogenome unveiling
differences from known conspecific mitogenomes. We propose the utility of the new
reference genome for further research and development of conservation-applied
genetic methods.

Introduction

The Siberian flying squirrel (*Pteromys volans* (LINNAEUS, 1758); Figure 1a) is the only native representative of its tribe (Pteromyini BRANDT, 1855) in Europe, with its characteristic patagium enabling a highly arboreal lifestyle, gliding between trees and largely avoiding the ground (Thorington et al., 2012; Koprowski et al., 2016). While being nocturnal, *P. volans* is active throughout the boreal seasons and therefore exposed to dramatic environmental and nutritional changes, adapting its activity and behaviour accordingly (Hokkanen et al., 1977; Törmälä et al., 1980; Hanski, 1998; Hanski et al., 2000). After the Eurasian red squirrel (*Sciurus vulgaris* LINNAEUS, 1758), *P. volans* has the second-largest continuous distribution range of any Sciuromorpha BRANDT, 1855 ranging from Northeast Europe throughout the Eurasian boreal forests until the East Asian Pacific coasts including Manchuria, the Korean Peninsula and the islands of Sakhalin and Hokkaidō southwards limited by arid regions of Central Asia (Thorington et al., 2012; Koprowski et al., 2016). The highest phylogenetic and (mitochondrial as well as nuclear) genetic variability can be found among the East Asian populations (Oshida et al., 2005; Lee et al., 2008; Ito Dos Santos et al., 2024), reflected in comprising all four described and currently accepted subspecies (Thorington et al., 2012; Holden-Musser et al., 2016; Hopkins, 2016; Koprowski et al., 2016). In large portions of its Eurasian range, including the last remaining European populations, it is represented by the nominate subspecies *P. v. volans* (Lee et al., 2008; Thorington et al., 2012; Koprowski et al., 2016; Ito Dos Santos et al., 2024).

While its conservation status is globally considered as *Least Concern* by the International Union for Conservation of Nature and Natural Resources (IUCN), populations experience declining trends throughout most of their distribution (Hokkanen et al., 1982; Shar et al., 2016; Zagorodniuk, 2022). Lately, this species has been listed as *Near Threatened* throughout its European distribution (Amori, 2024). Nationally, they are red-listed as *Vulnerable* in Finland (Liukko et al., 2019) and as *Critically Endangered* in Estonia (EELIS, 2025). In fact, the species went extinct in several European countries over the last centuries. Recently, it disappeared in Latvia and Lithuania (Pilāts, 2010, 2023), while a minor population was rediscovered in Northeast Belarus (Abramchuk, 2021). They are legally protected

within the European Union (EU) through the Habitats Directive (*Council Directive*
92/43/EEC on the Conservation of natural habitats and of wild fauna and flora) under
Annexes II and *IV* (European Commission, 2021). On the contrary, it is not
continuously protected throughout most of its distribution in Russia (Kurhinen et al.,
2011), which represents the majority of its range. Given its relatively large population
size, Finland holds special conservation responsibilities within the EU. Regardless of
its protection status, its populations are still decreasing mostly due to habitat loss and
fragmentation through extensive forestry (Lampila et al., 2009). Additionally, recent
changes in Finnish conservation and forestry legislations made the execution of
protection of this native rodent ineffective (Santangeli et al., 2013; Wistbacka et al.,
2018). Even though Northeast Europe is much less populated by humans compared
to the rest of the subcontinent, it is nevertheless highly exploited and thus causes the
loss of old-growth mixed forests. The foliage of conifers and cavities in large tree
trunks provide cover all year round, while the flying squirrel largely feeds on leaves
and catkins of birch and aspen during the warmer season (Hanski et al., 2000;
Thorington et al., 2012; Koprowski et al., 2016; Selonen and Wistbacka, 2016). Such
dependency on old-growth forests (Mönkkönen et al., 1997; Hanski, 1998) and its
charismatic, ecological and phylogenetic uniqueness in Europe makes *P. volans* also
an umbrella and flagship species for habitat conservation of intact boreal forests
(Hurme et al., 2008; Selonen and Mäkeläinen, 2017). Despite the current
conservation focus, the ecology and intraspecific phylogeography of the Siberian
flying squirrel is widely unknown.

High-contiguity genome assemblies serve as invaluable references for uncovering
such unknown aspects of a species' biology and advancing applied conservation
genetics (Brandies et al., 2019; Paez et al., 2022). Those assemblies not only
enhance our understanding of population structures but also enable the identification
of informative genetic markers, such as single-nucleotide polymorphisms (SNPs) and
microsatellites. With their high contiguity, these reference genomes lay the foundation
for developing genetic tools with broad applications in conservation efforts as needed
for the Siberian flying squirrel.

While a draft genome assembly of *Glaucomys volans* (LINNAEUS, 1758), native to
eastern and southern North America (Koprowski et al., 2016), was published (Wolf et
al., 2022), our study provides the first chromosome-level assembly of any Pteromyini.

Only such high-contiguity assemblies enable comprehensive analysis of
chromosomal structures, including improved SNP localisation, which facilitates
linkage disequilibrium mapping and the study of functional diversity in genes, as well
as the identification of runs of homozygosity and repeat regions. Here, we present
the first high-contiguity, chromosome-level *de novo* genome assembly of *Pteromys*
volans.

Material and Methods

Sample, DNA extraction and sequencing library preparations

High molecular weight DNA was extracted from muscle tissue taken from a whole-
frozen specimen. The specimen died of natural causes and was collected post-
mortem at 62.917950°N, 27.631830°E (WGS84) near Kuopio, Finland at the 26[th]
March 2020. It was morphologically assigned as a male of the nominate subspecies
of *Pteromys volans volans*. The specimen is stored at -20°C in the collection of the
Zoological Museum of the University of Oulu (ZMUO; Collection Management
System Kotka/FinBIF: http://id.zmuo.oulu.fi/OV.36069; GBIF:
https://www.gbif.org/occurrence/4977508301 (Finbif, 2024)).

We used the *NEB Monarch® HMW DNA Extraction Kit for Cells & Blood* (NEB, USA)
for the DNA extraction following the manufacturers protocol. Next, we measured DNA
concentration and length utilising the *Qubit dsDNA BR Assay* kit at a Qubit
Fluorometer 2.0 (Thermo Fisher Scientific), gel electrophoresis and the *Agilent High*
Sensitivity DNA Kit for the *Agilent 2100 Bioanalyzer* (Agilent Technologies, USA).
Two *SMRTbell* libraries were prepared according to the instructions of the *SMRTbell*
Express Prep Kit v3.0 and sequenced on a *PacBio Revio* system (PacBio, Menlo
Park, USA). Next, we generated a proximity ligation sequencing library from the
same individual using the *Dovetail Omni-C* Kit (Cantata Bio, USA) following the
manufacturers protocol, which was sequenced by *Novogene* in Munich on a
NovaSeq X platform (Illumina, Inc., San Diego, CA, USA).

Raw data processing, assembly and scaffolding

First, we extracted the HiFi reads from the BAM file and converted them to the
FASTQ format using *BamTools* v2.5.2 (Barnett et al., 2011) with the *-tag "rq":">=0.99"*

option and *BEDtools* v2.31.1 (Quinlan and Hall, 2010) with the *bamToFastq* option.
We then assembled baseline contig-level genome assemblies using the HiFi reads
and (1) the *Hifiasm* v0.19.6-r597 assembler (Cheng et al., 2021) and (2) the *Flye*
v2.9.4 assembler (Kolmogorov et al., 2019). Based on the *BUSCO* v5.4.7 (Manni et
al., 2021) scores and assembly continuity statistics (calculated using *abyss-fac*,
(Simpson et al., 2009)) we selected the *Hifiasm* assembly for scaffolding using the
Omni-C data. We ran *BUSCO* v5.4.7 in *euk_genome_met* mode and with the gene
predictor *metaeuk* (Levy Karin et al., 2020) and the lineage dataset
mammalia_odb10 (which included 24 genomes of 9,226 buscos).

We used Dovetail Genomics's library QC pipeline (https://omni-
c.readthedocs.io/en/latest/index.html) to assess the quality of the Omni-C library. For
the Omni-C scaffolding, we applied the Vertebrate Genome Project (VGP) Hi-C
scaffolding pipeline (Rhie et al., 2021). In brief, we first filtered the Omni-C reads
using the filter_five_end.pl *perl* script
(https://github.com/ignacio3437/HiC_mapping_pipeline/tree/master). Next, the reads
were merged and PCR duplicates were removed using *Sambamba* v1.0.1 (Tarasov
et al., 2015). We used *YaHS* v1.2 (Zhou et al., 2023) for the Omni-C scaffolding and
displayed the contact map using *JuicerTools* v1.22.01
(https://github.com/aidenlab/juicertools). Next, we carried out gapclosing using *TGS-*
GapCloser v2.0.0 (Xu et al., 2020). We then assessed the scaffolded assembly
quality using *BUSCO* v5.4.7 and abyss-fac. We then performed another round of
Omni-C scaffolding using the VGP Hi-C scaffolding pipeline and gapclosing using
TGS-GapCloser v2.0.0, followed by another assembly quality assessment (as
described above). Finally, we visualised the generated Omni-C contact map and
manually curated the scaffolding using *Juicebox* v2.17.00 (Dudchenko et al., 2018).

Quality assessment, Repeat and Gene Annotation

The final assembly was assessed with *QUAST* v5.2.0 (Gurevich et al., 2013),
compleasm v0.2.6, which was shown to be more accurate than *BUSCO* (Huang and
Li, 2023), and *BUSCO* v5.4.7, to be able to compare with previous runs. The latter
two each utilising the lineage dataset *mammalia_odb10*. In addition, we also
investigated the *k*-mer completeness (%; *k* = 21), the consensus quality value (QV;
Phred scale), and the *k*-mer multiplicity using *Meryl* v1.4.1 and *Merqury* v1.3 (Rhie et

al., 2020). The *k*-mer multiplicity was also esteiemented with *GenomeScope* v2.0
(Ranallo-Benavidez et al., 2020). We further estimated the genome-level and
scaffold-level coverage using *Qualimap* v2.3 (Okonechnikov et al., 2016) and
assessed potential contamination using *BlobTools* v1.1 (Laetsch et al., 2017; Laetsch
and Blaxter, 2017) and visualised in *BlobPlots* based on protein sequences found in
the database 'Swiss-Prot' (release: 04-2024; The UniProt Consortium, 2019) by
diamond v2.1.6 (Buchfink et al., 2021). To do so, we mapped the HiFi reads onto the
Omni-C scaffolded assembly using *minimap2* v2.24 (Li, 2018, 2021). We further
identified mitochondrial contigs in the assembly by blasting our assembled
mitochondrial genome (outlined in the next section) against the final genome
assembly using *NCBI BLAST* v2.16.0 (Boratyn et al., 2013) and removed them from
the assembly. The SRY gene (XM_047537220.1) extracted from an eastern grey
squirrel genome assembly (*Sciurus carolinensis* Gmelin, 1788; GCA_902686445.2
(Mead et al., 2020a)) was aligned to the novel genome assembly using blastn (*NCBI
BLAST* v2.16.0) to determine its genomic location.

To annotate repeats in the genome assembly, we first used *RepeatModeler* v2.0.5
(Flynn et al., 2020) with the search engine *rmblast* v2.14.1+ and *RepeatMasker*
v4.1.6 (Smit et al., 2023) with the *Repbase* 'rodentia' database (Jurka et al., 2005)
and the additional options of: *-nolow* to not hard-mask low complexity DNA or simple
repeats and *-gccalc* to calculate the GC content of each contig/scaffold individually.
Next, we applied *RepeatMasker* v4.1.6 to the masked genome but with the options: *-
xsmall* to change the masking mode from hard to soft-masked, *-noint* to only hard-
masks low complex/simple repeats, excluding intronic repeats, and *-gccalc.*

Genes in the masked assembly were predicted based on homology with Gene Model
Mapper *GeMoMa* v.1.9 (Keilwagen et al., 2016, 2018) using the following publicly
available assemblies at NCBI GenBank: human (*Homo sapiens* Linnaeus, 1758;
GCA_000001405.29)*,* house mouse (*Mus musculus* Linnaeus, 1758;
GCA_000001635.9), eastern grey squirrel (*Sciurus carolinensis*; GCA_902686445.2
(Mead et al., 2020a)), thirteen-lined ground squirrel (*Ictidomys tridecemlineatus*
(Mitchill, 1821); GCA_016881025.1 (Fu et al., 2021)), Arctic ground squirrel
(*Urocitellus parryii* Richardson, 1825; GCA_003426925.1), Alpine marmot (*Marmota
marmota marmota* (Linnaeus, 1758); GCA_001458135.2), and groundhog or
woodchuck (*Marmota monax* (Linnaeus, 1758); GCA_021218885.2 (Clarke and

Bader, 2024)). The busco completeness of the transcript annotation was assessed
with *BUSCO* v5.4.7 utilising the lineage dataset *mammalia_odb10*. Functional
annotation of the predicted proteins by *GeMoMa* was conducted by using *diamond*
(*blastp* mode) v2.1.6 (Buchfink et al., 2015) with an e-value significance cutoff of ≤
10^{-6} against the 'Swiss-Prot' database (release: 04-2024; The UniProt Consortium,
2019). Furthermore, we annotated gene ontology (GO) terms, domains, and motifs
using *InterProScan* v5.73-104.0 (Quevillon et al., 2005; Jones et al., 2014). The gene
annotation results were combined utilising modified functional annotation scripts from
MAKER v3.01.03 (Cantarel et al., 2008; https://github.com/schellt/maker-functional).
Gene annotation files are available on Dryad
(https://doi.org/10.5061/dryad.3xsj3txth).

Telomere regions were identified using *seqtk* with the option *telo*
(https://github.com/lh3/seqtk), and *quarTeT* v1.2.5 (Lin et al., 2023) with the options -
m 10 to *100*.

Lastly, we used the *circlize R* package v0.4.16 (Gu et al., 2014) to perform a six-track
circos plot for the largest 19 scaffolds. As an input, we used (A) the telomeres length
generated using -m 10, (B) the gene content clustered in 500 kbp windows, (C; D)
the repetitive elements (simple and complex repeats) were grouped according to their
type (the same TE type found sequentially) and plotted according to their total
relative length, and (E) GC content (%; compare Figure 2a).

Mitochondrial genome assembly

We used *MitoHifi* v3.2.2 (Uliano-Silva et al., 2023) to reconstruct and the *Flye* v2.9.4
polisher (Kolmogorov et al., 2019) to polish the mitochondrial genome.

Protein-coding genes, rRNAs and tRNAs were annotated and visualised with
Chlorobox online software tools (https://chlorobox.mpimp-golm.mpg.de/), *GeSeq*
(Tillich et al., 2017), and *OGDraw* (Greiner et al., 2019). For *GeSeq*, a circular
mitochondrial sequence source was selected, keeping only the best annotation.
Annotation via "*BLAT search*" (standard settings) with NCBI GenBank reference
MW722951.1 (*P. volans*) and third-party references from five Sciuridae Fɪꜱᴄʜᴇʀ ᴅᴇ
Wᴀʟᴅʜᴇɪᴍ, 1817 individuals: southern flying squirrel or assapan (*Glaucomys volans*;
NC_050026.1); particolored flying squirrel (*Hylopetes alboniger* (Hᴏᴅɢꜱᴏɴ, 1836);
NC_031847.1), red and white giant flying squirrel (*Petaurista alborufus* (Mɪʟɴᴇ-

EDWARDS, 1870); NC_023922.1), *P. volans* (NC_019612.1 (Ryu et al., 2013)), and

Eurasian red squirrel (*Sciurus vulgaris*; NC_002369.1 (Reyes et al., 2000)). GenBank

annotation was manually double-checked, and redundant replicates were removed

(keeping the version with best alignment scores). For *OGDraw*, a mitochondrial

sequence source was selected to visualise the generated GenBank file.

Synteny and phylogenetic analyses

Next, we assessed genome synteny to the chromosome-level reference genomes for

the Eurasian red squirrel (*Sciurus vulgaris*; GenBank: GCA_902686455.2 (Mead et

al., 2020b)), the eastern grey squirrel (*Sciurus carolinensis*; GenBank:

GCA_902686445.2 (Mead et al., 2020a)), and the European ground squirrel or

European souslik (*Spermophilus citellus* LINNAEUS, 1766; GenBank:

GCA_964194105.1) using JupiterPlot v1.1 (https://github.com/JustinChu/JupiterPlot)

using options: *ng = 101-110, m = 10000000*.

We then generated a phylogenetic tree for all available Sciuridae genomes and two

Muridae species as an outgroup on the NCBI GenBank: house mouse (*Mus

musculus*; GCA_000001635.9), black rat (*Rattus rattus* (LINNAEUS, 1758);

GCF_011064425.1 (Rane et al., 2020)), southern flying squirrel or assapan

(*Glaucomys volans*; GCA_020662805.1 (Wolf et al., 2022)), Eurasian red squirrel

(*Sciurus vulgaris*; GCA_902686455.2 (Mead et al., 2020b), eastern grey squirrel

(*Sciurus carolinensis*; GCA_902686445.2 (Mead et al., 2020a), eastern fox squirrel

(*Sciurus niger* LINNAEUS, 1758; GCA_020740815.1 (Kang et al., 2021)), unstriped

ground squirrel (*Xerus rutilus* (CRETZSCHMAR, 1828); GCA_028644305.1), Cape

ground squirrel (*Geosciurus inauris* (ZIMMERMAN, 1780); GCA_004024805.1),

Siberian chipmunk (*Eutamias sibiricus* (LAXMANN, 1769); GCA_025594165.1 (Li et al.,

2022)), Daurian ground squirrel (*Spermophilus dauricus* BRANDT, 1843;

GCA_002406435.1), thirteen-lined ground squirrel (*Ictidomys tridecemlineatus*;

GCA_016881025.1 (Fu et al., 2021)), Arctic ground squirrel (*Urocitellus parryii*;

GCA_003426925.1), Himalayan marmot (*Marmota himalayana* (HODGSON, 1841);

GCA_005280165.1 (Bai et al., 2019)), Alpine marmot (*Marmota marmota marmota*;

GCA_001458135.2), Vancouver Island marmot (*Marmota vancouverensis* (SWARTH,

1911); GCA_005458795.1), and yellow-bellied marmot or rock chuck (*Marmota

flaviventer* (AUDUBON & BACHMAN, 1841); GCA_003676075.3) using 3,269 busco

genes. To do so, we first obtained fasta files for each busco gene using

BUSCO_phylogenomics (https://github.com/jamiemcg/BUSCO_phylogenomics) with

default settings, and we aligned them individually using *Mafft* v7.505 (Katoh and

Standley, 2013). Next, we reconstructed phylogenetic trees for each gene separately

with *iqtree* v2.3.4 (Minh et al., 2020) utilising a GTR+I+G substitution model. Lastly,

we combined the gene trees into a species tree using a multi-species coalescent

model using *ASTRAL-III* v5.7.8 (Zhang et al., 2018). Phylogenetic discordance was

investigated and visualised using *DiscoVista* (Sayyari et al., 2018).

292 Results & Discussion

The 2.8 Gbp large genome of *Pteromys volans* is organised in the 19 largest

scaffolds (Figure 1b), representing the expected gonosome and 18 autosomes ($2n$ =

38; Oshida and Yoshida, 1996; Oshida et al., 2000; Table 1). Mapping the PacBio

HiFi reads back to the genome assembly indicates a mean coverage of 22.6×.

Scaffold CM117769.1 shows roughly half the mean genome coverage (12.1×;

Supplementary Figure S10, Supplementary Table S3) and minimal interchromosomal

contacts (Figure 1b), indicating weak or no interaction with other chromosomes. This

along with the chromosome assignment by NCBI GenBank and a blastn-based

search of the SRY gene against the novel genome assembly identities this scaffold

as the heteromorph gonosomes, which confirms the morphological sex determination

of the sequenced specimen. The total sequencing depth was not enough to

assemble the Y chromosome. Two runs of PacBio generated in total 64.7 Gbp in

filtered HiFi reads that were used for contig genome assembly (run 1: 63.9 Gbp,

7,143,058 reads; run 2: 836.4 Mbp, 90,539 reads). We were able to close two

assembly gaps with *TGS-GapCloser*. *Compleasm* scores (99.9% complete, 0.78%

duplicated, 0% fragmented and only 0.03% missing genes; Table 2) and *BUSCO*

scores (97% complete, 2% duplication, 0.6% fragmented and only 2.4% missing

genes; Table 2), a *k*-mer completeness of 97.04% (Consensus QV = 42.7294,

corresponding to an error rate of 5.33 × $10^{-\square}$), the *k*-mer multiplicity spectrum

(Supplementary Figure S9), and assembly statistics (scaffold-level N50 = 157 Mbp,

contig-level N50 = 60Mbp, scaffold-level L90 = 18, contig-level L90 = 73, etc.; Table

1) indicate a high contiguity. The *k*-mer multiplicity corresponds closely to the

expected genome coverage of ~22× (Supplementary Figure S9). *GenomeScope*

showed a *k*-mer spectrum of an individual with extremely low heterozygosity,

estimated as 0.2%, consistent with a previous study reporting very low
heterozygosities for the Finnish population of *P. volans* from which the sequenced
individual originates (Ito Dos Santos et al., 2024). The largest 19 scaffolds cover 92%
of the assembly and showed at least 10 copies of the telomere repeat TTAGGG on
either end. More than 100 copies were found on both ends for nine and at one end
for seven additional scaffolds. Only three scaffolds did not show telomeres with more
than 100 copies of TTAGGG (Figure 2a, Supplementary Figure S6). The 19 largest
scaffolds reached *compleasm* scores of 99.89% complete, 0.78% duplicated, 0.07%
fragmented and 0.04% missing genes, only one complete gene less than the full
assembly, and *BUSCO* scores equal to the whole assembly, indicating a high
completeness of the potential chromosomes. The GC-content with 40.77% lies
slightly below eutherian levels of 40.9 – 41.8% (Duret and Galtier, 2009). The
homology-based gene prediction with *GeMoMa* identified a total of 27,383 genes and
54,238 transcripts with a *BUSCO* completeness of 91.5% (Supplementary Table S2).
More than 97.26% (52,752) of those unique protein query sequences were annotated
to sequences from the 'Swiss-Prot' database with the *Diamond blastp*-option. The
functional annotation with *InterProScan* identified 53,845 unique protein sequences,
while 73.3% (39,487) could be assigned to at least one Gene Ontology (GO) term.
Blobtools identified 17 scaffolds belonging to viruses of the phylum Artverviricota
(Supplementary Figure S8), which were removed from the assembly. In total,
970,478,186 bp (34.07%; Table 3) were masked as repeat elements.

The novel genome assembly phylogenetically clusters with the genomic data of the
other published members of Sciuridae as expected based on other studies (Zelditch
et al., 2015; Menéndez et al., 2021; Figure 3).

Members of the sister tribe Sciurini LINNAEUS, 1758 showed only minor synteny
differences in their chromosomal architecture (Figure 1c). Noteworthy is the
apomorph chromosomal fission of scaffold 1 shown in both *Sciurus* species
investigated here (Figure 1c). A fused scaffold 1 seems to be a plesiomorph trait kept
in *P. volans* if compared to a member of Xerinae OSBORN, 1910 as a clade outside of
Sciurinae FISCHER DE WALDHEIM, 1817 (compare to the phylogenetic tree in Figure 3).
More distant taxa, such as the European ground squirrel, show a higher degree of
chromosomal rearrangements (Figure 1c).

As mentioned above, *Blobtools* identified scaffolds belonging to the viral family
Retroviridae. Due to this phylogenetic assignment, those sequences might originate
from proviral DNA rather than pararetroviral virions. But we cannot infer if the viruses
were active or rather in a proviral phase. Since, the cause for its natural death is
unknown and another adult male individual was found with the sequenced specimen,
it could shed light on a potential pathogenic relationship of *P. volans* with retroviruses.

The mitochondrial genome has a length of 16,511 bp, including 13 protein-coding
genes (CytB, ND1-6, COX1-3, ATP6, and ATP8), two ribosomal RNA genes (12S
rRNA and 16S rRNA), and 22 tRNA genes (Figure 2b). This new mitogenome, from
the westernmost distribution, contains two fewer base pairs than that of an individual
from the Korean Peninsula (Ryu et al., 2013), one of the easternmost ranges, mostly
considered to belong to the subspecies *Pteromys volans buechneri* SATUNIN, 1903
(Thorington et al., 2012; Koprowski et al., 2016). Previous genetic studies showed a
consistent differentiation between the East Asian mainland populations and the
northern Eurasian populations (Oshida et al., 2005; Koh et al., 2008; Ito Dos Santos
et al., 2024). However, a more comprehensive sampling and its genomic analysis is
required to further resolve the intraspecific phylogeography of the Siberian flying
squirrel (Ito Dos Santos et al., 2024), which could be facilitated by the newly
presented whole-genome assembly.

For the first time, we provide a chromosome-level genome assembly of a flying
squirrel species, namely *Pteromys v. volans* inhabiting extremely fluctuating
environments, which will help to understand many aspects of the species' biology,
such as phylogeny, ecological, and nutritional genomics (Ungerer et al., 2008). The
sequenced specimen is a representative of the species' westernmost population,
presumably only connected by a narrow corridor to the metapopulation (Kurhinen et
al., 2011) with a current conservation focus (Flying Squirrel LIFE; LIFE17
NAT/FI/000469). A previously shown low standing genetic variation within this
western population (Ito Dos Santos et al., 2024) is of considerable importance for its
adaptive capacity and inferred conservation strategies. Thus, necessary in-depth
investigations on inbreeding through runs of homozygosity (ROH) or the genetic
health looking at mutational load can only be accomplished with whole-genome
assemblies, as provided here. It will also constitute the reference to identify
informative genetic markers for conservation applications such as population

monitoring. Therefore, this novel genome assembly has the potential to not only shed

light on processes of natural selection and adaptation interesting beyond the species

context but will provide the basis for applied conservation-relevant studies on the

population genetics of the species itself.

386 References

Abramchuk, A.V., 2021. The Siberian flying squirrel (Pteromys volans) in Belarus:
distribution, abundance, threats, and conservation. Theriol. Ukr. 2021, 69–79.
https://doi.org/10.15407/TU2208
Amori, G., 2024. Pteromys volans (Europe assessment). The IUCN Red List of
Threatened Species 2024: e.T18702A221782992.
https://doi.org/10.2305/IUCN.UK.2024-2.RLTS.T18702A221782992.en
Bai, L., Liu, B., Ji, C., Zhao, S., Liu, S., Wang, R., Wang, W., Yao, P., Li, X., Fu, X.,
Yu, H., Liu, M., Han, F., Guan, N., Liu, H., Liu, D., Tao, Y., Wang, Z., Yan, S.,
Florant, G., Butcher, M.T., Zhang, J., Zheng, H., Fan, J., Liu, E., 2019. Hypoxic
and Cold Adaptation Insights from the Himalayan Marmot Genome. iScience
11, 505–507. https://doi.org/10.1016/j.isci.2019.01.019
Barnett, D.W., Garrison, E.K., Quinlan, A.R., Strömberg, M.P., Marth, G.T., 2011.
BamTools: a C++ API and toolkit for analyzing and managing BAM files.
Bioinformatics 27, 1691–1692. https://doi.org/10.1093/bioinformatics/btr174
Boratyn, G.M., Camacho, C., Cooper, P.S., Coulouris, G., Fong, A., Ma, N., Madden,
T.L., Matten, W.T., McGinnis, S.D., Merezhuk, Y., Raytselis, Y., Sayers, E.W.,
Tao, T., Ye, J., Zaretskaya, I., 2013. BLAST: a more efficient report with
usability improvements. Nucleic Acids Research 41, W29–W33.
https://doi.org/10.1093/nar/gkt282
Brandies, P., Peel, E., Hogg, C.J., Belov, K., 2019. The Value of Reference Genomes
in the Conservation of Threatened Species. Genes 10, 846.
https://doi.org/10.3390/genes10110846
Buchfink, B., Reuter, K., Drost, H.-G., 2021. Sensitive protein alignments at tree-of-
life scale using DIAMOND. Nat Methods 18, 366–368.
https://doi.org/10.1038/s41592-021-01101-x
Buchfink, B., Xie, C., Huson, D.H., 2015. Fast and sensitive protein alignment using
DIAMOND. Nat Methods 12, 59–60. https://doi.org/10.1038/nmeth.3176
Cantarel, B.L., Korf, I., Robb, S.M.C., Parra, G., Ross, E., Moore, B., Holt, C.,
Sánchez Alvarado, A., Yandell, M., 2008. MAKER: An easy-to-use annotation
pipeline designed for emerging model organism genomes. Genome Res. 18,
188–196. https://doi.org/10.1101/gr.6743907
Cheng, H., Concepcion, G.T., Feng, X., Zhang, H., Li, H., 2021. Haplotype-resolved
de novo assembly using phased assembly graphs with hifiasm. Nat Methods
18, 170–175. https://doi.org/10.1038/s41592-020-01056-5
Clarke, Z., Bader, G., 2024. Healthy woodchuck genome with viral sequences
appended used for single-cell RNA-seq analysis.
https://doi.org/10.5281/ZENODO.10855128
Dudchenko, O., Shamim, M.S., Batra, S.S., Durand, N.C., Musial, N.T., Mostofa, R.,
Pham, M., Glenn St Hilaire, B., Yao, W., Stamenova, E., Hoeger, M., Nyquist,
S.K., Korchina, V., Pletch, K., Flanagan, J.P., Tomaszewicz, A., McAloose, D.,
Pérez Estrada, C., Novak, B.J., Omer, A.D., Aiden, E.L., 2018. The Juicebox

Assembly Tools module facilitates *de novo* assembly of mammalian genomes with chromosome-length scaffolds for under $1000. https://doi.org/10.1101/254797

Duret, L., Galtier, N., 2009. Biased Gene Conversion and the Evolution of Mammalian Genomic Landscapes. Annu. Rev. Genom. Hum. Genet. 10, 285–311. https://doi.org/10.1146/annurev-genom-082908-150001

EELIS, 2025. EELIS (Eesti looduse infosüsteem). Keskkonnaagentuur. Pteromys volans (lendorav). https://www.eelis.ee/lnim/-646374363.

European Commission, 2021. Commission notice Guidance document on the strict protection of animal species of Community interest under the Habitats Directive.

Finbif, 2024. ZMUO Common vertebrate collection. https://doi.org/10.15468/U5MR6A

Flynn, J.M., Hubley, R., Goubert, C., Rosen, J., Clark, A.G., Feschotte, C., Smit, A.F., 2020. RepeatModeler2 for automated genomic discovery of transposable element families. Proc. Natl. Acad. Sci. U.S.A. 117, 9451–9457. https://doi.org/10.1073/pnas.1921046117

Fu, R., Gillen, A.E., Grabek, K.R., Riemondy, K.A., Epperson, L.E., Bustamante, C.D., Hesselberth, J.R., Martin, S.L., 2021. Dynamic RNA Regulation in the Brain Underlies Physiological Plasticity in a Hibernating Mammal. Front. Physiol. 11, 624677. https://doi.org/10.3389/fphys.2020.624677

Greiner, S., Lehwark, P., Bock, R., 2019. OrganellarGenomeDRAW (OGDRAW) version 1.3.1: expanded toolkit for the graphical visualization of organellar genomes. Nucleic Acids Research 47, W59–W64. https://doi.org/10.1093/nar/gkz238

Gu, Z., Gu, L., Eils, R., Schlesner, M., Brors, B., 2014. *circlize* implements and enhances circular visualization in R. Bioinformatics 30, 2811–2812. https://doi.org/10.1093/bioinformatics/btu393

Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G., 2013. QUAST: quality assessment tool for genome assemblies. Bioinformatics 29, 1072–1075. https://doi.org/10.1093/bioinformatics/btt086

Hanski, I.K., 1998. Home ranges and habitat use in the declining flying squirrel *Pteromys volans* in managed forests. Wildlife Biology 4, 33–46. https://doi.org/10.2981/wlb.1998.013

Hanski, I.K., Stevens, P.C., Ihalempiä, P., Selonen, V., 2000. Home-Range Size, Movements, and Nest-Site Use in the Siberian Flying Squirrel, *Pteromys volans*. Journal of Mammalogy 81, 798–809. https://doi.org/10.1644/1545-1542(2000)081<0798:HRSMAN>2.3.CO;2

Hokkanen, H., Törmälä, T., Vuorinen, H., 1982. Decline of the flying squirrel Pteromys volans l. populations in Finland. Biological Conservation 23, 273–284. https://doi.org/10.1016/0006-3207(82)90083-0

Hokkanen, H., Törmälä, T., Vuorinen, H., 1977. Seasonal changes in the circadian activity of Pteromys volans L. in central Finland. Annales Zoologici Fennici 14, 94–97.

Holden-Musser, M.E., Juškaitis, R., Musser, G., 2016. Family Dormice (Gliridae), in: Handbook of the Mammals of the World. Vol. 6. Lagomorphs and Rodents I. pp. 838–892.

Hopkins, S.S.B., 2016. Family Aplodontiidae (Mountain Beaver), in: Handbook of the Mammals of the World. Vol. 6. Lagomorphs and Rodents I. pp. 642–647.

Huang, N., Li, H., 2023. compleasm: a faster and more accurate reimplementation of BUSCO. Bioinformatics 39, btad595. https://doi.org/10.1093/bioinformatics/btad595

Hurme, E., Mönkkönen, M., Sippola, A.-L., Ylinen, H., Pentinsaari, M., 2008. Role of
the Siberian flying squirrel as an umbrella species for biodiversity in northern
boreal forests. Ecological Indicators 8, 246–255.
https://doi.org/10.1016/j.ecolind.2007.02.001
Ito Dos Santos, F., Lilley, T., Laine, V.N., Prost, S., Kurhinen, J., Susi, H., Gashev, S.,
Saveljev, A., Bondarchuk, S., Babina, S., Shishikin, A., Tirski, D., Vasina, A.,
Karpin, V., Kekkonen, J.A., 2024. Population structure of the endangered
Siberian flying squirrel (*Pteromys volans*) revealed by genomic and
mitochondrial data. https://doi.org/10.1101/2024.09.13.612918
Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H.,
Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas,
A., Scheremetjew, M., Yong, S.-Y., Lopez, R., Hunter, S., 2014. InterProScan
5: genome-scale protein function classification. Bioinformatics 30, 1236–1240.
https://doi.org/10.1093/bioinformatics/btu031
Jurka, J., Kapitonov, V.V., Pavlicek, A., Klonowski, P., Kohany, O., Walichiewicz, J.,
2005. Repbase Update, a database of eukaryotic repetitive elements.
Cytogenet Genome Res 110, 462–467. https://doi.org/10.1159/000084979
Kang, L., Michalak, P., Hallerman, E., Moncrief, N.D., 2021. A draft genome assembly
for the eastern fox squirrel, *Sciurus niger*. G3 Genes|Genomes|Genetics 11,
jkab315. https://doi.org/10.1093/g3journal/jkab315
Katoh, K., Standley, D.M., 2013. MAFFT Multiple Sequence Alignment Software
Version 7: Improvements in Performance and Usability. Molecular Biology and
Evolution 30, 772–780. https://doi.org/10.1093/molbev/mst010
Keilwagen, J., Hartung, F., Paulini, M., Twardziok, S.O., Grau, J., 2018. Combining
RNA-seq data and homology-based gene prediction for plants, animals and
fungi. BMC Bioinformatics 19, 189. https://doi.org/10.1186/s12859-018-2203-5
Keilwagen, J., Wenk, M., Erickson, J.L., Schattat, M.H., Grau, J., Hartung, F., 2016.
Using intron position conservation for homology-based gene prediction.
Nucleic Acids Res 44, e89–e89. https://doi.org/10.1093/nar/gkw092
Koh, H.-S., Jin, Y., Yang, B.-G., Lee, B.-K., Heo, S.-W., Jang, K.-H., 2008. Taxonomic
Status of Siberian Flying Squirrel from Korea (Pteromys volans aluco Thomas
1907). Animal Systematics, Evolution and Diversity 24, 169–172.
https://doi.org/10.5635/KJSZ.2008.24.2.169
Kokot, M., Długosz, M., Deorowicz, S., 2017. KMC 3: counting and manipulating k -
mer statistics. Bioinformatics 33, 2759–2761.
https://doi.org/10.1093/bioinformatics/btx304
Kolmogorov, M., Yuan, J., Lin, Y., Pevzner, P.A., 2019. Assembly of long, error-prone
reads using repeat graphs. Nat Biotechnol 37, 540–546.
https://doi.org/10.1038/s41587-019-0072-8
Koprowski, J., Goldstein, E., Bennett, K., Pereira, C., 2016. Family Sciuridae (Tree,
Flying and Ground Squirrels, Chipmunks, Prairie Dogs and Marmots), in:
Handbook of the Mammals of the World. Vol. 6. Lagomorphs and Rodents I.
pp. 648–837.
Kurhinen, J., Kulebyakina, E., Zadiraka, E., Mamontov, V., Muravskaya, E., Hanski,
I.K., 2011. Distribution of the Siberian flying squirrel (*Pteromys volans* L.) in
taiga isthmuses between Baltic and White Sea regions. Acta Zoologica
Lituanica 21, 306–310. https://doi.org/10.2478/v10043-011-0037-8
Laetsch, D.R., Blaxter, M.L., 2017. BlobTools: Interrogation of genome assemblies.
F1000Res 6, 1287. https://doi.org/10.12688/f1000research.12232.1
Laetsch, D.R., Koutsovoulos, G., Booth, T., Stajich, J., Sujai Kumar, 2017.
DRL/blobtools: BlobTools v1.0.1. https://doi.org/10.5281/ZENODO.845347

Lampila, S., Wistbacka, R., Mäkelä, A., Orell, M., 2009. Survival and population
growth rate of the threatened Siberian flying squirrel (*Pteromys volans*) in a
fragmented forest landscape. Écoscience 16, 66–74.
https://doi.org/10.2980/16-1-3184
Lee, Mu-Yeong, Park, S., Hong, Y., Kim, Y., Voloshina, I., Myslenkov, A., Saveljev,
A.P., Choi, T., Piao, R., An, J., Lee, Mun-Han, Lee, H., Min, M., 2008.
Mitochondrial genetic diversity and phylogenetic relationships of Siberian flying
squirrel *(Pteromys volans)* populations. Animal Cells and Systems 12, 269–
277. https://doi.org/10.1080/19768354.2008.9647182
Levy Karin, E., Mirdita, M., Söding, J., 2020. MetaEuk—sensitive, high-throughput
gene discovery, and annotation for large-scale eukaryotic metagenomics.
Microbiome 8, 48. https://doi.org/10.1186/s40168-020-00808-x
Li, H., 2021. New strategies to improve minimap2 alignment accuracy. Bioinformatics
37, 4572–4574. https://doi.org/10.1093/bioinformatics/btab705
Li, H., 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics
34, 3094–3100. https://doi.org/10.1093/bioinformatics/bty191
Li, R., Zhang, M., Cha, M., Xiang, J., Yi, X., 2022. Chromosome-level genome
assembly of the Siberian chipmunk (Tamias sibiricus). Sci Data 9, 783.
https://doi.org/10.1038/s41597-022-01910-5
Lin, Y., Ye, C., Li, X., Chen, Q., Wu, Y., Zhang, F., Pan, R., Zhang, S., Chen, S.,
Wang, X., Cao, S., Wang, Y., Yue, Y., Liu, Y., Yue, J., 2023. quarTeT: a
telomere-to-telomere toolkit for gap-free genome assembly and centromeric
repeat identification. Horticulture Research 10, uhad127.
https://doi.org/10.1093/hr/uhad127
Liukko, U.-M., Henttonen, H., Kauhala, K., Kojola, I., Kyheröinen, E.-M., Pitkänen, J.,
2019. Mammals, in: The 2019 Red List of Finnish Species. Ministry of the
Environment & Finnish Environment Institute. Helsinki.
Manni, M., Berkeley, M.R., Seppey, M., Simão, F.A., Zdobnov, E.M., 2021. BUSCO
Update: Novel and Streamlined Workflows along with Broader and Deeper
Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral
Genomes. Molecular Biology and Evolution 38, 4647–4654.
https://doi.org/10.1093/molbev/msab199
Mead, D., Fingland, K., Cripps, R., Portela Miguez, R., Smith, M., Corton, C., Oliver,
K., Skelton, J., Betteridge, E., Dolucan, J., Dudchenko, O., Omer, A.D., Weisz,
D., Lieberman Aiden, E., Fedrigo, O., Mountcastle, J., Jarvis, E., McCarthy,
S.A., Sims, Y., Torrance, J., Tracey, A., Howe, K., Challis, R., Durbin, R.,
Blaxter, M., 2020b. The genome sequence of the Eurasian red squirrel,
Sciurus vulgaris Linnaeus 1758. Wellcome Open Res 5, 18.
https://doi.org/10.12688/wellcomeopenres.15679.1
Mead, D., Fingland, K., Cripps, R., Portela Miguez, R., Smith, M., Corton, C., Oliver,
K., Skelton, J., Betteridge, E., Doulcan, J., Quail, M.A., McCarthy, S.A., Howe,
K., Sims, Y., Torrance, J., Tracey, A., Challis, R., Durbin, R., Blaxter, M.,
2020a. The genome sequence of the eastern grey squirrel, Sciurus
carolinensis Gmelin, 1788. Wellcome Open Res 5, 27.
https://doi.org/10.12688/wellcomeopenres.15721.1
Menéndez, I., Gómez Cano, A.R., Cantalapiedra, J.L., Peláez-Campomanes, P.,
Álvarez-Sierra, M.Á., Hernández Fernández, M., 2021. A multi-layered
approach to the diversification of squirrels. Mammal Review 51, 66–81.
https://doi.org/10.1111/mam.12215
Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., Von
Haeseler, A., Lanfear, R., 2020. IQ-TREE 2: New Models and Efficient

Methods for Phylogenetic Inference in the Genomic Era. Molecular Biology
and Evolution 37, 1530–1534. https://doi.org/10.1093/molbev/msaa015
Mönkkönen, M., Reunanen, P., Nikula, A., Inkeröinen, J., Forsman, J., 1997.
Landscape characteristics associated with the occurrence of the flying squirrel
Pteromys volans in old‑growth forests of northern Finland. Ecography 20,
634–642. https://doi.org/10.1111/j.1600-0587.1997.tb00432.x
Okonechnikov, K., Conesa, A., García-Alcalde, F., 2016. Qualimap 2: advanced multi-
sample quality control for high-throughput sequencing data. Bioinformatics 32,
292–294. https://doi.org/10.1093/bioinformatics/btv566
Oshida, T., Abramov, A., Yanagawa, H., Masuda, R., 2005. Phylogeography of the
Russian flying squirrel (*Pteromys volans*): implication of refugia theory in
arboreal small mammal of Eurasia. Molecular Ecology 14, 1191–1196.
https://doi.org/10.1111/j.1365-294X.2005.02475.x
Oshida, T., Yanagawa, H., Tsuda, M., Indue, S., Yoshida, M.C., 2000. Comparisons of
the banded karyotypes between the smallJapanese flying squirrel,Pteromys
momonga and the Russian flying squirrel, P. volans(Rodentia, Sciuridae).
Caryologia 53, 133–140. https://doi.org/10.1080/00087114.2000.10589188
Oshida, T., Yoshida, M.C., 1996. Banded karyotypes and the localization of ribosomal
RNA genes of Eurasian flying squirrel, *Pteromys volans orii* (Mammalia,
Rodentia). Caryologia 49, 219–225.
https://doi.org/10.1080/00087114.1996.10797366
Paez, S., Kraus, R.H.S., Shapiro, B., Gilbert, M.T.P., Jarvis, E.D., Vertebrate
Genomes Project Conservation Group, Al-Ajli, F.O., Ceballos, G., Crawford,
A.J., Fedrigo, O., Johnson, R.N., Johnson, W.E., Marques-Bonet, T., Morin,
P.A., Mueller, R.C., Ryder, O.A., Teeling, E.C., Venkatesh, B., 2022. Reference
genomes for conservation. Science 377, 364–366.
https://doi.org/10.1126/science.abm8127
Pilāts, V., 2023. Abstracts of the 81st Scientific Conference of the University of Latvia:
January – February 2023. EEB 21, 33–44. https://doi.org/10.22364/eeb.21.05
Pilāts, V., 2010. Results of flying squirrel survey in Latvia., Wildlife census methods:
reliability and application. Workshop proceedings. International Scientific
Practical Workshop. Kaunas.
Quevillon, E., Silventoinen, V., Pillai, S., Harte, N., Mulder, N., Apweiler, R., Lopez,
R., 2005. InterProScan: protein domains identifier. Nucleic Acids Research 33,
W116–W120. https://doi.org/10.1093/nar/gki442
Quinlan, A.R., Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics 26, 841–842.
https://doi.org/10.1093/bioinformatics/btq033
Ranallo-Benavidez, T.R., Jaron, K.S., Schatz, M.C., 2020. GenomeScope 2.0 and
Smudgeplot for reference-free profiling of polyploid genomes. Nat Commun
11, 1432. https://doi.org/10.1038/s41467-020-14998-3
Rane, R., Pichlmueller, F., Lee, R., Edwards, O., Gemmell, N., Tompkins, D., 2020.
Rattus rattus chromosomal assembly (New zealand strain).
https://doi.org/10.25919/5E4F60EA8B64E
Reyes, A., Gissi, C., Pesole, G., Catzeflis, F.M., Saccone, C., 2000. Where Do
Rodents Fit? Evidence from the Complete Mitochondrial Genome of Sciurus
vulgaris. Molecular Biology and Evolution 17, 979–983.
https://doi.org/10.1093/oxfordjournals.molbev.a026379
Rhie, A., McCarthy, S.A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-
Silva, M., Chow, W., Fungtammasan, A., Kim, J., Lee, C., Ko, B.J., Chaisson,
M., Gedman, G.L., Cantin, L.J., Thibaud-Nissen, F., Haggerty, L., Bista, I.,

Smith, M., Haase, B., Mountcastle, J., Winkler, S., Paez, S., Howard, J.,
Vernes, S.C., Lama, T.M., Grutzner, F., Warren, W.C., Balakrishnan, C.N.,
Burt, D., George, J.M., Biegler, M.T., Iorns, D., Digby, A., Eason, D.,
Robertson, B., Edwards, T., Wilkinson, M., Turner, G., Meyer, A., Kautt, A.F.,
Franchini, P., Detrich, H.W., Svardal, H., Wagner, M., Naylor, G.J.P., Pippel,
M., Malinsky, M., Mooney, M., Simbirsky, M., Hannigan, B.T., Pesout, T.,
Houck, M., Misuraca, A., Kingan, S.B., Hall, R., Kronenberg, Z., Sović, I.,
Dunn, C., Ning, Z., Hastie, A., Lee, J., Selvaraj, S., Green, R.E., Putnam, N.H.,
Gut, I., Ghurye, J., Garrison, E., Sims, Y., Collins, J., Pelan, S., Torrance, J.,
Tracey, A., Wood, J., Dagnew, R.E., Guan, D., London, S.E., Clayton, D.F.,
Mello, C.V., Friedrich, S.R., Lovell, P.V., Osipova, E., Al-Ajli, F.O., Secomandi,
S., Kim, H., Theofanopoulou, C., Hiller, M., Zhou, Y., Harris, R.S., Makova,
K.D., Medvedev, P., Hoffman, J., Masterson, P., Clark, K., Martin, F., Howe,
Kevin, Flicek, P., Walenz, B.P., Kwak, W., Clawson, H., Diekhans, M., Nassar,
L., Paten, B., Kraus, R.H.S., Crawford, A.J., Gilbert, M.T.P., Zhang, G.,
Venkatesh, B., Murphy, R.W., Koepfli, K.-P., Shapiro, B., Johnson, W.E., Di
Palma, F., Marques-Bonet, T., Teeling, E.C., Warnow, T., Graves, J.M., Ryder,
O.A., Haussler, D., O'Brien, S.J., Korlach, J., Lewin, H.A., Howe, Kerstin,
Myers, E.W., Durbin, R., Phillippy, A.M., Jarvis, E.D., 2021. Towards complete
and error-free genome assemblies of all vertebrate species. Nature 592, 737–
746. https://doi.org/10.1038/s41586-021-03451-0

Rhie, A., Walenz, B.P., Koren, S., Phillippy, A.M., 2020. Merqury: reference-free
quality, completeness, and phasing assessment for genome assemblies.
Genome Biol 21, 245. https://doi.org/10.1186/s13059-020-02134-9

Ryu, S.H., Kwak, M.J., Hwang, U.W., 2013. Complete mitochondrial genome of the
Eurasian flying squirrel Pteromys volans (Sciuromorpha, Sciuridae) and
revision of rodent phylogeny. Mol Biol Rep 40, 1917–1926.
https://doi.org/10.1007/s11033-012-2248-x

Santangeli, A., Wistbacka, R., Hanski, I.K., Laaksonen, T., 2013. Ineffective enforced
legislation for nature conservation: A case study with Siberian flying squirrel
and forestry in a boreal landscape. Biological Conservation 157, 237–244.
https://doi.org/10.1016/j.biocon.2012.09.012

Sayyari, E., Whitfield, J.B., Mirarab, S., 2018. DiscoVista: Interpretable visualizations
of gene tree discordance. Molecular Phylogenetics and Evolution 122, 110–
115. https://doi.org/10.1016/j.ympev.2018.01.019

Selonen, V., Mäkeläinen, S., 2017. Ecology and protection of a flagship species, the
Siberian flying squirrel. Hystrix, the Italian Journal of Mammalogy 28.
https://doi.org/10.4404/hystrix-28.2-12328

Selonen, V., Wistbacka, R., 2016. Siberian flying squirrels do not anticipate future
resource abundance. BMC Ecol 16, 51. https://doi.org/10.1186/s12898-016-
0107-7

Shar, S., Lkhagvasuren, D., Henttonen, H., Maran, T., Hanski, I.K., 2016. Pteromys
volans (errata version published in 2017). The IUCN Red List of Threatened
Species 2016: e.T18702A115144995. https://doi.org/10.2305/IUCN.UK.2016-
3.RLTS.T18702A22270935.en

Simpson, J.T., Wong, K., Jackman, S.D., Schein, J.E., Jones, S.J.M., Birol, İ., 2009.
ABySS: A parallel assembler for short read sequence data. Genome Res. 19,
1117–1123. https://doi.org/10.1101/gr.089532.108

Smit, A.F.A., Hubley, R., Green, P., 2023. RepeatMasker.

Tarasov, A., Vilella, A.J., Cuppen, E., Nijman, I.J., Prins, P., 2015. Sambamba: fast
processing of NGS alignment formats. Bioinformatics 31, 2032–2034.
https://doi.org/10.1093/bioinformatics/btv098
The UniProt Consortium, 2019. UniProt: a worldwide hub of protein knowledge.
Nucleic Acids Research 47, D506–D515. https://doi.org/10.1093/nar/gky1049
Thorington, R.W., Koprowski, J.L., Steele, M.A., Whatton, J.F. (Eds.), 2012. Squirrels
of the world. Johns Hopkins University Press, Baltimore.
Tillich, M., Lehwark, P., Pellizzer, T., Ulbricht-Jones, E.S., Fischer, A., Bock, R.,
Greiner, S., 2017. GeSeq – versatile and accurate annotation of organelle
genomes. Nucleic Acids Research 45, W6–W11.
https://doi.org/10.1093/nar/gkx391
Törmälä, T., Vuorinen, H., Hokkanen, H., 1980. Timing of circadian activity in the
flying squirrel in central Finland. Acta Theriologica 25, 461–474.
Uliano-Silva, M., Ferreira, J.G.R.N., Krasheninnikova, K., Darwin Tree of Life
Consortium, Blaxter, M., Mieszkowska, N., Hall, N., Holland, P., Durbin, R.,
Richards, T., Kersey, P., Hollingsworth, P., Wilson, W., Twyford, A., Gaya, E.,
Lawniczak, M., Lewis, O., Broad, G., Martin, F., Hart, M., Barnes, I., Formenti,
G., Abueg, L., Torrance, J., Myers, E.W., Durbin, R., Blaxter, M., McCarthy,
S.A., 2023. MitoHiFi: a python pipeline for mitochondrial genome assembly
from PacBio high fidelity reads. BMC Bioinformatics 24, 288.
https://doi.org/10.1186/s12859-023-05385-y
Ungerer, M.C., Johnson, L.C., Herman, M.A., 2008. Ecological genomics:
understanding gene and genome function in the natural environment. Heredity
100, 178–183. https://doi.org/10.1038/sj.hdy.6800992
Wistbacka, R., Orell, M., Santangeli, A., 2018. The tragedy of the science-policy gap
– Revised legislation fails to protect an endangered species in a managed
boreal landscape. Forest Ecology and Management 422, 172–178.
https://doi.org/10.1016/j.foreco.2018.04.017
Wolf, J.F., Bowman, J., Keobouasone, S., Taylor, R.S., Wilson, P.J., 2022. A *de novo*
genome assembly and annotation of the southern flying squirrel (*Glaucomys*
volans). G3 Genes|Genomes|Genetics 12, jkab373.
https://doi.org/10.1093/g3journal/jkab373
Xu, M., Guo, L., Gu, S., Wang, O., Zhang, R., Peters, B.A., Fan, G., Liu, X., Xu, X.,
Deng, L., Zhang, Y., 2020. TGS-GapCloser: A fast and accurate gap closer for
large genomes with low coverage of error-prone long reads. GigaScience 9,
giaa094. https://doi.org/10.1093/gigascience/giaa094
Zagorodniuk, I., 2022. The Siberian flying squirrel (Pteromys volans) in south of
Eastern Europe: distribution boundaries and its changes. Theriol. Ukr. 2022,
66–77. https://doi.org/10.15407/TU2308
Zelditch, M.L., Li, J., Tran, L.A.P., Swiderski, D.L., 2015. Relationships of diversity,
disparity, and their evolutionary rates in squirrels (Sciuridae): DIVERSITY AND
DISPARITY IN SQUIRRELS. Evolution 69, 1284–1300.
https://doi.org/10.1111/evo.12642
Zhang, C., Rabiee, M., Sayyari, E., Mirarab, S., 2018. ASTRAL-III: polynomial time
species tree reconstruction from partially resolved gene trees. BMC
Bioinformatics 19, 153. https://doi.org/10.1186/s12859-018-2129-y
Zhou, C., McCarthy, S.A., Durbin, R., 2023. YaHS: yet another Hi-C scaffolding tool.
Bioinformatics 39, btac808. https://doi.org/10.1093/bioinformatics/btac808
729

Table 1. Assembly statistics table on the scaffold-level as well was on the contig-level. (K)bp = (kilo) base pairs; #

= number; GC content = Guanine-Cytosine content; auN = area under the Nx curve. A more detailed assembly

statistics table can be found in the Supplementary Information (Supplementary Table S1).

Genome statistic	scaffold-level	contig-level
total length	2,848,339,732 bp	2,848,327,632 bp
# of sequences	638	759
# sequences (≥ 0 bp)	638	-
# sequences (≥ 1000 bp)	638	758
Total length (≥ 0 bp)	2,848,339,732 bp	-
Total length (≥ 1000 bp)	2,848,339,732 bp	2,848,326,863 bp
Largest sequence	214,577,690 bp	136,295,669 bp
GC level	40.77%	40.77%
N50	157,387,023 bp	60,698,384 bp
N90	42,248,379 bp	3,984,759 bp
auN	147,030,830.8 bp	62,127,644.0 bp
L50	8	16
L90	18	73
# N's per 100 kbp	0.42	0.00

733 **Table 1.** Busco gene scores generated with *compeasm* and *BUSCO* both based on the total number of 9,226
734 expected orthologs.

software	*compeasm*		*BUSCO*	
unit	*n*	*%*	*n*	*%*
Complete buscos (C = S + D)	9,217	99,90%	8,947	97%
Complete and single-copy buscos (S)	9,145	99.12%	8,763	95%
Complete and duplicated buscos (D)	72	0.78%	184	2%
Fragmented buscos, subclass 1 (F)	6	0.07%	51	0.6%
Fragmented buscos, subclass 2 (I)	0	0%	-	-
Missing buscos (M)	3	0.03%	228	2.4%

737 **Table 2.** Assembly assignment table estimated by *Repeatmodeler* and *Repeatmasker*. Most repeats fragmented
738 by insertions or deletions have been counted as one element (marked with an asterisk).

Type of elements	number of elements [*n*]*	length occupied [bp]	percentage of sequence [%]
Retroelements	3,255,026	886,574,991 bp	31.13%
SINEs:	1,483,628	169,453,665 bp	5.95%
Penelope:	62	11,402 bp	0.00%
LINEs:	878,225	522,064,913 bp	18.33%
CRE/SLACS	0	0 bp	0.00%
L2/CR1/Rex	86,215	23,004,669 bp	0.81%
R1/LOA/Jockey	0	0 bp	0.00%
R2/R4/NeSL	246	56,708 bp	0.00%
RTE/Bov-B	5,179	1,231,196 bp	0.04%
L1/CIN4	786,498	497,749,583 bp	17.48%
LTR elements:	893,173	195,056,413 bp	6.85%
BEL/Pao	0	0 bp	0.00%
Ty1/Copia	0	0 bp	0.00%
Gypsy/DIRS1	6,436	1,386,734 bp	0.05%
Retroviral	869,037	191,862,047 bp	6.74%
DNA transposons	224,420	46,280,608 bp	1.62%
hobo-Activator	163,126	32,188,953 bp	1.13%
Tc1-IS630-Pogo	55,186	13,103,816 bp	0.46%
En-Spm	0	0 bp	0.00%
MULE-MuDR	251	41,323 bp	0.00%
PiggyBac	521	160,128 bp	0.01%
Tourist/Harbinger	323	50,596 bp	0.00%
Other (Mirage, P-element, Transib)	0	0 bp	0.00%
Rolling-circles	879	195,645 bp	0.01%
Unclassified:	59,407	3,502,033 bp	0.12%
Total interspersed repeats:		936,369,034 bp	32.87%
Small RNA:	50,111	3,274,868 bp	0.11%
Satellites:	49,554	2,663,208 bp	0.09%
Simple repeats:	530,473	22,897,566 bp	0.80%
Low complexity:	106,166	0 bp	0.19%

Figure 1. (a) Adult specimen of *Pteromys volans volans* in Lapua, Finland. **(b)** Omni-C contact density map depicting the 19 distinct chromosome-level scaffolds. **(c)** Circos plots comparing the synteny of the chromosome-scale genome assembly of *Pteromys volans* (Uoulu_pteVol_1.0) with chromosome-scale assemblies of a Eurasian red squirrel (*Sciurus vulgaris*), an eastern grey squirrel (*Sciurus carolinensis*), and a European ground squirrel (*Spermophilus citellus*). All three synteny plots are available in a higher resolution in the supplementary information (Supplementary Figures S1-S3).

Figure 2. (a) Circos plot of the chromosome-level genome assembly of *Pteromys volans* (Uoulu_pteVol_1.0). The tracks show the following information for 19 scaffolds (length in Gbp) : Red [A]: Position of the telomeres; red asterisks indicate telomeric regions with length of > 100 copies measured with *quarTeT*. Turquoise [B]: Gene content within a 500,000 base pair windows. Blue [C]: Simple repeats. The length of the bars indicates the relative length of the simple repeats after clustering. Orange [D]: Transposable elements (TE) clustered by type. The length of the bars indicates the relative length of the TE after clustering. Brown [E]: GC content. **(b)** Circular gene map of the mitogenome of *Pteromys volans*. The different functional genes groups are shown in different colours, which are shown on the bottom right. The genes transcribed in clockwise and counterclockwise (indicated by arrows) are shown inside and outside of the external circle, respectively. The dark grey colour of inner circle shows the GC content. Figures 2a and 2b as well as the telomere overview plot from *quarTeT* can be found in the Supplementary Information (Supplementary Figures S4 – S6).

Figure 3. Coalescent phylogenetic tree from 17 rodent species based on 3,269 busco genes each with the assigned busco completeness [%]. Species with available chromosome-level genome assemblies were marked with an asterisk (*). The novel chromosome-level genome assembly from *Pteromys volans* is indicated with two asterisks (**). The outgroup and the according higher taxa are assigned on the right-hand side: families (darkest blue) as well as for the squirrel subfamilies (medium blue) and tribes (lightest blue). The branch quartet frequencies graphs generated by *DiscoVista* can be found in the Supplementary Information (Supplementary Figure S7).

Data Accessibility Statement

All underlying read data and the assembly are available at NCBI GenBank under BioProject PRJNA1141127. The hard-masked genome assembly and gene annotations are available at Dryad (https://doi.org/10.5061/dryad.3xsj3txth). More detailed results are available in the Supplementary Information.

Competing Interests Statement

The authors declare no competing interests.

Author Contributions

Gerrit Wehrenberg: Conceptualisation; Data Curation, Formal Analysis; Methodology; Visualisation; Writing – Original Draft Preparation (lead); Writing – Review & Editing (lead). **Angelika Kiebler**: Formal Analysis; Methodology; Visualisation; Writing – Original Draft Preparation; Writing – Review & Editing. **Carola Greve**: Formal Analysis; Methodology; Writing – Review & Editing. **Núria Beltrán-Sanz**: Data Curation, Formal Analysis; Methodology; Visualisation; Writing – Review & Editing. **Alexander Ben Hamadou**: Methodology. **René Meißner**: Formal Analysis; Methodology; Writing – Review & Editing. **Sven Winter**: Conceptualisation; Data Curation, Formal Analysis; Methodology; Supervision; Visualisation; Writing – Review & Editing. **Stefan Prost**: Conceptualisation; Data Curation, Formal Analysis; Funding Acquisition; Methodology; Project Administration; Supervision; Visualisation; Writing – Original Draft Preparation; Writing – Review & Editing.

Acknowledgments

G.W. and S.P. received funding from Biodiverse Anthropocenes (ANTS) through University of Oulu and the Research Council of Finland PROFI6 funding. We also thank Timo Perätie from Kuopion kaupungin alueelliset ympäristönsuojelupalvelut and the Zoological Museum of the University of Oulu, especially Mikko Vallinmäki, for sampling, sample provision, sample storage and database management. We thank Charlotte Gerheim of the LOEWE Translational Biodiversity Genomics (TBG) project based in the Senckenberg Research Institute and Museum, Frankfurt, for support with lab work. The authors wish to acknowledge CSC – IT Center for Science,

Finland, for computational resources. We also thank the Genome Technology Center
(RGTC) at Radboud University Medical Center (Radboudumc; Nijmegen, The
Netherlands) and the Bioscientia Institut für Medizinische Diagnostik GmbH
(Ingelheim, Germany) for providing the PacBio SMRT sequencing service on the
PacBio Revio platform. Yannis Schöneberg for helping with the data analysis. We are
grateful for the permission to use a wonderful photograph of a Finnish *Pteromys*
volans volans by Pekka Takamaa, as well as the scientific illustrations of various
rodent species provided by ALADA BOOKS, S.L. Kiitos paljon to Professor Marko
Mutanen for the abstract translation into the Finnish language.