ProSynTaxDB: A curated protein database and workflow for taxonomic classification of *Prochlorococcus* and *Synechococcus* in metagenomes

Allison Coe[1,*], James I. Mullet[1,*], Nhi N. Vo[1,*], Paul M. Berube[1], Maya I. Anjur-Dietrich[1], Eli Salcedo[1], Sierra M. Parker[1], Konnor VonEmster[1], Christina Bliem[1], Aldo A. Arellano[1], Kurt G. Castro[1], Jamie W. Becker[1], Sallie W. Chisholm[1,2]

1- Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA

2- Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, USA

*- These authors contributed equally to this work.

ABSTRACT

Prochlorococcus and *Synechococcus* are abundant marine picocyanobacteria that contribute significantly to ocean primary production. Recent genome sequencing efforts, including those presented here, have yielded a large number of high-quality reference genomes, enabling the classification of these picocyanobacteria in marine metagenomic sequence data at high phylogenetic resolution. When combined with environmental data, these classifications can guide cluster/clade/grade assignments and offer insights into niche differentiation within these populations. Here we present ProSynTaxDB, a curated protein sequence database and accompanying workflow aimed at enhancing the taxonomic resolution of *Prochlorococcus* and *Synechococcus* classification. ProSynTaxDB includes proteins from 1,260 genomes of *Prochlorococcus* and *Synechococcus*, including single-amplified genomes, high-quality draft genomes, and newly closed genomes. Additionally, ProSynTaxDB incorporates proteins from 27,799 genomes of marine heterotrophic bacteria, archaea, and viruses to assess microbial and viral communities surrounding *Prochlorococcus* and *Synechococcus*. This resource enables accurate classification of picocyanobacterial clusters/clades/grades in metagenomic data – even when present at 0.60% of reads for *Prochlorococcus* or 0.09% of reads for *Synechococcus*.

BACKGROUND / SUMMARY

Prochlorococcus and *Synechococcus* are oxygenic photosynthetic bacteria that are key microbes at the base of the marine microbial food web. Together, they have colonized much of the surface ocean[1], with *Synechococcus* having a broad geographic range and *Prochlorococcus* dominating the tropical and subtropical open ocean. *Prochlorococcus* and *Synechococcus* are part of a diverse community of marine phytoplankton, which have a key role in regulating the transformation of energy and matter in the oceans. Studies of marine ecosystems have generated large metagenomic datasets[2–5], which are a rich resource for investigating the distributions of microorganisms in the marine environment[6]. Many classification tools have been developed to process growing volumes of sequence data. Some use broad reference databases (e.g., NCBI and GTDB), while others are specialized to specific groups of microbes[7,8]. Accurate species identification is challenging due to the high diversity of natural samples and the limited availability of taxonomically classified genomes, particularly for distinguishing between very closely related organisms.

Determining the distributions of the picocyanobacteria *Prochlorococcus* and *Synechococcus* is of particular importance for understanding marine ecosystems. These organisms are exceptionally diverse[9–13], thus making the accurate differentiation of clusters/clades/grades within each genus difficult. As a result, *Prochlorococcus and Synechococcus* populations are often classified as a single strain[14], obscuring the finely-tuned niche partitioning that is well documented in these groups[15–20]. Over the past decade, a substantial amount of high-quality genomic reference data has become available to address these gaps. These data enable the linkage between the functional and taxonomic structure of *Prochlorococcus* and *Synechococcus* populations and the features of the environment in which they are embedded. They also include genomes from uncultivated single cells, dwarfing the number of available reference genomes from cultured cells by an order of magnitude[21–23]. Single-cell genomes have substantially expanded our understanding of picocyanobacterial diversity while mitigating biases inherent in cultivation-dependent methods, which tend to select for strains adaptable to existing isolation protocols.

Here, we leveraged these extensive datasets to present a comprehensive protein database – ProSynTaxDB – that has been manually curated to enable finely resolved taxonomic classification of *Prochlorococcus* and *Synechococcus*, along with the microbial and viral communities associated with these picocyanobacteria in short-read metagenomic datasets. The curated taxonomy is based on the core protein similarity of individual genomes as well as decades of research on niche partitioning and the diverse ecology and physiology of different phylogenetic branches of *Synechococcus* and *Prochlorococcus*[12,13,15,17]. Accompanying this protein database is a new workflow that links high-throughput quality control, taxonomic classification, and count normalization for short-read sequences[24,25]. Custom scripts are included for reporting both the absolute and normalized read abundances (the latter referred to as genome equivalents) of *Prochlorococcus* and *Synechococcus* at user-defined taxonomic levels in metagenomic samples, ranging from broad groupings of high-light and low-light adapted *Prochlorococcus* to more fine-grained distinctions between individual clades/grades. The workflow can accurately identify the major clusters/clades/grades/ecotypes of these picocyanobacteria in short-read sequence libraries consisting of a minimum of 1 million reads in which *Prochlorococcus* represents at least 0.60% of reads, and *Synechococcus* at least 0.09% of reads. *ProSynTaxDB* contains new closed circular genomes of 40 *Prochlorococcus,* 12 *Synechococcus*, and 10 marine heterotrophic bacterial strains that were co-isolated from *Prochlorococcus* cultures, including 21 previously partially assembled and 16 unpublished isolate *Prochlorococcus* genomes. Furthermore, ProSynTaxDB includes proteins from 27,799 genomes of marine heterotrophic bacteria, archaea, and viruses to analyze the microbial and viral communities associated with *Prochlorococcus* and *Synechococcus*. The closed genomes reported here will be particularly valuable for studies of microbial evolution that rely on fully assembled genomic islands and regions of the genome that are subject to remodeling.

METHODS
Enrichment and isolation of picocyanobacteria

Strains were isolated from seawater collected from 3 cruises in the North Pacific Subtropical Gyre and 1 cruise in the South Pacific Subtropical Gyre. Ten low-light (LL) adapted *Prochlorococcus* strains representing the LLI and LLIV clades and 4 heterotrophic bacteria representing classes Gammaproteobacteria and Alphaproteobacteria were isolated and sequenced (Fig. 1, Table 1). These 4 marine heterotrophic bacteria were co-isolated from *Prochlorococcus* cultures. *Synechococcus* strain Cu2B8 was also isolated and sequenced; however, records of its isolation location and depth are not available (Table. 1).

All *Prochlorococcus* strains were isolated from raw seawater, except MIT0913, MIT1227, and MIT1418 which were isolated from the filtrate of seawater put through a 1 or 1.2 μm filter. Different media formulations were used, with specific nutrient amendments aimed at selecting for various clades of *Prochlorococcus*, as described hereafter. Isolates MIT0912, MIT0913, and MIT0916 were amended with Pro2 nutrients[26], replacing the nitrogen source with 50 μM nitrate. Isolates MIT1011, MIT1012, and MIT1013 were isolated on media containing 20 μM ammonium chloride, 1 μM sodium phosphate, 0.1x Pro99 trace metal mix[26], and 1 mM sodium bicarbonate. Isolates MIT1201 and MIT1205 were isolated on media containing 16 μM ammonium chloride, 1 μM sodium phosphate, and 0.1x Pro99 trace metal mix. MIT1227 was isolated on media containing 32 μM ammonium chloride, 2 μM sodium phosphate, and 0.1x Pro99 trace metal mix. Finally, MIT1418 was isolated on media containing 0.1x Pro2 nutrient and metal concentrations. The chemicals used in media preparation were of the highest purity (i.e. Sigma BioUltra) to prevent contaminants that could affect culture growth.

Heterotrophic bacteria MIT1350, MIT1358, MIT1388, and MIT1392 were isolated on 0.3% agar pour plates consisting of Pro99 medium amended with 0.05% pyruvate and 3.75mM TAPS buffer. Colonies were picked and subcultured three times before being transferred into liquid Pro99 medium. MIT1350 was sequenced before freezing, while the remaining strains were directly frozen. Upon thawing MIT1358, MIT1388, and MIT1392 for this study, the cells were repurified using the same methods, with the exception that colonies were transferred into liquid ProMM medium[27] before pelleting. Purity was then assessed through genomic sequencing to verify the absence of contaminants. Heterotrophic bacteria strain MIT0238 was isolated from xenic *Prochlorococcus* MIT9313; however, the records detailing this isolation are not available.

Table 1. Culture isolates of *Prochlorococcus*, *Synechococcus*, and marine heterotrophic bacteria. Clade and cluster designations for *Prochlorococcus* and *Synechococcus* strains are provided, along with isolation location, depth, year, and cruise identifier.

Strain ID	Genus	Cluster/Clade	Cruise ID	Year Isolated	Coordinates	Depth (m)
MIT0912	*Prochlorococcus*	LLI	KM0915	2009	22.75°N 158°W	175
MIT0913	*Prochlorococcus*	LLI	KM0915	2009	22.75°N 158°W	175
MIT0916	*Prochlorococcus*	LLI	KM0915	2009	22.75°N 158°W	200
MIT1011	*Prochlorococcus*	LLI	MV1015	2010	26.25°S 103.96°W	150
MIT1012	*Prochlorococcus*	LLI	MV1015	2010	26.25°S 103.96°W	150
MIT1013	*Prochlorococcus*	LLI	MV1015	2010	26.25°S 103.96°W	150
MIT1201	*Prochlorococcus*	LLIV	KM1217	2012	22.75°N 158°W	124
MIT1205	*Prochlorococcus*	LLIV	KM1217	2012	22.75°N 158°W	175
MIT1227	*Prochlorococcus*	LLIV	KM1217	2012	22.75°N 158°W	124
MIT1418	*Prochlorococcus*	LLIV	KOK1404	2014	22.75°N 158°W	150
Cu2B8	*Synechococcus*	5.1 B - I	n/a	n/a	n/a	n/a
MIT0238	*Marinobacter*	-	n/a	2002	37.51°N 68.24°W	135
MIT1392	*Marinobacter*	-	KM1309	2013	22.75°N 158°W	150
MIT1358	*Marinobacter*	-	KM1309	2013	22.75°N 158°W	150
MIT1388	*Shimia*	-	KM1309	2013	22.75°N 158°W	150

Culture conditions and biomass preparation

Cultures were grown in 200 mL of Pro99 medium[26] at 24°C using continuous light or a 13:11 light-dark cycle at light intensities ranging from 10 to 50 µmol photons m^{-2} s^{-1}. Cells were harvested for sequencing during late exponential growth phase, with 0.003% Pluronic F-68 Polyol (MP Biomedicals, Cat# 2750049) added to promote cell aggregation and improve biomass yields during centrifugation. Cells were centrifuged at 7,197 x g for 15-20 min at room temperature. After centrifugation, the supernatant was decanted, and the resulting pellets were flash frozen in liquid nitrogen and stored at -80°C.

Extraction and sequencing

Genomic DNA was extracted from the frozen cell pellets using a modified version of a phenol:chloroform based extraction method[28] in order to obtain high molecular weight DNA suitable for sequencing on Pacific Biosciences (PacBio) platforms. Modifications included minimizing agitation to prevent DNA shear (e.g., gently shaking the sample tube by hand instead of vortexing), adding 1µl of 50 mg/mL lysozyme (cat# 90082, Thermo Scientific) to the cell pellet during the digestion step, performing a second chloroform/isoamyl alcohol step to remove residual phenol, and increasing the salt concentration by the addition of 0.1 volume of 3M sodium acetate or 3M ammonium acetate before precipitation with 0.6-0.8 volumes of isopropanol. The tube was then gently inverted multiple times and incubated at room temperature for 30 min. DNA was recovered by centrifugation at 10,000 x g for 3 min at 4°C. The supernatant was then carefully removed by pipetting. The pellet was then washed twice with 75% ethanol before allowing the pellet to dry at room temperature for approximately 10 min inside a biological safety hood to prevent contamination. It was then dissolved in 1x TE buffer (pH 8) and stored at -80°C.

Genomic DNA samples were diluted and fragmented to 10-12Kb using a gTube (cat# 520079, Covaris), followed by a 0.45X SPRIselect (cat# B23317, Beckman Coulter Life Sciences) bead cleanup. The samples were then made into indexed SMRTBell libraries following guidelines for the template prep kit v 1.0, 2.0, or 3.0 (cat# 100-259-100, 100-938-900, 102-141-700, respectively, Pacific Biosciences). These libraries were assembled into a single pool, and cleaned using MinElute reaction cleanup kit (cat# 28204, Qiagen), followed by additional cleaning using 0.4X SPRIselect beads. The libraries were then bound using Sequel Binding Kit v 2.1, Sequel II Binding Kit v 2.1, and Sequel II binding Kit v 3.2 (cat#100-369-800, 101-820-500, 102-194-100, respectively, Pacific Biosciences) and sequenced on PacBio Sequel I, Sequel II, or Sequel IIe flowcells at either the MIT BioMicro Center or Harvard Bauer core facilities.

Genome assembly

PacBio sequence data were demultiplexed by barcode using PacBio SMRTtools, generating both circular consensus sequencing reads and subreads. To maximize the likelihood of closing genomes, reads were assembled using multiple tools, including Flye v2.9.5, Canu v2.3, and SPAdes v4.1.0[29–31]. Contigs were classified using MMseqs2 v17.b804f against the GTDB v207 genome database[32,33]. The resulting genomes were submitted to NCBI and IMG, with accessions provided in ProSynTaxDB_genomes.tsv[34]. The GitHub repository, which includes documentation and code for genome assembly, is provided in the Code Availability section.

In addition to closing genomes of cultured strains, six heterotrophic bacterial genomes were successfully assembled from sequencing xenic cultures of *Prochlorococcus* and *Synechococcus*. However, since pure axenic isolates for these heterotroph strains are unavailable, we implemented a new nomenclature by appending an additional two-digit identifier ("01") to the picocyanobacterial strain name from which they were derived. For example, *Thalassospira* MIT121401 was obtained from sequencing a xenic culture of *Prochlorococcus* MIT1214.

Database construction

The database consists of single-amplified genomes, finished genomes of culture isolates, and high-quality draft genomes of cultured isolates. These genomes were obtained from several sources, including IMG-ProPortal[21], GORG-Tropics[35], NCBI RefSeq (February 2024), Cyanorak[36], and this study[34]. No metagenome-assembled genomes were included in the database to ensure only high-quality reference genomes were present. Additionally, we filtered the NCBI RefSeq database to only include only complete bacterial and archaeal genomes, selecting those with at least one reference from each taxon found in a marine environment to ensure the inclusion of only marine genomes. For any genomes without annotated protein sequences, prodigal v2.6.3[37] was used to identify open reading frames. Stop codons were removed from these predicted protein sequences to remain consistent across all genomes. All protein sequences were concatenated, and Kaiju v1.10.1 was used to construct the finalized database (kaiju-mkbwt and kaiju-mkfmi with default parameters)[24].

Picocyanobacteria taxonomy

The taxonomy of *Prochlorococcus* and *Synechococcus* was manually curated based on a concatenated multiple-sequence alignment of proteins encoded by any single-copy core genes annotated in each genome. Single-copy core genes were defined as those present in exactly one copy (i.e., genes with multiple copies within any genome were removed) in 100% of 92 genomes derived from cultures in the CyCOG v6.0 database[21] and estimated to be >99% complete using checkM[38]. All proteins encoded by *Prochlorococcus* and *Synechococcus* genomes were aligned against proteins in the CyCOG v6.0 database (blastp v2.16.0 -evalue 0.001), selecting the best hit for each gene. Following the annotation of all genomes using the CyCOG v6.0 database, protein sequences for the 424 single-copy core genes were aligned using ClustalOmega v1.2.4[39], concatenated with MEGA v11.0.13[40], and used to construct a phylogenetic tree with FastTreeMP v2.1.11 (-lg -boot 100)[41]. Genomes missing more than 95% data among positions with <50% gaps or genomes with less than 5 single-copy core genes were removed from the database and phylogenetic tree. The classified genomes were then analyzed using CheckM v1.2.3[38] to determine the completeness and contamination for each genome.

Figure 1. Phylogenetic tree of *Prochlorococcus* and *Synechococcus* included in ProSynTaxDB. The phylogeny includes 1,106 *Prochlorococcus* genomes and 154 *Synechococcus* genomes and is based on a concatenated protein alignment of the 424 single-copy core genes (see Materials and Methods). The tree is arbitrarily rooted at the *Synechococcus* cluster. The outer layer (colors) depicts the major cluster, clade, or grade defined in ProSynTaxDB for each genome. The middle layer (shades of gray) contains the associated genus for each genome, and the inner layer (black bars) highlights closed genomes that are presented for the first time in this work.

ProSynTaxDB Workflow
The ProSynTaxDB workflow assigns taxonomic classifications to metagenomic reads based on the taxonomic hierarchy in ProSynTaxDB and uses these classified reads to estimate the

genome equivalents of *Prochlorococcus* and *Synechococcus* clusters, clades, or grades within a metagenome sample. The workflow begins by trimming low-quality regions and removing adapter sequences from FASTQ paired-end read files using BBDuk v39.18 (bbduk -minlen 25 -trimq=10 maq=20 ktrim=r k=23 mink=11 hdist=1) from the BBTools suite[42]. Reads are then classified against the ProSynTaxDB using Kaiju v1.10.1 (-m 11 -s 65 -E 0.05 -e 5)[24]. A custom Python script[34] then summarizes raw read counts for each genus, and reads identified as *Prochlorococcus* and *Synechococcus* are extracted into separate FASTA files using Seqtk vr82[43]. To account for variations in average genome length across *Prochlorococcus* and *Synechococcus* clusters/clades/grades, raw read counts are aligned to single-copy core genes (SCCG) from the CyCOG v6.0 database[21] using DIAMOND Blastx v2.1.11[44]. The genome equivalents for each cluster/clade/grade are then estimated by normalizing the total SCCG read residues to the average SCCG length. The codes and documentation for the ProSynTaxDB workflow have been deposited on GitHub (see Code Availability).

Figure 2. ProSynTaxDB workflow for classifying metagenomic reads. The workflow consists of four stages: (1) quality filtering and adapter trimming using BBDuk (Bushnell et al., 2017); (2) classification of reads utilizing ProSynTaxDB[34] and Kaiju[24]; (3) summarization of raw read counts per genus (including unclassified reads); (4) binning of *Prochlorococcus* and *Synechococcus* reads using Seqtk[43], aligning the binned reads to protein sequences encoded by single-copy genes found in CyCOG6[21] using DIAMOND Blastx[44], and normalization calculations to obtain genome equivalents using a custom Python script[34].

DATA RECORDS

ProSynTaxDB data files are accessible online through a Zenodo repository[34], which includes the following files:

ProSynTaxDB_genomes.tsv

Table of genomes included in the ProSynTaxDB and their associated metadata. Data fields are as follows:

organism: The name of the organism recorded in NCBI when available. For genomes/organisms obtained from sources other than NCBI, the organism name is provided in NCBI format

genome_short_name: The genome name used in the ProSynTaxDB

domain: Bacteria, Archaea, Eukarya, or Virus

genus: The genus of the organism in NCBI

clade: The major cluster/clade/grade of *Prochlorococcus* or *Synechococcus* based on phylogenetic reconstruction using a concatenated alignment of proteins encoded by single-copy core genes

NCBI_BioProject: The NCBI BioProject accession number associated with the organism, when available

NCBI_BioSample: The NCBI BioSample accession number associated with the organism, when available

NCBI_GenBank: The NCBI GenBank accession number associated with the genome sequence data, when available

IMG_Genome_ID: The IMG Genome ID accession number, also known as the IMG Taxon ID, corresponds to the genome or organism in the Joint Genome Institute's (JGI) Integrated Microbial Genomes (IMG) repository, when available

ProSynTaxDB_names.dmp

Names taxonomy file for use with the ProSynTaxDB

ProSynTaxDB_nodes.dmp

Nodes taxonomy file for use with the ProSynTaxDB

ProSynTaxDB.fmi

Index file containing contents of ProSynTaxDB_v1.faa for use with ProSynTaxDB

CyCOG6.dmnd

Database containing orthologous groups of proteins used in the cluster/clade/grade normalization step

ProSynTaxDB_v1.faa
File containing protein sequences used by Kaiju for classification of reads. Each protein sequence contains a header starting with ">"

average_cycog_length.csv
Comma separated file containing the average length for each protein sequence used in the normalization step. Data fields are as follows:

cycog: name for single-copy core gene

mean_AA_length: the average length of amino acids in the protein sequence of the gene

ProSynTaxDB-workflow_benchmarking_genomes.tsv
This tab-delimited file contains a list of subsetted genomes used in each benchmarking experiment done in the Technical Validation section. Data fields are as follows:

Experiment Name: name of benchmarking experiment conducted

Subset ID: unique ID from the random genome subsetting

Genome Name: name of genome used in benchmarking experiment

ProSynTaxDB-workflow_benchmarking_composition.tsv
This tab-delimited file contains the taxon composition of all samples used in each benchmarking experiment done in the Technical Validation section. Data fields are as follows:

Experiment Name: name of benchmarking experiment conducted

Sample Name: unique sample name

Percent *Prochlorococcus*: percent of reads in simulated sample originating from *Prochlorococcus* genomes

Percent *Synechococcus*: percent of reads in simulated sample originating from *Synechococcus* genomes

Percent Heterotroph: percent of reads in simulated sample originating from marine heterotrophic bacterial genomes

Notes: additional information about the simulated sample

TECHNICAL VALIDATION

Generation of mock metagenome data

To assess the accuracy of cluster/clade/grade-level classification of metagenomic reads using ProSynTaxDB, we generated mock metagenome sequence datasets with known genome inputs. The mock dataset was generated using a custom script[34] to randomly select 20% of genomes from each *Prochlorococcus* and *Synechococcus* cluster/clade/grade and 20% of marine heterotrophs from the reference genome database for read simulation. These genomes were used as input for mason2 v2.0.0-beta1[45] to create simulated mock metagenomes with 1 million paired-end 150 nt sequences. The fasta file was modified to exclude sequences associated with the genomes used in the mock metagenomes. The .fmi file was then rebuilt using the subsetted .faa file using the Kaiju functionalities kaiju-mkbwt and kaiju-mkfmi[24]. This ensures that genomes used to create mock metagenomes were excluded from the set used for classification. Genomes were randomly subsetted 10 times for each technical validation experiment that consisted of a paired mock metagenome and a subsetted taxonomic database[34].

Assessment of misclassification rates

We evaluated the accuracy of the read classification workflow using the simulated dataset derived from a subset of the reference database (described above). Reads for each group were simulated separately, processed through the classification workflow, and evaluated based on the proportion classified as true positives or misclassified as false positives. The average results across 10 subsets are presented in Table 2.

Table 2. Misclassification rates of simulated metagenomic reads for *Prochlorococcus, Synechococcus,* and marine heterotrophic bacteria. Data represents the average and standard deviation from 10 random genome subsets. Grey rows represent true positive rates, while white rows represent false positive rates for each taxonomic group.

		Read Group		
		Prochlorococcus	*Synechococcus*	Heterotroph
Classified As	*Prochlorococcus*	84.56 ± 0.48	1.77 ± 0.20	0.025 ± 0.002
	Synechococcus	0.81 ± 0.09	76.79 ± 1.13	0.004 ± 0.002
	Heterotroph	14.63 ± 0.41	21.45 ± 0.95	99.97 ± 0.003

Limit of Detection

To determine the minimum detectable abundance of *Prochlorococcus* in a sample, we simulated datasets with varying proportions of *Prochlorococcus* and heterotrophs and

processed them through the classification workflow. The simulated heterotroph proportions are available on Zenodo (ProSynTaxDB-workflow_benchmarking_composition.tsv)[34]. The *Prochlorococcus* false positive rate was calculated by dividing the number of heterotroph reads misclassified as *Prochlorococcus* by the total reads classified as *Prochlorococcus*. To maintain a misclassification rate below 10%, *Prochlorococcus* abundance must exceed 0.28% of all classified reads (Fig. 3 A). Using the same approach, *Synechococcus* requires an abundance greater than 0.04% to keep the misclassification rate below 10% (Fig. 3 C). For more sensitive cluster/clade/grade-level identification, we recommend a misclassification rate below 5%, which requires a minimum *Prochlorococcus* abundance of 0.60% and *Synechococcus* abundance of 0.09% of all classified reads (Fig. 3 A and C).

To establish the minimum *Prochlorococcus*:*Synechococcus* ratio and vice versa, we simulated samples with varying proportions of the two genera and processed them through the classification workflow. False positive rates were calculated by dividing misclassified reads by the total classified reads for each taxon. To keep the *Prochlorococcus* misclassification rate below 10%, the *Prochlorococcus*:*Synechococcus* ratio must be greater than 0.24 (Fig. 3 B). Similarly, to maintain a *Synechococcus* misclassification rate below 10%, the *Synechococcus*:*Prochlorococcus* must be greater than 0.10 (Fig. 3 D). For a more sensitive classification with a 5% misclassification rate, the required ratios are 0.43 for *Prochlorococcus*:*Synechococcus* ratio and 0.20 for *Synechococcus*:*Prochlorococcus* (Fig. 3 B and D).

Figure 3. Misclassification rates and limit of detection for *Prochlorococcus* and *Synechococcus*. Misclassification rates examined across (A) *Prochlorococcus* abundances, (B) *Prochlorococcus:Synechococcus* ratios, (C) *Synechococcus* abundances, and (D) *Synechococcus:Prochlorococcus* ratios. Detection limits were determined at 5% (red dotted line) and 10% (blue dashed line) misclassification thresholds.

Cluster/Clade/Grade Accuracy

To evaluate the workflow's accuracy in estimating the composition of different *Prochlorococcus* and *Synechococcus* clusters/clades/grades, we processed simulated samples with equal proportions of all clusters/clades/grades with and without marine heterotrophic bacteria, following the composition outlined in the Zenodo file (ProSynTaxDB-workflow_benchmarking_composition.tsv)[34]. Specifically, we generated simulated samples containing only the respective picocyanobacterium, as well as mixtures where the respective picocyanobacterium was present at 2% with 98% marine heterotrophic bacteria, and at 1% with 98% marine heterotrophic bacteria plus 1% of the other picocyanobacterium. The classification

results for both *Prochlorococcus* and *Synechococcus* closely matched the expected values, with the exception of a slight increase in misclassification of *Prochlorococcus* grade LLVIII (Fig. 4 A) and *Synechococcus* cluster 5.3 (Fig 4 B). This discrepancy is likely due to their underrepresentation in the genome database, which contains only five reference genomes for each cluster/grade.

Figure 4. Accuracy of *Prochlorococcus* and *Synechococcus* cluster/clade/grade classification. Comparison of expected and classified percentages of *Prochlorococcus* (A) and *Synechococcus* (B) clusters/clades/grades in simulated samples of the respective picocyanobacterium alone, the respective picocyanobacterium at 2% with 98% marine heterotrophic bacteria, and the respective picocyanobacterium 1% with 98% marine heterotrophic bacteria and 1% of the other picocyanobacterium.

Validation with field data

To evaluate the accuracy and effectiveness of the ProSynTaxDB and workflow on field data, we analyzed the total abundance of *Prochlorococcus* and the clade/grade diversity throughout a depth profile at Station ALOHA using metagenomic data from HOT224-238[46]. By applying the 5% detection limit from Figure 3, we found that all reads exceeded this threshold, allowing us to determine the total *Prochlorococcus* read abundance for each depth (Fig. 5 A). We then classified clade/grade diversity at each depth and found predominantly high-light (HL) *Prochlorococcus* clades at <75m and low-light (LL) clades/grades at >125m (Fig. 5 B). These results are consistent with many studies on *Prochlorococcus* diversity at Station ALOHA[22,47–49], demonstrating that the ProSynTaxDB and workflow accurately capture ecological patterns consistent with previous studies.

Figure 5. *Prochlorococcus* clade/grade diversity at STATION ALOHA. Metagenomes from research cruise HOT224-238[46] were used to calculate (A) total *Prochlorococcus* read abundance above the 5% false positive rate (Figure 3) limit of detection (L.O.D., red dotted line) and (B) *Prochlorococcus* relative abundance, calculated using the mean genome equivalent derived from *Prochlorococcus* abundance at each depth.

USAGE NOTES

The code and dependencies for the ProSynTaxDB workflow are publicly available (https://github.com/jamesm224/ProSynTaxDB-workflow) with detailed documentation[34]. This workflow includes a built-in Snakemake[50] pipeline, enabling easy replication and processing of metagenomic data. This database can be customized by allowing users to add new genomes as needed as well as modify the taxonomy in order to improve classification accuracy and adapt the workflow to specific research needs.

To setup and run the ProSynTaxDB workflow, first clone the ProSynTaxDB workflow repository into a suitable directory. Then, download the protein database index file (ProSynTaxDB_v1.fmi), the names (ProSynTaxDB_names.dmp), the nodes (ProSynTaxDB_nodes.dmp), and the diamond blast database (CyCOG6.dmnd) from the Zenodo repository[34]. Additional required metadata files, such cycog_len.tsv and all_illumina_adapters.fa, can be found in the GitHub repository: https://github.com/jamesm224/ProSynTaxDB-workflow. Once the repository is cloned and necessary database files are downloaded, the workflow is ready for use. Before running the workflow, update the input files to specify the correct file paths and directories. After setup, navigate to the scripts directory and execute the run_classify_smk.sbatch script. Comprehensive tutorials are available in the public repository for further guidance[34]. Upon completion, the following files will be generated in the output directory: summary_read_count.tsv (overview of reads counts for *Prochlorococcus*, *Synechococcus*, marine viruses, and marine heterotrophs, as well as unclassified reads) and normalized_counts.tsv (normalized genome equivalents for each *Prochlorococcus* and *Synechococcus* cluster, clade, or grade).

After obtaining the results, we recommend filtering to remove samples with low abundances of *Prochlorococcus* or *Synechococcus* and adjusting the filtering parameters to account for potential errors with samples with lower sequencing coverage. Additionally, samples with large

percentages of unclassified reads or large imbalances of either *Prochlorococcus* or *Synechococcus* may require modification of the filtering parameters. The 10% false positive rate threshold is an effective cut-off for examining differences in high-level cluster/clade/grade differences, but we recommend the 5% false positive rate threshold for cluster/clade/grade-specific delineations.

CODE AVAILABILITY
All ProSynTaxDB files are available on Zenodo: https://zenodo.org/records/14889681

Code and documentation for ProSynTaxDB workflow, along with the accompanying technical validation are available on the GitHub repository:
https://github.com/jamesm224/ProSynTaxDB-workflow

Code and documentation for Genome Closing workflow are available on the GitHub repository:
https://github.com/konnorve/HIFI-genome-closing-improved.git

AUTHOR CONTRIBUTIONS

PMB, MAD, AC, SWC, JIM, NNV conceived and designed experiments. PMB and JWB isolated cultures and AC and PMB maintained cultures. AC, KGC, SMP, ES, CB, and AAA grew cultures and performed DNA extractions. NNV, PMB, and AC closed genomes and submitted data to repositories. JIM, PMB, and NNV developed code, workflow, and database. NNV and JIM performed validation of the database and MAD, AC, and PMB analyzed validation results. All authors contributed to writing and editing the manuscript.

COMPETING INTERESTS

The authors declare no competing interests. The funders had no role in the design or execution of the study nor the decision to submit the work for publication.

ACKNOWLEDGEMENTS

We thank the crews and research teams of the KM0915, MV1015, KM1217, KM1309, and KOK1404 cruises on the research vessels R/V Kilo Moana, R/V Melville, and R/V Kaʻimikai-O-Kanaloa for their support in facilitating the *Prochlorococcus* isolations used in this study. This work was supported by grants from the National Science Foundation (OCE-1153588 to S.W.C.; DBI-0424599 to S.W.C.; OCE-2048470 to P.M.B.), the Gordon and Betty Moore Foundation (GBMF495 to S.W.C.; GBMF4511 to S.W.C.), the Robert and Ardis James Foundation to S.W.C., and the Simons Foundation (Life Sciences Project Award ID 337262, S.W.C.; SCOPE Award ID 329108, S.W.C.; Marine Microbial Ecology Postdoctoral Fellowship 984601, M.A.D.). This paper is a contribution from the Simons Collaboration on Ocean Processes and Ecology (SCOPE).

REFERENCES

1. Flombaum, P. *et al.* Present and future global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*. PNAS **110**, 9824–9829 (2013).

2. Carradec, Q. *et al.* A global ocean atlas of eukaryotic genes. *Nat Commun* **9**, 373 (2018).

3. Clayton, S. *et al.* Bio-GO-SHIP: The Time Is Right to Establish Global Repeat Sections of Ocean Biology. *Front. Mar. Sci.* **8**, (2022).

4. Sunagawa, S. *et al.* Ocean plankton. Structure and function of the global ocean microbiome. *Science* **348**, 1261359 (2015).

5. Sunagawa, S. *et al.* Tara Oceans: towards global ocean ecosystems biology. *Nat Rev Microbiol* **18**, 428–445 (2020).

6. Caputi, L. *et al.* Community-Level Responses to Iron Availability in Open Ocean Plankton Ecosystems. *Global Biogeochemical Cycles* **33**, 391–419 (2019).

7. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927 (2019).

8. Groussman, R. D., Blaskowski, S., Coesel, S. N. & Armbrust, E. V. MarFERReT, an open-source, version-controlled reference library of marine microbial eukaryote functional genes. *Sci Data* **10**, 926 (2023).

9. Mazard, S., Ostrowski, M., Partensky, F. & Scanlan, D. J. Multi-locus sequence analysis, taxonomic resolution and biogeography of marine Synechococcus. *Environmental Microbiology* **14**, 372–386 (2012).

10. Kashtan, N. *et al.* Single-cell genomics reveals hundreds of coexisting subpopulations in wild *Prochlorococcus*. Science **344**, 416–420 (2014).

11. Farrant, G. K. *et al.* Delineating ecologically significant taxonomic units from global patterns of marine picocyanobacteria. *Proc Natl Acad Sci U S A* **113**, E3365-3374 (2016).

12. Doré, H. *et al.* Evolutionary Mechanisms of Long-Term Genome Diversification Associated With Niche Partitioning in Marine Picocyanobacteria. *Front Microbiol* **11**, 567431 (2020).

13. Becker, J. W. *et al.* Novel isolates expand the physiological diversity of *Prochlorococcus* and illuminate its macroevolution. *mBio* **15**, e0349723 (2024).

14. Tarn, J., Peoples, L. M., Hardy, K., Cameron, J. & Bartlett, D. H. Identification of Free-Living and Particle-Associated Microbial Communities Present in Hadal Regions of the Mariana Trench. *Front Microbiol* **7**, 665 (2016).

15. Johnson, Z. I. *et al.* Niche partitioning among *Prochlorococcus* ecotypes along ocean-scale environmental gradients. Science **311**, 1737–1740 (2006).

16. Larkin, A. A. *et al.* Niche partitioning and biogeography of high light adapted *Prochlorococcus* across taxonomic ranks in the North Pacific. *ISME J* **10**, 1555–1567 (2016).

17. Kent, A. G. *et al.* Parallel phylogeography of *Prochlorococcus* and *Synechococcus*. *ISME J* **13**, 430–441 (2019).

18. Thompson, A. W., Kouba, K. & Ahlgren, N. A. Niche partitioning of low-light adapted *Prochlorococcus* subecotypes across oceanographic gradients of the North Pacific Subtropical Front. *Limnology and Oceanography* **66**, 1548–1562 (2021).

19. Ustick, L. J., Larkin, A. A. & Martiny, A. C. Global scale phylogeography of functional traits and microdiversity in *Prochlorococcus*. *ISME J* **17**, 1671–1679 (2023).

20. Hunter-Cevera, K. R., Post, A. F., Peacock, E. E. & Sosik, H. M. Diversity of *Synechococcus* at the Martha's Vineyard Coastal Observatory: Insights from Culture Isolations, Clone Libraries, and Flow Cytometry. *Microb Ecol* **71**, 276–289 (2016).

21. Berube, P. M. *et al.* Single cell genomes of *Prochlorococcus*, *Synechococcus*, and sympatric microbes from diverse marine environments. *Sci Data* **5**, 180154 (2018).

22. Malmstrom, R. R. *et al.* Temporal dynamics of *Prochlorococcus* ecotypes in the Atlantic and Pacific oceans. *ISME J* **4**, 1252–1264 (2010).

23. Ulloa, O. *et al.* The cyanobacterium *Prochlorococcus* has divergent light-harvesting antennae and may have evolved in a low-oxygen ocean. *PNAS* **118**, (2021).

24. Menzel, P., Ng, K. L. & Krogh, A. Fast and sensitive taxonomic classification for metagenomics with Kaiju. *Nat Commun* **7**, 11257 (2016).

25. Bushnell, B. BBMap: A Fast, Accurate, Splice-Aware Aligner. *Conference: 9th Annual Genomics of Energy & Environment Meeting* (2014).

26. Moore, L. R. *et al.* Culturing the marine cyanobacterium *Prochlorococcus*. *Limnol. Oceanogr. Methods* **5**, 353–362 (2007).

27. Berube, P. M. *et al.* Physiology and evolution of nitrate acquisition in *Prochlorococcus*. *ISME J* **9**, 1195–1207 (2015).

28. Wilson, K. Preparation of Genomic DNA from Bacteria. *Current Protocols in Molecular Biology* **56**, 2.4.1-2.4.5 (2001).

29. Kolmogorov, M. *et al.* metaFlye: scalable long-read metagenome assembly using repeat graphs. *Nat Methods* **17**, 1103–1110 (2020).

30. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res* **27**, 722–736 (2017).

31. Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A. & Korobeynikov, A. Using SPAdes De Novo Assembler. *Curr Protoc Bioinformatics* **70**, e102 (2020).

32. Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic Acids Research* **50**, D785–D794 (2022).

33. Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat Biotechnol* **35**, 1026–1028 (2017).

34. Coe, A. *et al.* ProSynTaxDB: *Prochlorococcus* and *Synechococcus* Taxonomy Database. Zenodo https://doi.org/10.5281/zenodo.14889681 (2025).

35. Pachiadaki, M. G. *et al.* Charting the Complexity of the Marine Microbiome through Single-Cell Genomics. *Cell* **179**, 1623-1635.e11 (2019).

36. Garczarek, L. *et al.* Cyanorak v2.1: a scalable information system dedicated to the visualization and expert curation of marine and brackish picocyanobacteria genomes. *Nucleic Acids Research* **49**, D667–D676 (2021).

37. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**, 119 (2010).

38. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* **25**, 1043–1055 (2015).

39. Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol* **7**, 539 (2011).

40. Tamura, K., Stecher, G. & Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular Biology and Evolution* **38**, 3022–3027 (2021).

41. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree: computing large minimum evolution trees with profiles instead of a distance matrix. *Mol Biol Evol* **26**, 1641–1650 (2009).

42. Bushnell, B., Rood, J. & Singer, E. BBMerge – Accurate paired shotgun read merging via overlap. *PLOS ONE* **12**, e0185056 (2017).

43. Li, H. SeqTK, Toolkit for processing sequences in FASTA/Q formats v1.4lh3. *GitHub* https://github.com/lh3/seqtk (2023).

44. Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nat Methods* **18**, 366–368 (2021).

45. Holtgrewe, M. Mason - Tools for Biological Sequence Simulation (v3.0.0-beta1). *GitHub* https://github.com/seqan/seqan/blob/main/apps/mason2/README (2014).

46. Mende, D. R. *et al.* Environmental drivers of a microbial genomic transition zone in the ocean's interior. *Nat Microbiol* **2**, 1367–1373 (2017).

47. Mende, D. R., Boeuf, D. & DeLong, E. F. Persistent Core Populations Shape the Microbiome Throughout the Water Column in the North Pacific Subtropical Gyre. Front. Microbiol. **10**, (2019).

48. Coe, A. *et al.* Survival of *Prochlorococcus* in extended darkness. Limnology and Oceanography **61**, 1375–1388 (2016).

49. Thompson, A. W. *et al.* Dynamics of *Prochlorococcus* Diversity and Photoacclimation During Short-Term Shifts in Water Column Stratification at Station ALOHA. *Front. Mar. Sci.* **5**, (2018).

50. Mölder, F. *et al.* Sustainable data analysis with Snakemake. *F1000Res* **10**, 33 (2021).