The conserved two-component systems CutRS and CssRS control the protein secretion stress response in *Streptomyces* species

Thomas C. McLean[1,2*], Ainsley Beaton[1,2*], Neil A. Holmes [1,2], Carlo Martins[2,3], Gerhard Saalbach[2,3], Govind Chandra[1,2], Sibyl F.D. Batey[1], Jana China[1], Barrie Wilkinson[1,2] and Matthew I. Hutchings[1,2]

[1]Department of Molecular Microbiology, John Innes Centre, Norwich, Norwich Research Park, NR4 7UG, UK.
[2]The Centre for Microbial Interactions, Norwich, Norwich Research Park, NR4 7UG, UK.
[3]Department of Biochemistry and Metabolism, Proteomics Facility, John Innes Centre, Norwich, Norwich Research Park, NR4 7UH, UK.

Correspondence: barrie.wilkinson@jic.ac.uk and matt.hutchings@jic.ac.uk

[*]These authors contributed equally to this work

ORCiD information:
TCM = https://orcid.org/0000-0002-6467-4974
ADMB = https://orcid.org/0000-0002-3799-2600
NAH = https://orcid.org/0000-0002-4979-9680
CM = https://orcid.org/0000-0003-1287-8884
GS = https://orcid.org/0000-0002-9849-4875
GC = https://orcid.org/0000-0002-7882-6676
SFDB = https://orcid.org/0000-0001-6288-660X
JC = https://orcid.org/0009-0000-2143-4113
BW = https://orcid.org/0000-0001-7646-7174
MIH = https://orcid.org/0000-0001-6628-5940

ABSTRACT

Streptomyces secondary metabolites account for over half of all clinically used antibiotics, as well as numerous antifungal agents, anticancer compounds, and immunosuppressants. Two-component systems, which are widespread in bacteria, are key regulators of antibiotic production in *Streptomyces* species, yet their activating signals remain poorly understood. CutRS was the first two-component system identified in the genus *Streptomyces* and deletion of *cutRS* in *Streptomyces coelicolor* was shown to enhance antibiotic production, although its CutR regulon does not include any biosynthetic genes. Here, we used *Streptomyces venezuelae* to further investigate CutRS function. We show that deletion of *cutRS* leads to an increase in growth rate and a reversal of the typical glucose-mediated carbon catabolite repression typically observed in *Streptomyces* species. We also demonstrate that CutR DNA binding is glucose-dependent, but CutR does not directly regulate genes involved in growth, antibiotic biosynthesis, or glucose metabolism. The only CutR targets conserved in both *S. coelicolor* and *S. venezuelae* are the foldase genes *htrA3* and *htrB*, which are involved in the protein secretion stress response. Consistent with this, we show that CutS homologues all contain two conserved cysteine residues in their extracellular sensor domains and that changing these residues to serine constitutively activates *S. venezuelae* CutRS. We propose that failure of a disulfide bond to form between these cysteine residues indicates secretion stress and leads to activation of the CutRS system and the secretion stress response.

IMPORTANCE

Streptomyces bacteria are the primary source of clinically useful antibiotics. While many two-component systems have been linked to antibiotic biosynthesis in *Streptomyces* species, few have been well characterized. Here, we characterize a secretion stress sensing two-component system called CutRS and propose a model for how the sensor kinase detects extracellular protein misfolding via two highly conserved cysteine residues. Importantly, we also show that deletion of *cutRS* triggers antibiotic overproduction in the presence of glucose. Since glucose normally represses antibiotic biosynthesis in *Streptomyces* species through carbon catabolite repression, this finding reveals a simple genetic route to bypass this barrier. This has significant implications for antibiotic discovery pipelines and industrial production, where glucose-rich media are preferred for cost and scalability. Our results position CutRS as a key target for future strain improvement strategies.

INTRODUCTION

Streptomyces species are studied for their complex life cycles and prolific production of specialized metabolites, including many clinically important antibiotics (1). These metabolites are typically encoded by biosynthetic gene clusters (BGCs), but despite the wealth of genomic data available, only ~3% of BGCs have been linked to their corresponding natural products (2). The vast majority remain cryptic under standard laboratory conditions,
50   representing untapped reservoirs for novel bioactive compounds (3, 4). Unlocking these cryptic BGCs requires a better understanding of the regulatory mechanisms, including signal transduction pathways, that control specialized metabolism.

BGC expression is often regulated by transcription factors encoded within the cluster itself (cluster-situated regulators), as well as by global regulatory systems that integrate signals related to nutrient availability, growth phase, and environmental stress to specialized metabolite production (5). Among these global regulators, two-component systems (TCSs) are particularly important. TCSs are widespread in bacteria, fungi, and plants, and many have been shown to influence antibiotic production in *Streptomyces* species. However, the mechanisms by which they exert this control are often poorly understood (6–9).

60   TCSs typically consist of a membrane-bound sensor kinase (SK) and a cognate response regulator (RR). Upon detection of an external signal, the SK autophosphorylates and transfers the phosphate to the RR, which then dimerizes and binds DNA to modulate the transcription of target genes. *Streptomyces* genomes encode between 50 and 100 TCSs, yet only 15 are highly conserved across species (10). Of these, several have been shown to impact antibiotic biosynthesis, but only a small number have been functionally characterized, including their target regulons.

CutRS was the first TCS identified in the genus *Streptomyces* (11), and its disruption in *S. lividans* and *S. coelicolor* resulted in overproduction of the antibiotic actinorhodin (11, 12). However, a subsequent study in *S. coelicolor* revealed that none of the 88 genes directly
70   regulated by CutR were involved in actinorhodin biosynthesis (13). Instead, the CutRS system was implicated in the protein secretion stress response, with CutR activating *htrA3* and repressing *htrB*, both of which encode high-temperature requirement (Htr) chaperone-proteases that assist in folding or degradation of misfolded proteins in the extracellular space (13).

Here, we investigated the CutRS system in *Streptomyces venezuelae*, a genetically tractable and rapidly growing model organism that has emerged as a powerful system for studying *Streptomyces* development and regulation (14). We found that *S. venezuelae* CutR binds nine promoter regions, but the core CutR regulon shared with *S. coelicolor* includes only *htrA3* and *htrB*, reinforcing a conserved role in secretion stress regulation. Alignment of
80   >100 CutS homologues across *Streptomyces* species revealed two invariant cysteine residues as the only conserved features in their predicted extracellular sensor domains. We demonstrate that changing these cysteines to serine constitutively activates CutRS, suggesting that the system senses protein misfolding via disulfide bond status outside the cell.

Although CutRS is only conserved in *Streptomyces* species, analysis of SK sequences from 12,799 bacterial genomes revealed that 99% encode at least one SK with two or more cysteines in their predicted extracellular sensor domains. We therefore propose that extracellular redox sensing could be a widespread signaling mechanism across the bacterial domain.

RESULTS

Deletion of *cutRS* affects growth and antibiotic production in *S. venezuelae*

To investigate the function of CutRS, we deleted the *cutRS* operon in *S. venezuelae* and found that the Δ*cutRS* mutant exhibits a striking phenotype when grown on yeast peptone + D-glucose (YPD) agar. The mutant displays increased antibacterial activity and accelerated growth (Figures 1, 2A, and 2B). Further analysis revealed that the Δ*cutRS* strain produces 10-fold more biomass than the wild-type strain on YPD agar. This phenotype was not restricted to YPD medium; a similar increase in biomass was observed on MYM agar, but again only in the presence of D-glucose (Figure 2B). This increased growth correlates with the rapid

100   depletion of D-glucose from the medium (Figure 2C). Consistent with this, in the absence of D-glucose (YP agar), no phenotypic differences were observed between the wild-type and Δ*cutRS* strains (Figures 1A, 2A, and 2B).

When challenged with *Bacillus subtilis* on YP agar, both the wild-type and Δ*cutRS* strains produced small zones of inhibition (Figure 1A). However, on YPD agar, bioactivity was lost in the wild-type strain, consistent with the known inhibition of antibiotic biosynthesis by D-glucose via carbon catabolite repression (15), while the Δ*cutRS* strain exhibited a markedly larger zone of inhibition (Figure 1A). This suggests that D-glucose triggers antibiotic overproduction in the Δ*cutRS* strain, supporting previous observations that glucose activates actinorhodin production in the *S. coelicolor* Δ*cutRS* mutant (13). Notably, this is atypical, as

110   D-glucose generally represses antibiotic production in *Streptomyces* species (15, 16).

S. venezuelae is known to produce chloramphenicol and jadomycin, both of which inhibit *B. subtilis*. To determine which antibiotic was responsible for the observed activity, we deleted both BGCs to generate strain M1702. This strain failed to inhibit *B. subtilis* growth, indicating that the bioactivity of the parent strain was due to either chloramphenicol or jadomycin (Figure 1A). To further resolve this, we challenged the wild-type and Δ*cutRS* strains with a chloramphenicol-resistant *B. subtilis* strain. The Δ*cutRS* strain produced only a faint zone of inhibition, markedly smaller than that observed against the chloramphenicol-sensitive strain, confirming that the bioactivity was primarily due to chloramphenicol (Figure 1B).

Finally, we extracted cultures from YPD agar plates with ethyl acetate and analyzed the

120   samples by HPLC. The Δ*cutRS* strain produced approximately three times more chloramphenicol than the wild-type in the presence of D-glucose, confirming that increased chloramphenicol production is responsible for the enhanced antibacterial activity of the Δ*cutRS* strain (Figure 1C).

Identifying CutR target promoters and binding sites in *S. venezuelae*

Two-component systems typically regulate gene expression through their response regulators (RRs)—in this case, CutR. To define the CutR regulon, we complemented the *S. venezuelae* Δ*cutRS* strain with a modified *cutRS* operon encoding wild-type CutS and a C-terminally 3×Flag-tagged CutR. We then cultured both the wild-type and Flag-tagged strains on YP and YPD agar and performed ChIP-seq using monoclonal α-Flag antibodies.

130 Analysis of the sequencing data (accession number GSE225370) revealed a single significantly enriched peak in the Flag-tagged CutR strain grown on YP agar (absence of D-glucose), located upstream of the *cutRS* operon. In contrast, ten enriched peaks were detected in cultures grown on YPD agar (presence of D-glucose) (Table 1). These included peaks upstream of *htrA3* and *htrB*, which are also known CutR targets in *S. coelicolor* (Figure 3A), as well as a strong peak upstream of *vnz_08815*, a putative cell wall amidase gene.

To precisely define CutR binding sites, we expressed and purified hexahistidine-tagged CutR for use in a DNA footprinting method called Reusable DNA Capture Technology Surface Plasmon Resonance (ReDCaT SPR) (17). Using tiled, double-stranded oligonucleotide probes spanning the promoters of *htrA3* and *vnz_08815*, we measured CutR binding affinity via SPR.
140 A preliminary scan identified the core binding regions, and these probes were then truncated in 2 bp steps to determine the minimal binding sequences (Figure 3B). CutR bound the sequence TATATAAA in the *htrA3* promoter and TAAATAAA in the *vnz_08815* promoter (Figure 3B). From these and other enriched regions, we defined the CutR consensus binding sequence as TAWATAAA.

CutR is both a transcriptional activator and repressor

Having identified CutR binding sites at its target promoters (Table 1), we cross-referenced them with publicly available *S. venezuelae* transcription start site (TSS) data from http://streptomyces.org.uk (18). The CutR binding site at the *htrB* promoter is located 17 bp
150 upstream of the TSS, while at the *htrA3* promoter it lies 62 bp upstream (Table 1 and Figure 3A). The position of a regulator binding site relative to the TSS often determines whether it functions as an activator or repressor (19). Binding 17 bp upstream of the *htrB* TSS likely overlaps the core promoter region, interfering with RNA polymerase binding and resulting in transcriptional repression (20, 21). In contrast, binding further upstream, such as at the *htrA3* promoter, may facilitate transcriptional activation via interaction between CutR and RNA polymerase bound at the -35 site (21–23).

To assess the effect of *cutRS* deletion on CutR target gene products, we used tandem mass tag (TMT) quantitative proteomics. Of the ten predicted CutR-dependent proteins, six were detected (Table 1). These data show that CutR activates expression of HtrA3 (8-fold increase
160 in the wild-type strain compared to the Δ*cutRS* mutant) and the putative cell wall amidase Vnz_08815 (2.5-fold increase), consistent with the location of its binding sites upstream of their promoters (Figure 4, Table 1, and Supplementary Table 1). In contrast, HtrB production was repressed by CutR, showing a 19-fold increase in the Δ*cutRS* strain relative to the wild-type, consistent with CutR binding in a position that occludes RNA polymerase binding (Figure 3A).

The expression of the remaining detected target genes was not significantly affected by *cutRS* deletion. This may be because these genes are not directly regulated by CutR under the tested growth conditions, because CutR binding does not alter transcription in all cases, or because additional layers of regulation are involved (Table 1).

Table 1. Gene promoters with CutR binding sites that are enriched in the ChIP-seq dataset. The binding site positions are given relative to the transcript start sites (TSS). The DNA sequences of the CutR binding sites are shown, two of which (*htrA3* and *vnz_08815*) were verified *in vitro* using ReDCaT SPR (Figure 3).

Gene	Predicted function	CutR binding site relative to TSS	Binding site sequence	Predicted effect	Protein abundance (WT/Δ*cutRS)*
vnz_06385	Acyl coA dehydrogenase	+21	TAATTAAA	Repression	0.795
vnz_06390	TetR-family regulator	+28	TAATTAAA	Repression	0.947
vnz_08815	Cell wall amidase	-69	TAAATAAA	Activation	2.56
vnz_12615	Membrane protein	-4	TACATAAA	Repression	ND
htrA3	Quality control protease	-62	TATATAAA	Activation	8.134
htrB	Quality control protease	-17	CATATAAA	Repression	0.053
vnz_29845	NAD(P)/FAD-dependent oxidoreductase	-28	CAAATAAA	Repression	0.707
vnz_33990	Beta mannosidase	-21	TAAGTAAA	Repression	0.747
vnz_33995	Sugar binding protein for ABC transport	+1	TAAGTAAA	Repression	0.977

CutS activity is controlled by two conserved cysteines in its extracellular sensor domain.
The conservation of *htrA3* and *htrB* as CutR targets in the distantly related species *S. coelicolor* and *S. venezuelae* supports the hypothesis that CutRS plays a role in the protein secretion stress response. This further implies that CutRS complements the function of CssRS, which activates the expression of *htrA1*, *htrA2* and *htrB*, but not *htrA3* (24). We thus set out to investigate how the extracellular sensor domain, which is predicted to be between the two transmembrane helices of CutS, might detect defects in protein folding on the outside of the cell. We first aligned the extracellular sensor domains of >100 CutS proteins taken from complete, published *Streptomyces* genome sequences to look for conserved residues or motifs. This analysis identified two cysteine residues that are conserved across every CutS sensor domain (Supplementary Figure 3). These are the only conserved residues in the CutS sensor domain. Cysteines are important in extracellular protein folding in actinobacteria (25) and, consistent with this, the analysis of all proteins encoded in *S. venezuelae* with a Sec signal sequence revealed that 74% contain two or more cysteines that likely form disulfide bonds to help them fold. We thus hypothesized that CutS monitors disulfide bond formation in Sec-translocated proteins. Given the conservation of the two cysteine residues, we reasoned that this could be via the making or breaking of a disulfide bond in the CutS extracellular sensor domain.

Modelling of the *S. venezuelae* CutS extracellular sensor domain using AlphaFold indicated that these cysteine residues likely sit within 5Å of each other, the distance required to form disulfide bonds (Supplementary Figure 4). To test this, we designed a CutS variant where both cysteine residues were replaced with serine (CutS(C85S,C103S)) and introduced it into the *S. venezuelae* wild-type and Δ*cutRS* strains along with the wild-type *cutR* gene. This restored a wild-type growth phenotype to the Δ*cutRS* mutant (Figure 5A) which suggests the CutS(C85S,C103S) variant is active. However, previous work on the VanRS TCS demonstrated that it is possible for loss of a SK to render its cognate RR constitutively active (26). To discount this possibility, we introduced individual copies of either *cutS* or *cutR* under the control of the *ermE** promoter into the Δ*cutRS* mutant, but neither restored the wild-type phenotype, confirming that CutS is required for CutR activity. To further investigate the activity of the CutS(C85S,C103S) variant we used qRT-PCR to measure the expression of the CutR target genes *htrA3* and *htrB* in the wild-type, Δ*cutRS* and CutS(C85S,C103S) strains (Figure 5B). The expression of *htrA3* was higher in the strain harboring the CutS(C85S,C103S) variant compared to the wild-type strain, while the expression of *htrB* was reduced (Figure 5B) which is consistent with CutS(C85S,C103S) activating CutR which then directly activates the transcription of *htrA3* and represses *htrB*. Thus, we conclude that replacing the two extracellular sensor domain cysteines with serine residues results in a constitutively active CutRS system which supports the hypothesis that an inability to form disulfide bonds outside the cell activates CutRS and triggers the secretion stress response.

The wild-type phenotype is recovered by the deletion of *cssRS* in the Δ*cutRS* background

This work and previous reports suggest that the secretion stress response in *Streptomyces* species is controlled by the CssRS and CutRS two-component systems which together control the expression of *htrA1*, *htrA2*, *htrA3* and *htrB*, which encode the four conserved HtrA-family foldases in *Streptomyces* species (24, 27). Although the CssR regulon has not yet been defined, electrophoretic mobility shift assays alongside qRT-PCR were used previously in *S. lividans* (a very close relative of *S. coelicolor*) to show that CssR directly activates expression of *htrA1*, *htrA2* and *htrB*, but not *htrA3* (24). Instead, the expression of *htrA3* is directly activated by CutR, which also represses the expression of *htrB* in *S. coelicolor* and *S. venezuelae* (27). We attempted to replicate the Δ*cutRS* phenotype by deleting *htrA3* and overexpressing *htrB* in the same background (Supplementary Figure 2), but this did not result in the phenotypic change on YPD that we observed with Δ*cutRS*. We thus reasoned that the CssRS system might be compensating for the loss of CutRS. To further investigate the interplay of CssRS and CutRS we made single and double Δ*cssRS* and Δ*cutRS* mutants of *S. venezuelae* and observed their phenotypes on YP and YPD agar. In the absence of D-glucose, the wild-type, Δ*cssRS*, Δ*cutRS* and Δ*cssRS*-Δ*cutRS* strains all displayed the same phenotype (Figure 6). On YPD agar, growth is reduced in the wild-type strain by the addition of D-glucose to the medium, but the Δ*cutRS* mutant consumes the glucose and reverts to a phenotype like that observed on YP agar (Figures 1 and 5). Deletion of *cssRS* in the wild-type background has no obvious effect on growth on either YP or YPD agar, but deletion of *cssRS* in the Δ*cutRS* mutant restores the wild-type phenotype to the Δ*cutRS* mutant growing on YPD agar, suggesting that the altered growth of the Δ*cutRS* mutant on YPD is dependent on CssRS (Figure 6). Proteomics analysis of the wild-type and Δ*cutRS* strains revealed that the levels of CssRS are increased 5-fold in the Δ*cutRS* mutant, and as previously discussed, HtrB levels (activated by CssR and repressed by CutR) are increased 20-fold in the absence of *cutRS*. The *htrB* and *cssRS* genes are located adjacent to each other and likely form an operon under the direct control of both CutR (negative) and CssR (positive). Overproduction of HtrB could be due to the loss of CutR repression in the mutant strain combined with the overproduction of CssRS, with CssR further activating *htrB* expression. A future challenge will be to define the CssR regulon and determine how deletion of *cssRS* complements Δ*cutRS*.

Extracellular disulfide sensing could be conserved across bacteria. Analysis of the 15 conserved two-component systems in *Streptomyces* species (28–30) revealed that only CutS and CssS have two conserved extracellular cysteines. Given that these cysteines are implicated in the control of CutS activity we hypothesise the same may be true for CssS and perhaps other bacterial SKs. To investigate how widespread this might be in the bacterial kingdom, we analyzed 12,799 high-quality bacterial genomes, identifying 348,633 proteins predicted to function as SKs. Using DeepTMHMM (31) we predicted the transmembrane helices for all the SKs and then used this information to identify the putative sensor domains of these proteins and quantify the number of cysteine residues present in these extracellular domains. We observed that 98.9% of all the bacterial strains examined encode

at least one SK with two or more cysteines in their predicted extracellular sensor domains (Figure 7A).

Next, we examined the conservation of the CutS and CssS SKs beyond the genus *Streptomyces*. Reciprocal BLAST searches in 2,936 bacterial reference genomes revealed that 39% encoded a CutS homologue and 27% encoded a CssS homologue. Of the 2,936 genomes examined, 53% encode at least one of these two SKs and 12% encode both CutS and CssS. However, further inspection of the amino acid sequences of these homologues revealed that only 6.9% of CutS and 4.8% of CssS homologues contain two or more cysteines in their extracellular domains. Phylogenetic analysis using a maximum likelihood tree based on RpoB revealed a broad distribution of these homologues across bacterial taxa (Figure 7B). The presence of both CutS and CutR with two extracellular cysteines was universally conserved across all tested *Streptomyces* strains, with only three additional genomes, all from actinobacteria, also containing both. This may reflect the fact that actinobacteria are more likely to use disulfide bond formation as a mechanism to fold their secreted proteins (32). Taken together, our findings suggest that extracellular redox sensing via conserved cysteine residues in the sensor domains of sensor kinases (SKs) may represent a broadly important mechanism across the bacterial kingdom. While CutS and CssS appear to be largely restricted to the genus *Streptomyces*, this sensing function is likely carried out by a diverse array of SKs in other bacterial lineages.

DISCUSSION

In this study, we demonstrate that loss of the highly conserved CutRS TCS in *Streptomyces venezuelae* leads to dramatic changes in both growth and antibiotic production. To investigate the underlying mechanisms, we first defined the *S. venezuelae* CutR regulon. This revealed two conserved CutR targets—*htrA3* and *htrB*—shared between *S. venezuelae* and the distantly related *S. coelicolor*. These genes encode members of the high temperature requirement A (HtrA) family of proteases. In *S. venezuelae*, CutR represses *htrB* and activates *htrA3*. CutR also activates *vnz_08815*, a gene predicted to encode a cell wall amidase of unknown function. Using ReDCaT-SPR, we identified a consensus CutR binding motif (TAWATAAA) in the promoters of *htrA3* and *vnz_08815*, which was also found in five additional CutR-enriched promoter regions, thereby defining the complete CutR regulon. We further showed that the position of CutR binding relative to transcription start sites (TSSs) correlates with transcriptional outcome—either activation or repression. TMT proteomic analysis confirmed these regulatory effects, showing an increase in the levels of HtrB and a decrease in HtrA3 levels in the absence of *cutRS*. Given that CutRS coordinates the expression of two of the four conserved *htrA*-like genes in *S. venezuelae* and *S. coelicolor*, we conclude that CutRS functions in the secretion stress response, likely acting in parallel with the CssRS TCS .

While the CssR regulon remains undefined, previous work shows that CssR activates *htrA1*, *htrA2*, and *htrB* while having no effect on *htrA3* (33). Thus, CutRS and CssRS appear to act in opposition: they regulate distinct subsets of *htrA*-family genes and exert opposing effects on *htrB* expression. HtrA proteins support extracellular protein folding and degrade irreparably

misfolded proteins in both Gram-negative and Gram-positive bacteria (34). CssRS, conserved in Gram-positive species, is homologous to the envelope stress-sensing system CpxAR in *E. coli*. Our bioinformatic analysis shows that while CutRS and CssRS are widely conserved in *Streptomyces*, most other Actinobacteria encode one or the other, and homologs are also present across other bacterial phyla (Figure 7).

Proper folding of secreted proteins requires disulfide bond formation, catalyzed by DsbA-family proteins located on the extracellular face of the cytoplasmic membrane (35). The discovery that the sensor domain of CutS contains two conserved cysteines suggests a redox-based sensing mechanism. Indeed, mutating these residues to serines (C85S, C103S) 310 led to constitutive activation of CutR, with elevated *htrA3* expression and enhanced repression of *htrB*. This supports a model in which CutS monitors disulfide bond formation as a proxy for protein folding quality in the extracytoplasmic space.

Intriguingly, the extracellular domain of CssS also contains two conserved cysteines, and CutS and CssS are the only conserved *Streptomyces* sensor kinases (SKs) with this feature. This suggests that CssS may similarly sense disulfide bond status, which we intend to explore in future work. Moreover, *cssRS* is overexpressed in the Δ*cutRS* mutant, and deleting *cssRS* in this background restores wild-type growth, further supporting antagonistic functions for these two systems. Additional work is needed to define the CssR regulon and elucidate how CutRS and CssRS jointly regulate the HtrA proteases.

320 To our knowledge, this is the first detailed proposal of a mechanistic sensing model for a bacterial secretion or envelope stress response system. *Streptomyces* species are common in soil and the plant rhizosphere and endosphere (36), where protein secretion plays a vital role in breaking down complex organic material for nutrient acquisition. These environments are also oxygen-variable. We propose that, as illustrated in Figure 8, CutS senses oxygen availability via disulfide bond formation. Under anaerobic or microaerobic conditions, disulfide bonds cannot form, leading to an accumulation of misfolded extracellular proteins. We suggest that a disulfide bond fails to form in CutS under these conditions, activating the CutRS system and promoting expression of *htrA3*, which aids in resolving misfolded proteins.

330 Oxygen sensing via disulfide bond status has precedence. In *Shewanella* species, the MtrABC porin-cytochrome complex facilitates extracellular electron transfer under anoxic conditions (37). The periplasmic MtrC subunit contains a conserved CX_8C motif that forms a redox-active disulfide only in the presence of oxygen, allowing the cell to modulate reactive oxygen species and electron flux (38). Similarly, in *Pseudomonas aeruginosa*, the periplasmic protein YfiR contains four redox-active cysteines, two of which form a disulfide under oxidative conditions. YfiR modulates the activity of the diguanylate cyclase YfiN in response to oxygen availability (39).

The widespread conservation of sensor kinases with extracellular cysteines across diverse bacterial phyla suggests that redox and oxygen sensing outside the cell is a fundamental 340 feature of bacterial stress response systems. Future research into these pathways may reveal new regulatory strategies for secretion stress, redox adaptation, and virulence in a broad range of bacterial species.

METHODS

Growth media and supplements

Protocols for culturing and genetically manipulating *Streptomyces* species are all freely available from http://actinobase.org (40). Strains were grown in the following media (per L volume): Yeast Peptone (YP) agar (10 g yeast extract, 20 g bacto-peptone, 20 g agar); YPD (YP + D-glucose) agar (YP agar with the addition of 100 ml of a 40% filter-sterilized D-350 glucose); 2xYT (16 g tryptone, 10 g yeast extract, 5 g NaCl); SFM agar (20 g soy flour, 20 g mannitol, 20 g agar); lysogeny broth (LB) (10 g tryptone, 5 g yeast extract, 10 g NaCl) and LB soft agar (LB with the addition of 20 g agar). When appropriate, the media was supplemented with antibiotics at the following concentrations: hygromycin (50 μg/ml), apramycin (50 μg/ml) and nalidixic acid (25 μg/ml). NaCl was excluded from LB when hygromycin was used for selection.

Strains, plasmids, and primers

The bacterial strains, plasmids and primers used or generated in this study are listed in Supplementary Tables 2, 3 and 4, respectively.

Strain generation

Deletion mutants were complemented using the native gene and promoter cloned into the phage integrative vector pSS170 (40) using Gibson Assembly. Overexpression strains were generated by cloning the gene(s) of interest into pIJ10257 [28] using Gibson Assembly. CRISPR mediated gene deletions were carried out using pCRISPomyces2 as per the protocol described by Cobb et al.(41). In all cases the resulting construct was then used to transform *E. coli* ET12567/pUZ8002 (42, 43) and conjugated into the mutant strains as follows. Single colonies of ET12567/pUZ8002 containing the complementation construct were inoculated into 10 ml LB with appropriate selection and incubated overnight at 37°C with shaking at 370 220 rpm. The overnight culture was sub-cultured 1:100 in 50 ml LB with appropriate selection and grown to OD_{600} ~0.6. The cultures were washed with 10 ml ice-cold LB twice to remove the antibiotic and finally resuspended in 1 ml ice-cold LB. 500 μl of the *E. coli* cell suspension was mixed with 20 μl of *S. venezuelae* spores by inversion and then pelleted by centrifuging at 15,871 xg for 2 minutes. The supernatant was removed, and the pellet resuspended in 150 μl residual supernatant. Serial 10-fold dilutions were plated on SFM and incubated at room temperature for 16-20 hours, overlaid with 1 ml sterile distilled H_2O containing appropriate selection and then incubated at 30°C for 3-7 days until colonies appeared. Colonies were re-streaked on SFM agar with appropriate selection at least once before being plated for spore preparation.

Generation of *S. venezuelae* strain M1702

Deletion of the chloramphenicol biosynthetic gene cluster (BGC) and jadomycin BGC was achieved using the meganuclease system (44). The upstream flank for the chloramphenicol BGC was amplified with oligonucleotides NH112ChlPCR1F and NH115ChlPCR2R, whereas

the upstream flanking of the jadomycin cluster was amplified with NH120JadPCR1F and NH123JadPCR2R. The upstream flanking sequences were then cloned into the BglI and HindIII sites of pIJ12738 (44). The downstream flanking sequence of the chloramphenicol BGC was amplified with oligonucleotides NH116ChlPCR3F and NH119ChlPCR4R, whereas the upstream flanking sequence of the jadomycin BGC was amplified with NH124JadPCR3F

390 and NH127JadPCR4R. The downstream flanking sequences were combined with the corresponding upstream flanking sequences in pIJ12738 by cloning into the HindIII and SpeI sites. These pIJ12738 derivatives containing the flanking sequencing to each cluster followed by a I-SceI meganuclease cut site were then mobilized via conjugation through *E. coli* ET12567/pUZ8002 and single crossover recombinants were selected for with apramycin. We then conjugated pIJ12742 containing the meganuclease into the single crossover recombinants. Expression of the meganuclease results in a double strand break, which can be resolved by a second crossover event. Subsequent double crossovers were screened by PCR to determine if they were gene cluster deletion mutants or if they had reverted to wild-type. Gene cluster deletion mutants were incubated at 37°C for two generations to promote

400 loss of pIJ12742. The resulting strains were Δ*cml* (M1700) and Δ*jad* (M1701). M1700 was used as the starting material to introduce the jadomycin BGC deletion derivative of pIJ12738, and single crossover recombinants were used to introduce the meganuclease (on pIJ12742) followed by screening for loss of the jadomycin BGC and removal of the plasmid, generating the final double mutant of Δ*cml*Δ*jad* (M1702).

Antimicrobial bioassays

5 µl of spores of the relevant *S. venezuelae* strains were inoculated onto the centre of a YP or YPD agar plate and incubated for 5 days at 30°C. The *B. subtilis* indicator strain was grown in 10 ml LB at 37°C with 220 rpm shaking the night prior to the assay. Following overnight

410 incubation, the indicator strains were sub-cultured 1:100 into 50 ml LB and grown to OD_{600} ~0.6. and then diluted 1:10 into 50°C Soft LB agar. 5 ml of this inoculated soft agar was then pipetted over the *S. venezuelae*-containing plate ensuring complete coverage. The plates were air-dried and subsequently incubated at 37°C overnight before imaging.

Biomass and glucose quantification assays

5 µl of spores of the relevant *S. venezuelae* strains were inoculated on cellophane-covered YP or YPD agar plates for 5 days at 30°C. Following incubation, the cellophane disks were removed, and the mycelial mass was measured using an analytical balance. For glucose extraction the underlying agar was chopped into small pieces using a sterile scalpel and

420 transferred to a 250 ml conical flask. 20 ml sterile distilled H_2O was added, and samples were incubated for 2 hours at 25°C with shaking at 220 rpm. The resulting liquid was aspirated from the flask and diluted 1:1000 with sterile distilled H_2O before being assayed using the Glucose (GO) Assay Kit (Sigma Aldrich) following manufacturer's instructions.

Chromatin immunoprecipitation followed by sequencing (ChIP-Seq)

5 μl of spores of the relevant *S. venezuelae* strains were inoculated onto cellophane-covered YP or YPD agar plates in triplicate and grown for 5 days at 30°C. To cross-link proteins to DNA the cellophane disks were removed and submersed in 10 ml fresh 1% (v/v) formaldehyde solution at room temperature for 20 minutes. Following a 5-minute incubation in 10 ml 0.5 M glycine solution the mycelium was harvested and washed twice with 25 ml ice-cold PBS (pH 7.4) before being flash frozen in liquid nitrogen and stored at -80°C. For lysis, the pellets were resuspended in 2 ml lysis buffer (10 mM Tris-HCl pH 8.0, 50 mM NaCl, 10 mg/ml lysozyme, EDTA-free protease inhibitor) before 750 μl was transferred to a 2 ml microcentrifuge and samples incubated at 37°C for 30 minutes. For fragmentation 750 μl IP buffer (100 mM Tris-HCl pH 8.0, 250 mM NaCl, 0.5% v/v Triton X-100, 0.1% SDS, EDTA-free protease inhibitor) was added and samples sonicated on ice at 50 Hz for 20 cycles of 10 seconds with at least 1 minute of rest between each cycle. 25 μl of the crude lysate was combined with 75 μl TE Buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA) and extracted with 100 μl phenol:chloroform. 2 μl of 1 mg/ml RNase A was combined with 25 μl of the resulting extract, incubated at 37°C for 30 minutes and run on a 1% agarose gel to confirm a smear centred at 500 bp. The remaining crude lysate was centrifuged for 15 minutes at 15,871 xg at 4°C and supernatant saved. 750 μl of α-FLAG M2 magnetic beads (Sigma Aldrich) were washed in 3.75 ml 0.5% IP buffer (IP buffer - 150 mM NaCl, 1% Triton X-100, 50 mM Tris-HCl (pH 8.0), 2 mM EDTA, and Roche cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail tablet(s)) to prepare for them binding. The bead slurry was then mixed with 40 μl clarified lysate and incubated overnight on a vertical rotator at 4°C. The lysate was removed, and the beads washed four times with 500 μl 0.5 IP buffer with 10 minutes vertical rotation at 4°C each cycle. Washed beads were incubated in 100 μl elution buffer (50 mM Tris-HCl pH 7.6, 10 mM EDTA, 1% SDS) overnight at 65°C. Eluant was removed and saved before an additional 50 μl elution buffer was added and incubated at 65°C for a further 5 minutes. Once removed 3 μl of 10 mg/ml proteinase K was added to the combined eluants and incubated at 55°C for 2 hours. DNA was extracted with 200 μl phenol:chloroform and purified using a QIAquick PCR Purification Kit (QIAGEN) following the manufacturer's instructions. DNA was eluted in 50 μl Buffer EB (10 mM Tris-HCl pH 8.5) and 47 μl snap frozen in liquid nitrogen and stored at -80°C. The remaining 3 μl of sample was used to determine the DNA concentration and quality. The Qubit Fluorometer 2.0 with the high-sensitivity kit was used for DNA concentration whilst DNA purity was quantified using a Nanodrop™ 2000 UV-Vis Spectrophotometer. The stored DNA samples were the shipped to Novogene on dry ice for sequencing using the NovaSeq 6000 Sequencing System (Illumina). Raw sequencing data were received as FASTQ files from Novogene and processed as described previously (27). Raw files were aligned to the *S. venezuelae* NRRL B-65442 genome and enrichment normalised to a moving window of 30 nucleotides compared to a region of the surrounding 3000 nucleotides with the window moving in steps of 15 nucleotides.

Protein overexpression and purification

Protein preparation for ReDCaT SPR was performed as follows. N-terminally 6xHis-tagged CutR was expressed from the plasmid pET28a in *E. coli* Rosetta (BL21 DE3) pLysS competent cells. Overnight culture (40 ml) was used to inoculate 4 L of LB with selective antibiotics.

470 Cultures were incubated at 37°C with shaking at 220 rpm until OD_{600} ~0.6 before isopropyl-β-D-1-thiogalactopyranoside (IPTG) was added to a final concentration of 1 mM. Cultures were incubated for a further 4 hours then the cells were pelleted by centrifugation and stored at -80°C. Cell pellets were defrosted on ice for 30 minutes before being resuspended in 25 ml lysis buffer (20 mM Tris-HCl pH 8.0, 75 mM NaCl, 0.1% Triton X-100, 10 mg/ml lysozyme, EDTA-free protease inhibitor) and incubated at room temperature for a further 30 minutes. Suspensions were sonicated on ice 8 times at 50 Hz for 30 seconds per cycle with 1 minute off between each sonication cycle. Cell debris was removed by centrifugation at 20,600 xg for 30 minutes at 4°C and the supernatant filtered through a sterile 0.22 μm filter (Sartorius). Clarified lysate was loaded onto a 1 ml HisTrap HP column (Cytiva) pre-equilibrated with Buffer A (50 mM Tris-HCl pH 8.0, 200 mM NaCl, 5% glycerol). The loaded column was washed with increasing concentrations of imidazole (step 1: 10 mM; step 2: 20 mM; step 3: 30 mM) before the protein was eluted using 500 mM imidazole, all in the same buffer. Desired protein fractions were pooled and loaded onto a preparative-grade HiLoad 16/600 Superdex 200 pg gel filtration column (GE Healthcare) pre-equilibrated with gel filtration buffer (50 mM Tris-HCl pH 8.0, 200 mM NaCl, 10% glycerol, filtered). Desired fractions were identified and analysed for purity via SDS-PAGE before being pooled. Aliquots were flash frozen in liquid nitrogen and stored at -80°C.

Reusable DNA Capture Technology Surface Plasmon Resonance (ReDCaT SPR)

490 Protein-DNA binding site scanning and footprinting was performed using ReDCaT SPR as previously described (45). Preliminary scanning was performed using promoter regions of interest divided into a series of overlapping double stranded oligonucleotide probes (Supplementary Table 4). The total length of each fragment was 40 bp including 15 bp overlaps. To create the double-stranded probes each fragment was reverse-complimented and a ReDCaT biotinylated single-stranded linker added to the 3' end as an overhang. For each fragment DNA oligos (IDT) were resuspended to 100 μM in sterile distilled H_2O and 55 μl forward oligo was mixed with 45 μl reverse oligo. Oligos were annealed at 95°C for 5 minutes followed by ramping to 4°C at 0.1°C/s. The annealed DNA fragments were then diluted to a final concentration of 1 μM in HBS-EP+ buffer (150 mM NaCl, 3 mM EDTA,

500 0.05% v/v surfactant P20, 10 mM HEPES pH 7.4). Positive hits from site scanning were selected for footprinting. 2 bp truncations were made from the 3' end of the selected fragments and similarly from the 5' of the inverted fragment. All ReDCaT SPR experiments were performed using a single Sensor Chip SA (GE Healthcare) on a Biacore 8K SPR system (Cytiva) as follows. For each channel FC_1 was designated as the reference (FC_{REF}) and FC_2 as the test cell (FC_{TEST}). DNA capture at 400 RUs was achieved in FC_{TEST} with a contact time of 60 seconds at a flow rate of 10 μl/min. Protein analyte (Purified CutR; 2 μM or 1 μM) was then added over both FCs for 60 seconds at a flow rate of 50 μl/min followed by a buffer

only dissociation of 60 seconds. Regeneration of the chip was achieved using Regeneration buffer (1 M NaCl, 50 mM NaOH) injected for 60 seconds at a flow rate of 10 µl/min over both FCs. All experiments were run in duplicate and data analysis was performed using the Biacore Insight Evaluation software (Cytiva) and Microsoft Excel.

Tandem Mass Tagging proteomic analysis

5 µl of spores of the relevant *S. venezuelae* strains were inoculated on cellophane-covered agar plates as triplicate spots in duplicate. After 2- or 9-day incubation at 30°C the mycelium was scraped off the cellophanes into a 15 ml falcon and resuspended in 10 ml cell lysis buffer (50 mM TEAB buffer pH 8.0, 150mM NaCl, 2% SDS, EDTA-free protease inhibitor, PhosSTOP phosphatase inhibitor). The suspension was disrupted via French press three times before being boiled at 100°C for 10 minutes. Samples were sonicated at 50 Hz four times for 20 seconds per cycle and then pelleted at 3,220 xg for 30 minutes. Protein concentration was determined using the BCA assay (Thermo Fischer Scientific) and 1 mg of protein from each sample transferred to a fresh 15 ml falcon tube. The proteins were precipitated from the supernatant with chloroform-methanol (46) and washed with acetone.

The air-dried pellets were dissolved in 50-100 µl of 2.5% sodium deoxycholate (SDC; Merck). After quantification by BCA assay, 100 µg of protein was reduced, alkylated, and digested with trypsin in the presence of 0.2 M EPPS buffer (Merck) and 2.5% SDC according to standard procedures. After digestion, the SDC was precipitated by adjusting to 0.2% TFA, and the clarified supernatant subjected to C18 solid phase extraction (SPE; OMIX tips; Agilent). Tandem mass tagging (TMT) labelling was performed using a Thermo TMT10plex™ kit (lot TL274393) according to the manufacturer's instructions with slight modifications; the dried peptides were dissolved in 90 µl of 0.2 M EPPS buffer/10% acetonitrile, and 200 µg TMT10plex reagent dissolved in 22 µl of acetonitrile was added. Samples were assigned to the TMT channels in an order avoiding channel leakage between different samples, if possible, as detailed by Brenes et. al (47). After labelling, aliquots of 1.7 µl from each sample were combined and analysed on the mass spectrometer (detailed below) to check labelling efficiency and estimate total sample abundances. The main sample aliquots were then combined correspondingly and desalted using a 50 mg C18 Sep-pak cartridge (Waters). The eluted peptides were dissolved in 500 µl of 25 mM NH_4HCO_3 and fractionated by high pH reversed phase HPLC. Using an ACQUITY Arc Bio System (Waters), the samples were loaded to a Kinetex® 5 µm EVO C18 100 Å LC Column 250 x 4.6 mm (Phenomenex). Fractionation was performed with the following gradient of solvents A (water), B (acetonitrile), and C (25 mM NH_4HCO_3) at a flow rate of 1 ml/min: solvent C was kept at 10% throughout the gradient; solvent B: 0-5 min: 5%, 5-10 min: 5-13% curve 5, 10-70 min: 13-40%, 70-75 min: 40-50%, 75-80%: 50-80%; followed by 5 min at 80% B and re-equilibration to 5% for 24 minutes. Fractions of 1 ml were collected, dried down, and concatenated to produce 20 final fractions for MS analysis. Aliquots of 30-40 µl were removed and stored at -80°C for

global proteome quantification. Aliquots of all concatenated fractions were analysed by nanoLC-MS/MS on an Orbitrap Fusion™ Tribrid™ mass spectrometer coupled to an

550 UltiMate® 3000 RSLCnano LC system (Thermo Fisher Scientific). The samples were loaded onto a trap column (nanoEase M/Z Symmetry C18 Trap Column, Waters) with 0.1% TFA at 15 µl/min for 3 minutes. The trap column was then switched in-line with the analytical column (nanoEase M/Z column, HSS C18 T3, 1.8 µm, 100 Å, 250 mm x 0.75 µm, Waters) for separation using the following gradient of solvents A (water, 0.1% formic acid) and B (80% acetonitrile, 0.1% formic acid) at a flow rate of 0.25 µl/min; for global protein analysis: 0-4 min 3% B (parallel to trapping); 4-17 min increase B curve 4 to 14%; 17-80 min linear increase B to 35%; 80-102 min linear increase B to 55% followed by a ramp to 99% B and re-equilibration to 3% B for 23 minutes; for enriched phosphopeptides: 0-4 min 3% B (parallel to trapping); 4-11 min increase B curve 4 to 12%; 11-52 min linear increase B to 43%; 52-61

560 min linear increase B to 55%, 61-64 min linear increase B to 65%, followed by a ramp to 99% B and re-equilibration to 3% B for 23 minutes. Data were acquired with the following parameters in positive ion mode: MS_1/OT: resolution 120K, profile mode, mass range m/z 400-1800, AGC target 2e5, max inject time 50 ms; MS_2/IT: data dependent analysis with the following parameters: top10 in IT Turbo mode, centroid mode, quadrupole isolation window 1.6 Da, charge states 2-5, threshold 1.9e4, CE=35 , AGC target 2e4, max. inject time 70 ms, dynamic exclusion 1 count/7 s with ±7 ppm, multistage activation on with neutral loss 97.98 Da (for phosphopeptides only); MS_3 synchronous precursor selection (SPS): 10 SPS precursors, isolation window 1.6 Da, HCD fragmentation with CE=65, Orbitrap resolution 50k, AGC target 1.5e5, max inject time 175 ms.

570 The acquired raw data were processed and quantified in Proteome Discoverer 3.1 (Thermo Fisher Scientific); all mentioned tools of the following workflow are nodes of the proprietary Proteome Discoverer (PD) software. The *Streptomyces venezuelae* (http://strepdb.streptomyces.org.uk/, 7420 entries) protein fasta database was imported into PD adding a reversed sequence database for decoy searches; a database for common contaminants (maxquant.org, 245 entries) was also included. For global protein quantification, the database search was performed using the incorporated search engine CHIMERYS (MSAID, Munich, Germany). The processing workflow also included spectrum selection, the TopN Peak Filter with 20 peaks per 100 Da, and the reporter ion quantification by most confident centroid (20 ppm). The inferys_3.0.0. fragmentation prediction model

580 was used with fragment tolerance of 0.5 Da, enzyme trypsin with 1 missed cleavage, variable modification oxidation (M), fixed modifications carbamidomethyl (C) and TMT10plex on N-terminus and K. Identifications were calculated for False Discovery Rate (FDR) 0.01 (strict) and 0.05 (relaxed). The results were exported into a Microsoft Excel table including data for normalized and un-normalized abundances, ratios for the specified conditions, the corresponding p-values and adjusted p-values, number of unique peptides, q-values, PEP-values, identification scores from both search engines; FDR confidence filtered

for high confidence (strict FDR 0.01) only. Further filtering included removal of contaminants and single unique peptide matches.

590 **Chemical extraction of chloramphenicol from solid media**
Two 5 μl aliquots of spore stock were used to inoculate cellophane-covered 30 ml agar plates with *Streptomyces* spp. before incubation at 30°C for 2 days. Subsequently the cellophane was removed. Agar was cut into 1 cm^3 pieces using a sterile razor blade and transferred to a 100 ml Duran bottle. 30 ml of HPLC-grade ethyl acetate was added to the agar, briefly shaken, and left at room temperature for 1 hour. The top 25 ml of the ethyl acetate was then transferred to a sterile 50 ml falcon tube and the sample was dried under vacuum. The sample was then resuspended in 500 μl HPLC-grade methanol and transferred to a 1.5 ml microcentrifuge tube. Insoluble particulates were removed via centrifugation at 15,871 xg for 10 minutes with the uppermost 200 μl of solution transferred to a 2 ml HPLC

600 vial containing a 200 μl glass insert for downstream chloramphenicol quantification.

Chloramphenicol Quantification by High-performance Liquid Chromatography
Analytical HPLC was performed on a 1290 Infinity II LC System (Agilent). A chloramphenicol standard curve was prepared in triplicate with serial dilutions of chloramphenicol in methanol from 1000 – 0.01 μg/ml. Chloramphenicol standards and extracted samples were chromatographed over a Kinetex 2.6 μm XB-C18 110 Å, 100 x 4.6mm column (Phenomenex) eluting with a linear gradient: mobile phase A, 0.1% (v/v) trifluoroacetic acid; mobile phase B, acetonitrile; 0 min, 10%B; 1 min, 10%B; 11 min, 100%B; 13 min, 100%B; 13.2 min, 10%B; 15 min, 10%B; flow rate 1 ml/min; injection volume 10 μL. UV absorbance was measured at

610 278 nm and the peak area at this wavelength was measured for analysis. The minimum concentration of chloramphenicol that could accurately be detected under these conditions was 0.5 μg/ml.

Quantitative reverse transcription PCR (qRT-PCR)
To perform quantitative RT-PCR (qRT-PCR), *S. venezuelae* wild-type, Δ*cutRS,* Δ*htrA3* and Δ*htrA3* + pIJ10257-*htrB* strains were grown on top of sterilized cellophane discs on YP or YPD agar at 30°C for 8-10 days. Colonies were removed from the cellophanes kept in liquid nitrogen whilst being crushed using a sterile pestle and mortar on dry ice. Crushed samples were resuspended in 1 ml RLT Buffer (Qiagen) supplemented with 1 % 2-mercaptoethanol.

620 This suspension was added to a QIA-shredder column (Qiagen) with the flowthrough transferred to a new 1.5 ml microfuge tube (leaving the pellet). Then, 700 μl acidic phenol:chloroform was added and the sample incubated at room temperature for 3 minutes and subsequently centrifuged at 15,871 xg. for 20 min. The resulting upper phase was mixed with 0.5 volumes 95 % ethanol. This was then applied to a RNeasy Mini spin column (Qiagen) and processed following the manufacturer's protocol. The eluant was treated with the Turbo-DNase kit (Invitrogen) followed by a RNeasy mini clean-up kit (Qiagen) both used according to manufacturer's instructions. The samples were then flash frozen in liquid

nitrogen for storage at −80°C. RNA was quantified with Nanodrop and Qubit Fluorometer and~1 mg converted to cDNA using the LunaScript RT SuperMix Kit (NEB) as per the manufacturer's instructions, including No-RT control reactions. Primers for each transcript were optimized using serial dilutions of template DNA and checked for efficiency by generating standard curves and calculating as follows: $E=(10slope−1)−1$. Primer sets with an efficiency between 90 and 110 % and within 5 % of each other were selected. qRT-PCR was run using the QuantStudio1 (Applied Biosystems) with Luna Universal qPCR Master Mix according to the manufacturer's guidelines for a 20 µl reaction with 2 µl template cDNA and a final concentration of 0.25 mM of each primer. Controls with no reverse transcriptase were run to ensure the absence of contaminating gDNA. ΔC_T values were normalized to the *hrdB* gene, which encodes the housekeeping RNA polymerase sigma factor.

Phylogenetic analysis

Genomes for use in phylogenetic analysis were obtained from the NCBI database(48), in all cases only complete and 'reference' RefSeq genomes were selected to avoid duplication. The accession numbers for genomes used to generate Figure 6A and 6B are given in Supplementary Information 1 and 2 respectively. Protein sequence alignments were carried out using MAFFT version 7 (mafft --memsavetree --retree 1 –treeout) (49, 50). Aligned sequences were used to produce a maximum likelihood tree using FASTTREE (FastTree -jtt alignment_file > output_tree) (51). The resulting trees were annotated in Interactive Tree Of Life (iTOL) v7 (52). Homologues of SvCutS and SvCssS were identified by reciprocal BLAST search, where reciprocal BLAST best hits were used in further analysis.

Protein Sequence Alignments

Protein sequences for alignment were obtained via reciprocal blast searches using *S. venezuelae* GCF_001886595.1 CutS and CssS as the query sequences. The sequences obtained for CutS and CssS homologues are available in Supplemental information 1.7 and 1.8 respectively. Sequences were aligned with Clustal Omega (53) and visualised in Jalview (54).

Data availability

The ChIP-seq data generated in this study have been deposited in the GEO database under the accession code GSE225370. The proteomics data generated in this study have been deposited in the ProteomeXchange database with identifier PXD051851. The genome sequence accession numbers and protein identifiers used to generate the data in Figure 7A and 7B available as Supplementary Information 1 and 2 respectively. Protein identifiers used to generate Supplementary Figure 2 and Supplementary Figure 4 are available as Supplementary Information .

REFERENCES

1. de Lima Procópio RE, da Silva IR, Martins MK, de Azevedo JL, de Araújo JM. 2012. Antibiotics produced by Streptomyces. The Brazilian Journal of Infectious Diseases 16:466–471.

2. Gavriilidou A, Kautsar SA, Zaburannyi N, Krug D, Müller R, Medema MH, Ziemert N. 2022. Compendium of specialized metabolite biosynthetic diversity encoded in bacterial genomes. Nat Microbiol 7:726–735.

3. Melnyk S, Stierhof M, Bratiichuk D, Fries F, Müller R, Rebets Y, Luzhetskyy A, Ostash B. 2025. Uncovering the genetic basis of antiviral polyketide limocrocin biosynthesis through heterologous expression. Microb Cell Fact 24:17.

4. Nguyen CT, Dhakal D, Pham VTT, Nguyen HT, Sohng JK. 2020. Recent advances in strategies for activation and discovery/characterization of cryptic biosynthetic gene clusters in streptomyces. Microorganisms 8.

5. McLean TC, Wilkinson B, Hutchings MI, Devine R. 2019. Dissolution of the Disparate: Co-ordinate Regulation in Antibiotic Biosynthesis. Antibiotics 8:83.

6. Cruz-Bautista R, Ruíz-Villafán B, Romero-Rodríguez A, Rodríguez-Sanoja R, Sánchez S. 2023. Trends in the two-component system's role in the synthesis of antibiotics by Streptomyces. Appl Microbiol Biotechnol 107:4727.

7. Burkovski A, Sprenger GA, Sánchez De La Nieta R, Santamaría RI, Díaz M. 2022. Two-Component Systems of Streptomyces coelicolor: An Intricate Network to Be Unraveled. International Journal of Molecular Sciences 2022, Vol 23, Page 15085 23:15085.

8. Cruz-Bautista R, Zelarayan-Agüero A, Ruiz-Villafán B, Escalante-Lozada A, Rodríguez-Sanoja R, Sánchez S. 2024. An overview of the two-component system GarR/GarS role on antibiotic production in Streptomyces coelicolor. Appl Microbiol Biotechnol 108:1–16.

9. Sánchez de la Nieta R, Santamaría RI, Díaz M. 2022. Two-Component Systems of Streptomyces coelicolor: An Intricate Network to Be Unraveled. Int J Mol Sci 23.

10. McLean TC, Lo R, Tschowri N, Hoskisson PA, Al Bassam MM, Hutchings MI, Som NF. 2019. Sensing and responding to diverse extracellular signals: An updated analysis of the sensor kinases and response regulators of streptomyces species. Microbiology (United Kingdom) 165:929–952.

11. Chang HM, Chen MY, Shieh YT, Bibb MJ, Chen CW. 1996. The cutRS signal transduction system of Streptomyces lividans represses the biosynthesis of the polyketide antibiotic actinorhodin. Mol Microbiol 21:1075–85.

12. The cutRS signal transduction system of Streptomyces lividans represses the biosynthesis of the polyketide antibiotic actinorhodin - PubMed. https://pubmed.ncbi.nlm.nih.gov/8885276/. Retrieved 26 June 2025.

13. McLean TC, Beaton ADM, Martins C, Saalbach G, Chandra G, Wilkinson B, Hutchings MI. 2023. Evidence of a role for CutRS and actinorhodin in the secretion stress response in Streptomyces coelicolor M145. Microbiology (United Kingdom) 169.

14. Jordan ML, Schlimpert S. 2025. Microbe Profile: Streptomyces venezuelae – a model species to study morphology and differentiation in filamentous bacteria. Microbiology (N Y) 171:001541.

15. Görke B, Stülke J. 2008. Carbon catabolite repression in bacteria: Many ways to make the most out of nutrients. Nat Rev Microbiol 6:613–624.

16. Romero-Rodríguez A, Rocha D, Ruiz-Villafán B, Guzmán-Trampe S, Maldonado-Carmona N, Vázquez-Hernández M, Zelarayán A, Rodríguez-Sanoja R, Sánchez S. 2017. Carbon catabolite regulation in Streptomyces: new insights and lessons learned. World J Microbiol Biotechnol 33.

17. Stevenson CEM, Lawson DM. 2021. Analysis of Protein–DNA Interactions Using Surface Plasmon Resonance and a ReDCaT Chip. Methods in Molecular Biology 2263:369–379.

18. Fernández-Martínez LT, Del Sol R, Evans MC, Fielding S, Herron PR, Chandra G, Dyson PJ. 2011. A transposon insertion single-gene knockout library and new ordered cosmid library for the model organism Streptomyces coelicolor A3(2). Antonie van Leeuwenhoek, International Journal of General and Molecular Microbiology 99:515–522.

19. Rojo F. 1999. Repression of Transcription Initiation in Bacteria. J Bacteriol 181:2987.

20. Bertrand-Burggraf E, Hurstel S, Daune M, Schnarr M. 1987. Promoter properties and negative regulation of the uvrA gene by the LexA represser and its amino-terminal DNA binding domain. J Mol Biol 193:293–302.

21. Liu B, Hong C, Huang RK, Yu Z, Steitz TA. 2017. Structural basis of bacterial transcription activation. Science (1979) 358:947–951.

22. Busby SJW, Browning DF. 2024. Transcription activation in Escherichia coli and Salmonella . EcoSal Plus https://doi.org/10.1128/ECOSALPLUS.ESP-0039-2020/SUPPL_FILE/ECOSALPLUS.ESP-0039-2020-S0001.DOCX.

23. Shi J, Wang L, Wen A, Wang F, Zhang Y, Yu L, Li F, Jin Y, Feng Z, Li J, Yang Y, Gao F, Zhang Y, Feng Y, Wang S, Zhao W, Lin W. 2022. Structural basis of three different transcription activation strategies adopted by a single regulator SoxS. Nucleic Acids Res 50:11359–11373.

24. Vicente RL, Gullón S, Marín S, Mellado RP. 2016. The three streptomyces lividans HtrA-Like proteases involved in the secretion stress response act in a cooperative manner. PLoS One 11.

25. Daniels R, Mellroth P, Bernsel A, Neiers F, Normark S, Von Heijne G, Henriques-Normark B. 2010. Disulfide bond formation and cysteine exclusion in gram-positive bacteria. Journal of Biological Chemistry 285:3300–3309.

26. Hutchings MI, Hong HJ, Buttner MJ. 2006. The vancomycin resistance VanRS two-component signal transduction system of Streptomyces coelicolor. Mol Microbiol 59.

27. McLean TC, Beaton ADM, Martins C, Saalbach G, Chandra G, Wilkinson B, Hutchings MI. 2023. Evidence of a role for CutRS and actinorhodin in the secretion stress response in Streptomyces coelicolor M145. Microbiology (United Kingdom) 169.

28. McLean TC, Lo R, Tschowri N, Hoskisson PA, Al Bassam MM, Hutchings MI, Som NF. 2019. Sensing and responding to diverse extracellular signals: an updated analysis of the sensor kinases and response regulators of Streptomyces species. Microbiology (Reading) 165:929–952.

29. Sánchez de la Nieta R, Antoraz S, Alzate JF, Santamaría RI, Díaz M. 2020. Antibiotic Production and Antibiotic Resistance: The Two Sides of AbrB1/B2, a Two-Component System of Streptomyces coelicolor. Front Microbiol 11.

30. Som NF, Heine D, Holmes NA, Munnoch JT, Chandra G, Seipke RF, Hoskisson PA, Wilkinson B, Hutchings MI. 2017. The conserved actinobacterial two-component system MtrAB coordinates chloramphenicol production with sporulation in Streptomyces venezuelae NRRL B-65442. Front Microbiol 8.

31. Hallgren J, Tsirigos KD, Damgaard Pedersen M, Juan J, Armenteros A, Marcatili P, Nielsen H, Krogh A, Winther O. 2022. DeepTMHMM predicts alpha and beta transmembrane proteins using deep neural networks. bioRxiv.

32. Daniels R, Mellroth P, Bernsel A, Neiers F, Normark S, Von Heijne G, Henriques-Normark B. 2009. Disulfide Bond Formation and Cysteine Exclusion in Gram-positive Bacteria. J Biol Chem 285:3300.

33. Noone D, Botella E, Butler C, Hansen A, Jende I, Devine KM. 2012. Signal Perception by the Secretion Stress-Responsive CssRS Two-Component System in Bacillus subtilis. J Bacteriol 194:1800.

34. Clausen T, Kaiser M, Huber R, Ehrmann M. 2011. HTRA proteases: Regulated proteolysis in protein quality control. Nat Rev Mol Cell Biol 12:152–162.

35. Reardon-Robinson ME, Ton-That H. 2016. Disulfide-bond-forming pathways in Gram-positive bacteria. J Bacteriol 198:746–754.

36. Viaene T, Langendries S, Beirinckx S, Maes M, Goormachtig S. 2016. Streptomyces as a plant's best friend? FEMS Microbiol Ecol 92:119.

37. Edwards MJ, White GF, Butt JN, Richardson DJ, Clarke TA. 2020. The Crystal Structure of a Biological Insulated Transmembrane Molecular Wire. Cell 181:665-673.e10.

38. Norman MP, Edwards MJ, White GF, Burton JAJ, Butt JN, Richardson DJ, Louro RO, Paquete CM, Clarke TA. 2023. A Cysteine Pair Controls Flavin Reduction by Extracellular Cytochromes during Anoxic/Oxic Environmental Transitions. mBio 14.

39. Yang X, Yang XA, Xu M, Zhou L, Fan Z, Jiang T. 2015. Crystal structures of YfiR from Pseudomonas aeruginosa in two redox states. Biochem Biophys Res Commun

820 461:14–20.

40. Feeney MA, Newitt JT, Addington E, Algora-Gallardo L, Allan C, Balis L, Birke AS, Castaño-Espriu L, Charkoudian LK, Devine R, Gayrard D, Hamilton J, Hennrich O, Hoskisson PA, Keith-Baker M, Klein JG, Kruasuwan W, Mark DR, Mast Y, McHugh RE, McLean TC, Mohit E, Munnoch JT, Murray J, Noble K, Otani H, Parra J, Pereira CF, Perry L, Pintor-Escobar L, Pritchard L, Prudence SMM, Russell AH, Schniete JK, Seipke RF, Sélem-Mojica N, Undabarrena A, Vind K, van Wezel GP, Wilkinson B, Worsley SF, Duncan KR, Fernández-Martínez LT, Hutchings MI. 2022. ActinoBase: tools and protocols for researchers working on Streptomyces and other filamentous

830 actinobacteria. Microb Genom 8.

41. Cobb RE, Wang Y, Zhao H. 2015. High-Efficiency Multiplex Genome Editing of *Streptomyces* Species Using an Engineered CRISPR/Cas System. ACS Synth Biol 4:723–728.

42. MacNeil DJ, Gewain KM, Ruby CL, Dezeny G, Gibbons PH, MacNeil T. 1992. Analysis of Streptomyces avermitilis genes required for avermectin biosynthesis utilizing a novel integration vector. Gene 111:61–68.

840 43. Larcombe DE, Braes RE, Croxford JT, Wilson JW, Figurski DH, Hoskisson PA. 2024. Sequence and origin of the Streptomyces intergenetic-conjugation helper plasmid pUZ8002. Access Microbiol 6:000808.v3.

44. Fernández-Martínez LT, Bibb MJ. 2014. Use of the Meganuclease I-SceI of Saccharomyces cerevisiae to select for gene deletions in actinomycetes. Sci Rep 4.

45. Stevenson CEM, Lawson DM. 2021. Analysis of Protein–DNA Interactions Using Surface Plasmon Resonance and a ReDCaT ChipMethods in Molecular Biology.

850 46. Wessel D, Flügge UI. 1984. A method for the quantitative recovery of protein in dilute solution in the presence of detergents and lipids. Anal Biochem 138.

47. Brenes A, Hukelmann J, Bensaddek D, Lamond AI. 2019. Multibatch TMT Reveals False Positives, Batch Effects and Missing Values. Molecular & Cellular Proteomics 18:1967–1980.

48. Sayers EW, Beck J, Bolton EE, Brister JR, Chan J, Comeau DC, Connor R, DiCuccio M, Farrell CM, Feldgarden M, Fine AM, Funk K, Hatcher E, Hoeppner M, Kane M, Kannan S, Katz KS, Kelly C, Klimke W, Kim S, Kimchi A, Landrum M, Lathrop S, Lu Z, Malheiro A, Marchler-Bauer A, Murphy TD, Phan L, Prasad AB, Pujar S, Sawyer A, Schmieder E, Schneider VA, Schoch CL, Sharma S, Thibaud-Nissen F, Trawick BW, Venkatapathi T, Wang J, Pruitt KD, Sherry ST. 2024. Database resources of the National Center for Biotechnology Information. Nucleic Acids Res 52:D33–D43.

49. Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Mol Biol Evol 30:772–780.

50. Katoh K, Misawa K, Kuma KI, Miyata T. 2002. MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res 30:3059–3066.

51. Price MN, Dehal PS, Arkin AP. 2009. FastTree: Computing Large Minimum Evolution Trees with Profiles instead of a Distance Matrix. Mol Biol Evol 26:1641–1650.

52. Letunic I, Bork P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree display and annotation tool. Nucleic Acids Res 52:W78–W82.

53. Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R, Butcher S. 2024. The EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024. Nucleic Acids Res 52:W521–W525.

54. Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview Version 2—a multiple sequence alignment editor and analysis workbench. Bioinformatics 25:1189–1191.

Acknowledgements

We would like to thank Professor Mervyn Bibb (John Innes Centre) for his contributions to the conceptualisation and generation of *S. venezuelae* strain M1702. We also thank past and present members of the Hutchings and Wilkinson groups (John Innes Centre) and Dr Alison MacFadyen (The Sainsbury Laboratory) and Professor Tracy Palmer (Newcastle University) for useful discussions. We thank John Innes Centre support services for growth media, glassware, plasticware and waste disposal and Phil Robinson for his photography expertise.

This work was supported by the BBSRC doctoral training programme grant BB/M011216/1 (T.C.M.); by the BBSRC-funded Institute Strategic Programme Project BBS/E/J/000PR9790 to the John Innes Centre and by the BBSRC-funded Institute Strategic Program Harnessing Biosynthesis for Sustainable Food and Health (HBio) grant BB/X01097X/1 to the John Innes Centre.

Author Contributions.

Conceptualization – T.C.M. and M.I.H.

Data curation – A.D.M.B., G.C., C.M. and G.S.

Formal analysis – T.C.M., A.D.M.B., G.C., C.M. and G.S.

Funding acquisition– M.I.H. and B.W.

Investigation – T.C.M., A.D.M.B., N.A.H., J.C., G.C., C.M. and G.S.

Methodology – T.C.M., A.D.M.B., G.C., C.M. and G.S.

Project administration – T.C.M., M.I.H. and B.W.

910 Resources – C.M., G.S., M.I.H. and B.W.

Software – G.C., C.M. and G.S.

Supervision – M.I.H. and B.W.

Validation – T.C.M. and A.D.M.B.

Visualization– T.C.M. and A.D.M.B.

Writing – original draft – T.C.M., A.D.M.B. and M.I.H.

Writing – review and editing – T.C.M., A.D.M.B., G.C., C.M., G.S., B.W. and M.I.H.

Figure 1. | Deletion of *cutRS* results in the overproduction of chloramphenicol. A. Antimicrobial overlay assays of the *S. venezuelae* wild-type, Δ*cutRS*, M1702 (Δ*cml*Δ*jad*) and M1702 Δ*cutRS* strains with *B. subtilis.* **B.** Antimicrobial overlay assays of *S. venezuelae* wild-type and Δ*cutRS* on YPD agar. (left) Overlays with the chloramphenicol-sensitive strain *B. subtilis* JH642. (right) Overlays with the chloramphenicol-resistant strain *B. subtilis* JH642 Cml[R] **C.** Quantification of chloramphenicol extracted from *S. venezuelae* wild-type and Δ*cutRS* strains on YP and YPD agar by HPLC.

Figure 2. | The CutRS two-component system is involved in colony growth and glucose consumption. A. *S. venezuelae* wild-type, Δ*cutRS and* Δ*cutRS* +pSS170-*cutRS* colonies grown on YP (- glucose) and YPD (+ glucose) agar for 10 days at 30°C. **B.** The colony dry weight (mg) of *S. venezuelae* wild-type and Δ*cutRS* after 10 days of growth at 30°C on YP, YPD, MYM and MYM+glucose. **C.** The remaining glucose concentration (mg/ml) in spent media after 10 days of growth at 30°C (*S. venezuelae* wild-type or Δ*cutRS*).

Figure 3. | The defined DNA binding site for the CutR response regulator. A. ChIP-seq peaks for biological replicates of CutR-3xFLAG (blue lines) and the wild-type control (grey lines). CutR binds upstream of the genes *htrA3*, *vnz_08815* and *htrB*. **B.** ReDCaT SPR was used to determine the binding site of 6xHis-CutR to the promoter regions of *htrA3* and the putative target gene *vnz_08815*. Sequential truncations of 2 bp were used to define the precise binding motif of TAWATAAA. The %R_{MAX} was determined using the molecular masses of the annealed oligo duplex and monomeric 6xHis-CutR. This binding motif was then used to identify the CutR binding site upstream of *htrB* (A).

Figure 4. | The deletion of *cutRS* causes a global shift in the proteome. Volcano plot of the Fold Change (Log_2) protein abundance detected by TMT-proteomics against the calculated adjusted P-value (Log_{10}) for *S. venezuelae* wild-type (wild-type) and Δ*cutRS* strains on YPD agar. The abundance of CutRS in the wild-type samples appears exaggerated due to the data processing against the Δ*cutRS* samples and does not represent normal levels.

Figure 5. | The conserved extracellular cysteine residues control the activity of CutS. A. The Δ*cutRS* mutant can be partially complemented by *in trans* expression of an operon encoding wild-type CutR and a CutS protein in which the extracellular cysteines are altered to serine (CutS(C85S,C103S)). **B.** Quantitative RT-PCR analyzing the abundance of *htrA3* (light grey bars) and *htrB* (dark grey bars) transcripts normalized to constitutively expressed *hrdB* and compared to wild-type. Error bars represent standard deviation across replicates.

Figure 6. | The *S. venezuelae* wild-type phenotype is recovered by deletion of *cssRS* in the Δ*cutRS* background. *S. venezuelae* wild-type, Δ*cutRS*, Δ*cssRS and* Δ*cutRS* Δ*cssRS* strains grown on YP (- glucose) and YPD (+ glucose) agar for 10 days at 30°C.

A

% of SK EC Domains with ≥ 2 Cysteines

- 0
- 10
- 20
- 30
- 40
- 50
- 60
- 70
- 80
- 90
- 100

Class

- Deinococci
- Alphaproteobacteria
- Betaproteobacteria
- Actinomycetes
- Desulfovibrionia
- Gammaproteobacteria
- Flavobacteria
- Spirochaetia
- Bacilli
- Desulfuromonadia
- Opitutia
- Epsilonproteobacteria
- Clostridia
- Cytophagia
- Deltaproteobacteria
- Bacteroidia
- Cyanophyceae
- Sphingobacteriia
- Saprospiria
- Erysipelotrichia
- Chitinophagia
- Fusobacteriia
- Negativicutes
- Thermotogae
- Planctomycetia
- Tissierellia
- Anaerolineae
- Terriglobia
- Aquificae
- Verrucomicrobiia
- Thermodesulfovibrionia
- Lentisphaeria
- Coriobacteriia
- Armatimonadia
- Oligoflexia
- Desulfobulbia
- Mollicutes
- Chlamydiia
- Desulfobacteria
- Acidimicrobiia
- Synergistia

B

Outer Ring Labels

- Streptomyces
- CutS RBH
- CutS RBH 2+ EC Cysteines
- CusS RBH
- CusS RBH 2+ EC Cysteines

Figure 7. | Phylogenetic and cysteine residue analysis of bacterial sensor kinases. A. Maximum likelihood phylogenetic tree of 348,633 predicted SKs from 12,799 high-quality bacterial genomes. The outer ring is color-coded by bacterial class, as indicated in the legend. The heatmap overlay in the inner ring shows the proportion of SKs with two or more cysteine residues in their extracellular domains, as predicted by DeepTMHMM. Dark blue indicates 0% of the SKs in that genome can form a disulphide bond in their extracellular domain, increasing to dark red, which indicates 100% of the SKs in that genome can form a disulphide bond in their extracellular domain. **B.** Maximum likelihood phylogenetic tree based on RpoB protein sequences from 2,940 bacterial reference genomes. The outer rings represent the following: Genomes in the *Streptomyces* genus (light green); CutS reciprocal BLAST best hits (RBBH) (dark blue); CutS homologs with two or more cysteine residues in their extracellular domains, as predicted by DeepTMHMM (light blue); CssS reciprocal BLAST best hits (RBBH) (dark pink); CssS homologs with two or more cysteine residues in their extracellular domains, as predicted by DeepTMHMM (light pink). Trees were constructed using MAFFT for sequence alignment of RpoB protein sequences and FASTTREE for maximum likelihood inference and visualised in iTOL.

Figure 8. | A proposed model for the extracellular redox-sensing activity of the CutRS system in _Streptomyces_ spp. When the extracellular space is conducive to protein folding (left side) the cysteine residues in the extracellular sensor domain of the SK CutS (pink) form a disulphide bond. In this state the CutRS system is inactive, _htrA3_ is expressed and _htrB_ is repressed. When the extracellular space in not conducive to protein folding (right side) the disulphide bond in the extracellular sensor domain of CutS is unable to form. In this state the SK CutS is activated, autophosphorylates and transfers this phosphate to the cognate RR CutR (light green). Phosphorylated CutR homodimerizes and binds DNA, repressing the transcription of _htrA3_ and activating _htrB_. These genes encode dual function chaperone / proteases which can act to restore the cell to homeostasis.

1000

SUPPLEMENTARY FIGURES

1010 **Supplementary Figure 1. | Overexpression of *htrB* in the *S. venezuelae* Δ*htrA3* background is unable to reproduce the Δ*cutRS* phenotype. A.** *S. venezuelae* wild-type, Δ*htrA3* and Δ*htrA3* with *htrB* overexpression colonies grown on YP (- glucose) and YPD (+ glucose) for 10 days at 30°C.

Supplementary Figure 2. | Residue conservation in the extracellular sensor domain of CutS. Clustal OMEGA alignment of 105 CutS homologous from *Streptomyces* spp. showing two highly conserved cysteine residues in the extracellular sensor domain. Alignment view taken from Jalview giving the alignment conservations, quality, consensus and occupancy over one section of the alignment. Sequences use to generate this figure can be found as Supplementary File 3.

1020

Supplementary Figure 3. | CutS is predicted to form disulphide bonds between the two cysteine residues present in the extracellular sensor domain. A. AlphaFold2-predicted structure of homodimeric *S. venezuelae* CutS. The extracellular sensor domain, transmembrane domain and DHp / catalytic domain are annotated. (Inset) Close-up view of the predicted extracellular sensor domain highlighting the intra-chain disulphide bonds and predicted bond lengths. Chain A: green; Chain B: grey. **B.** AlphaFold2-predicted structure of homodimeric *S. venezuelae* CutS coloured by confidence score (pIDDT): very high (pIDDT > 90; dark blue), confident (90 > pIDDT > 70; light blue), low (70 > pIDDT > 50; yellow) and very low (pIDDT < 50; red).

1030

Supplementary Figure 4. | Residue conservation in the extracellular sensor domain of CssS of _Streptomyces_ spp. Clustal OMEGA alignment of 105 CssS homologous from _Streptomyces_ species showing two highly conserved cysteine residues in the extracellular sensor domain. Alignment view taken from Jalview giving the alignment conservations, quality, consensus and occupancy over one section of the alignment. Sequences use to generate this Figure can be found as Supplementary File 3

1040