Title: Type I Interferon production in myeloid cells is regulated by factors independent of *Ptpn22*

<u>Running title:</u> *Ptpn22* does not regulate IFN-I production

Jenna R. Barnes, Anam Fatima Shaikh, Alec M. Bevis, Tammy R. Cockerham, Robin C. Orozco

Department of Molecular Biosciences, University of Kansas, Lawrence, KS 66045 United States

of America

Corresponding Author: Robin C. Orozco, 1200 Sunnyside Dr, University of Kansas, Lawrence,

KS, 66045, United States of American, <u>orozco.robin@ku.edu</u>

Keywords (3-5): type I Interferon, Ptpn22, myeloid cells, autoimmunity-associated allele

ABSTRACT

The immune regulatory gene *PTPN22* is expressed in all immune cells and encodes Lyp in humans and the ortholog PEP in mice. The *PTPN22* alternative allele, 1858C>T, is expressed in 5-15% of the North American population and is strongly associated with the development of autoimmune disease while simultaneously capable of providing protection during virus infection and cancer. In murine models, significant progress has been made in elucidating the molecular mechanisms that PEP and its pro-autoimmune variant (PEP-R619W) modulate T cell function, yet their influence on non-T cell pathways, such as antigen presenting cell cytokine production, remains less defined. Previously, it was reported that PEP promotes type I interferon (IFN-I) production in dendritic cells (DCs) and macrophages following TLR4 stimulus. Here, we show that contrary to previous results, both PEP-WT and the PEP-R619W variant do not promote IFN-I production in DCs and macrophages following exposure to LPS, 3p-hpRNA, or coronavirus MHV A59. We attribute the prior findings to mouse strain-specific differences and conclude that something other than PEP may be regulating IFN-I production in these studies. We further show that PEP and its R619W variant distinctly modulate the production of TNFα, IL-12 and IL-2, in DCs following LPS stimulus. Taken together, our results challenge the current understanding of PEP's role during inflammation while providing new insight into how the PEP-R619W variant may alter myeloid cell function during disease.

INTRODUCTION

Allelic variation in immune-regulatory genes can significantly impact disease outcomes [1, 2]. One such variant of interest is the 1858C>T allele of the Protein Tyrosine Phosphatase Non-Receptor Type 22 gene (*PTPN22*). *PTPN22* is constitutively expressed in all immune cells and encodes for the phosphatase lymphoid protein Lyp, and its murine ortholog, PEST-domain enriched phosphatase (PEP). The *PTPN22* 1858C>T allele, resulting in an amino acid substitution of arginine (R) to tryptophan (W) at position 620 in human, is expressed in 5-15% of the North American population and is considered the highest non-HLA risk allele for the development of autoimmunity, including type I diabetes, rheumatoid arthritis, and systemic lupus erythematous[3-11].

To investigate the role of *PTPN22* and its autoimmunity-associated variant during inflammation, researchers have employed mouse models which lack expression of PEP (PEP-null), the murine ortholog of Lyp. Through these studies, it was shown that PEP-null mice have enhanced persistent viral clearance and T cell function[12, 13]. However, because the LYP-R620W, or the murine equivalent PEP-R619W mutation, does not cause a loss of protein expression, but rather altered function[14, 15], using PEP-null models to investigate how the *PTPN22* autoimmunity associated allele drives inflammation is not always appropriate. Furthermore, PEP-null and PEP-R619W mice have been reported to exhibit differing phenotypes [16-20]. As an example, PEP-null mice exhibit an increased frequency and potency of Tregs compared to PEP-WT mice, but this phenotype is not observed in PEP-R619W mice [19, 21, 22]. Using CRISPR-Cas9 generated PEP-R619W mice, we were the first to report that the PEP-R619W variant is similarly protective

during chronic virus infection and promotes enhanced T cell function [1]. Through adoptive

transfer studies, we and others showed that the enhanced antiviral T cell function in PEP-null and

PEP-R619W mice is due to both T cell intrinsic and extrinsic mechanisms [1-3]. However,

additional studies are needed to understand the specific antiviral molecular mechanisms that PEP

and the PEP-R619W variant may influence. Significant progress has been made in elucidating

the molecular mechanisms that PEP and its R619W variant modulate T cell function [12-17, 19, 23-28].

However, the role of PEP and its R619W variant on non-T cell pathways that may influence T

cell function, such as antigen presenting cell (APC) cytokine production, remains less

understood.

APCs, such as dendritic cells (DCs), produce a variety of cytokines capable of broadly

promoting inflammation or directing lymphocyte activation and differentiation [4-6]. Type-I

interferons (IFN-I) are potent cytokines produced by DCs which exhibit significant antiviral

activity by modulating the activation and effector function of other immune cells, including T

cells [29, 30]. Although a short burst of IFN-I secretion is critical for effective clearance of viruses

including Lymphocytic Choriomeningitis virus clone 13 (LCMV-cl13)[31-33], prolonged IFN-I

production has been shown to contribute to immune dysfunction, such as T cell exhaustion, and

persistent viral infection [31, 34]. Thus, tight regulation of IFN-I and other pro-inflammatory

cytokines during infection is crucial and mechanisms which alter the production of cytokines

have the potential to significantly impact the outcome of infection. Given the versatile role of

PEP and its R619W variant during a variety of inflammatory processes, researchers have set out

to evaluate the potential impact of PEP and its R619W variant on the production of cytokines by

APCs during infection.

Previously, it was reported that PEP promotes IFN-I production in DCs and macrophages

following stimulus with bacterial lipopolysaccharide (LPS) or Poly(I:C)), Toll Like Receptor 4

(TLR4) and Toll Like Receptor 3 (TLR3) agonists, respectively [35]. However, these studies were

performed using PEP-null mice available at The Jackson Labs which have been found to have

numerous SNPs associated with C57BL/6N (B6N) mice [36]. The recommended control for this

PEP-null strain are C57BL/6J PEP-WT mice and historically, the majority of *Ptpn22* literature in

C57BL/6 mice has also utilized Jackson Labs PEP-null mice with C57BL/6J PEP-WT mice as

controls. This poses concern that our current understanding of *Ptpn22* and how it regulates

inflammation may have been influenced by mouse strain specific differences driven by genetic

variation. To enable a more rigorous investigation into defining how *PTPN22* and its

autoimmunity associated allele regulate inflammation, our research team employs C57BL/6J

CRISPR-Cas9-generated PEP-null and PEP-R619W mouse models. Using these genetic models,

we first set out to determine if previous data regarding the impact of PEP on IFN-I production

could be recapitulated. We also further investigated the impact of the PEP-R619W variant on

IFN-I production in dendritic cells. In this study, we report that neither PEP nor the PEP-R619W

variant regulate IFNβ production in DCs or macrophages following LPS or viral RNA

stimulation. However, we found that both PEP and PEP-R619W differentially modulate the

production of other pro-inflammatory cytokines by dendritic cells, including IL-2, IL-12, and

TNFα.

RESULTS

Strain-specific differences impact IFNβ production in PEP-null bone marrow dendritic cells (BMDCs)

Previously, it was shown that PEP-null BMDCs and bone marrow macrophages (BMMs) produce reduced levels of IFN-I when stimulated with bacterial lipopolysaccharide (LPS) or Poly(I:C)), TLR4 and TLR3 agonists respectively [35]. Using our C57BL/6J CRISPR-Cas9 generated PEP-null model (PEP-null CRISPR) and the commercially available Jackson Labs PEP-null mice (PEP-null JAX), we tested if the B6N SNPs present in the Jackson Labs PEP-null strain were driving this reported phenotype, as opposed to PEP expression. Using qRT-PCR, we confirmed that our CRISPR-Cas9 generated PEP-null mice do not express quantifiable amounts of *Ptpn22* mRNA (Figure 1A). Using FLT3-ligand differentiated BMDCs from C57BL/6J PEP-WT, CRISPR-Cas9 generated PEP-null, and PEP-null Jackson Labs mice we measured for IFNβ production following LPS stimulation. While IFNβ concentration was comparable between PEP-WT and our PEP-null CRISPR mice, BMDCs from the commercial PEP-null JAX strain showed significantly less IFNβ production (Figure 1B). These data successfully recapitulate prior results, but through the use of our CRISPR-Cas9 generated PEP-null model, we observe that PEP does not promote IFNβ production. These strain-specific differences in IFN-I production suggest that other genetic differences associated with the B6N background, rather than *Ptpn22* deficiency alone, may account for the previously reported phenotype.

In addition to the PEP-null JAX model, studies have also utilized chemical inhibition of PEP to evaluate its role in IFN-I production and activity [28, 35]. However, varying results regarding PEP inhibitors and their specificity have been reported [37]. To evaluate the specificity of the PEP inhibitor LTV-1 during IFN-I production, we stimulated PEP-WT and CRISPR-Cas9 generated

PEP-null bone marrow macrophages (BMMs) with LPS in the presence or absence of LTV-1 to

inhibit PEP's enzymatic activity. We observed reduced IFNβ production in both PEP-WT and

PEP-null BMMs treated with LTV-1, but no significant differences in IFNβ production between

the two genotypes (Figure 1C). This data suggests that LTV-1 has off target effects which

reduces IFNβ production, rather than the loss of PEP's enzymatic activity.

PEP and PEP-R619W do not mediate IFNβ production in myeloid cells

Our initial results showed comparable IFNβ production between PEP-WT and PEP-null

CRISPR BMDCs following TLR4 stimulation via LPS. However, beyond TLRs, additional

pattern recognition receptors (PRRs) such as RIG-I Like Receptors (RLRs) induce large

quantities of IFN-I in response to virus infection [38-40]. To determine if PEP promotes IFN-I

production induced by TLR and RLR signaling in macrophages, we cultivated PEP-WT and

PEP-null BMMs and then stimulated them with LPS or infected with the murine coronavirus

Murine Hepatitis Virus Strain A59 (MHV A59), a MDA5 agonist [40]. There were no differences

detected in IFNβ production between PEP-null and PEP-WT BMMs following LPS exposure or

MHV A59 infection (Figure 2A). These results suggest that PEP does not promote IFN-I

production in BMMs following TLR4 or MDA5 activation.

Further, it is unknown if the autoimmunity-associated PEP-R619W variant, which alters

protein function rather than expression, uniquely affects IFN-I production compared to PEP-WT.

To determine if PEP-R619W either phenocopies PEP-null CRISPR DCs or distinctly alters IFN-I

production following TLR or RLR activation, we cultivated FLT3-ligand (FLT3-L) BMDCs

from C57BL/6J PEP-WT, PEP-null, and PEP-R619W mice and stimulated them with LPS or 5'

Triphosphate hairpin ribonucleic acid (3p-hpRNA), a RIG-I agonist. We detected no significant

differences in IFNβ production between all three genotypes (PEP-WT, PEP-null, and PEP-

R619W) following either LPS or 3p-hpRNA stimulation (Figure 2 B-C). Taken together, these

results indicate that neither PEP nor the PEP-R619W variant modulate IFNβ production in

dendritic cells mediated by TLR4 or RIG-I activation.

PEP and the PEP-R619W variant distinctly modulate the production of pro-inflammatory

cytokines in dendritic cells

Our findings show that PEP nor the PEP-R619W variant impact IFN-I production in DCs

and macrophages during TLR4, MDA5, or RIG-I stimulation. However, DCs produce a variety

of other cytokines which can promote widespread inflammation or influence T cell activation

and differentiation. Considering that the PEP-R619W variant promotes non-T cell mechanisms

which enhance T cell function during virus infection, it is possible that PEP regulates the

production of other inflammatory cytokines relevant to T cell function. Further, the PEP-R619W

variant may distinctly alter these regulatory mechanisms. Previous studies which have

investigated the role of PEP and its variant on cytokine production in myeloid cells have reported

varied findings[35, 41, 42]. Using PEP-R619W knock-in mouse models, one group reported increased

IL-12 production in Granulocyte-macrophage colony stimulating factor (GM-CSF) differentiated

PEP-R619W BMDCs post LPS exposure[41]. Another group reported that GM-CSF differentiated

PEP-null Jax BMDCs exhibited enhanced secretion of IL-6, IL-8 and TNFα compared to PEP-

WT following treatment with muramyl dipeptide (MDP)[42]. They also reported that the loss of

PEP reduced while the PEP-R619W variant increased IL-1β secretion by BMDCs following

ultra-pure LPS exposure[20]. However, a different group later reported no differences in TNFα, IL-

1β, or IL-12 production following treatment with LPS between FLT3-ligand differentiated PEP-

WT and PEP-null JAX BMDCs [35]. These contrary results align with previous reports of PEP-null

and PEP-R619W mice exhibiting different phenotypes but could also be attributed to the usage

of different mouse models, type of stimulus, and selected assays. Using a consistent genetic

model and experimental design, we set out to investigate the role of PEP and the PEP-R619W

variant in modulating cytokine production. To accomplish this, we again employed FLT3-ligand

differentiated BMDCs from PEP-WT, PEP-null CRISPR, and PEP-R619W mice. BMDCs were

stimulated with LPS, and the supernatant was collected and analyzed for TNFα, IL-12 (p70), and

IL-2 production. We found that compared to PEP-WT, TNFα production was reduced in both

PEP-null and PEP-R619W DCs (Figure 3A). Additionally, we observed a reduction in IL-12

production by PEP-R619W DCs compared to PEP-WT and PEP-null (Figure 3B). Finally, we

observed increased IL-2 production by both PEP-null and PEP-R619W DCs compared to PEP-

WT (Figure 3C). Taken together, these findings suggest that PEP and the PEP-R619W variant

distinctly modulate cytokine production in dendritic cells.

DISCUSSION

The *PTPN22* 1858C>T allele is strongly associated with the development of

autoimmunity, but the murine equivalent mutation has also been shown to be protective during

virus infection and cancer [16, 21]. While the impact of the PEP-R619W variant is well-defined in T

cells, the molecular mechanisms by which PEP and its variant impact the function of other

immune cells, such as the ability of APCs to produce pro-inflammatory cytokines, remain less

understood. Previously, it was suggested that PEP promotes IFN-I production in macrophages

and dendritic cells [35]. However, these data were generated using PEP-null mice on a B6N/B6J

mixed background. In this paper, we tested if the B6N SNPs present in this PEP-null model were

driving this reported phenotype by comparing BMDCs derived from our C57BL/6J CRISPR-

Cas9-generated PEP-null model to BMDCs from the B6N/B6J PEP-null mice available at the

Jackson labs. While the PEP-null JAX BMDCs had reduced IFN-I production, we did not detect

a difference between our PEP-null CRISPR BMDCs and C57BL/6J PEP-WT BMDCs (Figure

1B). These strain-specific differences in IFN-I production recapitulate previous findings, while

also revealing that unknown genetic variation in the B6N/B6J PEP-null mice may have

influenced the previously observed phenotype. This interpretation is further supported by reports

of reduced inflammatory responses during virus infection in B6N mice compared to C57BL/6J

mice [43, 44]. However, our study results do not address why transgenic PEP-null JAX BMMs

which over-express the human protein Lyp have restored IFNβ production [35]. Identifying the

mechanism which drives the reduced IFNβ production observed in the B6N/B6J Jackson lab's

available PEP-null mice but not in the CRISPR-Cas9 generated PEP-null mice will require

further investigation. Regardless, we are not the first to comment on or identify discrepancies

regarding previous data reported on PEP's role during IFN-I production in myeloid cells.

However, these prior publications have largely focused on differences identified in human

samples, as well as in the context of systemic lupus erythematous (SLE) [45-49]. Thus, to the best of

our knowledge, we are the first to show that in a mouse model there are strain specific

differences affecting IFN-I production by DCs and BMMs.

Next, when evaluating the specificity of the PEP inhibitor LTV-1 during IFN-I production

in BMMs, we found reduced IFNβ production in both PEP-WT and PEP-null macrophages

treated with LTV-1 (Figure 1C). These findings suggest that the observed reduction of IFNβ may

not be due to the deficiency of PEP nor the inhibition of its enzymatic function, but rather

because of non-specific effects of LTV-1. Taken together, our findings reveal additional

confounding variables that could impact the results of studies which aim to discern the role of

PEP during inflammation through the use of either genetic models or chemical inhibition. Our

observations add to the numerous conflicting findings already identified in the field regarding

PEP, PEP-R619W, and their effects on cytokine production [20, 35, 41, 42]. These differences likely

stem from genetic variation present in both human and animal studies, as well as the utilization

of diverse experimental approaches including over-expression, transgenic models, knockdown

techniques, and chemical inhibition. Each of these systems offer distinct advantages yet also

varying limitations, yielding inconsistent results. To address this lack of clarity, our lab employs

CRISPR/Cas9-generated PEP-null and PEP-R619W mouse models on a C57BL/6J background.

By using this model, we can precisely evaluate the mechanisms in which PEP and the PEP-

R619W variant drive inflammation while minimizing confounding genetic variables.

Collectively, our results indicate that IFNβ production by DCs and macrophages is not

promoted by PEP following TLR4 activation (Figure 1B, 2A, and 2B). Because IFN-I is induced

by additional PRRs such as RLRs, we also investigated if PEP regulates IFN-I production

induced by RLR signaling but observed no significant differences in IFNβ production between

PEP-WT and PEP-null macrophages or dendritic cells following MHV A59 infection or viral

RNA stimulus (Figure 2 A,C). Finally, since the PEP-R619W variant still has PEP expression,

but causes altered protein binding capabilities, it is possible that PEP-R619W DCs could have

distinctive changes in IFN-I production compared to PEP-null cells. Our results suggest that

similar to PEP-WT, PEP-R619W does not promote IFNβ production in dendritic cells post LPS

or viral RNA stimulus (Figure 2 B-C).

Although our findings indicate that PEP and PEP-R619W do not promote IFNβ

production in myeloid cells, we found that PEP and its variant distinctly modulate the production

of other pro-inflammatory cytokines. More specifically, we found that compared to PEP-WT,

TNFα production was reduced in both PEP-null and PEP-R619W DCs. Additionally, we

observed a reduction in IL-12 production by PEP-R619W DCs compared to PEP-WT and PEP-

null (Figure 3). Previously it was reported that PEP-R619W GM-CSF differentiated DCs have

increased IL-12 production compared to PEP-WT [41]. These conflicting findings may be due to a

variety of factors such as differences in the PEP-R619W mouse models and LPS dosage utilized,

as well as functional differences between FLT-ligand differentiated and GM-CSF differentiated

BMDCs. Finally, we observed increased IL-2 production by both PEP-null and PEP-R619W

DCs compared to PEP-WT. Since DC derived IL-2 can stimulate the growth and activation of T

cells, it is possible that the increased IL-2 produced by PEP-null and PEP-R619W DCs is a

contributing factor to the enhanced T cell function observed during virus infection.

In summary, PEP-R619W DCs have reduced TNFα and IL-12, but increased IL-2

production compared to PEP-WT. However, the molecular mechanisms by which PEP and its

variant distinctly modulate the production of pro-inflammatory cytokines in dendritic cells

remain unknown. Previously, PEP has been shown to bind to the E3 ubiquitin ligase, TNF

receptor-associated factor 3 (TRAF3) [35]. TRAF proteins are essential components of cell

signaling pathways and TRAF3 is uniquely capable of both positively regulating IFN-I

production and negatively regulating the canonical and non-canonical signaling pathways of

Nuclear Factor-kappa B (NF-κB) [50]. Based on prior study results [35], the interaction between PEP

and TRAF3 is necessary for the induction of IFN-I. However, our data suggests that PEP is not

required for the production of IFN-I. Instead, we report that PEP and the PEP-R619W variant

distinctly modulate the production of other pro-inflammatory cytokines produced downstream of

NF-κB signaling. A possible explanation for these observations is that PEP regulates cytokine

production by interacting with TRAF3 or other key signaling proteins involved in the NF-κB

signaling cascade, and the R619W mutation potentially alters these interactions. The impact of

the PEP-R619W variant on the production of cytokines critical for T cell activation, such as IL-2

and IL-12, may begin to further explain how the PEP-R619W variant promotes T cell extrinsic

effects which ultimately lead to enhanced immune function and viral clearance. However,

additional rigorous studies will be required to tease out the precise mechanisms in which PEP

and its PEP-R619W variant regulate these inflammatory processes in DCs and macrophages.

Altogether, our findings challenge the current understanding of PEP while providing new

insight into how the PEP-R619W variant is capable of simultaneously promoting autoimmunity

and conferring protection against persistent viral infection and cancer. While our results reveal

that changes in IFN-I production are not a key factor in how PEP regulates inflammation as

previously regarded, our data indicates that PEP and the PEP-R619W variant have a novel role in

modulating the production of other cytokines by dendritic cells which are critical for a variety of

inflammatory processes. Future studies will need to evaluate how changes mediated by PEP and

the PEP-R619W variant in dendritic cells and other antigen presenting cells are relevant in order

to accurately define the broad consequences they have on disease pathogenesis.

MATERIAL AND METHODS

Ethics:

All animal studies were reviewed and approved by University of Kansas Animal Care

and Use Committee (IACUC) (protocol number: 278-01).

Mice:

Mice were housed in general housing conditions and all animal studies were reviewed

and approved by the University of Kansas Animal Care and Use Committee (protocol number:

278-01). Both males and females ranging from 5-12 weeks of age were used in this study. No

sex-based differences were observed in our study results. C57BL/6J WT (PEP-WT) mice were

originally purchased from Jackson labs, then bred and maintained in the University of Kansas

Animal Care Unit. Ptpn22 $^{-/-}$ (PEP-null CRISPR) mice were generated using CRISPR/Cas9

technology on a C57BL/6 background as described previously [7]. Ptpn22 $^{-/-}$ mice (PEP-null

JAX) were gifted from Dr. Won Jin Ho (John Hopkins University School of Medicine). These

mice are available for purchase from the Jackson Laboratory, and their generation has been

previously described [23]. Mice expressing the Ptpn22 pro-autoimmune allele (PEP-R619W) were

generated using CRISPR/Cas9 technology on a C57BL/6 background using methods previously

reported [16, 21, 51].

Bone Marrow Dendritic Cell (BMDC) Culture:

Bone marrow cells were isolated from PEP-WT, PEP-null, and PEP-R619W mouse femurs. Following isolation, bone marrow cells were cultured in Advanced DMEM containing 10% FBS, 1% penicillin streptomycin, 1% L-Glutamine, and 100ng/mL FLT3L (Stem Cell Technologies) for 8 days. On day 8, FLT3-ligand differentiated dendritic cells were harvested, counted, and re-plated.

BMDC Stimulation:

BMDCs were resuspended and plated at 1×10^6 cells/mL in Advanced DMEM (Gibco) containing 10% FBS, 1% penicillin streptomycin and 1% L-Glutamine. The cells were stimulated with either LPS or 3p-hpRNA. Cells stimulated with LPS (500 ng/mL) (Sigma) were incubated overnight (16-18 hours). The supernatant and cells (resuspended in TRIzol (Invitrogen)) were collected and stored frozen at -81 degrees Celsius for future assays. Cells stimulated with 3p-hpRNA (2 ng/mL) complexed with LyoVec (InvivoGen) were incubated overnight (16-18 hours). The supernatant and cells (resuspended in TRIzol (Invitrogen)) were collected and stored frozen at -81 degrees Celsius for future assays.

Enzyme-Linked Immunosorbent Assay (ELISA):

Levels of IFNβ, TNFα, IL-12 (p70), and IL-2 in cell culture supernatant were measured using ELISA Kits (IFNβ (PBL Assay Science), TNFα, IL-12 and IL-2 (BioLegend)) according to the manufacturer's instructions, followed by analysis with a BioTek EON microplate reader at 450 and/or 570 nm.

qRT-PCR:

RNA was extracted using TRIzol according to the manufacturer's instructions

(Invitrogen). cDNA was synthesized using the High-Capacity cDNA Reverse Transcription Kit

(Applied Biosystems), according to the manufacturer's instructions. Real-time PCR was

performed on a Quant Studio 3 system using PowerTrack™ SYBR™ Green Master Mix

(Applied Biosystems). Each reaction was measured at least triplicate, and data was normalized to

the expression levels of the house keeping gene GAPDH. The designed primers for each selected

murine gene are listed:

Gene	Forward Primer	Reverse Primer
Gapdh	AACGACCCCTTCATTGACCTC	ACTGTGCCGTTGAATTTGCC
Ptpn22	AGCTGATGAAAATGTCCTATTGTGA	GTCCCACTGCATTCTGGTGA

Statistical analysis and graphing:

All statistical analysis was performed using GraphPad Prism Software. The type of

statistical test is listed in figure legends. Data was considered statistically significant if the p

value<0.05. Graphs were made in GraphPad Prism (La Jolla, CA). Figure legends indicate if data

shown is pooled from multiple studies or is from representative study.

ACKNOWLEDGEMENTS

353   **CONFLICTS OF INTEREST**

354   None Declared

REFERENCES

1.   Conway JR, Kofman E, Mo SS, Elmarakeby H, Van Allen E. Genomics of response to immune checkpoint therapies for cancer: implications for precision medicine. Genome medicine. 2018;10(1):93-. doi: 10.1186/s13073-018-0605-7. PubMed PMID: 30497521.

2.   Sivapalan L, Anagnostou V. Genetic variation in antigen presentation and cancer immunotherapy. Immunity. 2022;55(1):3-6. doi: 10.1016/j.immuni.2021.12.010. PubMed PMID: 35021056.

3.   Metzler G, Dai X, Thouvenel CD, Khim S, Habib T, Buckner JH, Rawlings DJ. The Autoimmune Risk Variant PTPN22 C1858T Alters B Cell Tolerance at Discrete Checkpoints and Differentially Shapes the Naive Repertoire. Journal of immunology (Baltimore, Md : 1950). 2017;199(7):2249-60. Epub 2017/08/11. doi: 10.4049/jimmunol.1700601. PubMed PMID: 28801357.

4.   Cao Y, Liu K, Tian Z, Hogan SL, Yang J, Poulton CJ, Falk RJ, Li W. PTPN22 R620W polymorphism and ANCA disease risk in white populations: a metaanalysis. J Rheumatol. 2015;42(2):292-9. Epub 2014/12/01. doi: 10.3899/jrheum.131430. PubMed PMID: 25448792.

5.   Bottini N, Vang T, Cucca F, Mustelin T. Role of PTPN22 in type 1 diabetes and other autoimmune diseases. Seminars in Immunology. 2006;18(4):207-13. doi: 10.1016/j.smim.2006.03.008.

6.   Wang X-H, Ma A-G, Han X-X, Chen L, Liang H, Aishan L, Abudumijit A, Xue F. Protein tyrosine phosphatase nonreceptor type 22 (PTPN22) gene single nucleotide polymorphisms and its interaction with T2DM on pulmonary tuberculosis in Chinese Uygur population. Oncotarget. 2017;8(39):65601-8. doi: 10.18632/oncotarget.19274. PubMed PMID: 29029456.

7.   Wang H, Wang Z, Rani PL, Fu X, Yu W, Bao F, Yu G, Li J, Li L, Sun L, Yue Z, Zhao Q, Pan Q, Cao J, Wang C, Chi X, Wang Y, Yang Q, Mi Z, ..., Zhang F. Identification of PTPN22, ST6GAL1 and JAZF1 as psoriasis risk genes demonstrates shared pathogenesis between psoriasis and diabetes. Exp Dermatol. 2017;26(11):1112-7. Epub 20170825. doi: 10.1111/exd.13393. PubMed PMID: 28603863.

8.   Wang W, Meng X, Liu Y, Ma X, Zhang Q, Li C, Li C, Ren L. Association Between Protein Tyrosine Phosphatase Non-Receptor Type 22 (PTPN22) Polymorphisms and Risk of Ankylosing Spondylitis: A Meta-analysis. Med Sci Monit. 2017;23:2619-24. doi: 10.12659/msm.901083. PubMed PMID: 28555069.

9.   Armitage LH, Wallet MA, Mathews CE. Influence of PTPN22 Allotypes on Innate and Adaptive Immune Function in Health and Disease. Front Immunol. 2021;12:636618-. doi: 10.3389/fimmu.2021.636618. PubMed PMID: 33717184.

10.   Gomez LM, Anaya JM, Gonzalez CI, Pineda-Tamayo R, Otero W, Arango A, Martín J. PTPN22 C1858T polymorphism in Colombian patients with autoimmune diseases. Genes & Immunity. 2005;6(7):628-31. doi: 10.1038/sj.gene.6364261.

11.   Burn GL, Svensson L, Sanchez-Blanco C, Saini M, Cope AP. Why is *PTPN22* a good candidate susceptibility gene for autoimmune disease? FEBS Letters. 2011;585(23):3689-98. doi: 10.1016/j.febslet.2011.04.032.

12.   Maine CJ, Teijaro JR, Marquardt K, Sherman LA. PTPN22 contributes to exhaustion of
T lymphocytes during chronic viral infection. Proc Natl Acad Sci U S A. 2016;113(46):E7231-
E9. Epub 2016/10/31. doi: 10.1073/pnas.1603738113. PubMed PMID: 27799548.
13.   Jofra T, Di Fonte R, Hutchinson TE, Dastmalchi F, Galvani G, Battaglia M, Salek‐
Ardakani S, Fousteri G. Protein tyrosine phosphatase PTPN22 has dual roles in promoting
pathogen versus homeostatic‐driven CD8 T‐cell responses. Immunology & Cell Biology.
2016;95(2):121-8. doi: 10.1038/icb.2016.92.
14.   Bottini N, Musumeci L, Alonso A, Rahmouni S, Nika K, Rostamkhani M, MacMurray J,
Meloni GF, Lucarelli P, Pellecchia M, Eisenbarth GS, Comings D, Mustelin T. A functional
variant of lymphoid tyrosine phosphatase is associated with type I diabetes. Nature
Genetics. 2004;36(4):337-8. doi: 10.1038/ng1323.
15.   Begovich AB, Carlton VEH, Honigberg LA, Schrodi SJ, Chokkalingam AP, Alexander
HC, Ardlie KG, Huang Q, Smith AM, Spoerke JM, Conn MT, Chang M, Chang S-YP, Saiki RK,
Catanese JJ, Leong DU, Garcia VE, McAllister LB, Jeffery DA, ..., Gregersen PK. A missense
single-nucleotide polymorphism in a gene encoding a protein tyrosine phosphatase
(PTPN22) is associated with rheumatoid arthritis. American journal of human genetics.
2004;75(2):330-7. Epub 2004/06/18. doi: 10.1086/422827. PubMed PMID: 15208781.
16.   Orozco RC, Marquardt K, Pratumchai I, Shaikh AF, Mowen K, Domissy A, Teijaro JR,
Sherman LA. Autoimmunity-associated allele of tyrosine phosphatase gene PTPN22
enhances anti-viral immunity. PLoS Pathog. 2024;20(3):e1012095-e. doi:
10.1371/journal.ppat.1012095. PubMed PMID: 38512979.
17.   Knipper JA, Wright D, Cope AP, Malissen B, Zamoyska R. PTPN22 Acts in a Cell
Intrinsic Manner to Restrict the Proliferation and Differentiation of T Cells Following
Antibody Lymphodepletion. Front Immunol. 2020;11:52-. doi: 10.3389/fimmu.2020.00052.
PubMed PMID: 32047502.
18.   Fousteri G, Jofra T, Debernardis I, Stanford SM, Laurenzi A, Bottini N, Battaglia M.
The protein tyrosine phosphatase PTPN22 controls forkhead box protein 3 T regulatory cell
induction but is dispensable for T helper type 1 cell polarization. Clin Exp Immunol.
2014;178(1):178-89. doi: 10.1111/cei.12393. PubMed PMID: 24905474.
19.   Maine CJ, Marquardt K, Cheung J, Sherman LA. PTPN22 controls the germinal center
by influencing the numbers and activity of T follicular helper cells. Journal of immunology
(Baltimore, Md : 1950). 2014;192(4):1415-24. Epub 2014/01/22. doi:
10.4049/jimmunol.1302418. PubMed PMID: 24453256.
20.   Spalinger MR, Kasper S, Gottier C, Lang S, Atrott K, Vavricka SR, Scharl S, Raselli T,
Frey-Wagner I, Gutte PM, Grütter MG, Beer HD, Contassot E, Chan AC, Dai X, Rawlings DJ,
Mair F, Becher B, Falk W, ..., Scharl M. NLRP3 tyrosine phosphorylation is controlled by
protein tyrosine phosphatase PTPN22. J Clin Invest. 2016;126(11):4388. Epub 20161101.
doi: 10.1172/JCI90897. PubMed PMID: 27801679; PMCID: PMC5096914.
21.   Orozco RC, Marquardt K, Mowen K, Sherman LA. Proautoimmune Allele of Tyrosine
Phosphatase, PTPN22, Enhances Tumor Immunity. J Immunol. 2021;207(6):1662-71. Epub
20210820. doi: 10.4049/jimmunol.2100304. PubMed PMID: 34417261; PMCID:
PMC8428750.

22.   Dai X, James RG, Habib T, Singh S, Jackson S, Khim S, Moon RT, Liggitt D, Wolf-Yadlin
A, Buckner JH, Rawlings DJ. A disease-associated PTPN22 variant promotes systemic
autoimmunity in murine models. J Clin Invest. 2013;123(5):2024-36. Epub 2013/04/24. doi:
10.1172/JCI66963. PubMed PMID: 23619366.
23.   Hasegawa K, Martin F, Huang G, Tumas D, Diehl L, Chan AC. PEST Domain-Enriched
Tyrosine Phosphatase (PEP) Regulation of Effector/Memory T Cells. Science.
2004;303(5658):685-9. doi: 10.1126/science.1092138.
24.   Sanchez-Blanco C, Clarke F, Cornish GH, Depoil D, Thompson SJ, Dai X, Rawlings
DJ, Dustin ML, Zamoyska R, Cope AP, Purvis HA. Protein tyrosine phosphatase PTPN22
regulates LFA-1 dependent Th1 responses. J Autoimmun. 2018;94:45-55. Epub 2018/07/24.
doi: 10.1016/j.jaut.2018.07.008. PubMed PMID: 30054208.
25.   Anderson W, Barahmand-Pour-Whitman F, Linsley PS, Cerosaletti K, Buckner JH,
Rawlings DJ. PTPN22 R620W gene editing in T cells enhances low-avidity TCR responses.
Elife. 2023;12:e81577. doi: 10.7554/eLife.81577. PubMed PMID: 36961507.
26.   Vang T, Liu WH, Delacroix L, Wu S, Vasile S, Dahl R, Yang L, Musumeci L, Francis D,
Landskron J, Tasken K, Tremblay ML, Lie BA, Page R, Mustelin T, Rahmouni S, Rickert RC,
Tautz L. LYP inhibits T-cell activation when dissociated from CSK. Nat Chem Biol.
2012;8(5):437-46. doi: 10.1038/nchembio.916. PubMed PMID: 22426112.
27.   Salmond RJ, Brownlie RJ, Morrison VL, Zamoyska R. The tyrosine phosphatase
PTPN22 discriminates weak self peptides from strong agonist TCR signals. Nat Immunol.
2014;15(9):875-83. Epub 2014/08/10. doi: 10.1038/ni.2958. PubMed PMID: 25108421.
28.   Hornick EL, Wallis AM, Bishop GA. TRAF3 enhances type I interferon receptor
signaling in T cells by modulating the phosphatase PTPN22. Science signaling.
2022;15(753):eabn5507-eabn. Epub 2022/09/27. doi: 10.1126/scisignal.abn5507. PubMed
PMID: 36166512.
29.   Swiecki M, Colonna M. Type I interferons: diversity of sources, production pathways
and effects on immune responses. Curr Opin Virol. 2011;1(6):463-75. Epub 2011/11/25.
doi: 10.1016/j.coviro.2011.10.026. PubMed PMID: 22440910.
30.   Huber JP, Farrar JD. Regulation of effector and memory T-cell functions by type I
interferon. Immunology. 2011;132(4):466-74. Epub 2011/02/14. doi: 10.1111/j.1365-
2567.2011.03412.x. PubMed PMID: 21320124.
31.   Lukhele S, Boukhaled GM, Brooks DG. Type I interferon signaling, regulation and
gene stimulation in chronic virus infection. Seminars in immunology. 2019;43:101277-.
Epub 2019/05/30. doi: 10.1016/j.smim.2019.05.001. PubMed PMID: 31155227.
32.   Teijaro JR, Ng C, Lee AM, Sullivan BM, Sheehan KCF, Welch M, Schreiber RD, de la
Torre JC, Oldstone MBA. Persistent LCMV infection is controlled by blockade of type I
interferon signaling. Science (New York, NY). 2013;340(6129):207-11. doi:
10.1126/science.1235214. PubMed PMID: 23580529.
33.   Wilson EB, Yamada DH, Elsaesser H, Herskovitz J, Deng J, Cheng G, Aronow BJ, Karp
CL, Brooks DG. Blockade of chronic type I interferon signaling to control persistent LCMV
infection. Science (New York, NY). 2013;340(6129):202-7. doi: 10.1126/science.1235208.
PubMed PMID: 23580528.
34.   Trinchieri G. Type I interferon: friend or foe? J Exp Med. 2010;207(10):2053-63. Epub
2010/09/13. doi: 10.1084/jem.20101664. PubMed PMID: 20837696.

35.   Wang Y, Shaked I, Stanford SM, Zhou W, Curtsinger JM, Mikulski Z, Shaheen ZR,
Cheng G, Sawatzke K, Campbell AM, Auger JL, Bilgic H, Shoyama FM, Schmeling DO,
Balfour HH, Jr., Hasegawa K, Chan AC, Corbett JA, Binstadt BA, ..., Peterson EJ. The
autoimmunity-associated gene PTPN22 potentiates toll-like receptor-driven, type 1
interferon-dependent immunity. Immunity. 2013;39(1):111-22. Epub 2013/07/18. doi:
10.1016/j.immuni.2013.06.013. PubMed PMID: 23871208.
36.   Laboratory J. B6.Cg-Ptpn22tm2Achn/J. Available from:
https://www.jax.org/strain/028977.
37.   Jassim BA, Lin J, Zhang Z-Y. PTPN22: structure, function, and developments in
inhibitor discovery with applications for immunotherapy. Expert Opin Drug Discov.
2022;17(8):825-37. Epub 2022/06/07. doi: 10.1080/17460441.2022.2084607. PubMed
PMID: 35637605.
38.   Rehwinkel J, Gack MU. RIG-I-like receptors: their regulation and roles in RNA
sensing. Nat Rev Immunol. 2020;20(9):537-51. Epub 2020/03/13. doi: 10.1038/s41577-
020-0288-3. PubMed PMID: 32203325.
39.   Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity:
update on Toll-like receptors. Nature Immunology. 2010;11(5):373-84. doi:
10.1038/ni.1863.
40.   Zalinger ZB, Elliott R, Rose KM, Weiss SR. MDA5 Is Critical to Host Defense during
Infection with Murine Coronavirus. J Virol. 2015;89(24):12330-40. Epub 2015/09/30. doi:
10.1128/JVI.01470-15. PubMed PMID: 26423942.
41.   Zhang J, Zahir N, Jiang Q, Miliotis H, Heyraud S, Meng X, Dong B, Xie G, Qiu F, Hao Z,
McCulloch CA, Keystone EC, Peterson AC, Siminovitch KA. The autoimmune disease–
associated PTPN22 variant promotes calpain-mediated Lyp/Pep degradation associated
with lymphocyte and dendritic cell hyperresponsiveness. Nature Genetics. 2011;43(9):902-
7. doi: 10.1038/ng.904.
42.   Spalinger MR, Lang S, Vavricka SR, Fried M, Rogler G, Scharl M. Protein tyrosine
phosphatase non-receptor type 22 modulates NOD2-induced cytokine release and
autophagy. PLoS One. 2013;8(8):e72384-e. doi: 10.1371/journal.pone.0072384. PubMed
PMID: 23991106.
43.   Eisfeld AJ, Gasper DJ, Suresh M, Kawaoka Y. C57BL/6J and C57BL/6NJ Mice Are
Differentially Susceptible to Inflammation-Associated Disease Caused by Influenza A
Virus. Front Microbiol. 2019;9:3307-. doi: 10.3389/fmicb.2018.03307. PubMed PMID:
30713529.
44.   Malm Tillgren S, Nieto-Fontarigo JJ, Cerps S, Ramu S, Menzel M, Mahmutovic
Persson I, Meissner A, Akbarshahi H, Uller L. C57Bl/6N mice have an attenuated lung
inflammatory response to dsRNA compared to C57Bl/6J and BALB/c mice. J Inflamm
(Lond). 2023;20(1):6-. doi: 10.1186/s12950-023-00331-4. PubMed PMID: 36810092.
45.   Wang Y, Ewart D, Crabtree JN, Yamamoto A, Baechler EC, Fazeli P, Peterson EJ.
PTPN22 Variant R620W Is Associated With Reduced Toll-like Receptor 7–Induced Type I
Interferon in Systemic Lupus Erythematosus. Arthritis & Rheumatology. 2015;67(9):2403-
14. doi: https://doi.org/10.1002/art.39211.

46.   Ivashkiv LB. PTPN22 in autoimmunity: different cell and different way. Immunity.
2013;39(1):91-3. doi: 10.1016/j.immuni.2013.07.007. PubMed PMID: 23890067; PMCID:
PMC3785409.
47.   Kariuki SN, Crow MK, Niewold TB. The PTPN22 C1858T polymorphism is associated
with skewing of cytokine profiles toward high interferon-alpha activity and low tumor
necrosis factor alpha levels in patients with lupus. Arthritis Rheum. 2008;58(9):2818-23.
doi: 10.1002/art.23728. PubMed PMID: 18759295; PMCID: PMC2621106.
48.   Hamerman JA, Pottle J, Ni M, He Y, Zhang Z-Y, Buckner JH. Negative regulation of TLR
signaling in myeloid cells—implications for autoimmune diseases. Immunological
Reviews. 2016;269(1):212-27. doi: https://doi.org/10.1111/imr.12381.
49.   Berggren O, Rönnblom L, Eloranta M-L. Effect of PTPN22 Gene Variant R620W on
Type I Interferon Production Stimulated by Different Toll-like Receptor 7 Agonists:
Comment on the Article by Wang et al. Arthritis & Rheumatology. 2016;68(4):1045-. doi:
https://doi.org/10.1002/art.39571.
50.   Lin M, Ji X, Lv Y, Cui D, Xie J. The Roles of TRAF3 in Immune Responses. Dis Markers.
2023;2023:7787803-. doi: 10.1155/2023/7787803. PubMed PMID: 36845015.
51.   Lin X, Pelletier S, Gingras S, Rigaud S, Maine CJ, Marquardt K, Dai YD, Sauer K,
Rodriguez AR, Martin G, Kupriyanov S, Jiang L, Yu L, Green DR, Sherman LA. CRISPR-Cas9–
Mediated Modification of the NOD Mouse Genome With Ptpn22R619W Mutation Increases
Autoimmune Diabetes. Diabetes. 2016;65(8):2134-8. doi: 10.2337/db16-0061.

ABBREVIATIONS

547	**ABBREVIATIONS**	
548	IFN-I	Type I Interferon
549	Ptpn22	Protein tyrosine phosphatase non-receptor type 22
550	Lyp	Lymphoid protein
551	PEP	PEST-domain enriched phosphatase
552	DCs	Dendritic Cells
553	TLR4	Toll-Like Receptor 4
554	LPS	Lipopolysaccharide
555	3p-hp-RNA	5' Triphosphate hairpin ribonucleic acid
556	MHV A59	Mouse Hepatitis virus strain A59
557	TNFα	Tumor Necrosis Factor alpha
558	IL	Interleukin
559	R	Arginine
560	W	Trypotophan
561	HLA	Human Leukocyte Antigen
562	Tregs	T regulatory cell
563	CRISPR-Cas9	Cluster regularly interspaced short palindromic repeats Cas9
564	APC	Antigen Presenting Cell
565	LCMV-cl13	Lymphocytic Choriomeningitis Virus Clone 13
566	TLR3	Toll-like receptor 3
567	SNP	single nucleotide polymorphism
568	RLR	Rig I like receptor
569	MDA5	Melanoma Differentiation Associated protein 5

570	IFNβ	Interferon Beta
571	RNA	Ribonucleic Acid
572	BMDC	Bone Marrow Derived Dendritic cells
573	BMM	Bone Marrow Derived Macrophage
574	JAX	Jackson Labs
575	mRNA	messenger ribonucleic acid
576	FLT3-L	FMS-like tyrosine kinase 3- Ligand
577	GM-CSF	Granulocyte- Macrophage Colony Stimulating factor
578	RIG-I	Retinoic acid inducible gene 1
579	MDP	Muramyl dipeptide
580	ELISA	Enzyme Linked Immunosorbent Assay
581	ANOVA	Analysis of Variance
582	NFκB	Nuclear Factor kappa B
583	TRAF3	TNF receptor-associated factor 3
584	IACUC	Institutional Animal Care and Use Committee
585	DMEM	Dulbecco's Modified Eagle Medium
586	FBS	Fetal Bovine Serum
587	MOI	Multiplicity of Infection
588	Ns	not significant
589	qRT-PCR	Quantitative Reverse Transcriptase Polymerase Chain Reaction

 FIGURE LEGENDS

Figure 1. Strain specific differences impact IFNβ production in PEP-null BMDCs. Gene

expression values of *Ptpn22* relative to *GAPDH* were determined by qRT-PCR for naïve PEP-

WT (black bars) and PEP-null (CRISPR) (red) splenocytes, each dot represents a mouse **(A)**.

FLT3-L BMDCs were cultivated and differentiated from C57BL/6J PEP-WT (black),

CRISPR/Cas9-generated PEP-null (red), and PEP-null mice from the Jackson Laboratory (JAX

stock #028977) (gray). Cultures were incubated with LPS (500 ng/mL) for 18 hours. The

supernatant was used to determine IFNβ concentration (ELISA) **(B)**. PEP-WT and PEP-null

(CRISPR) BMMs were incubated with LPS (500 ng/mL) and 10 uM LTV-1 for 18 hours. The

supernatant was used to determine IFNβ concentration (ELISA) **(C)**. ns= no significance;

*p<0.01; **p<0.001; ****p<0.0001; Two Way ANOVA with Tukey post hoc analysis. Data is

pooled from 2 independent experiments **(B)**. Data is representative of 3 biological replicates **(C)**.

Figure 2. PEP and PEP-R619W do not mediate IFNβ production in myeloid cells following

LPS, MHV A59, or viral RNA stimulation. C57BL/6J PEP-WT (black, square symbol) and

PEP-null (red, circle symbol) BMMs were cultivated and then stimulated with LPS (500 ng/mL)

or infected with MHV A59 (MOI 0.1) for 18 hours **(A)**. FLT3-L BMDCs were cultivated from

C57BL/6J PEP-WT, PEP-null, and PEP-R619W mice. Cultures were incubated with LPS (500

ng/mL) **(B)** or 3p-hpRNA (2 ng/mL) **(C)** for 18 hours. The supernatant was used to determine

IFNβ concentration (ELISA). ns= no significance. Two Way ANOVA with Tukey post hoc

analysis. Data is representative of 2 biological replicates **(A)**. Data is pooled from 3 independent

experiments **(B-C)**.

Figure 3. PEP and the PEP-R619W variant distinctly modulate the production of pro-inflammatory cytokines in dendritic cells. FLT3-L BMDCs were cultivated from C57BL/6J PEP-WT (black), PEP-null (red), and PEP-R619W (blue) mice. Cultures were incubated with LPS (500 ng/mL) for 18 hours. The supernatant was used to determine IL-2 **(A)**, IL-12 **(B)**, and TNFα **(C)** concentration (ELISA). ns= no significance;**$p<0.01$, ****$p<0.0001$; Two Way ANOVA with Tukey post hoc analysis. Data is representative of 2 independent experiments.

Figure 1

Figure 2

Figure 3