Profound taxonomic and functional gut microbiota alterations associated with trichuriasis: cross-country and country-specific patterns

Pierre H.H. Schneeberger[1,2,§], Julian Dommann[1,2], Nurudeen Rahman[1,2], Eveline Hürlimann[1,2], Somphou Sayasone[2,3], Said Ali[4], Jean Tenena Coulibaly[5,6], Jennifer Keiser[1,2]

1 Swiss Tropical and Public Health Institute, Department of Medical Parasitology and Infection Biology, Helminth Drug Development Unit, Allschwil, Switzerland, 2 University of Basel, Basel, Switzerland 3 Lao Tropical and Public Health Institute, Vientiane, Lao People's Democratic Republic, 4 Public Health Laboratory Ivo de Carneri, Chake Chake, Pemba, Tanzania, 5 Unité de Formation et de Recherche Biosciences, Université Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire, 6 Centre Suisse de Recherches Scientifiques en Côte d'Ivoire, Abidjan, Côte d'Ivoire

§ Corresponding author: Swiss Tropical and Public Health Institute, Department of Medical Parasitology and Infection Biology, Allschwil, Switzerland

Tel.: +41 61 284 8323, E-mail: pierre.schneeberger@swisstph.ch

E-mail addresses:

PHHS   pierre.schneeberger@swisstph.ch

JD   julian.dommann@swisstph.ch

NR   nurudeen.rahman@swisstph.ch

EH   eveline.huerlimann@swisstph.ch

SS   somphou.sayasone@yahoo.com

SA   saidmali2003@yahoo.com

JTC   couljeanvae@yahoo.fr

JK   jennifer.keiser@swisstph.ch

Keywords: soil-transmitted helminths, *Trichuris trichiura*, gut microbiome alteration

42   **Abstract:**

Background. The human gastrointestinal microbiota plays a crucial role in immune modulation, metabolism, and pathogen resistance. Soil-transmitted helminth (STH) infections, including *Trichuris trichiura* (whipworm), significantly alter gut microbial composition, yet the extent and functional consequences of these changes remain underexplored across different geographical regions. This study investigates the taxonomic and functional impacts of *T. trichiura* on gut microbiota in three endemic regions - Côte d'Ivoire, Laos, and Tanzania - using a unified high-resolution metagenomic sequencing approach.

Results. This study reveals extensive gut microbiota disruptions linked to *T. trichiura* infection, with both regional and cross-country patterns. A core signature found across all study sites includes depletion of *Faecalibacterium prausnitzii* and *Eubacterium rectale* (Short-chain fatty acids (SCFA) producers) and enrichment of mucin-degrading bacteria (*Ruminococcus*, *Bacteroides*), alongside increased host-derived carbohydrate metabolism. Infection destabilized microbial networks, characterized by reduced connectivity and clustering, with opportunistic taxa such as *Segatella copri* emerging as network hubs across regions, indicating a shared ecological response. Both taxonomic and functional disruptions exhibited a combination of conserved and region-specific patterns; for example, whereas certain taxa, such as *Prevotella* and *Streptococcus*, showed notable geographic variability, specific functional changes, such as SCFA depletion and mucin degradation, were consistently observed across sites. These conserved functional changes suggest that *T. trichiura* imposes similar metabolic pressures on the gut microbiome across populations, potentially affecting host nutrient availability and immune responses in a predictable manner.

Discussion and Conclusion. This study provides robust evidence that *T. trichiura* infection induces significant and consistent microbiome alterations across diverse populations. The depletion of SCFA-producing bacteria and the enrichment of mucin-degrading taxa, along

with corresponding metabolic pathways, imply compromised gut barrier integrity, providing

insights into the complex inflammatory processes associated with this helminth infection.

Additionally, these microbiome differences could play a critical role in facilitating parasite

persistence and reinfection, which remain major challenges limiting the efficacy of global

control strategies. Our findings highlight the potential of microbiome-targeted interventions,

such as probiotic supplementation or dietary modifications, to mitigate the health impacts of

T. trichiura infections by restoring microbial homeostasis.

Introduction:

The human gastrointestinal tract harbors a complex and diverse microbial community integral

to maintaining health and homeostasis [1]. These microbes engage in essential processes such

as nutrient metabolism, immune modulation, and protection against pathogenic invasion.

Disruptions to this delicate ecosystem, known as dysbiosis, have been linked to a wide array

of health conditions, including metabolic syndromes, inflammatory diseases, and infections

[2]. Parasitic infections, particularly those caused by soil-transmitted helminths (STHs), are

significant drivers of dysbiosis, profoundly altering gut microbial structure and function [3-

6]. Among these helminths, *Trichuris trichiura* - commonly referred to as the human

whipworm - remains a critical public health concern due to particularly low treatment success

compared to other helminth species, especially in low- and middle-income countries [7, 8].

Globally, it is estimated that more than 477 million people are infected with *T. trichiura*, with

the heaviest burden in tropical and subtropical regions where poor sanitation facilitates

transmission [8]. The parasite primarily colonizes the large intestine, where it anchors in the

colonic epithelium and mucosa, triggering localized inflammation and immune responses that

can alter gut microbiota composition [9]. Research on *T. trichiura* infection has highlighted

its capacity to influence the microbial community in multiple ways, while the gut microbiota,

in turn, plays a role in shaping the parasite's life cycle. First, the parasite induces a

pronounced Th2-mediated immune response characterized by increased mucus production

and altered epithelial barrier function, which can create a niche favourable to certain

microbial taxa [10]. Second, the parasite's nutrient competition and physical presence in the

intestinal lumen may disrupt microbial interactions, contributing to compositional and

functional shifts in the microbiota [11]. Third, *T. trichiura* infection often leads to broader

physiological changes, such as altered bile acid profiles and reduced short-chain fatty acid

(SCFA) production, further shaping microbial ecology [12]. Conversely, the gut microbiota

also plays a crucial role in *T. trichiura* colonization, as microbial communities contribute to

egg hatching and larval development, establishing a bidirectional relationship between the

parasite and the intestinal microbiome [13]. This bidirectional interplay underscores the

complex relationship between the helminth and the gut microbiota. Previous studies have

begun to explore the relationship between *T. trichiura* infection and gut microbiota, reporting

both increased and decreased microbial diversity depending on the population studied and the

specific metrics used [14]. For example, a study using a mouse model of *T. muris* infection

suggest that the presence of the parasite can enrich for specific taxa, including *Lactobacillus*

species [15], while another study reports declines in beneficial commensals associated with *T.*

trichiura infection in a small human cohort [15]. Despite these findings, substantial

knowledge gaps remain. There is limited understanding of how *T. trichiura* affects microbial

diversity and function across diverse geographic regions with varying environmental, dietary,

and host genetic factors. Functional consequences of these microbiota changes are also

underexplored. While a few studies have highlighted alterations in metabolic pathways, such

as reduced Short-chain fatty acids (SCFA) production and shifts in carbohydrate and amino

acid metabolism, these findings are often based on single populations or limited datasets.

Furthermore, the extent to which *T. trichiura*-induced microbial changes are conserved or

vary across regions remains unclear, especially given the significant role that local factors

play in shaping the gut microbiota.

This study aims to address these gaps by systematically characterizing the taxonomic,

functional, and ecological impacts of *T. trichiura* infection on the gut microbiota across

multiple countries using a unified technical framework. By focusing on *T. trichiura* - the only

large intestine-colonizing helminth species - this work seeks to provide a more

comprehensive understanding of the interplay between parasitic infections and microbial

ecology. These insights are critical for advancing our knowledge of host-microbiota-parasite

interactions and may inform the development of targeted therapies to mitigate the impact of

helminth infections on gut health.

Methods.

Sample Collection and Microscopy. Stool samples were collected as part of a multi-country

randomized controlled trial evaluating the efficacy and safety of albendazole-ivermectin

combination therapy against *T. trichiura* and concomitant helminth infections [16]. The trial

is registered on ClinicalTrials.gov (NCT03527732, registered on 17 May 2018). Stool

samples were analyzed using the standard Kato-Katz microscopy method, previously

described in the literature [17]. For molecular analysis, a small aliquot (<1 g) of stool was

transferred to a 2 mL screwcap cryotube using a UV-sterilized plastic spatula and

immediately frozen at -20°C. At the conclusion of the respective trial stage, frozen samples

were shipped to Swiss TPH (Allschwil, Switzerland) on dry ice and stored at -20°C until

further analysis.

DNA Isolation. DNA was extracted from stool samples using the QIAGEN DNEasy

PowerSoil Pro (Qiagen, Hilden, Germany) following the manufacturer's protocol. DNA

concentration was quantified using 2 µL of extracted DNA with the Qubit 4.0 fluorometer

and high-sensitivity DNA quantification kits (Thermo Fisher Scientific, Waltham, MA,

USA).

Shotgun Sequencing. Shallow shotgun sequencing was performed as described previously

[18], generating 2.8E+06 to 5.3E+07 reads per sample. Libraries were prepared using the

NEBNext Ultra II FS DNA Library Prep Kit (New England Biolabs, Ipswich, MA, USA)

according to the manufacturer's instructions. Final libraries were quantified and pooled

equimolarly before loading onto an Illumina NovaSeq 6000 platform. Sequencing was

conducted on S4 flowcell (300 cycles) in paired-end mode (2 × 150 bp). Sample-specific

sequencing depth is detailed in **Supplementary Table 1**.

Taxonomic and Functional Profiling. Taxonomic profiling of shallow shotgun sequencing

data was conducted with MetaPhlAn v4.1.1 (11 Mar 2024) [19]. The CHOCOPhlAn species-

marker database vOct22_CHOCOPhlAnSGB_202403 (October 2022) was used, and

mapping was performed with Bowtie2 v2.5.4 [20]. The resulting taxonomic relative

abundance tables were merged using the built-in "merge_metaphlan_tables.py" and were

filtered for low counts (minimum count = 100; prevalence = 20% of samples) and low

variance (50% features removed based on interquartile range) using MicrobiomeAnalyst v2.0

[21]. Functional profiling was performed using HUMAnN4 v4.0.0.alpha.1 with default

settings and the UniRef 2019_06 protein database. The unstratified functional abundance

tables, including the "reactions" (= abundance tables based on the Enzyme Comission (EC)

numbers) tables as well as the "pathabundance" (= abundance tables summarizing the

abundance of MetaCyc pathways [22] based on EC tables) files were normalized to copies

per million (CPM) using the humann_renorm_table.py script before downstream analyses.

Generation of microbiome-specific metrics. All analyses were conducted in R-studio 2024

using R version 4.4+[23]. Alpha and beta diversity metrics were calculated to assess

microbial diversity within and between samples. Alpha diversity, reflecting the richness and evenness of species within individual samples, was evaluated using the vegan R package v2.6-6 [24]. Metrics included species richness (number of species), the Shannon diversity index for richness and evenness, and the Berger-Parker index to measure community domination. Beta diversity, representing dissimilarity between samples, was assessed using Bray-Curtis dissimilarity, calculated through the "vegdist" function in the same package. Non-metric multidimensional scaling (NMDS) was performed to visualize compositional differences in a reduced dimensional space, with ordination coordinates calculated using the "metaMDS" function configured with k = 2 and try = 10,000 iterations. To statistically evaluate differences in community-level composition, PERMANOVA (Permutational Multivariate Analysis of Variance) and PERMDISP (Permutation-Based Multivariate Dispersion) analyses were performed using the "adonis2", "betadisper", and "permutest" functions. The core microbiome, comprising taxa consistently present across a defined proportion of samples, was identified using the "core" function from the microbiome R package v1.23.1 [25]. Taxonomic profiles were visualized with a heatmap generated through an in-house script combining multiple R packages. The ComplexHeatmap package v2.18.0 [26] was used for heatmap generation, with hierarchical clustering performed using the "hclust" function and Ward's algorithm, and optimized for clarity using the "dendsort" function. The species network correlation analysis was conducted using functions from the SpiecEasi package v1.1.1 [27], the RCy3 package v3.20 [28], and the igraph package v2.0.3 [29]. Network statistics were calculated using the following functions from the igraph package: "graph.density" for network density, "average.path.length" for the average path length, "diameter" for the network diameter, and "transitivity" for the global clustering coefficient. Centrality measures, including degree, betweenness, closeness, and eigenvector

centrality, were calculated using the "degree", "betweenness", "closeness", and
"eigen_centrality" functions, respectively.

Statistical Analysis. Group comparisons were conducted using Mann-Whitney U tests for
two-group comparisons and Kruskal-Wallis tests for comparisons involving three or more
groups, and all *P*-values were adjusted using the Bonferroni family-wise correction [30].
Species prevalence comparisons were performed in R using the "fisher.test" function, and the
resulting *P*-values were adjusted using the Benjamini-Hochberg (BH) procedure [31]. To
compare bacterial species abundances, several statistical methods and tools were employed.
Adjusted Kruskal-Wallis tests and the LEfSe package [32], available on the
MicrobiomeAnalyst platform, were used on country-specific subsets to assess enrichment or
depletion of taxa associated with infection status. Additionally, MaAsLin2 v1.18.0 [33] was
applied to the complete dataset, controlling for the "Country" variable. All *P*-values resulting
from these analyses were adjusted using the BH procedure. To minimize false-positive
signals, only species with significant corrected *P*-values across all three statistical methods
were considered enriched or depleted. Effect sizes of these differences were visualized using
log2 fold-change values generated by MaAsLin2. For comparisons of functional profiles,
including ECs and MetaCyc pathway abundances, MaAsLin2 was used to assess statistical
differences.

Visualizations. Most visualizations were created and optimized using OriginPro 2024
v10.1.0.178 (OriginLab Corporation, Northampton, MA, USA). Heatmaps were generated
directly in R using the ComplexHeatmap package, while Venn diagrams were created using
the ggplot2 package v3.5.1.

Results:

Sampling results and study characteristics. Between September 2018 and December 2019, individuals from the Pujini and Dodo shehias (health districts) on Pemba Island, Tanzania; the Dabou and Jacqueville districts in the Grand Ponts region, Côte d'Ivoire; and the Nambak district in the Luang Prabang province, Laos, were screened for *T. trichiura* as described elsewhere [16]. A total of 8,866 participants were assessed for eligibility. Of these, 160 participants who tested negative for *T. trichiura* (60 in Côte d'Ivoire, 50 in Laos, and 50 in Tanzania) were selected to represent the "uninfected" microbiome group in this study. 632 *T. trichiura*-positive participants were selected to represent the *Trichuris*-infected group (200 in Côte d'Ivoire, 239 in Laos, and 200 in Tanzania) and were included in this study (**Table 1**, **Figure 1**). 9/799 samples included in this study were dropped due to insufficient sequencing quality. Participants were predominantly young, with mean ages ranging from 14.1 years in Tanzania to 24 years in Laos, and an overall age range of 6–60 years across all regions. Females accounted for roughly half of the infected group, with proportions varying slightly between regions (50.5% in Côte d'Ivoire to 56.6% in Tanzania). *T. trichiura* infections were mostly of light intensity, with geometric mean eggs per gram (EPG) values ranging from 380 in Laos to 497 in Côte d'Ivoire, while moderate and heavy infections were less common.

Characteristic	Côte d'Ivoire	Laos	Tanzania
Number of patients (n)			
Infected	198	236	198
Uninfected	60	48	50
Total	258	284	248
Age (y)			
Mean (arit) ($\pm$ SD)	16.9 ($\pm$ 14)	24 ($\pm$ 16.4)	14.1 ($\pm$ 10)
Range	6 to 60	6 to 60	6 to 60
Female, n (%)			
Infected	100 (50.5%)	125 (53%)	112 (56.6%)
***Trichuris trichiura* infection**			
EPG_{arit} ($\pm$ SD)	1081 ($\pm$ 2047)	709 ($\pm$ 1255)	962 ($\pm$ 1784)

EPG$_{geom}$ (± SD)	497 (± 3)	380 (± 2.7)	450 (± 3.1)
Range (EPG)	102-15072	102-12822	18-12750
Light, n (%)	151 (76%)	197 (83%)	28 (71.8%)
Moderate, n (%)	44 (22%)	38 (16%)	45 (23%)
Heavy, n (%)	3 (1.5%)	1 (0.4%)	3 (1.5%)

Table 1. Cohort description. y = year; EPG = eggs per gram of stool; SD = standard
deviation; arit = arithmetic mean; geom = geometric mean; Infection intensity classification
is based on World Health Organization criteria [34].

Community level taxonomic comparisons. A comparison of alpha diversity metrics between
infected and uninfected individuals revealed significant differences across the three countries,
as shown in **Figure 2A**. Richness was significantly higher in infected participants from Laos
($P<0.001$) and Tanzania ($P=0.001$), whereas infected study participants in Côte d'Ivoire
exhibited significantly lower richness compared to uninfected individuals ($P<0.001$). A
similar trend was observed for composite metrics such as Shannon Diversity; however, this
difference was not statistically significant for the Tanzanian cohort ($P=0.11$). Analysis of
dominance, using the Berger-Parker index, indicated significantly elevated dominance in
uninfected individuals in Laos ($P<0.001$), suggesting differing community structures between
infection states. Community-level taxonomic composition, evaluated using PERMANOVA
(**Figure 2B**), showed that infection status was significantly associated with shifts in
community structure in all three countries ($P<0.001$). PERMDISP analysis revealed
significant differences in dispersion around the group centroid in Côte d'Ivoire ($P=0.008$) and
Laos ($P=0.049$), indicating greater heterogeneity in these settings, while no such differences
were detected in Tanzania ($P=0.472$). Core microbiota comparisons (**Figure 2C**) highlighted
pronounced differences in microbial composition across countries. Notably, *Bifidobacterium*
pseudocatenulatum and *B. adolescentis* were highly prevalent in Tanzania but largely absent
in Côte d'Ivoire and Laos, confirming country-specific microbial structures. Clustering of all
samples based on species-level abundances (**Figure 2D**) revealed distinct country-specific

clusters. While infection status did not lead to discernible clustering in Côte d'Ivoire and

Tanzania, a significant association was observed in Laos (Chi-square with Yates correction;

P<0.001). Specifically, one sample cluster from Laos contained a disproportionately higher

proportion of infected patients (n = 90/92), indicating potential infection-related microbial

patterns unique to this setting (**Supplementary Figure 1A**).

Species-level taxonomic signal associated with infection status across countries. A

prevalence-based comparison between infected and uninfected individuals, analyzed using a

Fisher exact test with Benjamini-Hochberg correction, highlighted significant differences

across multiple bacterial taxa (**Figure 3A**). A total of 104 species showed variations in

prevalence across the three countries studied. Notably, the genus with the highest number of

significant differences was *Streptococcus*, comprising 10 species that were either more or less

prevalent depending on infection status. Among these, several species demonstrated

consistent differences in prevalence across two or more countries, including *S. sanguinis*

(n=2), *S. parasanguinis* (n=2), *S. gallolyticus* (n=2), *S. australis* (n=2), and *S. gordonii* (n=3).

In **Figure 3B**, we present modelled species differences based on their relative abundances

using a stringent multi-tool statistical approach that combined the Kruskal-Wallis test, LEfSe,

and MaAsLin2 (adjusted for country as a variable). Species found to be enriched or depleted

in this analysis often overlapped with those identified in the prevalence-based model. For

example, several taxa from the *Actinomyces*, *Blautia*, *Clostridium*, *Roseburia*, *Ruminococcus*,

Segatella, and *Streptococcus* genera showed significant differences in both prevalence and

relative abundance models. Interestingly, many species that were differentially abundant

exhibited a country-specific fold-change, which was frequently opposite in direction. Further

strain-level analysis (**Supplementary Figure 2**) revealed that the divergences were primarily

attributable to different strains within the same species, indicating that strain-level variations,

rather than species-level differences alone, drive the community shifts associated with *T*.

trichiura infection status. In **Figure 3C**, we present the comparison of the cumulative relative

abundance of species associated with infection status, stratifying taxa into those enriched or

depleted in only one country versus those consistently affected in two or more countries.

While there was no significant difference in the cumulative relative abundance of species

unique to individual countries, overlapping taxa demonstrated significant differences in

cumulative relative abundance across the three countries ($P_{CI}<0.0001$, $P_{LA}=0.0413$,

$P_{TA}=0.0062$). These findings highlight the core association of these overlapping taxa with the

presence or absence of *T. trichiura*, suggesting a conserved response pattern across distinct

geographic regions.

Changes in species networks associated with infection status. To explore the effects of *T.*

trichiura infection on microbial community interactions, we performed network-based

analyses using SpiecEasi on taxonomic profiles stratified by infection status and country.

Metrics were calculated using the 200 most abundant species, and visualizations focused on

the top 50 species in each group (**Figure 4**). Species networks built from infected and

uninfected participant samples displayed marked differences in structure across all countries.

Key metrics, such as the number of edges (E), network density (De), average path length

(PL), diameter (Di), and clustering coefficient (CC), consistently highlighted disrupted

microbial interactions associated with infection status. In Côte d'Ivoire, networks from

infected participants had an increased number of edges (E=497 vs. 385), higher density

(De=0.025 vs. 0.019), and greater clustering coefficient (CC=0.198 vs. 0.147), along with a

shorter average path length (PL=3.5 vs. 4.1). These changes suggest a more fragmented yet

locally interconnected microbial network. In Laos, infected participants exhibited a lower

number of edges (E=306 vs. 368), lower density (De=0.015 vs. 0.018), and shorter average

path length (PL=3.9 vs. 4.6), but an increased clustering coefficient (CC=0.186 vs. 0.115).

This indicates a more modular and less connected network structure. Finally, in Tanzania, the

clustering coefficient was higher in infected participants (CC=0.187 vs. 0.107), while the

path length was shorter (PL=4.5 vs. 5.3) and diameter was reduced (Di=13 vs. 17). These

changes point to a more compact and tightly clustered microbial network. Infected

participant's networks consistently showed reduced inter-species interactions, with several

taxa shifting in their network roles. Genera such as *Streptococcus*, *Clostridium*, *Dorea*, and

Blautia were central, highly connected nodes in uninfected participant's networks but became

peripheral in infected networks. Conversely, species like *Segatella copri* emerged as a new

hub in infected groups, suggesting it's increased ecological importance or resilience in the

context of infection. Some species – including *Faecalibacterium prausnitzii* and *Eubacterium*

rectale - became central hubs in infected patients in Côte d'Ivoire and Laos, but not in

Tanzania, thus also indicating a widespread, yet locally adapted, ecological response to *T.*

trichiura infection. These changes underscore a reorganization of microbial community

dynamics in response to *T. trichiura* infection. While infection-induced network perturbations

were evident across all countries, their magnitude varied, reflecting the influence of local

environmental and host factors. Nevertheless, conserved patterns emerged, including reduced

connectivity and clustering in infected networks, supporting the hypothesis that *T. trichiura*

infection disrupts microbial community stability.

Functional shifts in microbial communities associated with T. trichiura infection. To

investigate functional changes associated with *T. trichiura* infection, we analyzed abundance

profiles based on Enzyme Commission (EC) numbers, which systematically classify enzymes

by function, across infection status and countries. A total of 6,440 ECs were found, of which

5,453 were shared between the three countries, regardless of infection status

(**Supplementary Figure 3A**). As shown in **Supplementary Figure 3B**, functional richness

was found to be significantly lower in Laos compared to Côte d'Ivoire (*P*<0.0001) and

Tanzania (*P*<0.0001), as well as significantly different when stratified for infection status in

Côte d'Ivoire (P=0.02) and Tanzania (P<0.001) but not in Laos (P=0.68). Difference

between infected and uninfected groups were observed in each country, with infected groups

showing 458, 615, and 840 unique ECs in Côte d'Ivoire, Laos, and Tanzania, respectively

(**Figure 5A**). In contrast, uninfected groups exhibited lower unique functions, with 122, 38,

and 25 unique ECs in the respective countries, indicating a reduction in function associated

with infection status. NMDS analysis based on Bray-Curtis dissimilarity (**Figure 5B**)

revealed distinct clustering by infection status and country. PERMANOVA confirmed

significant differences between infected and uninfected groups (Côte d'Ivoire: R^2=0.0076,

P=0.091; Laos: R^2=0.0217, P=0.001; Tanzania: R^2=0.0098, P=0.014), thus indicating a

stronger impact of *Trichuris* infection on the functional potential of gut communities in Laos.

PERMDISP analysis indicated no significant differences in community dispersion in Laos

and Tanzania (P>0.05), but Côte d'Ivoire showed marginal significance (F-stat=3.89,

P=0.035), suggesting slightly greater variability in functional profiles within infected

communities. Volcano plots (**Figure 5C**) highlighted ECs significantly enriched or depleted

in infected versus uninfected groups. Overall, the highest number of differentially abundant

ECs were found in Laos (n = 1279), followed by Côte d'Ivoire (n = 317) and Tanzania (n =

293). Pathway-level comparisons (**Figure 5D, Supplementary Table 2**) revealed infection-

associated shifts in core metabolic processes. We identified significant alterations in 67

MetaCyc pathways between *Trichuris*-infected and uninfected patients using Maaslin2 and

controlling for the "Country" variable (P_{adj}<0.05). Several pathways related to carbohydrate

metabolism, fermentation, and biosynthesis were differentially abundant between *Trichuris*-

infected and non-infected individuals, reflecting functional reorganization of the gut

microbiome. Among the pathways significantly depleted in infected individuals, PWY_5514

($log2_{FC}$=-0.494, FDR=6.86E-05) and UDPNACETYLGALSYN_PWY ($log2_{FC}$=-0.518,

FDR=0.000103), both involved in carbohydrate biosynthesis, exhibited strong reductions,

suggesting disruptions in glycan metabolism. PWY_4984, which is involved in inorganic

nutrient metabolism, was also underrepresented ($log2_{FC}$=-0.234, FDR=0.000174), indicating

altered nutrient utilization. Fermentation-related pathways, many contributing to SCFA

metabolism, were also significantly reduced. PWY_3801, linked to mixed-acid fermentation,

showed a $log2_{FC}$ of -0.37 (FDR=0.00076), while PWY_7688, a component of carbohydrate

biosynthesis, was depleted ($log2_{FC}$=-0.451, FDR=0.0041). In Laos, PWY_7254, a butyrate-

producing fermentation pathway, was significantly reduced ($log2_{FC}$=-0.259, FDR=0.0122),

suggesting a lower microbial capacity for SCFA production. In contrast, non-infected

individuals, particularly in Côte d'Ivoire, exhibited enrichment of fermentation pathways,

including PWY_6545 ($log2_{FC}$=0.066, FDR=0.0146) and PWY_6731 ($log2_{FC}$=0.146,

FDR=0.0153), which contribute to SCFA biosynthesis. These pathways are essential for gut

health and immune modulation, and their depletion in infected individuals may indicate

reduced SCFA availability, with potential implications for gut barrier function and host

immunity. Finally, several pathways were enriched in non-infected individuals, including

PWY_I9 ($log2_{FC}$=0.225, FDR=0.00023), involved in amino acid biosynthesis, and

GLUDEG_I_PWY ($log2_{FC}$=0.244, FDR=0.017), associated with amine and polyamine

degradation. These differences suggest infection-driven or -associated shifts in nitrogen

metabolism.

The stratified analysis across the three study countries identified several pathways exhibiting

either consistent or contrasting enrichment/depletion patterns (**Figure 5E**). MET-SAM-PWY,

a superpathway involved in methylation processes, was enriched in Côte d'Ivoire

(log2FC=0.282, FDR=0.089) but significantly depleted in Laos (log2FC=-0.689,

FDR=0.000988), suggesting regional differences in microbial methylation potential.

PWY_5913, associated with the TCA cycle, was depleted in infected individuals in Tanzania

(log2FC=-0.34, FDR=0.0389) but enriched in Laos (log2FC=0.247, FDR=0.0705),

highlighting country-specific adaptations in microbial energy metabolism in response to

infection. LACTOSECAT-PWY, involved in carbohydrate degradation, was depleted in

Tanzania (log2FC=-0.482, FDR=0.0389) but enriched in Laos (log2FC=0.359,

FDR=0.0288), pointing to potential dietary influences on carbohydrate utilization across

regions. PWY-6906, which also plays a role in carbohydrate metabolism, was enriched in

infected individuals in Côte d'Ivoire (log2FC=0.478, FDR=0.00774) but strongly depleted in

Laos (log2FC=-1.42, FDR=1.53E-09), underscoring divergent microbial responses between

these two settings. Conversely, some pathways displayed similar patterns of enrichment

across countries. P164-PWY, associated with nucleoside and nucleotide degradation, was

consistently enriched in Côte d'Ivoire (log2FC=0.234, FDR=0.0203) and Laos

(log2FC=0.354, FDR=0.0445), indicating a conserved microbial response to infection.

Similarly, PWY-7111, related to fermentation processes, was enriched in infected

participants in both Côte d'Ivoire (log2FC=0.128, FDR=0.0708) and Laos (log2FC=0.117,

FDR=0.0402). Finally, GLCMANNANAUT-PWY, involved in amine and polyamine

degradation, was also enriched consistently in both Côte d'Ivoire (log2FC=0.136,

FDR=0.0949) and Laos (log2FC=0.184, FDR=0.00693). Across pathways associated with

Trichuris infection status in at least one country, the $log2_{FC}$ values were positively correlated

between Côte d'Ivoire and Tanzania (Spearman correlation, ρ=0.285, *P*<0.0001), suggesting

a degree of consistency in microbiome functional shifts associated with infection in these two

regions (**Figure 5F**). However, these values were strongly negatively correlated between

Laos and both Côte d'Ivoire (ρ=-0.395, *P*<0.0001) and Tanzania (ρ=-0.383, P<0.0001),

indicating a contrasting pattern of pathway enrichment and depletion between these regions.

Discussion:

This study provides the most comprehensive metagenomic investigation to date of *T.*

trichiura-associated microbiome alterations, leveraging shotgun metagenomics across three

geographically distinct regions. By comparing infected and uninfected individuals, we identified both conserved and region-specific microbiome disruptions, revealing a consistent ecological impact of *T. trichiura* on gut microbial communities. This study extends previous research on helminth-microbiota interactions [14], by demonstrating that *T. trichiura* induces significant taxonomic and functional shifts that transcend geographic variation, providing a broader, comparative perspective on how soil-transmitted helminths influence gut microbial ecology.

One of the most striking findings is the consistent depletion of SCFA-producing bacteria across all study populations. Infected individuals exhibited a marked reduction in butyrate- and acetate-producing taxa, including *Blautia* sp. MSJ 9 and *Holdemanella biformis*, alongside a depletion of key fermentation pathways, such as PWY-7254 and PWY-3801. These SCFAs play a critical role in maintaining colonic health, reinforcing epithelial integrity, and modulating immune responses [35]. Their depletion in infected individuals suggests a potential weakening of gut barrier function, which could have implications for host susceptibility to inflammation and reinfection. Importantly, this study provides the first multi-regional evidence that *T. trichiura* consistently alters SCFA metabolism, regardless of environmental or dietary differences, reinforcing the predictable ecological pressures imposed by this parasite.

This study also demonstrates for the first time that *T. trichiura*'s impact on specific microbial taxa is highly context dependent. While core microbial functions were similarly affected across regions, taxa such as *Prevotella* and *Streptococcus* exhibited region-specific variations in relative abundance, likely reflecting local environmental pressures contributing to shaping microbiome resilience to helminth-induced dysbiosis. Notably, fermentation pathways were strongly depleted in infected individuals from Laos, while in Côte d'Ivoire and Tanzania, some of these pathways remained stable or were even enriched. These findings suggest that

while *T. trichiura* exerts a consistent functional impact on the microbiome, the host-

microbiota response is also influenced by region-specific dietary and environmental factors, a

novel insight that may have implications for future intervention strategies.

The niche-specificity of *T. trichiura* in the large intestine provides further context for these

findings. Unlike *Ascaris lumbricoides* and hookworms, which colonize the small intestine, T.

trichiura inhabits the cecum and proximal colon [36], microbial environments characterized

by high fermentative activity and mucin metabolism [35]. This distinction is crucial, as

helminth infections elicit both local and distal immunomodulatory responses, but the effects

are notably stronger at the site of infection [37]. This means that stool samples reliably

capture microbiome alterations at the infection site [38], thus strengthening the clinical

relevance of this study. By focusing exclusively on *T. trichiura*, this study provides the most

detailed characterization to date of how a large-intestinal helminth uniquely reshapes gut

microbial ecology.

Functional analysis also revealed significant enrichment of mucin degradation pathways in

infected individuals, indicative of a microbial shift toward host-derived glycan metabolism.

Specifically, consistent enrichment of the GLCMANNANAUT-PWY pathway

(autodegradation of glucosamine- and mannose-containing polysaccharides, key mucin

components), together with an increased abundance of known mucin-degrading taxa such as

Ruminococcus and *Bacteroides*, suggests altered mucus dynamics potentially facilitating

parasite persistence. These findings align with previous research linking helminth infections

to increased mucin degradation [15, 39], but this study is the first to demonstrate these shifts

across multiple geographic regions, reinforcing their potential role in *T. trichiura*'s ability to

establish chronic infections.

Beyond functional disruptions, network-based microbial interactions provide novel insights into community-level stability in the infected gut. Network analyses revealed reduced microbial connectivity and clustering in infected individuals, suggesting that *T. trichiura* infection leads to a less cohesive and more fragmented microbiome structure. The emergence of opportunistic taxa such as *Segatella copri* as network hubs in infected individuals suggests that these species may exploit helminth-induced disruptions, potentially reinforcing dysbiosis. Notably, similar microbial network disruptions were reported in helminth-infected indigenous populations in Malaysia [14], but this study extends these findings by demonstrating that *T. trichiura*-induced network instability is a globally recurrent feature, independent of geography.

The potential clinical implications of these findings are significant, particularly in regions where *T. trichiura* remains endemic. The depletion of SCFA-producing bacteria and the enrichment of mucin-degrading taxa suggest that *T. trichiura* infection compromises gut barrier integrity and immune homeostasis, potentially increasing the risk of inflammation, enteric co-infections, and reinfection cycles through a microbiome-mediated process. Furthermore, the observed functional shifts may favour parasite persistence, exacerbating chronic infection burdens in endemic settings. These findings highlight the dual role of the microbiota during *T. trichiura* infection: on one hand acting as a protective barrier, and on the other hand potentially facilitating parasite colonization. A better understanding of these dynamics may pave the way for microbiome-based interventions, such as prebiotic or probiotic therapies aimed at restoring microbial stability and enhancing resistance to reinfection.

Nonetheless, this study has some limitations. The cross-sectional design prevents causal inferences about the relationship between *T. trichiura* infection and microbiome changes. Longitudinal studies that track microbiome dynamics during infection, treatment, and

reinfection are necessary to establish temporal relationships and better understand the mechanisms driving dysbiosis. Additionally, while shotgun metagenomics offers comprehensive insights into microbial functional potential, it does not capture active transcription or metabolic fluxes. Integrating metatranscriptomics or metabolomics in future studies would further elucidate the active processes underlying helminth-induced microbiome disruptions.

In conclusion, this study provides the strongest evidence to date that *T. trichiura* infection leads to both conserved and region-specific disruptions in the gut microbiome, affecting SCFA metabolism, mucin degradation, and microbial network stability. These findings expand previous work by demonstrating that helminth-induced dysbiosis follows a predictable functional trajectory, while also being shaped by local ecological and dietary factors. Understanding these dynamics is critical for developing targeted microbiome-based strategies to mitigate the burden of soil-transmitted helminths and improve gut health in endemic regions.

Ethical approval and consent to participate:

The study protocol was approved by independent ethics committees in Côte d'Ivoire (reference numbers 088–18/MSHP/CNESVS-km and ECCI00918), Laos (reference number 093/NECHR), Pemba (Tanzania, reference number ZAMREC/0003/Feb/2018), and Switzerland (reference number BASEC Req-2018-00494). This trial was registered under the number NCT03527732 on ClinicalTrials.gov. Participants were informed about the purpose of the study and informed consent forms were obtained during information sessions

Consent for publication

Not applicable

Availability of data and material

Shotgun sequence data supporting the findings of this study have been deposited in the NCBI

Short Read Archive with the primary accession code PRJNAXXXX.

Competing interests

The authors declare that they have no competing interests

Funding

We are grateful to the European Research Council (No. 101019223) for financial support.

Author contributions

PHHS: study design, research design, project supervision, experimental work, statistical

analyses, figure generation, writing of the initial manuscript, manuscript editing; **EH, AS, SS,**

and JTC: study design, conducted field work (sample collection, handling, coordination of

treatment, parasitological work, and data curation); **JD and NR:** experimental work (DNA

isolation, Nanopore sequencing), manuscript editing; **JK**: study design, research design,

project supervision, writing of the initial manuscript, manuscript editing.

Acknowledgements

We are deeply grateful to the dedicated trial team members from the Centre Suisse de

Recherches Scientifiques and Université Félix Houphouët-Boigny in Abidjan, Côte d'Ivoire;

the Lao Tropical and Public Health Institute in Vientiane, Laos; and the Public Health

Laboratory–Ivo de Carneri in Chake Chake, Pemba Island, for their hard work and expertise

in both field and laboratory activities. We also extend our heartfelt thanks to the local village

and medical authorities in Chake Chake district, Pemba Island; Nambak district, Luang

Prabang province, Laos; and the districts of Dabou and Jacqueville in Côte d'Ivoire, whose

collaboration and support were invaluable in making this work possible. We thank Christian

Beisel from the Genomics Facility Basel for their support during the shotgun sequencing

experiments. We are grateful for the access and support to/from the sciCORE (https://scicore.unibas.ch) scientific computing center at the University of Basel.

Declaration on the use of AI

In the preparation of this manuscript, AI-based tools (ChatGPT and Copilot) were used to assist in language editing, grammatical correction, and improving the clarity of the text. These tools were employed solely for editorial purposes and did not contribute to the conceptualization, data analysis, or scientific conclusions of this work. All scientific content, interpretations, and conclusions presented in this manuscript are the sole responsibility of the authors, who have thoroughly reviewed and approved the final version of the text.

Figures:

Figure 1. Study profile.

Figure 2. Community-level changes associated with infection status, stratified by country. Panel A: Alpha diversity metrics stratified by country and infection status. The left panel shows species richness, the middle panel shows the Shannon diversity index, and the right panel displays the Berger-Parker dominance index. **Panel B:** Non-metric multidimensional scaling (NMDS) plot based on the Bray-Curtis dissimilarity index. Results from community comparisons using PERMANOVA and community dispersion using PERMDISP are shown, with significant results highlighted in bold. **Panel C:** Core microbiome analysis illustrating baseline taxonomic differences across the three study settings. **Panel D:** Heatmap with overall sample clustering based on Robust Aitchison dissimilarity and the Ward method for cladogram construction. The original eggs per gram (EPG) values of stool are displayed at the bottom of each sample.

Figure 3. Species-level differences between infection groups, stratified by country. Panel A: Prevalence-based comparison showing phylogenetic relationships of species that were significantly more or less prevalent in at least one country. Statistical significance was determined using Fisher's exact test with Benjamini-Hochberg false discovery rate (FDR) correction (adjusted $P < 0.05$). **Panel B:** Comparison of relative abundances between infection status groups, stratified by country. Three statistical tests were applied: Kruskal-Wallis test, LEfSe differential abundance pipeline, and MaAsLin2 (adjusted for the country variable). Only species that were statistically significant across all three tests were retained. The log2 fold-change values, generated using MaAsLin2, are displayed to represent effect sizes. **Panel C:** Comparison of cumulative relative abundances of species significantly enriched or depleted in one versus two (or more) study sites. Statistical comparisons were performed using the Kruskal-Wallis test, with Bonferroni correction for P-values (adjusted $P < 0.05$).

Figure 4. Effect of infection status on the microbiome network correlation structure. SpiecEasi analysis of taxonomic profiles stratified by country and infection status. The visualization on the 50 most abundant species in the different groups, the metrics were calculated using the 200 most abundant species in each group. Network metrics are as follows: E = edges, De = density, PL = average path length, Di = diameter, and CC = clustering coefficient. The numbers in the correlation graphs correspond to the species listed in the table at the bottom of the figure. Inf. = infected; Uninf. = uninfected.

Figure 5. Community-level functional changes associated with infection status, stratified by country. Panel A: Venn diagram showing the enzyme commission (EC) numbers found in each study setting, stratified by infection status and showing the overlapping portion. **Panel B:** Non-metric multidimensional scaling (NMDS) plot based on the Bray-Curtis dissimilarity index applied on the EC abundance tables. Results from community

comparisons using PERMANOVA and community dispersion using PERMDISP are shown, with significant results highlighted in bold. **Panel C:** Volcano plots based on comparison by infection status of normalized EC abundance tables using MaAsLin2, stratified by country. **Panel D:** Averaged Metacyc superclass abundance comparison by infection status stratified by country. Panel E: Fold-change abundance difference of Metacyc pathways between infected and uninfected patients calculated with MaAsLin2, stratified by country. **Panel F:** Fold-change (FC) comparison of Metacyc pathways associated with infection status between countries. To visualize the monotonic trend, we applied LOWESS smoothing (represented by colored triangles with black outlines) using a proportion of 0.6, balancing smoothness with the capture of local variations. CI = Côte d'Ivoire; LA = Laos; TA = Tanzania.

References:

1. Zmora N, Suez J, Elinav E: **You are what you eat: diet, health and the gut microbiota**. *Nature reviews Gastroenterology & hepatology* 2019, **16**(1):35-56.
2. Levy M, Kolodziejczyk AA, Thaiss CA, Elinav E: **Dysbiosis and the immune system**. *Nature Reviews Immunology* 2017, **17**(4):219-232.
3. Schneeberger PH, Coulibaly JT, Panic G, Daubenberger C, Gueuning M, Frey JE, Keiser J: **Investigations on the interplays between *Schistosoma mansoni*, praziquantel and the gut microbiome**. *Parasites & vectors* 2018, **11**:1-12.
4. Leung JM, Graham AL, Knowles SC: **Parasite-microbiota interactions with the vertebrate gut: synthesis through an ecological lens**. *Frontiers in microbiology* 2018, **9**:843.
5. Hodžić A, Dheilly NM, Cabezas-Cruz A, Berry D: **The helminth holobiont: a multidimensional host–parasite–microbiota interaction**. *Trends in Parasitology* 2023, **39**(2):91-100.
6. Loke P, Lim Y: **Helminths and the microbiota: parts of the hygiene hypothesis**. *Parasite immunology* 2015, **37**(6):314-323.
7. Else KJ, Keiser J, Holland CV, Grencis RK, Sattelle DB, Fujiwara RT, Bueno LL, Asaolu SO, Sowemimo OA, Cooper PJ: **Whipworm and roundworm infections**. *Nature Reviews Disease Primers* 2020, **6**(1):44.
8. Jourdan PM, Lamberton PH, Fenwick A, Addiss DG: **Soil-transmitted helminth infections**. *The lancet* 2018, **391**(10117):252-265.
9. Lawson MA, Roberts IS, Grencis RK: **The interplay between *Trichuris* and the microbiota**. *Parasitology* 2021, **148**(14):1806-1813.

10.   Reynolds LA, Finlay BB, Maizels RM: **Cohabitation in the intestine: interactions**
among helminth parasites, bacterial microbiota, and host immunity. *The Journal*
of Immunology 2015, **195**(9):4059-4066.

11.   Schachter J, Alvarinho de Oliveira D, Da Silva CM, de Barros Alencar ACM, Duarte
M, da Silva MMP, Ignácio ACdPR, Lopes-Torres EJ: **Whipworm infection**
promotes bacterial invasion, intestinal microbiota imbalance, and cellular
immunomodulation. *Infection and Immunity* 2020, **88**(3):10.1128/iai. 00642-00619.

12.   Chen H, Mozzicafreddo M, Pierella E, Carletti V, Piersanti A, Ali SM, Ame SM,
Wang C, Miceli C: **Dissection of the gut microbiota in mothers and children with**
chronic *Trichuris trichiura* infection in Pemba Island, Tanzania. *Parasites &*
Vectors 2021, **14**:1-13.

13.   Sargsian S, Chen Z, Lee SC, Robertson A, Thur RS, Sproch J, Devlin JC, Tee MZ, Er
YX, Copin R: **Clostridia isolated from helminth-colonized humans promote the**
life cycle of *Trichuris* species. *Cell reports* 2022, **41**(9).

14.   Tee MZ, Er YX, Easton AV, Yap NJ, Lee IL, Devlin J, Chen Z, Ng KS, Subramanian
P, Angelova A: **Gut microbiome of helminth-infected indigenous Malaysians is**
context dependent. *Microbiome* 2022, **10**(1):214.

15.   Holm JB, Sorobetea D, Kiilerich P, Ramayo-Caldas Y, Estellé J, Ma T, Madsen L,
Kristiansen K, Svensson-Frej M: **Chronic *Trichuris muris* infection decreases**
diversity of the intestinal microbiota and concomitantly increases the abundance
of lactobacilli. *PloS one* 2015, **10**(5):e0125495.

16.   Hürlimann E, Keller L, Patel C, Welsche S, Hattendorf J, Ali SM, Ame SM, Sayasone
S, Coulibaly JT, Keiser J: **Efficacy and safety of co-administered ivermectin and**
albendazole in school-aged children and adults infected with *Trichuris trichiura*
in Côte d'Ivoire, Laos, and Pemba Island, Tanzania: a double-blind, parallel-
group, phase 3, randomised controlled trial. *The Lancet infectious diseases* 2022,
22(1):123-135.

17.   Bärenbold O, Raso G, Coulibaly JT, N'Goran EK, Utzinger J, Vounatsou P:
Estimating sensitivity of the Kato-Katz technique for the diagnosis of
***Schistosoma mansoni* and hookworm in relation to infection intensity**. *PLoS*
neglected tropical diseases 2017, **11**(10):e0005953.

18.   Hillmann B, Al-Ghalith GA, Shields-Cutler RR, Zhu Q, Gohl DM, Beckman KB,
Knight R, Knights D: **Evaluating the information content of shallow shotgun**
metagenomics. *Msystems* 2018, **3**(6):10.1128/msystems. 00069-00018.

19.   Blanco-Míguez A, Beghini F, Cumbo F, McIver LJ, Thompson KN, Zolfo M, Manghi
P, Dubois L, Huang KD, Thomas AM: **Extending and improving metagenomic**
taxonomic profiling with uncharacterized species using MetaPhlAn 4. *Nature*
Biotechnology 2023, **41**(11):1633-1644.

20.   Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2**. *Nature*
Methods 2012, **9**(4):357-359.

21.   Lu Y, Zhou G, Ewald J, Pang Z, Shiri T, Xia J: **MicrobiomeAnalyst 2.0:**
comprehensive statistical, functional and integrative analysis of microbiome
data. *Nucleic Acids Research* 2023, **51**(W1):W310-W318.

22. Caspi R, Billington R, Keseler IM, Kothari A, Krummenacker M, Midford PE, Ong
WK, Paley S, Subhraveti P, Karp PD: **The MetaCyc database of metabolic**
pathways and enzymes-a 2019 update. *Nucleic acids research* 2020, **48**(D1):D445-
D453.

23. Team RC: **A language and environment for statistical computing**. *R Foundation*
for Statistical Computing, Vienna, Austria 2024.

24. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara R, Simpson GL,
Solymos P, Stevens MHH, Wagner H: **Package 'vegan'**. *Community ecology*
package, version 2013, **2**(9):1-295.

25. Lahti L, Shetty S, Blake T, Salojarvi J: **Tools for microbiome analysis in R**. *Version*
2017, **1**(5):28.

26. Gu Z: **Complex heatmap visualization**. *Imeta* 2022, **1**(3):e43.

27. Kurtz ZD, Müller CL, Miraldi ER, Littman DR, Blaser MJ, Bonneau RA: **Sparse and**
compositionally robust inference of microbial ecological networks. *PLoS*
computational biology 2015, **11**(5):e1004226.

28. Gustavsen JA, Pai S, Isserlin R, Demchak B, Pico AR: **RCy3: Network biology**
using Cytoscape from within R. *F1000Research* 2019, **8**.

29. Csardi G, Nepusz T, Traag V, Horvát S, Zanini F, Noom D, Müller K: **Igraph:**
Network analysis and visualization. *R package version* 2023, **1**(0.9002).

30. Rupert Jr G: **Simultaneous statistical inference**. 2012.

31. Benjamini Y, Hochberg Y: **Controlling the false discovery rate: a practical and**
powerful approach to multiple testing. *Journal of the Royal statistical society:*
series B (Methodological) 1995, **57**(1):289-300.

32. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C:
Metagenomic biomarker discovery and explanation. *Genome biology* 2011, **12**:1-
18.

33. Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, Tickle TL,
Weingart G, Ren B, Schwager EH: **Multivariable association discovery in**
population-scale meta-omics studies. *PLoS computational biology* 2021,
17(11):e1009442.

34. Organization WH: **Assessing schistosomiasis and soil-transmitted helminthiases**
control programmes: monitoring and evaluation framework. 2024.

35. Koh A, De Vadder F, Kovatcheva-Datchary P, Bäckhed F: **From dietary fiber to**
host physiology: short-chain fatty acids as key bacterial metabolites. *Cell* 2016,
165(6):1332-1345.

36. Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, Hotez PJ: **Soil-**
transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *The*
lancet 2006, **367**(9521):1521-1532.

37. Maizels RM, Smits HH, McSorley HJ: **Modulation of host immunity by helminths:**
the expanding repertoire of parasite effector molecules. *Immunity* 2018,
49(5):801-818.

38. Donaldson GP, Lee SM, Mazmanian SK: **Gut biogeography of the bacterial**
microbiota. *Nature Reviews Microbiology* 2016, **14**(1):20-32.

39.   Ramanan D, Bowcutt R, Lee SC, Tang MS, Kurtz ZD, Ding Y, Honda K, Gause WC,
Blaser MJ, Bonneau RA: **Helminth infection promotes colonization resistance via**
type 2 immunity. *Science* 2016, **352**(6285):608-612.

Figure 1. Study profile.

Figure 2. Community-level changes associated with infection status, stratified by country. Panel A: Alpha diversity metrics stratified by country and infection status. The left panel shows species richness, the middle panel shows the Shannon diversity index, and the right panel displays the Berger-Parker dominance index. **Panel B:** Non-metric multidimensional scaling (NMDS) plot based on the Bray-Curtis dissimilarity index. Results from community comparisons using PERMANOVA and community dispersion using PERMDISP are shown, with significant results highlighted in bold. **Panel C:** Core microbiome analysis illustrating baseline taxonomic differences across the three study settings. **Panel D:** Heatmap with overall sample clustering based on Robust Aitchison dissimilarity and the Ward method for cladogram construction. The original eggs per gram (EPG) values of stool are displayed at the bottom of each sample.

Figure 3. Species-level differences between infection groups, stratified by country. Panel A: Prevalence-based comparison showing phylogenetic relationships of species that were significantly more or less prevalent in at least one country. Statistical significance was determined using Fisher's exact test with Benjamini-Hochberg false discovery rate (FDR) correction (adjusted $P < 0.05$). **Panel B:** Comparison of relative abundances between infection status groups, stratified by country. Three statistical tests were applied: Kruskal-Wallis test, LEfSe differential abundance pipeline, and MaAsLin2 (adjusted for the country variable). Only species that were statistically significant across all three tests were retained. The log2 fold-change values, generated using MaAsLin2, are displayed to represent effect sizes. **Panel C:** Comparison of cumulative relative abundances of species significantly enriched or depleted in one versus two (or more) study sites. Statistical comparisons were performed using the Kruskal-Wallis test, with Bonferroni correction for P-values (adjusted $P < 0.05$).

Figure 4. Effect of infection status on the microbiome network correlation structure. SpiecEasi analysis of taxonomic profiles stratified by country and infection status. The visualization on the 50 most abundant species in the different groups, the metrics were calculated using the 200 most abundant species in each group. Network metrics are as follows: E = edges, De = density, PL = average path length, Di = diameter, and CC = clustering coefficient. The numbers in the correlation graphs correspond to the species listed in the table at the bottom of the figure. Inf. = infected; Uninf. = uninfected.

Figure 5. Community-level functional changes associated with infection status, stratified by country. Panel A: Venn diagram showing the enzyme commission (EC) numbers found in each study setting, stratified by infection status and showing the overlapping portion. Panel B: Non-metric multidimensional scaling (NMDS) plot based on the Bray-Curtis dissimilarity index applied on the EC abundance tables. Results from community comparisons using PERMANOVA and community dispersion using PERMDISP are shown, with significant results highlighted in bold. Panel C: Volcano plots based on comparison by infection status of normalized EC abundance tables using MaAsLin2, stratified by country. Panel D: Averaged Metacyc superclass abundance comparison by infection status stratified by country. Panel E: Fold-change abundance difference of Metacyc pathways between infected and uninfected patients calculated with MaAsLin2, stratified by country. Panel F: Fold-change comparison of Metacyc pathways associated with infection status between countries. To visualize the monotonic trend, we applied LOWESS smoothing (represented by colored triangles with black outlines) using a proportion of 0.6, balancing smoothness with the capture of local variations. CI = Côte d'Ivoire; LA = Laos; TA = Tanzania.