In vitro detection of canine anti-human antibodies following intratumoral injection

of the hu14.18-IL2 immunocytokine in spontaneous canine melanoma

Short title: Analysis of canine anti-human antibodies

Andrew T. Kosharek[1¶], Cindy L. Zuleger[2,3¶], William S. Glass[4], Jens Eickhoff [3,5], Paul M.

Sondel[3,6,7], David M. Vail[3,8], Mark R. Albertini[2,3,9,10]*

[1]University of Wisconsin School of Medicine and Public Health, Madison, WI, United
States of America
11
12 [2]University of Wisconsin School of Medicine and Public Health, Department of Medicine,
13 Madison, WI, United States of America
14
15 [3]University of Wisconsin Carbone Cancer Center, Madison, WI, United States of
16 America
17
18 [4]University of Wisconsin-Madison, Madison, WI, United States of America
19
20 [5]University of Wisconsin School of Medicine and Public Health, Department of
21 Biostatistics & Medical Informatics, Madison, WI, United States of America
22
23 [6]University of Wisconsin School of Medicine and Public Health, Department of Human
24 Oncology, Madison, WI, United States of America
25
26 [7]University of Wisconsin School of Medicine and Public Health, Department of
27 Pediatrics, Madison, WI, United States of America
28
29 [8]University of Wisconsin School of Veterinary Medicine, Department of Medical
30 Sciences, Madison, WI, United States of America
31
32 [9]University of Wisconsin School of Medicine and Public Health, Department of
33 Dermatology, Madison, WI, United States of America
34
35 [10]The Medical Service, William S. Middleton Memorial Veterans Hospital, Madison,
36 Wisconsin, United States of America
37

38   * Corresponding author

39   E-mail: mralbert@wisc.edu (MRA)

40   ¶These authors contributed equally to this work.

Abstract

Background: Canine and human malignant melanoma are naturally occurring cancers with many similarities, making the dog an important parallel patient population to study both diseases. However, development of canine anti-human antibodies (CAHA) needs to be considered when evaluating humanized biotherapeutics in dogs.

Objectives: Characterize CAHA in sera from dogs with spontaneous melanoma receiving radiotherapy and intratumoral immunocytokine (IT-IC) with humanized 14.18-IL2.

Methods: Serum samples were obtained pre-treatment and at several post-treatment times from 12 dogs with locally advanced or metastatic melanoma treated with radiotherapy to the primary site and regional lymph nodes (when clinically involved) followed by IT-IC of humanized 14.18-IL2. Two CAHA assays were developed. A sandwich enzyme-linked immunosorbent assay (ELISA) was developed to detect antibodies against the humanized IgG component of hu14.18-IL2. A flow cytometry assay was developed to determine the ability of CAHA to inhibit binding of a mouse anti-GD2 monoclonal antibody to its target.

Results: Post-treatment sera from 7 of 12 dogs developed CAHA levels over pre-treatment that were identified by ELISA as significant increases at Day 30 and/or Day 60. Day 10, Day 30, and Day 60 post-treatment sera from 10 of 12 dogs significantly inhibited the binding of anti-GD2 monoclonal antibody to its target compared to pre-treatment. Significant binding inhibition was also detected in 2 of 12 dogs after local RT but before IT-IC (Day 1). Normal canine sera did not mediate binding inhibition.

63   **Conclusions:** This study advances CAHA detection strategies and reports the kinetics of

64   CAHA following IT-IC in dogs with spontaneous melanoma.

Introduction

Companion (pet) dogs with spontaneous tumors provide unique and clinically relevant insights into human tumor biology, therapy, and drug development [1, 2]. Dogs develop spontaneous tumors, similar to humans [3]. This contrasts with many laboratory mouse models where tumors are experimentally induced [4]. Moreover, the molecular and genetic characteristics of many canine cancers resemble those of human cancers [5]. While no model perfectly recapitulates human cancer, the companion dog melanoma patient population holds a distinctive position in oncology research and has received attention from the scientific community as being somewhat parallel to human melanoma patients, particularly for evaluation of immunotherapy studies [6-10]. In that regard, development of antidrug antibodies (ADAs) against immunotherapeutic monoclonal antibodies (mAbs) is well documented in human patients, and the presence of ADAs can impact the safety and efficacy of immunotherapy [11-15]. Moreover, the immune response is highly species-specific and relatively few immunotherapeutic mAbs are caninized, i.e., canine speciated [16]. Therefore, it will be critical to monitor ADAs in the setting of canine immunotherapy, particularly when the immunotherapeutic is a mAb of a species 'foreign' to the dog (e.g., a humanized mAb).

The humanized mAb Hu14.18-IL2 and the chimeric ch14.18-IL2, both GD2-targeting mAbs linked to IL2 and known as immunocytokines (IC), have been studied *in vitro* and in preclinical mouse models with GD2-expressing tumors [17-20]. Clinical trials evaluating intravenous (IV) delivery of hu14.18-IL2 IC in patients with melanoma or neuroblastoma have demonstrated immune activation and had manageable safety

profiles [21-25]. Although hu14.18-IL2 is a humanized mAb, some patients still

developed antibodies to the therapeutic IC [22, 26]. In these patients, antibodies were

detected to both the idiotype, i.e., the antigen binding end, and the Fc-IL2 end of the IC

[22, 26]. A substantial fraction of patients develop a detectible anti-IC antibody

response, and some patients develop a strong anti-IC response and functional, in vivo

"neutralization" of detectible IC. Namely, following detection of a strong anti-IC

response, blood obtained immediately after an IC infusion does not contain the

anticipated level of detectible IC, based on the IC level in the blood obtained from

samples collected prior to development of the anti-IC response [22, 26]. A neutralizing

anti-human antibody (HAHA) or anti-chimeric antibody (HACA) has been reported for

some patients treated with humanized or chimeric hu14.18 mAb [27, 28].

The immunogenicity of the humanized hu14.18-IL2 is potentially a greater

concern when administering this molecule to a "foreign" species, e.g., to a mouse or

dog, as ADAs could interfere with the pharmacokinetics of this therapy, potentially

reducing its efficacy [29]. When huKS1/4-IL2, a humanized antibody directed against

the human epithelial cell adhesion molecule, was administered intravenously to mice

with adenocarcinoma, the mice mounted significant ADAs in the form of mouse anti-

human antibodies (MAHAs) resulting in rapid clearance of additional doses of the

therapeutic [18]. Delivering hu14.18-IL2 intratumorally in mouse models also induced a

MAHA response, similar to what was seen following IV-IC therapy [30]. In addition,

ADAs in dogs have been observed in other studies, including one report following

subcutaneous administration of a therapeutic IC [31-33]. However, the development of

canine anti-human antibodies (CAHAs) following intratumoral immunotherapy with an IC

has not been reported.

Here we present CAHA assay development and CAHA monitoring of companion

dogs with spontaneous melanoma enrolled in a clinical trial of local RT plus intratumoral

hu14.18-IL2. An in-house ELISA was developed and used to monitor CAHA against the

IgG portion of the IC. Previous studies have demonstrated the ability of ELISA to detect

ADAs against antibody-based biopharmaceuticals [26, 34-37]. Our ELISA is

customizable, cost-effective, and uses common laboratory reagents and equipment. We

also examined if CAHA interferes with the ability of the GD2-binding mAb part of the IC

to recognize GD2 in a cell-based flow cytometry assay. Details about safety, clinical

activity, and tumor biopsy analyses from this trial, as well as another trial, are separately

reported [38, 39].

Material and Methods

Trial eligibility and enrollment

Client-owned companion (pet) dogs weighing ≥5 kg with a readily palpable and

accessible oral malignant melanoma (histologically confirmed) of at least 2.0 cm

(longest diameter) were included in the study. Both treatment naïve dogs and dogs with

surgically recurrent tumors were allowed, provided the recurrence met the minimum

size criteria. All dogs were required to have a pretreatment constitutional clinical sign

status of 0 or 1 (normal or mild lethargy over baseline; diminished activity from pre-

disease level but able to function as an acceptable pet) according to VCOG-CTCAE v2

criteria at study entry [40]. Dogs determined to have any significant co-morbid illness

were excluded.

Study design

Client-owned companion dogs with locally advanced or metastatic melanoma

were randomized (6 dogs per Arm) to receive either a single 8 Gy fraction (Arm A) or

three 8 Gy fractions delivered over 1 week (Arm B) to the primary site and regional

lymph nodes (when clinically involved) with the single or last fraction 5 days prior to IT-

IC at 12 mg/m^2 on 3 consecutive days. Prior to treatment, a baseline physical

examination with tumor measurements was performed, blood was collected for clinical

assessments, and blood was collected for banking of serum, plasma, and peripheral

blood mononuclear cells (PBMC) for subsequent *in vitro* analysis. External beam

radiation therapy (RT) was delivered by image-guided intensity modulated radiation

using helical tomotherapy (TomoTherapy HiArt Treatment System, Accuray Inc.,

Sunnyvale, CA, USA). The hu14.18-IL2 was provided as lyophilized vials (4 mg/vial;

each ml prior to lyophilization contained 4 mg/ml hu14.18-IL2, 2% sucrose, 80mM L-

arginine, 10mM citric acid, 0.2% polysorbate 20, pH 5.5) and was reconstituted with

either 0.5 ml or 0.25 ml 0.9% (w/v) NaCl for injection at a concentration of either 8

mg/ml or 16 mg/ml. The total volume was administered using multiple needle re-

directions into the target lesion within and surrounding the lesion. Injections were

performed on 3 consecutive days. PBMC were obtained at pre-treatment and various

times post-treatment. Etomidate (1-2 mg/kg, IV) and/or Propofol (4-5 mg/kg IV) were

used to induce anesthesia in dogs receiving RT. Isoflurane (1.2-1.3%, inhalation) and/or

Sevoflurane (2.4%, inhalation) were used to maintain anesthesia and the gas

percentages were adjusted depending on the canine patient's clinical response to

anesthesia under guidance of the clinician. The duration of anesthesia was

approximately 30 minutes. Butorphanol (0.2-0.4 mg/kg IM or IV) and/or midazolam (0.2

mg/kg IV) were used to sedate the dogs during IT-IC administration, the duration of

which was approximately 45 minutes. Pulse, respiratory and heart rates were monitored

throughout anesthesia and at least every 15 minutes until the dog was ambulatory.

Lidocaine was administered at the site of the IT-IC administration for pain (0.5-0.75 ml

2% Lidocaine, intradermal). Following IT-IC administration, pain was alleviated with

analgesics for 3 days or longer if the clinician or owner perceived that the dog was

experiencing continued discomfort. Analgesics included either Carprofen (2.2 mg/kg,

orally, 2x daily), Deracoxib (3-4 mg/kg, orally, 1x daily), Meloxicam (0.1-0.2 mg/kg,

orally, 1x daily), and/or Tramadol (1-5 mg/kg, orally, every 8 hours). Physical

examination and canine patient history were assessed at each clinic visit. Dogs were

monitored by the dog's owner and clinician for animal health and behavior, as well as

the development of adverse events (AE). AEs were graded and attributed according to

VCOG-CTCAE v2 [40]. The attending clinician prescribed appropriate treatment and/or

supportive care if an AE was observed. All efforts were made to minimize suffering. The

study endpoint was the time the dog developed progressive disease as determined by

physical examination and/or thoracic radiographs. Owners could remove a dog from the

study for any reason, or the attending clinician could remove a dog from the study if it

was determined there were severe AEs not manageable with supportive care.

Euthanasia was not a study endpoint nor was it part of this study and was only

performed at the request of a dog's owner. If requested, the American Veterinary

Medical Association guidelines were followed [41]. The treatment schema and blood

collection times are shown in Fig 1.

Fig 1. Treatment Schema. All dogs receive radiation therapy (RT) to the local site

either in 1x 8 Gy (Arm A) or 3x 8 Gy fractions over 1 week (Arm B). RT was delivered on

Day -4 (Arm A) or on Days -8, -6 -4 (Arm B). Intratumoral (IT) injection of IC is

administered (IT-IC) on three consecutive days starting five days after completion of RT.

All days were allowed $\pm$ 1-2 days leeway for flexible scheduling around holidays, subject

vacations, and clinic scheduling.

IACUC approvals

The authors confirm that the appropriate ethical review committee approval has

been received. All procedures and treatments performed on client-owned companion

dogs were approved by the Institutional Animal Care and Use Committees of the

University of Wisconsin-Madison School of Veterinary Medicine (Approval V006037)

and by the Animal Care Committee at the William S. Middleton Memorial Veterans

Hospital. Written informed consent was obtained from all companion dog caregivers

prior to entry into this trial. Protocol treatments were administered at the University of

Wisconsin Veterinary Care (UWVC) hospital.

Canine Anti-Human ELISA

A standard protocol was adapted to develop an ELISA to detect canine

antibodies against human IgG [42]. Nunc MaxiSorp™ Flat-Bottom 96-well plates

(Thermo Scientific Inc., USA) were coated with ChromPure Human IgG (Jackson

ImmunoResearch Laboratories, Inc., USA) [10μg/mL, 50μL/well in ELISA Coating Buffer

(Bio-Rad Laboratories, Inc., USA)], which served as the "capture" agent, and incubated

overnight at 4°C. The coating solution was removed, and the wells washed three times

(3x) with distilled water (DI). The plate was blocked with Neptune™ Block

(ImmunoChemistry Technologies, USA) (200µL/well) for 1 hour at 25°C followed by

washing as above. Canine serum was diluted 1:20 in PBS and added to wells in

quadruplicate at 50µL/well and incubated for 1 hour at 25°C. A murine mAb against

human IgG (ab436, Abcam, USA) was used as a positive control, as we are not aware

of any commercially available canine anti-human IgG to serve as a positive control. The

mAb was diluted 1:1,000 in PBS and added to wells in quadruplicate at 50µL/well and

incubated for 1 hour at 25°C. PBS was added to wells as a negative control. Wells were

washed as above. Rabbit anti-mouse IgG (H+L) antibody, affinity purified against human

serum proteins conjugated to horseradish peroxidase (HRP) (# 315-035-045, Jackson

ImmunoResearch Laboratories Inc., USA) was the secondary (i.e.: "detection") antibody,

was diluted 1:40,000 in PBS, added to wells at 75μL/well, and incubated for 30 minutes

(min.) at 25°C. Wells were washed as above. The plate was developed with1-Step™

Ultra TMB-ELISA solution (Thermo Scientific Inc., USA) (75μL/well) and incubated for

15 min. at 25°C. The reaction was stopped by addition of TMB stop solution (Sera Care,

USA) (75μL/well), and the absorbance at 450nm read on a SpectraMax M3 Micro-Plate

reader.

Cell culture

224   The GD2+ M21 human melanoma cell line was cultured in RPMI-1640

225   supplemented with 10% fetal bovine serum (GeminiBio, BenchMark™), 25 mM HEPES,

226   2 mM L-glutamine, 1 mM sodium pyruvate, and 1X non-essential amino acids

227   (Mediatech). M21 was obtained as a gift from Dr. Ralph Reisfeld (Scripps Research

228   Institute, La Jolla, CA, USA).

Detection of GD2 binding inhibition by CAHA using flow

cytometry

Mouse anti-GD2 mAb 14G2a-PE (BioLegend, #357303) (100 μl) was added to

2.5 μl canine sera from this study (Baseline, Day 1, Day 10, Day 30, Day 60) or to

serum from healthy dogs (2.5 μl or 20 μl) (Antibodies Inc, #54-490-0005) and adjusted

to a total volume of 120 μl using 1% FBS/PBS flow cytometry wash buffer (WB). A

sample without canine sera was used as a staining control. The light-protected mixture

was incubated for 30 min. at 4°C. The serum-14G2a mixture was added to GD2+ M21

cells aliquoted to flow cytometry tubes and incubated for 30 min.at 4°C, followed by 2x

washes with WB. Cells were fixed with Cytofix Fixation Buffer (BD, #554655) for 15 min.

at 25°C, washed 1x with WB, resuspended in WB, and held at 4°C until analysis. Data

were acquired on a BD LSR Fortessa, single cells were gated, and the Median

Fluorescence Intensity (MFI) was determined with FlowJo™ Software (BD Life

Sciences). Binding inhibition was calculated as follows where X equals Day 1, Day 10,

Day 30, or Day 60:

$$\% \text{ Binding Inhibition} = 100 \times \frac{\text{Baseline MFI} - \text{Day X MFI}}{\text{Baseline MFI}}$$

247 Statistical Analysis and Considerations

Animal specific kinetic levels of CAHA following IT-IC were summarized in terms of

means and standard errors. Profile plots were generated to examine changes over time.

Changes from baseline to Day 10, 30, and 60 were evaluated using a generalized linear

model. Model assumptions were validated by examining residual and normal

probabilities plots. Dunnett's test was used to control the type I error (<0.05) when

conducting multiple comparisons for changes from baseline to Day 10, 30, and 60. All

reported P values are two-sided and P<0.05 was used to define statistical significance.

Statistical analyses were conducted using SAS software (SAS Institute, Cary NC),

version 9.4. Plots were generated using R software version 4.2.1 with graphical

packages [43-45].

259 Results

260 Companion dog characteristics

261 Twelve dogs met eligibility criteria for the trial and were enrolled and randomized into

262 one of the two treatment groups from May 2020 to December 2021 [38]. Companion

263 dog breed, tumor characteristics, and randomization results are summarized in Table 1.

264 All 12 dogs completed the treatment phase of the protocol, and none experienced

265 significant or unexpected adverse events [38].

Table 1. Characteristics and status of dogs receiving 12 mg/m^2 IC and randomized to RT Arm A or RT Arm B

Patient ID	Breed	Sex	Age (yr)	Wt (kg)	Tumor Site & Stage at Diagnosis	RT Arm[a]	Sera assayed from all timepoints?
ITIC-04	Labrador Retriever	SF	10	38.9	Left Maxilla/Pulmonary nodule (T2N0M1)	A	Yes
ITIC-05	Burnese Mountain	NM	7	43.4	Right hindlimb subcutaneous (T2N2M0)	B	Yes
ITIC-06	Min. Poodle mix	NM	10	6.6	Right Mandible (T2N1M0)	A	No[b]
ITIC-07	Aust. Cattle mix	NM	13	31.2	Left Maxilla (T2N0M0)	B	Yes
ITIC-08	Beagle mix	NM	11	15.8	Soft palate (T2N1M0)	B	Yes
ITIC-09	Great Dane	SF	10	51.7	Left Maxilla (T2N1M1)	A	Yes
ITIC-10	German Shepherd	NM	9	26.3	Right Maxilla (T2N0M0)	A	Yes
ITIC-11	Mix Breed	NM	11	26.5	Right Mandible (T2N0M1)	B	Yes
ITIC-12	Pom mix	SF	11	9.0	Left Maxilla (T3N1M0)	A	No[b]
ITIC-13	Mix Breed	NM	10	7.0	Cutaneous (TxN1M1)	A	Yes
ITIC-14	Collie mix	SF	11	25.0	Left Maxilla (T2N1M0)	B	Yes
ITIC-15	Mix Breed	NM	9	36.1	Lingual (T2N0M0)	B	Yes

IC, immunocytokine; RT, radiation therapy; SF, spayed female; NM, neutered male
[a]Arm A, a single 8 Gy fraction to the primary tumor followed 5 days later with hu14.18-IL2 IC (12 mg/m^2 for 3 consecutive days); Arm B, three 8 Gy fractions delivered on a

Detection of CAHAs by ELISA

We developed an in-house sandwich ELISA to detect antibodies in canine sera

specific to human IgG — the antibody portion of hu14.18 is humanized IgG. For this

assay to work, a secondary antibody specific for, or cross-reactive to, canine

immunoglobulin is required. Additionally, to avoid the need for multiple secondary

reagents, reactivity to mouse immunoglobulin, the species the positive control is raised

in, was preferred. We first tested our potential secondary reagents by optimizing their

ability to generate a strong signal when detecting mouse anti-human IgG mAb as the

positive control (in place of the canine serum), as the mouse anti-human IgG was

serving as the positive control reagent, namely the analyte. Our initial "checkerboard"

trials tested various concentrations of human IgG coating antigen, mouse anti-human

IgG mAb as the analyte, and rabbit anti-dog HRP secondary antibody. As expected, the

measured absorbance was dependent on the concentration of the rabbit anti-dog HRP

secondary, i.e., absorbance decreased as concentration of the secondary decreased. In

contrast, the absorbance was not dependent on the concentration of either the coating

human IgG or the mouse anti-human IgG mAb (data not shown). To address these

issues, we first troubleshot reagents and/or steps, e.g., testing different blocking buffers,

including 5% normal rabbit sera, as well as non-mammalian protein buffers, increasing

the blocking time, and testing different wash buffers. These changes did not resolve the

high background issue, which was also observed in negative control wells, i.e., in wells

incubated with PBS instead of the mouse anti-human IgG mAb.

These data suggested that the rabbit anti-dog HRP secondary may be binding to

the human IgG coating on the plate. Immunoglobulins from different species share

similar protein structures [42]. Therefore, antibodies against one species are likely to

cross-react with a number of other species — a property which, as discussed below,

was leveraged to optimize the assay. We adsorbed the rabbit anti-dog secondary

antibody with excess human IgG; however, recognition of human IgG by this adsorbed

reagent was only slightly minimized (data not shown). We are not aware of a

commercially available anti-dog secondary antibody adsorbed to human serum proteins.

Using the cross-reactivity of different species' immunoglobulins, we next tested a

commercially available rabbit anti-mouse IgG secondary antibody adsorbed to ensure

minimal cross-reactivity with human serum proteins. The human adsorbed rabbit anti-

mouse IgG did not bind to plate-bound human IgG while the cross-reactivity to the

positive control mouse IgG was maintained. This one reagent then permitted the use of

a single secondary antibody to effectively detect both CAHA and mouse-anti-human

antibody (MAHA).

Post-treatment sera from seven of twelve dogs tested positive for anti-human IgG

antibodies determined via our in-house CAHA ELISA (Fig 2). Pre-treatment sera was

used as baseline and as an internal control for each dog's post-treatment samples. Two

of six and five of six dogs in Arm A and Arm B, respectively, demonstrated significant

increases in anti-human IgG compared to the pre-treatment absorbance (Fig 2). The

post-treatment anti-human IgG was significantly increased compared to pre-treatment

anti-human IgG at Day 30 for five dogs and at Day 60 for five dogs. For Arm A, both

samples reaching significance were from Day 30 (K906 and K909). The sera from

K909 no longer had a significant increase in CAHA at Day 60 and no sera was available

for Day 60 CAHA analysis from K906. For Arm B, anti-human IgG antibody levels in

three dogs were elevated similarly at Day 30 and Day 60 (K905, K911, K915), one dog

had an elevated level at Day 30 that declined by Day 60 (K914), and one had an

elevated level that was only detected at Day 60 (K908). No dogs had detectable anti-

human IgG antibodies following RT but before IT-IC (Day 1) or at Day 10 post-

treatment. There were no substantial differences in anti-human IgG levels between

negative control wells with PBS only and pre-treatment samples for the 12 dogs

enrolled in this study.

Fig 2. CAHA is induced in some dogs receiving IT-IC injections. Mean CAHA levels,

measured as O.D. values by ELISA, from sera assayed in quadruplicate collected at

Baseline (before radiation), Day 1 (after radiation, but prior to IT-IC), and Days 10, 30

and 60 (after IT-IC) with standard deviation error bars are shown. Day 60 timepoint was

not available for two dogs, K906 and K912, in Arm A. Increases in absorbance from

Baseline to Day 1, Day 10, Day 30, or Day 60 for each dog were analyzed and

significance indicated as * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Sera from dogs treated with hu14.18-IL2 inhibit in vitro

binding to GD2

We modified a flow cytometry assay to determine the potential for CAHA in

canine sera to inhibit the binding of IC to GD2 on cells [30]. In this assay, sera are first

incubated with 14G2a-PE, a mouse mAb that recognizes GD2 at a dilution of ~1:48.

The mixture is subsequently added to GD2+ M21 cells, and the level of PE fluorescence

is determined. 14G2a is an IgG2a-class switch variant of the mouse IgG3 mAb clone

14.18, and thus has the same idiotype as that of the humanized immunocytokine

hu14.18-IL2 [46]. Consequently, some of the CAHA in dog sera that recognizes 14G2a

may in effect recognize the idiotype shared by 14G2a and hu14.18-IL2, while some of

the CAHA might also recognize other regions of the IC. Thus, in this assay, the PE

fluorescence of the 14G2a-PE mAb bound to the M21 cells will be decreased if any of

the CAHA sin sera from treated subjects recognize the idiotype of 14G2a and effectively

block its binding to GD2 on the M21 cells. In contrast, substantial PE fluorescence will

be detected if the serum antibodies do not block the binding of 14G2a to GD2.

In preliminary experiments, we assayed healthy canine sera for components in

dog serum that might interfere with the flow cytometry assay. Serum is a complex

biologic that contains carbohydrates, proteins, and lipids that can interfere with the

ability of antibodies to bind their targets. Healthy serum was spiked into PBS and

incubated with the 14G2a anti-GD2 mAb before adding the mixture to M21 cells, as

described above. Compared to M21 stained in the absence of healthy sera, the addition

of sera substantially inhibited 14G2a binding to GD2 in a concentration dependent

manner. Substantial inhibition (~50.2%) was observed at 1:6 ratio of normal sera to PBS

whereas at a 1:48 ratio inhibition was largely undetected (> 0.1%) (S1 Fig). As the

observed interference of healthy control serum on 14.G2a-PE binding to M21 was

ameliorated at the 1:48 ratio, we used this same ratio to test sera from the clinical trial.

We included heathy donor sera diluted 1:6 and 1:48 with PBS as positive and negative,

respectively, controls in each assay. Each dog's pre-treatment sample was used as its

baseline.

Varying levels of binding inhibition were observed in 10 of 12 dogs (all but K913

and K914) when canine sera were mixed with 14G2a mAb before the mixture was used

to stain M21 cells (Table 2, Fig 3, and S2 Fig for select data). The binding was

significantly inhibited at Days 10, 30 and 60 compared to pre-treatment in all 10 of these

same dogs (save for subjects K906 and K912, that did not have day 60 serum sample

collected). GD2 binding was also significantly inhibited by Day 1 sera from 2 of these

same 10 dogs, one dog in each Arm, (K910 and K915), but for these 2 dogs, the

calculated binding inhibition on Day 1 was not as high as at any of the other days tested

for those same 2 dogs. In the remaining 10 dogs, that did not have significant binding

inhibition on Day 1, the binding of 14G2a to GD2 was increased at Day 1 for 8 of them

(all but K905 and K908), i.e., the % inhibition calculated using the formula in the

methods section was negative, corresponding to a slight increase in binding at Day 1

compared to the pre-treatment serum. However, the increase in binding seen on Day 1

(i.e., the negative binding inhibition value) was significant in only one of these eight

dogs (K909) (median -8.7%, range -0.3 to -21.2%) (Fig 3 and Table 2). These relatively

small positive or negative binding inhibition values detected on Day 1 (prior to any IT-IC

treatment, but after RT), which are significant only for 3 dogs (two positive and one

negative), were not anticipated and are discussed below; for all 3 of these dogs, all 3

subsequent time points for each dog show higher and statistically significant binding

inhibition. In contrast to the 10 dogs that did show significant binding inhibition, sera

from the other 2 dogs (K913 and K914), did not inhibit GD2 binding at any timepoint (Fig

3 and Table 2). Select flow cytometry data are shown in S2 Fig.

Table 2. Binding Inhibition (%) of anti-GD2 mAb to GD2 target in the presence of canine serum

Patient ID	RT Arm	Day 1	Day 10	Day 30	Day 60
ITIC-04	A	-7.6	88.3	98.2	57.2
ITIC-05	B	-1.5	92.8	96.6	98.8
ITIC-06	A	-3.5	74.2	97.3	n/a
ITIC-07	B	-5.6	12.5	91.3	40.6
ITIC-08	B	-0.3	26.4	96.4	89.9
ITIC-09	A	-21.2	93.3	96.1	97.0
ITIC-10	A	17.3	36.1	91.4	53.2
ITIC-11	B	-9.7	23.1	47.1	34.0
ITIC-12	A	-10.4	37.0	91.0	n/a
ITIC-13	A	-11.0	-23.2	-0.6	-6.9
ITIC-14	B	-10.7	13.1	10.2	14.6
ITIC-15	B	12.5	20.9	85.9	75.8

mAb, monoclonal antibody; RT, radiation therapy
Fluorescently labelled anti-GD2 antibody was incubated with canine serum and then used to stain GD2+ M21 tumor cells. % Binding Inhibition of post-treatment samples compared to baseline samples was determined by flow cytometry as described in Materials and Methods.

Fig 3. In vitro inhibition of binding of anti-GD2 antibody to GD2 target. The %

binding inhibition at various timepoints compared to Baseline. Data shown are

calculated from the mean of triplicates as described in Materials and Methods. Sera was

collected at Baseline (before radiation), Day 1 (after radiation, but just prior to IT-IC),

and Days 10, 30 and 60 (after IT-IC). Timepoints for each dog are presented and

connected by a line. Day 60 timepoint was not available for two dogs, K906 and K912,

in Arm A. Significance indicated is based on mean of triplicates at Baseline compared to

post-treatment timepoints, * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Comparison of CAHA Detection by ELISA and binding

inhibition by *in vitro* flow cytometry assay

Significant CAHA at various timepoints was detected by ELISA in 7 of 12 dogs: i)

K906 and K909 from Arm A at Day 30; ii) K908 from Arm B at Day 60; iii) K914 from Arm

B at Day 30; and iv) K905, K911, and K9 15 from Arm B at Day 30 and Day 60 (Fig 2).

Except for the Day 30 sera from K914, the above sera also significantly inhibited 14G2a

from binding to GD2 in the flow cytometry assay (Fig 3). To evaluate the relationship

between the ELISA and binding inhibition, the values for each serum sample for each

subject are plotted in Fig 4, with the ELISA O.D. value plotted on the Y axis, and the %

binding inhibition value plotted on the X-axis. There were no significant correlations

between the two assays within or for both arms combined detected. Significant binding

inhibition was detected earlier than significant CAHA: i) at Day 1 in 2 dogs; and ii) at

Day 10 in 10 dogs. Overall, inhibition was detected in 10 of 12 dogs: i) 5 of 6 Arm A

dogs and 5 of 6 Arm B dogs at Day 10; ii) 5 of 6 Arm A dogs and 5 of 6 Arm B dogs at

Day 30; iii) 3 of 4 Arm A dogs and 5 of 6 Arm B dogs at Day 60. Sera from the two dogs

that did not inhibit binding in the flow cytometry assay (K913 and K914) were also

CAHA negative by ELISA except for the Day 30 sera from K914 (Figs 2 and 3). Healthy

canine sera at the same dilution as sera from the clinical trial dogs were negative for

CAHA and did not inhibit the binding of 14G2a to M21 (Fig B in S1 Fig).

Fig 4. The relationship of CAHA and binding inhibition by *in vitro* flow cytometry

assay. The % binding inhibition related to CAHA at various timepoints is shown.

Timepoints for each dog are presented. Timepoints with significant CAHA responses are

shown with significance indicated, these timepoints also had significant binding

inhibition except for K914 Day 30. Sera from 10 of 12 dogs significantly inhibited binding

of anti-GD2 antibody to GD2 target at Day 10, 30 and 60 with the exceptions of K913

(shown as open symbols) and K914 (shown as black symbols). Day 1 sera from two

dogs (K910 and K915) inhibited binding. A single serum sample, K914 Day 30, showed

a significant CAHA response but did not inhibit binding (shown as the black circle).

Discussion

We developed two distinct assays to monitor and assess the CAHA response in

dogs receiving treatment of a readily palpable oral malignant melanoma with local RT

plus IT injection of a humanized IC. An ELISA was designed to detect CAHAs

specifically targeting human IgG – which is analogous to the Fc portion of hu14.18-IL2

[46]. We also utilized a flow cytometry assay to investigate the inhibition of anti-GD2

mAb binding to GD2 by sera from dogs receiving RT and IT-IC. This approach

employed ELISA as a tool to identify the presence of ADA in serum, followed by a

subsequent flow cytometry analysis to provide additional characterization of the ADA

response.

Although there are commercially available assays to detect various anti-species

antibodies, e.g., mouse anti-human (MAHA) and human anti-mouse (HAMA), an off-the-

shelf assay is not available to detect CAHA. The absence of a commercially available

ELISA specifically designed to detect CAHAs is an unmet need within the comparative

immuno-oncology field. This gap becomes increasingly relevant as the canine model

gains recognition for its value in studying human disease. The primary objective of

developing our in-house CAHA ELISA was to create a monitoring tool that would be

easy to use in veterinary and translational research laboratories. Additionally, we sought

to develop a cost-effective and flexible assay. The binding specificity of the CAHA could

also be explored in more depth by modifying the whole human IgG coating reagent that

was used in this report (e.g., whole IgG, Fc fragment, Fab or F(ab')$_2$).

A challenge in developing an ELISA to detect canine antibody that binds human

IgG was the lack of commercially available anti-canine IgG HRP conjugates that were

cross absorbed against human IgG. This cross-absorption step is crucial for reducing

the assay background and ensuring specificity. To address this limitation, we used a

rabbit anti-mouse IgG HRP conjugate that had been cross absorbed against human

IgG. Our testing during assay optimization demonstrated sufficient cross-reactivity

between mouse and canine-derived IgG, enabling us to detect CAHA to human IgG

(data not shown). In addition, we included a commercially available mouse IgG specific

for human IgG as a control analyte sample for detection by the rabbit anti-mouse HRP

in each assay. The selection of antibodies and the incorporation of cross-absorption

steps contributed to the assay's specificity and adaptability for potential use in both

veterinary and translational research lab settings.

Our CAHA ELISA is primarily designed to detect free ADAs in serum, by

detecting their binding to plate-bound human IgG. In contrast, the in vitro binding

inhibition flow cytometry assay allows the ADAs to interact with the soluble mouse

14.G2a mAb (which shares the idiotype of the hu14.18-IL2 IC) in solution, and

evaluates detection of GD2 binding in a cellular context, enabling the detection of ADAs

that inhibit or interfere with the binding of a therapeutic drug to its specific target in its

natural form on the cell surface, e.g., GD2 on tumor cells. This distinction is critical

because antigens immobilized to plastic, as in an ELISA, may not accurately mimic the

natural conformations of antigens on cell surfaces [47], potentially leading to an

underestimation of certain ADAs by this ELISA that flow cytometry can detect.

It is of interest that two dogs (K910 and K915) inhibited the binding of the anti-

GD2 mAb to GD2 on target cells at Day 1. The Day 1 sample was collected the day of,

but prior to, the first injection of hu14.18-IL2. Therefore, these responses following RT

may reflect pre-existing canine anti-human or anti-mouse antibodies, cross species-

reactive antibodies, or assay interference [48-50]. The first step in the flow cytometry

binding inhibition assay is the incubation of canine sera with the mouse anti-GD2 mAb

14G2a prior to incubating the mixture with the GD2+ M21 cells. Therefore, a possible

explanation for the Day 1 binding inhibition is pre-existing canine anti-mouse antibodies

as previously reported [49], but these would have needed to appear, or be boosted,

following baseline, as the positive binding inhibition is detected in comparison to the

baseline value for that same dog. We did not prospectively survey clients regarding their

pet's known allergies or exposures to other mammalian species, e.g., mouse or rabbit,

which could potentially induce a humoral response. mAbs of both mouse and rabbit

origin are used in our in-house assays and could result in assay background or

interference. However, neither of these two dogs were CAHA+ by ELISA at Day 1. Anti-

human IgG was not detected by ELISA until Day 30 in any dog, nor from healthy canine

sera, suggesting exposure to hu14.18-IL2 as the origin of the antibodies. However, we

have not formally tested for pre-existing anti-mouse or anti-rabbit antibodies in these

dogs. These seemingly discordant data could reflect the different assay sensitivities and

different assay modalities. Moreover, serum from healthy humans and healthy dogs can

both contain detectable levels of autoantibodies including rheumatoid factors and anti-

idiotypic antibodies which may cause assay interference [48, 50, 51].

Additionally, activation of B cells, generation of anti-tumor antibodies and

establishment of B cell memory following a treatment regimen similar to that used in this

trial have been reported in a mouse melanoma model [52]. As IgM typically arises early

in an immune response, the CAHA response detected at early timepoints (e.g., Day 1,

Day 10) in this study may be of IgM isotype which our CAHA ELISA may also detect. in

contrast, the flow cytometry assay does not detect a particular antibody isotype and

thus may detect ADA despite lack of CAHA detected by ELISA in the same serum. The

detection of binding inhibition at early time points may also result from the isotype

agnostic nature of the flow cytometry assay. The increase in binding inhibition observed

in two dogs (K910 and K915) from baseline to Day 1 was unexpected as the dogs had

not been exposed to hu14.18-IL2 at the time of serum collection on either day.

Moreover, the timeframe from baseline to Day 1 in this study was between 5-9 days,

representing rapid kinetics for a *de novo* humoral response. However, we have not

determined the fine specificity of the antibody responses in this study. All dogs treated

in this study received radiation, thus there is potential for radiation-induced antigen

shedding by tumor to trigger a humoral response [53]. Tumor-specific and self-specific

antibodies have been found in sera of cancer patients in several studies [54, 55]. Thus,

the binding inhibition at Day 1 in these two dogs could be due to endogenous cross-

reactive antibodies or those recognizing cryptic epitopes or neoantigens [56].

The IC has humanized IL2 linked to the mAb Fc region, thus the IL2 and/or the

linker between the Fc and the IL2 may be recognized as foreign in the dog triggering an

immune response. None of the dogs in this study experienced significant or unexpected

adverse events [38]. However, as dogs received only a single 3-day course of IT-IC, we

could not evaluate the impact of induced ADAs on subsequent pharmacokinetics of

hu14.18-IL2 or on the potential toxicity during subsequent courses. It will be important to

evaluate peak serum levels of hu14.18-IL2 as well as monitor anti-IL2 and anti-linker

responses to determine if CAHA+ sera affect antibody-mediated cellular cytotoxicity, as

seen in some humans receiving IC treatment [26].

Additional studies could further characterize the binding specificity in terms of

distinct IgG regions, e.g., Fc receptor, Fab hinge region [49]. Further, as in humans, IgG

is the most prevalent Ig in canine sera, however IgM and IgA are present at lower

concentrations [57]. The rabbit anti-mouse IgG HRP secondary reagent used in the

ELISA is reactive to both the heavy and light chain of mouse IgG. As such, this reagent

may also react with other immunoglobulin classes, e.g., IgM, IgA, since these all share

the same light chain. Therefore, subsequent testing could determine the isotype of the

CAHA responses, as well as species cross-reactivity – in particular to mouse or rabbit

IgG.

The timeline of the anti-human IgG CAHA ELISA response observed in our study

(7 of 12 dogs at Day 30) is similar to that observed in a previous study of dogs treated

with NHS-IL12, a humanized antibody against necrotic tumor regions linked to

humanized IL12. Dogs treated with NHS-IL12 developed ADA against both human IgG

(8 of 12 dogs) and against human IL12 (5 of 14 dogs) between 15 to 29 days after

treatment [31]. Factors that may explain differences in the ADA timeline between our

study and that of Paoloni et al., include the composition and target of the IC, IC dose

level, administration route, as well as the assays used to monitor the development of

ADA.

Conclusion

The CAHA detection assays in this report provide tools to monitor parallel canine

patient populations receiving treatment with human biopharmaceuticals. These assays

should be translatable to other treatment regimens delivering "foreign" protein

molecules *in vivo*, as well as to species other than the dog, provided appropriate

controls are available. The recent availability of immune checkpoint blockade in the form

of a caninized anti-PD1 therapeutic mAb (Gilvetmab, Merck) expands the treatment

options for dogs with melanoma. To this end, we are evaluating the addition of immune

checkpoint blockade, i.e., anti-PD1, to our novel treatment regimen of RT and IT-IC with

hu14.18-IL2 in dogs with melanoma. Local anti-tumor responses as well as systemic

immune responses, including the development of CAHA using the assays described

herein, will be investigated.

Acknowledgements

546 We are grateful for the dogs with melanoma, their caregivers, and referring

547 veterinarians who participated in the clinical trial. We are also thankful for the healthy

548 dogs and their caregivers for donating peripheral blood samples. We thank Rubi Hayim,

549 Emily M. Tomesek, and the UW Veterinary Care Oncology team for caring for the dogs

550 in this study.

References

1.   Gardner HL, Fenger JM, London CA. Dogs as a Model for Cancer. Annu Rev

Anim Biosci. 2016;4:199-222. Epub 20151109. doi: 10.1146/annurev-animal-022114-

110911. PubMed PMID: 26566160; PubMed Central PMCID: PMCPMC6314649.

2.   Paoloni M, Khanna C. Translation of new cancer treatments from pet dogs to

humans. Nat Rev Cancer. 2008;8(2):147-56. doi: 10.1038/nrc2273. PubMed PMID:

18202698.

3.   Vail DM, MacEwen EG. Spontaneously occurring tumors of companion animals

as models for human cancer. Cancer Invest. 2000;18(8):781-92. doi:

10.3109/07357900009012210. PubMed PMID: 11107448.

4.   Onaciu A, Munteanu R, Munteanu VC, Gulei D, Raduly L, Feder RI, et al.

Spontaneous and Induced Animal Models for Cancer Research. Diagnostics (Basel).

2020;10(9). Epub 20200831. doi: 10.3390/diagnostics10090660. PubMed PMID:

32878340; PubMed Central PMCID: PMCPMC7555044.

5.   Oh JH, Cho JY. Comparative oncology: overcoming human cancer through

companion animal studies. Exp Mol Med. 2023;55(4):725-34. Epub 20230403. doi:

10.1038/s12276-023-00977-3. PubMed PMID: 37009802; PubMed Central PMCID:

PMCPMC10167357.

6.   Crossley RA, Matz A, Dew T, Kalinauskas A, Faucette N, Poff B, et al. Safety

Evaluation of Autologous Tissue Vaccine Cancer Immunotherapy in a Canine Model.

Anticancer Res. 2019;39(4):1699-703. doi: 10.21873/anticanres.13275. PubMed PMID:

30952708.

7.   Dow S. A Role for Dogs in Advancing Cancer Immunotherapy Research. Front

Immunol. 2019;10:2935. Epub 20200117. doi: 10.3389/fimmu.2019.02935. PubMed

PMID: 32010120; PubMed Central PMCID: PMCPMC6979257.

8.   Maekawa N, Konnai S, Nishimura M, Kagawa Y, Takagi S, Hosoya K, et al. PD-

L1 immunohistochemistry for canine cancers and clinical benefit of anti-PD-L1 antibody

in dogs with pulmonary metastatic oral malignant melanoma. NPJ Precis Oncol.

2021;5(1):10. Epub 20210212. doi: 10.1038/s41698-021-00147-6. PubMed PMID:

33580183; PubMed Central PMCID: PMCPMC7881100.

9.   Park JS, Withers SS, Modiano JF, Kent MS, Chen M, Luna JI, et al. Canine

cancer immunotherapy studies: linking mouse and human. J Immunother Cancer.

2016;4:97. Epub 20161220. doi: 10.1186/s40425-016-0200-7. PubMed PMID:

28031824; PubMed Central PMCID: PMCPMC5171656.

10.   Zuleger CL, Kang C, Ranheim EA, Kurzman ID, Macklin MD, Newton MA, et al.

Pilot study of safety and feasibility of DNA microseeding for treatment of spontaneous

canine melanoma. Vet Med Sci. 2017;3(3):134-45. Epub 2017/10/27. doi:

10.1002/vms3.65. PubMed PMID: 29067210; PubMed Central PMCID:

PMCPMC5645840.

11.   Davda J, Declerck P, Hu-Lieskovan S, Hickling TP, Jacobs IA, Chou J, et al.

Immunogenicity of immunomodulatory, antibody-based, oncology therapeutics. J

Immunother Cancer. 2019;7(1):105. Epub 20190415. doi: 10.1186/s40425-019-0586-0.

PubMed PMID: 30992085; PubMed Central PMCID: PMCPMC6466770.

12.   Kim C, Yang H, Kim I, Kang B, Kim H, Kim H, et al. Association of High Levels of

Antidrug Antibodies Against Atezolizumab With Clinical Outcomes and T-Cell

Responses in Patients With Hepatocellular Carcinoma. JAMA Oncol. 2022;8(12):1825-

9. doi: 10.1001/jamaoncol.2022.4733. PubMed PMID: 36264560; PubMed Central

PMCID: PMCPMC9585463.

13. Sasson SC, Wilkins LE, Watson RA, Jolly C, Brain O, Klenerman P, et al.

Identification of neutralising pembrolizumab anti-drug antibodies in patients with

melanoma. Sci Rep. 2021;11(1):19253. Epub 20210928. doi: 10.1038/s41598-021-

98700-7. PubMed PMID: 34584157; PubMed Central PMCID: PMCPMC8478874.

14. Vaisman-Mentesh A, Gutierrez-Gonzalez M, DeKosky BJ, Wine Y. The Molecular

Mechanisms That Underlie the Immune Biology of Anti-drug Antibody Formation

Following Treatment With Monoclonal Antibodies. Front Immunol. 2020;11:1951. Epub

20200818. doi: 10.3389/fimmu.2020.01951. PubMed PMID: 33013848; PubMed Central

PMCID: PMCPMC7461797.

15. van Brummelen EM, Ros W, Wolbink G, Beijnen JH, Schellens JH. Antidrug

Antibody Formation in Oncology: Clinical Relevance and Challenges. Oncologist.

2016;21(10):1260-8. Epub 20160720. doi: 10.1634/theoncologist.2016-0061. PubMed

PMID: 27440064; PubMed Central PMCID: PMCPMC5061540.

16. Klingemann H. Immunotherapy for Dogs: Still Running Behind Humans. Front

Immunol. 2021;12:665784. Epub 20210805. doi: 10.3389/fimmu.2021.665784. PubMed

PMID: 34421888; PubMed Central PMCID: PMCPMC8374065.

17. Hank JA, Surfus JE, Gan J, Jaeger P, Gillies SD, Reisfeld RA, Sondel PM.

Activation of human effector cells by a tumor reactive recombinant anti-ganglioside GD2

interleukin-2 fusion protein (ch14.18-IL2). Clin Cancer Res. 1996;2(12):1951-9. Epub

1996/12/01. PubMed PMID: 9816154.

18. Imboden M, Murphy KR, Rakhmilevich AL, Neal ZC, Xiang R, Reisfeld RA, et al.

The level of MHC class I expression on murine adenocarcinoma can change the

antitumor effector mechanism of immunocytokine therapy. Cancer Res.

2001;61(4):1500-7. PubMed PMID: WOS:000167255400047.

19. Morris ZS, Guy EI, Francis DM, Gressett MM, Werner LR, Carmichael LL, et al.

In Situ Tumor Vaccination by Combining Local Radiation and Tumor-Specific Antibody

or Immunocytokine Treatments. Cancer Res. 2016;76(13):3929-41. Epub 2016/05/20.

doi: 10.1158/0008-5472.CAN-15-2644. PubMed PMID: 27197149; PubMed Central

PMCID: PMCPMC4930687.

20. Neal ZC, Imboden M, Rakhmilevich AL, Kim KM, Hank JA, Surfus J, et al. NXS2

murine neuroblastomas express increased levels of MHC class I antigens upon

recurrence following NK-dependent immunotherapy. Cancer Immunol Immunother.

2004;53(1):41-52. Epub 20030918. doi: 10.1007/s00262-003-0435-2. PubMed PMID:

14504825; PubMed Central PMCID: PMCPMC11032988.

21. Albertini MR, Hank JA, Gadbaw B, Kostlevy J, Haldeman J, Schalch H, et al.

Phase II trial of hu14.18-IL2 for patients with metastatic melanoma. Cancer Immunol

Immunother. 2012;61(12):2261-71. Epub 2012/06/09. doi: 10.1007/s00262-012-1286-5.

PubMed PMID: 22678096; PubMed Central PMCID: PMCPMC3502633.

22. Albertini MR, Yang RK, Ranheim EA, Hank JA, Zuleger CL, Weber S, et al. Pilot

trial of the hu14.18-IL2 immunocytokine in patients with completely resectable recurrent

stage III or stage IV melanoma. Cancer Immunol Immunother. 2018;67(10):1647-58.

Epub 2018/08/04. doi: 10.1007/s00262-018-2223-z. PubMed PMID: 30073390; PubMed

Central PMCID: PMCPMC6168354.

23.   King DM, Albertini MR, Schalch H, Hank JA, Gan J, Surfus J, et al. Phase I

clinical trial of the immunocytokine EMD 273063 in melanoma patients. J Clin Oncol.

2004;22(22):4463-73. Epub 2004/10/16. doi: 10.1200/JCO.2004.11.035. PubMed PMID:

15483010; PubMed Central PMCID: PMCPMC2367368.

24.   Osenga KL, Hank JA, Albertini MR, Gan J, Sternberg AG, Eickhoff J, et al. A

phase I clinical trial of the hu14.18-IL2 (EMD 273063) as a treatment for children with

refractory or recurrent neuroblastoma and melanoma: a study of the Children's

Oncology Group. Clin Cancer Res. 2006;12(6):1750-9. Epub 2006/03/23. doi:

10.1158/1078-0432.CCR-05-2000. PubMed PMID: 16551859; PubMed Central PMCID:

PMCPMC2587020.

25.   Shusterman S, London WB, Gillies SD, Hank JA, Voss SD, Seeger RC, et al.

Antitumor activity of hu14.18-IL2 in patients with relapsed/refractory neuroblastoma: a

Children's Oncology Group (COG) phase II study. J Clin Oncol. 2010;28(33):4969-75.

Epub 2010/10/06. doi: 10.1200/JCO.2009.27.8861. PubMed PMID: 20921469; PubMed

Central PMCID: PMCPMC3020698.

26.   Hank JA, Gan J, Ryu H, Ostendorf A, Stauder MC, Sternberg A, et al.

Immunogenicity of the hu14.18-IL2 immunocytokine molecule in adults with melanoma

and children with neuroblastoma. Clin Cancer Res. 2009;15(18):5923-30. Epub

2009/09/10. doi: 10.1158/1078-0432.Ccr-08-2963. PubMed PMID: 19737959; PubMed

Central PMCID: PMCPMC2745522.

27.   Navid F, Sondel PM, Barfield R, Shulkin BL, Kaufman RA, Allay JA, et al. Phase I

trial of a novel anti-GD2 monoclonal antibody, Hu14.18K322A, designed to decrease

toxicity in children with refractory or recurrent neuroblastoma. J Clin Oncol.

2014;32(14):1445-52. Epub 20140407. doi: 10.1200/jco.2013.50.4423. PubMed PMID:

24711551; PubMed Central PMCID: PMCPMC4017710.

28.   Yu AL, Gilman AL, Ozkaynak MF, Naranjo A, Diccianni MB, Gan J, et al. Long-

Term Follow-up of a Phase III Study of ch14.18 (Dinutuximab) + Cytokine

Immunotherapy in Children with High-Risk Neuroblastoma: COG Study ANBL0032. Clin

Cancer Res. 2021;27(8):2179-89. Epub 20210127. doi: 10.1158/1078-0432.Ccr-20-

3909. PubMed PMID: 33504555; PubMed Central PMCID: PMCPMC8046731.

29.   Schellekens H. Factors influencing the immunogenicity of therapeutic proteins.

Nephrol Dial Transplant. 2005;20 Suppl 6:vi3-9. doi: 10.1093/ndt/gfh1092. PubMed

PMID: 15958824.

30.   Baniel CC, Sumiec EG, Hank JA, Bates AM, Erbe AK, Pieper AA, et al.

Intratumoral injection reduces toxicity and antibody-mediated neutralization of

immunocytokine in a mouse melanoma model. J Immunother Cancer. 2020;8(2). doi:

10.1136/jitc-2020-001262. PubMed PMID: 33115944; PubMed Central PMCID:

PMCPMC7594540.

31.   Paoloni M, Mazcko C, Selting K, Lana S, Barber L, Phillips J, et al. Defining the

Pharmacodynamic Profile and Therapeutic Index of NHS-IL12 Immunocytokine in Dogs

with Malignant Melanoma. PLoS One. 2015;10(6):e0129954. Epub 20150619. doi:

10.1371/journal.pone.0129954. PubMed PMID: 26091536; PubMed Central PMCID:

PMCPMC4474860.

32.   Parng C, Bolt M, Pittman DD, Caiazzo T, Dyleski L, Gorovits B, Webster R.

Induction and Impact of Anti-Drug Responses Elicited by a Human Recombinant

Coagulation Factor FXa(I16L) in Preclinical Species. Aaps j. 2019;21(3):52. Epub

20190411. doi: 10.1208/s12248-019-0324-z. PubMed PMID: 30976993.

33.   Wong C, Muguiro DH, Lavergne S, Behling-Kelly E, Goggs R. Pharmacokinetics

of human recombinant C1-esterase inhibitor and development of anti-drug antibodies in

healthy dogs. Vet Immunol Immunopathol. 2018;203:66-72. Epub 20180823. doi:

10.1016/j.vetimm.2018.08.006. PubMed PMID: 30243377.

34.   Jani M, Isaacs JD, Morgan AW, Wilson AG, Plant D, Hyrich KL, et al. Detection of

anti-drug antibodies using a bridging ELISA compared with radioimmunoassay in

adalimumab-treated rheumatoid arthritis patients with random drug levels.

Rheumatology (Oxford). 2016;55(11):2050-5. Epub 20160825. doi:

10.1093/rheumatology/kew299. PubMed PMID: 27565176; PubMed Central PMCID:

PMCPMC5088626.

35.   Stubenrauch K, Wessels U, Birnboeck H, Ramirez F, Jahreis A, Schleypen J.

Subset analysis of patients experiencing clinical events of a potentially immunogenic

nature in the pivotal clinical trials of tocilizumab for rheumatoid arthritis: Evaluation of an

antidrug antibody ELISA using clinical adverse event-driven immunogenicity testing.

Clin Ther. 2010;32(9):1597-609. doi: 10.1016/j.clinthera.2010.07.021. PubMed PMID:

20974317.

36.   Suh K, Kyei I, Hage DS. Approaches for the detection and analysis of antidrug

antibodies to biopharmaceuticals: A review. J Sep Sci. 2022;45(12):2077-92. Epub

20220310. doi: 10.1002/jssc.202200112. PubMed PMID: 35230731; PubMed Central

PMCID: PMCPMC9520450.

37.   Vande Casteele N, Buurman DJ, Sturkenboom MG, Kleibeuker JH, Vermeire S,

Rispens T, et al. Detection of infliximab levels and anti-infliximab antibodies: a

comparison of three different assays. Aliment Pharmacol Ther. 2012;36(8):765-71. Epub

20120828. doi: 10.1111/apt.12030. PubMed PMID: 22928581.

38.   Albertini MR, Zuleger CL, Ranheim EA, Shiyanbola O, Sondel PM, Morris ZS, et

al. Administration of intratumoral GD2-directed interleukin-2 immunocytokine and local

radiation therapy to activate immune rejection of spontaneous canine melanoma.

Melanoma Res. 2024;34(4):307-18. Epub 20240520. doi:

10.1097/CMR.0000000000000975. PubMed PMID: 38768442.

39.   Magee K, Marsh IR, Turek MM, Grudzinski J, Aluicio-Sarduy E, Engle JW, et al.

Safety and feasibility of an in situ vaccination and immunomodulatory targeted

radionuclide combination immuno-radiotherapy approach in a comparative (companion

dog) setting. PLoS One. 2021;16(8):e0255798. Epub 20210812. doi:

10.1371/journal.pone.0255798. PubMed PMID: 34383787; PubMed Central PMCID:

PMCPMC8360580.

40.   LeBlanc AK, Atherton M, Bentley RT, Boudreau CE, Burton JH, Curran KM, et al.

Veterinary Cooperative Oncology Group-Common Terminology Criteria for Adverse

Events (VCOG-CTCAE v2) following investigational therapy in dogs and cats. Vet Comp

Oncol. 2021;19(2):311-52. Epub 20210218. doi: 10.1111/vco.12677. PubMed PMID:

33427378; PubMed Central PMCID: PMCPMC8248125.

41.   Leary S, Members of the Panel on Euthanasia:. AVMA guidelines for the

euthanasia of animals: 2020 edition. Schaumburg, IL: American Veterinary Medical

Association; 2020.

42.   Hornbeck PV. Enzyme-Linked Immunosorbent Assays. Curr Protoc Immunol.

2015;110:2.1.-2.1.23. Epub 20150803. doi: 10.1002/0471142735.im0201s110. PubMed

PMID: 26237010.

43.   R Core Team. R: A language and environment for statistical computing. The R

Foundation for Statistical Computing, Vienna, Austria. 2022.

44.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New

York; 2016.

45.   Wickham H. stringr: Simple, Consistent Wrappers for Common String

Operations. 2022.

46.   Navid F, Santana VM, Barfield RC. Anti-GD2 antibody therapy for GD2-

expressing tumors. Curr Cancer Drug Targets. 2010;10(2):200-9. doi:

10.2174/156800910791054167. PubMed PMID: 20201786; PubMed Central PMCID:

PMCPMC2888262.

47.   Butler JE, Ni L, Brown WR, Joshi KS, Chang J, Rosenberg B, Voss EW, Jr. The

immunochemistry of sandwich ELISAs--VI. Greater than 90% of monoclonal and 75% of

polyclonal anti-fluorescyl capture antibodies (CAbs) are denatured by passive

adsorption. Mol Immunol. 1993;30(13):1165-75. doi: 10.1016/0161-5890(93)90135-x.

PubMed PMID: 8413321.

48.   Bergman D, Bäckström C, Hansson-Hamlin H, Larsson A, Holst BS. Pre-existing

canine anti-IgG antibodies: implications for immunotherapy, immunogenicity testing and

immunoassay analysis. Sci Rep. 2020;10(1):12696. Epub 20200729. doi:

10.1038/s41598-020-69618-3. PubMed PMID: 32728049; PubMed Central PMCID:

PMCPMC7391631.

49. Bergman D, Larsson A, Hansson-Hamlin H, Åhlén E, Holst BS. Characterization

of canine anti-mouse antibodies highlights that multiple strategies are needed to combat

immunoassay interference. Sci Rep. 2019;9(1):14521. Epub 20191010. doi:

10.1038/s41598-019-51228-3. PubMed PMID: 31601945; PubMed Central PMCID:

PMCPMC6787031.

50. Goldberg JL, Navid F, Hank JA, Erbe AK, Santana V, Gan J, et al. Pre-existing

antitherapeutic antibodies against the Fc region of the hu14.18K322A mAb are

associated with outcome in patients with relapsed neuroblastoma. J Immunother

Cancer. 2020;8(1). doi: 10.1136/jitc-2020-000590. PubMed PMID: 32169872; PubMed

Central PMCID: PMCPMC7069273.

51. Harding FA, Stickler MM, Razo J, DuBridge RB. The immunogenicity of

humanized and fully human antibodies: residual immunogenicity resides in the CDR

regions. MAbs. 2010;2(3):256-65. Epub 20100501. doi: 10.4161/mabs.2.3.11641.

PubMed PMID: 20400861; PubMed Central PMCID: PMCPMC2881252.

52. Baniel CC, Heinze CM, Hoefges A, Sumiec EG, Hank JA, Carlson PM, et al. In

situ Vaccine Plus Checkpoint Blockade Induces Memory Humoral Response. Front

Immunol. 2020;11:1610. Epub 20200724. doi: 10.3389/fimmu.2020.01610. PubMed

PMID: 32849544; PubMed Central PMCID: PMCPMC7396490.

53. Nabrinsky E, Macklis J, Bitran J. A Review of the Abscopal Effect in the Era of

Immunotherapy. Cureus. 2022;14(9):e29620. Epub 20220926. doi:

10.7759/cureus.29620. PubMed PMID: 36321062; PubMed Central PMCID:

PMCPMC9604762.

54.   Brichory FM, Misek DE, Yim AM, Krause MC, Giordano TJ, Beer DG, Hanash

SM. An immune response manifested by the common occurrence of annexins I and II

autoantibodies and high circulating levels of IL-6 in lung cancer. Proc Natl Acad Sci U S

A. 2001;98(17):9824-9. doi: 10.1073/pnas.171320598. PubMed PMID: 11504947;

PubMed Central PMCID: PMCPMC55537.

55.   Reuschenbach M, von Knebel Doeberitz M, Wentzensen N. A systematic review

of humoral immune responses against tumor antigens. Cancer Immunol Immunother.

2009;58(10):1535-44. Epub 20090628. doi: 10.1007/s00262-009-0733-4. PubMed

PMID: 19562338; PubMed Central PMCID: PMCPMC2782676.

56.   Fischer E, Kobold S, Kleber S, Kubuschok B, Braziulis E, Knuth A, et al. Cryptic

epitopes induce high-titer humoral immune response in patients with cancer. J Immunol.

2010;185(5):3095-102. Epub 20100726. doi: 10.4049/jimmunol.0902166. PubMed

PMID: 20660712.

57.   German AJ, Hall EJ, Day MJ. Measurement of IgG, IgM and IgA concentrations

in canine serum, saliva, tears and bile. Vet Immunol Immunopathol. 1998;64(2):107-21.

doi: 10.1016/s0165-2427(98)00132-9. PubMed PMID: 9661261.

Supporting information

S1 Fig. Binding inhibition assay controls. PE-conjugated anti-GD2 antibody 14G2a

was mixed with either (A) 20 μl (open histogram with solid line) or (B) 2.5 μl (open

histogram with solid line) healthy canine donor sera before staining M21 cells. M21

stained with 14G2a-PE mixed with 20 μl or 2.5 μl PBS as a no-serum control is

represented as a gray histogram in both (A) and (B). Data are representative of

triplicates. Values represent % binding inhibition compared to no-serum control.

S2 Fig. Serum from dogs treated with hu14.18-IL2 shows varied level of inhibition

of 14G2a binding to target GD2. Prior to staining M21 cells, PE-conjugated anti-GD2

antibody 14G2a was mixed with baseline/pre-treatment sera or post-treatment sera

from different timepoints from (A) K914 or (B) K915. Baseline/pre-treatment data are

represented as gray histograms; post-treatment data are represented as open

histograms with solid lines. Data are representative of triplicates. Values represent %

binding inhibition compared to baseline/pre-treatment control.

Data availability statement

All relevant data are within the manuscript and its Supporting Information files.

Funding

Supported by Merit Review Award I01 BX003916 from the Biomedical Laboratory Research and Development Service of the United States (U.S.) Department of Veterans Affairs (MRA), the University of Wisconsin Carbone Cancer Center (UWCCC) Cancer Pharmacology Laboratory and Flow Cytometry Laboratory (grant 1S10OD018202-01) supported by P30 CA014520 (NIH/HHS), the Barbara A. Suran Comparative Research Endowment (DMV), melanoma research gifts to the UWCCC (Tim Eagle Memorial (MRA), gift from Jill and Bob Freiermuth (MRA)), and use of facilities at the William S. Middleton Memorial Veterans Hospital, Madison, WI. The contents do not represent the views of the U.S. Department of Veterans Affairs or the United States Government. Anyxis Immuno-Oncology Gmbh is the current license holder for hu14.18-IL2 and provided hu14.18-IL2 for this program of research. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests

None