Critical Environmental Limits: Assessing the Limitations of Core Temperature Inflection Point (CTIP) and Biophysical Modeling

Faming Wang[1,2,3], Huijuan Xu[3], Tze-Huan Lei[2], Yi Xu[2], Haojian Wang[2], Lijuan Wang[4]

[1]One Health Research Group, School of Environmental Science and Engineering, Tianjin University, 300072, Tianjin, China

[2]Centre for Molecular Biosciences and Non-Communicable Diseases, School of Safety Science and Engineering, Xi'an University of Science and Technology, Xi'an 710054, China

[3]Hebei Key Laboratory of Man-machine Environmental Thermal Control Technology and Equipment, Hebei Vocational University of Technology and Engineering, Xingtai, 054000, China

[4]College of Urban Planning and Municipal Engineering, Xi'an Polytechnic University, Xi'an, Shaanxi, 710048, China

Running head: Limitations of CTIP and Biophysical Modeling

*Correspondence:

Prof. Faming Wang, PhD

School of Environmental Science and Engineering

Tianjin University

Tianjin, China

E-mail: faming.wang@xust.edu.cn

Word count: 4,439 words (references excluded)

Critical Environmental Limits: Assessing the Limitations of Core Temperature Inflection Point (CTIP) and Biophysical Modeling

ABSTRACT (230 WORDS)

The Core Temperature Inflection Point (CTIP) method and biophysical modeling are widely used to determine critical environmental limits (CEL), yet their validity under prolonged heat exposure remains untested. This study evaluated their predictive accuracy by exposing 36 healthy young adults (20 males, 16 females; age: 20.9–22.4 yr) to five counterbalanced 8-hour heat trials in a controlled chamber (36°C/74.5% RH, 40°C/55.0% RH, 44°C/29.2% RH, 47°C/35.6% RH, 50°C/24.5% RH). These conditions were selected based on prior CTIP and biophysical model predictions of CELs. Participants engaged in sedentary office tasks (1.29–1.67 METs), wore standardized summer clothing (0.39–0.40 clo), and had ad libitum access to an electrolyte drink, with a 500-kcal sandwich provided at midday. Rectal temperature (T_{rec}) was continuously monitored. Contrary to CTIP and biophysical model predictions, all five conditions remained compensable (T_{rec} rise rate ≤ 0.1°C/h), with mean peak T_{rec} staying well below heatstroke thresholds (38.2 $\pm$ 0.4°C). At 44°C/29.2% RH, females had significantly lower T_{rec} than males ($p < 0.05$), but steady-state T_{rec} responses were similar between sexes (all $p > 0.10$). Collectively, CTIP and biophysical models substantially underestimated CELs, leading to overpredicted heat risk across all trials. These findings challenge the reliability of current predictive methods, suggesting human tolerance may exceed existing estimates. Refining these models is essential for improving heat risk assessment during real-world heatwaves and informing public and occupational health guidelines in a warming climate.

Keywords: compensability; survivability; heat tolerance; core temperature inflection; biophysical model; prolonged heat exposure

NEW & NOTEWORTHY

This study demonstrates that healthy young adults maintained heat balance over 8 hours across five extreme heat conditions (36–50°C), despite the Core Temperature Inflection Point (CTIP) method and biophysical modeling predicting uncompensability. These findings challenge the reliability of these tools for assessing prolonged heat exposure, highlighting the need for refined models to accurately predict critical environmental limits in real-world scenarios, such as extreme heatwaves.

INTRODUCTION

As global temperatures rise due to climate change, extreme heat events are becoming more frequent, intense, and prolonged, posing significant risks to human health and productivity (Mora et al., 2017). Prolonged exposure to extreme heat can lead to heat-related illnesses, ranging from heat exhaustion to life-threatening heat stroke, particularly in vulnerable populations such as the elderly, children, individuals with chronic illnesses, and outdoor workers (Luber and McGeehin, 2008). Understanding human heat tolerance limits is critical for developing effective mitigation strategies (Hanna and Brown, 1983; Sherwood and Ramsay, 2023). To assess these limits, researchers have developed three fast approaches to estimate the environmental conditions beyond which humans can no longer maintain thermal balance, often referred to as the uncompensability limit. These methods include the Core Temperature Inflection Point (CTIP) method, biophysical modeling, and thermoregulatory modeling (Belding and Kamon, 1973; Kamon and Avellini, 1976; Vecellio et al., 2022; Wolf et al., 2023; Vanos et al., 2023; Xu et al., 2024). While widely used, these methods require further validation through prolonged heat exposure studies in real-world or controlled laboratory settings to ensure their accuracy and applicability (Wang, 2025).

The CTIP method, introduced by Belding and Kamon (1973), identifies the critical environmental limits where heat stress transitions from compensable to uncompensable, marked by a distinct inflection in core temperature as heat gain exceeds heat loss (Avellini et al., 1980). Biophysical modeling, advanced by Vanos et al. (2023), integrates physiological and environmental variables, such as metabolic heat production and evaporative cooling, to estimate heat stress under diverse conditions (Wang et al., 2023). Thermoregulatory modeling (Kang et al., 2019; Xu et al., 2024), exemplified by the six-cylinder thermoregulatory model (SCTM), simulates dynamic physiological responses (e.g., sweating, vasodilation) to estimate critical environmental limits (CELs) for human survivability and livability. Each of these methods aims to refine the theoretical wet-bulb temperature (T_w) threshold of 35°C, originally proposed by Sherwood and Huber (2010) as the upper limit for human survival, by accounting for individual variability and real-world factors. However, their validation has primarily been limited to short-term controlled exposures or computer simulations, leaving their accuracy for prolonged heat stress untested.

Prolonged heat exposure, such as an 8-hour workday in hot environments, introduces unique physiological challenges, including cumulative heat storage, hydration deficits, and cardiovascular strain (Kenny et al., 2018; Meade et al., 2024). These factors elevate the risk of

heat-related illnesses, yet the CTIP method and biophysical modeling remain unvalidated under such conditions (Wang, 2025). This gap restricts their applicability in occupational safety, public health, and climate adaptation planning. Validating these methods against prolonged heat exposure is essential to ensure they reliably predict human heat tolerance in real-world scenarios.

In this study, we validated the CTIP method and biophysical modeling against 8-hour heat exposure in a controlled climatic chamber. By comparing their predicted uncompensability limits with observed physiological responses, we assess their reliability and accuracy for prolonged heat stress. Our findings will clarify the strengths and limitations of these methods, improving their applicability for assessing human heat tolerance during prolonged heat exposure and enhancing their applicability for real-world predictions in a warming world.

METHODS

Ethical approval and human participants

This study was approved by the Institutional Review Board of Xi'an University of Science and Technology (approval number: XUST-IRB225003) and registered in the Chinese Clinical Trial Registry (ChiCTR) database (registration number: ChiCTR2300071853). All participants provided verbal and written informed consent prior to enrollment, in accordance with the Declaration of Helsinki. Thirty-six healthy young adults were recruited: 20 males (age: 22.4 ± 0.9 years, body surface area: 1.83 ± 0.11 m², body mass index [BMI]: 22.3 ± 2.1 kg/m²) and 16 females (age: 20.9 ± 2.0 years, body surface area: 1.60 ± 0.12 m², BMI: 21.1 ± 2.2 kg/m²). Female participants were tested during the early follicular phase of their menstrual cycle (5 ± 3 days), a period characterized by the lowest concentrations of estrogen and progesterone (Bouman et al., 2005). Participants engaged in lower to moderate physical activities, with males exercising 4.9 ± 2.0 times per week (activities included jogging, badminton, and strength training) and females exercising 3.0 ± 1.5 times per week (activities included jogging, tennis, basketball, and badminton). This ensured a study population with comparable activity levels while maintaining generalizability to young, physically active adults. Participants were also screened for cardiovascular, respiratory, and metabolic disorders, and were free from heat-related illnesses or medication use affecting thermoregulation.

Test conditions

Figure 1 Environment conditions for validating the CTIP method and biophysical modeling. The five validation conditions covered wet-bulb temperatures (T_w) ranging from 28.4°C to 33.0°C. The metabolic rates reported in the CTIP study by Vecellio et al. (2022) and the biophysical modeling simulations by Vanos et al. (2023) were 1.3-1.5 METs and 1.5 METs, respectively. Filled circles with a pink border and yellow interior represent the three conditions selected for validating the CTIP method, while filled circles with a cyan border and yellow interior denote the two conditions chosen for validating the biophysical modelling. Additionally, clothing insulation in the CTIP method by Vecellio et al. (2022) and the biophysical modeling simulations by Vanos et al. (2023) was approximately 0.36 clo.

To validate the Core Temperature Inflection Point (CTIP) method and biophysical modeling for prolonged heat exposure, environmental conditions were selected based on prior predictions of uncompensability limits. Vecellio et al. (2022) defined critical relative humidity (RH) thresholds using the CTIP method: 74.0% at 36°C, 64.5% at 38°C, 53.5% at 40°C, 29.2% at 44°C, 21.0% at 47.3°C, and 13.2% at 50.6°C dry-bulb temperature. Three conditions were chosen for validation: 36°C with 74.5% RH, 40°C with 55.0% RH, and 44°C with 29.2% RH, representing moderate to extreme heat stress scenarios. Separately, Vanos et al. (2023) used biophysical modeling to predict survivability limits, with critical RH values ranging from 88.9%

at 35°C to 16.1% at 52°C dry-bulb temperature. Two conditions were selected to test these predictions: 47°C with 35.6% RH and 50°C with 24.5% RH, where Vanos et al. (2023) estimated young adults (18–40 years) would reach a core temperature of 43.0°C, indicative of heatstroke, within 6 hours under indoor shaded conditions (Pal and Eltahir, 2016). These five conditions (see Figure 1) were implemented in a controlled environmental chamber to assess the reliability of CTIP and biophysical modeling predictions over an 8-hour exposure period. The air velocity inside the test chamber across the five conditions was 0.20 ± 0.05 m/s, representing a static environment. The CO_2 concentration was maintained at 947 ± 32 ppm to ensure optimal air quality during the trials (Satish et al., 2012).

Protocol overview

Participants completed five counterbalanced 8-hour heat exposure trials, each simulating one of the environmental conditions outlined above (36°C/74.5% RH, 40°C/55.0% RH, 44°C/29.2% RH, 47°C/35.6% RH, 50°C/24.5% RH). Trials were conducted in a controlled environmental chamber (Espec Corp., Osaka, Japan) between November 2024 and March 2025 to minimize the potential effects of seasonal heat acclimatization on core temperature fluctuation (Buono et al., 1998). Each trial ran from 9:00 a.m. to 5:00 p.m., separated by a 5-day washout period to ensure recovery.

Before each trial, participants provided urine samples to verify a urine specific gravity (USG) below 1.020, indicating adequate hydration (Oppliger et al., 2005). If the threshold was exceeded, participants consumed an isotonic electrolyte drink (Nuotelande: 10.8 mg zinc, 0.5 g carbohydrates, 283 mg sodium; Jinan, China) until compliance was achieved. Pre-trial measurements included nude body weight and venous blood sampling. During trials, participants were seated in office chairs and engaged in sedentary office activities (e.g., reading, computer work). They wore standardized clothing, short-sleeve T-shirts, sports bras (females), underwear, trousers, socks, and sports shoes, yielding clothing insulation values of 0.39 clo (males) and 0.40 clo (females) (ASHRAE, 2023).

Hydration and dietary standardization

Participants had ad libitum access to an electrolyte drink throughout the 8-hour heat exposure to maintain hydration. Participants were instructed to drink as needed to avoid any development of thirst sensation, rather than follow a fixed intake schedule. To ensure consistency, the electrolyte drink was stored in a temperature-controlled chamber at 37°C, matching body

temperature to minimize potential cooling effects. Participants were also permitted restroom breaks as needed to maintain comfort and normal fluid regulation.

Dietary intake was standardized across all physiological trials, with each participant receiving a 500-kcal (2092 kJ) sandwich at 12:00 p.m. This controlled energy intake helped minimize dietary influences on metabolic rate and physiological responses (Lei et al., 2025).

Measurements

Body temperature

Core temperature (T_{rec}) was monitored continuously using a rectal thermistor (YSI401, Yellow Spring Instrument, Yellow Springs, OH; accurate: ±0.1°C) inserted 10 cm beyond the anal sphincter. The rectal thermistor was chosen as an index of deep body temperature because it remains unaffected by fluid intake and allows participants to consume the standardize meal during the 8-hour heat exposures without compromising the accuracy of the measurements. Mean skin temperature (T_{sk}) was calculated from four iButton dataloggers (DS1992L, Maxim Integrated Product Inc., San Jose, CA) placed at the chest, upper arm, thigh, and calf, averaged using Ramanathan's formula ($T_{sk} = 0.3 \times (T_{chest} + T_{upper\ arm}) + 0.2 \times (T_{thigh} + T_{calf})$) (Ramanathan, 1964).

Metabolic rate

Metabolic rate was measured using a COSMED K5 wearable metabolic system (COSMED srl, Rome, Italy) at two time points during the 8-hour prolonged heat exposure: at the beginning (15 minutes after exposure onset to allow participants to stabilize while engaging in sedentary office tasks such as reading and computer work) and at the end of the exposure. Each measurement was conducted continuously for 8 minutes, with data from the final 5 minutes used to calculate the metabolic rate. This approach ensured a representative assessment of participants' energy expenditure under controlled environmental conditions while minimizing transient fluctuations in metabolic data. This system was calibrated daily with high-grade calibration gases (GASCO: 5% CO_2, 21% O_2, balance N_2; Airgas Therapeutics LLC, Oldsmar, FL) and a 3-liter syringe (Hans Rudolph 3L Calibration Syringe), following the manufacturer's instructions to ensure accurate measurements. Metabolic rates were expressed in metabolic equivalents (METs) by dividing the volume of oxygen consumption by 3.5 mL·kg^{-1}·min^{-1} (Jetté et al., 1990).

Hydration status

The hydration status of participants was carefully monitored throughout the entire 8-hour exposure period. Participants were regularly reminded to maintain a sensation of not feeling thirsty, ensuring that they remained adequately hydrated during the trial. In addition to these verbal reminders, urine samples were collected both pre- and post-exposure to assess the participants' hydration levels. Urine specific gravity (USG) was measured using a urine specific gravity refractometer (Atago PAL-10S, cat. No. 4410, Atago Co. Ltd., Tokyo, Japan), which has an accuracy of ±0.001 and a measurement range of 1.000 to 1.060.

Data processing and statistical analysis

Individual rectal temperature data were processed using a smooth spline function in R software (version 4). The core temperature rate of change was calculated at 5-minute intervals throughout the 8-hour exposure to identify the core temperature steady state. Steady state was determined using one of two criteria: either the core temperature rate of change was ≤ 0.10°C/h after 3 hours of exposure, or it remained ≤ 0.1°C for the last 30 minutes of the 8-hour heat exposure.

Statistical analyses were performed using IBM SPSS Statistics (version 20, IBM Corp., Armonk, NY), with figures generated in GraphPad Prism (version 10.3.1, GraphPad Software LLC, Boston, MA). Data normality was tested with the Kolmogorov-Smirnov test, and variance homogeneity with Levene's test. A one-way mixed model analysis of variance (ANOVA) assessed the effects of timepoints (9:00 a.m.–5:00 p.m.) on T_{rec} with sex as a between-subjects factor. Significant main or interaction effects were followed by Bonferroni-corrected post hoc pairwise comparisons. Data are reported as means ± standard deviation (SD) unless otherwise noted. Statistical significance was set at $p \leq 0.05$.

RESULTS

Hydration status

Mean urine specific gravity (USG) values remained within the euhydrated range (≤ 1.020, see Table S1 in the **Supplemental Material**) across all five experimental conditions, indicating that participants maintained adequate hydration throughout the 8-hour heat exposure. While minor fluctuations in USG were observed between pre- and post-exposure measurements, these variations remained within normal limits, suggesting that fluid balance was effectively regulated. No evidence of dehydration was detected in either males or females at any tested condition.

Environmental conditions inside the chamber

The actual environmental conditions recorded for validating the CTIP method were $36.1 \pm 0.1°C$ with $74.1 \pm 0.2\%$ RH, $39.9 \pm 0.3°C$ with $55.7 \pm 0.7\%$ RH, and $44.0 \pm 0.1°C$ with $29.2 \pm 0.2\%$ RH, corresponding to wet-bulb temperatures of 32.0°C, 32.0°C and 28.7°C, respectively. Similarly, the two conditions chosen for validating the biophysical modeling were $47.1 \pm 0.1°C$ with $35.8 \pm 0.5\%$ RH and $50.1 \pm 0.2°C$ with $24.0 \pm 0.3\%$ RH, corresponding to wet-bulb temperatures of 33.1°C and 31.3°C.

Metabolic rate

The metabolic rate increased significantly by the end of the 8-hour heat exposure across all environmental conditions ($p < 0.05$; Table 1). The rise in MET from pre-exposure to post-exposure ranged from 15.4% to 20.1% in males and 17.8% to 20.3% in females, depending on the environmental conditions. Notably, males had consistently higher MET values than females throughout the trials ($p < 0.01$), with sex differences being statistically significant at both time points in all conditions ($p < 0.05$). The largest increase in MET was observed in the 50°C, 24.5% RH condition for males (20.1% increase) and the 47°C, 35.6% RH condition for females (20.3% increase), highlighting a trend where both sexes experienced greater metabolic elevations in extreme heat conditions.

Table 1 Metabolic equivalent (MET) before and at the end of the exposure for both sexes (males: n=20; females: n=16). Data were analyzed using one-way repeated ANOVA. *, indicates a significant difference from Pre-exposure ($p < 0.05$); #, indicates a significant difference between sexes ($p < 0.05$).

Environmental conditions	Males		Females	
	Pre-exposure (METs)	Post-exposure (METs)	Pre-exposure (METs)	Post-exposure (METs)
36°C, 74.5% RH	$1.35 \pm 0.10^{\#}$	$1.58 \pm 0.11^{*\#}$	1.29 ± 0.06	$1.52 \pm 0.09^{*}$
40°C, 55.0% RH	1.36 ± 0.08	$1.57 \pm 0.07^{*\#}$	1.30 ± 0.10	$1.51 \pm 0.06^{*}$
44°C, 29.2% RH	1.40 ± 0.09	$1.63 \pm 0.07^{*\#}$	1.28 ± 0.10	$1.53 \pm 0.09^{*}$
47°C, 35.6% RH	$1.39 \pm 0.08^{\#}$	$1.65 \pm 0.11^{*\#}$	1.28 ± 0.10	$1.54 \pm 0.09^{*}$
50°C, 24.5% RH	1.39 ± 0.08	$1.67 \pm 0.12^{*\#}$	1.30 ± 0.10	$1.56 \pm 0.05^{*}$

Body temperatures

All five experimental conditions represented compensable heat stress for both sexes ([Figure 2]). Among male participants, the time to reach thermoregulatory steady state across the five conditions (36°C/74.5% RH, 40°C/55.0% RH, 44°C/29.2% RH, 47°C/35.6% RH, 50°C/24.5% RH) was 278 ± 61, 288 ± 100, 216 ± 38, 291 ± 89, and 260 ± 64 minutes, respectively. For females, the corresponding times were 262 ± 48, 256 ± 33, 221 ± 56, 249 ± 70, and 227 ± 50 minutes. Core temperatures during the 8-hour exposure were generally comparable between sexes, except in the 44°C/29.2% RH condition ([Figure 2C], $p < 0.01$). No significant sex differences in core temperature were observed in other conditions (all $p > 0.10$). Individual core temperature responses and rates of core temperature change are provided in the **Supplemental Material**.

Skin temperature (T_{sk}) increased by the end of the 8-hour prolonged heat exposure across all five experimental conditions (all $p < 0.01$). Significant sex differences in the mean skin temperature were observed at 40°C/55.0% RH, 44°C/29.2% RH, 47°C/35.6% RH and 50°C/24.5% RH environments. Specifically, males exhibited higher post-exposure T_{sk} values than females at 40°C/55.0% RH, 44°C/29.2%, while females had higher post-exposure T_{sk} than males at 47°C/35.6% RH and 50°C/24.5%RH (interaction: all $p < 0.05$).

Figure 2 Time course of core temperature and time to reach steady-state (T_{rec} rise rate ≤ 0.10 °C/h) under five different environmental conditions. A, 36°C/74.5% RH; B, 40°C/55.0% RH; C, 44°C/29.2% RH; D, 47°C/35.6% RH; E, 50°C/24.5% RH; F, time to reach steady-state for both sexes (males: n=20; females: n=16). Data were analysed by one-way mixed model ANOVA. *, $p < 0.05$.

DISCUSSION

Maintaining strict humidity control during physiological trials is crucial for accurately determining the critical environmental limit (or uncompensability limit). If the actual relative humidity exceeds the target value but remains undetected, it could lead to an underestimation of the uncompensability limit, and vice versa. Theoretically, even a 5% fluctuation in RH can significantly alter the wet-bulb temperature (T_W). For instance, at 40°C, increasing RH from 45% to 50% raises T_W by 1.1°C, while at 50°C, increasing RH from 15% to 20% results in a 2.4°C rise in T_W. The precise environmental control in our study ensures accurate assessments of environmental compensability and provides reliable validation for the predictive methods being evaluated (i.e., the CTIP and biophysical modeling). Additionally, when determining critical environmental limits, environmental temperature and relative humidity at a minimum of four heights (in our study, we used 0.1 m, 0.6 m, 1.1 m, and 1.7 m) inside the test chamber should be continuously monitored (ISO7730, 2005) and adjusted throughout the prolonged exposure to ensure the strict maintenance of the target wet-bulb temperature.

The observed metabolic rates in our study were comparable to MiniAct (minimal activity, average metabolic rate: 1.34-1.51 METs) measured in Vecellio et al.'s study (2022), as well as that used for biophysical model simulations (i.e., 1.5 METs) by Vanos et al. (2023). Additionally, we observed a 15.4% to 20.3% increase in participants' metabolic rate after the 8-hour heat exposure, which aligns well with existing literatures reporting a comparable metabolic rate increase (i.e., 7-23%) per degree Celsius rise in core temperature (Saxton, 1981; Cramer et al., 2022). It should be noted that the participant's metabolic rate was assumed to be constant when employing the CTIP method and biophysical modeling to estimate critical environmental limits. This fails to replicate the actual metabolic process for both methods and thereby, introducing errors to the estimated critical environmental limits (or uncompensability limit).

Our findings reveal critical discrepancies between empirical results and predictions from biophysical models and the CTIP method, exposing significant limitations in accurately defining the "uncompensability limit". Biophysical models, which rely on simplified heat exchange equations, often fail to account for the complexities of human thermoregulation, behavioral adaptations, and sex differences (see Figure 3). For instance, adaptive behaviors such as periodic sweat drying, which enhances evaporative cooling, play a vital role in mitigating heat stress, particularly in high-humidity environments (Piantadosi, 2023). Our study found that participants were able to endure all five environmental conditions for

prolonged durations, in contrast to model predictions that suggested a rapid onset of heatstroke. Notably, the biophysical model estimated that participants could reach a core temperature (T_{core}) of 43°C within 6 hours under the 47°C with 35.6% RH and 50°C with 24.5% RH conditions. However, the actual physiological responses demonstrated greater tolerance than predicted, highlighting the limitations of the model in capturing human adaptive behaviors and thermoregulatory responses. A significant limitation of biophysical models is their lack of real-world validation under survival threshold conditions. To the best of our knowledge, models such as those used by Vanos et al. (2023) have not been tested in the extreme environments examined in our study. This absence of empirical validation likely contributes to the substantial overestimation of heat stress and underestimation of human survivability observed in their predictions.

Figure 3 Critical gaps in biophysical models in predicting human responses to extreme heat stress. (A) Lack of consideration for behavioral thermoregulation strategies, such as hydration status and periodic skin drying (Tansey and Johnson, 2015). (B) Neglect of sex differences, including variations in sweating response, body physique, and cardiovascular capacity (Kaciuba-Uscilko and Grucza, 2001). (C) Omission of cellular response mechanisms, including mitochondrial function, cytokine secretion (e.g., IL-6, IL-1β), and heat shock protein (HSP) expression (Leon et al., 1998). These responses can significantly influence the rate and extent of core temperature (T_{core}) rise. Pro-

inflammatory cytokines may increase metabolic heat production and impair heat dissipation, while HSPs stabilize proteins and enhance metabolic efficiency (Fields, 2001). Additionally, adaptive mitochondrial and vascular changes improve heat tolerance over time. (D) Insufficient model validation under extreme conditions, with no testing at T_W=35°C or during prolonged extreme heat exposures.

Many models neglect sex differences in thermoregulatory responses. Our results showed that females exhibited slower core temperature rises than males (Figure 2C), likely due to variations in body composition, metabolic heat production, and vascular adaptations (Piantadosi, 2023). These differences suggest that females may have greater heat tolerance during prolonged uncompensable exposure. However, treating humans as a homogenous group, as most biophysical models do, leads to inaccuracies in predicting heat tolerance for mixed-sex populations. Incorporating sex-specific data into these heat tolerance models is therefore essential to improve their accuracy and applicability.

The core temperature inflection point (CTIP) method also has notable shortcomings. This method assumes that a transient rise in core temperature over a short timeframe (5 minutes) indicates a loss of heat balance, thereby signaling uncompensable heat stress and often predicting heatstroke at lower wet-bulb temperatures (e.g., critical T_W averaging 30.55±0.98°C) (Vecellio et al., 2022). However, such short-duration fluctuations more likely reflect a delay in thermoeffector responses rather than true heat balance (Meade et al., 2024; Wang, 2025). Our prolonged 8-hour trials at the three chosen conditions (36°C/74.5% RH, 40°C/55.0% RH, and 44°C/29.2% RH) demonstrated that all these conditions remain compensable in both sexes with full hydration (see Figure 2F), directly contradicting the CTIP's reported critical wet-bulb temperatures and highlighting its limitation in real-world conditions. Additionally, our results indicate that adaptive mechanisms, such as enhanced sweating efficiency, fluid intake, and increased skin blood flow, can effectively slow the core temperature rise. While not the primary focus of our study, previous research supports the role of these adaptive mechanisms in delaying heatstroke onset and mitigating heat stress risks (Wang et al., 2024). Taken together, the CTIP method fails to account for the delayed effects of thermoregulation during prolonged exposure, potentially underestimating human survivability. Future studies should incorporate real-world factors, such as hydration status, behavioral adaptations, and physiological variability, to refine the definition of uncompensable thresholds and improve predictive accuracy of human heat tolerance.

FUTURE PERSPECTIVES

Advancing heat tolerance research necessitates addressing current limitations by incorporating empirical data from human survival studies and capturing the full spectrum of physiological and behavioral responses to prolonged heat stress. Real-world survival mechanisms such as periodic sweat drying with tissues or towels and fanning with hands, should be incorporated into models to enhance their accuracy. Furthermore, accounting for physiological differences across sexes and populations is essential to ensure applicability across diverse scenarios and prevent overgeneralization. Refining biophysical models with these considerations will significantly improve their predictive power for human heat tolerance under extreme conditions. While the CTIP and biophysical modeling methods may offer rapid estimations, it lacks the precision and reliability of prolonged physiological exposure trials in defining uncompensable stress and human survival thresholds. Traditional prolonged physiological trials remain the gold standard for accurately assessing human resilience to extreme heat. Future physiological research should prioritize prolonged heat exposure trials to better replicate real-world conditions. Such studies would enable a deeper understanding of adaptive mechanisms, including enhanced sweating efficiency, fluid intake management, and vascular responses, which may not be evident in shorter trials (Wang, 2025). Beyond acute heat tolerance, it is also critical to investigate the chronic health impacts of prolonged heat exposure on cardiovascular, renal, and cognitive systems. Addressing these long-term risks is essential for developing effective public health interventions and climate adaptation strategies in response to escalating climate change-driven heatwaves. By tackling research priorities, future research will provide the comprehensive insights needed to mitigate health risks and enhance human resilience to extreme heat.

LIMITATIONS

Our study did not aim to define uncompensable heat stress limits, and thus such critical environmental limits remain undetermined in this study. While our focus was validating the Core Temperature Inflection Point (CTIP) method and biophysical modeling, future research is warranted to establish precise uncompensability limits. Prolonged heat exposure trials across a range of wet-bulb temperatures could better elucidate human heat tolerance, survivability, and adaptability during extreme heatwaves, enhancing the applicability of these methods to real-world scenarios.

CONCLUSIONS

This study demonstrates that all five tested conditions (36°C/74.5% RH, 40°C/55.0% RH, 44°C/29.2% RH, 47°C/35.6% RH, 50°C/24.5% RH) were compensable over 8 hours, with participants maintaining stable core temperatures under extreme heat stress. Contrary to predictions, both the Core Temperature Inflection Point (CTIP) method and biophysical modeling significantly underestimated the critical environmental limits (CELs), suggesting that these approaches may not accurately reflect human heat tolerance during prolonged exposure. These findings reveal a critical discrepancy between model-based thresholds and observed physiological responses, particularly under real-world-relevant durations (such as the 8-hour exposure used in this study), as many methods typically reply on only 1-3 hours of exposure. Our results emphasize the need to refine these methods to account for extended heat exposure, enhancing their reliability for occupational and public health applications. Future studies should investigate higher wet-bulb temperatures or longer exposure durations to more accurately define the true uncompensability threshold (or CELs) across diverse populations.

ORCID

Faming Wang: https://orcid.org/0000-0002-2945-4685

Tze-Huan Lei: https://orcid.org/0000-0003-3157-099X

Haojian Wang: https://orcid.org/0009-0004-1469-7539

Data availability statement

The data supporting the present findings are provided in the Supplemental Material.

Acknowledgements

This research was financially supported by the National Excellent Young Scientist Program (grant number: 6119924022, to FW). We would also like to express our gratitude to all participants who took part in the study for their invaluable contributions to this research.

Author contributions

All authors from this study meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship, and all those who qualify are listed. All authors had full access to and accept responsibility for the data presented in the study. FW conceived the research question and acquired funding. FW designed the trial. FW, TL, HW, YX, and HX collected data. FW, HW, TL and HX processed data. TL performed statistical analysis. FW, TL and YX created the data visualizations. FW drafted the manuscript. All authors revised and approved the final version of manuscript.

Declaration of interests

The authors declare no competing interests.

REFERENCES

American Society of Heating, Refrigeration and Air Conditioning Engineers. Thermal Environmental Conditions for Human Occupancy. ASHRAE Standard 55, ASHRAE, USA, 2023.

Avellini BA, Kamon E, Krajewski JT. Physiological responses of physically fit men and women to acclimation to humid heat. J. Appl. Physiol. Respir. Environ. Exerc. Physiol. 49:254-261, 1980. doi: 10.1152/jappl.1980.49.2.254.

Belding HS, Kamon E. Evaporative coefficients for prediction of safe limits in prolonged exposures to work under hot conditions. Fed. Proc. 32(5):1598-1601, 1973.

Bouman A, Heineman MJ, Faas MM. Sex hormones and the immune response in humans. Hum. Reprod. Update. 11(4):411-423, 2005. doi: 10.1093/humupd/dmi008.

Buono MJ, Heaney JH, Canine KM. Acclimation to humid heat lowers resting core temperature. Am. J. Physiol. 274(5):R1295-R1299, 1998. doi: 10.1152/ajpregu.1998.274.5.R1295.

Cramer MN, Gagnon D, Laitano O, Crandall CG. Human temperature regulation under heat stress in health, disease, and injury. Physiol. Rev. 102(4):1907-1989, 2022. doi: 10.1152/physrev.00047.2021.

Fields PA. Review: Protein function at thermal extremes: balancing stability and flexibility. Comp. Biochem. Physiol. A. Mol. Integr. Physiol. 129(2-3):417-431, 2001. doi: 10.1016/s1095-6433(00)00359-7.

Hanna JM, Brown DE. Human heat tolerance: an anthropological perspective. Annu. Rev. Anthropol. 12:259–284, 1983. doi: 10.1146/annurev.an.12.100183.001355.

International Organization for Standardization. Ergonomics of the thermal environment — Analytical determination and interpretation of thermal comfort using calculation of the PMV and PPD indices and local thermal comfort criteria. ISO Standard 7730, ISO, Geneva, Switzerland, 2005.

Jetté M, Sidney K, Blümchen G. Metabolic equivalents (METS) in exercise testing, exercise prescription, and evaluation of functional capacity. Clin. Cardiol. 13(8):555-565, 1990. doi: 10.1002/clc.4960130809.

Kaciuba-Uscilko H, Grucza R. Gender differences in thermoregulation. Curr. Opin. Clin. Nutr. Metab. Care. 4(6):533-536, 2001. doi: 10.1097/00075197-200111000-00012.

Kamon E, Avellini B. Physiologic limits to work in the heat and evaporative coefficient for women. J. Appl. Physiol. 41(1):71-76, 1976. doi: 10.1152/jappl.1976.41.1.71.

Kang Z, Wang F, Udayraj. An advanced three-dimensional thermoregulation model of the human body: development and validation. Int. Commun. Heat Mass Transfer 107:34-43, 2019. doi: 10.1016/j.icheatmasstrasnfer.2019.05.006.

Kenny GP, Wilson TE, Flouris AD, Fujii N. Heat exhaustion. Handb. Clin. Neurol. 157:505-529, 2018. doi: 10.1016/B978-0-444-64074-1.00031-8.

Lei TH, Lu C, Lei Y, Wang F. Electric fan use and fluid replacement fail to prevent systemic inflammatory response syndrome (SIRS) during prolonged humid heat in young men. Build. Environ. 270:112539, 2025. doi:10.1016/j.buildenv.2025.112539.

Leon LR, White AA, Kluger MJ. Role of IL-6 and TNF in thermoregulation and survival during sepsis in mice. Am. J. Physiol. 275(1):R269-R277, 1998. doi: 10.1152/ajpregu.1998.275.1.R269.

Luber G, McGeehin M. Climate change and extreme heat events. Am. J. Prev. Med. 35(5):429-435, 2008. doi: 10.1016/j.amepre.2008.08.021.

Meade RD, Notley SR, Kenny GP. Time to reach equilibrium deep body temperatures in young and older adults resting in the heat: a descriptive secondary analysis. Am. J. Physiol. Regul. Integr. Comp. Physiol. 327: R369–R377, 2024. doi:10.1152/ajpregu.00089.2024.

Mora C, Dousset B, Caldwell I. et al. Global risk of deadly heat. Nature Clim Change 7:501–506, 2017. https://doi.org/10.1038/nclimate3322.

Oppliger RA, Magnes SA, Popowski LA, Gisolfi CV. Accuracy of urine specific gravity and osmolality as indicators of hydration status. Int. J. Sport. Nutr. Exerc. Metab. 15(3):236-251, 2005. doi: 10.1123/ijsnem.15.3.236.

Pal J, Eltahir E. Future temperature in southwest Asia projected to exceed a threshold for human adaptability. Nature Clim. Change 6:197–200, 2016. https://doi.org/10.1038/nclimate2833.

Piantadosi CA. Tolerance to Heat. In *The Biology of Human Survival: Life and Death in Extreme Environments*. Oxford University Press, Oxford, 2023.

Ramanathan NL. A new weighting system for mean surface temperature of the human body. J. Appl. Physiol. 19:531-533, 1964. doi: 10.1152/jappl.1964.19.3.531.

Satish U, Mendell MJ, Shekhar K, Hotchi T, Sullivan D, Streufert S, Fisk WJ. Is CO2 an indoor pollutant? Direct effects of low-to-moderate CO2 concentrations on human decision-making performance. Environ. Health Perspect. 120(12):1671-1677, 2012. doi: 10.1289/ehp.1104789.

Saxton C. Effects of severe heat stress on respiration and metabolic rate in resting man. Aviat. Space Environ. Med. 52(5):281-286, 1981.

Sherwood SC, Huber M. An adaptability limit to climate change due to heat stress. Proc. Natl. Acad. Sci. U. S. A. 107(21):9552-9555, 2010. doi: 10.1073/pnas.0913352107.

Sherwood SC, Ramsay EE. Closer limits to human tolerance of global heat. Proc. Natl. Acad. Sci. U. S. A. 120(43):e2316003120, 2023. doi: 10.1073/pnas.2316003120.

Tansey EA, Johnson CD. Recent advances in thermoregulation. Adv. Physiol. Educ. 39(3):139-148, 2015. doi: 10.1152/advan.00126.2014.

Vanos J, Guzman-Echavarria G, Baldwin JW, Bongers C, Ebi KL, Jay O. A physiological approach for assessing human survivability and liveability to heat in a changing climate. Nat. Commun. *14*, 7653, 2023. doi: 10.1038/s41467-023-43121-5.

Vecellio DJ, Wolf ST, Cottle RM, Kenney WL. Evaluating the 35°C wet-bulb temperature adaptability threshold for young, healthy subjects (PSU HEAT Project). J. Appl. Physiol. 132:340–345, 2022. doi:10.1152/japplphysiol.00738.2021.

Wang F. Rethinking the core temperature inflection method: misjudging thresholds of human adaptability to extreme heat. Am. J. Physiol. Regul. Integr. Comp. Physiol. 2025. doi:10.1152/ajpregu.000xx.2025.

Wang F, Deng Q, Lei TH, Wang X, Wang AR. Biophysical modelling predicts unreliable core temperature response on healthy older adults using electric fans at residential homes during heatwaves. Build. Environ. 228:109888, 2023. doi: 10.1016/j.buildenv.2022.109888.

Wang F, Lu C, Lei TH, Lei Y. Fluid replacement and acute systemic inflammatory response syndrome during prolonged eight-hour extreme heat exposure in young men. bioRxiv. 2024, doi:10.1101/2024.12.12.628082.

Wolf ST, Cottle RM, Fisher KG, Vecellio DJ, Kenney WL. Heat stress vulnerability and critical environmental limits for older adults. Commun. Earth Environ. **4**:486, 2023. doi: 10.1038/s43247-023-01159-9.

Xu X, Rioux TP, Castellani JW, Montain SJ, Charkoudian N. Validation of livability environmental limits to heat and humidity. J. Appl. Physiol. 137(6):1642-1648, 2024. doi: 10.1152/japplphysiol.00225.2024.