The bacterial chaperone CsgC inhibits functional amyloid CsgA formation by

promoting the intrinsically disordered pre-nuclear state

Anthony Balistreri[1]*, Divya Kolli[1]*, Sanduni Wasana Jayaweera[3], Daniel Lundahl[3], Yilin Han[2], Lily Kalcec[1], Emily
Goetzler[1], Rachel Alessio[1], Brandon Ruotolo[2], Anders Olofsson[3], Matthew R. Chapman[1]†.

1 Department of Molecular, Cellular and Developmental Biology, University of Michigan, Ann Arbor, MI 48109,
USA

2 Department of Chemistry, University of Michigan, Ann Arbor, MI 48109, USA

3 Department of Medical Biochemistry and Biophysics, Umeå University, 901 87 Umeå, Sweden.

† Correspondence: Matthew R. Chapman

Email: champmanm@umich.edu

Abstract

E. coli secretes a functional amyloid called curli during biofilm formation. Curli fibers are

composed of polymers of the CsgA protein, which adopts a beta-sheet rich fold upon

fibrillization. A chaperone-like protein called CsgC inhibits CsgA amyloid formation. Like other

amyloidogenic proteins, CsgA undergoes a 3-stage aggregation process: an initial lag phase

where a beta-rich nucleus forms, an exponential elongation phase, and a plateau phase. It is

currently not known if CsgC inhibits amyloid formation by inhibiting formation of the pre-fibril

nucleus, or rather, if CsgC inhibits a later stage of amyloid formation by blocking monomer

addition to a growing fiber. Here, CsgC homologs from *C. youngae*, *C. davisae*, and *H. alvei*

were purified and characterized for their ability to interrogate CsgA amyloid formation. Each of

the CsgC homologs prolonged the lag phase of *E. coli* CsgA amyloid formation in a similar

fashion as *E. coli* CsgC. Additionally, we found that *E. coli* CsgC interacted transiently and

weakly with a monomeric, pre-nucleus species of CsgA and that this interaction delayed amyloid

formation. A transient CsgC-CsgA heterodimer was observed using ion mobility-mass

spectrometry. When CsgC was added to actively polymerizing CsgA, exponential growth

commonly associated with nucleation-dependent amyloid formation was lost. However, the

addition of preformed CsgA seeds did not rescue exponential growth indicating that CsgC also

has inhibitory activity during fibril elongation. Indeed, CsgC interacted strongly with CsgA

fibers, suggesting that the interaction between CsgC and CsgA fibers can slow new fiber growth.

CsgC displays a unique inhibitory activity at multiple stages of amyloid formation. CsgC acts as

an energy-independent chaperone that transiently interacts with prefibrillar CsgA as well as an

amyloid fiber.

Introduction

Amyloids are fibril protein aggregates that are characterized by their distinct morphology,

stability, and tinctorial properties (1). Abnormal accumulation of amyloids can lead to a wide

range of pathologies, including diseases such as Parkinson's Disease, Alzheimer's Disease, and

Type II Diabetes (2). The proteins that comprise amyloid fibers are often intrinsically disordered,

or have intrinsically disordered regions, which adopt a repetitive cross-β strand structure upon

fibrilization (2). Amyloid intermediates are cytotoxic and have been shown to be more toxic than

mature amyloid fibers (3, 4). While inhibiting disease-associated amyloid formation remains a

significant focus of research, a growing body of work is focused on utilizing functional amyloids

as a model for amyloid formation and regulation.

Functional amyloids are a class of amyloid that provide a useful function for the cell, the list

of which continues to grow (5–7). The most widely studied functional amyloid is curli, the main

protein component of the *E. coli* biofilm matrix (6). Curli fibrils surround cells in the biofilm and

provide protection from desiccation, predation, and viral infection (6). The *E. coli csg* operons

contain seven curli specific genes that function as the producer and regulator for curli formation,

with each protein playing a significant role (8). The main amyloid protein, CsgA, is translocated

into the periplasm by the SecYEG pore (6). Periplasmic CsgA is then shepherded to the

nonameric CsgG outer membrane pore by CsgE, a periplasmic chaperone (9). Two auxiliary

proteins, CsgB and CsgF, are also exported through the CsgG pore and act as a nucleator and

anchor, respectively, to allow CsgA amyloid fibers to form and attach to the cell's outer surface

(10, 11). Lastly, the operon contains a second periplasmic protein, CsgC, which functions as a

potent inhibitor of CsgA and CsgB amyloid formation (12).

CsgC is a 110 amino acid protein with an immunoglobulin-like β-sandwich structure and

proposed chaperone-like activity (13). *E. coli* CsgC is an efficient sub-stoichiometric inhibitor of

amyloid formation against its native client *E. coli* CsgA (12). CsgC can also inhibit amyloid

formation by *E. coli* CsgA homologs like the *Pseudomonas* amyloid FapC, and the human

disease-associated amyloid alpha-synuclein (12). Little is known about the CsgC-client protein

interaction, except i) the interaction is guided by electrostatic interactions (14) and ii) there is a

weak client sequence determinant (12). Many details about CsgC-mediated amyloid inhibition

remain poorly understood. CsgC is a simple protein with no predicted active sites, domains, or

significant amino acids. CsgC requires no hydrolysable substrate, cofactor, or cellular energy to

perform its activity. CsgC does not take up a substrate or produce a quantifiable product; the

only evidence of CsgC activity is the inhibition of amyloid formation. Therefore, CsgC

represents a potentially novel class of anti-amyloid chaperone proteins that do not utilize an

energy source and do not maintain a stable interaction with the client protein.

Amyloid formation occurs in a series of steps, during which soluble, disordered monomers

join an insoluble, ordered amyloid fiber. In a solution containing amyloid-competent monomeric

proteins, a low abundance species arises from the pool of disordered monomers which acts as a

nucleation site for fiber formation (15). This nucleation event represents the rate-limiting step for

amyloid formation. Disordered monomers add on to these "nuclei" and begin the progressive and

thermodynamically favorable process of forming amyloid fibers (15). Fiber formation marks the

transition of amyloid proteins from the soluble state to the insoluble, and in the case of curli

fibrils, a transition from an intrinsically disordered monomeric conformation to a stable and

ordered β-sheet rich aggregate.

Previous work suggested that CsgC inhibits CsgA amyloid formation via electrostatic

interactions (14), however, it is unclear what stage(s) of amyloid formation is inhibited by CsgC

and what conformer of CsgA that CsgC binds to (14, 16–18). Here, we used several biochemical

and biophysical techniques to provide direct evidence that CsgC can inhibit CsgA amyloid

formation through two different mechanisms. We tested CsgC homologs from four

Gammaproteobacteria and found that they inhibit *E. coli* CsgA amyloid formation kinetics

through more than one mechanism. CsgC transiently binds a sub-population of monomeric CsgA

on-pathway to form amyloid and prolongs the formation of the aggregation-prone β-sheet rich

fold state characteristic of amyloid proteins. CsgC strongly binds CsgA fibrils and blocks fibril

elongation. We propose that these chaperone-like mechanisms are in accordance with the *in vivo*

function of CsgC: preventing formation of cytotoxic amyloid aggregates in the bacterial

periplasm and acting as an amyloid fibril formation inhibitor.

Experimental procedures

Bacterial Growth

All overnight cultures were grown in LB supplemented with 100 μg/mL ampicillin and/or 50

μg/mL kanamycin at 37 °C with shaking at 220 rpm. When necessary, LB plates were

supplemented with ampicillin 100 μg/mL or kanamycin 50 μg/mL.

Strains and Plasmids

The full list of strains, plasmids, and primers can be found in the Supporting Information.

Gibson Assembly was used to construct plasmids using NEBuilder® HiFi DNA Assembly

Cloning Kit (Cat. No. E5520S). Primers were designed using the NEBuilder webtool

(https://nebuilder.neb.com) and purchased from IDT (https://www.idtdna.com). Mutagenized

plasmids were constructed in the MC1061 cell background. Correct mutations were confirmed

using Sanger sequencing provided by Eurofins (https://www.eurofins.com/genomic-services/our-

services/custom-dna-sequencing/). Plasmids were extracted from transformants using Promega

PureYield™ Plasmid Miniprep System (Cat No. PRA1223). Miniprepped plasmids were

transformed into an expression strain (NEB3016) for purification.

Protein Purification

CsgA was purified as described previously (19). CsgC and its variants were purified as

described previously (20). Size exclusion chromatography was performed for CsgC purification

as previously described (21). Briefly, Ni-NTA affinity chromatography elution fractions were

pooled, concentrated, and passed through a 0.22 μm filter. The sample underwent gel filtration

using a Superdex 75 10/300 GL column (Cat No.45-002-903) attached to an Äkta pure protein

purification system. Elution fractions were assayed for protein concentration using A220 and/or

Pierce™ BCA Protein Assay Kit (Cat No. 23225). Samples corresponding to elution peaks were

analyzed using SDS-PAGE.

Denaturing Gel Electrophoresis and Dot Blot

Purified protein samples were diluted in 4X SDS loading buffer and run on a 15% SDS

PAGE gel with 10 μL loaded into each well (21). The gels were stained with Coomassie blue

dye to visualize protein bands. Dot blots were performed by spotting a nitrocellulose membrane

with 2 μL of protein and allowing the spots to fully dry. Blots were blocked for one hour in a

mixture of TBST and skim milk. Blocked blots were then washed three times with TBST and

probed with a primary antibody solution against CsgA (1:12,000) (21) or CsgC (1:4000) (17).

Primary antibody was removed from blots which were then washed three times with TBST prior

to the addition of a secondary antibody solution. Secondary antibodies against rabbit IgG and

conjugated with IRDye 800CW (Cat No. NC9401842) were used to image the blots in a Licor

Odyssey FC.

Surface Plasmon Resonance

Surface Plasmon Resonance (SPR) experiments were performed on a BIAcore™ 3000 (GE

Healthcare) using a CM5 chip for the immobilization of sonicated CsgA fibrils and gel filtered

CsgA monomers of a stable CsgA mutant (22). Immobilization was carried out following

protocols from previous fibrillar SPR studies (23), with CsgA fibrils immobilized at 1300 RU

and monomers at 500 RU. After immobilization, the flow cells were probed with gel-filtered

CsgC at concentrations ranging from 4 μM to 0.25 μM (two-fold dilutions). CsgC was injected

for 5 minutes over the CsgA fibrils and 7.5 minutes over the CsgA monomers. The experiment

was conducted in phosphate buffer (pH 7.5) containing 0.05% Tween, at a temperature of 25°C.

Prior to immobilization, CsgA fibrils were sonicated for 1 minute in a water bath.

Preparation of modified NHS columns

CsgC was purified as described previously (20). NHS-activated agarose beads were buffer

exchanged into 50 mM potassium phosphate buffer (pH = 7.4) and reacted with purified CsgC

(1mg CsgC/ml NHS-activated agarose beads, Cat. No. 26200) overnight at 4 °C with gentle

rocking. The reacted resin was then washed with 50 mM potassium phosphate buffer, pH = 7.4

to remove any unbound CsgC and cleaved NHS-groups. Reaction progress was tracked by 260

nm and 220/280 nm absorbance values to measure increase in cleaved NHS-groups and decrease

in free CsgC protein, respectively. SDS-PAGE and Pierce™ BCA Protein Assay Kit (Cat No.

23225) were used to ensure complete binding of CsgC to the resin. Unreacted NHS groups still

bound to agarose beads were cleaved by addition of a quenching buffer (1M Tris HCl, pH 7.5)

and incubated for 30 mins at room temperature with gentle rocking. The fully reacted resin was

buffer exchanged into 50 mM potassium phosphate buffer, pH = 7.4 to remove quenching buffer.

260 nm and 220/280 nm absorbance values were measured to ensure complete quenching of

unreacted NHS-linked agarose beads. Modified NHS columns were stored at 4 °C until use.

Individual columns were only used once for each assay.

Sample Preparation for Ion mobility-mass spectrometry (IM-MS)

Purified CsgA and CsgC were buffer exchanged into 20 mM ammonium acetate (pH 7.4)

using Thermo Scientific Zeba™ Spin Desalting Columns 7k MWCO (Cat No. 89892). The

protein concentration after buffer exchange was assayed using Thermo Scientific Pierce™ Rapid

Gold BCA Protein Assay Kit. CsgA and CsgC were each diluted to 20 µM with 20 mM

ammonium acetate, mixed at 1:1 ratio. The mixture was incubated at 37°C for 23 hours. Time

points were taken at 0 hr, 3 hr, 6hr and 23hr.

IM-MS

IM-MS data was collected on a quadrupole ion-mobility time-of-flight (TOF) mass

spectrometer (Synapt G2 HDMS, Waters, Milford, MA, USA) with a nano-electrospray

ionization (nESI) source. The source was operated at positive mode with the nESI voltage set at

1.0-1.2 kV, the sampling cone was set to 15 V and the bias was set to 42 V. The source

temperature was set to 20°C. The traveling–wave ion mobility separator operated at a pressure of

approximately 3.4 mbar with wave height and wave velocity set at 30V and 500 m/s,

respectively. The m/z window was set from 100 – 8000 m/z with a TOF pressure of 1.5e-6 mbar.

Mass spectra were analyzed using MassLynx 4.1 and Driftscope 2.0 software (Waters, Milford,

MA, USA). CCS (Ω) measurements were externally calibrated using a database of known values

in helium. We reported the standard deviations from replicate measurements of CCS and an

additional ±3% to incorporate the errors involved in the calibration process.

Thioflavin T Binding Assay

Assays were performed as previously described (19). Briefly, freshly purified CsgA was

diluted with phosphate buffer to 20 µM or 10 µM and combined with an excess of the amyloid-

specific dye thioflavin-T (ThT) (Cat No. AC211760250). Amyloid formation was monitored by

measuring an increase in ThT fluorescence at 495 nm (450 nm excitation). Assays were

performed in triplicate at microscale within 96-well plates and measured with Infinite Pro M200

or Infinite Nano⁺ F200 Tecan plate readers. CsgC proteins were purified, diluted with phosphate

buffer, and added to specified assays in the reported stoichiometric ratio. Fiber stimulating

"seeds" were produced by obtaining previously purified WT CsgA fibers and were sonicated

directly before adding to a specified assay. Time to half-maximum ($T_{1/2}$) is calculated by the

time at which the Thioflavin-T fluorescence reaches half of the maximum Thioflavin-T

fluorescence. Duration of lag phase (T_{lag}) was calculated as described by Arosio et al (24).

Briefly, T_{lag} is calculated by $T_{1/2}$ subtracted by half of the maximum of the first derivative of the

sigmoidal ThT fluorescence.

Circular Dichroism

CsgA was diluted with phosphate buffer to a final concentration of 15 μM. Secondary

structure was measured at room temperature every 60 minutes (in triplicate) with the JASCO-

1500 using default parameters. Measurements were averaged, the baseline signal of phosphate

buffer was subtracted, and the curve was smoothed using a Savitzky-Golay filter. Final curves

were normalized by concentration and path length.

Results

E. coli CsgC and three CsgC homologs affect multiple stages in the CsgA amyloid formation

Pathway

CsgC was first identified in *E. coli* (EC), but *csg* operons from other Gammaproteobacteria

contain CsgC homologs (25). *Citrobacter youngae* (CY), *Cedecea davisae* (CD), and *Hafnia*

alvei (HA) are three Gammaproteobacteria that have *csg* operons containing homologs to EC

CsgC. These four CsgC homologs have similar structures, as predicted by Alphafold 2.0 (26),

however, they share a range of sequence identity with CsgC EC: CY (69%), CD (51%), and HA

(35%) (**Fig. 1A-D and Fig. S1**). CsgC and each homolog was tested for inhibitory activity

against *E. coli* CsgA using ThT binding assays. CsgC EC was the most efficient inhibitor of *E.*

coli CsgA amyloid formation, with full inhibition at a 1:500 stoichiometric ratio (40 nM) (**Fig.**

1E). CsgC CY, CsgC CD, and CsgC HA all inhibited CsgA aggregation to a lesser extent (**Fig.**

1F-H). All ThT assay results were fit with a sigmoidal function curve. We derived the duration

of the lag phase for each curve as a proxy for the nucleation rate of CsgA polymerization (24).

The lag phase of *E. coli* CsgA was significantly prolonged upon addition of CsgC homologs in

11 of 12 assays (**Fig. 1I**). The lag phase was not determined in two cases because the ThT signal

never increased within the time frame of the experiment (**Fig. 1I**). The only condition where

Fig. 1. CsgC EC and its homologs are effective at extending the lag phase CsgA amyloid formation. A-D) AlphaFold 2.0 predicted structures for CsgC EC, CY, CD, and HA. **E-H)** 4 different CsgC homologs were purified and tested for their ability to inhibit CsgA in vitro. In all cases CsgA was freshly purified and mixed in the stated stoichiometric ratio with CsgC homologs. The data points shown represent the average of triplicate experiments, the error bars show the SEM, and the curves were fit using the sigmoidal logistic function described in (24). **I)** The calculated lag phase for each homolog according to the fit curves. The error bars represent the SEM for each value. "nd" was used to denote values which were not defined by the equation used. Significance was attributed using a one-way ANOVA analysis comparing all values to the uninhibited CsgA condition; *p < 0.05, ****p < 0.00005

addition of CsgC did not affect the lag phase was with addition of the lowest ratio (1:1000) of

CsgC HA homolog, which has the lowest sequence identity to CsgC EC (**Fig. 1H and I**).

Amylofit was used for additional analysis of the ThT curves produced in **Fig. 1**. Amylofit is a

global fitting software that fits ThT binding assay curves, calculates estimated values for reaction

orders, rate constants, and more to describe amyloid aggregation kinetics (27). Meisl and

coworkers suggest that Amylofit can be used to determine the molecular species or stage along

an aggregation pathway that is affected by an inhibitor (27). The rate expressions used to fit ThT

curves describe the formation of CsgA amyloid fibers starting with unfolded CsgA monomers:

$$CsgA_{Disordered} \overset{k_n}{\underset{}{\rightleftarrows}} CsgA_{Nucleus} \quad (1)$$

$$CsgA_{Nucleus} + CsgA_{Monomer} \overset{k_+}{\underset{}{\rightleftarrows}} CsgA_{Fiber} \quad (2)$$

A primary nucleus forms in solution from an unknown number of folded CsgA monomers (k_n,

eq.1). After a critical concentration of nuclei are present, CsgA monomers use nuclei as folding

templates, sparking rapid fiber elongation (k_+, eq.2). For some amyloid proteins, the presence of

amyloid fibers in solution stimulates new fiber growth (this could be described by rate constant

k_2).

We inputted the same data displayed in **Fig. 1E-H** into Amylofit and normalized each curve

to reflect the final aggregate amount in the uninhibited CsgA condition. The "secondary

nucleation dominated" model was used to fit the ThT curves because, under these aggregation

conditions, this model provided the best fit. At first, all rate constant parameters (k_n, k_2, and k_+)

were allowed to be fit individually, providing the best fit curves for each condition (**Fig. S2A,**

S3A, S4A, and S5A). Next, all parameters except one were set to a global constant using the

values derived from CsgA fibril formation in the absence of inhibitor. This was repeated for all

rate constant parameters. Using this method, each rate constant parameter was set to individually

fit each inhibited condition (**Fig. S2B-D, S3B-D, S4B-D, and S5B-D**). This method illustrated

the degree to which each rate constant can explain changes in the ThT curve upon addition of

increasing concentrations of inhibitor (27). Best fit curves were created for each condition and

the mean squared residual error (MRE) results from each condition provided a global metric for

goodness of fit for all curves (27). Low MRE values are an indication that the fitted rate constant

is affected by the addition of the inhibitor. Interestingly, the rate constant that provided the

lowest MRE value was different between each

homolog, but MRE values were generally

close to the universal best fit MRE value (**Fig.**

S2E, S3AE, S4E, and S5E). The rate of

secondary nucleation (k_2) and elongation rate

(k_+) often resulted in lower MRE values than

the primary nucleation rate (k_n), suggesting

that CsgC could be affecting CsgA

aggregation at the level of fibril elongation.

Taken together, **Fig. 1-2** and **Fig. S2-5**

indicated that CsgC and its homologs may

affect CsgA amyloid formation at multiple

stages.

CsgC partially inhibits CsgA amyloid

formation during the elongation phase

CsgA amyloid formation follows a

primary nucleation-dependent polymerization

mechanism that has three distinct phases: an

initial lag phase, an elongation phase, and a

stationary phase. The elongation phase is

marked by fibril elongation directed by

soluble protein monomer addition onto fiber

ends (28, 29). Importantly, even during the

elongation phase there are intrinsically

disordered CsgA monomers still in solution

capable of either primary nucleation or

addition onto a fiber (30).

CsgC was added at different time points to

actively polymerizing CsgA to determine how

Fig. 2. CsgC inhibits amyloid formation during the fiber elongation stage. A) 200 nM (1:100) CsgC was added to actively polymerizing CsgA at the times indicated and amyloid aggregation was monitored in a ThT binding assay. **B)** In a similar assay, CsgC was added to a 2% seeded reaction of CsgA at various ratios as indicated in the key above (All olive conditions also contain 2% seeds).

285 CsgC inhibitory activity changes throughout amyloid formation. First, CsgC was added to a final

286 concentration of 200 nM (1:100 CsgC:CsgA molar ratio) at different time points after the start of

CsgA amyloid formation (0 hours) up to 4 hours (**Fig. 2A**). When CsgC was added at time 0 hours the ThT signal remained very low compared to CsgA samples without CsgC, indicating nearly total inhibition. When CsgC was added to CsgA after 1 hour, the ThT signal increased linearly (**Fig. 2A**). When CsgC was added after the lag phase had ended, the result was a linear increase in ThT signal followed by a gradual decrease reaching a plateau in signal (**Fig. 2A**) indicating that CsgC was capable of slowing fiber elongation. However, it is not clear whether the inhibition is due to prevention of new nuclei formation or prevention of fibril elongation.

To determine if CsgC inhibitory activity is solely based on preventing nuclei formation, CsgC was added to CsgA that had been seeded by the addition of preformed CsgA fibers (**Fig. 2B**). During a seeded reaction, CsgA is presented with preformed fibers and the fibers act as a template for amyloid formation, effectively diminishing the need for primary nucleation and the lag phase entirely (**Fig. 2B, dark blue squares compared to black circles**) (31). When CsgC was added to a seeded CsgA reaction, the result was a linear increase in ThT signal (**Fig. 2B**). In both cases (**Fig. 2A and B**), the addition of CsgC into solution when fibril formation is largely dependent on CsgA monomer addition to actively growing amyloid fibrils resulted in partial inhibition of amyloid formation indicating that CsgC inhibited fibril elongation.

CsgC binds to CsgA fibrils and not to CsgA monomers

Surface plasmon resonance was used to determine the binding kinetics between CsgC and CsgA monomers or CsgA fibrils. Surface plasmon resonance can interrogate interactions between proteins and calculate a K_D (32). Either CsgA fibrils (**Fig. 3A**) or CsgA monomers (**Fig. 3B**) were bound to a CM5 chip using standard amino-coupling reaction. A double cysteine variant of CsgA that retains an intrinsically disordered structure under oxidizing conditions was used to prevent bound monomeric CsgA from aggregating during the course of the SPR experiment (22). Solutions containing freshly purified monomeric CsgC were flowed over the sensors.

The sensogram data with curve fits is shown in **Fig. 3**. The residuals indicated a generally good fit, with only minor deviations observed at certain concentrations. Notably, concentrations below 250 nM were excluded from the kinetic analysis due to detection limitations, as the responses at these lower concentrations were too weak to be reliably measured by SPR. Sensogram data indicated that monomeric CsgA had no measurable binding to monomeric CsgC

(**Fig. 3B**). However, monomeric CsgC interacted with CsgA fibrils with a dissociation rate of K_d

$= (1,18 \pm 0.05) \times 10^{-3}$ s^{-1} and an equilibrium dissociation constant of $K_D = 360 \pm 10$ nM (**Fig.**

3A).

Transient interaction between

monomeric CsgC and non-fibrillar

CsgA is sufficient to delay amyloid

formation

Previous studies have shown that

CsgA incubated with CsgC does not

readily form higher order fibrillar

aggregates (14, 17). This data, in

conjunction with data in **Fig. 1** and

Fig. S2-5, indicated that CsgC can also

inhibit the initial conversion of

monomeric CsgA into an oligomeric

amyloid. Due to the undetectable

binding between monomeric CsgC and

CsgA from **Fig. 3,** we utilized a

modified pull-down assay to determine

if CsgC interacts transiently with

prefibrillar CsgA.

N-hydroxy-succinimide (NHS)

resin beads were coupled to CsgC

protein for the modified pull-down

assay. Another batch of *N*-hydroxy-

succinimide (NHS) resin beads were

quenched with Tris to use as a control.

Immediately after Ni-NTA and SEC

purification, a CsgA sample was split

Fig. 3. Sensograms of CsgA fibrils and monomers probed by CsgC. A) Interaction between CsgA fibrils and CsgC, showing an equilibrium dissociation constant (K_D) of 360 ± 10 nM. The concentrations labeled "A" represents the first injection, while "B" denotes the secondary injection. The fitted curve represents the curve fitting performed using Scrubber2. **B)** No detectable interaction between CsgA monomers and CsgC across all tested concentrations. Residual standard deviation = 3,194 RU.

into equal parts. One aliquot of the purified CsgA was left on ice, while the other aliquot was

passed through a column containing either the CsgC-linked beads or the Tris quenched beads

(**Fig. 4A**). CsgA eluent from the column was then diluted into the wells of a ThT binding assay

to monitor amyloid formation. The CsgA from the Tris-quenched resin condition began

aggregating at 2 hours, consistent with CsgA that had been left on ice (**Fig. S6A**). Interestingly,

the CsgA that passed by the CsgC-linked column had a diminished propensity to form amyloid,

displaying an approximately 77 min increase in ThT lag phase and a 75 minute increase in the

time it takes to reach half maximum ThT fluorescence compared to CsgA that had been left on

ice (**Fig. S6B**). CsgA eluted from the CsgC-linked column displayed a 51 minute increase in the

lag time when directly compared to CsgA eluted from the Tris-quenched column and a 52 min

increase in the time it takes to reach half maximum ThT fluorescence, displaying that transient

interaction between CsgA and CsgC is sufficient for delayed amyloid formation **(Fig. 4B and**

D). Buffer was eluted from the Tris-quenched column and the CsgC-linked column and added

into a ThT binding assay with freshly purified CsgA to ensure that no CsgC, NHS, or Tris eluted

from the resin that could affect CsgA aggregation (**Fig. 4C**).

CsgA protein concentration was quantified prior to, and following, elution from each resin to

determine how much CsgA was retained on the CsgC-linked beads. Little to no difference was

observed between the eluent concentration of CsgA eluted from the control Tris-linked resin

beads (31.2 μM) and the eluent concentration of CsgA eluted from the CsgC-linked resin beads

(25.7 μM). To determine if the small difference in concentration was due to CsgA retention on

the CsgC-linked resin column, the CsgC-linked resin and Tris-linked resin was treated with

hexafluoroisopropanol (HFIP) to digest CsgA aggregates left on the resin beads. Samples were

then analyzed by a dot blot and SDS-PAGE to determine if any CsgA is retained on the beads

following elution. Both the SDS-PAGE gel and the dot blot indicated that no detectable CsgA

was retained on the resin following elution (**Fig. S7**). This further supports that CsgC transiently

interacts with CsgA, and that this interaction is sufficient to observe the extended lag phase in

Fig. 4.

Additionally, the change in CsgA secondary structure from disordered to beta-sheet rich was

visualized using circular dichroism **(Fig. S8)**. Amyloid protein folding is often assessed using

CD by measuring the formation of a spectral minimum at 220 nm indicating the formation of a

Fig. 4. Passing monomeric CsgA by immobilized CsgC delayed the formation of ThT positive aggregates. A) Affinity purified CsgA was passed through a column of NHS agarose resin that was either amino coupled to CsgC or Tris base to quench the free NHS binding spots. **B)** Eluent CsgA was then diluted to 10 µM in phosphate buffer for a ThT binding assay to observe amyloid aggregation kinetics. **C)** Quenched and CsgC resin was incubated in phosphate buffer for 24 hours. The incubated phosphate buffers were used to dilute freshly purified CsgA to 10 µM and added to a ThT binding assay. **D)** A measurement of the lag phase (T_{lag}) and time to half maximum fluorescence ($T_{1/2}$) was calculated using a sigmoidal curve function. Error bars represent the standard error of the mean across three technical replicates.

beta-sheet rich and amyloidogenic secondary structure. No significant effect of CsgC transient

interaction on beta-sheet structure formation was detected by circular dichroism. With the

indication of an interaction between CsgC and CsgA monomers by ThT but not by CD, we

decided to look more closely at their interaction using a highly sensitive native mass

spectrometry technique.

A transient 1:1 CsgC:CsgA complex is detected

Native mass spectrometry (native MS) uses electrospray ionization conditions to preserve

non-covalent protein-protein and protein-ligand complexes (33). The added dimension of the ion

mobility (IM) spectrometry allows ions to be separated based on their size, shape, and charge

(**Fig. 5A**) (34). We purified both CsgC monomers and CsgA monomers, added them together in

a 1:1 mixture, and analyzed the resulting solution for any interactions between the two proteins.

During our first time point, we observed monomeric and dimeric species of both proteins,

including low abundance heterodimeric complexes of CsgA and CsgC (**Fig. 5B and C**). CsgA

exhibited a broad charge state distribution ranging from 5+ to 13+, as typically observed for

intrinsically disordered proteins (**Fig. 5B**) (35). The CsgA:CsgC mixture remained soluble

throughout the 23 hour time course while apo CsgA, serving as a control sample, formed micro-

scale aggregates that led to the clogging of the nESI emitters and prevented the collection of both

6 hour and 23 hour time point data. An analysis of collisional cross-section (CCS) values of

CsgA monomers showed very little difference between time 0 and 3 hour (**Fig. S9A**). Similarly,

we also saw no changes in CCS for CsgC monomer ions through the 23 hour time course (**Fig.**

S9B).

We then analyzed the IM arrival time distributions (ATDs) to assess conformational changes

within CsgA and CsgC over time. The ATD data contains information regarding both ion shape

and the number of conformer families present for a single protein ion population (36). Data

shown in **Fig. 5D** displays three overlaid ATD profiles for CsgA 11+ monomer ions recorded

after 0 and 3 hour incubations. In addition, similar IM ATD data is shown for CsgA/CsgC

mixtures following a 3 hour incubation. All three datasets yield a bimodal ATD profile,

consisting of both a more compacted (shorter IM drift times) and a more extended

conformational family (longer IM drift times). A comparison of CsgA IM data collected at both

0 and 3 hour time points revealed the peak corresponding to the more compact conformation

decreased significantly in intensity, resulting in overwhelmingly a more extended protein ion

population. Overall, our IM ATD data suggested that the conformational ensemble of CsgA

becomes more extended as protein aggregation progresses, an observation common to other

amyloid proteins (37, 38). Importantly, when we compare our IM ATD data recorded for CsgA

at 0 hour with ATDs recorded for CsgA/CsgC mixtures following 3 hours of incubation, we

detected few significant differences, indicating that the initial, intrinsically disordered state of

Fig. 5. IM-MS reveals a 1:1 CsgA and CsgC heterodimer complex in solution and the stabilization of intrinsically disordered CsgA monomers. A) Proteins were passed through an ion mobility drift cell that separates based on size. Next, proteins pass through a time-of-flight mass analyzer. In combination, IM-MS can provide three modes of information for a given ion: mass to charge ratio, drift time, and the intensity of species with those properties. **B)** Mass spectra for CsgA incubated with CsgC in a 1:1 molar ratio at 37 °C. Monomeric CsgA (blue circles), dimeric CsgA (blue double circles), monomeric CsgC (purple triangles), dimeric CsgC (double purple triangles) and 1:1 CsgA:CsgC complexes (blue circle and purple triangle) are annotated. A magnified MS spectrum **C)** showed that the complex is seen flanked on either side by dimeric CsgA and dimeric CsgC. **D)** Arrival time distribution of the 11+ monomer of CsgA in three experimental conditions: Apo CsgA in solution at 0 hr (olive trace), apo CsgA in solution after 3 hr (light purple trace), and CsgA in solution with CsgC after 3 hr (dark grey trace). Traces are overlapped to show the changes in ATD after 3 hr of incubation.

CsgA is better preserved in the presence of CsgC. The results discussed above were consistent

across multiple CsgA charge states (**Fig. S9C and D**).

Discussion

Amyloid formation can lead to a wide range of pathologies. Therefore, it is of great interest

to discover and characterize molecules that can prevent pathogenic amyloid formation. Synthetic

amyloid inhibitor peptides occupy a large biochemical space, from β-blocking D-amino acid

peptides (39) to camelid nanobodies (40, 41). There is a short list of naturally occurring proteins

which putatively inhibit functional amyloid formation. This list includes molecular chaperones

and a diverse set of human and bacterial proteins which have similar 3D structures (18).

However, CsgC differs from the rest of the known amyloid inhibitor proteins in one significant

way: CsgC remains the only example of an amyloid inhibitor protein that exists exclusively to

moderate the amyloid formation of specific proteins. CsgC is part of the CsgBAC operon,

therefore, when amyloid proteins CsgA and CsgB are expressed, the CsgC protein will also be

expressed. Studying CsgC provides a unique opportunity to answer questions about how cells

prevent amyloid formation and utilize functional amyloids, as well as how other structurally

similar amyloid inhibitors may function.

CsgC is a highly effective anti-amyloid chaperone protein (17), however, CsgC's mechanism

of action has remained unclear. Early work found that CsgC could inhibit CsgA without the need

for a hydrolysable energy source and at a low sub-stoichiometric ratio (17). Key to

understanding the mechanism of CsgA amyloid inhibition is the nature of the CsgC interacting

partner. It has previously been suggested that CsgC interacts with elongating CsgA fibers (42),

and there are other amyloid inhibitor proteins that act to inhibit monomer addition to growing

fiber ends (43). However, molecular chaperones that interfere with amyloid formation have been

observed to interact with a variety of binding partners and inhibit aggregation at different steps

along the amyloid formation process (44). Given the multiplex role of monomers during amyloid

formation, interactions between an inhibitor and an amyloid-competent monomer results in a

decrease in the rate of several steps along amyloid formation (44). Our current data is consistent

with an interaction between CsgC and a monomeric, prefibrillar species of CsgA as well as

fibrillar CsgA at an extremely sub-stoichiometric ratio.

CsgC extends CsgA Lag Phase

When proteins self-assemble into amyloid fibers through a nucleation-dependent process

they typically display sigmoidal kinetics (24). There are three phases to polymerization

beginning with the so called "lag phase", a rate-limiting period of protein folding and initial

oligomerization (24). After the lag phase, a critical mass of nuclei has formed, and aggregation

increases exponentially as monomers quickly add onto fiber ends and new fibers continue to

sprout from nuclei. CsgA is known to act mostly through a primary nucleation pathway and not

through secondary nucleation where new fibers sprout from the lateral edge of an existing fiber
(18, 45, 46).

CsgA requires a longer time to begin rapid amyloid formation when CsgC is present in
solution. We saw lag phase extension in almost every scenario we tested, which included four
CsgC homologs in a wide range of substrate:CsgC concentrations (**Fig. 1**). The CsgC homologs
we tested share between 68% and 33% sequence identity with *E. coli* CsgC and yet, they all
affect CsgA in a similar manner. Moreover, there is a distinct correlation between the
concentration of the CsgC proteins and how much the lag phase was extended (**Fig. 1**). This
would indicate a true dose-response relationship between CsgC proteins and the extension of
CsgA lag phase and thereby the extent of primary nucleation. If CsgC is affecting nuclei
formation, CsgC could be interacting with a prenuclear species of CsgA.

CsgC inhibits new nuclei formation and fibril elongation during ThT elongation phase

CsgC remains an active amyloid inhibitor during the rapid fiber formation phase of
aggregation. We show two similar experiments in **Fig. 2** where CsgC was presented to CsgA
during the elongation phase. In the first experiment, CsgC was added to CsgA at different times
throughout aggregation and we monitored the effect of CsgC addition on the increase in ThT
fluorescence (**Fig. 2A**). CsgA alone takes approximately 1.3 hours to transition from the lag to
the elongation phase (**Fig. 2A**). When CsgC is added at time 0, we see little to no increase in ThT
fluorescence throughout the time course of the experiment, consistent with total inhibition of
nuclei formation (**Fig. 2A**). For all conditions tested where CsgC was added after time 0 there
was a linear increase in ThT signal, until the signal eventually reached a plateau and flattened out
(**Fig. 2A**). This indicates that the addition of CsgC after initial formation of CsgA nuclei yields
partial inhibition of CsgA amyloid formation. Linear or isodesmic polymerization is consistent
with a model of aggregation that is independent of nucleation (47). Nucleation-dependent
polymerization requires the formation of high energy nuclei, a bottleneck for polymerization,
followed by the very favorable addition of monomers to a growing fiber (48). CsgC-inhibited
CsgA could present an isodesmic polymerization profile because the formation of new nuclei is
being hindered while CsgA unfolded monomers can still add to fiber ends. CsgC could be
fighting the early process of CsgA transitioning between the disordered-denatured state and

disordered-collapsed state as described by Frieden (47). However, this result can also arise due to

inhibition of new nuclei formation as well as inhibition of fibril elongation.

To determine whether CsgC inhibition is solely based on inhibition of nuclei formation, a

similar experiment was performed in the presence of seeds to rescue CsgA amyloid formation.

Adding preformed fiber seeds to monomeric CsgA provides a template for the monomers to

quickly adopt an aggregation prone state and therefore add onto a growing fiber end. Seeds act as

catalysts, effectively lowering the free energy requirements to fibril formation. A linear increase

in ThT signal can be seen when CsgC is added to a seeded CsgA aggregation reaction (**Fig. 2B**)

indicating that the addition of seeds does not recapitulate un-inhibited CsgA amyloid formation.

Therefore, CsgC may be acting to inhibit nuclei formation as well as fibril elongation.

The surface plasmon resonance findings indicate robust binding between CsgC monomer and

CsgA fibrils (**Fig. 3**). Additionally, Amylofit data fitted to CsgA ThT binding assays in the

presence of varying concentrations of CsgC homologs indicate that addition of CsgC and CsgC

homologs drastically effect the rate of fibril elongation (**Fig. S2-5**). Taken together, it is likely

that CsgC inhibits at later stages in amyloid formation through a stable interaction with fibrils,

sterically preventing addition of CsgA monomers for fibril elongation.

It is worth noting that a "secondary-nucleation dominant" model was chosen to fit the

experimental data analyzed by Amylofit. Secondary nucleation is less prevalent in functional

amyloids as compared to disease associated amyloids and has not been observed in wild-type and

mutant CsgA fibril formation (46, 49). Despite this, CsgA aggregation best resembles an

aggregation model that accounts for both fibril growth and a form of self-replication (50). Future

directions include single-molecule imaging of CsgA amyloid fibril growth to quantify individual

fibril growth rates in the presence or absence of CsgC (51).

CsgC Transiently Interacts with the CsgA Monomer

While CsgC may inhibit CsgA fibril growth, its function *in vivo* is likely to prevent any

initiation of amyloid assembly which would be cytotoxic to the cell. Therefore, we sought to

determine the mechanism of CsgC inhibition of pre-nuclear CsgA. Surface plasmon resonance

indicated no measurable binding between CsgC monomers and CsgA monomers (**Fig. 3B**).

However, after quickly passing through a CsgC-linked column, freshly purified CsgA monomers

require approximately 50 additional minutes to begin forming ThT positive species (**Fig. 4B and**

D). The CsgA monomers were quickly pushed through the column by hand, therefore, any

interaction with immobilized CsgC was over a matter of seconds. And yet, there was lasting

impression on the CsgA, since aggregation was delayed (**Fig. 4D**). This suggests that a fast

CsgC-CsgA monomer interaction delays CsgA monomers from adopting an aggregation-prone

state.

Using IM-MS, we detected 1:1 CsgC and CsgA heterodimers, which indicates that CsgA and

CsgC monomers can interact with each other (**Fig. 5B**). The heterodimer peaks were in low

abundance, consistent with the concept of a transient interaction between CsgA and CsgC

monomers. Indeed, the heterodimer peaks are the first direct observation of an interaction

between full length CsgC and CsgA monomers.

CsgC Maintains Intrinsically Disordered Monomeric CsgA

Much can be gleamed from focusing on the IM-MS signals associated with CsgA monomers.

Multiple charge states of CsgA monomers can be seen in the mass spectrum, consistent with an

intrinsically disordered protein (**Fig. 5B**). We analyzed how the CsgA monomer's shape changed

by examining the ATD over time (**Fig. 5D, S9C-D**). The ATD profile suggests that the

conformation of CsgA became more extended as aggregation progresses, a change commonly

observed with other amyloid proteins as well (**Fig. 5D, S9C-D**) (37, 38). Remarkably, the

presence of CsgC in solution causes the ATD of CsgA monomers to remain consistent

throughout the time course of our experiment (**Fig. 5D, S9C-D**). Since CsgA monomers start as

intrinsically disordered, CsgC must be pushing CsgA monomers away from the aggregation-

prone state by promoting or stabilizing CsgA's intrinsically disordered fold. This is consistent

with the findings from **Fig. 4**, illustrating that transient interaction between CsgA and CsgC leads

to a memorable but temporary inhibition of CsgA amyloid formation.

Conclusions

We utilized an array of biophysical methods to show that CsgC interacts transiently with an

intrinsically disordered, monomeric CsgA to disrupt new nuclei formation in an ATP-

independent fashion. This model of inhibition fits within the greater context of curli biogenesis:

CsgC is a periplasmic protein tasked with inhibiting amyloid formation by CsgA within the

intermembrane space to prevent the formation of cytotoxic intracellular aggregates while also

allowing for efficient secretion of monomeric and intrinsically disordered CsgA for subsequent

amyloid formation on the bacterial surface. We also show that CsgC is capable of inhibiting

CsgA fibril formation even in the presence of CsgA nuclei, likely through stably interacting with

the amyloid fibril preventing further growth. Additionally, we show that CsgC utilizes a non-

canonical chaperone mechanism by interacting with its client protein transiently, remodeling

protein structure independently of a hydrolysable energy source, and promoting the unstructured

fold state. We propose that CsgC, and other similarly structured amyloid inhibitors, may

represent a new class of chaperone proteins capable of potent protein remodeling activity.

Data availability

551   All data generated or analyzed during this study are included in this article or are available from

552   the corresponding author upon reasonable request.

Supporting information

554   This article contains supporting information.

Acknowledgments

556   We thank members of the Chapman, Olofsson, and Ruotolo lab for their helpful suggestions and

557   discussions. We also thank Georg Meisl for advice on using the Amylofit program correctly.

Author Contributions

559   AB, DK, MC contributed conception and design of the study. AB, DK, YH, SJ, DL, LK, EG,

560   and RA performed the experiments. AO and BR provided materials and expertise. AB and DK

561   wrote the manuscript; AB, DK, and MC contributed to editing and revising the manuscript.

Funding and additional information

563   This work was supported by the National Institutes of Health grant no. R01 GM118651 and R21

564   AI137535 to M.R.C., R01 GM095832 to B.R., and NIH T32 AI007528 to DK.

Conflict of Interest

The authors declare that they have no conflicts of interest with the contents of this article.

Abbreviations

SPR, Surface Plasmon Resonance; SDS-PAGE, sodium dodecyl sulphate-polyacrylamide gel

electrophoresis; IM-MS, ion mobility-mass spectrometry; TOF, time-of-flight; ThT, Thioflavin-

T; EC, *E. coli*; CY, *Citrobacter youngae*; CD, *Citrobacter youngae*; HA, *Hafnia alvei*; MRE,

mean residual error; HFIP, hexafluoroisopropanol; CCS, collisional cross section; ATD, arrival

time distribution.

References

1.   Chiti, F., and Dobson, C. M. (2017) Protein Misfolding, Amyloid Formation, and Human Disease:
A Summary of Progress Over the Last Decade. *Annu Rev Biochem*. **86**, 27–68

2.   Chiti, F., and Dobson, C. M. (2006) Protein misfolding, functional amyloid, and human disease.
Annu Rev Biochem. **75**, 333–366

3.   Bucciantini, M., Giannoni, E., Chiti, F., Baroni, F., Taddei, N., Ramponi, G., Dobson, C. M., and
Stefani, M. (2002) Inherent toxicity of aggregates implies a common mechanism for protein
misfolding diseases. *Nature 2002 416:6880*. **416**, 507–511

4.   Haass, C., and Selkoe, D. J. (2007) Soluble protein oligomers in neurodegeneration: lessons from
the Alzheimer's amyloid β-peptide. *Nature Reviews Molecular Cell Biology 2007 8:2*. **8**, 101–112

5.   Balistreri, A., Goetzler, E., and Chapman, M. (2020) Functional Amyloids Are the Rule Rather
Than the Exception in Cellular Biology. *Microorganisms*. **8**, 1951

6.   Chapman, M. R., Robinson, L. S., Pinkner, J. S., Roth, R., Heuser, J., Hammar, M., Normark, S.,
and Hultgren, S. J. (2002) Role of Escherichia coli curli operons in directing amyloid fiber
formation. *Science (1979)*. **295**, 851–855

7.   Otzen, D. (2010) Functional amyloid: turning swords into plowshares. *Prion*. **4**, 256–64

8.   Deshmukh, M., Evans, M. L., and Chapman, M. R. (2018) Amyloid by Design: Intrinsic
Regulation of Microbial Amyloid Assembly. *J Mol Biol*. **430**, 3631–3641

9.   Nenninger, A. A., Robinson, L. S., Hammer, N. D., Epstein, E. A., Badtke, M. P., Hultgren, S. J.,
and Chapman, M. R. (2011) CsgE is a curli secretion specificity factor that prevents amyloid fibre
aggregation. *Mol Microbiol*. **81**, 486–99

10. Hammer, N. D., Schmidt, J. C., and Chapman, M. R. (2007) The curli nucleator protein, CsgB,
contains an amyloidogenic domain that directs CsgA polymerization. *Proc Natl Acad Sci U S A*.
104, 12494–12499

11. Nenninger, A. A., Robinson, L. S., and Hultgren, S. J. (2009) Localized and efficient curli
nucleation requires the chaperone-like amyloid assembly protein CsgF. *Proc Natl Acad Sci U S A*.
106, 900–905

12. Evans, M. L., Chorell, E., Taylor, J. D., Åden, J., Götheson, A., Li, F., Koch, M., Sefer, L.,
Matthews, S. J., Wittung-Stafshede, P., Almqvist, F., and Chapman, M. R. (2015) The Bacterial
Curli System Possesses a Potent and Selective Inhibitor of Amyloid Formation. *Mol Cell*. **57**, 445–
455

13. Taylor, J. D., Zhou, Y., Salgado, P. S., Patwardhan, A., McGuffie, M., Pape, T., Grabe, G.,
Ashman, E., Constable, S. C., Simpson, P. J., Lee, W., Cota, E., Chapman, M. R., and Matthews,
S. J. (2011) Atomic Resolution Insights into Curli Fiber Biogenesis. *Structure*. **19**, 1307–1316

14. Taylor, J. D., Hawthorne, W. J., Lo, J., Dear, A., Jain, N., Meisl, G., Andreasen, M., Fletcher, C.,
Koch, M., Darvill, N., Scull, N., Escalera-Maurer, A., Sefer, L., Wenman, R., Lambert, S., Jean, J.,
Xu, Y., Turner, B., Kazarian, S. G., Chapman, M. R., Bubeck, D., de Simone, A., Knowles, T. P.
J., and Matthews, S. J. (2016) Electrostatically-guided inhibition of Curli amyloid nucleation by
the CsgC-like family of chaperones. *Sci Rep*. **6**, 24656

15. Rochet, J. C., and Lansbury, P. T. (2000) Amyloid fibrillogenesis: themes and variations. *Curr
Opin Struct Biol*. **10**, 60–68

16. Van Gerven, N., Klein, R. D., Hultgren, S. J., and Remaut, H. (2015) Bacterial amyloid formation:
structural insights into curli biogensis. *Trends Microbiol*. **23**, 693–706

17. Evans, M. L., Chorell, E., Taylor, J. D., Åden, J., Götheson, A., Li, F., Koch, M., Sefer, L.,
Matthews, S. J., Wittung-Stafshede, P., Almqvist, F., and Chapman, M. R. (2015) The Bacterial
Curli System Possesses a Potent and Selective Inhibitor of Amyloid Formation. *Mol Cell*. **57**, 445–
455

18. Nagaraj, M., Najarzadeh, Z., Pansieri, J., Biverstål, H., Musteikyte, G., Smirnovas, V., Matthews,
S., Emanuelsson, C., Johansson, J., Buxbaum, J. N., Morozova-Roche, L., and Otzen, D. E. (2022)
Chaperones mainly suppress primary nucleation during formation of functional amyloid required
for bacterial biofilm formation. *Chem Sci*. **13**, 536

19. Zhou, Y., Smith, D., Leong, B. J., Brannstrom, K., Almqvist, F., and Chapman, M. R. (2012)
Promiscuous cross-seeding between bacterial amyloids promotes interspecies biofilms. *J Biol
Chem*. **287**, 35092–35103

20. Salgado, P. S., Taylor, J. D., Cota, E., and Matthews, S. J. (2011) Extending the usability of the
phasing power of diselenide bonds: SeCys SAD phasing of CsgC using a non-auxotrophic strain.
Acta Crystallogr D Biol Crystallogr. **67**, 8–13

21. Evans, M. L., Gichana, E., Zhou, Y., and Chapman, M. R. (2018) Bacterial amyloids. in *Methods
in Molecular Biology*, pp. 267–288, Humana Press Inc., **1779**, 267–288

22. Balistreri, A., Kahana, E., Janakiraman, S., and Chapman, M. R. (2020) Tuning Functional
Amyloid Formation Through Disulfide Engineering. *Front Microbiol*. 10.3389/fmicb.2020.00944

23. Brännström, K., Öhman, A., and Olofsson, A. (2011) Aβ peptide fibrillar architectures controlled
by conformational constraints of the monomer. *PLoS One*. 10.1371/journal.pone.0025157

24. Arosio, P., Knowles, T. P. J., and Linse, S. (2015) On the lag phase in amyloid fibril formation.
Physical Chemistry Chemical Physics. **17**, 7606–7618

25. Dueholm, M. S., Albertsen, M., Otzen, D., and Nielsen, P. H. (2012) Curli functional amyloid
systems are phylogenetically widespread and display large diversity in operon and protein
structure. *PLoS One*. **7**, e51274

26. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool,
K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J.,
Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D.,
Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver,
D., Vinyals, O., Senior, A. W., Kavukcuoglu, K., Kohli, P., and Hassabis, D. (2021) Highly
accurate protein structure prediction with AlphaFold. *Nature 2021 596:7873*. **596**, 583–589

27. Meisl, G., Kirkegaard, J. B., Arosio, P., Michaels, T. C. T., Vendruscolo, M., Dobson, C. M.,
Linse, S., and Knowles, T. P. J. (2016) Molecular mechanisms of protein aggregation from global
fitting of kinetic models. *Nature Protocols 2016 11:2*. **11**, 252–272

28. Collins, S. R., Douglass, A., Vale, R. D., and Weissman, J. S. (2004) Mechanism of prion
propagation: Amyloid growth occurs by monomer addition. *PLoS Biol*.
10.1371/JOURNAL.PBIO.0020321

29. Jahn, T. R., and Radford, S. E. (2008) Folding versus aggregation: Polypeptide conformations on
competing pathways. *Arch Biochem Biophys*. **469**, 100–117

30. Dueholm, M. S., Nielsen, S. B., Hein, K. L., Nissen, P., Chapman, M., Christiansen, G., Nielsen,
P. H., and Otzen, D. E. (2011) Fibrillation of the major curli subunit CsgA under a wide range of
conditions implies a robust design of aggregation. *Biochemistry*. **50**, 8281–8290

31. Wang, X., Smith, D. R., Jones, J. W., and Chapman, M. R. (2007) In vitro polymerization of a
functional Escherichia coli amyloid protein. *J Biol Chem*. **282**, 3713–3719

32. Homola, J. (2008) Surface plasmon resonance sensors for detection of chemical and biological
species. *Chem Rev*. **108**, 462–493

33. Tamara, S., Den Boer, M. A., and Heck, A. J. R. (2021) High-Resolution Native Mass
Spectrometry. *Chem Rev*.
10.1021/ACS.CHEMREV.1C00212/ASSET/IMAGES/LARGE/CR1C00212_0030.JPEG

34. May, J. C., and McLean, J. A. (2015) Ion mobility-mass spectrometry: Time-dispersive
instrumentation. *Anal Chem*. **87**, 1422–1436

35. Santambrogio, C., Natalello, A., Brocca, S., Ponzini, E., and Grandori, R. (2019) Conformational
Characterization and Classification of Intrinsically Disordered Proteins by Native Mass
Spectrometry and Charge-State Distribution Analysis. *Proteomics*. **19**, 1800060

36. Baumketner, A., Bernstein, S. L., Wyttenbach, T., Bitan, G., Teplow, D. B., Bowers, M. T., and
Shea, J.-E. (2006) Amyloid β-protein monomer structure: A computational and experimental
study. *Protein Sci*. **15**, 420

37. Hyung, S. J., Detoma, A. S., Brender, J. R., Lee, S., Vivekanandan, S., Kochi, A., Choi, J. S.,
Ramamoorthy, A., Ruotolo, B. T., and Lim, M. H. (2013) Insights into antiamyloidogenic
properties of the green tea extract (-)-epigallocatechin-3-gallate toward metal-associated amyloid-
β species. *Proc Natl Acad Sci U S A*. **110**, 3743–3748

38.   Soper, M. T., DeToma, A. S., Hyung, S. J., Lim, M. H., and Ruotolo, B. T. (2013) Amyloid-beta-
678   neuropeptide interactions assessed by ion mobility-mass spectrometry. *Phys Chem Chem Phys*. **15**,
679   8952–8961

680   39.   Sievers, S. A., Karanicolas, J., Chang, H. W., Zhao, A., Jiang, L., Zirafi, O., Stevens, J. T., Münch,
681   J., Baker, D., and Eisenberg, D. (2011) Structure-based design of non-natural amino-acid
682   inhibitors of amyloid fibril formation. *Nature 2011 475:7354*. **475**, 96–100

683   40.   Chan, P. H., Pardon, E., Menzer, L., De Genst, E., Kumita, J. R., Christodoulou, J., Saerens, D.,
684   Brans, A., Bouillenne, F., Archer, D. B., Robinson, C. V., Muyldermans, S., Matagne, A.,
685   Redfield, C., Wyns, L., Dobson, C. M., and Dumoulin, M. (2008) Engineering a camelid antibody
686   fragment that binds to the active site of human lysozyme and inhibits its conversion into amyloid
687   fibrils. *Biochemistry*. **47**, 11041–11054

688   41.   Giorgetti, S., Greco, C., Tortora, P., and Aprile, F. A. (2018) Targeting amyloid aggregation: An
689   overview of strategies and mechanisms. *Int J Mol Sci*. 10.3390/IJMS19092677

690   42.   Sleutel, M., Van den Broeck, I., Van Gerven, N., Feuillie, C., Jonckheere, W., Valotteau, C.,
691   Dufrêne, Y. F., and Remaut, Han. (2017) Nucleation and growth of a bacterial functional amyloid
692   at single-fiber resolution. *Nat Chem Biol*. **13**, 902–908

693   43.   Gharibyan, A. L., Jayaweera, S. W., Lehmann, M., Anan, I., and Olofsson, A. (2022) Endogenous
694   Human Proteins Interfering with Amyloid Formation. *Biomolecules 2022, Vol. 12, Page 446*. **12**,
695   446

696   44.   Arosio, P., Michaels, T. C. T., Linse, S., Månsson, C., Emanuelsson, C., Presto, J., Johansson, J.,
697   Vendruscolo, M., Dobson, C. M., and Knowles, T. P. J. (2016) Kinetic analysis reveals the
698   diversity of microscopic mechanisms through which molecular chaperones suppress amyloid
699   formation. *Nature Communications 2016 7:1*. **7**, 1–9

700   45.   Van Gerven, N., Klein, R. D., Hultgren, S. J., and Remaut, H. (2015) Bacterial amyloid formation:
701   structural insights into curli biogensis. *Trends Microbiol*. **23**, 693–706

702   46.   Andreasen, M., Meisl, G., Taylor, J. D., Michaels, T. C. T., Levin, A., Otzen, D. E., Chapman, M.
703   R., Dobson, C. M., Matthews, S. J., and Knowles, T. P. J. (2019) Physical determinants of amyloid
704   assembly in biofilm formation. *mBio*. 10.1128/MBIO.02279-18/SUPPL_FILE/MBIO.02279-18-
705   S0003.PDF

706   47.   Frieden, C. (2007) Protein aggregation processes: In search of the mechanism. *Protein Sci*. **16**,
707   2334

708   48.   Jain, S., and Udgaonkar, J. B. (2011) Prion protein aggregation. *Curr Sci*

709   49.   Olsen, W. P., Courtade, G., Peña-Díaz, S., Nagaraj, M., Sønderby, T. V., Mulder, F. A. A., Malle,
710   M. G., and Otzen, D. E. (2024) CsgA gatekeeper residues control nucleation but not stability of
711   functional amyloid. *Protein Science*. **33**, e5178

712   50.   Meisl, G., Xu, C. K., Taylor, J. D., Michaels, T. C. T., Levin, A., Otzen, D., Klenerman, D.,
713   Matthews, S., Linse, S., Andreasen, M., and Knowles, T. P. J. (2022) Uncovering the universality
714   of self-replication in protein aggregation and its link to disease. *Sci Adv*.
715   10.1126/SCIADV.ABN6831

51. Olsen, W. P., Christensen, J. L., Malle, M. G., and Otzen, D. E. (2025) SMARTQ: single-
molecule amyloid fibRil tracking and quantification. A method for accurately imaging, tracking,
and quantifying the growth of individual amyloid fibrils using TIRF. *The Three Functional States*
of Proteins: Structured, Intrinsically Disordered, and Phase Separated. 10.1016/B978-0-443-
21809-5.00006-5

52. Baker, N. A., Sept, D., Joseph, S., Holst, M. J., and McCammon, J. A. (2001) Electrostatics of
nanosystems: Application to microtubules and the ribosome. *Proceedings of the National*
Academy of Sciences. **98**, 10037–10041

53. Katoh, K., Kuma, K. I., Toh, H., and Miyata, T. (2005) MAFFT version 5: improvement in
accuracy of multiple sequence alignment. *Nucleic Acids Res*. **33**, 511–518

54. Livingstone, C. D., and Barton, G. J. (1993) Protein sequence alignments: a strategy for the
hierarchical analysis of residue conservation. *Bioinformatics*. **9**, 745–756

55. Casadaban, M. J., and Cohen, S. N. (1980) Analysis of gene control signals by DNA fusion and
cloning in Escherichia coli. *J Mol Biol*. **138**, 179–207