Electrical stimulation of the superior temporal gyrus evokes rapid responses in human visual cortex

Emily C. Cunningham[1] & David Brang[1]

[1]Department of Psychology
University of Michigan
530 Church St.
Ann Arbor, MI, 48109

Correspondence regarding this article should be addressed to Emily Cunningham at
ecngham@umich.edu

COI statement: The authors have no conflicts of interest to disclose.

Abstract

Sounds produce rapid responses in human visual cortex. Animal research suggests that these responses are enabled by corticocortical projections from auditory to visual cortex. To test for the presence of such pathways in humans, we examined intracranial responses to direct electrical stimulation. Stimulation over auditory cortex elicited rapid responses in early visual cortex (particularly near the visual periphery), providing causal support for corticocortical transmission of cross-modal information.

Early visual cortex is increasingly recognized as a site of rapid audiovisual interaction. For
example, sounds can modulate activity in the anterior portion of human V1 within 30ms of onset.[1]
Although rapid visual responses to auditory stimuli have been repeatedly demonstrated,[1–4] we know
relatively little regarding the pathways by which these signals are communicated in humans. Nonhuman
primate tracing studies point to several possibilities, including subcortical, corticothalamocortical, and
direct corticocortical transmission[5–10]. Of these candidates, the most interest/debate surrounds the
possibility that human visual cortex may receive direct anatomical inputs from auditory cortex.
The features of human sound-evoked visual responses are broadly consistent with the existence
of such pathways. Their distribution across visual cortex mirrors nonhuman primate anatomy, and their
speed of onset suggests rapid communication. Human structural connectivity also provides some support
for corticocortical communication.[11,12] However, these observations are indirect, and given the sparse
nature of the projections identified in macaques, the question of how these pathways extend to humans
remains unresolved (similar responses could be produced, for example, by tectopulvinar transmission[10]).
Here, we provide a causal evaluation of the hypothesis that rapid sound-evoked visual responses
originate in auditory cortex. Specifically, we examine how human visual cortex responds to single-pulse
electrical stimulation of regions in and around auditory cortex using intracranial EEG (iEEG) recordings
from patients with epilepsy (open data from Van Blooijs et al., 2023[13]). This technique provides an
estimate of effective connectivity between cortical areas through the distribution/latency of evoked
responses (corticocortical evoked potentials, or CCEPs[14]) following current passed between adjacent
intracranial contacts. CCEPs typically consist of a rapid early deflection (<50ms) followed by a
subsequent slower wave (~80-250ms; Figure 1A). The first-emerging peak in the signal ('N1') is thought
to result from corticocortical signal propagation,[14–16] and its distribution/timing may provide insight into the
presence of directional, corticocortical projections from auditory to visual cortex.
Occipital CCEPs following stimulation of the superior temporal gyrus (STG) were examined in 23
patients, with a focus on the presence and latency of N1 deflections occurring within 80ms of stimulation
(Figure 1A). Stimulation pairs were grouped according to HCP-MMP atlas location of the center of the pair
along the STG (Figure 1A). For summary analyses, recording sites were grouped by location within
occipital cortex (V1/V2, V3/V4, hMT+, and 'remaining occipital contacts' or 'other'). Two of these regions,
V1/V2 and hMT+ (anatomically defined), reflect hypothesized loci of early cross-modal input[5,6,17,18] and
are the subject of detailed examination; the remaining two provide points of comparison. For analyses of
response prevalence, null distributions were created from 1000 surrogate datasets based on time-series
drawn from pre-stimulation windows and processed identically to stimulation-evoked responses.
Overall, 112/231 occipital sites (across 19/23 patients) exhibited early responses to STG
stimulation (Figure 1B-C; p < .001 vs. response rate in permuted data). The proportion of responses
detected was greatest following stimulation of the posterior STG, and declined to near chance with
increasingly anterior stimulation (Figure 1B). Within V1/V2, response rates were greatest following mid-
STG stimulation (i.e., when stimulation was centered over likely auditory association areas accessible to
contacts placed on the outer surface of mid-STG), declining with more anterior/posterior stimulation
(Figure 1D; note that only the reduction in response rate between A4/PBelt and Posterior STG was robust
to FDR correction), an effect which appears largely driven by responses in anterior portions of the
calcarine sulcus and lingual gyrus (Figure 1C). Within lateral occipital regions near hMT+, the proportion
of detected responses per patient matched the overall trend, increasing with proximity as stimulation
moved from anterior (6%) to posterior STG (26%; Figure 1D). Over the remaining recording sites,
responses were often detected at levels above chance, but did not show clear differentiation by
stimulation location (Figure 1D).
In nonhuman primates, auditory projections preferentially target anterior regions of V1
corresponding to representation of the peripheral visual field[5,6]. In this dataset, electrode coverage of
multiple locations along V1 provided a rare opportunity to explore the presence of a similar bias in the
human CCEP response (Figure 2). Responses were observed in the most anterior portion of the calcarine

sulcus (with an average response rate of 33% across 3 patients with anterior calcarine sites; Figure 2A),
and to some extent the occipital pole (9%; below the corresponding chance threshold, Figure 2A). There
were no detectable early responses in mid-calcarine locations typically associated with parafoveal
processing (Figure 2A). The latency of early anterior visual responses ranged from 18-61ms, with 5/7
peaks emerging within 30ms of stimulation (β = 33 ms, 95% CI [22, 45]; Figure 2B). The latency of
occipital polar responses ranged from 34-76ms (β = 45 ms, 95% CI [29, 62]). The spectral representation
of CCEPs across V1/V2 sites (Figure 2D) was characterized by transient low-frequency activity together
with no detectable change in high-gamma power (Figure 2C). This pattern is consistent with modulatory
effects produced by feedback-type signal propagation, with no evidence of stimulation-elicited local
neuronal spiking typically indexed by increased high-frequency activity;[19,20] It maps well to the anatomy of
primate corticocortical projections, which also have a feedback/modulatory character[21].
Patterns of sound-evoked activity[2] and human/nonhuman primate connectivity[11,17,18] also highlight
V5/hMT+ as a likely site of early cross-modal input, particularly from auditory motion-sensitive areas
along the planum temporale. This is consistent with our general observation of posterior-dominant STG
responses in the broad hMT+ ROI (Figure 1D). To clarify the specificity of responses within this region
(which was covered in n = 22 patients), we examined whether the relation between posterior STG
stimulation and hMT+ activation persisted in the most highly probable hMT+ sites (Figure 3; Figure S3
displays surrounding sites). Notably, the response to posterior STG stimulation was maintained in these
sites, but responses to stimulation from other locations along the STG were comparatively limited (i.e.,
responses in probable hMT+ showed some selectivity to posterior STG; Figure 3A). The latency of
probable hMT+ responses ranged from 18-74ms (β = 44 ms, 95% CI [33 54]; Figure 3B). The spectral
content of CCEPs in the greater hMT+ area was, as with V1/V2, largely dominated by low-frequency
activity, with little evidence of activity in the high-gamma range (Figure 3C).
Overall, these results provide causal support for the presence of direct pathways from auditory to
visual cortex in humans. Early visual cortex responded rapidly (18ms+) to electrical stimulation over mid-
and posterior STG, with a distribution that largely matched patterns of connectivity observed in humans
(using non-invasive methods) and nonhuman primates. Although functional localizers were not available
for these patients, the distribution of stimulation-evoked responses also corresponded well to reports of
sound-evoked responses in anterior V1[1,3], and reports of auditory motion-related activation of V5/hMT+
and adjacent cortex[22]. STG stimulation also evoked possible responses over the occipital pole. These
responses, which were weaker than those over anterior calcarine (see Figure 2A, subject A), have limited
support in nonhuman primate data[5], but are consistent with human DTI[12] and cross-modal effects
reported at the occipital pole[23,24], and may warrant additional investigation. This profile would be
additionally strengthened with investigation of the effects of stimulation over contacts adjacent to A1 and
auditory belt/parabelt regions where many of the visual projections identified in macaques originate[5,6,21].
These results indicate that the rapid response to sound in human visual cortex can occur through
corticocortical as opposed to solely subcortical transmission. Functionally, the observation that STG
stimulation produced low- but not high-frequency changes in visual activity is also consistent with the
suggestion that sounds modulate visual cortical excitability without eliciting action potentials[4,23,25].
However, these data do not rule out thalamocortical and corticothalamocortical contributions to the
sound-evoked visual response. Respective contributions of these pathways remain an open question.
Although the current dataset contained only ipsilateral stimulation/recording combinations, we expect that
early cross-modal corticocortical pathways are restricted to the same hemisphere, given animal work and
prior research in humans suggesting primacy for spatially-aligned auditory-visual stimuli[2,26,27].
Early cross-modal interactions mediated by corticocortical projections not only support visual
processing in the sighted, but also provide avenues for rapid cortical reorganization following vision loss.
The use of CCEPs to examine effective connectivity may provide insight into patterns of plasticity
following disease/injury by illustrating where cross-modal corticocortical inputs in the sighted are most
prevalent. This is likely most relevant for the late blind, but our results also bear similarities to sound-

evoked responses in the occipital cortex of early or congenitally blind individuals. For example, patterns of
auditory scene decoding in the early visual cortex of congenitally blind participants also show a
pronounced eccentricity bias (present to a lesser extent in sighted subjects)[28], and early blindness is
associated with enhanced recruitment of visual motion-sensitive cortex in response to auditory
motion[29,30]. Although the current data was obtained from patients with epilepsy, the compatibility of our
observations with reports of sound-evoked responses in other patients[1–4], connectivity in nonpatient
samples[11,12], and tractography in nonhuman primates[6,17,18,21] provides a measure of confidence in the
results.
201
202

203
204
Figure 1*. Effects of stimulation over STG on responses in occipital cortex. (A) Upper: Illustration of a*
CCEP response with prominent early negative deflection (N1). Lower: superior temporal and occipital
regions of interest. Stimulation locations with limited representation were combined with neighboring
locations to create four summary regions: posterior STG, PBelt/A4, A5, and anterior STG (B) Global
CCEP response to STG stimulation (center). Black line represents averaged CCEP across all sites &
patients (averaged within then across patients); All waveforms with detected responses were rectified

prior to averaging. Grey ribbon indicates patient-level 95% CI. Black bars at the bottom highlight
timepoints at which the observed mean exceeded the 95th percentile of a matched permutation null
distribution (FDR-corrected, Benjamini & Hochberg). (outer) all occipital CCEP waveforms split by
stimulation location along the STG. Black lines above each plot indicate timepoints at which early peaks
were identified. Inset plots: diamonds indicate the average proportion of early responses detected per
patient, and grey point-clouds indicate the corresponding proportion of false-positive peaks identified in
each of 1000 iterations of shuffled surrogate data; Black lines indicate the 99th percentile of this null
distribution. Proportions and surrogate p-values are: Anterior STG: p(resp) = .11, p < .001; A5: p(resp) =
.11, p = .002; A4/PBelt: p(resp) = .14, p < .001; Posterior STG: p(resp) = .25, p < .001 (C) Distribution of
responsive electrodes as a function of stimulation location along the STG (HCP-MMP atlas). RH
electrodes were translated to LH coordinates for visualization. Stimulation pairs within each region are
plotted in black. Color indicates latency of the earliest detected peak for a given electrode. Electrodes
with no detectable response in the first 80ms are grey. Electrode size is scaled by the effect size of the
associated peak (Cohen's d_z). Bar plots indicate the number of patients with at least one responsive
electrode (red) relative to the total number of patients (grey) for a given combination of stimulation
location and recording region. (D) Average proportion of responses detected per patient (diamonds
indicate average; grey dots indicate single patient data) by stimulation and recording region (relative to
the total number of CCEP waveforms for a given patient and grouping; see also Figure S4). Error bars
are SEM. White diamonds indicate an observed proportion less than the 99th percentile of the associated
surrogate null distribution. Black horizontal bars indicate effects of stimulation location on response
proportion (after FDR correction across all 24 pairwise combinations). Effects of stimulation location were
evaluated using linear mixed-effects models with patient as a random effect; surrogate p-values were
computed as the proportion of fixed effects from models fit to null data ≥ the observed fixed effect (V1/V2:
PostSTG-A4/PBelt, p = .004, all others n.s.; hMT+: PostSTG-A5, p = .004, PostSTG-AntSTG, p = .002, all
others n.s.; V3/V4: PostSTG-A4/PBelt, p = .002, all others n.s.; Other: all n.s.).

fixed effects (Wald). The area +/- half of the standard deviation of the gaussian window of the longest
wavelet is masked (note: over this window, interpolation of the stimulation artifact artificially reduces high-
gamma power). Lower: Heat maps depict time-frequency decomposition (2-40 Hz) via Morlet wavelets.
Color ('a.u.') indicates fixed intercept estimates from mixed-effects models of single-trial power for each
time/frequency point, accounting for patient- and electrode-level variance. Faded central bars indicate the
window over which the stimulation artifact was interpolated prior to time-frequency decomposition (-5 to
12ms). For illustrative purposes, black lines indicate FDR-corrected regions showing significant difference
from baseline. The curve of the line following stimulation onset highlights the standard deviation of the
gaussian window for the wavelet at each frequency (i.e., the region of potential interference from the
stimulation artifact, over which statistics were not evaluated).

261
Figure 3. *Examination of responses in the most probable MT/MST sites. (A) All CCEP waveforms from*
probable hMT+ sites. Electrode locations within the greater ROI are highlighted in the upper left, with
inset plot indicating probability of membership. Each subplot contains data for a separate patient
(identified by number). Bold lines highlight waveforms for which early responses were detected. Color of
the waveform indicates stimulation location along the STG. Lower right: Average proportion of responses
detected per patient for each stimulation location (diamonds) plotted against the associated surrogate
null/chance distribution (grey point clouds). Number of patients contributing data for each stimulation
location is indicated above the plot. Black bars indicate 99th percentile of the null distribution. (B) Peak

latencies for all responsive sites in probable hMT+ colored according to stimulation location. Relative size of each point is scaled by effect size at the peak (Cohen's dz; range: 0.23-6.07). Linear mixed-effects models with random effects of subject were applied to derive estimates of peak latency (main text). (C) Spectral representations of all CCEP waveforms from the broad hMT+ roi. Upper: high gamma (70-150 Hz) activity time-series. Y-axis ('a.u.') indicates fixed intercept estimates from mixed-effects models of single-trial power for each timepoint, accounting for patient- and electrode-level variance. Error ribbons indicate 95% CI surrounding fixed effects (Wald). The area +/- half of the standard deviation of the gaussian window of the longest wavelet is masked (note: over this window, interpolation of the stimulation artifact artificially reduces high-gamma power). Lower: Heat maps depict time-frequency decomposition (2-40 Hz) via Morlet wavelets. Color ('a.u.') indicates fixed intercept estimates from mixed-effects models of single-trial power for each time/frequency point, accounting for patient- and electrode-level variance. Faded central bars indicate the window over which the stimulation artifact was interpolated prior to time-frequency decomposition (-5 to 12ms). For illustrative purposes, black lines indicate FDR-corrected regions showing significant difference from baseline. The curve of the line following stimulation onset highlights the standard deviation of the gaussian window for the wavelet at each frequency (i.e., the region of potential interference from the stimulation artifact, over which statistics were not evaluated).

References

1. Brang D, Towle VL, Suzuki S, Hillyard SA, Di Tusa S, Dai Z, Tao J, Wu S, Grabowecky M.
Peripheral sounds rapidly activate visual cortex: evidence from electrocorticography. Journal of
Neurophysiology. 2015 Nov 1;114(5):3023–3028.

2. Plass J, Ahn E, Towle VL, Stacey WC, Wasade VS, Tao J, Wu S, Issa NP, Brang D. Joint encoding
of auditory timing and location in visual cortex. J Cogn Neurosci. 2019 Jul;31(7):1002–1017.
PMCID: PMC6551606

3. Ferraro S, Van Ackeren MJ, Mai R, Tassi L, Cardinale F, Nigri A, Bruzzone MG, D'Incerti L,
Hartmann T, Weisz N, Collignon O. Stereotactic electroencephalography in humans reveals
multisensory signal in early visual and auditory cortices. Cortex. 2020 May 1;126:253–264.

4. Mercier MR, Foxe JJ, Fiebelkorn IC, Butler JS, Schwartz TH, Molholm S. Auditory-driven phase
reset in visual cortex: Human electrocorticography reveals mechanisms of early multisensory
integration. NeuroImage. 2013 Oct 1;79:19–29.

5. Falchier A, Clavagnier S, Barone P, Kennedy H. Anatomical Evidence of Multimodal Integration in
Primate Striate Cortex. J Neurosci. Society for Neuroscience; 2002 Jul 1;22(13):5749–5759. PMID:
12097528

6. Rockland KS, Ojima H. Multisensory convergence in calcarine visual areas in macaque monkey.
International Journal of Psychophysiology. 2003 Oct 1;50(1):19–26.

7. Borra E, Rockland KS. Projections to Early Visual Areas V1 and V2 in the Calcarine Fissure from
Parietal Association Areas in the Macaque. Front Neuroanat. Frontiers; 2011 Jun 22

8. Clavagnier S, Falchier A, Kennedy H. Long-distance feedback projections to area V1: Implications
for multisensory integration, spatial awareness, and visual consciousness. Cognitive, Affective, &
Behavioral Neuroscience. 2004 Jun 1;4(2):117–126.

9. Cappe C, Rouiller EM, Barone P. Multisensory anatomical pathways. Hearing Research. 2009 Dec
1;258(1):28–36.

10. Lyon DC, Nassi JJ, Callaway EM. A Disynaptic Relay from Superior Colliculus to Dorsal Stream
Visual Cortex in Macaque Monkey. Neuron. 2010 Jan 28;65(2):270–279.

11. Gurtubay-Antolin A, Battal C, Maffei C, Rezk M, Mattioni S, Jovicich J, Collignon O. Direct Structural
Connections between Auditory and Visual Motion-Selective Regions in Humans. J Neurosci.
Society for Neuroscience; 2021 Mar 17;41(11):2393–2405. PMID: 33514674

12. Beer AL, Plank T, Greenlee MW. Diffusion tensor imaging shows white matter tracts between
human auditory and visual cortex. Exp Brain Res. 2011 Sep 1;213(2):299–308.

13. van Blooijs D, van den Boom MA, van der Aar JF, Huiskamp GM, Castegnaro G, Demuru M,
Zweiphenning WJEM, van Eijsden P, Miller KJ, Leijten FSS, Hermes D. Developmental trajectory of
transmission speed in the human brain. Nat Neurosci. Nature Publishing Group; 2023
Apr;26(4):537–541.

14. Matsumoto R, Nair DR, LaPresto E, Najm I, Bingaman W, Shibasaki H, Lüders HO. Functional
connectivity in the human language system: a cortico-cortical evoked potential study. Brain. 2004
Oct;127(10):2316–2330.

15.   Conner CR, Ellmore TM, DiSano MA, Pieters TA, Potter AW, Tandon N. Anatomic and electro-
physiologic connectivity of the language system: A combined DTI-CCEP study. Computers in
Biology and Medicine. 2011 Dec 1;41(12):1100–1109.

16.   Matsumoto R, Kunieda T, Nair D. Single pulse electrical stimulation to probe functional and
pathological connectivity in epilepsy. Seizure. 2017 Jan 1;44:27–36.

17.   Palmer SM, Rosa MGP. Quantitative Analysis of the Corticocortical Projections to the Middle
Temporal Area in the Marmoset Monkey: Evolutionary and Functional Implications. Cerebral Cortex.
2006 Sep 1;16(9):1361–1375.

18.   Majka P, Rosa MGP, Bai S, Chan JM, Huo BX, Jermakow N, Lin MK, Takahashi YS, Wolkowicz IH,
Worthy KH, Rajan R, Reser DH, Wójcik DK, Okano H, Mitra PP. Unidirectional monosynaptic
connections from auditory areas to the primary visual cortex in the marmoset monkey. Brain Struct
Funct. 2019 Jan 1;224(1):111–131.

19.   Bastos AM, Vezoli J, Bosman CA, Schoffelen JM, Oostenveld R, Dowdall JR, De Weerd P,
Kennedy H, Fries P. Visual Areas Exert Feedforward and Feedback Influences through Distinct
Frequency Channels. Neuron. Elsevier; 2015 Jan 21;85(2):390–401. PMID: 25556836

20.   Buffalo EA, Fries P, Landman R, Buschman TJ, Desimone R. Laminar differences in gamma and
alpha coherence in the ventral stream. Proceedings of the National Academy of Sciences.
Proceedings of the National Academy of Sciences; 2011 Jul 5;108(27):11262–11267.

21.   Falchier A, Cappe C, Barone P, Schroeder CE. Sensory Convergence in Low-Level Cortices. In:
Stein BE, editor. The New Handbook of Multisensory Processing. The MIT Press; 2012

22.   Rezk M, Cattoir S, Battal C, Occelli V, Mattioni S, Collignon O. Shared Representation of Visual and
Auditory Motion Directions in the Human Middle-Temporal Cortex. Current Biology. Elsevier; 2020
Jun 22;30(12):2289-2299.e8. PMID: 32442465

23.   Romei V, Murray MM, Cappe C, Thut G. Preperceptual and Stimulus-Selective Enhancement of
Low-Level Human Visual Cortex Excitability by Sounds. Current Biology. 2009 Nov 17;19(21):1799–
1805.

24.   Romei V, Murray MM, Merabet LB, Thut G. Occipital Transcranial Magnetic Stimulation Has
Opposing Effects on Visual and Auditory Stimulus Detection: Implications for Multisensory
Interactions. J Neurosci. Society for Neuroscience; 2007 Oct 24;27(43):11465–11472. PMID:
17959789

25.   Iurilli G, Ghezzi D, Olcese U, Lassi G, Nazzaro C, Tonini R, Tucci V, Benfenati F, Medini P. Sound-
Driven Synaptic Inhibition in Primary Visual Cortex. Neuron. Elsevier; 2012 Feb 23;73(4):814–828.
PMID: 22365553

26.   Wallace MT, Wilkinson LK, Stein BE. Representation and integration of multiple sensory inputs in
primate superior colliculus. Journal of Neurophysiology. American Physiological Society Bethesda,
MD; 1996

27.   McDonald JJ, Teder-Sälejärvi WA, Hillyard SA. Involuntary orienting to sound improves visual
perception. Nature. Nature Publishing Group; 2000 Oct;407(6806):906–908.

28.   Vetter P, Bola Ł, Reich L, Bennett M, Muckli L, Amedi A. Decoding Natural Sounds in Early "Visual"
Cortex of Congenitally Blind Individuals. Current Biology. Elsevier; 2020 Aug 3;30(15):3039-
3044.e2. PMID: 32559449

29.   Poirier C, Collignon O, Scheiber C, Renier L, Vanlierde A, Tranduy D, Veraart C, De Volder AG.
Auditory motion perception activates visual motion areas in early blind subjects. NeuroImage. 2006
May 15;31(1):279–285.

30.   Strnad L, Peelen MV, Bedny M, Caramazza A. Multivoxel Pattern Analysis Reveals Auditory Motion
Information in MT+ of Both Congenitally Blind and Sighted Individuals. PLOS ONE. Public Library of
Science; 2013 Apr 30;8(4):e63198.

Method

Participants
Data were drawn from the open repository associated with van Blooijs et al., 2023. A subset (N = 23; mean age = 25 years, range 8-51) of the original lifespan sample of 74 participants had combinations of STG stimulation and occipital recording sites that were relevant for the current question (see Exclusion/Inclusion criteria below), but which were not considered in the initial paper. In all cases, sites were positioned ipsilaterally, with the majority on the left hemisphere (17/23; Figure S1).

Exclusion/Inclusion criteria. Data were screened to identify patients with some combination of STG and occipital electrode coverage, defined in terms of Destrieux atlas labels from individual cortical parcellations (Destrieux et al., 2010). Patients with relevant coverage were included if they had (a) at least one stimulation pair in which both electrodes were located on the STG, and (b) at least 10 trials for analysis for at least one relevant stimulation pair.

Data Acquisition
For details of data acquisition, readers are referred to van Blooijs et al. (2023). In brief, single-pulse electrical stimulation was conducted as iEEG data were recorded from subdural electrode strips/grids with 1 cm spacing. Data was initially acquired at a sampling rate of between 512-2048 Hz. For a given pair of adjacent electrodes, stimulation (pulse width: 1ms, current intensity: 8mA) was applied up to 10 times at intervals of 5s.

Processing & Analysis
Selection of stimulation pairs and recording sites. For patients meeting inclusion criteria, stimulation pairs were included in the following analyses if both members of a pair were located over the STG, or if one member was located on the STG and the other was located on the inferior portion of the supramarginal gyrus. The region 'PSL' from the Glasser (2016) parcellation was used to define the upper border for inclusion of supramarginal electrodes. Pairs that crossed the superior temporal sulcus or lateral fissure were not included. Stimulation pairs were also excluded if they contained any channels marked for exclusion by the criteria described in van Blooijs et al. (2023), or if the total number of trials for a pair was less than 10. All trials for a given stimulation pair were included, except in rare instances in which stimulation occurred within 3s of another trial. Occipital recording sites were included if their locations fell within the set of Destrieux atlas labels {G&S_occipital_inf, G_cuneus, G_oc-temp_med-Lingual, G_occipital_middle, G_occipital_sup, Pole_occipital, S_calcarine, S_oc_middle&Lunatus, S_oc_sup&transversal, S_occipital_ant, S_parieto_occipital}. Additional exclusion criteria based on data quality are described below.

Preprocessing and measurement of evoked responses. As in Van Blooijs et al (2023), no modification of the continuous signal was performed (i.e., data were neither rereferenced nor filtered). For measurement of evoked responses, timepoints between -12 and 12ms were excluded from analysis to limit contamination from stimulation artifacts. CCEPs were evaluated relative to a prestimulus average baseline of [-200 ms to -12 ms]. For peak latency measurement, the averaged ERP for a given combination of electrode site and stimulation pair was transformed by z-scoring the waveform relative to the mean and standard deviation over the baseline window (after the approach employed by Trebaul et al., 2018; Lemarechal et al., 2022). Peaks were identified as the first minimum/maximum in the interval [12 to 250ms] with an absolute value exceeding $z = 5$ and a minimum peak prominence of 2 (i.e. peaks were required to have a drop of $z = 2$ on either side before encountering a neighboring peak of greater amplitude). Electrodes were considered to show an early response if the CCEP waveform contained a detectable peak in the range 12-80ms following stimulation. Note that the use of an 80ms window risks

497 inclusion of some peaks later than the N1 component (which typically peaks in the first 50ms). However,
498 extending the range of inquiry to 80ms ensures more complete description of early responses.
499
Saturated responses (defined by post-stimulation amplitudes exceeding 2000 microvolts) were excluded
from analysis (n = 3 electrodes, 1 patient), as were responses that showed extreme time-locked line-
noise contamination in the averaged waveform (defined as having a mean power spectral density over
the range 50-60 Hz, prior to stimulation, that was greater than 10x the median value over all stimulation
pairs and electrode sites; n = 62 responses over 8 patients; this exclusion criterion was data-driven and
implemented to reduce the occurrence of false-positive 'peaks' resulting from line noise). Overall, after
these data quality exclusions, 1305 combinations of stimulation pair and recording site (231 total
recording sites) were considered for analysis. Included stimulation pairs and recording sites for each
patient are presented in Figure S1.
509
Given that the z-scored waveforms used to compute peak latency are scored relative to variation in the
averaged waveform over the baseline window, the z-scores alone contain no information regarding
variance in the waveform from trial to trial. To provide a coarse index of stability/variation in amplitude
across repeated stimulations, peak latency values were supplemented with Cohen's d_z values computed
for peak amplitude (computed by dividing peak amplitude by the standard deviation of the waveform
across trials at that timepoint).
516
Two permutation approaches were adopted to provide chance baselines against which to compare the
observed data. First, for time series evaluation, the following procedure was conducted to provide a
simple summary assessment and visualization of the occipital response to STG stimulation. First, 1000
null datasets were generated using a sign-based permutation approach, in which the values for half of the
trials associated with each CCEP waveform were randomly inverted prior to averaging. For each null
dataset, CCEP waveforms were generated and peaks evaluated exactly as in the observed data. Then,
for both the null and observed data, to account for the fact that CCEP polarity can vary by stimulation pair
and recording site, all waveforms for which a negative early peak was detected were inverted to place all
peaks (for both null and comparison distributions) in a consistent direction. All waveforms across all
electrodes (regardless of whether a response was detected) were then averaged within-subject and
across subjects to generate a summary CCEP reflecting the entire occipital ROI. Surrogate p-values were
derived from comparisons of the observed deviation from baseline and the associated permutation
distribution at a given timepoint (FDR-corrected, Benjamini & Hochberg, 1995).
530
Second, to generate a chance baseline that best reflects the expected false-positive rate in the absence
of stimulation, we generated a comparison distribution out of 1000 repetitions of the following: For each
CCEP time-series, we extracted a matched null time-series centered around a point within the 2 seconds
prior to that trial (selected randomly for each repetition). These null time-series were then processed in
the same way as the observed data described above, resulting in 1000 surrogate datasets identical to the
main dataset except for the use of null data. In this way, it was possible to evaluate expected false
positive rates assuming no effect of stimulation for every observed statistic. Observed values were
evaluated against this null distribution, and surrogate p-values were computed as the proportion of null
values matching or exceeding the magnitude of the observed statistic. The 99th percentile of the null
distribution was used as a benchmark for evaluating response rates in a given region.
541
542 ***Summary statistics by stimulation location and region.*** Stimulation pairs were grouped by the HCP-
543 MMP (Glasser et al., 2016) parcellation label associated with the nearest coordinate in Freesurfer
544 average space to the center of the pair. Stimulation locations with limited representation were combined

545 with neighboring locations to create four summary regions: posterior STG, PBelt/A4, A5, and anterior
546 STG (see Figure S2 for a full breakdown of the distribution of responses for all STG labels).
547
548 For summary analyses over visual locations, we defined sites near probable V5/hMT+ as those where
549 nearest coordinates in Freesurfer fsaverage space were positioned within the bounds of a mask drawn by
550 applying a >5% threshold to the probabilistic atlas developed by Huang et al., 2019 (version 'thr3_snrH').
551 Remaining locations in and around V1/V2 and V3/V4 were defined by proximity of nearest coordinates in
552 Freesurfer average space to HCP-MMP (Glasser et al., 2016) parcellation labels (Figure 1).
553
For comparisons of response rates (N1 prevalence) by stimulation location within each visual location,
linear mixed-effects models (maximum likelihood estimation) were generated for each pair of stimulation
locations within a region using the lme4 package in R (Bates et al., 2015), with random intercepts for
patient ID to address situations in which the same patient received stimulation from multiple STG
locations. The fixed effect of stimulation location was compared against the fixed effects derived from
identical models fit to the null data. The surrogate p-values obtained from these comparisons (proportion
of null responses matching or exceeding the magnitude of the observed fixed effect) were corrected
across the full set of 24 comparisons (6 comparisons within each of four visual ROIs; FDR, Benjamini &
Hochberg, 1995).
563
564 **_Exploratory analyses of V1 and hMT+._** To explore responses within early visual cortex as a function of
565 eccentricity, V1 sites were divided into anterior/middle/posterior regions, split based on fsaverage y-
566 coordinate (divided at -94 and -80, chosen by visual inspection). To explore specificity of hMT+
567 responses, the initial hMT+ ROI was reduced to highlight only those locations with maximum probability
568 MT/MST labels on the HCP-MMP atlas (Glasser et al., 2016).
569
570 To summarize peak latency within these locations, accounting for variance in number of responsive sites
571 across patients, linear mixed-effects models with random effects of patient identity were generated for
572 each region. These simple models contained the random effects and a fixed intercept, the value of which
573 was used as an estimate of the population mean peak latency (Bates et al., 2015). Confidence intervals
574 around the fixed effect (95%) computed using the Wald method provide estimates of uncertainty.
575
Spectral analyses. For the detailed descriptions of V1/V2 and hMT+ responses, spectral decomposition
was performed to characterize the frequency content of the signal over all CCEPs in the region. The
intent of this decomposition was descriptive. Trials were epoched from -3 to 3s to allow a sufficient
window for event-related spectral perturbation (ERSP) computation. The stimulation artifact was removed
by interpolating the time-series between -5 and 12 ms for each trial with piecewise cubic hermite
interpolation. Single-trial spectrograms were computed by convolving the iEEG time series with complex
Morlet wavelets (for lower frequencies: 39 center frequencies linearly spaced between 2-40 Hz, with
number of cycles per frequency increasing linearly from 3 to 12; for high-gamma power: 16 center
frequencies linearly spaced between 75 and 150 Hz, with number of cycles per frequency increasing
linearly from 12 to 24). A time step of ~4ms was applied to produce time-frequency plots sampled at 256
Hz. Following convolution, power was extracted as squared amplitude for each trial, timepoint, and
frequency. Spectrograms were baseline-corrected and normalized using the 'vssum' approach: for each
timepoint the average power over time and trials in the interval [-500 to -200ms] at that frequency was
subtracted, and values were then normalized by dividing by the sum of the value at that timepoint and the
associated baseline value. The resulting values are bounded between -1 and 1 and scaled such that +/-
0.33 indicates a 2x increase/decrease in power from baseline. High-gamma power values were averaged
across all frequencies to produce a single time-series representation of high-frequency activity.
593

Single-trial spectral information was modeled with a mixed-effects approach to generate fixed-effect
estimates of power relative to baseline (used to create time-frequency visualizations), and associated
confidence intervals (used in representations of high-gamma time series) for each time/frequency point.
For these models (with trial nested within electrode nested within subject), the random effects structure
included crossed random effects of electrode, run (for patients with multiple runs), and stimulation pair.
Confidence intervals around the fixed effect (95%) were computed using the Wald method for high-
gamma time-series. For time-frequency maps, to visualize uncertainty, p-values extracted using lmerTest
were evaluated over the post-stimulation interval (FDR corrected, Benjamini-Hochberg). To minimize
contamination by the interpolated stimulation artifact, these analyses were conducted over the interval
starting at the end of the interpolation window (12ms) plus half the width of the standard deviation of the
gaussian window applied at a given frequency. Note that interpolation of the stimulation artifact produces
an artificial reduction in high-gamma power in the peristimulus interval.
612