High rate of mutation and efficient removal by selection of structural variants from natural

populations of *Caenorhabditis elegans*

Ayush Shekhar Saxena[1,2*] and Charles F. Baer[1,3]

1 – Department of Biology, University of Florida, Gainesville, FL, USA

2 – Present address – Regeneron Pharmaceuticals Inc., Tarrytown, NY, USA

3 – University of Florida Genetics Institute, Gainesville, FL, USA

*Correspondence to:

Ayush Shekhar Saxena

ayushsaxena.iitg@gmail.com

Abstract

The importance of genomic structural variants (SVs) is well-appreciated, but much less is known about their mutational properties than of single nucleotide variants (SNVs) and short indels.  The reason is simple: the longer the variant, the less likely it will be covered by a single sequencing read, thus the harder it is to map unambiguously to a unique genomic location.

Here we report SV mutation rate estimates from six mutation accumulation (MA) lines from two strains of *C. elegans* (N2 and PB306) using long-read (PacBio) sequencing.  The inferred SV mutation rate is ~1/10 the SNV rate and ~1/4 the short indel rate.  We identified 40 mutations and 52 false positive (FP) variants by manual inspection of each SV call.  Excluding one atypical line (5 mutations, 35 FPs), the signal (mutant) to noise (FP) ratio is approximately 2:1.  False negative rates were determined by simulating variants in the reference genome and observing 'recall'.  Recall rate ranges from >90% for short indels and declines as SV length increases.  Small deletions have nearly the same recall rate as small insertions (~100bp), but deletions have higher recall rates than insertions as size increases.  The reported SV mutation rate is likely a lower bound.

A quarter of identified SV mutations occur in SV hotspots that harbor pre-existing low complexity repeat variation.  Comparison of the spectrum of spontaneous SVs to wild isolates implies that natural selection is not only efficient at removing SVs in exons but also removes roughly half of SVs in intergenic regions.

**Introduction**

Based on the total number of bases affected, structural variants (SV) — large insertions and deletions, repeat expansions and contractions, inversions, and translocations — are the largest source of genetic diversity in the genomes of multicellular organisms, and they have the largest average phenotypic and fitness effects of any type of mutation (Collins et al. 2020; Beyter et al. 2021). SVs are associated with a host of human disorders, including autism (Marshall et al. 2008; Weiss et al. 2008; Pinto et al. 2010; Sanders et al. 2011), schizophrenia (Walsh et al. 2008; Karayiorgou et al. 2010), and several types of cancers (Friedman et al. 1994; Wooster et al. 1995; Meyerson and Pellman 2011; Stransky et al. 2014; Nattestad et al. 2018). The functional consequences of SVs are well-documented (Hurles et al. 2008; Conrad et al. 2010; Yi and Ju 2018; Loegler et al. 2025).

In recent years, human population geneticists have converged on the idea that the genetic basis of complex disease can be largely accounted for by the combined influence of very rare, recently arisen variants of large effect (Wang et al. 2021) and variants of small effect at very many loci (the "omnigenome", Boyle et al. 2017). Both classes of variants imply a significant role for mutation, the former for obvious reasons and the latter because, even if the per-locus mutation rate is low, the cumulative genetic variance contributed by a large fraction of the genome at mutation-selection-drift balance may be substantial.

Given the outsized contribution of SVs to genetic diversity on the one hand, and the apparently important contribution of mutation to complex disease on the other, understanding the mutational properties of SVs is an important problem in biomedical genetics. More broadly, characterizing SV mutation is necessary for an unbiased estimate of the distribution of fitness effects (DFE) of new mutations, which is a longstanding goal of evolutionary genetics (Fisher 1922; Eyre-Walker and Keightley 2007). Nearly all estimates of the DFE are restricted to the effects of SNPs and small indels (and most are restricted to SNPs, e.g., Kousathanas and Keightley 2013; Kim et al. 2017; Tataru et al. 2017; Böndel et al. 2019; Gilbert et al. 2021; James et al. 2023; Crombie et al. 2024). Because SVs are

ignored, all the variation in fitness is ascribed to small indels and SNPs, artificially inflating

their importance by an unknown amount that may be quite large.

The reason that the mutational properties of SVs are poorly characterized is not

because their biological relevance is not appreciated; it obviously is.  Rather, the problem is

technical: SVs are challenging to unambiguously identify and characterize with short-read

sequencing technology, which remains the dominant mode of whole-genome sequencing.

Long-read sequencing is superior to short-read sequencing for SV calling for several

reasons.  Longer reads may span the entire region of interest in one contiguous molecule,

including problematic regions such as low complexity repeats.  Even when a single read

does not span the entire region of interest, longer reads result in superior local *de novo*

assemblies that can span regions much larger than the read length itself.  The obvious

disadvantages include cost and base-calling error rate with respect to base-substitution and

small indel calls, although the accuracy of the latest generation of some long-read platforms

approaches that of short-read sequencing (Harvey et al. 2023).

A complete *de novo* assembly (whole genome assembly, WGA) of an individual is

the only method certain to capture the full array of structural variants in a genome (Chaisson

et al. 2015).  As the cost of long-read technology declines, genome assemblies from single

individuals have become economically feasible.  Two recent studies highlight the rapidity

and magnitude of advances in the field. Loegler et al. (2025) reported the genotype to

phenotype relationship of >1000 yeast strains based on long-read sequencing.  They found

that including SVs (and small indels) in the analyses increased the average explained

heritability of >8000 traits by ~14% above that explained by SNPs alone.  Schloissnig et al.

(2025) estimated the frequency of SVs in >1000 human genomes, resulting in a ~4X

increase in the estimated frequency of SVs in the average human genome.  However, the

application of long-read sequencing technology to characterize spontaneous mutations is

still nascent, although it is evident that long-read sequencing picks up de novo variants that

are missed with short-read sequencing (Noyes et al. 2022; Lopez-Cortegano et al. 2023;

Zhou et al. 2023; Hénault et al. 2024; López-Cortegano et al. 2025; Porubsky et al. 2025).

Here we present estimates of the spontaneous mutation rate and spectrum of SVs in

a set of *C. elegans* mutation accumulation (MA) lines propagated under minimal selection for

~250 generations.  A schematic diagram of the MA experiment is depicted in Figure 1.  Four

MA lines derived from the N2 strain and two MA lines derived from the PB306 strain, and

their ancestral progenitors, were sequenced using Pacific Biosciences Sequel and Sequel 2

technology.  In addition, we report estimates of the standing SV diversity derived from long-

read sequences of the genomes of four wild isolates of *C. elegans.*  Comparison of the

relative proportions of SVs to other variants (SNPs and small indels) among spontaneous

mutations to the same proportions in the standing variation provides an estimate of the

relative strength of selection against SVs in nature.

**Results**

*Workflow*

We define a structural variant (SV) as a variant larger than 30 base pairs (bp), using a

combination of alignment and assembly, followed by conventional variant calling.  We chose

30 bp as our lower bound for designating a variant as an SV rather than the more common

50 bp because preliminary analysis revealed that long-read sequencing allowed us to detect

indels between 30-50 bp that were missed by short-read sequencing.  The difference is only

meaningful when it comes to comparison of SV rates between this study and studies that

use 50 bp as the lower bound; we note the discrepancy where appropriate.

Recognizing the inherent challenges in SV detection, we attempted to automate the

process of discovering the optimal parameters for the variant calling pipeline using

segregating SVs as 'ground truth', under the assumption that a variant called across many

wild isolates is likely real.  We discovered that the signal-to-noise ratio of spontaneous SVs

is less than that of segregating SVs, complicating the development of an appropriate

pipeline.  These complications are detailed in section 2 of the Supplementary Discussion.

We ultimately adopted a workflow loosely based on the methodology of Audano et al.

(2019), with three key modifications:

**Figure 1.** Schematic diagram of the mutation accumulation (MA) experiment. A cryopreserved sample of the ancestor (N2 or PB306) was thawed onto an agar plate (G0, left) and grown for a few days until the cryopreserved worms had reached the L4 larval stage. L4-stage worms were transferred singly to new NGM agar plates; those were the founders of the MA lines (MA.1, MA.2,...MA.n; n=100). At four-day intervals (one generation), a single immature worm was picked at random onto a new plate. This procedure was repeated until the experiment reached Gmax=250 generations. Individual MA lines experienced fewer than Gmax generations, Gave=238 generations. The different colored worms represent (the set of) unique mutations fixed in each MA line, which contribute to the among-line (mutational) variance.

1. Utilization of a superior aligner, minimap2 (Li 2018), instead of blasr (https://github.com/jcombs1/blasr);

2. Variant calling with PBSV (https://github.com/PacificBiosciences/pbsv); and, critically,

3. Manual ("by eye") verification of each SV call by inspection in IGV (Robinson et al. 2017).

This workflow involves aligning error-prone PacBio raw reads ('subreads') to the *C.*

*elegans* reference genome, followed by *de novo* assembly of aligned reads in 60-kb tiles.

Base calling errors in raw subreads can exceed 10% (Weirather et al. 2017; Zhang et al.

2020). The assembled contigs — henceforth referred to as "pseudo-reads"— are then re-

aligned to the reference genome, followed by conventional variant calling using PBSV. The

pseudo-reads exhibit higher sequence quality than raw PacBio subreads, and aligned

pseudo-reads are free from alignment errors that may occur with subread alignment. In a

conventional SV-calling workflow, variants are genotyped only in the sample relative to the

reference genome. In contrast, identifying *de novo* (spontaneous) SVs requires accurate

genotype calls in <u>both</u> the MA progenitor and the derived MA line. Because false positives

can arise from genotyping error in either sample (parent or MA), the spontaneous-mutation

variant calling workflow inherently carries a higher false-positive rate than conventional SV

calling. Consequently, each final mutation call is manually verified using IGV.

Supplementary Table 1 summarizes the base-calling error rates of the raw PacBio

subreads and the assembly-derived pseudo-reads. The error rate of raw reads, calculated

from all N2-derived lines, is 14%, while the pseudo-reads exhibit a significantly reduced per

base error rate of 2.2%. Base-call error rates were calculated after alignment using the 'NM'

tag in the BAM file (See Methods), which represents the Levenshtein distance between the

aligned segment of the read and the reference. Notably, pseudo-reads also exhibit fewer

unaligned segments, measured as soft or hard clips after alignment. The percentage of

clipped bases upon alignment is approximately 5.3% for raw reads, compared to 4.3% for

pseudo-reads. To minimize the effect of reference bias on base-call error rate estimates, we

only report results from N2-derived lines, from which the reference genome is derived,

thereby providing more reliable estimates. The gains to alignment quality are even higher

for wild isolates and PB306-derived MA lines. Base-calling error rates and clipping

percentages for the remaining lines are likewise presented in Supplementary Table 1.

Raw sequence data generated in this study are archived in the NCBI Short Read Archive

(PRJNA1233366).  We also utilized publicly available long-read data of *C. elegans* wild

isolates from the following SRA repositories (PRJNA523481, and PRJNA1025857).

*SV mutation rate*

The mutation rate estimated from observed SVs (i.e., not accounting for false negatives) is

~7 SVs per MA line, over approximately 238 generations per MA line, on average.  In

comparison, we identified an average of 62 single nucleotide variants (SNV) and 25 short

indel mutations per line in the same set of lines.  The inferred SV mutation rate is roughly

10% of the SNV rate and ~30% of the short indel rate, indicating that SVs comprise about

7.5% of new mutations (~6% by the more traditional >50 bp criterion), or roughly one new

SV mutation per-genome every ~30 generations.  Summary statistics of SV mutations are

presented in Table 1.  The full set of SVs and their attributes are given in Supplementary

Tables 2 (N2) and 3 (PB306).

*Estimation of False Discovery Rates*

(*a*) False positives.  The accuracy of SV mutation rate estimates is contingent upon the

accuracy of the variant calling workflow.  False positives can be identified only by either (*i*)

"by eye" validation of putative variants using Integrative Genomics Viewer (IGV), or (*ii*) by

using a different sequencing technique, ideally by PCR amplification of the putative mutant

locus followed by Sanger sequencing of the resulting amplicon.  However, those approaches

become impractical as the number of samples and/or putative variants increases.

We scrutinized each putative variant in the MA lines by visualizing the aligned

pseudo-reads, raw PacBio subreads, and Illumina reads from a separate study in IGV.  We

identified 52 false positives in the original set of called putative SVs, resulting in a signal

(inferred mutant, n=40) to noise (false positive, n=52) ratio of 40/52 = 0.77.  However, 2/3 of

the false positives occurred in one MA line (N2 line MA517).  MA517 had lower sequencing

coverage than the other lines (17X vs an average of 51X for other MA lines), which

**Table 1. Summary statistics**. Column headings: *Line* (strain.MA line # or wild isolate ID); *Type* (MA or Wild isolate); *#Muts* (number of inferred structural variants > 30bp in size); *Man. Corr* (# of SVs that required manual correction; see Methods for explanation); *#False pos.* (# of false positives); *%Recall (100 bp, 1Kb, 10Kb)* (% of simulated SVs of that length recalled; see Methods for explanation); *Ave. Cov.* (average sequencing coverage); *Raw error rate* (Error rate of raw PacBio subreads calculated after alignment to reference); *Pseudo-reads Error rate* (Error rate of assembled contigs after alignment to reference); *#gens MA* (number of MA generations; see Methods); *Rel. fit (%)* (Fitness of MA lines relative to ancestor N2; unpublished data from Saxena et al.); #SNVs (number of base-substitution mutations);   #Indels (number of indel mutations ≤ 30 bp).  "Mutations" in wild isolates are variant compared to the N2 reference genome.  For a more detailed discussion of false positives in wild isolates, see section 2 of Supplementary Discussion.

| Line | Type | #Muts | Man. Corr | # False pos. | Recall rate (100-bp) | Recall rate (1 kb) | Recall rate (10 kb) | Ave. Cov. | Raw Error Rate (%) | Pseudo-read Error Rate (%) | #gens MA | Rel. Fit (%) | # SNVs | # Indels |
|------|------|-------|-----------|--------------|----------------------|--------------------|---------------------|-----------|---------------------|------------------------------|----------|--------------|--------|----------|
| MA517 | N2 - MA | 5 | 3 | 35 | 49 | 53 | 29 | 17 | 14 | 1% | 242 | 72.6 | 53 | 17 |
| MA530 | N2 - MA | 7 | 2 | 1 | 65 | 77 | 47 | 56 | 13 | 3% | 243 | 122.2 | 51 | 26 |
| MA563 | N2 - MA | 5 | 1 | 0 | 63 | 73 | 43 | 33 | 15 | 3% | 237 | 66.0 | 70 | 24 |
| MA566 | N2 - MA | 8 | 4 | 6 | 62 | 73 | 45 | 37 | 13 | 2% | 238 | 58.6 | 71 | 21 |
| MA445 | PB306 - MA | 10 | 5 | 6 | 57 | 68 | 42 | 50 | 16 | 6% | 232 | 126.8 | 76 | 30 |
| MA459 | PB306 - MA | 5 | 2 | 4 | 55 | 62 | 42 | 79 | 16 | 6% | 235 | 61.4 | 61 | 31 |
| PB306 | WI | 7691 | - | - | 59 | 69 | 43 | 188 | 15 | 7% | - | - | 189278 | 73876 |
| CB4856 | WI | 11028 | - | - | 56 | 65 | 37 | 59 | 18 | 7% | - | - | 285787 | 104591 |
| QX1211 | WI | 16528 | - | - | 50 | 49 | 30 | 54 | 18 | 8% | - | - | 505065 | 164826 |
| JU1088 | WI | 6181 | - | - | 60 | 69 | 39 | 52 | 17 | 6% | - | - | 151734 | 61013 |
| N2 | Baer - N2 | - | - | - | 62 | 76 | 54 | 83 | 15 | 2% | - | - | - | - |

potentially introduced alignment and/or assembly artifacts.  When line MA517 is excluded,

the signal-to-noise ratio increases to ~2:1.

Among the true positive calls, minor corrections were made after visualizing the SVs

in IGV.  Corrections are required when an MA SV occurs at a locus that already carries an

ancestral SV.  Including other minor reasons for adjustments, corrections were necessary for

17 of the 40 true positive calls.  The reasons for corrections are provided in Supplementary

Tables 2 and 3.

(*b*) False negatives.  In contrast to identification of false positives, which both can and must

be done by brute force, false negatives can be inferred using bioinformatics (Sedlazeck et al.

2018b).  We estimated the false negative rate by (*i*) simulating SVs ("pseudo-variants") in

the reference genome and (*ii*) attempting to identify ("recall") the pseudo-variant in the

sequenced sample.  For example, a pseudo-deletion introduced in the reference genome

should appear as a variant pseudo-insertion in the sequenced sample.  We measured recall

in both the MA lines and wild isolates, with the latter providing information about the extent to

which reference bias impacts false negative rates.

Recall rates differ for different classes of SV (Figure 2, Supplementary Table 4).  The

average recall rate is high (~90%) for deletions up to 10 kb.  The recall rate of 100 bp

insertions is also roughly 90% but declines as length increases. Recall rates were lower in

wild isolates compared to the N2 strain, presumably because the *C. elegans* reference

genome is N2; the difference is greater in the highly divergent QX1211 strain (Figure 2,

Table 2).  As a result, slightly more (~4%) de novo SVs were missed in the PB306 MA lines

than in the N2 MA lines.  For reasons that are unclear, we miss almost all small inversions

(100-bp), whereas we recall roughly a quarter of 100-kb inversions.

**Figure 2**. Recall rates by genotype, SV type, and SV size.  N2 has the highest recall rate
across SV types and sizes.  Small deletions (100 bp) have the highest recall rates, comparable
to 1-kb and 10-kb deletions. Recall rates of insertions drop sharply beyond 100 bp.  Recall
rate decreases with increasing SV size and genetic divergence from N2, a trend also observed
in PB306-derived MA lines (not shown). Inversions have the lowest recall rates. QX1211, the
most divergent genotype from N2, is the most challenging for SV detection. Minor horizontal
jitter was added to the x-axis to improve the visual separation of overlapping points.

*SV mutation spectrum*

Frequencies of SVs are listed by type and average size in Table 2.  The total number of
bases affected is approximately 16 kb in the four N2-derived lines, with a net loss of 1.5 kb
across four lines (7 insertions and 18 deletions).  For the two PB306-derived lines,
approximately 15 kb of bases were affected in total, with a net gain of 7 kb (5 insertions and
10 deletions).  Deletions outnumber insertions 2:1, the same bias observed among smaller
indels in a larger set of N2 and PB306 MA lines (Rajaei et al. 2021).  Among wild isolates,
the number of insertions and deletions are nearly equal, with large insertions outnumbering

237 large deletions by roughly 10% among SVs, and insertions and deletions being nearly equal

238 among smaller indels.

239 **Table 2**. SV spectrum by variant type and number of bases affected.
240

| Line | Type | Insertions | Deletions | Inversions | Net bases affected | Net base gain (loss) |
|------|------|-----------|-----------|------------|-------------------|---------------------|
| N2.517 | MA | 2 | 3 | 0 | 5,342 | 184 |
| N2.530 | MA | 3 | 4 | 0 | 1,509 | 843 |
| N2.563 | MA | 2 | 3 | 0 | 7,606 | (309) |
| N2.566 | MA | 0 | 8 | 0 | 2,166 | (2,166) |
| PB.445 | MA | 3 | 7 | 0 | 3,501 | (1,397) |
| PB.459 | MA | 2 | 3 | 0 | 12,184 | 8,536 |
| PB306 | WI | 4347 | 3280 | 64 | 4,027,284 | 1,791,544 |
| CB4856 | WI | 5887 | 5056 | 85 | 5,205,431 | 1,562,847 |
| QX1211 | WI | 9148 | 7279 | 101 | 7,949,707 | 2,849,281 |
| JU1088 | WI | 3597 | 2525 | 59 | 3,331,096 | 1,465,676 |

Although our variant caller (PBSV) reports all SVs as insertions or deletions (or

inversions, although we did not find any), several of these events represent tandem repeat

changes or potentially translocation events.  A slight majority of spontaneous SVs (22/40)

result from simple repeat expansions or deletions; the average size of the repeat family

across 22 independent events is 93 bp.  We annotate these cases manually in

Supplementary Tables 2 and 3, based on sequence context and alignment features.  For

example, translocations are typically flagged when inserted sequences show high similarity

to distant genomic loci, and tandem repeat changes are identified when insertions or

deletions occur within repetitive motifs.  These classifications are conservative, as formal

proof of translocation or repeat structure would require defining thresholds for sequence

identity and repeat purity.

Spontaneous SV mutations are marginally larger in size compared to segregating

SVs identified in wild isolates (mean SV size: MA=808 bp, WI=602 bp; median SV size:

MA=215 bp, WI=125 bp; Kolmogorov-Smirnov test, p<0.1).  This difference is primarily

driven by the size of insertions (mean insertion size: MA=1583 bp, WI=799 bp; median

insertion size: MA=441 bp, WI=162; K-S test, p<0.05), whereas deletions show no significant

difference in size (mean deletion size: MA=475 bp, WI=375 bp; median deletion size:

MA=152 bp, WI=105 bp; K-S test, p>0.37).  Although the difference in deletion sizes is not

statistically significant, we observe a heavier tail in the size distribution of spontaneous

deletions, similar to that of spontaneous insertions. This distribution exhibits a multimodal

pattern, indicative of larger deletions in the MA lines (Supplementary Figures S1-S3).

Four of the 40 SVs are associated with transposon-related events. In the PB306 MA

line 445, a CER1 transposon insertion was identified on chromosome 1, resulting in an 8.8-

kb insertion within a short 80-bp intron of the *smg-1* gene.  The inserted sequence contains

3' end of the *plg-1* gene and the full CER1 transposon sequence.  Similarly, in the N2 MA

line 566, an RTE transposon insertion was observed, leading to a 3.2-kb insertion at the 5'

end of the *alg-1* gene.  3/4 TE events are >3 Kb, whereas the largest non-TE event is ~2700

bp.  The *C. elegans* genome is replete with TEs (Larrichia et al. 2017; Bush et al. 2025), so

the greater average size of spontaneous SVs relative to standing variants is unlikely due to

an atypically high degree of TE activity in our MA lines.  On the contrary, the size

discrepancy implies that most TEs are effectively removed by purifying selection.

*Distribution of SVs across the genome*

The distribution of spontaneous SV mutations among exons, introns, and intergenic regions

does not deviate significantly from expectations based on genomic composition (10 SVs in

exon, which constitute ~32% of the genome; 19 SVs in intron, ~35%; 11 SVs in intergenic

regions, ~33%; 3x2 goodness-of-fit test, $\chi^2$=2.76, df=2, N=40 SVs, p>0.25).  Seven of the 40

SVs are located within hyperdivergent genomic regions, consistent with their frequency in

the *C. elegans* genome (~20%; Lee et al. 2021).

In the wild isolates, the distribution of segregating SVs across the genome mirrors

the SNV distribution – higher in chromosome arms and lower in the centers (Figure 3); see

Figure 3 of Andersen et al. (2012) for the distribution of SNVs along *C. elegans*

chromosomes.  The recombination rate is higher in chromosomal arms (of autosomes) than

centers in *C. elegans* (Rockman and Kruglyak 2009). The greater genetic diversity in

chromosomal arms is consistent with reduced Hill-Robertson interference in the arms

compared to the chromosome centers. We do not see this pattern on the X chromosome,

**Figure 3.** Distribution of SVs within and among chromosomes in wild isolates (density

shown in blue) and MA lines (vertical black lines). The overall distributions between these

two groups are not significantly different (goodness-of-fit test; See text for details).

which is also consistent with expectation, since the recombination rate is nearly uniform on

the X.

It is possible, however, that the chromosomal distribution of genetic diversity is not

solely due to variation in the efficacy of purifying selection; it may be that the SV mutation

rate is higher in the arms.  Indeed, we have previously shown that the base-substitution

mutation rate is higher in the arms of the chromosomes than the center in *C. elegans*

(Saxena et al. 2019).  This view is supported by the fact that low complexity DNA is

overrepresented in the arms of *C. elegans* chromosomes (see Supplementary Figures S4

and S5).  Even with the small sample size of this study, we observe significantly more

spontaneous SVs in the arms of the chromosomes than in the centers (26 SVs in arms

which comprise 46% of the genome vs. 10 SVs in the center which comprise 47% of the

genome, excluding tips; $\chi^2$=7.5, df=1, N=36, p<0.007).  In contrast, the distribution of SVs in

wild isolates did not differ significantly from the distribution of SVs in MA lines (20,174 SVs in

arms, and 6,723 SVs in center, $\chi^2$=0.14, df=1, N=26,993, p>0.7). Including chromosomal tips

did not alter these conclusions.  Genomic boundaries for chromosomal arms and centers

follow Table 1 of Rockman et al. (2009).  It may be that the distribution of SVs across the

genome can be explained by a combination of mutation (more in the arms than the centers)

and selection (stronger Hill-Robertson interference in the centers).

*Fitness effects of SVs*

To infer the fitness effects of SVs, we compared the spectrum of spontaneous SVs to that of

wild isolates (Table 3).  Spontaneous SVs are marginally larger than segregating SVs,

consistent with the expectation that larger SVs are more strongly deleterious and thus

preferably removed by natural selection.

For a more fine-scaled assessment of fitness effects, we compared relative diversity

(*w*\*; see Equation 1 in the Methods) of different categories of variants between MA lines and

wild isolates (Table 3).  SNVs in introns, intergenic regions, and synonymous SNVs in exons

were pooled into an (assumed) neutral class of variants for this analysis.  Unsurprisingly,

SVs in exons are under strong purifying selection; $w^*$ in exons is approximately 14% of the

neutral expectation (p<0.0001, 2x2 Fisher's exact test). $w^*$ in introns is ~30% of the neutral

expectation (p<0.0001, 2x2 Fisher's exact test).  Perhaps more surprisingly, $w^*$ of SVs in

intergenic regions is approximately half of the neutral expectation ($w^*$=0.52, p<0.04; 2x2

Fisher's exact test), suggesting strong purifying selection against SVs even in intergenic

regions.  This observation is, to our knowledge, the first of its kind and warrants further

investigation into the functional impacts of intergenic SVs, which may have significant

implications for human genetics research.  We discuss the implications of this result to the

concept of 'junk DNA' later.

The preceding calculations are potentially biased, because the MA lines (4 N2, 2

PB306) are more similar to the (N2) reference genome than are the wild isolates, and the

failure to recall rate (FTR) increases with divergence from the reference genome (Figure 2).

To investigate the possible role of reference bias, we repeated the calculations including

only standing variants in the PB306 genome.  PB306 is relatively close to N2 and shares 2/6

MA lines, so the effects of reference bias should be less in PB306 than in the average of the

wild isolates.  The results are qualitatively similar (Table 3), although the values of $w^*$ are

slightly greater (i.e., closer to the neutral expectation) in PB306 than in the full set of

standing variants, and the relative paucity of SVs in intergenic regions no longer meets the

criterion of statistical significance ($w^*$=0.62, p>0.09)

**Table 3**. Distribution of variant types by genomic context. See Equation 1 in the Methods for details of calculation of relative diversity $w^*$. The subscripts "*All*" and "*PB306*" for $w^*$ and *p*-values represent values calculated from all four wild isolates and PB306 only, respectively. Values of $w^*$ in bold font are statistically significant ($\square = 0.05$); "*p*" is the *p*-value of the 2x2 right-sided Fisher's exact test of the hypothesis that the focal class is neutral (i.e., $w^*=1$); see Methods for details.

| Type | Context | All Wild Isolates | PB306 only | PB306 MA | N2 MA | All MA | $w^*_{All}$ / $w^*_{PB306}$ | $p_{All}$ / $p_{PB306}$ |
|------|---------|-------------------|------------|----------|-------|--------|------------------------------|--------------------------|
| SV | Exon | 2866 | 630 | 4 | 6 | 10 | **0.14 / 0.17** | <0.0001 / <0.0001 |
| | Intron | 11818 | 3031 | 5 | 14 | 19 | **0.31 / 0.44** | <0.0001 / <0.002 |
| | Intergenic | 11292 | 2470 | 6 | 5 | 11 | **0.52** / 0.62 | <0.04 / >0.09 |
| SNV | Splice Variant | 2840 | 515 | 2 | 1 | 3 | 0.48 / 0.48 | >0.17 / >0.17 |
| | Disruptive Start/Stop Codon | 1830 | 313 | 2 | 2 | 4 | **0.23 / 0.22** | <0.02 / <0.02 |
| | Missense Variant | 78191 | 14202 | 11 | 30 | 41 | 0.96 / 0.96 | >0.42 / >0.42 |
| | Intergenic Region | 243014 | 42215 | 53 | 88 | 141 | Neutral | - |
| | Intron Variant | 274542 | 50937 | 52 | 94 | 146 | Neutral | - |
| | Synonymous Variant | 82326 | 15411 | 6 | 8 | 14 | Neutral | - |
| Indel | Frameshift Variant | 8446 | 1339 | 4 | 5 | 9 | **0.47 / 0.41** | <0.04 /<0.02 |
| | In-frame Indel | 2905 | 463 | 0 | 4 | 4 | 0.36 / **0.32** | >0.06 / <0.05 |
| | Intergenic Region | 74279 | 13577 | 20 | 26 | 46 | 0.81 / 0.82 | >0.10 / >0.12 |
| | Intron Variant | 100183 | 19053 | 32 | 45 | 77 | **0.65 / 0.69** | <0.001 / <0.003 |

**Discussion**

*Rate, Spectrum, and Selective Effects of Structural Variants*

The two primary findings of this study are (1) SVs constitute at least 7.5% of spontaneous mutations in the *C. elegans* genome, and (2) SVs are efficiently removed by selection, apparently even from non-coding regions of the genome.  But also (3), it is evident that accurately calling de novo SVs is a difficult task that requires a lot of ad hoc curation, even on such friendly terrain as long sequence reads at high coverage of a relatively small, extremely well-characterized, nearly completely homozygous genome.

Robust estimates (i.e., from long-read sequencing) of spontaneous SV mutation rates are now available from *E. coli* (Zhou et al. 2023), the unicellular green alga *Chlamydomonas reinhardtii* (Lopez-Cortegano et al. 2023), mouse (López-Cortegano et al. 2025) and humans (Noyes et al. 2022; Porubsky et al. 2025).  *E. coli* and humans span approximately four orders of magnitude of effective population size (Ne ≈ $10^4$ for humans, ≈ $10^8$ for E. coli; Sung et al. 2012), with the accompanying difference in the effectiveness of selection (the "drift barrier"; $s>1/N_e$).  The per-genome, per-generation rate of single nucleotide (SNV) mutations also varies by roughly four orders of magnitude between *E. coli* (~$10^{-3}$/genome/generation) and humans (~30/haploid genome/generation).  SVs constitute approximately 30% of spontaneous mutations in *E. coli* (Zhou et al. 2023) and 6-12% in *Chlamydomonas* (Lopez-Cortegano et al. 2023), but only about 2-3% in humans (Porubsky et al. 2025) and mouse (López-Cortegano et al. 2025).  The same calculations for *C. elegans* reveal a per-genome, per-generation SNV mutation rate of ~0.25/genome/generation and an SV mutation rate of ~0.03/genome/generation, which corresponds to ~7.5% of all spontaneous mutations (~6% using the >50bp threshold) observed across the six MA lines in this study (Table 3; Supplementary Tables 2 and 3). The data are obviously sparse, but the trend is evidently toward a positive relationship between $N_e$ and the fraction of spontaneous mutations that are SVs.  Given that SVs are surely more deleterious on average than SNVs, the apparent higher proportion of SV mutations in taxa with more stringent selection is intriguing and seems counterintuitive.

Comparison of the relative diversity ($w^*$) between classes of variants provides a

heuristic test of the relative strengths of selection. For neutral variants, $E[w^*] = 1$ (Equation

1 in the Methods); the smaller $w^*$ is, the more effectively are variants removed from the

population by selection. Comparison of relative diversity ($w^*$) across genomic features

(Table 3) shows that structural variants (SVs) exhibit substantially reduced diversity in exons

compared to missense SNVs, consistent with stronger purifying selection against exon-

disrupting SVs. This is not a surprise; a large indel in an exon is undoubtedly more

deleterious on average than an amino acid replacement. Selection also effectively removes

SVs from introns. Somewhat less obviously, selection apparently is also effective at

removing SVs from intergenic regions. $N_e$ of *C. elegans* is on the order of $10^4$ (Teterina et al.

2023), which implies that many intergenic SVs experience selection $s>10^{-4}$.

*Limitations in Estimating the Contribution of Structural Variants to Fitness Decline*

Over the course of an average of 238 generations of mutation accumulation (MA), the six

MA lines in this study exhibited an average decline in absolute fitness of ~15% (Baer et al.

2006). Structural variants (SVs) constitute only ~7.5% of all detected variants. However,

accurately quantifying their contribution to fitness decline is challenging without detailed

knowledge of the distribution of fitness effects (DFE), which is likely influenced by factors

such as variant size and genomic context.

For instance, SVs occurring in exons represent a mutation class with particularly low

diversity in natural populations. After adjusting for differences in mutation rates across

variant classes, SV diversity in exons is approximately 14% of the neutral expectation. Even

that is an underestimate, however, given that the standing variants potentially include

relatively new deleterious mutations that have persisted long enough to be observed but are

on their way to being selected out of the population (i.e. mutation-selection balance).

In contrast, small indels in introns, the most abundant non-neutral mutation class in

our dataset, exhibit diversity at ~65% of the neutral expectation. Despite this, the MA lines

contain approximately eight times as many small indels in introns than SVs in exons. While

it is difficult to estimate relative selection coefficients (*s*) for these two classes without a well-

characterized DFE, their relative abundance suggests that small indels in introns could

contribute substantially to fitness decline, even if their average effect is weaker than that of

SVs in exons.  Ignoring variant frequency in favor of a focus on average effects risks

underestimating the evolutionary consequences of more frequent but potentially less

deleterious mutations, such as small indels.

*Effect of False Negatives in Inference of SV Spectrum and Estimates of Selection*

Interpretation of the SV spectrum must also be considered in light of the detection limits of

our variant-calling pipeline.  Our workflow reliably detects variants between ~30 bp and 2 kb,

with recall rates >80% for 1 kb deletions and ~60–80% for 1 kb insertions (Figure 2;

Supplementary Figures S1–S3).  These recall rates are similar in both N2 and the wild

isolates, with the exception of the QX1211 line, which reduces concern about reference-

strain bias within this size range.  Detection declines sharply for insertions >10 kb, though

such large events account for only a tiny fraction of all calls.

Because the overwhelming majority of variants in our dataset fall within the 50 bp–2

kb range (~90% of spontaneous variants and ~95% of segregating SVs), our conclusions

about the SV spectrum and the strength of selection necessarily pertain to this size class.

Because we did not detect many variants much larger than ~2 kb, we avoid making

inferences about mutation or selection acting on larger SVs for which our detection power is

limited.

*Structural Variants in Low-Complexity Mutational Hotspots*

A significant proportion (~22.5%; 9 out of 40) of structural variants (SVs) identified in

mutation accumulation (MA) lines occur in low-complexity mutational hotspots.  Previously,

Konrad et al. (2018) reported a similar estimate (~31%) for duplications and a much higher

estimate (~81%) for deletions occurring in loci with pre-existing CNVs in the ancestor, based

on short read sequencing of N2-dervied MA lines in *C. elegans*.  Among the SV calls in MA

lines, the majority (22 out of 40) are simple repeat expansions or deletions.  These

annotations were determined through manual inspection of variants using the JBrowse

genome browser as well as IGV rather than automated annotation, which, while scalable,

may introduce noise.  The repeating units range in size from as small as dinucleotide

repeats (e.g., AT, CA, CT) to larger structural events, such as the deletion of a single copy

within a four-copy 360-mer repeat.

An unexpected observation is that 9 of the 22 SVs identified in repeat-rich regions

required a 'manual correction' due to a common misclassification in the workflow – where

the MA variant call occurs on a locus with a pre-existing variant in the ancestor.  This

suggests that these loci represent mutational hotspots.  Notably, the variant caller PBSV did

not correctly handle these cases, even in joint genotyping mode.  Supplementary Table S2

and S3 provides annotations for SV calls in MA lines, detailing the necessary corrections

following manual verification using IGV and genome browser inspection.

For example, a variant call on chromosome I of PB306-derived MA line MA445 (I:

1567493), illustrated in Figure 4a with a pedigree and 4b with an IGV snapshot highlights

this issue.  In the PB306 ancestor, this region contains a 3.1 kb insertion composed of

perfectly repeating copies of a 173-mer interspersed with additional sequence.  Some of this

additional sequence contains similar repeating motifs, but of smaller units, and precise

annotation is difficult due to sequencing errors.  The sequence of this insertion, along with its

repeating unit, is provided in Supplementary Sequence File S1 in FASTA format.  We show

the alignment of raw PacBio reads in Supplementary Figure S6 and show the repeating

copies of 173-mer highlighted in text in Supplementary Figure S7.  The same 3.1 kb

insertion is also present in the PB306-derived MA line MA459, consistent with the

expectation that a variant present in the ancestor is inherited by its descendant.  This

insertion is absent in the N2 ancestor, indicating that it is polymorphic in *C. elegans.*

Interestingly, in MA445, this region harbors a 1.9 kb insertion, effectively representing a 1.2

kb deletion relative to the PB306 ancestor.  However, variant calling software failed to

annotate this as a deletion event, necessitating extensive manual validation of both raw and

assembled reads.  Because some repeat copies are perfectly conserved while others exhibit

minor variations, the identification of this type of event remains beyond the resolution of

current variant calling algorithms.

The need for manual correction in 9 out of 22 repeat-associated SVs raises an

important question: Are certain repeat regions disproportionately prone to structural

variation?  The *C. elegans* genome contains a vast number of repeat-associated sequences.

We observed 122,161 distinct repeat-associated sequences in the WS270 GFF3 file that

span roughly 19.5% of the genome (See Methods).  Despite this substantial genomic

footprint, only a small subset of these regions harbored recurrent SVs in our lines.  Our

findings suggest that loci already carrying an SV in the ancestor are prone to recurrent SVs,

making them significant hotspots for genetic variation.  Analogous SV mutational hotspots

have been identified in the human genome (Porubsky et al. 2025).  Furthermore, these

regions are likely underrepresented or misannotated by automated variant calling pipelines.

With only 9 such identified events, it is beyond the scope of this study to understand the

factors that cause some repeat sequences to be hotspots.  Notably, 2 of the 9 manually

corrected repeat-associated SVs occur in hyperdivergent regions—an observation worth

noting, though we refrain from drawing broader conclusions.

**(4a)**

**(4b)**

**Figure 4**. (**a**) A schematic representation of the variants identified at the locus (I: 1567493) in
the pedigree of MA lines.  PBSV initially classified the 1.9 kb insertion and the 3.1 kb
insertion as two independent mutations.  However, a deeper analysis—accounting for
sequencing errors—reveals that a 1.2 kb segment was deleted within the inserted sequence in
the ancestor, leading to the observed 1.9 kb insertion. (**b**) IGV snapshot of the 1.2 kb deletion
in PB306-derived MA 445.  The IGV snapshot shows pseudo-read alignments, which are
assembled contigs representing longer sequences.  Some pseudo-reads fail to fully capture the
inserted segment and appear as soft-clipped sequences.  If the inserted segment is not entirely
covered within the pseudo-read, it is clipped from the alignment, appearing as base calls
(colored bars) instead of clean (gray bars), fully aligned sequences.  Some insertion
sequences are misaligned because of the repetitive nature of the locus; however, all pseudo-
reads carry the insertion.  Supplementary Figure S6 presents the same analysis using error-
prone PacBio raw reads.

*Estimating the Fraction of the Genome under Evolutionary Constraint*

We observe that selection efficiently removes most SVs, regardless of their genomic

locations (genic versus intergenic).  This finding challenges the conventional view of "junk

DNA," which is often assumed to evolve free from the constraints of purifying selection.  Our

results suggest that a larger fraction of the genome may be functionally relevant than

previously inferred from studies focusing on base substitutions.  Unconstrained genomic

regions are typically defined by the absence of purifying selection.  However, it is possible

that some genomic regions can tolerate base substitutions but not SVs.  For instance,

consider a hypothetical low-complexity region between an enhancer and a promoter

(Bateman and Johnson 2022).  Base substitutions in such regions may not disrupt gene

function, whereas SVs that alter the distance between the enhancer and promoter might,

which could in turn negatively impact fitness.  Similarly, regulatory sequences flanking genes

may be more tolerant of base substitutions than of small indels or larger SVs.  In a similar

vein, Payer et al. (2017) found that polymorphic *Alu* elements in the human genome were

significantly associated with Trait Associated SNPs (TASs) in human GWAS.  Larger-scale

studies, capable of identifying more SV mutations, could reveal many such regions that

appear to evolve freely based on SNP variation but are subject to purifying selection against

SVs.

Taken together with recent findings from other organisms, the results reported here

have implications for the design and interpretation of studies of complex traits in humans

(e.g., GWAS).  The most recent estimates of $N_e$ in *C. elegans* – on the order of $10^4$ (Teterina

et al. 2023) – are surprisingly close to estimates of $N_e$ in humans (Tenesa et al. 2007), given

that $10^8$ *C. elegans* could fit comfortably inside a single human.  A recent study with yeast

(*S. cerevisiae*) revealed that including SVs in GWAS explained an additional ~14% of

heritable variance in thousands of traits, relative to the variance explained by SNVs alone

(Loegler et al. 2025).  If even intergenic SVs are under effective selection ($s>1/N_e$) in *C.*

*elegans*, it implies the same may be true for humans, all else equal.  Which in turn implies

that SVs are likely to explain significant additional heritable variance for human complex

traits, but also that those variants are likely to be rare.

**Conclusion**

The spontaneous SV mutation rate is inferred to be approximately 7.5% of the SNV/small

indel rate, but that is likely an underestimate.  SVs are efficiently removed from natural

populations, indirectly implying a function (biochemical or otherwise) to a larger fraction of

the genome than is typically assumed.

**Material and Methods**

*MA Protocol*

The MA protocol has been described in detail previously (Baer et al. 2005) and is depicted in

Figure 1.  The maximum number of MA generations (Gmax) is 250.  The method by which

we estimated the number of generations of MA for a line (below Gmax) is reported in

Saxena et al. (2019).

*DNA extraction, library preparation, and sequencing*

Cryopreserved samples of the MA lines were thawed onto 100 mm NGMA agar plates

seeded with a lawn of *E. coli* OP50 and grown for 2-3 days, until there were many gravid

worms, at which time worms were harvested, "bleached" to remove bacterial and fungal

contamination (Stiernagle 2006), and embryos transferred to a new, seeded 100 mm plate.

L1 larvae were collected the next day and transferred to seven seeded 100 mm plates and

grown until the plates were nearly starved.  Worms were washed from the plates, double-

rinsed in M9 buffer, and pelleted for DNA extraction.  DNA was extracted from each sample

with the Quiagen Gentra Puregene kit, following the manufacturer's instructions.  Extracted

DNA samples were sent to the University of Maryland genomics core facility for library

preparation and sequencing on the Pacific Biosciences Sequel platform.

*Variant Calling Pipeline*

We evaluated multiple variant calling strategies, including direct alignment-based

approaches and de novo assembly followed by genome-to-genome alignment using

Assemblytics (Nattestad and Schatz 2016). Ultimately, we adopted a modified version of the

SMRT-SV2 (Audano et al. 2019) pipeline, as it provided a balance between sensitivity and

specificity while minimizing false positives. A detailed discussion of our rationale for

selecting this approach over others is provided in section 1 of the Supplementary

Discussion.

The SMRT-SV2 pipeline processes error-prone PacBio subreads by aligning them to

the reference genome using blasr (Chaisson and Tesler 2012). These alignments are then

segmented into 60-kb tiles, which are assembled using Canu (Koren et al. 2017). This

localized assembly approach reduces interference from reads originating from other

genomic regions. Importantly, the assembled contigs effectively "clean" the PacBio

subreads, leading to improved downstream alignments. While SMRT-SV2 automatically re-

aligns these contigs using blasr, we opted to replace blasr with minimap2 (Li 2018), which

provided superior alignment performance for our dataset.

SMRT-SV2 includes its own variant caller. However, it led to an unacceptably high

false positive rate for de novo mutations. To improve accuracy, we benchmarked several

alternative structural variant (SV) callers, including Sniffles (Sedlazeck et al. 2018a) and

PBSV (Wenger et al. 2019), ultimately selecting PBSV due to its optimized performance on

high-quality PacBio reads. Further details on the performance of SMRT-SV2, Sniffles, and

SMRT-SV2 variant calling are provided in section 3 of the Supplementary Discussion.

In cases where a potential variant was detected, SMRT-SV2 produced multiple

assemblies—referred to as "pseudo-reads"—for the candidate locus. These pseudo-reads

were then aligned with minimap2 and variants were called using PBSV. Despite this

refinement, we still observed many false positives, albeit a manageable number, further

necessitating manual curation. Each candidate variant was inspected in IGV (Robinson et

al. 2017) and JBrowse genome browser (Diesh et al. 2023), comparing error-prone raw

PacBio subread alignments, pseudo-read alignments, and independent Illumina sequencing

data generated for a separate study.  Only variants present in the mutation accumulation

(MA) line but absent in both the MA ancestor and other MA lines were considered true de

novo mutations.  Once validated, variants were functionally annotated using genome

browser annotations to classify their genomic context (e.g., exon, intron, UTR, repetitive

element, hyperdivergent region, potential regulatory elements).

For identifying segregating SVs, we followed the same pipeline with one key

difference: instead of an ancestor-descendant comparison, all wild isolates were jointly

called using PBSV.

*Estimating Sequencing Base-Call Error rate*

Base-call errors arise during PacBio sequencing and manifest as nucleotide-level

mismatches between raw subreads and the reference genome.  We quantified these errors

using Levenshtein distance, which measures the minimal number of single-character edits

(insertions, deletions, substitutions) required to transform one sequence (read) into another

(aligned segment of reference).  Raw subreads were aligned to the reference genome using

minimap2 (Li 2018), and Levenshtein distance was computed for each alignment using the

'NM' tag in the bam file.  We estimated the base-call error rate of pseudo-reads generated

after alignment and assembly similarly by aligning the pseudo-read to reference, and parsing

the 'NM' tag in the bam file.  This analysis ignores soft-clipped regions which could contain

worse base-call quality on the edges of the reads, hence this analysis provides a

conservative estimate of base-call error rate.

*Variant Annotation*

Annotation of structural variants (SVs) identified in the mutation accumulation (MA) lines was

performed manually.  Basic genomic classifications—such as exon, intron, and intergenic

regions—were straightforward to assign.  Similarly, regulatory annotations, including DNase

hypersensitive sites, transcription factor binding sites, and hyperdivergent regions, could be

systematically determined using existing datasets.  However, annotating repetitive elements

within SVs was significantly more time-consuming.  Many SVs overlapped repetitive

sequences that were not explicitly annotated by RepeatMasker, requiring manual inspection

of the variant and read sequences to deduce repeat structure.

In contrast, annotation of SVs in wild isolates was automated using GFF3-based

genome annotations.  Variant calls were intersected with genomic features using *bedtools*

(Quinlan and Hall 2010), allowing for systematic classification. Automated annotation tools

such as *snpEff* (Cingolani et al. 2012) were not suitable for SV annotation, as they do not

properly account for the full extent of large structural variants.  For example, an 8-kb deletion

may only be annotated at its breakpoints rather than across its entire span.  To ensure

accurate functional classification, we assigned the highest-impact annotation observed

across the deleted region.  If a deletion started upstream of a gene but extended into coding

sequences, it was annotated based on the most functionally significant feature it disrupted

(e.g., an exon rather than an upstream region).

Short variants (SNPs and small indels) in both MA and wild isolates were annotated

using *snpEff* with the following flags: -no-downstream -no-upstream -no-utr.  As a result,

upstream, downstream, and UTR annotations were instead categorized as intergenic,

maintaining consistency with our SV annotation approach.  *snpEff* produces a diverse set of

annotations, many of which are redundant or overly specific for our purposes.  To

standardize annotation categories, we applied a systematic simplification approach using a

predefined mapping (Supplementary Table 5).  This mapping condensed detailed annotation

labels into a manageable number of biologically meaningful categories, ensuring consistency

across variant types.

*Estimating the relative strength of selection for different categories of mutations*

To estimate the relative strength of selection we first defined a neutral class of variants,

including intergenic SNVs, intronic SNVs, and synonymous SNVs within exons.  We

included intronic and intergenic variants in addition to synonymous variants to increase the

sample size of MA variants in the neutral class, given the small number of synonymous MA

variants in our dataset (n=14).  The relative diversity ($w^*$) of a focal class of variant (e.g.,

SVs in exons) was calculated from the equation:

$$w^* = \frac{\frac{\#\ Segregating\ variants\ in\ focal\ class}{\#\ Segregating\ variants\ in\ neutral\ class}}{\frac{\#\ Spontaneous\ mutations\ in\ focal\ class}{\#\ Spontaneous\ mutations\ in\ neutral\ class}} \qquad \text{[Equation 1]}$$

This approach quantifies the standing diversity within a particular class of (focal) variant

relative to the diversity in the neutral class, accounting for the mutation rates in both

categories.  If the focal class of variants is neutral, $E(w^*) = 1$.  If the focal class of variant is

subject to purifying selection, there will be fewer segregating variants than expected given

the mutation rate and $w^*<1$.  Statistical significance is assessed by a 2x2 Fisher's exact test.

Counts of variants in each category in the wild isolates and MA lines are given in Table 3.

*Simulation of Reference Genomes and Estimation of False Negative Rates*

To evaluate the sensitivity of our variant calling pipeline, we simulated structural variants in

the *C. elegans* reference genome (WS270) using the R package *RSVSim* (Bartenhagen and

Dugas 2013).  For each variant type—insertions, deletions, and inversions—we generated

three independent simulated genomes.  Within each variant type, we introduced 100 variants

of lengths 100 bp, 1,000 bp, and 10,000 bp, as well as 50 variants of 100,000 bp.

Following simulation, we applied our variant calling pipeline to assess recall rates.

Introducing structural variants into the reference genome shifts the positions of subsequent

variants, making direct locus-based comparisons dependent on tracking the precise order of

variant introduction.  While this could be addressed if the software explicitly recorded variant

placement order, doing so would introduce additional dependencies into the analysis.  To

avoid this complexity, we instead used a size-based evaluation approach, considering a

variant correctly detected if its length fell within predefined tolerance ranges: (98–102) bp,

(995–1,005) bp, (9,995–10,005) bp, and (99,995–100,005) bp for the respective size

categories. This approach simplified the evaluation while providing a systematic estimate of

false negative rates.

*Estimating number of repeat-associated entries in C.elegans genome*

To quantify the genomic burden of repetitive DNA, we merged overlapping annotations from

the WS270 GFF3 file, including "repeat_region", "inverted_repeat", and "tandem_repeat"

entries. This yielded 122,161 distinct repeat-associated regions, collectively spanning

approximately 19.5 Mb or 19.5% of the *C. elegans* genome.

*Data Analysis*

All statistical analyses (e.g., Kolmogorov-Smirnov test, Fisher's exact test) were performed

in Python using SciPy (Virtanen et al. 2020). Data visualization was conducted using

Matplotlib (Hunter 2007) and Seaborn (Waskom 2021).

Acknowledgments

We thank Tim Crombie, Joanna Dembek, Lindsay Johnson, and Mike Snyder for assistance

in the lab, and Erik Andersen and Annalise Paaby for providing the wild isolate data. Two

anonymous reviewers provided insightful comments on a previous version. Support was

provided by NIH award GM107227 to CFB.

Declaration of interests

A.S. contributed to this article as a Ph.D student at the University of Florida and in his

personal capacity. The views expressed do not necessarily represent the views of

Regeneron Pharmaceuticals Inc.

Figure Legends

Figure 1. Schematic diagram of the mutation accumulation (MA) experiment. A

cryopreserved sample of the ancestor (N2 or PB306) was thawed onto an agar plate (G0,

left) and grown for a few days until the cryopreserved worms had reached the L4 larval

stage. L4-stage worms were transferred singly to new NGM agar plates; those were the

founders of the MA lines (MA.1, MA.2,...MA.n; n=100). At four-day intervals (one

generation), a single immature worm was picked at random onto a new plate. This

procedure was repeated until the experiment reached Gmax=250 generations. Individual

MA lines experienced fewer than Gmax generations, Gave=238 generations. The different

colored worms represent (the set of) unique mutations fixed in each MA line, which

contribute to the among-line (mutational) variance.

Figure 2. Recall rates by genotype, SV type, and SV size. N2 has the highest recall rate

across SV types and sizes. Small deletions (100 bp) have the highest recall rates,

comparable to 1-kb and 10-kb deletions. Recall rates of insertions drop sharply beyond 100

bp. Recall rate decreases with increasing SV size and genetic divergence from N2, a trend

also observed in PB306-derived MA lines (not shown). Inversions have the lowest recall

rates. QX1211, the most divergent genotype from N2, is the most challenging for SV

detection. Minor horizontal jitter was added to the x-axis to improve the visual separation of

overlapping points.

Figure 3. Distribution of SVs within and among chromosomes in wild isolates (density shown

in blue) and MA lines (vertical black lines). The overall distributions between these two

groups are not significantly different (goodness-of-fit test; See text for details).

Figure 4. (Top) A schematic representation of the variants identified at the locus (I: 1567493)

in the pedigree of MA lines. PBSV initially classified the 1.9 kb insertion and the 3.1 kb

insertion as two independent mutations.  However, a deeper analysis—accounting for

sequencing errors—reveals that a 1.2 kb segment was deleted within the inserted sequence

in the ancestor, leading to the observed 1.9 kb insertion.

(Bottom) IGV snapshot of the 1.2 kb deletion in PB306-derived MA 445.  The IGV snapshot

shows pseudo-read alignments, which are assembled contigs representing longer

sequences.  Some pseudo-reads fail to fully capture the inserted segment and appear as

soft-clipped sequences.  If the inserted segment is not entirely covered within the pseudo-

read, it is clipped from the alignment, appearing as base calls (colored bars) instead of clean

(gray bars), fully aligned sequences.  Some insertion sequences are misaligned because of

the repetitive nature of the locus; however, all pseudo-reads carry the insertion.

Supplementary Figure S6 presents the same analysis using error-prone PacBio raw reads.

References

Andersen EC, Gerke JP, Shapiro JA, Crissman JR, Ghosh R, et al. 2012. Chromosome-scale selective
723            sweeps shape Caenorhabditis elegans genomic diversity. Nature Genetics 44:285-U283.
724    Audano PA, Sulovari A, Graves-Lindsay TA, Cantsilieris S, Sorensen M, et al. 2019. Characterizing the
725            Major Structural Variant Alleles of the Human Genome. Cell 176:663-+.
726    Baer CF, Phillips N, Ostrow D, Avalos A, Blanton D, et al. 2006. Cumulative effects of spontaneous
727            mutations for fitness in caenorhabditis: Role of genotype, environment and stress. Genetics
728            174:1387-1395.
729    Baer CF, Shaw F, Steding C, Baurngartner M, Hawkins A, et al. 2005. Comparative evolutionary
730            genetics of spontaneous mutations affecting fitness in rhabditid nematodes. Proceedings of
731            the National Academy of Sciences of the United States of America 102:5785-5790.
732    Bartenhagen C, Dugas M. 2013. RSVSim: an R/Bioconductor package for the simulation of structural
733            variations. Bioinformatics 29:1679-1681.
734    Bateman JR, Johnson JE. 2022. Altering enhancer-promoter linear distance impacts promoter
735            competition in cis and in trans. Genetics 222.
736    Beyter D, Ingimundardottir H, Oddsson A, Eggertsson HP, Bjornsson E, et al. 2021. Long-read
737            sequencing of 3,622 Icelanders provides insight into the role of structural variants in human
738            diseases and other traits. Nature Genetics 53:779-+.
739    Böndel KB, Kraemer SA, Samuels T, McClean D, Lachapelle J, et al. 2019. Inferring the distribution of
740            fitness effects of spontaneous mutations in *Chlamydomonas reinhardtii*. PLoS Biology
741            17:e3000192.
742    Boyle EA, Li YI, Pritchard JK. 2017. An Expanded View of Complex Traits: From Polygenic to
743            Omnigenic. Cell 169:1177-1186.
744    Bush Z, Naftaly A, Dinwiddie D, Albers C, Hillers K, et al. 2025. Transposable elements and
745            heterochromatic regions are enriched for structural variation and sequence divergence in
746            the genome of wild-type Caenorhabditis elegans. G3-Genes Genomes Genetics 15.
747    Chaisson MJ, Tesler G. 2012. Mapping single molecule sequencing reads using basic local alignment
748            with successive refinement (BLASR): application and theory. Bmc Bioinformatics 13:238.
749    Chaisson MJP, Wilson RK, Eichler EE. 2015. APPLICATIONS OF NEXT-GENERATION SEQUENCING
750            Genetic variation and the de novo assembly of human genomes. Nature Reviews Genetics
751            16:627-640.
752    Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, et al. 2012. A program for annotating and
753            predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of
754            Drosophila melanogaster strain w(1118); iso-2; iso-3. Fly 6:80-92.
755    Collins RL, Brand H, Karczewski KJ, Zhao X, Alfoldi J, et al. 2020. A structural variation reference for
756            medical and population genetics. Nature 581:444-+.
757    Conrad DF, Pinto D, Redon R, Feuk L, Gokcumen O, et al. 2010. Origins and functional impact of copy
758            number variation in the human genome. Nature 464:704-712.
759    Crombie TA, Rajaei M, Saxena AS, Johnson LM, Saber S, et al. 2024. Direct inference of the
760            distribution of fitness effects of spontaneous mutations from recombinant inbred
761            *Caenorhabditis elegans* mutation accumulation lines. Genetics 228.
762    Diesh C, Stevens GJ, Xie P, De Jesus Martinez T, Hershberg EA, et al. 2023. JBrowse 2: a modular
763            genome browser with views of synteny and structural variation. Genome Biology 24:74.
764    Eyre-Walker A, Keightley PD. 2007. The distribution of fitness effects of new mutations. Nature
765            Reviews Genetics 8:610-618.
766    Fisher RA. 1922. On the Dominance Ratio. Proceedings of the Royal Society of Edinburgh 42:321-341.
767    Friedman LS, Ostermeyer EA, Szabo CI, Dowd P, Lynch ED, et al. 1994. CONFIRMATION OF BRCA1 LAY
768            ANALYSIS OF GERMLINE MUTATIONS LINKED TO BREAST AND OVARIAN-CANCER IN 10
769            FAMILIES. Nature Genetics 8:399-404.

Gilbert KJ, Zdraljevic S, Cook DE, Cutter AD, Andersen EC, et al. 2021. The distribution of mutational
effects on fitness in *Caenorhabditis elegans* inferred from standing genetic variation.
Genetics 220.
Harvey WT, Ebert P, Ebler J, Audano PA, Munson KM, et al. 2023. Whole-genome long-read
sequencing downsampling and its effect on variant-calling precision and recall. Genome
Research 33:2029-2040.
Hénault M, Marsit S, Charron G, Landry CR. 2024. The genomic landscape of transposable elements
in yeast hybrids is shaped by structural variation and genotype-specific modulation of
transposition rate. Elife 12.
Hunter JD. 2007. Matplotlib: A 2D Graphics Environment. Computing in Science & Engineering 9:90-
95.
Hurles ME, Dermitzakis ET, Tyler-Smith C. 2008. The functional impact of structural variation in
humans. Trends in Genetics 24:238-245.
James J, Kastally C, Budde KB, González-Martínez SC, Milesi P, et al. 2023. Between but not within
species variation in the Distribution of Fitness Effects. Mol Biol Evol 13.
Karayiorgou M, Simon TJ, Gogos JA. 2010. 22q11.2 microdeletions: linking DNA structural variation
to brain dysfunction and schizophrenia. Nature Reviews Neuroscience 11:402-416.
Kim BY, Huber CD, Lohmueller KE. 2017. Inference of the Distribution of Selection Coefficients for
New Nonsynonymous Mutations Using Large Samples. Genetics 206:345-361.
Konrad A, Flibotte S, Taylor J, Waterston RH, Moerman DG, et al. 2018. Mutational and
transcriptional landscape of spontaneous gene duplications and deletions in *Caenorhabditis*
*elegans*. Proceedings of the National Academy of Sciences 115:7386-7391.
Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, et al. 2017. Canu: scalable and accurate long-
read assembly via adaptive *k*-mer weighting and repeat separation. Genome Research
27:722-736.
Kousathanas A, Keightley PD. 2013. A comparison of models to Infer the distribution of fitness
effects of new mutations. Genetics 193:1197-1208.
Larrichia KM, Zdraljevic S, Cook DE, Andersen E. 2017. Natural variation in the distribution and
abundance of transposable elements across the *Caenorhabditis elegans* species. Molecular
Biology and Evolution.
Lee D, Zdraljevic S, Stevens L, Wang Y, Tanny RE, et al. 2021. Balancing selection maintains hyper-
divergent haplotypes in Caenorhabditis elegans. Nat Ecol Evol 5:794-807.
Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34:3094-3100.
Loegler V, Thiele P, Teyssonnière E, Tsouris A, Brach G, et al. 2025. From genotype to phenotype
with 1,086 near telomere-to-telomere yeast genomes. Nature.
López-Cortegano E, Chebib J, Jonas A, Vock A, Künzel S, et al. 2025. The rate and spectrum of new
mutations in mice inferred by long-read sequencing. Genome Research 35:43-54.
Lopez-Cortegano E, Craig RJ, Chebib J, Balogun EJ, Keightley PD. 2023. Rates and spectra of de novo
structural mutations in Chlamydomonas reinhardtii. Genome Research 33:45-60.
Marshall CR, Noor A, Vincent JB, Lionel AC, Feuk L, et al. 2008. Structural variation of chromosomes
in autism spectrum disorder. American Journal of Human Genetics 82:477-488.
Meyerson M, Pellman D. 2011. Cancer Genomes Evolve by Pulverizing Single Chromosomes. Cell
144:9-10.
Nattestad M, Goodwin S, Ng K, Baslan T, Sedlazeck FJ, et al. 2018. Complex rearrangements and
oncogene amplifications revealed by long-read DNA and RNA sequencing of a breast cancer
cell line. Genome Research 28:1126-1135.
Nattestad M, Schatz MC. 2016. Assemblytics: a web analytics tool for the detection of variants from
an assembly. Bioinformatics 32:3021-3023.
Noyes MD, Harvey WT, Porubsky D, Sulovari A, Li RY, et al. 2022. Familial long-read sequencing
increases yield of *de novo* mutations. American Journal of Human Genetics 109:631-646.

Payer LM, Steranka JP, Yang WR, Kryatova M, Medabalimi S, et al. 2017. Structural variants caused
821            by Alu insertions are associated with risks for many human diseases. Proceedings of the
822            National Academy of Sciences of the United States of America 114:E3984-E3992.
823     Pinto D, Pagnamenta AT, Klei L, Anney R, Merico D, et al. 2010. Functional impact of global rare copy
824            number variation in autism spectrum disorders. Nature 466:368-372.
825     Porubsky D, Dashnow H, Sasani T, Logsdon G, Hallast P, et al. 2025. Human de novo mutation rates
826            from a four-generation pedigree reference. Nature 643.
827     Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
828            Bioinformatics 26:841-842.
829     Rajaei M, Saxena AS, Johnson LM, Snyder MC, Crombie TA, et al. 2021. Mutability of mononucleotide
830            repeats, not oxidative stress, explains the discrepancy between laboratory-accumulated
831            mutations and the natural allele-frequency spectrum in *C. elegans*. Genome Research
832            31:1602-1613.
833     Robinson JT, Thorvaldsdóttir H, Wenger AM, Zehir A, Mesirov JP. 2017. Variant Review with the
834            Integrative Genomics Viewer. Cancer Research 77:e31-e34.
835     Rockman MV, Kruglyak L. 2009. Recombinational Landscape and Population Genomics of
836            Caenorhabditis elegans. Plos Genetics 5.
837     Sanders SJ, Ercan-Sencicek AG, Hus V, Luo R, Murtha MT, et al. 2011. Multiple Recurrent De Novo
838            CNVs, Including Duplications of the 7q11.23 Williams Syndrome Region, Are Strongly
839            Associated with Autism. Neuron 70:863-885.
840     Saxena AS, Salomon MP, Matsuba C, Yeh SD, Baer CF. 2019. Evolution of the Mutational Process
841            under Relaxed Selection in Caenorhabditis elegans. Molecular Biology and Evolution 36:239-
842            251.
843     Schloissnig S, Pani S, Ebler J, Hain C, Tsapalou V, et al. 2025. Structural variation in 1,019 diverse
844            humans based on long-read sequencing. Nature 644:442-452.
845     Sedlazeck FJ, Rescheneder P, Smolka M, Fang H, Nattestad M, et al. 2018a. Accurate detection of
846            complex structural variations using single-molecule sequencing. Nature Methods 15:461-
847            468.
848     Sedlazeck FJ, Rescheneder P, Smolka M, Fang H, Nattestad M, et al. 2018b. Accurate detection of
849            complex structural variations using single-molecule sequencing. Nature Methods 15:461-+.
850     Stiernagle T. 2006. Maintenance of *C. elegans*. In: Community T*Ce*R, editor. Wormbook.
851     Stransky N, Cerami E, Schalm S, Kim JL, Lengauer C. 2014. The landscape of kinase fusions in cancer.
852            Nature Communications 5.
853     Sung W, Ackerman MS, Miller SF, Doak TG, Lynch M. 2012. Drift-barrier hypothesis and mutation-
854            rate evolution. Proceedings of the National Academy of Sciences 109:18488-18492.
855     Tataru P, Mollion M, Glemin S, Bataillon T. 2017. Inference of Distribution of Fitness Effects and
856            Proportion of Adaptive Substitutions from Polymorphism Data. Genetics 207:1103-1119.
857     Tenesa A, Navarro P, Hayes BJ, Duffy DL, Clarke GM, et al. 2007. Recent human effective population
858            size estimated from linkage disequilibrium. Genome Res 17:520-526.
859     Teterina A, Willis J, Lukac M, Jovelin R, Cutter A, et al. 2023. Genomic diversity landscapes in
860            outcrossing and selfing Caenorhabditis nematodes. Plos Genetics 19.
861     Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, et al. 2020. SciPy 1.0: fundamental
862            algorithms for scientific computing in Python. Nature Methods 17:261-272.
863     Walsh T, McClellan JM, McCarthy SE, Addington AM, Pierce SB, et al. 2008. Rare structural variants
864            disrupt multiple genes in neurodevelopmental pathways in schizophrenia. Science 320:539-
865            543.
866     Wang QL, Dhindsa RS, Carss K, Harper AR, Nag A, et al. 2021. Rare variant contribution to human
867            disease in 281,104 UK Biobank exomes. Nature 597:527-+.
Waskom ML. 2021. seaborn: statistical data visualization. Journal of Open Source Software 6:3021.

Weirather JL, de Cesare M, Wang Y, Piazza P, Sebastiano V, et al. 2017. Comprehensive comparison
of Pacific Biosciences and Oxford Nanopore Technologies and their applications to
transcriptome analysis. F1000Res 6:100.
Weiss LA, Shen YP, Korn JM, Arking DE, Miller DT, et al. 2008. Association between microdeletion
and microduplication at 16p11.2 and autism. New England Journal of Medicine 358:667-675.
Wenger AM, Peluso P, Rowell WJ, Chang P-C, Hall RJ, et al. 2019. Accurate circular consensus long-
read sequencing improves variant detection and assembly of a human genome. Nature
Biotechnology 37:1155-1162.
Wooster R, Bignell G, Lancaster J, Swift S, Seal S, et al. 1995. IDENTIFICATION OF THE BREAST-
CANCER SUSCEPTIBILITY GENE BRCA2. Nature 378:789-792.
Yi K, Ju YS. 2018. Patterns and mechanisms of structural variations in human cancer. Experimental
and Molecular Medicine 50.
Zhang H, Jain C, Aluru S. 2020. A comprehensive evaluation of long read error correction methods.
Bmc Genomics 21:889.
Zhou X, Pan J, Wang YH, Lynch M, Long HA, et al. 2023. De Novo structural variations of *Escherichia*
*coli* detected by Nanopore long-read sequencing. Genome Biology and Evolution 15.

Variant Density by Chromosome