<u>Common non-antibiotic drugs enhance selection for antimicrobial resistance</u>
<u>in mixture with ciprofloxacin</u>

April Hayes[1*], Lihong Zhang[1], Jason Snape[2], Ed Feil[3], Barbara Kasprzyk-Hordern[4],
William H Gaze[1], Aimee K Murray[1]

1 European Centre for Environment and Human Health, University of Exeter,
Penryn, UK

2 Department of Environment and Geography, University of York, UK

3 Department of Life Sciences, University of Bath, Bath, UK

4 Department of Chemistry, University of Bath, Bath, UK

*Corresponding author – a.hayes4@exeter.ac.uk

<u>Abstract</u>

Antimicrobial resistance (AMR) is a major health concern, and a range of antibiotic
concentrations and non-antibiotic agents can select for AMR. Selection for AMR is
often investigated using single compounds, however, in the natural environment and
the human body, pharmaceuticals will be present as mixtures, including both non-
antibiotic drugs (NADs), and antibiotics. Here, we assessed the effects of one of
three NADs in combination with ciprofloxacin, a commonly used antibiotic that is
often found at concentrations in global freshwaters sufficiently high to select for
AMR. We used a combination of growth assays and qPCR to determine selective
concentrations of mixtures and used metagenome sequencing to identify changes to
the resistome and community composition. The selective concentration of
ciprofloxacin was reduced in mixtures, and some changes in selection for specific
genes or species were observed. Results show that NADs and ciprofloxacin are
more selective than ciprofloxacin alone and these mixtures can cause distinct
changes to the community composition. This indicates that future work should on the

maintenance and spread of AMR should consider combinations of antibiotics and
NADs.

<u>Introduction</u>

Antimicrobial resistance (AMR) is a global health threat, with 1.27 million deaths in
2019 directly caused by antibiotic resistant bacterial infections [1]. Traditionally,
resistance to antibiotics has been determined by identifying concentrations that
inhibit growth. However, research shows that low concentrations of antibiotics can
select for antibiotic resistance in both single species [2, 3], and in bacterial
communities [4-9]. Additionally, other non-antibiotic compounds can co-select for
antibiotic resistance, including metals and biocides [10, 11]. Non-antibiotic drugs
(NADs) have previously been shown to reduce bacterial growth [12, 13], increase
horizontal gene transfer rates [14-17], and select for antibiotic resistance [18-20] in
single species experiments at therapeutic concentrations. There is some evidence
suggesting that NADs at lower, more environmentally relevant concentrations may
not select, or select less strongly for AMR, in both single species and mixed
communities [21, 22]. We previously tested three NADs commonly found in the
environment - diclofenac, metformin and 17-β-estradiol – on their selective potential
for AMR [22]. Diclofenac, a non-steroidal anti-inflammatory drug [32], is one of the
five most frequently detected pharmaceuticals in the aquatic environment [23].
Metformin is used in front-line diabetes treatments [24] and 17-β-estradiol is a
naturally produced hormone but is also used in hormone replacement therapy [25,
26]. Diclofenac, metformin and 17-β-estradiol did not strongly select for antibiotic
resistance genes within a bacterial community, or affect bacterial diversity [22].
However, they did have antimicrobial activity in terms of impact on growth rate and
additionally had some effects on metal resistance gene abundance/diversity [22].

Pharmaceuticals are present in both the human body and in the aquatic environment
at a range of concentrations [24, 27-30]. NADs will be present alongside antibiotics
in these environments as both simple and complex mixtures of multiple
pharmaceuticals [31]. Environmentally relevant concentrations of ciprofloxacin can
select for AMR [2, 9] and previous environmental risk assessments have shown that

there is a risk of AMR selection by ciprofloxacin in various wastewater environments,
even in high income countries [32].

Several studies have studied the effects of mixtures of NADs and antibiotics with a
specific focus on their capacity to reduce or completely inhibit bacterial growth or
increase antibiotic susceptibility [33, 34]. For example, diclofenac and metformin
have been shown to act both synergistically and antagonistically with a range of
antibiotics across a range of classes and species [35] including the pathogen
Pseudomonas aeruginosa. Diclofenac can also increase the inhibitory activity of
ciprofloxacin in *Proteus mirabilis* [36]. Alternatively, metformin has been shown to
increase the inhibitory activity of tetracycline antibiotics, and restored tetracycline
susceptibility in a resistant *Escherichia coli* strain [37]. Overall, there is not enough
evidence to conclusively suggest mechanistic insights into combinations of these
pharmaceuticals. These results may be species specific, or related to altered gene
expression, and there is less evidence when considering complex communities.

Furthermore, most research has focused on testing the effects of mixtures of
antibiotics. Work in this area has illustrated within-species variability in responses to
antibiotic mixtures [38], and shown that the responses of single species do not
predict the responses of a more complex community [39]. Furthermore, there is
evidence to suggest that complex microbial communities may be more resilient to
mixtures than individual species [39], which also occurs with single antibiotic
compounds [40]. Additionally, research has shown that when antimicrobials are
combined in complex mixtures, a variety of interactions can occur. This includes
increased and decreased evolution of resistance [41-43]. Overall, mixtures are likely
to lead to different selection dynamics in bacterial communities compared to single
compounds alone due to increased variability in both species and genes.

In this study, we aimed to understand the effects of simple mixtures in a complex
microbial community, since this is first step to understanding complex mixture
effects. We experimentally spiked single concentrations of diclofenac, metformin,
and 17-β-estradiol alongside a range of ciprofloxacin concentrations to see if this
affected the minimal selective concentration of ciprofloxacin. Firstly, we determined if
there was a significant reduction in the growth of the community in the mixture
compared to ciprofloxacin alone, and secondly, if the minimal selective concentration

of ciprofloxacin changed in the presence of the NAD. We used *intl1* as the selective
concentration endpoint, since this has previously been shown to increase with
ciprofloxacin selection [9], and *intl1* has been suggested as a proxy for antimicrobial
resistance acquisition in environmental monitoring [44]. Finally, we determined
whether the community resistome or composition changed in the mixtures using
metagenome sequencing.

Methods

Pharmaceuticals

Diclofenac (Sigma Aldritch), metformin (Enzo), 17-β-estradiol (Sigma Aldritch), and
ciprofloxacin (Sigma Aldritch) were acquired and dissolved in water, water, ethanol,
and 0.8mol HCl and 1.2mL water respectively and filter sterilised. Aliquots were kept
at -20^{o}C for up to two weeks before use. Stock concentrations of pharmaceuticals
were diluted in filter sterilised water to concentrations for use.

Wastewater influent

Wastewater influent was collected from Falmouth (UK) wastewater treatment plant,
in June 2022. The wastewater was collected in clean 1L glass Duran bottles and
processed the same day. Wastewater was mixed 1:1 with 40% glycerol and kept
frozen at -70^{o}C until use.

Growth Assays

Combinations of ciprofloxacin and either diclofenac, metformin, or 17-β-estradiol
were tested across a range of ciprofloxacin concentrations with a spiked
concentration of the NAD (diclofenac: 50µg/L and 25µg/L, metformin: 26µg/L and
13µg/L, 17-β-estradiol: 24.8µg/L and 12.4µg/L). The higher set of these
concentrations had previously been shown to significantly reduce growth compared
to a no-NAD control, and the lower concentration did not significantly reduce growth
compared to a no-NAD control, within the same experimental set up [22].

A 96 well plate was filled with 180µL Iso-Sensitest broth (Oxoid). For the
ciprofloxacin gradient, 180µL Iso-Sensitest broth with 15.6µg/L ciprofloxacin was
added to the top 12 wells of the plate, and 180µL was serially diluted down the plate,

leaving one row as a no-antibiotic control. The higher NAD concentration was added
to four columns, the lower NAD concentration to four columns, and the remaining
four columns were left as a no-NAD control (Figure 1). Wastewater influent was
thawed, and washed with 0.85% NaCl twice to remove contaminants and nutrient
carry over, and 10µL of this was inoculated into each well. The plate was sealed with
a MicroAmp Optical seal and optical density was measured at 600nm (OD600) every
ten minutes for 24 hours in a BioTek plate reader (Agilent), with five seconds of
shaking at 180 rpm every ten minutes.

Figure 1. 96 well plate layout for the growth assays for mixture experiments.

Selection Experiments for qPCR analysis

Selection experiments were carried out across a ciprofloxacin gradient, with the
addition of single spiked NAD concentrations. Firstly, the lowest observed effect
concentration (LOEC) of ciprofloxacin was determined. Ciprofloxacin was tested at
40µg/L, 20µg/L, 10µg/L, 5µg/L, and 2.5µg/L and 0 µg/L. This range was informed by
a previous study where the LOEC was calculated to be 15.6µg/L [9]. Then, mixture
experiments using the NADs were performed over this ciprofloxacin concentration
range, but including either diclofenac spiked at 50µg/L, metformin at 26µg/L or 17-β-
estradiol at 24.8µg/L [22].

To set up the experiments, wastewater influent was thawed and washed twice with
0.85% NaCl and inoculated with 10% vol/vol into Iso-Sensitest broth. This inoculated
broth was separated into 30mL aliquots, which were spiked with the ciprofloxacin
and NAD concentrations. These 30mL aliquots were then separated into 5mL

microcosms. Day zero samples were taken at this time point. For this, 1mL of each
microcosm was taken, centrifuged at 2100rpm for two minutes and the pellet
resuspended with 20% glycerol. Samples were kept at -70^{o}C until use. The
microcosms were incubated at 37^{o}C with 180rpm shaking, with transfers into fresh
media with fresh NAD and ciprofloxacin daily for seven days. On day seven, samples
were taken by mixing 0.5mL culture 1:1 with 40% glycerol and kept at -70^{o}C until
use.

QPCR analysis

QPCR was performed using day zero and day seven samples to determine effects
on selection after the weeklong experiment. DNA was extracted using a Qiagen
DNeasy UltraClean Microbial Kit. All steps were carried out to manufacturer's
instructions with the initial centrifugation step extended to one minute. Extracted
DNA was diluted 5X with TE and stored at 4^{o}C before use. QPCR was performed
using *intI1* primers and standardised to *16S* rRNA copy number using a Thermo
Fisher QuantStudio 7 Real-Time PCR machine (Thermo Fisher). Prevalence was
calculated by dividing the *intI1* copy number by the 16S rRNA copy number*.* The
reaction mix included 10µL SYBR MasterMix with ROX and SYBR (PrimerDesign),
1µL forward primer (Integrated DNA Technologies), 1µL reverse primer (Integrated
DNA Technologies), 0.2µL Bovine Serum Albumin and 2.8µL nuclease free water
(Ambion). The cycling protocol was as follows – 120 second hold at 95^{o}C, 50 rounds
of cycling with 10 seconds at 95^{o}C for denaturation and 60 seconds at 60^{o}C for data
collection. Only runs with an efficiency of 90-11%, and an R^2 of greater than 0.99
were used in analyses. Primer and gblock sequences are presented in
Supplementary Table 1.

Metagenome Sequencing

For Illumina metagenome sequencing, day seven samples were thawed, and DNA
extracted using a Qiagen DNeasy UltraClean Microbial Kit, with all steps carried out
to the manufacturer's instructions save for two exceptions. The initial centrifugation
step was elongated to 2 minutes, and the centrifugation of the PowerBead tubes was
increased to one minute at 12,000G. Extracted DNA was purified using a standard
RNase A and standard Ampure XP bead protocol. DNA was eluted in 10mM Tris-

HCl and stored at 4^{o}C until being sent for sequencing. NEB PCR-free library prep
was carried out by the Exeter Sequencing Centre prior to sequencing using a
NovaSeq SP to a depth of up to 20GB per sample.

Metagenome Analyses

Trimmed reads from the Exeter Sequencing Service were used in all analyses. All
reads were checked for quality using FastQC and MultiQC [45]. AMR++ was used to
process the reads [46]. Low quality reads and reads mapping to host (human) were
removed. For the resistome analysis, reads were aligned to the MEGARES 3.0
database [46], which includes multiple resistome databases including BacMet [47],
ResFinder [48], and CARD [49]. For the microbiome analysis, *kraken2* [50] was used
to identify taxonomy as part of the AMR++ pipeline using the minikraken database
[51, 52].

Outputs from these pipelines were input into R, and converted into phyloseq objects
using *phyloseq* v1.48.0 [53]. Reads were normalised using *metagenomeSeq* v1.46.0
[54, 55]. Relative abundances (proportions) were then created using these
normalised data.

Data Analysis

All statistical analyses were carried out in R version 4.4.1 [56]. All figures were
generated using *ggplot2* 3.5.1 [57] and used *MetBrewer* v0.2.0 [58]. For all models,
the most parsimonious model was used, determined by sequentially deleting terms
and comparing model fits using X^2 tests. Only those models with residuals fitting
assumptions were used. Fit of residuals were checked using *DHARMa* v0.4.6 [59].

Growth Analyses

To determine the minimal selective concentrations or LOECs of the mixtures using
growth, a method previously used was applied [60]. In summary, the time point in
exponential phase with the largest dose response was determined using either
Spearman's or Pearson's correlation test, as determined by the normality fit of the
data. Then, at this time point, a Dunn's test (*dunn.test* v1.3.5) was used to identify
which concentrations significantly differed from the control growth (i.e. no
ciprofloxacin and no NAD). The lowest concentration that was significantly different
to the control was determined as the LOEC

To determine total growth capacity or productivity, total area under the curve (AUC)
was used. AUC was determined using the *growthcurver* v0.3.1 package [61]. The
AUC from exponential growth phase was used in linear mixed effect models using
lme4 v1.1.31 [62], with concentration of ciprofloxacin and treatment of NAD as fixed
effects, and microcosm as a random effect. Pairwise comparisons were determined
using *emmeans* v1.8.2 [63], and p values adjusted for multiple comparisons using
false discovery rate.

QPCR Analyses

Intl1 prevalence was calculated by dividing the *intl1* quantity by the 16S rRNA
quantity. These prevalences were used to calculate LOECs. LOECs were
determined using linear mixed effect models, with time and treatment as fixed
effects, and microcosm was included as a random effect. Pairwise comparisons
were calculated as above. LOECs were determined to be the lowest concentration
that showed a significantly increased *intl1* prevalence compared to the day seven
control.

Diversity analyses

Alpha diversity or richness was used to identify the total number of taxa and genes
present in each sample. *Phyloseq* was used to estimate Shannon's index [64], which
was used to identify the evenness of taxa and genes. Tests for significant differences
in treatments for alpha diversity were tested using linear models, with richness as the
response variable, and mixture type and ciprofloxacin concentration as explanatory
variables. Pairwise comparisons were calculated as above. For beta diversity
analyses, Bray-Curtis ordinations were calculated using *vegan* v2.6.6.1 [65] for both
the resistome and the taxonomy. Changes to the diversity were determined using
Analysis of Similarity (ANOSIM) tests.

Changes to resistance gene classes or species

Kruskal-Wallis tests were used to identify differences in gene classes, or taxonomic
order by mixture type, using the normalised gene counts. Genes or taxonomic orders
that differed significantly in at least one treatment were tested to determine if mixture
type and/or ciprofloxacin concentration significantly affected gene abundance by
linear models and ANOVAs.

Log2-fold change in resistance genes or species

For both the resistome and the community taxonomy, significant log2-fold changes
between mixtures and the ciprofloxacin alone treatments were identified using
DESeq2 v 1.44.0 [66].

<u>Results</u>

NAD and ciprofloxacin mixtures significantly reduced community productivity

To test differences in community productivity, we tested changes to the AUC, which
can be considered a proxy for overall productivity or growth capacity. The AUC of the
mixtures were compared to in-plate AUC data of ciprofloxacin alone. Firstly, none of
the lower concentrations of the NADs significantly reduced growth compared to the
no-NAD control ($p<0.05$). However, all three NADs at the higher concentration in
mixture with ciprofloxacin significantly reduced the productivity of communities
compared to growth in the no NAD control (Figure 2), across the entire ciprofloxacin
concentration gradient ($p<0.001$). We also found that as expected, total AUC
reduced with increased ciprofloxacin concentration ($p<0.001$). More details of the
outputs for these models are detailed in the Supplementary File, section 1.

257
Figure 2. Area under the curve during exponential phase across a
ciprofloxacin concentration gradient in mixture with A) diclofenac, B)

metformin, and C) 17-β-estradiol. Pale points indicate each individual replicate,
larger circles indicate the mean at each concentration. Model line shown,
including upper and lower model predictions of 95% confidence intervals in
shaded ribbon.

Secondly, we determined the growth-based LOEC for the mixtures containing the
higher NAD concentrations, using a low-cost method previously published [60].
Previous work has indicated that reduction in growth is the strongest indicator of
selection for AMR [67]. We found that all mixtures reduced the estimated selective
concentration of ciprofloxacin. The diclofenac-ciprofloxacin mixture reduced the
LOEC of ciprofloxacin from 3.7µg/L to 0.12µg/L, a 32-fold decrease. The metformin-
ciprofloxacin mixture reduced the LOEC of ciprofloxacin from 0.98µg/L to 0.24µg/L, a
4-fold decrease. The 17-β-estradiol-ciprofloxacin mixture reduced the LOEC of
ciprofloxacin from 1.95µg/L to 0.24µg/L, an 8-fold decrease.

NAD mixtures reduce the selective concentration of ciprofloxacin for *intI1*

Next, we determined whether there was specific selection for the commonly used
AMR marker *intI1.* We hypothesised that the mixtures of NADs and ciprofloxacin
would have increased selectivity compared to ciprofloxacin alone.

Ciprofloxacin alone selected for *intI1* at 40µg/L (Figure 3A). We found that *intI1*
prevalence significantly increased with treatment (treatment main effect, X^2=18.35,
df=4, p=0.0011) and that only 40µg/L ciprofloxacin had a day seven prevalence that
was significantly greater than the prevalence at day seven control population
(p=0.012).

We then tested the mixtures of NADs and ciprofloxacin and found that they all
reduced the selective concentration to 10µg/L (Figure 3). In all cases, the
concentration of *intI1* either increased or stayed constant with time (Supplementary
Figure 1), so we only show day seven data in the following plots.

Firstly, 50µg/L diclofenac in mixture with ciprofloxacin (Figure 3B) reduced the
selective concentration for *intI1*, with the communities exposed to 10µg/L
(p=0.0085), 20µg/L (p=0.028) and 40µg/L (p=0.0085) showing significantly increased
intI1 prevalence compared to the control. Secondly, metformin at 26µg/L reduced the
selective concentration of ciprofloxacin to 10µg/L (Figure 3C). We see that 10µg/L

(p=0.0016), and 20µg/L (p=0.024) in the mixture significantly increased *intI1*
prevalence, however this increase was not seen with 40µg/L ciprofloxacin (p=0.063).
Finally, we found that 24.4µg/L 17-β-estradiol-ciprofloxacin mixture also reduced the
selective concentration to 10µg/L (p<0.0001), with communities exposed to 20µg/L
(p<0.0001) and 40µg/L (p<0.0001) also showing significantly increased *intI1*
prevalence compared to the ciprofloxacin only community.

Despite mixtures of NADs and ciprofloxacin selecting for *intI1* at a lower ciprofloxacin
concentration, the overall strength of selection in the mixtures was lower than that
with ciprofloxacin alone, indicating some suppression of selection within the
mixtures.

Figure 3. *IntI1* prevalence as a function of ciprofloxacin concentration in both ciprofloxacin alone (A) or in combination with diclofenac (B), metformin (C), or 17-β-estradiol (D). Five biological replicates shown. Significant differences of day seven prevalences are shown from pairwise comparisons to the day seven prevalence at 0µg/L. NS = non-significant, * = p<0.05, ** = p<0.01, *** = p<0.001.

Mixtures did not strongly alter microbiome or resistome diversity

Next, we analysed the metagenomes of the evolved communities to understand how the mixtures might have affected the community composition and the resistome. We hypothesised that there would likely be increases in gene abundances occurring at lower ciprofloxacin concentrations in the mixtures, since the mixtures appeared to be more selective. We also hypothesised that it was likely that the mixtures had selected for different species, and that the mixtures would have decreased richness and diversity, since the communities were exposed to multiple pharmaceutical stressors.

There were no significant differences in richness of the taxa present in each sample. However, there were differences in the richness of the AMR genes between the treatments (Figure 4). Exposure to the 17-β-estradiol mixture decreased resistome richness compared to the diclofenac treatment (p=0.021). This mixture also showed a non-significant decrease in richness compared to the ciprofloxacin only treatment (p=0.071). Exposure to the metformin mixture significantly reduced resistome richness compared to the diclofenac mixture (p=0.0087), and the ciprofloxacin alone treatment (p=0.021).

Figure 4. Resistome richness (number of observed AMR genes) of the evolved samples. Colour of the points indicates ciprofloxacin concentration. Asterisks indicate significance values where * = p<0.05, and ** = p<0.001.

There was no significant difference in the evenness of the resistome or the taxonomy between the mixture treatments. There was also no significant difference in the beta diversity of either the resistome or community composition (ANOSIM, p>0.05). (Supplementary Figures 2 and 3).

Mixtures significantly altered the abundance of three AMR genes

Next, we tested all genes within the resistome to see if they differed in at least one treatment and found that ten genes were significantly different in at least one sample (p<0.05) (Supplementary Table 2). Of these genes, six were significantly affected by mixture type either as main or interaction effect: *aph3-DPRIME, aph6, fecE, qnrB, tetA, tetQ,* and *tolC* (Figure 5). All details on the model outputs can be found in the Supplementary File in section 2.

Figure 5. Normalised abundance of all genes that were significantly different in at least one sample and were significantly affected by NAD presence. Colours indicate mixture treatments. Lines show best model fit, and the shaded areas show 95% confidence intervals. Points indicate the individual replicates.

FecE, *qnrB*, and *tolC* showed the largest changes in gene abundances. For these three genes, the same trends appeared with the metformin mixture showing the largest increase in gene abundances, followed by the diclofenac mixture. The 17-β-estradiol mixture either decreased in gene abundance (*fecE* and *tolC*) or showed the smallest increase (*qnrB*). For the other four genes, abundances changed very slightly and often had a unimodal response (e.g. *tetA*).

Mixtures significantly altered the abundances of multiple bacterial taxa

We tested whether any order of microorganism was significantly different in at least one treatment. Two orders showed strong alterations across either concentration or mixture type: Caulobacterales and Nitrososphaerales. Caulobacterales abundance in the community decreased after exposure to ciprofloxacin only but increased in abundance in all the mixture treatments (interaction effect: $F_{3,110}=3.71$, p=0.014) (Figure 6). Nitrososphaerales increased in abundance with ciprofloxacin concentration (concentration main effect: $F_{1,54}=17.52$, p=0.00011), and had a

decreased abundance in the mixtures compared to ciprofloxacin alone treatment
(mixture main effect: $F_{3,54}=5.81$, p=0.0016) (Figure 6).

Figure 6. Normalised abundance of Caulobacterales and Nitrososphaerales as
a function of ciprofloxacin concentration. Colours indicate mixture treatments.
Lines show the best model fit, and shaded areas show 95% confidence
intervals. Points represent the individual replicates.

Additionally, we grouped the community based on abundances greater than 10%. In
these high abundance genera, there were no clear trends in changes to these
genera, or phyla across the mixtures (Supplementary Figure 4). This indicates that
was no dominance effect, and the communities were mostly stable across
treatments.

However, over 100 species of microorganism had a log2-fold increase or decrease in
each of the mixtures compared to the ciprofloxacin alone (Supplementary Figure 5).
43 species that either significantly increased or decreased in all three of the mixtures
compared to the control (Figure 7). For these 43 species, the diclofenac and
metformin mixtures showed the opposite response to the 17-β-estradiol mixture, i.e.
if the diclofenac and metformin mixture showed an increased abundance of a
particular species, the 17-β-estradiol mixture showed a decrease, and vice versa.

Some of these species are known pathogens (e.g. *Pseudomonas. aeruginosa,* which

increased in the 17-β-estradiol mixture).

Figure 7. Log2-fold change of species for mixtures compared to ciprofloxacin

alone treatment. Those species with a positive change are coloured orange,

those with a negative are coloured blue.

<u>Discussion</u>

Here, we showed that addition of a NAD at a single spiked concentration can reduce

the selective concentration of ciprofloxacin in a mixed bacterial community. This is

despite previous work [22] indicating that these NADs have little to no selective effect
when on their own. Further, this work indicates that mixtures of NADs and antibiotics
might be significantly altering the composition of microbial communities and may
select for specific AMR genes. We base this on our observations that both growth-
based and qPCR-based effect concentrations reduced when NADs were present in
mixture with ciprofloxacin. In addition, across all the mixtures, the productivity of the
community decreased, indicating that these mixtures are significantly impairing
community growth capacity.

Sequencing indicated that mixtures of NADs and ciprofloxacin can select for AMR
genes, and can lead to changes to the community composition, although these
changes are NAD- and mixture-specific. The concentrations used in this study are
sub-MIC, and are much lower than would be used clinically, so the effects we
observed might be more profound at higher concentrations of either the antibiotic,
NAD, or both. Two of the AMR genes that were significantly altered by the mixtures
encode for part of a membrane transport system (*fecE:* an ABC iron transporter [68],
and *tolC:* a multi-drug efflux channel [69]). These efflux pump mechanisms may be
important in resistance in the metformin and diclofenac mixtures, but not in the 17-β-
estradiol mixture. Most intriguing is the finding that the log2-fold changes in species
seen differ between diclofenac/metformin mixtures, and the 17-β-estradiol mixture.
Coupled with the findings above relating to *fecE* and *tolC,* this might indicate that the
17-β-estradiol mixture is selecting for taxa that are intrinsically resistant or contain
unannotated resistance mechanisms. It might also indicate that the diclofenac and
metformin mixture is interacting with the community at a molecular level in a similar
way. Additionally, we can suggest that the mixtures are acting differently to
ciprofloxacin alone with regards to selection for or against different species, with the
Gram negative Caulobacterales selected for across the mixtures and negatively
selected against by ciprofloxacin alone. This order may be more adaptable towards
growth in antibiotic-NAD mixtures, although there is yet no clear evidence as to why.

As mentioned, previous work indicated that diclofenac can reduce the MIC of
ciprofloxacin [36], and the data presented here indicates that it can also reduce the
MSC of ciprofloxacin. The proposed mechanism of diclofenac is inhibition of DNA
synthesis [70]. As ciprofloxacin also interferes with DNA synthesis [71, 72], it is likely
the combined effects of the two compounds in mixture will significantly reduce

replication rate, reflected in the reduction in growth observed in this study. Diclofenac
at 10µg/L has also been demonstrated to increase mutation frequency of *E. coli* [19],
and to upregulate *sigB*, a gene involved in the stress response [73]. An increased
stress response can, in turn, increase mutation rates and integron activation [74, 75].
This might explain some of the *intI1* abundance increases observed in the mixture,
and the increases in *tolC* and *fecE* abundances.

Metformin has previously been shown to reduce the MIC of various antibiotics
(including ampicillin, doxycycline, and levofloxacin) [37, 76], and it has been
suggested that this could be due to the ability of metformin to disrupt cell membranes
[37]. This proposed mechanism may allow for increased ciprofloxacin influx into
cells, which may lead to selection for any resistance genes present. This is might
explain why the metformin mixture had the largest increase in *fecE, qnrB*, and *tolC*
abundance, since increased ciprofloxacin influx would exert a stronger selection
pressure. Metformin acted synergistically with ciprofloxacin, reducing the selective
concentration to 10µg/L compared to 40µg/L with ciprofloxacin alone (and it was not
selective when tested alone [22]). However, there was an antagonistic effect at
40µg/L. This antagonistic effect may be due to chemical interactions between the
two pharmaceuticals. This has been demonstrated to occur previously with
ciprofloxacin and zinc [77]. Future work could aim to understand if this effect holds
true for a larger range of concentrations, and whether these pharmaceuticals were
indeed interacting chemically.

There is no previous experimental evidence investigating the mixture effect of 17-β-
estradiol with an antibiotic. Therefore, these data are presented as novel. Results
suggest that 17-β-estradiol may act additively or synergistically with ciprofloxacin to
reduce bacterial growth and increase selection for AMR by reducing the minimal
selective concentration. In previous work, 17-β-estradiol has been shown to select
for *intI1* (from 7µg/L to 5400µg/L), and also selected for metal resistance genes [22].
In this study, the communities exposed to the 17-β-estradiol mixture often responded
in the opposite way to the diclofenac and metformin mixtures (for example, the
communities exposed to the 17-β-estradiol mixtures showed a decreased *tolC* and
fecE abundance). Perhaps these communities are relying upon a shift to more
tolerant species (as indicated by the fold change in various species), or increased
gene expression. Future work could aim to unpick changes to gene expression in

response to various NAD and antibiotic mixtures, which might illuminate the changes
identified here.

Overall, these results provide some concern for human health. These NADs might
be present in the human gut alongside antibiotics, particularly if patients with long-
term health conditions (such as diabetes, or those requiring hormone replacement
therapy) acquired an infection that required antibiotic treatment. The human gut
microbiome can take weeks to recover after antibiotic treatment, and some species
never recolonise [78]. In addition to this, after antibiotic treatment there is evidence
to suggest that there is an increase in AMR gene abundance in the human gut [79].
Therefore, mixtures of pharmaceuticals in the gut may increase the selective
potential of antibiotics, potentially increasing selection for AMR within the gut
microbiome. Additionally, of concern is that in the human gut, lower concentrations of
ciprofloxacin are required to induce the expression of class I integron [80] (which can
contain AMR gene cassettes), and mixtures of pharmaceuticals may exacerbate this.
In general, these data suggest that mixtures of NADs and antibiotics lead to
selection for AMR at lower concentrations than seen with antibiotics alone, and
selectively lead to changes in abundance of species, or specific resistance genes.
This could lead to increased selection, maintenance, and dissemination of antibiotic
resistance genes between species in the human gut, and increased shedding of
these genes and resistant bacteria into the environment.

Furthermore, these pharmaceuticals, alongside others, will be present in the
freshwater aquatic, or wastewater environments as micropollutants [23, 31], where
they may be affecting growth of the natural communities, and impacting ecosystem
functioning. This includes in wastewater treatment plants where they may directly
impact wastewater treatment plant functioning if these pharmaceuticals are affecting
microorganisms involved in sludge digestion. In our study we found a short-term
overall reduction in community growth, but a longer-term increase in
Nitrososphaerales abundance, archaea that may play a role in the nitrogen and
carbon cycles. Further investigation is needed to determine whether the growth
effects are short- or long-term, and what this might mean for ecosystems.

These mixtures may lead to increased selection, maintenance, and dissemination of
antibiotic resistance genes throughout environmental compartments, or increased

input of resistant bacteria to the receiving waters. This then may lead to increased
risk of infection or colonisation with resistant strains to people and animals
interacting with these environments (e.g., surfers and swimmers in the sea [81]).

Taken together, these data confirm findings that mixtures of NADs and antibiotics
can be more selective than the antibiotic or NAD alone. This is of particular concern
since previous work [22] has indicated that two of these NADs did not select for AMR
in a similar community, and so may have been discounted as potentially selective
agents. Many compounds may be disregarded for further study if individually they do
not select for AMR. However, data here indicates that inclusion of these compounds
in mixture investigations is imperative to understand selection for AMR in more
complex mixtures and environments.

<u>Conclusion</u>

Mixtures of diclofenac, metformin, or 17-β-estradiol with ciprofloxacin both increased
the growth inhibitory effects and reduced the selective concentration of ciprofloxacin
in a complex bacterial community. Additionally, the mixtures led to selective
increases or decreases in specific AMR genes or specific species, some of which
are known human pathogens. The effects of antibiotics are traditionally considered in
isolation, particularly in terms of selection for AMR. However, antibiotics are present
alongside other pharmaceuticals, both in the gut, and in the environment. Further,
the data here pertain to simple mixtures, and the effects of more complex mixtures
should be considered in future studies, including co-occurring pharmaceuticals which
may not be selective in isolation.

<u>Data Availability</u>

All data for this project will be made publicly available upon publication.

<u>References</u>

1.   Murray, C.J.L., et al., *Global burden of bacterial antimicrobial resistance in 2019: a*
systematic analysis. The Lancet, 2022. **399**(10325): p. 629-655.

2. Gullberg, E., et al., *Selection of resistant bacteria at very low antibiotic concentrations.* PLOS Pathogens, 2011. **7**(7): p. e1002158.
3. Westhoff, S., et al., *The evolution of no-cost resistance at sub-MIC concentrations of streptomycin in Streptomyces coelicolor.* The ISME Journal, 2017. **11**(5): p. 1168-1178.
4. Kraupner, N., et al., *Selective concentration for ciprofloxacin resistance in Escherichia coli grown in complex aquatic bacterial biofilms.* Environment International, 2018. **116**: p. 255-268.
5. Kraupner, N., et al., *Selective concentrations for trimethoprim resistance in aquatic environments.* Environment International, 2020. **144**: p. 106083.
6. Lundström, S.V., et al., *Minimal selective concentrations of tetracycline in complex aquatic bacterial biofilms.* Science of The Total Environment, 2016. **553**: p. 587-595.
7. Murray, A.K., et al., *Novel Insights into Selection for Antibiotic Resistance in Complex Microbial Communities.* mBio, 2018. **9**(4): p. e00969-18.
8. Sanchez-Cid, C., et al., *Gentamicin at sub-inhibitory concentrations selects for antibiotic resistance in the environment.* ISME Communications, 2022. **2**(1): p. 29.
9. Stanton, I.C., et al., *Evolution of antibiotic resistance at low antibiotic concentrations including selection below the minimal selective concentration.* Communications Biology, 2020. **3**(1): p. 467.
10. Murray, L.M., et al., *Co-selection for antibiotic resistance by environmental contaminants.* npj Antimicrobials and Resistance, 2024. **2**(1): p. 9.
11. Baker-Austin, C., et al., *Co-selection of antibiotic and metal resistance.* Trends in Microbiology, 2006. **14**(4): p. 176-82.
12. Maier, L., et al., *Extensive impact of non-antibiotic drugs on human gut bacteria.* Nature, 2018. **555**(7698): p. 623-628.
13. Younis, W., S. Thangamani, and M.N. Seleem, *Repurposing Non-Antimicrobial Drugs and Clinical Molecules to Treat Bacterial Infections.* Current Pharmaceutical Design, 2015. **21**(28): p. 4106-4111.
14. Lu, J., et al., *Antidepressants promote the spread of extracellular antibiotic resistance genes via transformation.* ISME Communications, 2022. **2**(1): p. 63.
15. Wang, Y., et al., *Non-antibiotic pharmaceuticals enhance the transmission of exogenous antibiotic resistance genes through bacterial transformation.* The ISME Journal, 2020: p. 2179–2196.
16. Ding, P., et al., *Antidepressants promote the spread of antibiotic resistance via horizontally conjugative gene transfer.* Environmental Microbiology, 2022. **24**(11): p. 5261–5276.
17. Wang, Y., et al., *Non-antibiotic pharmaceuticals promote the transmission of multidrug resistance plasmids through intra- and intergenera conjugation.* The ISME Journal, 2021. **15**(9): p. 2493-2508.
18. Jin, M., et al., *Antidepressant fluoxetine induces multiple antibiotics resistance in Escherichia coli via ROS-mediated mutagenesis.* Environment International, 2018. **120**: p. 421-430.
19. Li, X., et al., *Nonsteroidal anti-inflammatory drug diclofenac accelerates the emergence of antibiotic resistance via mutagenesis.* Environmental Pollution, 2023. **326**: p. 121457.
20. Wei, Z., et al., *Emerging pollutant metformin in water promotes the development of multiple-antibiotic resistance in Escherichia coli via chromosome mutagenesis.* Journal of Hazardous Materials, 2022. **430**: p. 128474.
21. Hall, R.J., et al., *Non-antibiotic pharmaceuticals are toxic against Escherichia coli with no evolution of cross-resistance to antibiotics.* npj Antimicrobials and Resistance, 2024. **2**(1): p. 11.
22. Hayes, A., et al., *Antimicrobial effects, and selection for AMR by non-antibiotic drugs on bacterial communities.* bioRxiv, 2024: p. 2024.04.23.590690.
23. aus der Beek, T., et al., *Pharmaceuticals in the environment--Global occurrences and perspectives.* Environmental Toxicology and Chemistry, 2016. **35**(4): p. 823-35.

24. Blair, B.D., et al., *Pharmaceuticals and personal care products found in the Great Lakes above concentrations of environmental concern.* Chemosphere, 2013. **93**(9): p. 2116-2123.

25. den Heijer, M., A. Bakker, and L. Gooren, *Long term hormonal treatment for transgender people.* BMJ, 2017. **359**: p. j5027.

26. Nichols, K.C., L. Schenkel, and H. Benson, *17 beta-estradiol for postmenopausal estrogen replacement therapy.* Obstetrical & Gynecological Survey, 1984. **39**(4): p. 230-45.

27. Ebele, A.J., M. Abou-Elwafa Abdallah, and S. Harrad, *Pharmaceuticals and personal care products (PPCPs) in the freshwater aquatic environment.* Emerging Contaminants, 2017. **3**(1): p. 1-16.

28. Fent, K., A.A. Weston, and D. Caminada, *Ecotoxicology of human pharmaceuticals.* Aquatic Toxicology, 2006. **76**(2): p. 122-159.

29. UKWIR, *Chemicals Investigation Programme Phase 2 Database*, UKWIR, Editor. 2020.

30. Stephanie Graumnitz, D.J., *The database "Pharmaceuticals in the Environment" Update for the period 2017-2020*. 2021.

31. Wilkinson, J.L., et al., *Pharmaceutical pollution of the world's rivers.* Proceedings of the National Academy of Sciences, 2022. **119**(8): p. e2113947119.

32. Hayes, A., et al., *Predicting selection for antimicrobial resistance in UK wastewater and aquatic environments: Ciprofloxacin poses a significant risk.* Environment International, 2022. **169**: p. 107488.

33. Gil, D., et al., *Synergistic antibacterial effects of analgesics and antibiotics against Staphylococcus aureus.* Diagnostic Microbiology and Infectious Disease, 2020. **96**(4): p. 114967.

34. Chan, E.W.L., et al., *Synergistic effect of non-steroidal anti-inflammatory drugs (NSAIDs) on antibacterial activity of cefuroxime and chloramphenicol against methicillin-resistant Staphylococcus aureus.* Journal of Global Antimicrobial Resistance, 2017. **10**: p. 70-74.

35. Annadurai, S., et al., *Experimental studies on synergism between aminoglycosides and the antimicrobial antiinflammatory agent diclofenac sodium.* Journal of Chemotherapy, 2002. **14**(1): p. 47-53.

36. Hegazy, W.A.H., *Diclofenac inhibits virulence of Proteus mirabilis isolated from diabetic foot ulcer.* African Journal of Microbiology Research, 2016. **10**: p. 733-743.

37. Liu, Y., et al., *Metformin Restores Tetracyclines Susceptibility against Multidrug Resistant Bacteria.* Advanced Science, 2020. **7**(12): p. 1902227-1902227.

38. Tang, P.-C., et al., *Within-species variability of antibiotic interactions in Gram-negative bacteria.* mBio, 2024. **15**(3): p. e00196-24.

39. Smith, T.P., et al., *High-throughput characterization of bacterial responses to complex mixtures of chemical pollutants.* Nature Microbiology, 2024. **9**(4): p. 938-948.

40. Klümper, U., et al., *Selection for antimicrobial resistance is reduced when embedded in a natural microbial community.* The ISME Journal, 2019. **13**(12): p. 2927-2937.

41. Singh, N. and P.J. Yeh, *Suppressive drug combinations and their potential to combat antibiotic resistance.* J Antibiot (Tokyo), 2017. **70**(11): p. 1033-1042.

42. Tekin, E., et al., *Prevalence and patterns of higher-order drug interactions in Escherichia coli.* NPJ Syst Biol Appl, 2018. **4**: p. 31.

43. Sutradhar, I., et al., *Effects of antibiotic interaction on antimicrobial resistance development in wastewater.* Sci Rep, 2023. **13**(1): p. 7801.

44. Gillings, M.R., et al., *Using the class 1 integron-integrase gene as a proxy for anthropogenic pollution.* The ISME Journal, 2015. **9**(6): p. 1269-79.

45. Ewels, P., et al., *MultiQC: summarize analysis results for multiple tools and samples in a single report.* Bioinformatics, 2016. **32**(19): p. 3047-3048.

46. Bonin, N., et al., *MEGARes and AMR++, v3.0: an updated comprehensive database of antimicrobial resistance determinants and an improved software pipeline for*

627  *classification using high-throughput sequencing.* Nucleic Acids Res, 2023. **51**(D1): p.
628  D744-d752.

629 47. Pal, C., et al., *BacMet: antibacterial biocide and metal resistance genes database.*
630  Nucleic acids research, 2014. **42**(Database issue): p. D737-D743.

631 48. Florensa, A.F., et al., *ResFinder - an open online resource for identification of*
632  *antimicrobial resistance genes in next-generation sequencing data and prediction of*
633  *phenotypes from genotypes.* Microb Genom, 2022. **8**(1).

634 49. Alcock, B.P., et al., *CARD 2023: expanded curation, support for machine*
635  *learning, and resistome prediction at the Comprehensive Antibiotic Resistance*
636  *Database.* Nucleic Acids Research, 2023. **51**(D1): p. D690-d699.

637 50. Wood, D.E., J. Lu, and B. Langmead, *Improved metagenomic analysis with Kraken*
638  *2.* Genome Biology, 2019. **20**(1): p. 257.

639 51. O'Leary, N.A., et al., *Reference sequence (RefSeq) database at NCBI: current*
640  *status, taxonomic expansion, and functional annotation.* Nucleic Acids Res, 2016.
641  **44**(D1): p. D733-45.

642 52. Pruitt, K.D., T. Tatusova, and D.R. Maglott, *NCBI reference sequences (RefSeq): a*
643  *curated non-redundant sequence database of genomes, transcripts and proteins.*
644  Nucleic Acids Res, 2007. **35**(Database issue): p. D61-5.

645 53. McMurdie, P.J. and S. Holmes, *phyloseq: An R Package for Reproducible Interactive*
646  *Analysis and Graphics of Microbiome Census Data.* PLOS ONE, 2013. **8**(4): p.
647  e61217.

648 54. Paulson JN, O.N., Braccia DJ, Wagner J, Talukder H, Pop M, Bravo HC,
649  *metagenomeSeq: Statistical analysis for sparse high-throughput sequncing.* 2031:
650  Bioconductor package.

651 55. Paulson, J.N., et al., *Differential abundance analysis for microbial marker-gene*
652  *surveys.* Nature Methods, 2013. **10**(12): p. 1200-1202.

653 56. R Core Team, *R: A Language and Environment for Statistical Computing.* 2019, R
654  Foundation for Statistical Computing.

655 57. Wickham, H., *ggplot2: Elegant Graphics for Data Analysis.* 2016: Springer-Verlag
656  New York.

657 58. Mills, B.R., *MetBrewer: Color Palettes Inspired by Works at the Metropolitan Museum*
658  *of Art.* 2022.

659 59. Hartig, F., *DHARMa: residual diagnostics for hierarchical (multi-level/mixed)*
660  *regression models.* R package version 0.3, 2020. **3**.

661 60. Murray, A.K., et al., *The SELection End points in Communities of bacTeria (SELECT)*
662  *Method: A Novel Experimental Assay to Facilitate Risk Assessment of Selection for*
663  *Antimicrobial Resistance in the Environment.* Environmental Health Perspectives,
664  2020. **128**(10): p. 107007.

665 61. Sprouffske, K. and A. Wagner, *Growthcurver: an R package for obtaining*
666  *interpretable metrics from microbial growth curves.* BMC Bioinformatics, 2016. **17**(1):
667  p. 172.

668 62. Bates, D., et al., *Fitting Linear Mixed-Effects Models Using {lme4}.* Journal of
669  Statistical Software, 2015: p. 1-48.

670 63. Lenth, R., et al., *Emmeans: Estimated marginal means, aka least-squares means. R*
671  *Package Version 1 (2018).* 2021.

672 64. Shannon, C.E., *A mathematical theory of communication.* The Bell system technical
673  journal, 1948. **27**(3): p. 379-423.

674 65. Dixon, P., *VEGAN, a package of R functions for community ecology.* Journal of
675  Vegetation Science, 2003. **14**(6): p. 927-930.

676 66. Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*
677  *dispersion for RNA-seq data with DESeq2.* Genome Biology, 2014. **15**(12): p. 550.

678 67. Greenfield, B.K., et al., *Modeling the Emergence of Antibiotic Resistance in the*
679  *Environment: an Analytical Solution for the Minimum Selection Concentration.*
680  Antimicrob Agents Chemother, 2018. **62**(3).

68. Van Hove, B., H. Staudenmaier, and V. Braun, *Novel two-component transmembrane transcription control: regulation of iron dicitrate transport in Escherichia coli K-12.* J Bacteriol, 1990. **172**(12): p. 6749-58.

69. Paulsen, I.T., et al., *A family of Gram-negative bacterial outer membrane factors that function in the export of proteins, carbohydrates, drugs and heavy metals from Gram-negative bacteria.* FEMS Microbiology Letters, 1997. **156**(1): p. 1-8.

70. Dastidar, S.G., et al., *The anti-bacterial action of diclofenac shown by inhibition of DNA synthesis.* International Journal of Antimicrobial Agents, 2000. **14**(3): p. 249-251.

71. Nakamura, S., et al., *gyrA and gyrB mutations in quinolone-resistant strains of Escherichia coli.* Antimicrobial Agents and Chemotherapy, 1989. **33**(2): p. 254-5.

72. Yoshida, H., et al., *Quinolone-resistant mutations of the gyrA gene of Escherichia coli.* Molecular Genetics and Metabolism, 1988. **211**(1): p. 1-7.

73. Riordan, J.T., et al., *Alterations in the transcriptome and antibiotic susceptibility of Staphylococcus aureus grown in the presence of diclofenac.* Annals of Clinical Microbiology and Antimicrobials, 2011. **10**: p. 30-30.

74. Guerin, É., et al., *The SOS Response Controls Integron Recombination.* Science, 2009. **324**(5930): p. 1034-1034.

75. Hocquet, D., et al., *Evidence for Induction of Integron-Based Antibiotic Resistance by the SOS Response in a Clinical Setting.* PLOS Pathogens, 2012. **8**(6): p. e1002778.

76. Masadeh, M.M., et al., *Metformin as a Potential Adjuvant Antimicrobial Agent Against Multidrug Resistant Bacteria.* Clinical Pharmacology : Advances and Applications, 2021. **13**: p. 83-90.

77. Vos, M., et al., *Zinc can counteract selection for ciprofloxacin resistance.* FEMS Microbiology Letters, 2020. **367**(3): p. fnaa038.

78. Palleja, A., et al., *Recovery of gut microbiota of healthy adults following antibiotic exposure.* Nature Microbiology, 2018. **3**(11): p. 1255-1265.

79. MacPherson, C.W., et al., *Gut Bacterial Microbiota and its Resistome Rapidly Recover to Basal State Levels after Short-term Amoxicillin-Clavulanic Acid Treatment in Healthy Adults.* Scientific Reports, 2018. **8**(1): p. 11192.

80. Baltazar, M., et al., *Activation of class 1 integron integrase is promoted in the intestinal environment.* PLOS Genetics, 2022. **18**(4): p. e1010177.

81. Leonard, A.F.C., et al., *Exposure to and colonisation by antibiotic-resistant E. coli in UK coastal water users: Environmental surveillance, exposure assessment, and epidemiological study (Beach Bum Survey).* Environment International, 2018. **114**: p. 326-333.

717