Environmental predictability favours adaptive behavioural plasticity and relaxes selection against deleterious alleles

Lynne Caddy[1,2], Tessy Munoz[1], Julian Baur[1], Loke von Schmalensee[1],
Mareike Koppik[1,3] & David Berger[1*]

1. Department of Ecology and Genetics, Uppsala University. Uppsala 75236, Sweden
2. Plymouth High School for Girls, St Lawrence Road, Plymouth, PL4 6HT, UK.
3. Department of Zoology, Animal Ecology, Martin-Luther University Halle-Wittenberg, Halle (Saale), Germany

Corresponding author: david.berger@ebc.uu.se

Abstract

Behavioural plasticity can play a key role in evolution by either facilitating or impeding genetic adaptation. The latter occurs when behaviours mitigate selection pressures that otherwise would target associated traits. Therefore, environments that facilitate the evolution of adaptive behavioural plasticity are predicted to relax natural selection overall, but experimental evidence for this prediction remains scarce. Here, we first demonstrate that adaptive behavioural plasticity in maternal care in the beetle *Callosobruchus maculatus* is dependent on reliable environmental cues that allow females to reduce larval competition via learning and informed oviposition choices. By comparing survival of larvae of low and high genetic quality in competition with the conspecific *C. phaseoli*, we show that predictable maternal environments relax selection on deleterious alleles in offspring. We further find that females of low genetic quality generally provide poorer care. However, in the predictable environment, the increased opportunity for learning reduced genetic differences in female care, further relaxing selection against deleterious alleles. We illustrate how the identified link between adaptive behavioural plasticity and the strength of natural selection can have important implications for the build-up of cryptic genetic loads and outcomes of interspecific competition across variable environments that differ in their predictability.

Introduction

Phenotypic plasticity plays a fundamental role in shaping organismal responses to changing environments (1–6). Plasticity can help organism to survive and reproduce despite initial adversities. This can allow populations to persist in novel environments and respond to selection, leading to increased genetic change over time (1,3,7–13). However, by mitigating selection pressures, phenotypic plasticity can also reduce the necessity for genetic adaptation (3,14,15). This forgiving nature of plasticity should relax purifying selection and lead to the build-up of genetic variance in quantitative traits that would be deleterious in the absence of plasticity (16–21). If this process continues over time, a population could become dependent on plasticity to survive, as without it, the built up load of conditionally deleterious alleles would become exposed to selection (18,19,22–24).

The evolution of adaptive plasticity is contingent on the organism's ability to change phenotype in accord with changes in the environment (17,25–28). This leads to the expectation that variable environments that provide reliable cues of future change should favour the evolution of adaptive plasticity, whereas unpredictable environments may hinder it when environmental fluctuations are too fast, and/or costs of plasticity are too high, to allow accurate phenotype-environment matching (17,27,29–31). Many behavioural phenotypes represent striking forms of plasticity, where a range of cognitive processes allows the organism to use and process information through experience and learning to fine-tune its strategies to fit the perceived environment (4,32,33). Indeed, adaptive behavioural plasticity in the form of learning has been argued to be a key determinant of evolution in fluctuating environments (3,4,17,34–36). Like for other forms of plasticity, however, learning requires reliable environmental cues to result in accurate behavioural adjustments (17,25,27,37–39). This central role of cue reliability suggests that populations inhabiting environments that differ in their predictability should exhibit systematic differences in learning and genetic load in associated traits. These dynamics are expected to have important consequences for both the levels of segregating genetic variation in natural populations (16,40–42) and for predicting responses to future environmental change (1,3,4,8,17,28,43–45).

Here, we investigated how environmental predictability impacts the rate of learning and the resulting strength of selection on deleterious genetic variation in the seed beetle *Callosobruchus maculatus* (**Fig. 1**). We studied behavioural plasticity in maternal care and females' ability to use information to assess future egg-laying opportunities and learn the spatial location of high-quality hosts. The immediate survival benefits of maternal care to offspring are obvious (46–48), but the flip-side of the coin is that by relaxing selection in offspring, maternal care should, at least theoretically, allow deleterious genetic variation to accumulate. Indeed, in burying beetles, populations that are deprived of parental care suffer immediate increases in juvenile mortality and inbreeding depression (49,50), but at the same time seem to more efficiently purge genetic variation in the long run (21,22,51).

Moreover, given that cognitive processes are costly and governed by many genes (17,52–56), it could be envisioned that the build-up of genetic load via relaxed selection in offspring also feeds back on parents' ability to learn and provide care. Genetic variation in parental care is well-known to impact the expression of genetic variation and selection on traits in offspring via indirect genetic effects (IGEs: the effect of genetic variation expressed in one individual on trait expression in interacting individuals (40–42,57–59)). Imagine that parents of high genetic quality provide better care for their young, who also have intrinsically higher survival due to their superior genetic quality. This positive covariance between parental and offspring breeding values increases genetic variance in offspring survival and is predicted to speed up responses to selection compared to a situation where IGEs are absent (40,57,58,60). Thus, if environmental predictability affects the amount of expressed covariance between adult and juvenile traits, it could have strong effects on evolutionary dynamics across variable environments (24,39,43,61,62).

Female seed beetles provide care for their offspring by preferentially placing eggs on uninfested host seeds, a behaviour that can be finetuned by experience (63–66). First, we gave females the opportunity to choose between uninfested and infested hosts in a spatially complex environment that either provided reliable or unreliable information on infestation rates of future hosts. We show that learning is faster, and care behaviour more accurate, in females receiving reliable information about future hosts, providing evidence that adaptive behavioural plasticity is facilitated by environmental predictability. Second, we find that viability selection in offspring is significantly reduced by faster rates of female learning and improved maternal care in the predictable environment. Third, by comparing the accuracy of maternal care behaviour in mothers of different genetic quality, we find that improved learning in the predictable environment also reduces selection on deleterious alleles in adults. Together, this renders substantially weakened IGEs between maternal care and offspring survival in the predictable environment. Finally, we adopt an existing theoretical model and parametrize it with our experimental data to illustrate potential population-level consequences of the evolutionary feedback between adaptive behavioural plasticity, the strength of natural selection, and the accumulation of conditionally deleterious genetic variation. The results imply that populations evolving exclusively in predictable environments may build up cryptic genetic loads to such extent that they would be unable to colonize unpredictable environments with resident competitor species.

Figure 1: Study design.
A) Path diagram showing hypothesized relationships between environmental predictability, learning, natural selection, and genetic load. Environmental predictability is expected to favour adaptive behavioural plasticity in form of learning. Behavioural plasticity can mitigate genetic differences in trait expression and relax the strength of natural selection, thereby increasing the build-up of deleterious genetic variation over time (i.e. genetic load). The build-up of genetic load could in turn compromise adaptive behavioural plasticity that necessitates cognitive learning (broken arrow). However, the degree to which genetic load impairs learning could also depend on the available environmental cues used in the learning response. To test these hypotheses, we assessed how environmental predictability affected learning of maternal care, and how this in turn affected selection in larval offspring. **B)** Female *C. maculatus* were challenged to find and discriminate among hosts seeds in a repeated spatial learning task. The seeds were either uninfested or infested with eggs of the competitor *C. phaseoli.* We manipulated environmental predictability by giving females three different treatment cues before they performed the task; a reliable cue giving accurate information about the presence of uninfested hosts, an unreliable cue giving false information about the level of infestation of future hosts, and no cue at all. We measured the fitness consequence of maternal care behaviour and learning in each treatment by combining data on female oviposition choice with estimates of larval survival of *C. maculatus* under different levels of competition with *C. phaseoli*. **C)** Three highly inbred strains (blue stripes) were used and compared to themselves when outcrossed (orange) (base stock population; black). Fecundity estimates per line are given as mean ± 2se. The inbred strains show a recessive genetic load amounting to a reduction in fecundity of 55-60% relative to when outcrossed (grey striped bars). The comparison between the three strains in inbred and outbred state was used to assess the relative strength of viability selection in larvae (*s*) and the effect of recessive genetic load on female learning and maternal care, across the three treatments. Based on these estimates, we predicted the genetic load at mutation-selection balance in a meta-population experiencing the predictable, *i*, and unpredictable environment, *j*, at fraction *p* and *k* as a function of the deleterious mutation rate, *u*, and the strength of selection, *s*. See main text for further details.

Methods

Study species

Callosobruchus maculatus is a capital breeding bruchid beetle and a cosmopolitan pest of leguminous crops common in the tropical and subtropical regions. It is facultatively aphagous, i.e. adults do not require food or water to reproduce at high rates (67,68). Adults typically die 7-14 days after emergence in the absence of food and water in standard laboratory conditions of 29°C and 50%RH (69). Both sexes can start reproducing on the day of adult eclosion and females lay 70-90% of their eggs during the first few days of their life (67). The juvenile phase is completed in 3-4 weeks, and egg-to adult-survival rate on preferred hosts *Vigna unguiculata* (black eyed bean) and *V. radiata* (mung bean) is typically around 90% in standard laboratory conditions (70–72). Female host plant search and discrimination is complex. *C. maculatus* has a wide repertoire of fabaceus host plants (73) but females have a clear host hierarchy and preference, and they discriminate between high and low quality species (74,75). Females also heavily discriminate against egg-laden seeds and actively avoid competition with other egg-laying females (63,65), as high larval density limits survival and size at maturity, and thus comes at substantial fitness costs (75–77). *C. maculatus* utilizes both natural habitats where host plant patches used for both egg laying and adult nectar feeding are widely distributed, as well as grain storages where adult food is absent but egg laying substrate is abundant and population density is higher (73,75,76). Thus, selection on spatial cognition and chemosensory cues is likely always strong in natural populations but may take different forms, which is predicted to maintain genetic variation in the cognitive traits under study, as seen for other characters related to these alternative environments (78).

Inbred lines

The three genetic lines used in the experiments were derived from a population isolated from *Vigna unguiculata* (black-eyed bean) seed pods collected at a small-scale agricultural field close to Lomé, Togo (06°10'N 01°13'E) in October and November, 2010 (79). Virgin males and females hatching out of beans were paired randomly and each pair founded an isofemale line that was expanded to a population size of approximately 200 adults over the first two generations. In total, 41 lines were established. These lines have previously been shown to harbor abundant genetic variation for life-history traits and fitness correlates (80). The three inbred genetic lines were created by successive full-sib matings of the 41 lines, resulting in loss of inbred lines and at the end, following more than 30 generations of full-sib mating, the three most productive lines were kept and used for the experiments described below. Thus, these lines represent extremely inbred lines that are next to completely homozygous. Therefore, the comparisons of these lines in inbred and outcrossed state do not correspond to the effect of natural levels of inbreeding. The near complete homozygosity prevents further evolution during the experiments and limits within-line variation due to segregation. These lines should therefore be seen as a powerful tool to inspect the fitness consequences of

genome-wide homozygosity and estimate the strength of selection across genomic sites harboring deleterious alleles.

To provide an estimate of the inbreeding load in the used lines, we estimated the fecundity (number of hatched eggs produced over the first week of reproduction) in females that were either produced by within- (inbred) or between- (outbred) line-crosses. Outcrossing between the three lines was performed in a round-robin fashion following a particular schedule (**Fig. 1C**) and should hide the fitness effects of segregating recessive deleterious alleles, given that these are relatively rare and unlikely to be shared among the three lines. To ensure that bad sperm from inbred males played no part in the fecundity estimates, all females were mated with males from the outbred Lome stock population. We assayed 15 females from each of the six crosses (three inbred and three outbred). For comparison we also scored the fecundity of 15 females from the Lome stock population. Each female was placed into a 35mm petri dish filled with mung beans. These petri dishes were placed in a climate cabinet set at standard conditions and the females were left to lay for the next 7 days after which females were removed. After another 7 days beans were removed, and the number of hatched eggs were counted. This showed that inbreeding has led to a decrease in female fecundity of 55-60% relative to the outbred state (**Fig. 1C,** Supplementary 1).

The Competitor
To estimate the fitness effects of variation in maternal care of *C. maculatus*, we let females choose between uninfested mung beans and mung beans infested with larvae of the closely related competitor, *C. phaseoli*, which has strongly overlapping geographical distribution and host plant preference with *C. maculatus* (81–83). The two species also have similar life-histories (84), although the lab stock of *C. phaseoli* has a 20-30% smaller body size than *C. maculatus* (85). The relative success during interspecific competition inside host seeds could be quantified by counts of relative adult offspring numbers emerging from beans as the species differ in colour patterning. To estimate how variation in maternal care affects offspring fitness, all experiments were performed using mung beans, which is a preferred host of both species but small enough to typically allow for only one or two adults to emerge (**Fig. 1A**).

Learning and maternal care in predictable and unpredictable environments
To decide on the spatial and temporal context in which to assess the ability of *C. maculatus* females to learn the location of host seeds and discriminate their quality, we performed several pilot experiments informed by previous findings showing that females improve their host choice throughout repeated learning trials (66). Based on the previous findings and our pilot data, we constructed arenas from 150 mm diameter petri dishes with the bottom covered in paper to improve traction for females exploring their spatial surroundings. Each arena consisted of a 'starting zone' at

one side of the petri dish, where the females were later to be released. Within the petri dish 35 mung beans, each infested with 2-4 *C. phaseoli* eggs, were spread out in the area outside of the starting zone. Approximately 35 uninfested mung beans were placed into a small 35mm petri dish so that its bottom was covered by a single layer of beans. This petri dish was then inverted into the arena in a designated area opposite to the starting zone after which the dish was removed leaving just the beans (**Fig. 1B**).

Virgin females from the three lines were collected upon adult emergence from mung beans that had been individually isolated in 24-welled chambers. These adult emergences were staggered to allow a constant supply of individuals over a period of a few weeks, so when used in trials females were always between 2 and 4 days old. Virgin females were placed *en masse* in 90mm petri dishes on warming plates set to 30°C. Newly emerged males from the same inbred line were also placed in the petri dish and mating pursued. Mating couples were removed from the petri dish and once mating was over, the female was moved to a small 35mm petri dish and left there for at least 24 hours without beans (29°C and 12:12LD).

Based on a previous study (66), we decided to assay females in groups of five, as this stimulates locomotor activity. A trial started by dividing fifteen mated females equally between three 35mm petri dishes (hereafter known as a 'holding dishes'). Arenas with their associated holding dishes were placed on a single heating plate set to keep the temperature at about 30°C. Females spent five minutes in their holding dish, and approximately 1 minute before being placed in the arena, an environmental cue treatment was added to the holding dish. To one holding dish, 5 non-infested seeds were added (reliable cue), to another holding dish 5 highly infested seeds with over 10 eggs on them were added (unreliable cue), and the final holding dish was left empty (no cue) (**Fig. 1A**). After the final minute with the environmental cue, the females were removed from their holding dish and placed in the starting zone of the arena and left for 10 minutes during which their location (on uninfested beans or not) was recorded every 30 seconds. At the end of the 10 minutes, the females were placed back into their holding dish for five minutes and the procedure to introduce the environmental cues was repeated. During this time, the uninfested beans in the arena were also replaced by new beans. The five females in each of the three environmental cue treatments were exposed to the arena five times (from hereon: "runs"). Previous studies have shown that females of *C. maculatus* adjust their expectations and egg-laying behaviour based on the quality of previously encountered host seeds (63,65,66). We therefore predicted that the environment with the reliable cue would make females perform more accurate host choices as these females had gotten correct information about the presence of uninfested seeds (later found at the other end of the arena). From hereon, we refer to this treatment as the "predictable environment". For the no cue and unreliable cue environments, we expected females to take longer to find the uninfested seeds. For females receiving no cue, we predicted that these females would be less stimulated to start their search for

hosts, as they yet had received any information about the presence of hosts, and that they might regard the hosts with competitor eggs acceptable in the absence of other obvious alternatives. For females receiving the unreliable cue, we predicted that these females would spend more time inspecting the infested seeds found in the arena and also would be more willing to lay eggs on them, as these seeds are of comparatively good quality relative to the seeds that the females had been exposed to in the holding dish before each run. From hereon we refer to these two treatments as the "unpredictable environments".

The first 12 trials were conducted with two environmental treatments simultaneously (6 with reliable versus unreliable cue, and 6 with reliable versus no cue). Once the observer was used to the protocol, we included all three environmental treatments simultaneously, to run an additional 24 trials. The observations were conducted without blinding the observer (LC) due to practical limitations. However, to assess the potential for observer-bias, we later repeated the behavioral assays with a blind observer (TM), conducting another 12 trials using the Lome base population. This showed that all three experiments (2-treatment, 3-treatment and 3-treatment blinded trials) produced qualitatively the same result (summarized in Supplementary 1). We therefore combined all data for the final analysis presented in the main text, based on 66,000 observations of behavior from 660 females in total.

Learning was quantified as the improvement between runs (from 1 to 5) in the rate that females found the patch with uninfested seeds. Thus, to evaluate the effect of environmental predictability on learning, we first analyzed the number of females found on uninfested beans in each arena as a binomial response (maximum value of all five females and minimum value of 0 females) using mixed effects generalized linear models using the lme4 package (86) for R (87). We included fixed effects of time (10 minutes within each run) and run (1-5) crossed with treatment (reliable/unreliable/no cue). Time and run were both added as continuous variables fitted with a second-degree polynomial to account for non-linearities in their effect on the response. Additionally, we included female age and the line ID as additional fixed effects. We included the trial ID as a random effect crossed with treatment. We note that we did not analyze line ID as a random term crossed with treatment, although strictly this could be seen as a relevant level of replication in our experiment, especially in latter analyses comparing the lines in inbred and outbred state (see further below). However, fitting line as a crossed random term with only three levels proved difficult as the mixed models had problems converging. We therefore instead estimated the consistency of the responses in the three lines with reference to treatment and inbreeding status. This showed that the three lines behaved qualitatively the same, and we report a summary of all results from line comparisons in Supplementary 1.

We also quantified differences in learning between the three environments by comparing the number of eggs laid on infested and uninfested seeds from the two non-blinded experiments for a total of

480 females scored in fives over 96 trials. One model was run with a binomial response variable to inspect the fraction of eggs laid on the two types of seeds at the end of each trial. This model included environmental cue and line ID as fixed effects. The second model analyzed the number of eggs laid on the uninfested (high-quality) hosts during each of the five consecutive runs of each trial as a Poisson distributed response variable. This model included environmental cue, run, and their interactions as fixed effects, and trial as a random term. Note that eggs on infested seeds were only counted after the end of each trial and not every run within trials.

Maternal care and the strength of selection on recessive deleterious alleles
To estimate the fitness consequences of variation in maternal care, ultimately leading to either correctly placed eggs onto seeds without competitors, or misplaced eggs onto seeds with many competitors, we tracked survival of *C. maculatus* larvae competing with *C. phaseoli*. To estimate the strength of selection on recessive deleterious alleles at different levels of competition, we compared survival of inbred and outbred *C. maculatus* larvae. We achieved this by mating parents either within (inbred) or between (outbred) lines (**Fig .1C**), reasoning that survival differences between inbred and outbred larvae should reflect natural selection against recessive deleterious alleles revealed by homozygosity in inbred individuals (88). The experiment was carried out in two temporal blocks. Two females (mated either within or between lines), were placed in a 35mm petri dish with mung beans. The seeds were either uninfested or infested with a known number (1, 2, 3 or 4) of *C. phaseoli* eggs that had been laid on the seeds a week earlier (as in the behavioural trials). After 72 hours, the females were removed from the petri dishes and the beans were taken out and checked for eggs. If new eggs were found, they were counted, and beans were placed individually into 24-welled chambers. These chambers were placed into climate cabinets set at standard conditions and were later checked to see how many adults that had emerged of each species.

The data consisted of three replicates (each of two females) per line, inbreeding status, and competition level (0-4 *C. phaseoli* eggs), for a total of 60 female duos, laying 8323 eggs on 2450 mung beans. We analysed the survival of *C. maculatus* as a binomially distributed response, with inbreeding status (inbred/outbred), the number of competitor eggs, and their interaction as fixed effects. The ID of the female duo was added as a random effect. Note that we chose to sum the eggs of both *C. maculatus* and *C. phaseoli* as a measure of competition intensity, as mung beans had 1-3 *C. maculatus* egg laid on them, with one egg being most common and three eggs rare. Additionally, seven seeds had more than three *C. maculatus* eggs on them. We chose to discard these seeds from the analysis since these were unevenly distributed with six of seven seeds from outbred females. A summary of the distribution of eggs laid by both species is presented in Supplementary 7.

To predict the strength of selection, we refitted the mixed effect model within a Bayesian framework using the MCMCglmm package (89) for R. Based on the posterior estimates of model coefficients transformed from log-odds back to original scale, we calculated the strength of selection (with 95% credible intervals) for different levels of competition, c, as : $\Delta\omega_c = 1 - \frac{v_{c,I}}{(v_{c,O})}$, where v_I and v_O is viability of inbred and outbred offspring, respectively. For all Bayesian analyses in this study, we ran models for 100,000 iterations, preceded by 50,000 burn-in iterations, and stored every 100[th] iteration, resulting in 1000 uncorrelated posterior estimates of model coefficients. We used flat and weak priors for the random effects.

Learning and maternal care by mothers of different genetic quality

Using the same set-up as for the original behavioural assays, we quantified the effect of inbreeding load on females' ability to use the environmental cues and adjust their maternal care behaviour. For this we compared inbred and outbred females resulting from within- and between line crosses of the three inbred strains. To minimize the risk that inbred males' sperm would negatively affect female fecundity and their motivation for host search, all females were mated with outbred males from the Lome stock population. The data consisted of 18 trials. In each trial, the three cue treatments were included. Nine trials were run with inbred females and nine with outbred females (i.e. three replicates of five females per line and inbreeding status). This resulted in a total of 27,000 behavioural observations from 270 females. The data was analysed by the same model fitting procedure as for the first experimental data, analysing behavioural responses and the fraction and total number of eggs laid on infested and uninfested seeds, with the addition of adding inbreeding status (inbred/outbred) as a fully crossed fixed effect.

Genetic load in predictable and unpredictable environments

To quantify the fitness cost associated with the revealed recessive deleterious alleles in each of the three environments of our experiment, we refitted the mixed models on juvenile survival (see Fig. 4) and female behavioural plasticity (see Fig. 6) in inbred and outbred individuals using a Bayesian framework, which allowed us to calculate the predicted strength of selection against recessive deleterious alleles based on model posterior distributions. Fitness (ω) of inbred and outbred genotypes could thus be calculated by weighting the number of eggs laid on infested and uninfested seeds by the predictions for larval survival at no competition (uninfested seeds) and three competitors (the mean number of *C. phaseoli* eggs on infested seeds in the behavioural experiments). The genome-wide strength of selection against deleterious alleles in environment j was then calculated as: $s_j = 1 - \omega_{i,j} / \omega_{o,j}$. Using the estimates of s for each environment we then predicted the long-term fitness consequences and genetic load at mutation-selection balance in our predictable and unpredictable environments using a previously published model by Kawecki et al. (1997) (23). Details are described in Supplementary 6.

To further explore potential consequences of the build-up of genetic load for the outcome of interspecific competition, we analysed how the survival of the competitor, *C. phaseoli*, depended on the density of eggs and the inbreeding status of *C. maculatus* (the later as a surrogate measure of genetic load) in the experiment on larval survival (Fig. 4). We also estimated the competitive outcome between the two species by comparing the relative numbers of adults emerging from coinhabited beans. Both models thus fitted survival as a binomial response as a function of the number of *C. phaseoli* and *C. maculatus* eggs, with the latter variable crossed with inbreeding status of *C. maculatus*. The ID of the female duo that laid the *C. maculatus* eggs was added as a random effect in both models. To assess the predicted outcome of interspecific competition depending on the genetic load of *C. maculatus*, we also calculated interspecific competition coefficients between the species, separately for inbred and outbred *C. maculatus*, based on how each species' larval survival was affected by competing eggs of its own and the other species. We approximated each species' carrying capacity for a single mung bean based on the survival data (Supplementary 7; K ≈ 2 for both species). Based on these approximations, we calculated interspecific competition isoclines and mapped them on the model predictions of relative survival of the two species (Fig. 7E).

Results

Learning and maternal care in predictable and unpredictable environments

We first explored whether environmental predictability affected females' ability to learn and express maternal care. Within each of the five runs of a trial, females explored the arena, resulting in more females finding the patch with uninfested seeds as time progressed ($\chi_2^2 = 4851.9$, p < 0.001, Fig. 2). There were clear signs of learning during the trials, signified by a general increase in the proportion of females finding the uninfested beans between the five consecutive runs ($\chi_2^2 = 5818.0$, p < 0.001, Fig. 2). There was evidence for increased learning in the predictable environment compared to the two unpredictable environments providing an unreliable cue or no cue (Fig. 2). This was evident as an effect of environment on the proportion of females finding the uninfested seeds ($\chi_2^2 = 108.8$, p < 0.001) and as a difference between environments in how this proportion increased between runs (Environment:Run interaction: $\chi_4^2 = 72.9$, p < 0.001, Fig. 2). Because females in the predictable environment found the uninfested seeds faster compared to females in the two unpredictable environments, there was also a stronger saturation in the proportion of females found on the uninfested seeds with time in the predictable environment (Environment:Run:Time interaction: $\chi_4^2 = 116.7$, p < 0.001, Fig. 2). Summary of results in Supplementary Table 2.

Figure 2: The effect of environmental predictability on female host search
With time, more females located the patch with uninfested host seeds both within and between runs. The improvement over runs demonstrates learning during the experiment. The rate of learning was highest in females exposed to the reliable cue (blue) compared to those exposed to the unreliable cue (yellow) or no cue (brown).

Females in the predictable environment also laid a larger fraction of eggs (χ_2^2 = 204.3, p < 0.001, Fig. 3, Supplementary Table 3A), and more eggs in total (χ_2^2 = 224.4, p < 0.001, Fig. 3, Supplementary Table 3B), on uninfested seeds. Thus, environmental predictability, providing the opportunity for learning of maternal care, is likely to reduce competition in larval offspring.

Figure 3: The effect of environmental predictability on female host choice
Females laid more eggs on uninfested hosts in later runs of the experiment, showing that they got better at locating the patch with uninfested seeds. The rate of improvement was highest in females exposed to the reliable cue (blue) compared to those exposed to the unreliable cue (yellow) or no cue (brown). This also resulted in females receiving the reliable cue placing a higher proportion of their eggs on uninfested seeds relative to females receiving the other treatments.

Maternal care and the strength of selection on recessive deleterious alleles

To estimate the fitness consequences of variation in maternal care, ultimately leading to either correctly placed eggs onto seeds without competitors, or misplaced eggs onto seeds with many competitors, we tracked survival of *C. maculatus* larvae competing with its close relative, *C. phaseoli*. As expected, the density of competitor eggs had a negative impact on egg-to-adult survival of *C. maculatus* (χ_1^2 = 52.6, p < 0.001, Fig 4). To estimate the strength of selection on recessive deleterious alleles expressed in offspring, we compared the effect of larval competition in inbred and outbred *C. maculatus larvae*. As expected, inbreeding reduced larval survival (χ_1^2 = 49.2, p < 0.001, Fig. 4). Moreover, inbred larvae suffered more from the effect of competition compared to outbred offspring (Inbreeding:Egg density interaction: χ_1^2 = 3.72, p = 0.035, Fig. 4, Supplementary Table 4), confirming that competition increases selection on genetic quality.

Interaction coefficients in logistic regression models represent deviations from multiplicative effects on odds ratios. However, the relevant currency in the eyes of natural selection is given by relative differences in survival among genotypes. Therefore, we estimated the strength of natural selection against recessive deleterious alleles by comparing the relative survival of outbred and inbred larvae at low density (no competition) and high density (six competitors – the maximum observed within our data range) based on estimates from a Bayesian mixed effect model, with the strength of selection at the level of competition, *c*, proportional to: $\Delta\omega_c = 1 - \frac{v_{I,c}}{(v_{O,c})}$, where v_I and v_O is viability of inbred and outbred larvae, respectively. This yielded a fitness cost of deleterious recessives equal to: $\Delta\omega_{C=0}$ = 0.20 (95% CI: 0.05-0.37) with no competition, and: $\Delta\omega_{C=6}$ = 0.84 (0.67-0.94) at high competition, implying a roughly four times greater strength of purifying selection.

Figure 4: The fitness consequences of larval competition
Eggs placed on seeds with more competitor eggs suffered lower survival. This effect was stronger in inbred (grey) compared to outbred (black) larvae, resulting in stronger selection against deleterious alleles at high competition.

Learning and maternal care by mothers of different genetic quality

We compared the ability of inbred and outbred females to assess the environmental cue and improve maternal care by learning. A larger proportion of the females in the predictable environment were found in the patch with uninfested seeds compared to females in the two unpredictable environments (χ_2^2 = 177.2, p < 0.001), replicating the results seen in the previous experiment (compare Fig. 2 and Fig. 5). Outbred females performed better care compared to inbred females (χ_1^2 = 16.6, p < 0.001). Interestingly, however, the difference diminished with time both within (χ_2^2 = 117.8, p < 0.001) and between (χ_2^2 = 223.0, p < 0.001) runs, suggesting that inbred females took longer to find the patch with uninfested seeds, but then performed like outbred females in the last run of each trial (Fig. 5). This indicates that learning can diminish innate genetic differences in maternal care, given enough time. Importantly, this ameliorating effect of learning was strongest in the predictable environment (Environment:Inbreeding interaction: χ_2^2 = 14.6, p < 0.001; Environment:Inbreeding:Run interaction: χ_4^2 = 24.3, p < 0.001, Fig. 5, Supplementary Table 5).

Figure 5: Host search in females of different genetic quality
Females learnt the location of the patch with uninfested host seeds as runs progressed. The rate of learning was again highest in females exposed to the reliable cue (blue) compared to those exposed to the unreliable cue (yellow) or no cue (brown). Inbred females were poorer at finding uninfested seeds. However, learning reduced this difference between outbred and inbred females, especially if females received the reliable cue.

When inspecting the number of eggs laid on host seeds, we found that females in the predictable environment laid both a larger fraction of eggs (χ^2_2 = 72.8, p < 0.001) and overall more eggs (χ^2_2 = 98.3, p < 0.001) on uninfested seeds compared to females in the two unpredictable environments, again replicating the results seen in the previous experiment (compare Fig. 3 and Fig. 6). Outbred females performed better compared to inbred females, both with regards to the fraction (χ^2_1 = 13.4, p < 0.001) and total number (χ^2_1 = 33.2, p < 0.001) of eggs laid on uninfested seeds. For the fraction of eggs on uninfested seeds, we found no effect of the environmental cue on the difference between inbred and outbred females (Environment:Inbreeding interaction: χ^2_2 = 3.88, p = 0.14). However, for the total number of eggs on uninfested seeds, the difference between inbred and outbred females was smaller in the predictable environment (Environment:Inbreeding interaction: χ^2_2 = 11.7, p = 0.003, Fig. 6). This result lends further supports to the hypothesis that predictable environments that leave opportunity for learning can ameliorate individual fitness differences in maternal care owing to genetic quality. Thus, our results show that adaptive behavioural plasticity in the predictable environment diminishes genetic differences in both maternal care and offspring survival, reducing the strength of selection and weakening IGEs between mothers and their offspring.

Figure 6: Host choice in females of different genetic quality
Females got better at locating the patch with uninfested seeds as runs progressed. The rate of improvement was again highest in females exposed to the reliable cue (blue). Inbred females laid fewer eggs on the uninfested seeds. However, when receiving the reliable cue, inbred and outbred females showed smaller differences in the number and proportion of eggs placed on uninfested seeds compared to when receiving the unreliable (yellow) or no (brown) cue.

Genetic load in predictable and unpredictable environments

We quantified the relative fitness cost associated with recessive deleterious alleles in each of the three environments by comparing fitness of inbred and outbred genotypes. Fitness was calculated by weighting the total number of eggs laid on infested and uninfested seeds (Fig. 6) by the predictions for larval survival at no competition (uninfested seeds) and three competitors (the mean number of *C. phaseoli* eggs on infested seeds) (Fig. 4). Expected fitness was higher in the predictable environment compared to the two unpredictable environments (Fig. 7A). Moreover, and echoing the previous findings, the reduction in fitness due to recessive deleterious alleles was lower in the predictable environment with a reliable cue ($\Delta\omega$= 48%, CI: 29-61%) than in the environments with a unreliable cue ($\Delta\omega$= 60%, CI: 45-70%, P_{MCMC} = 0.062) and no cue ($\Delta\omega$= 64%, CI: 52-74%, P_{MCMC} = 0.006), although the difference was marginally non-significant in the first comparison (Fig. 7B, C).

Hence, in accordance with theory, the predictable environment rendered higher fitness and population growth, while at the same time relaxing selection against deleterious alleles. Based on our experimental estimates of selection, we predicted the long-term fitness consequences of behavioural plasticity in the predictable (reliable cue) and unpredictable (no cue) environment by parameterizing an existing model by Kawecki et al. (1997) (23) with our experimental data (full details are given in Supplementary 6). This model estimates the genetic load at mutation-selection balance in a meta-population experiencing two different environments that differ in the strength of selection on deleterious alleles. Based on our experimental data, we set the mean strength of selection in the unpredictable environment to be 0.64/0.48 times stronger in the unpredictable environment (i.e. s_k = 1.33s_j, Fig. 7B, C). The results how that, while fitness is always greater in the predictable environment with a reliable cue, the frequency of deleterious alleles increases in the population as the environment becomes more predictable due to the alleviating effect of adaptive behavioral plasticity on the strength of selection (Fig. 7D). The resulting fitness effects of this accumulated genetic load are more pronounced at high mutation rate and in the unpredictable environment. Thus, in certain scenarios (high predictability of environmental variation and high mutation rate), it could be envisioned that species evolving in predictable environments will be unable to colonize unpredictable environments due to competitive exclusion by resident species.

To illustrate this point further, we modelled the relative survival of the competitor, *C. phaseoli*, in the survival experiment (Fig. 4), depending on the density of eggs and the inbreeding status of *C. maculatus*. This showed that the survival of *C. phaseoli* depends on if *C. maculatus* is inbred or not (Inbreeding:egg density interaction: χ^2_1 = 5.38, p = 0.020, Supplementary Fig. X). To quantify this dependence in more detail, we estimated the competitive outcome between the two species when coinhabiting the same seed. This model confirmed that the outcome of the species-interaction depends on the inbreeding status of *C. maculatus* in such as way that *C. phaseoli* has a competitive

disadvantage against outbred *C. maculatus* larvae, but an advantage when *C. macultatus* is inbred (Inbreeding:egg density: χ_1^2 = 7.83, p = 0.005, Fig. 7E). Based on estimates of interspecific competition coefficients and carrying capacities for each species based on the survival data, we calculated competition isoclines and mapped them on the model predictions of competitive outcome between the two species (Fig. 7E). When *C. phaseoli* is competing against outbred *C. maculatus*, a non-stable equilibrium is reached, with the more abundant species excluding the rare colonizing species. However, when *C. phaseoli* is competing against inbred *C. maculatus* it is the superior competitor and is predicted to exclude *C. maculatus*, irrespective of starting population sizes.

Figure 7: Genetic load in predictable and unpredictable environments.
In A) Expected offspring production is highest in the predictable environment providing a reliable cue compared to the other two environments. B) The strength of selection against deleterious recessive alleles in inbred individuals is reduced in the predictable environment relative to the two unpredictable environments. C) Bayesian posterior densities showing the increase in the strength of selection in the two unpredictable environments compared to the predictable environment. D) Evolution in the predictable environment is predicted to relaxe selection on deleterious alleles, leading to higher fitness but also an accumulation of genetic load in the meta-population. This effect increases for higher deleterious mutation rates (thick lines: *U* = 2, thin lines *U* = 1, set to capture the range of probable values in eukaryotes (90)). Load increases further when the predictable environment is more productive (full lines: B_j/B_k = 1.96 based on the difference in expected fitness of inbred individuals in the environment with a reliable cue relative to those in the environment with no cue, hatched lines: B_j/B_k = 1.30 based on the same difference between outbred individuals, and dotted lines: $B_j = B_k$ for comparison). Further details are given in Supplementary 6. Given that the deleterious mutation rate is not negligible, behavioral plasticity and evolution in the predictable environment will increase load in the unpredictable environment to levels with potentially drastic effects on the outcome of interspecific competition. E) This effect is illustrated by comparing the relative success of *C. phaseoli* in the survival experiment when competing against either outbred or inbred *C. maculatus*. Interspecific competition isoclines are drawn based on approximations of each species' carrying capacity (K) and interspecific competition coefficients (α) on a single mung bean, based on the survival data. When *C. phaseoli* is competing against outbred *C. maculatus*, a non-stable equilibrium is predicted, with the competitive outcome dependent on the starting abundance of each species. Thus, the more abundant species is expected to exclude the colonizing species. When *C. phaseoli* is competing against inbred *C. maculatus* it is the superior competitor and will exclude *C. maculatus*, irrespective of starting frequencies.

Discussion

Finetuning of behaviours can allow organisms to avoid perils and shield them from selection. Behavioural plasticity is therefore regarded as a central mechanism allowing animals to cope with fluctuating environments (3,14,34,40). However, by relaxing the strength of selection overall, behavioural plasticity may lead to increases in conditionally deleterious alleles, which' effects may become exposed in the absence of mitigating behaviours. If this process continues for long enough, organism practising extensive behavioural plasticity are predicted to build up substantial conditionally deleterious genetic loads (17,42,58), perhaps to the extent that they no longer will be competitive in situations where adaptive plasticity is restricted by too rapid and unpredictable environmental changes (90,91). Indeed, to accurately finetune behaviours by learning, the organism needs to be able to assess and predict environmental variability (17,26,27), leading to the expectation that organisms occupying environments that differ in their predictability should show systematic differences in the expression of learning and conditional genetic loads.

Here, we sought an experimental proof-of-principle for this hypothesis. By manipulating the predictability of environmental cues given to egg-laying female seed beetles, we show that learning is strongly dependent on reliable information on future host quality and location. This led to adaptive behavioural plasticity and greatly improved maternal care in the predictable environment, in turn reducing genetic selection in juveniles. This relaxed selection is expected to lead to the build-up of a cryptic genetic load over time. We show that increased genetic load is likely to feed back on the female's ability to perform maternal care; inbred mothers were significantly worse at finding uninfested hosts. However, inbred females learned the location of uninfested hosts with time, diminishing genetic differences in maternal care. Importantly, learning was facilitated by reliable cues, contributing further to relaxing selection on genetic load in the predictable environment. Hence, our study presents convincing evidence for a positive feedback loop between environmental predictability, adaptive behavioural plasticity, and the strength of purifying selection.

Our study also represents a rare experimental demonstration of how the strength of indirect genetic effects (IGEs) can be moulded by the environment. We identified effects of recessive deleterious alleles on both the ability of mothers to provide care and on their offsprings' survival, such that (outbred) mothers of high genetic quality provided better care and increased survival of their offspring, who also had higher intrinsic viability. This positive genetic covariance between the mothers trait (care) and the expression of her offsprings' trait (survival) increases genetic variance in offspring fitness relative to a situation without IGEs (40,57,58,60), and should lead to more efficient purging of genetic load. However, this positive covariance was considerably weakened in the predictable environment that reduced genetic differences between both adults and offspring,

demonstrating that environmental variation can be a key component of evolutionary dynamics driven by IGEs.

C. maculatus females use the presence of eggs to choose whether or not to oviposit (63). However, even though a mother can use cues to make informed decision on where to oviposit, she cannot protect the offspring from the choices made subsequently by other mothers ovipositing on the same host. Likewise, larval genotypes that compete will constitute an evolving environment that may favour the evolution of even more (or less) pronounced competitive traits. This illustrates that environmental predictability in this system is likely to evolve as a function of genes and the IGEs exerted between both competing females and their juvenile offspring (77). Similar dynamics have been elegantly demonstrated in burying beetles (22,51,92,93), and as we argue here, the increased opportunity for learning of maternal care may have strong impact on these processes. It is possible that the interplay between environmental predictability and IGEs may be a key driver of geographic differentiation in strategies of host use in this system. Indeed, different geographic strains of *C. maculatus* show genetic differentiation in foraging strategies (94) but see: ((95)) and novel host use (71). Host specialization has been shown to evolve rapidly in this (74,96) and closely related species (97) and future studies could more directly explore the role of IGEs in driving local adaptation in this system, which may provide important insights into the rapid geographic spread of this insect pest.

While our chosen environmental treatments represent realistic situations for egg-laying *C. maculatus* females, they are unlikely to perfectly match the level of environmental predictability and the frequencies of informed and uninformed oviposition choices in natural populations. Nevertheless, because female *C. maculatus* are faced with ephemeral host availability and varying population densities and levels of competition (63,70,71,71,75,76,78), unpredictability in host quality and abundance is likely ubiquitous. Thus, our treatments were chosen to capture this natural variability, yet the results should not be taken literally but as a proof-of-principle that environmental predictability is key in determining the adaptive value of behavioural plasticity and potential for learning, with cascading effects on the strength of selection and genetic load. To illustrate this point further, we also analysed effects on juvenile survival of the competitor *C. phaseoli* when coinhabiting seeds with inbred and outbred *C. maculatus* larvae. This showed that also competing species can be affected by the dynamics between environmental predictability and behavioural plasticity (Fig. 7). Indeed, our adoption and parameterization of the model on genetic load in heterogenous environments by Kawecki et al. (1997) (23) illustrates that behavioural plasticity in predictable environments can build up a sizeable conditionally deleterious genetic load that becomes exposed to selection upon contact with competing species evolving without plasticity. By parameterizing the model with our rather extreme experimental scenario, comparing an environment giving perfect cue reliability with an environment giving no cue at all, in a situation where heavily infested hosts were presented before high quality hosts (thus skewing the placement of eggs towards infested hosts, all else equal), we

likely overestimated the cost associated with environmental unpredictability. Thus, the model predictions should be taken as an illustration of the principle rather than as an attempt to predict genetic load in *C. maculatus*. Nevertheless, it is worth noting that by assuming homogeneous selection across genomic loci within each environment, and by focusing only on deleterious alleles (i.e. no antagonistic pleiotropy across environments), our calculations underestimate the magnitude of the (hidden) genetic load built up in the predictable environment (98).

Another potential caveat of our study is that our lines were almost completely isogenic (highly inbred), and hence, our estimates of the genetic load exposed by homozygosity is in this sense arbitrary and does not reflect the level of exposure of recessive deleterious alleles that would occur by inbreeding in natural populations. However, our rationale was that these lines would serve as a powerful tool to estimate the relative strength of selection across the predictable and unpredictable environments by manipulating mothers and their larvae to be assayed in outbred and inbred state; completely hiding or expressing all the differentially fixed recessive deleterious alleles across the three lines. Hence, all our estimates are meant to be interpreted in relative terms, giving indications of how the strength of purifying natural selection is affected by correlated changes in maternal care and offspring competition. By doing so, our study illustrates the important role that environmental predictability can play in facilitating learning and adaptive behavioural plasticity, in turn leading to correlated changes in natural selection on associated traits and affecting the strength of IGEs. In support of our findings, a recent comparative study across bird species found that unpredictable environments are associated with reduced parental care (38). Another recent study on burying beetles found that parental care led to a build-up of deleterious mutations over 20 generations of laboratory evolution compared to populations evolving without care (22). However, a study on the same species found the contrasting result that parental care increased phenotypic variance in offspring size (99), which is opposite to general expectations (21,47,49,100,101). This highlights the importance of considering the exact type traits and context of parental provisioning and thus motivates further study of the relationship between environmental predictability and IGEs involving parents and offspring (62). In summary, our study highlights the important link between environmental variation, adaptive behavioural plasticity and genetic load, and that IGEs between parental provisioning and offspring fitness can be highly sensitive to environmental conditions. This in turn has important implications for rates of local adaptation and outcomes of species competition across environments that differ in their predictability.

References

1. Chevin LM, Lande R, Mace GM. Adaptation, Plasticity, and Extinction in a Changing Environment: Towards a Predictive Theory. Kingsolver JG, editor. PLoS Biol [Internet]. 2010 Apr 27 [cited 2018 Feb 16];8(4):e1000357. Available from: http://dx.plos.org/10.1371/journal.pbio.1000357

2. West-Eberhard MJ. Plasticity. In: West-Eberhard MJ, editor. Developmental Plasticity and Evolution [Internet]. Oxford University Press; 2003 [cited 2025 Feb 21]. p. 0. Available from: https://doi.org/10.1093/oso/9780195122343.003.0008

3. Price TD, Qvarnström A, Irwin DE. The role of phenotypic plasticity in driving genetic evolution. Proc R Soc Lond B Biol Sci [Internet]. 2003 Jul 22 [cited 2019 Jan 17];270(1523):1433–40. Available from: http://www.royalsocietypublishing.org/doi/10.1098/rspb.2003.2372

4. Badyaev AV. Stress-induced variation in evolution: from behavioural plasticity to genetic assimilation. Proc R Soc B Biol Sci [Internet]. 2005 May 7 [cited 2020 Nov 21];272(1566):877–86. Available from: https://royalsocietypublishing.org/doi/full/10.1098/rspb.2004.3045

5. Pfennig DW, editor. Phenotypic Plasticity & Evolution: Causes, Consequences, Controversies [Internet]. Taylor & Francis; 2021 [cited 2025 Jan 16]. Available from: https://library.oapen.org/handle/20.500.12657/47881

6. Gibert P, Debat V, Ghalambor CK. Phenotypic plasticity, global change, and the speed of adaptive evolution. Curr Opin Insect Sci [Internet]. 2019 Oct 1 [cited 2025 Jan 16];35:34–40. Available from: https://www.sciencedirect.com/science/article/pii/S2214574519300355

7. Lande R. Adaptation to an extraordinary environment by evolution of phenotypic plasticity and genetic assimilation. J Evol Biol [Internet]. 2009 [cited 2020 Jan 30];22(7):1435–46. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1420-9101.2009.01754.x

8. Snell-Rood EC, Kobiela ME, Sikkink KL, Shephard AM. Mechanisms of Plastic Rescue in Novel Environments. Annu Rev Ecol Evol Syst [Internet]. 2018 [cited 2019 Dec 27];49(1):331–54. Available from: https://doi.org/10.1146/annurev-ecolsys-110617-062622

9. Chevin Luis-Miguel, Hoffmann Ary A. Evolution of phenotypic plasticity in extreme environments. Philos Trans R Soc B Biol Sci [Internet]. 2017 Jun 19 [cited 2019 Feb 5];372(1723):20160138. Available from: https://royalsocietypublishing.org/doi/full/10.1098/rstb.2016.0138

10. Levis NA, Pfennig DW. Evaluating 'Plasticity-First' Evolution in Nature: Key Criteria and Empirical Approaches. Trends Ecol Evol. 2016;31(7):563–74.

11. Fox RJ, Donelson JM, Schunter C, Ravasi T, Gaitán-Espitia JD. Beyond buying time: the role of plasticity in phenotypic adaptation to rapid environmental change. Philos Trans R Soc B Biol Sci [Internet]. 2019 Jan 28 [cited 2024 May 6];374(1768):20180174. Available from: https://royalsocietypublishing.org/doi/full/10.1098/rstb.2018.0174

12. Wcislo WT. Behavioral Environments and Evolutionary Change. Annu Rev Ecol Evol Syst [Internet]. 1989 Nov 1 [cited 2025 Feb 27];20(Volume 20, 1989):137–69. Available from: https://www.annualreviews.org/content/journals/10.1146/annurev.es.20.110189.001033

13. Maklakov AA, Immler S, Gonzalez-Voyer A, Ronn J, Kolm N. Brains and the city: big-brained passerine birds succeed in urban environments. Biol Lett [Internet]. 2011 Oct 23 [cited 2018 Nov 17];7(5):730–2. Available from: http://rsbl.royalsocietypublishing.org/cgi/doi/10.1098/rsbl.2011.0341

14. Huey RB, Hertz PE, Sinervo B. Behavioral Drive versus Behavioral Inertia in Evolution: A Null Model Approach. Am Nat [Internet]. 2003;161(3):357–66. Available from: http://www.jstor.org/stable/3473156

15. Muñoz MM. The Bogert effect, a factor in evolution. Evolution [Internet]. 2022 Feb 1 [cited 2024 Feb 8];76(s1):49–66. Available from: https://doi.org/10.1111/evo.14388

16. Whitlock MC. The Red Queen Beats the Jack-Of-All-Trades: The Limitations on the Evolution of Phenotypic Plasticity and Niche Breadth. Am Nat [Internet]. 1996 [cited 2020 Jan 30];148:S65–77. Available from: https://www.jstor.org/stable/2463048

17. Snell-Rood EC. An overview of the evolutionary causes and consequences of behavioural plasticity. Anim Behav [Internet]. 2013 May 1 [cited 2018 Nov 26];85(5):1004–11. Available from: http://www.sciencedirect.com/science/article/pii/S0003347213000080

18. Dyken JDV, Wade MJ. The Genetic Signature of Conditional Expression. Genetics [Internet]. 2010 Feb 1 [cited 2019 Mar 6];184(2):557–70. Available from: http://www.genetics.org/content/184/2/557

19. Snell-Rood EC, Van Dyken JD, Cruickshank T, Wade MJ, Moczek AP. Toward a population genetic framework of developmental evolution: the costs, limits, and consequences of phenotypic plasticity. BioEssays [Internet]. 2010 Jan [cited 2019 Jan 17];32(1):71–81. Available from: http://doi.wiley.com/10.1002/bies.200900132

20. Walters RJ, Blanckenhorn WU, Berger D. Forecasting extinction risk of ectotherms under climate warming: an evolutionary perspective. Woods A, editor. Funct Ecol [Internet]. 2012 Dec [cited 2018 Feb 19];26(6):1324–38. Available from: http://doi.wiley.com/10.1111/j.1365-2435.2012.02045.x

21. Snell-Rood EC, Burger M, Hutton Q, Moczek AP. Effects of parental care on the accumulation and release of cryptic genetic variation: review of mechanisms and a case study of dung beetles. Evol Ecol [Internet]. 2016 Apr 1 [cited 2019 Oct 15];30(2):251–65. Available from: https://doi.org/10.1007/s10682-015-9813-4

22. Pascoal S, Shimadzu H, Mashoodh R, Kilner RM. Parental care results in a greater mutation load, for which it is also a phenotypic antidote. Proc R Soc B Biol Sci [Internet]. 2023 May 24 [cited 2025 Feb 20];290(1999):20230115. Available from: https://royalsocietypublishing.org/doi/10.1098/rspb.2023.0115

23. Kawecki TJ, Barton NH, Fry JD. Mutational collapse of fitness in marginal habitats and the evolution of ecological specialisation. J Evol Biol [Internet]. 1997 May 1 [cited 2019 Jun 20];10(3):407–29. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1420-9101.1997.10030407.x

24. Badyaev AV, Uller T. Parental effects in ecology and evolution: mechanisms, processes and implications. Philos Trans R Soc B Biol Sci [Internet]. 2009 Apr 27 [cited 2020 Jan 30];364(1520):1169–77. Available from: https://royalsocietypublishing.org/doi/full/10.1098/rstb.2008.0302

25. Komers PE. Behavioural plasticity in variable environments. Can J Zool [Internet]. 1997 Feb [cited 2025 Feb 21];75(2):161–9. Available from: https://cdnsciencepub.com/doi/10.1139/z97-023

26. Forsman A. Rethinking phenotypic plasticity and its consequences for individuals, populations and species. Heredity [Internet]. 2015 Oct [cited 2019 Jan 17];115(4):276–84. Available from: http://www.nature.com/articles/hdy201492

27. Gabriel W, Luttbeg B, Sih A, Tollrian R. Environmental Tolerance, Heterogeneity, and the Evolution of Reversible Plastic Responses. Am Nat [Internet]. 2005 Sep [cited 2025 Feb 21];166(3):339–53. Available from: https://www.journals.uchicago.edu/doi/abs/10.1086/432558

28. Bonamour S, Chevin LM, Charmantier A, Teplitsky C. Phenotypic plasticity in response to climate change: the importance of cue variation. Philos Trans R Soc B Biol Sci [Internet]. 2019 Mar 18 [cited 2021 Mar 22];374(1768):20180178. Available from: https://royalsocietypublishing.org/doi/10.1098/rstb.2018.0178

29. Vinton AC, Gascoigne SJL, Sepil I, Salguero-Gómez R. Plasticity's role in adaptive evolution depends on environmental change components. Trends Ecol Evol [Internet]. 2022 Dec 1 [cited 2024 Sep 30];37(12):1067–78. Available from: https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347(22)00216-6

30. Burton NO, Willis A, Fisher K, Braukmann F, Price J, Stevens L, et al. Intergenerational adaptations to stress are evolutionarily conserved, stress-specific, and have deleterious trade-offs. Tautz D, editor. eLife [Internet]. 2021 Oct 8 [cited 2021 Oct 9];10:e73425. Available from: https://doi.org/10.7554/eLife.73425

31. Dupont L, Thierry M, Zinger L, Legrand D, Jacob S. Beyond reaction norms: the temporal dynamics of phenotypic plasticity. Trends Ecol Evol [Internet]. 2024 Jan 1 [cited 2024 Mar 31];39(1):41–51. Available from: https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347(23)00225-2

32. Dukas R. Evolutionary Biology of Animal Cognition. Annu Rev Ecol Evol Syst [Internet]. 2004 Dec 15 [cited 2018 Nov 17];35(1):347–74. Available from: http://www.annualreviews.org/doi/10.1146/annurev.ecolsys.35.112202.130152

33. Shettleworth SJ. Animal cognition and animal behaviour. Anim Behav [Internet]. 2001 Feb 1 [cited 2018 Nov 27];61(2):277–86. Available from: http://www.sciencedirect.com/science/article/pii/S0003347200916063

34. Baldwin JM. A New Factor in Evolution. Am Nat [Internet]. 1896 Jun 1 [cited 2018 Dec 21];30(354):441–51. Available from: https://www.journals.uchicago.edu/doi/10.1086/276408

35. Mayr E. Behavior Programs and Evolutionary Strategies: Natural selection sometimes favors a genetically 'closed' behavior program, sometimes an 'open' one. Am Sci [Internet]. 1974 [cited 2025 Feb 27];62(6):650–9. Available from: https://www.jstor.org/stable/27845170

36. Snell-Rood EC, Steck MK. Behaviour shapes environmental variation and selection on learning and plasticity: review of mechanisms and implications. Anim Behav [Internet]. 2019 Jan 1 [cited 2019 Apr 9];147:147–56. Available from: http://www.sciencedirect.com/science/article/pii/S0003347218302586

37. Piersma T, Drent J. Phenotypic flexibility and the evolution of organismal design. Trends Ecol Evol [Internet]. 2003 May 1 [cited 2025 Feb 21];18(5):228–33. Available from: https://www.sciencedirect.com/science/article/pii/S0169534703000363

38. Caro SM, Griffin AS, Hinde CA, West SA. Unpredictable environments lead to the evolution of parental neglect in birds. Nat Commun [Internet]. 2016 Mar 29 [cited 2025 Jan 23];7(1):10985. Available from: https://www.nature.com/articles/ncomms10985

39. Leimar O, McNamara JM. The Evolution of Transgenerational Integration of Information in Heterogeneous Environments. Am Nat [Internet]. 2015 Mar [cited 2025 Feb 27];185(3):E55–69. Available from: https://www.journals.uchicago.edu/doi/full/10.1086/679575

40. Bailey NW, Marie-Orleach L, Moore AJ. Indirect genetic effects in behavioral ecology: does behavior play a special role in evolution? Simmons L, editor. Behav Ecol [Internet]. 2018 Jan 13 [cited 2019 Jan 17];29(1):1–11. Available from: https://academic.oup.com/beheco/article/29/1/1/4737155

41. Bijma P. The quantitative genetics of indirect genetic effects: a selective review of modelling issues. Heredity [Internet]. 2014 Jan [cited 2024 Sep 30];112(1):61–9. Available from: https://www.nature.com/articles/hdy201315

42. Linksvayer TA, Wade MJ. GENES WITH SOCIAL EFFECTS ARE EXPECTED TO HARBOR MORE SEQUENCE VARIATION WITHIN AND BETWEEN SPECIES. Evolution [Internet]. 2009 Jul [cited 2019 Jan 17];63(7):1685–96. Available from: http://doi.wiley.com/10.1111/j.1558-5646.2009.00670.x

43. Fisher DN, Kilgour RJ, Siracusa ER, Foote JR, Hobson EA, Montiglio PO, et al. Anticipated effects of abiotic environmental change on intraspecific social interactions. Biol Rev [Internet]. 2021 [cited 2025 Feb 27];96(6):2661–93. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/brv.12772

44. Sih A. Understanding variation in behavioural responses to human-induced rapid environmental change: a conceptual overview. Anim Behav [Internet]. 2013 May 1 [cited 2025 Feb 28];85(5):1077–88. Available from: https://www.sciencedirect.com/science/article/pii/S0003347213000936

45. Greggor AL, Berger-Tal O, Blumstein DT. The Rules of Attraction: The Necessary Role of Animal Cognition in Explaining Conservation Failures and Successes. Annu Rev Ecol Evol Syst [Internet]. 2020 Nov 2 [cited 2024 Sep 30];51(Volume 51, 2020):483–503. Available from: https://www.annualreviews.org/content/journals/10.1146/annurev-ecolsys-011720-103212

46. Mousseau T. The adaptive significance of maternal effects. Trends Ecol Evol [Internet]. 1998 Oct 1 [cited 2019 Jan 17];13(10):403–7. Available from: http://linkinghub.elsevier.com/retrieve/pii/S0169534798014724

47. Royle NJ, Smiseth PT, Kölliker M. The Evolution of Parental Care. Oxford University Press; 2012. 377 p.

48. Clutton-Brock TH. The Evolution of Parental Care. Princeton University Press; 1991. 372 p.

49. Pilakouta N, Jamieson S, Moorad JA, Smiseth PT. Parental care buffers against inbreeding depression in burying beetles. Proc Natl Acad Sci [Internet]. 2015 Jun 30 [cited 2025 Feb 20];112(26):8031–5. Available from: https://www.pnas.org/doi/full/10.1073/pnas.1500658112

50. Bladon EK, Pascoal S, Kilner RM. Can recent evolutionary history promote resilience to environmental change? Behav Ecol [Internet]. 2024 Nov 1 [cited 2024 Dec 19];35(6):arae074. Available from: https://doi.org/10.1093/beheco/arae074

51. Mashoodh R, Trowsdale AT, Manica A, Kilner RM. Parental care shapes the evolution of molecular genetic variation. Evol Lett [Internet]. 2023 Sep 5 [cited 2023 Sep 14];qrad039. Available from: https://doi.org/10.1093/evlett/qrad039

52. Kotrschal A, Rogell B, Bundsen A, Svensson B, Zajitschek S, Brännström I, et al. Artificial Selection on Relative Brain Size in the Guppy Reveals Costs and Benefits of Evolving a Larger Brain. Curr Biol [Internet]. 2013 Jan [cited 2019 Jan 17];23(2):168–71. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0960982212014388

53. Christie ST, Schrater P. Cognitive cost as dynamic allocation of energetic resources. Front Neurosci [Internet]. 2015 Aug 24 [cited 2025 Feb 28];9:289. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4547044/

54. Isler K, van Schaik CP. The Expensive Brain: A framework for explaining evolutionary changes in brain size. J Hum Evol [Internet]. 2009 Oct 1 [cited 2018 Dec 14];57(4):392–400. Available from: http://www.sciencedirect.com/science/article/pii/S0047248409001286

55. Aiello LC, Wheeler P. The Expensive-Tissue Hypothesis: The Brain and the Digestive System in Human and Primate Evolution. Curr Anthropol [Internet]. 1995 [cited 2018 Nov 26];36(2):199–221. Available from: https://www.jstor.org/stable/2744104

56. Mery F, Burns JG. Behavioural plasticity: an interaction between evolution and experience. Evol Ecol [Internet]. 2010 May 1 [cited 2025 Feb 21];24(3):571–83. Available from: https://doi.org/10.1007/s10682-009-9336-y

57. Moore AJ, Brodie ED, Wolf JB. INTERACTING PHENOTYPES AND THE EVOLUTIONARY PROCESS: I. DIRECT AND INDIRECT GENETIC EFFECTS OF SOCIAL INTERACTIONS. Evol Int J Org Evol. 1997 Oct;51(5):1352–62.

58. Wolf JB, Iii EDB, Cheverud JM, Moore AJ, Wade MJ. Evolutionary consequences of indirect genetic effects. Trends Ecol Evol [Internet]. 1998 Feb 1 [cited 2024 Sep 30];13(2):64–9. Available from: https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347(97)01233-0

59. Feldman MW, Laland KN. Gene-culture coevolutionary theory. Trends Ecol Evol [Internet]. 1996 Nov 1 [cited 2021 Jan 9];11(11):453–7. Available from: http://www.sciencedirect.com/science/article/pii/0169534796100525

60. Kirkpatrick M, Lande R. THE EVOLUTION OF MATERNAL CHARACTERS. Evolution [Internet]. 1989 May 1 [cited 2024 Sep 30];43(3):485–503. Available from: https://doi.org/10.1111/j.1558-5646.1989.tb04247.x

61. Bell AM, Hellmann JK. An Integrative Framework for Understanding the Mechanisms and Multigenerational Consequences of Transgenerational Plasticity. Annu Rev Ecol Evol Syst

[Internet]. 2019 Nov 2 [cited 2025 Feb 27];50(Volume 50, 2019):97–118. Available from: https://www.annualreviews.org/content/journals/10.1146/annurev-ecolsys-110218-024613

62. Räsänen K, Kruuk LEB. Maternal effects and evolution at ecological time-scales. Funct Ecol [Internet]. 2007 [cited 2024 Sep 30];21(3):408–21. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2435.2007.01246.x

63. Messina FJ, Renwick J a. A. Mechanism of egg recognition by the cowpea weevil Callosobruchus maculatus. Entomol Exp Appl [Internet]. 1985 [cited 2025 Feb 20];37(3):241–5. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1570-7458.1985.tb03481.x

64. Messina FJ, Morrey JL, Mendenhall M. Why do host-deprived seed beetles ?dump? their eggs? Physiol Entomol [Internet]. 2007 Sep [cited 2019 Jan 17];32(3):259–67. Available from: http://doi.wiley.com/10.1111/j.1365-3032.2007.00579.x

65. Horng SB, Lin HC, Wu WJ, Godfray HCJ. Behavioral processes and egg-laying decisions of the bean weevil, Callosobruchus maculatus. Res Popul Ecol [Internet]. 1999 Dec 1 [cited 2025 Feb 21];41(3):283–90. Available from: https://doi.org/10.1007/s101440050033

66. Baur J, Nsanzimana J d'Amour, Berger D. Sexual selection and the evolution of male and female cognition: A test using experimental evolution in seed beetles*. Evolution [Internet]. 2019 [cited 2020 Jan 2];73(12):2390–400. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/evo.13793

67. Fox CW. Multiple Mating, Lifetime Fecundity and Female Mortality of the Bruchid Beetle, Callosobruchus maculatus (Coleoptera: Bruchidae). Funct Ecol [Internet]. 1993 [cited 2018 Feb 19];7(2):203–8. Available from: http://www.jstor.org/stable/2389888

68. Messina FJ, Fry JD. Environment-dependent reversal of a life history trade-off in the seed beetle Callosobruchus maculatus: Stability of life history trade-offs. J Evol Biol [Internet]. 2003 Apr 17 [cited 2019 Jan 17];16(3):501–9. Available from: http://doi.wiley.com/10.1046/j.1420-9101.2003.00535.x

69. Berger D, Stångberg J, Grieshop K, Martinossi-Allibert I, Arnqvist G. Temperature effects on life-history trade-offs, germline maintenance and mutation rate under simulated climate warming. Proc R Soc B. 2017 Nov 15;284(1866):20171721.

70. Fox CW. A Quantitative Genetic Analysis of Oviposition Preference and Larval Performance on Two Hosts in the Bruchid Beetle, Callosobruchus Maculatus. Evolution [Internet]. 1993 Feb 1 [cited 2018 Nov 28];47(1):166–75. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1558-5646.1993.tb01207.x

71. Messina FJ, Lish AM, Gompert Z. Variable Responses to Novel Hosts by Populations of the Seed Beetle Callosobruchus maculatus (Coleoptera: Chrysomelidae: Bruchinae). Environ Entomol [Internet]. 2018 Oct 3 [cited 2022 Mar 15];47(5):1194–202. Available from: https://doi.org/10.1093/ee/nvy108

72. Berger D, Stångberg J, Baur J, Walters RJ. Elevated temperature increases genome-wide selection on de novo mutations. Proc R Soc B Biol Sci [Internet]. 2021 Feb 10 [cited 2022 Jan 5];288(1944):20203094. Available from: https://royalsocietypublishing.org/doi/full/10.1098/rspb.2020.3094

73. Tuda M, Kagoshima K, Toquenaga Y, Arnqvist G. Global Genetic Differentiation in a Cosmopolitan Pest of Stored Beans: Effects of Geography, Host-Plant Usage and Anthropogenic Factors. PLOS ONE [Internet]. 2014 Sep [cited 2018 Nov 28];9(9):e106268. Available from: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0106268

74. Fricke C, Arnqvist G. RAPID ADAPTATION TO A NOVEL HOST IN A SEED BEETLE (*CALLOSOBRUCHUS MACULATUS*): THE ROLE OF SEXUAL SELECTION: ADAPTATION AND SEXUAL SELECTION. Evolution [Internet]. 2007 Feb [cited 2018 Feb 19];61(2):440–54. Available from: http://doi.wiley.com/10.1111/j.1558-5646.2007.00038.x

75. Mitchell R. The Evolution of Oviposition Tactics in the Bean Weevil, Callosobruchus maculatus (F.). Ecology [Internet]. 1975 May 1 [cited 2018 Nov 28];56(3):696–702. Available from: https://esajournals.onlinelibrary.wiley.com/doi/abs/10.2307/1935504

76. Credland PF, Dick KM, Wright AW. Relationships between larval density, adult size and egg production in the cowpea seed beetle, Callosobruchus maculatus. Ecol Entomol [Internet]. 1986 Feb 1 [cited 2018 Nov 28];11(1):41–50. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2311.1986.tb00278.x

77. Fox CW, Savalli UM. Inheritance of Environmental Variation in Body Size: Superparasitism of Seeds Affects Progeny and Grandprogeny Body Size Via a Nongenetic Maternal Effect. Evolution [Internet]. 1998 [cited 2024 Sep 30];52(1):172–82. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1558-5646.1998.tb05150.x

78. Utida S. Density dependent polymorphism in the adult of Callosobruchus maculatus (Coleoptera, Bruchidae). J Stored Prod Res [Internet]. 1972 Jun 1 [cited 2018 Nov 28];8(2):111–25. Available from: http://www.sciencedirect.com/science/article/pii/0022474X72900288

79. Berger D, Grieshop K, Lind MI, Goenaga J, Maklakov AA, Arnqvist G. INTRALOCUS SEXUAL CONFLICT AND ENVIRONMENTAL STRESS: SEX, GENES, AND CONFLICT IN STRESSFUL ENVIRONMENTS. Evolution [Internet]. 2014 May [cited 2018 Feb 19];n/a-n/a. Available from: http://doi.wiley.com/10.1111/evo.12439

80. Berger D, Martinossi-Allibert I, Grieshop K, Lind MI, Maklakov AA, Arnqvist G. Intralocus Sexual Conflict and the Tragedy of the Commons in Seed Beetles. Am Nat [Internet]. 2016 Oct [cited 2018 Nov 28];188(4):E98–112. Available from: https://www.journals.uchicago.edu/doi/10.1086/687963

81. Temreshev II, Kazenas VL. Callosobruchus phaseoli (Gyllenhal, 1833) (Coleoptera, Chrysomelidae, Bruchinae): a new invasive species in Kazakhstan. Acta Biol Sib [Internet]. 2020 Jul 23 [cited 2024 Oct 15];6:87–92. Available from: https://abs.pensoft.net/article/53070/

82. Daglish GJ, Erbacher JM, Eelkema M. Efficacy of protectants against *Callosobruchus phaseoli* (Gyll.) and *C. maculatus* (F.) (Coleoptera: Bruchidae) in mungbeans. J Stored Prod Res [Internet]. 1993 Dec 1 [cited 2024 Oct 15];29(4):345–9. Available from: https://www.sciencedirect.com/science/article/pii/0022474X9390050E

83. Lambrides CJ, Imrie BC. Susceptibility of mungbean varieties to the bruchid species Callosobruchus maculatus (F.), C. phaseoli (Gyll.), C. chinensis (L.), and Acanthoscelides obtectus (Say.) (Coleoptera:Chrysomelidae). Aust J Agric Res [Internet]. 2000 [cited 2024 Oct 15];51(1):85–90. Available from: https://www.publish.csiro.au/ar/ar99051

84. Arnqvist G, Rönn J, Watson C, Goenaga J, Immonen E. Concerted evolution of metabolic rate, economics of mating, ecology, and pace of life across seed beetles. Proc Natl Acad Sci [Internet]. 2022 Aug 16 [cited 2024 Jun 12];119(33):e2205564119. Available from: https://www.pnas.org/doi/abs/10.1073/pnas.2205564119

85. Bagchi B, Corbel Q, Khan I, Payne E, Banerji D, Liljestrand-Rönn J, et al. Sexual conflict drives micro- and macroevolution of sexual dimorphism in immunity. BMC Biol [Internet]. 2021 Jun 2 [cited 2021 Jun 4];19(1):114. Available from: https://doi.org/10.1186/s12915-021-01049-6

86. Bates D. Mixed models in R using the lme4 package Part 4: Inference based on profiled deviance. :23.

87. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2020. Available from: https://www.R-project.org/

88. Charlesworth D, Willis JH. The genetics of inbreeding depression. Nat Rev Genet [Internet]. 2009 Nov [cited 2020 Dec 5];10(11):783–96. Available from: https://www.nature.com/articles/nrg2664

89. Hadfield JD. MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. J Stat Softw. 2010;33(2):1–22.

90. Agrawal AF, Whitlock MC. Mutation Load: The Fitness of Individuals in Populations Where Deleterious Alleles Are Abundant. Annu Rev Ecol Evol Syst [Internet]. 2012 Dec [cited 2018 Feb 19];43(1):115–35. Available from: http://www.annualreviews.org/doi/10.1146/annurev-ecolsys-110411-160257

91. Tienderen PHV. Evolution of Generalists and Specialists in Spatially Heterogeneous Environments. Evolution [Internet]. 1991 [cited 2021 Mar 22];45(6):1317–31. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1558-5646.1991.tb02638.x

92. Rebar D, Bailey NW, Jarrett BJM, Kilner RM. An evolutionary switch from sibling rivalry to sibling cooperation, caused by a sustained loss of parental care. Proc Natl Acad Sci [Internet]. 2020 Feb 4 [cited 2025 Mar 4];117(5):2544–50. Available from: https://www.pnas.org/doi/abs/10.1073/pnas.1911677117

93. Bladon EK, Pascoal S, Bird N, Mashoodh R, Kilner RM. The evolutionary demise of a social interaction: experimentally induced loss of traits involved in the supply and demand of care. Evol Lett [Internet]. 2023 Jun 1 [cited 2025 Mar 4];7(3):168–75. Available from: https://doi.org/10.1093/evlett/qrad016

94. Credland PF, Dick KM. Food consumption by larvae of three strains of Callosobruchus maculatus (Coleoptera: Bruchidae). J Stored Prod Res [Internet]. 1987 Feb [cited 2022 Mar 15];23(1):31–40. Available from: https://linkinghub.elsevier.com/retrieve/pii/0022474X87900336

95. Burc E, Girard-Tercieux C, Metz M, Cazaux E, Baur J, Koppik M, et al. Life-history adaptation under climate warming magnifies the agricultural footprint of a cosmopolitan insect pest. Nat Commun [Internet]. 2025 Jan 18 [cited 2025 Jan 19];16(1):827. Available from: https://www.nature.com/articles/s41467-025-56177-2

96. Rêgo A, Messina FJ, Gompert Z. Dynamics of genomic change during evolutionary rescue in the seed beetle Callosobruchus maculatus. Mol Ecol. 2019 May;28(9):2136–54.

97. Savković U, Đorđević M, Šešlija Jovanović D, Lazarević J, Tucić N, Stojković B. Experimentally induced host-shift changes life-history strategy in a seed beetle. J Evol Biol [Internet]. 2016 Apr [cited 2019 Jan 17];29(4):837–47. Available from: http://doi.wiley.com/10.1111/jeb.12831

98. Hermisson J, Wagner GP. The Population Genetic Theory of Hidden Variation and Genetic Robustness. Genetics [Internet]. 2004 Dec [cited 2018 Jan 29];168(4):2271–84. Available from: http://www.genetics.org/lookup/doi/10.1534/genetics.104.029173

99. Schrader M, Jarrett BJM, Kilner RM. Parental care and sibling competition independently increase phenotypic variation among burying beetle siblings. Evolution [Internet]. 2018 Nov 1 [cited 2025 Feb 20];72(11):2546–52. Available from: https://doi.org/10.1111/evo.13607

100. Hunt J, Simmons LW. The genetics of maternal care: Direct and indirect genetic effects on phenotype in the dung beetle Onthophagus taurus. Proc Natl Acad Sci [Internet]. 2002 May 14 [cited 2024 Sep 30];99(10):6828–32. Available from: https://www.pnas.org/doi/10.1073/pnas.092676199

101. Patterson C, Pilakouta N. Effects of Parental Care on the Magnitude of Inbreeding Depression: A Meta-Analysis in Fishes. Am Nat [Internet]. 2024 Feb 1 [cited 2024 Nov 8]; Available from: https://www.journals.uchicago.edu/doi/10.1086/728001