Warming and species richness weaken eco-phenotypic feedback loop in

long-term natural ecosystems.

Running title: Warming weakens eco-phenotypic feedback

Qinghua Zhao[1,5], Yingying X. G. Wang[2,5], Chi Xu[3], George Sugihara[4], Frederik

De Laender[1,5]

[1]Research Unit of Environmental and Evolutionary Biology (URBE), Institute of

Complex Ecosystems (naXys), Institute of Life, Earth and the Environment

(ILEE), University of Namur, Namur, Belgium.

[2]School of Veterinary Medicine, University of California-Davis, Davis, CA, USA

[3]School of Life Sciences, Nanjing University, Nanjing, 210023, China

[4]Scripps Institution of Oceanography, University of California-San Diego, La

Jolla, CA, USA

[5]These authors contributed equally

Correspondence: Yingying X. G. Wang (Yingying.xgz.wang@gmail.com)

Abstract

Eco-phenotypic feedback—reciprocal interactions between phenotypic traits and ecological dynamics—is increasingly recognized as a driver of biodiversity patterns, species interactions, and ecosystem functioning. Through this feedback, phenotypic traits such as body size can rapidly respond to environmental variation through plastic or evolutionary changes, altering population abundance, which in turn feeds back to shape the trait dynamics. Yet, whether the integrity of this feedback remains stable under environmental change remains unclear. Using long-term monitoring data from 101 aquatic and terrestrial ecosystems encompassing multiple generations, we provide the first synthesis showing that warming and species richness systematically weakens the eco-phenotypic feedback loop, while mean species body size strengths it. Our findings reveal that climate change can erode key trait–demography couplings and highlight the importance of integrating eco-phenotypic frameworks into global change research.

Keywords: Eco-phenotypic feedback, Climate Warming, Phenotypic trait, Eco-evolutionary feedback

Introduction

Environmental change can rapidly alter both phenotypic traits and population abundance (Gardner *et al.* 2011; McLean *et al.* 2016). For instance, warming can reduce body size through increased metabolic costs while lowering population density via reduced survival or recruitment (Gardner *et al.* 2011); similarly, predation often favor smaller-bodied individuals while directly decreasing abundance through higher mortality (Bergquist *et al.* 1985; Wellborn 1994). More broadly, changes in phenotypic traits and population abundance are not independent but often interdependent (Gibert *et al.* 2022; Govaert & Klauschies 2024; Han *et al.* 2023). Trait shifts can modify demographic rates, influencing population density, which in turn feeds back to shape trait dynamics through plasticity or selection (Govaert & Klauschies 2024). When these reciprocal influences occur together, they form a coupled loop in which ecological and phenotypic changes reinforce and modify each other over time (Govaert & Klauschies 2024; Han *et al.* 2023). This coupled dynamic gives rise to eco-phenotypic feedback (Govaert & Klauschies 2024; Han *et al.* 2023).

Unlike eco-evolutionary feedback that requires heritable genetic changes, eco-phenotypic feedback can operate through both heritable genetic changes and nonheritable mechanisms, including behavioral shifts, and physiological adjustments (Govaert & Klauschies 2024). Eco-phenotypic feedback has been documented in a range of taxa and environments, from single species to food webs, regulating population persistence, trophic interactions, community structure, and ecosystem resilience under ongoing environmental changes (Gibert *et al.* 2022; Govaert & Klauschies 2024; Han *et al.* 2023). For instance, in fish populations subject to intense harvesting, reduced body size can lower fecundity and recruitment, driving population decline; lower densities then relax intraspecific competition for resources, potentially allowing body size to increase in subsequent generations (Conover & Munch 2002; Hutchings & Reynolds 2004; Law 2000). Similarly, in mussel beds, high density increases

competition for planktonic food and space, limiting growth and reducing

average shell size (Bayne & Worrall 1980; Okamura 1986). Smaller mussels

have reduced fecundity, lowering recruitment and abundance over time (Bayne

& Worrall 1980; Gosling 2003). Lower density then reduces competition and

increases resource availability, enabling larger bodied shell growth in surviving

individuals and recruits (Bayne & Worrall 1980; Okamura 1986). This feedback

shapes mussel bed structure, competitive interactions with other sessile

invertebrates, and the resilience of intertidal communities (Rietkerk *et al.* 2021;

Van De Koppel *et al.* 2008).

However, strength of eco-phenotypic feedback is not fixed but context

dependent, shifting with environmental conditions (Govaert & Klauschies 2024;

Han *et al.* 2023). Climate warming—a pervasive anthropogenic driver—can

alter both trait dynamics and population abundance (McLean *et al.* 2016;

Schwarz *et al.* 2017). Rising temperatures can influence phenotypic traits

through metabolic constraints, developmental acceleration, and shifts in life

history strategies (McLean *et al.* 2016). Concurrently, warming can modify

population abundance by affecting reproductive rates, mortality, dispersal, and

interspecific interactions (Parmesan 2006; Walther *et al.* 2002). Although

numerous studies have examined these effects independently, few have

explicitly investigated how warming disrupts their coupling, with existing

evidence largely coming from experiments and no studies from natural

ecosystems. Beyond temperature, eco-phenotypic feedback strength may also

be shaped by biodiversity such as species richness. Greater richness increases

the likelihood of diverse interspecific interactions—competition, facilitation,

predation—that can alter both traits and population abundance trajectories

(Tilman 1999). Trait variation within populations, such as mean body size, can

further buffer or amplify the feedback strength; species with smaller mean size

often exhibit shorter generation times and greater plasticity, potentially

tightening or weakening the feedback (Kingsolver & Pfennig 2007; Peters 1983).

However, no study has explicitly documented the effect of species richness and
mean body size on the integrity of eco-phenotypic feedback.

We address these gaps by conducting a synthesis evaluating the eco-
phenotypic feedback integrity under warming, biodiversity and trait variation in
natural ecosystems. We compiled long-term, continent-wide data from 101
ecological systems spanning aquatic and terrestrial environments, including
plankton, fish, and ground beetle communities, spanning 10 to 47 years and
encompassing multiple generations per species and enabling robust detection
of trait–phenotypic feedback. Specifically. We use Convergent Cross Mapping
(CCM), a model-free technique for inferring causal relationships, to quantify
bidirectional feedback between the phenotypic trait (body size) and population
dynamics (Fig 1). We then assess how the prevalence of this feedback varies
with climate warming and biotic drivers such as species richness and average
trait values. By linking eco-phenotypic feedback to key abiotic and biotic
drivers, this study strengthens the conceptual and empirical basis for
incorporating eco-phenotypic feedback into ecological theory and forecasting.
113

Methods

We compiled a long-term dataset encompassing 101 ecosystems: 51
aquatic planktonic systems monitored for 10–40 years, 30 fish systems
spanning 31–47 years, and 20 terrestrial systems (beetles) spanning 9–11
years. The time series span multiple generations for each species, providing a
sufficient temporal window for the phenotypic trait (body size) to change
through plastic or evolutionary processes. This duration also enables the
detection of concurrent shifts in abundance and body size, allowing robust
assessment of eco-phenotypic feedback. We use body size as a proxy for
phenotypic traits, following previous studies on eco-phenotypic feedbacks
(Govaert & Klauschies 2024; Han *et al.* 2023), as body size is one of the most
influential traits, exerting pervasive effects across levels of biological
organization and underpinning numerous physiological, ecological, and

evolutionary processes that often scale with it (Gardner *et al.* 2011; White *et al.*
2007).

Aquatic plankton Data. Body size and abundance data were compiled
from 51 long-term monitoring sites across aquatic planktonic ecosystems,
comprising 20 lakes, 8 rivers, and 23 marine systems (Fig 2a). These datasets
were accessed by North Temperate Lakes Long-Term Ecological Research
(Magnuson *et al.* 2023a, 2024a, 2022, 2023b), Pangaea (Felden *et al.* 2023),
USGS (Nejad *et al.* 2017), Greenland Ecosystem Monitoring (Christensen *et al.*
2017) and Data Observation Network for Earth (Michener *et al.* 2011). The 51
ecosystems have been consistently sampled for 10–40 years. The resident
species include phytoplankton and zooplankton, with maximum generation
times typically less than a week and two months, respectively (Elbrächter
1976; Gillooly 2000a; Litchman 2000). Therefore, the 10–40 years of sampling
encompass numerous generations of these planktonic species. For each of the
51 aquatic planktonic ecosystems, there were three types of variables: (1)
species abundance; (2) species size (i.e. body length or biovolume); (3) water
temperature. For species size (2), a single metric is reported for each species
and subsequently used for further analysis.

Next, for consistency, each species' time series were standardized to
seasonal sampling frequencies (if multiple samplings were conducted per
season, e.g., monthly and bimonthly sampling data sets, those were averaged).
Here, the seasonal resolution (trimonthly) is the shared consensus that can be
applied to all 51 data sets, and the seasonal average here is also the most
representative measure across all data sets, because the equal sampling
interval is necessary for Convergent Cross Mapping (CCM) analysis. After the
seasonal average, there were 0.05% missing points across the whole planktonic
dataset (41 out of 85161 data points). Those missing data points were linearly
interpolated using *na.approx* function in the package of zoo (Karakoç *et al.*
2020; Zhao *et al.* 2023). Then, the species abundance across all data sets was

scaled to the same unit (individuals per litre). Species length and biovlume

were standardized to the same unit (μm) and (μm^3) respectively.

Prior to CCM analysis, we removed long-term linear trends and seasonal

cycles from each time series to avoid spurious causality from shared external

drivers (Chang *et al.* 2020, 2022). Specifically, we first removed the long-term

linear trend from each time series by using the residuals from a linear

regression against time (Chang *et al.* 2020, 2022). Next, we removed

seasonality by scaling against the mean and standard deviation of values

occurring in the same year, $D_{season}(t) = (O(t) - \mu_{season})/\sigma_{season}$, where $D_{season}(t)$

is the deseasoned time series, $O(t)$ is the original time series, μ_{season} is the year

mean, and the σ_{season} is the year standard deviation (Chang *et al.* 2020, 2022).

Finally, each time series (both species abundance and size) was scaled to zero

mean and unit variance.

Fish Data. Fish size and abundance data were compiled from 30

monitoring sites, including six rivers, five lakes and 19 marines. These datasets

were accessed by North Temperate Lakes Long-Term Ecological Research

(Magnuson *et al.* 2024c, b), International Council for the Exploration of the Sea

(ICES) (Fish trawl surveys 2025), and Reef Life Survey (Edgar & Stuart-Smith

2014). These ecosystems have been consistently sampled for 31–47 years.

Again, for each Fish ecosystem, there were three types of variables: (1) species

abundance; (2) species size (i.e. body length); (3) water temperature.

For consistency each species time series of abundance and size were

standardized to the yearly sampling frequency (if multiple samplings were

conducted per year, e.g., quarterly sampling data sets, those were averaged).

Again, the yearly resolution is shared across all data sets. After the yearly

average, there were no missing points across the fish dataset. Then, the fish

abundance across all data sets was standardized to the same unit (caught per

effort). Species length was standardized to the same unit (mm). As all fish data

were standardized to yearly intervals, there was no need to remove long-term

linear or seasonal trends from each time series. Therefore, the original data

were utilized for further analysis. Finally, each time series (both species

abundance and size) were scaled to zero mean and unit variance.

Terrestrial species. Terrestrial species size and abundance data were

compiled from 20 monitoring sites, all focused on ground beetles. All data were

retrieved from BioTIME (Dornelas *et al.* 2018). These ground beetles were

consistently sampled for 10–11 years. Given a maximum generation time of

typically smaller than a year, the sampling period encompasses numerous

generations across all species (Kirk 1975; Matalin 2007; Rivard 1964). Again,

for each site, there were three types of variables: (1) species abundance; (2)

species size (i.e. mg); (3) surface temperature. Each site reports both

abundance and size. The daily temperature and precipitation for each site were

retrieved from NOAA's National Centers for Environmental Information (GHCN-

Daily, Version 3 2012).

Next, to ensure consistency, each species' time series were standardized

to seasonal sampling frequencies (if multiple samplings were conducted per

season, those were averaged). After calculating seasonal averages, 0.52% of

data points were missing in the dataset. These missing values were

subsequently linearly interpolated using the *na.approx* function from the zoo

package (Karakoç *et al.* 2020; Zhao *et al.* 2023). Then, long-term linear and

seasonal trends were removed from each time series using the same

methodology applied to the plankton data. Finally, each time series, including

both species abundance and size, was standardized to have a mean of zero and

a unit variance.

Presence/absence of eco-phenotypic feedback linkages under biotic

and abiotic drivers. For each species (plankton, fish or terrestrial species), we

identified the presence or absence of the feedback linkages (i.e. size

→abundance, abundance →size) using CCM (Fig 1). CCM is grounded in

Takens's theorem for dynamic systems, which infers causal relationships

among variables from empirical time series (Deyle & Sugihara 2011; Sugihara

et al. 2012). If a causal variable X (e.g., species abundance) and a response

variable Y (e.g., body size) belong to the same dynamical system, information

about X is embedded in Y. Thus, the state of X can be predicted from Y through
a process known as cross mapping (cross-prediction) (Sugihara *et al.* 2012).
CCM quantifies causality by assessing the degree to which X leaves a signature
in the time series of Y. In this study, the appropriate embedding dimensions for
cross-mapping E were determined by the optimized the hindcast cross-
mapping in which X(t) projected one-step backward to Y(t-1), examining values
of E from 2 to square root of *n*, where n is the length of the time series (Karakoç
et al. 2020). Thus, we computed *n* as the geometric mean time series length
across all data sets. E was finally examined from 2 to 7 across all data sets. We
used simplex projection to select the best E that gave the highest prediction
skill (Hsieh *et al.* 2005).
To accommodate the fact that the causal variables (e.g., species
abundance) can exhibit time-delayed effects on the affected variables (e.g.,
species size), we carried out 0 to 1 year (i.e. 0~4 time point for plankton and
terrestrial species but 1 time point for fish) lagged CCM analyses, in which we
retained the CCM with the time lag resulting in the highest cross mapping skill
ρ (Ye *et al.* 2019). The criterion for testing causality is the convergence toward
higher cross-mapping skill with increasing library length (i.e., the number of
points used for state space reconstruction) (Sugihara *et al.* 2012). An increase
in library length enhances the density of points in the reconstructed attractor,
facilitating more precise identification of nearest neighboring points for
predictions. This, in turn, leads to improved predictive accuracy (Sugihara &
May 1990). Convergence can be assessed by evaluating whether there is a
significant monotonic increasing trend in cross-mapping skill ρ as library
length increases using Kendall's τ test (Chang *et al.* 2020), and whether the ρ
at the largest library length is significantly greater than that the ρ at the
smallest library length using Fisher's Z test (Chang *et al.* 2020). In this study,
library length was set from the minimum (E) to the maximum length (i.e., the
length of the entire time series). Throughout this study, an interaction link
(e.g., X→Y) was considered causal if both Kendall's τ test and Fisher's Z test

were statistically significant ($P < 0.05$) for testing convergence (Chang *et al.*
2020).

Then, the warming rate, representing the long-term warming trend of
water temperature or land surface temperature in each ecosystem, was
measured using the Theil–Sen median-based trend estimator, implemented via
'sens.slope' from 'trend' package (Chang *et al.* 2020; Mohsin & Gough 2010).
This method was chosen due to its widespread use in quantifying warming
rates (Pinsky *et al.* 2025; Tong *et al.* 2023), its robustness to episodic extreme
events, and its ability to provide stable estimates of long-term trends in the
presence of outliers (Mohsin & Gough 2010). Species mean body mass was
computed as average across whole time series, estimated from biovolume,
assuming that organisms have the same density as water, if unavailable, was
obtained from published sources (Cohen & Lough 1981; Huntley 1992).
Similarly, mean temperature was computed as average across whole time
series. Species richness for each of the 101 ecosystems was calculated as the
total number of species present within each system.

Next, we employed a generalized linear mixed model (glmmTMB function
from the glmmTMB package (Brooks *et al.* 2017)) with a binomial error
distribution to test the main effects of warming, mean temperature, species
richness, mean species body mass on presence/absence of the feedback
linkages—either from phenotypic trait (body size) to abundance (size
→abundance) or from abundance to body size (abundance → size) (table S1).
To ensure normality, mean temperature, species richness, and mean species
body mass were log-transformed using base 10. Each of these predictors in the
model were normalized using the scale function from base R. The response
variable in this model was binary, with 1 indicating the presence of a feedback
linkage identified using CCM and 0 indicating its absence. Last, we set species
groups (e.g. plankton) nested within sample locations (e.g. Trout Lake) as
random intercepts. Details of statistics see Table S1.

Strength of eco-phenotypic feedback linkages under biotic and

abiotic drivers. The strength of feedback linkages (i.e. size →abundance or

abundance →size) was quantified by CCM based on cross-mapping skill ρ (i.e.,

correlation coefficient between observations and CCM predictions) using the

maximal training set (Sugihara *et al.* 2012). To ensure comparability across

101 ecosystems, we accounted for variation in cross-map skill ρ arising from

differences level of noise in time-series observed in each ecosystem. Following

(Chang *et al.* 2020), linkage strength was therefore standardized by dividing

each value by the maximum cross-map ρ within a system, yielding scaled

strengths between 0 and 1 that reflect the relative importance of each causal

link (Chang *et al.* 2022).

Next, we employed a generalized linear mixed model using the glmmTMB

function from the *glmmTMB* package (Brooks *et al.* 2017) with a Gaussian error

distribution to test the main effects of warming, mean temperature, species

richness, and mean species body mass on linkage strength—either from

phenotypic traits (body size) to abundance (size → abundance) or from

abundance to body size (abundance → size) (Table S2). Again, to ensure

normality, mean temperature, species richness, and mean species body mass

were log10-transformed, and all of these predictors were normalized using the

scale function in base R. Species groups (e.g., plankton) nested within

sampling locations (e.g., Trout Lake) were included as random intercepts. Full

statistical results are provided in Table S2.

All analyses were done with R (ver. 4.1.2). The CCM analysis was

examined using the rEDM package (version 1.2.3).

Results

We first used a binary response variable to indicate the presence or

absence of feedback linkages between the phenotypic trait (body size) and

population abundance, with links identified by Convergent Cross Mapping

(CCM). Across 101 ecological systems monitored for 10 to 47 years, our

analysis showed that warming weakened eco-phenotypic feedback loop, as
evidenced by a negative association between warming and the proportion of
both linkages (abundance → size and size → abundance) (Fig 2a, 2e). Species
richness likewise weakened the feedback loop (Fig 2b, 2f), whereas mean
species body mass strengthened it (Fig 2c, 2g). However, mean temperature
showed no effect on either linkage of abundance → size or size → abundance
(Fig 2d, 2h).

313   Within taxonomic groups (plankton, fish, or terrestrial species), the
314   effects of species richness and mean species body mass on eco-phenotypic
315   feedback loops were taxon-specific, whereas warming consistently weakened
316   feedback strength across all taxa (Fig 2). In most taxa, species richness
317   weakened the eco-phenotypic feedback loop (Fig 2b, 2f), whereas mean species
318   body mass strengthened it (Fig 2c, 2g).

319   Last, we used the strength of feedback linkages, as quantified by CCM,
as the response variable. Across 101 ecological systems, mean temperature
only increased the strength of the size → abundance linkage (Fig 3d). In
contrast, warming, species richness, and mean species body mass had no
effects on the strength of either linkage (abundance → size or size →
abundance) (Fig 3). Taken together, these results indicate that warming and
species richness weakened eco-phenotypic feedback loops by reducing the
proportion of both linkage types rather than altering their strength, whereas
mean species body mass strengthened the feedback loop by increasing them.
328

Discussion

330   Our study provides the first systematic evaluation of eco-phenotypic
feedback loop in response to climate warming and biotic drivers. Using long-
term field data from 101 ecosystems—each spanning multiple generations per
species and providing sufficient temporal durations to detect changes in both
phenotypic traits and population abundance—we found that warming and
species richness generally weakened eco-phenotypic feedback loops, whereas
mean species body size strengthened them. These effects were driven by

changes in the proportion of both linkage types (abundance → size or size →
abundance).

The phenotypic trait of body size respond rapidly to environmental
change through phenotypic plasticity or evolutionary adaptation (Merckx *et al.*
2018) and exerts a substantial influence on population abundance (Martins *et*
al. 2023). Previous experimental studies have documented that the eco-
phenotypic feedback are tightly linked in single-species populations (Gibert *et*
al. 2022, 2023), a two-species competitive interactions (Govaert & Klauschies
2024), and a microbial food web (Han *et al.* 2023). Our study further showed
that warming tended to weaken the feedback loop (Fig 2).

Previous evidence has documented weaker eco-phenotypic feedback at
higher temperatures from an experimental protist food web (Han *et al.* 2023).
This may be attributed to the differing response rates of phenotypic trait of
body size and abundance to warming (Genner *et al.* 2010). Higher
temperatures can lead to a significant reduction in species body size
(Audzijonyte *et al.* 2020; Terry *et al.* 2022), while it can have slight effect on
population abundance (Sebastian *et al.* 2012). Alternatively, higher
temperatures have shown to accelerate growth, leading to an rapid increase in
population abundance (Brandenburg *et al.* 2019; Han *et al.* 2023), while
simultaneously either leaving body size unaffected or slightly altered body size
(Crozier *et al.* 2010; Genner *et al.* 2010). These different responses rate among
body size and abundance may weaken the feedback between them (Han *et al.*
2023).

We found that higher species richness also weakens eco-phenotypic
feedback loop. Previous experimental studies have similarly shown that
increased species richness reduces the proportion of links from phenotypic
traits body size to abundance (body size→ abundance) (Govaert & Klauschies
2024). Here, enhanced species richness can subject organisms to a broader
spectrum of ecological interactions, including interspecific competition and
predation, which may disrupt the feedback between size and abundance

(Govaert & Klauschies 2024). Evidence has documented that adding

interspecific competition and predation can weaken the link strength between

them (Govaert & Klauschies 2024; Han *et al.* 2023).

We found that mean species body size strengthens the eco-phenotypic

feedback loop. To date, no direct empirical evidence has been documented to

validate this trend. Species with larger body mass exhibit longer generation

times and slower population turnover (Gillooly 2000b; White *et al.* 2007),

allowing both body size and population abundance to respond more gradually

to environmental changes. This extended adaptation period may contribute to a

more stable and synchronized relationship between body size and population

abundance over time.

In summary, by using long-term field data covering multiple generations

for each species, our findings show that climate warming weakens eco-

phenotypic feedback loops, highlighting their vulnerability to anthropogenic

pressures. Understanding how eco-phenotypic feedback responds to

environmental change may help predict trait and population abundance

dynamics in a warming world.

Competing interests

The authors declare no competing interests.

Author Contributions

Q.H.Z., F.D.L. and Y.X.G.W designed the research. Q.H.Z. and Y.X.G.W

performed the study and analyzed the data. Q.H.Z and G.S performed the CCM

analysis. Q.H.Z., F.D.L. and Y.X.G.W led the writing, with contributions from

C.X. and G.S.

Data and code availability

The data reproducing all the results in this study are available on GitHub

(https://github.com/QZhao16/link.abundance-size).

Acknowledgements

We thank all participants in each long-term monitoring site. Q.Z. acknowledges

funding from the concerted research action (ARC) from the special research

fund from the University of Namur, Belgium (ARC grant DIVERCE, Convention

18/23-095).

Figures

Figure 1. Overview of methods and sample locations. a, Sample locations

for plankton, fish, and terrestrial species. **b**, Conceptual illustration of the eco-

phenotypic feedback loop. Evolutionary and plastic shifts in phenotypic traits

such as body size influence demographic processes—survival, reproduction,

and competitive ability—that can ultimately alter population abundance. In

turn, changes in abundance modify phenotypic traits through exerting

selection pressures and inducing plastic responses. This reciprocal interaction

between phenotypic traits and ecological dynamics constitutes the eco-

phenotypic feedback. In this study, we investigate eco-phenotypic feedback at

the level of individual focal species (e.g., *Gambusia affinis*) rather than across

species.

 Figure 2. Presence/absence of eco-phenotypic feedback linkages under

 biotic and abiotic drivers. a–h, Presence (1) or absence (0) of feedback

 linkages in response to warming, species richness, mean species body mass

 and mean temperature. Link presence was identified using convergent cross

 mapping (CCM) and coded as 1, while absence was coded as 0. Solid black

 lines with shaded error bands indicate significant best-fit trendlines and

 corresponding 95% confidence intervals from two-sided logistic regressions

 across 101 ecosystems; dashed black lines indicate nonsignificant

 relationships. Non-bold colored dashed lines show logistic model trendlines

 within each taxon (plankton, fish, or terrestrial species). Points represent

 binary link values (0 or 1) between abundance and the phenotypic trait (body

 size) for each species across 101 ecosystems, with vertical jitter added to

 reduce overlap. Statistical results are provided in Table S1.

 Figure 3. Strength of eco-phenotypic feedback linkages under biotic and abiotic drivers. a–h, Strength of feedback linkages in response to warming, species richness, mean species body mass and mean temperature. Link strength was quantified using convergent cross mapping (CCM), with larger values indicating stronger feedback linkages. Solid black lines with shaded error bands indicate significant best-fit trendlines and corresponding 95% confidence intervals from two-sided regressions across 101 ecosystems; dashed black lines indicate nonsignificant relationships. Statistical results are provided in Table S2.

Supplementary Tables

Table S1. **Presence/absence of eco-phenotypic feedback linkages under biotic**

and abiotic drivers. A generalized linear mixed model (glmmTMB function in the

glmmTMB package) with a binomial error distribution was used to test the main effects

of warming, mean temperature, species richness, and mean species body mass on the

presence/absence of feedback linkages (size → abundance or abundance → size). The

response variable was binary, with 1 indicating the presence of a feedback linkage

identified using convergent cross mapping (CCM) and 0 indicating its absence. Species

groups (e.g., fish) nested within sampling locations (e.g., Trout Lake) were included as

random intercepts. Mean temperature, species richness, and mean species body mass

were log10-transformed to improve model performance. ***P< 0.001, **P<0.01,

*P<0.05.

	presence/absence of body size → abundance			presence/absence of abundance → body size		
	coefficient	Z	P value	coefficient	Z	P value
(Intercept)	-1.10	-4.95	**<8e-8*****	-1.27	-11.92	**<2e-16*****
Species richness	-0.99	-3.31	**<0.0001*****	-3.44	-9.45	**<2e-16*****
Warming	-0.99	-4.65	**<4e-6*****	-2.87	-11.72	**<2e-16*****
Mean body mass	0.81	2.66	**0.008****	3.10	8.59	**<2e-16*****
Mean Temperature	0.12	1.09	0.28	0.11	1.12	0.26

Table S2. **Strength of eco-phenotypic feedback linkages under biotic and abiotic**

drivers. A generalized linear mixed model (glmmTMB function in the *glmmTMB*

package) with a gaussian error distribution was used to test the main effects of

warming, mean temperature, species richness, and mean species body mass on the

strength of feedback linkages (size → abundance or abundance → size). Strength of

feedback linkages was quantified using convergent cross mapping (CCM), with larger

values indicating stronger feedback linkages. Species groups (e.g., fish) nested within

sampling locations (e.g., Trout Lake) were included as random intercepts. Mean

temperature, species richness, and mean species body mass were log10-transformed

to improve model performance. ***P< 0.001, **P<0.01, *P<0.05.

	Strength of body size → abundance			Strength of abundance → body size		
	coefficient	Z	P value	coefficient	Z	P value
(Intercept)	1.80	15.75	**<2e-16*****	2.06	16.60	**<2e-16**
Species richness	-0.17	-1.53	0.13	0.01	0.05	0.96
Warming	-0.14	-1.04	0.30	-0.25	-1.61	0.11
Mean body mass	-0.02	-0.14	0.89	0.15	1.04	0.30
Mean Temperature	0.33	3.00	**0.002****	0.17	1.40	0.16

References

Audzijonyte, A., Richards, S.A., Stuart-Smith, R.D., Pecl, G., Edgar, G.J., Barrett, N.S., *et al*. (2020). Fish body sizes change with temperature but not all species shrink with warming. *Nat Ecol Evol*, 4, 809–814.

Bayne, B.L. & Worrall, C.M. (1980). Growth and Production of Mussels Mytilus edulis from Two Populations. *Marine Ecology Progress Series*, 3, 317–328.

Bergquist, A.M., Carpenter, S.R. & Latino, J.C. (1985). Shifts in phytoplankton size structure and community composition during grazing by contrasting zooplankton assemblages. *Limnology and Oceanography*, 30, 1037–1045.

Brandenburg, K.M., Velthuis, M. & Van de Waal, D.B. (2019). Meta-analysis reveals enhanced growth of marine harmful algae from temperate regions with warming and elevated CO2 levels. *Glob Chang Biol*, 25, 2607–2618.

Brooks, M., Bolker, B., Kristensen, K., Maechler, M., Magnusson, A., Skaug, H., *et al*. (2017). glmmTMB: Generalized Linear Mixed Models using Template Model Builder.

Chang, C.-W., Miki, T., Ye, H., Souissi, S., Adrian, R., Anneville, O., *et al*. (2022). Causal networks of phytoplankton diversity and biomass are modulated by environmental context. *Nat Commun*, 13, 1140.

Chang, C.-W., Ye, H., Miki, T., Deyle, E.R., Souissi, S., Anneville, O., *et al*. (2020). Long-term warming destabilizes aquatic ecosystems through weakening biodiversity-mediated causal networks. *Global Change Biology*, 26, 6413–6423.

Christensen, T.R., Topp-Jørgensen, E., Sejr, M.K. & Schmidt, N.M. (2017). Foreword: Synthesis of the Greenland Ecosystem Monitoring program. *Ambio*, 46, 1–2.

Cohen, R.E. & Lough, R.G. (1981). Length-Weight Relationships for Several Copepods Dominant in the Georges Bank-Gulf of Maine Area. *J. Northw. Atl. Fish. Sci.*, 2, 47–52.

Conover, D.O. & Munch, S.B. (2002). Sustaining Fisheries Yields Over Evolutionary Time Scales. *Science*, 297, 94–96.

Crozier, L.G., Zabel, R.W., Hockersmith, E.E. & Achord, S. (2010). Interacting effects of density and temperature on body size in multiple populations of Chinook salmon. *Journal of Animal Ecology*, 79, 342–349.

Deyle, E.R. & Sugihara, G. (2011). Generalized Theorems for Nonlinear State Space Reconstruction. *PLOS ONE*, 6, e18295.

Dornelas, M., Antao, L., Moyes, F., Bates, A., Magurran, A., Adam, D., *et al*. (2018). BioTIME: A database of biodiversity time series for the Anthropocene.

Edgar, G.J. & Stuart-Smith, R.D. (2014). Systematic global assessment of reef fish communities by the Reef Life Survey program. *Sci Data*, 1, 140007.

Elbrächter, M. (1976). Population dynamic studies on phytoplankton cultures. *Marine Biology*, 35, 201–209.

Felden, J., Möller, L., Schindler, U., Huber, R., Schumacher, S., Koppe, R., *et al*. (2023). PANGAEA - Data Publisher for Earth & Environmental Science. *Sci Data*, 10, 347.

Fish trawl surveys. (2025). . Available at: https://www.ices.dk/data/dataset-collections/Pages/Fish-trawl-survey.aspx. Last accessed 30 April 2025.

Gardner, J.L., Peters, A., Kearney, M.R., Joseph, L. & Heinsohn, R. (2011). Declining body
size: a third universal response to warming? *Trends in Ecology & Evolution*, 26, 285–
291.
Genner, M.J., Sims, D.W., Southward, A.J., Budd, G.C., Masterson, P., Mchugh, M., *et al.*
(2010). Body size-dependent responses of a marine fish assemblage to climate
change and fishing over a century-long scale. *Global Change Biology*, 16, 517–527.
GHCN-Daily, Version 3. (2012). Global Historical Climatology Network - Daily (GHCN-
Daily), Version 3.
Gibert, J.P., Han, Z.-Y., Wieczynski, D.J., Votzke, S. & Yammine, A. (2022). Feedbacks
between size and density determine rapid eco-phenotypic dynamics. *Functional*
Ecology, 36, 1668–1680.
Gibert, J.P., Wieczynski, D.J., Han, Z.-Y. & Yammine, A. (2023). Rapid eco-phenotypic
feedback and the temperature response of biomass dynamics. *Ecology and*
Evolution, 13, e9685.
Gillooly, J.F. (2000a). Effect of body size and temperature on generation time in
zooplankton. *Journal of Plankton Research*, 22, 241–251.
Gillooly, J.F. (2000b). Effect of body size and temperature on generation time in
zooplankton. *Journal of Plankton Research*, 22, 241–251.
Gosling, E. (2003). *Bivalve Molluscs: Biology, Ecology and Culture*. Blackwell Publishing
Ltd.
Govaert, L. & Klauschies, T. (2024). Eco-phenotypic feedback loops differ in multistressor
environments. *Ecology*, e4480, e4480.
Han, Z.-Y., Wieczynski, D.J., Yammine, A. & Gibert, J.P. (2023). Temperature and nutrients
drive eco-phenotypic dynamics in a microbial food web. *Proceedings of the Royal*
Society B: Biological Sciences, 290, 20222263.
Hsieh, C., Glaser, S.M., Lucas, A.J. & Sugihara, G. (2005). Distinguishing random
environmental fluctuations from ecological catastrophes for the North Pacific
Ocean. *Nature*, 435, 336–340.
Huntley, M.E. (1992). Temperature-Dependent Production of Marine Copepods: A Global
Synthesis. *The American Naturalist*, 140, 201–242.
Hutchings, J.A. & Reynolds, J.D. (2004). Marine Fish Population Collapses: Consequences
for Recovery and Extinction Risk. *BioScience*, 54, 297–309.
Karakoç, C., Clark, A.T. & Chatzinotas, A. (2020). Diversity and coexistence are influenced
by time-dependent species interactions in a predator–prey system. *Ecology Letters*,
23, 983–993.
Kingsolver, J.G. & Pfennig, D.W. (2007). Patterns and Power of Phenotypic Selection in
Nature. *BioScience*, 57, 561–572.
Kirk, V.M. (1975). Biology of Pterostichus chalcites1, a Ground Beetle of Cropland2. *Ann*
Entomol Soc Am, 68, 855–858.
Law, R. (2000). Fishing, selection, and phenotypic evolution. *ICES J Mar Sci*, 57, 659–668.
Litchman, E. (2000). Growth rates of phytoplankton under fluctuating light. *Freshwater*
Biology, 44, 223–235.
Magnuson, J., Carpenter, S. & Stanley, E. (2023a). North Temperate Lakes LTER:
Phytoplankton - Trout Lake Area 1984 - 2006.

Magnuson, J., Carpenter, S. & Stanley, E. (2024a). North Temperate Lakes LTER: Zooplankton - Madison Lakes Area 1997 - current.

Magnuson, J.J., Carpenter, S.R. & Stanley, E.H. (2022). North Temperate Lakes LTER: Phytoplankton - Madison Lakes Area 1995 - current.

Magnuson, J.J., Carpenter, S.R. & Stanley, E.H. (2023b). North Temperate Lakes LTER: Zooplankton - Trout Lake Area 1982 - current.

Magnuson, J.J., Carpenter, S.R. & Stanley, E.H. (2024b). North Temperate Lakes LTER: Fish Abundance 1981 - current.

Magnuson, J.J., Carpenter, S.R. & Stanley, E.H. (2024c). North Temperate Lakes LTER: Fish Lengths and Weights 1981 - current.

Martins, I.S., Schrodt, F., Blowes, S.A., Bates, A.E., Bjorkman, A.D., Brambilla, V., *et al*. (2023). 0. Widespread shifts in body size within populations and assemblages. *Science*, 381, 1067–1071.

Matalin, A.V. (2007). Typology of life cycles of ground beetles (Coleoptera, Carabidae) in Western Palaearctic. *Entomological Review*, 87, 947–972.

McLean, N., Lawson, C.R., Leech, D.I. & van de Pol, M. (2016). Predicting when climate-driven phenotypic change affects population dynamics. *Ecology Letters*, 19, 595–608.

Merckx, T., Souffreau, C., Kaiser, A., Baardsen, L.F., Backeljau, T., Bonte, D., *et al*. (2018). Body-size shifts in aquatic and terrestrial urban communities. *Nature*, 558, 113–116.

Michener, W., Vieglais, D., Vision, T., Kunze, J., Cruse, P. & Janée, G. (2011). DataONE: Data Observation Network for Earth - Preserving Data and Enabling Innovation in the Biological and Environmental Sciences. *D-Lib Magazine*, 17, 3.

Mohsin, T. & Gough, W.A. (2010). Trend analysis of long-term temperature time series in the Greater Toronto Area (GTA). *Theor Appl Climatol*, 101, 311–327.

Nejad, E.S., Cloern, J. & Schraga, T. (2017). Phytoplankton Species Composition, Abundance and Cell Size in San Francisco Bay: Microscopic Analyses of USGS Samples Collected 1992-2014.

Okamura, B. (1986). Group living and the effects of spatial position in aggregations of Mytilus edulis. *Oecologia*, 69, 341–347.

Parmesan, C. (2006). Ecological and Evolutionary Responses to Recent Climate Change. *Annual Review of Ecology, Evolution, and Systematics*, 37, 637–669.

Peters, R.H. (1983). *The Ecological Implications of Body Size*. Cambridge Studies in Ecology. Cambridge University Press, Cambridge.

Pinsky, M.L., Hillebrand, H., Chase, J.M., Antão, L.H., Hirt, M.R., Brose, U., *et al*. (2025). Warming and cooling catalyse widespread temporal turnover in biodiversity. *Nature*, 1–5.

Rietkerk, M., Bastiaansen, R., Banerjee, S., van de Koppel, J., Baudena, M. & Doelman, A. (2021). Evasion of tipping in complex systems through spatial pattern formation. *Science*, 374, eabj0359.

Rivard, I. (1964). Observations on the breeding periods of some ground beetles (coleoptera: carabidae) in eastern ontario. *Can. J. Zool.*, 42, 1081–1084.

Schwarz, B., Barnes, A.D., Thakur, M.P., Brose, U., Ciobanu, M., Reich, P.B., *et al*. (2017).
Warming alters energetic structure and function but not resilience of soil food webs.
Nature Clim Change, 7, 895–900.

Sebastian, P., Stibor, H., Berger, S. & Diehl, S. (2012). Effects of water temperature and
mixed layer depth on zooplankton body size. *Mar Biol*, 159, 2431–2440.

Sugihara, G., May, R., Ye, H., Hsieh, C., Deyle, E., Fogarty, M., *et al*. (2012). Detecting
causality in complex ecosystems. *Science*, 338, 496–500.

Sugihara, G. & May, R.M. (1990). Nonlinear forecasting as a way of distinguishing chaos
from measurement error in time series. *Nature*, 344, 734–741.

Terry, J.C.D., O'Sullivan, J.D. & Rossberg, A.G. (2022). No pervasive relationship between
species size and local abundance trends. *Nat Ecol Evol*, 6, 140–144.

Tilman, D. (1999). The Ecological Consequences of Changes in Biodiversity: A Search for
General Principles. *Ecology*, 80, 1455–1474.

Tong, Y., Feng, L., Wang, X., Pi, X., Xu, W. & Woolway, R.I. (2023). Global lakes are warming
slower than surface air temperature due to accelerated evaporation. *Nat Water*, 1,
929–940.

Van De Koppel, J., Gascoigne, J.C., Theraulaz, G., Rietkerk, M., Mooij, W.M. & Herman,
P.M.J. (2008). Experimental Evidence for Spatial Self-Organization and Its Emergent
Effects in Mussel Bed Ecosystems. *Science*, 322, 739–742.

Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T.J.C., *et al*. (2002).
Ecological responses to recent climate change. *Nature*, 416, 389–395.

Wellborn, G.A. (1994). Size-Biased Predation and Prey Life Histories: A Comparative Study
of Freshwater Amphipod Populations. *Ecology*, 75, 2104–2117.

White, E., Ernest, S.K.M., Kerkhoff, A. & Enquist, B. (2007). Relationships between body size
and abundance in ecology. *Trends in ecology & evolution*, 22, 323–30.

Ye, L., Chang, C.-W., Matsuzaki, S.S., Takamura, N., Widdicombe, C.E. & Hsieh, C. (2019).
Functional diversity promotes phytoplankton resource use efficiency. *Journal of
Ecology*, 107, 2353–2363.

Zhao, Q., Van den Brink, P.J., Xu, C., Wang, S., Clark, A.T., Karakoç, C., *et al*. (2023).
Relationships of temperature and biodiversity with stability of natural aquatic food
webs. *Nat Commun*, 14, 3507.

626