Warming is associated with temporal decoupling of species' body size and abundance

Qinghua Zhao[1,2,3,7], Yingying X. G. Wang[4,7], Chi Xu[5], George Sugihara[6], Frederik De

Laender[1,2,3,7]

[1]Research Unit of Environmental and Evolutionary Biology (URBE), University of Namur,

Namur, Belgium.

[2]Institute of Complex Ecosystems (naXys), University of Namur, Namur, Belgium.

[3]Institute of Life, Earth and the Environment (ILEE), University of Namur, Namur, Belgium.

[4]School of Veterinary Medicine, University of California-Davis, Davis, CA, USA

[5]School of Life Sciences, Nanjing University, Nanjing, 210023, China

[6]Scripps Institution of Oceanography, University of California-San Diego, La Jolla, CA, USA

[7]These authors contributed equally

Email: yingying.xgz.wang@gmail.com

Abstract

Across species, body size and population abundance are often assumed to be tightly linked, with

smaller species reaching higher abundance. Consequently, either size or abundance is commonly

used as a proxy to assess responses to abiotic and biotic environmental changes and to inform

conservation strategies. However, whether the tight relationship holds within species remains

unclear. Here, we analyzed 41 natural ecosystems spanning 10 to 48 years and found that species

abundance and size are not tightly linked over time. Only 14.0% of species abundance was

causally linked to size, while 21.5% of size was causally linked to abundance. Only 7.7% of

species had bidirectional causal links among size and abundance. Warming was associated with

decoupling between abundance and size. Our findings suggest that species' abundance and size

tend to be decoupled over time under climate change. Thus, abundance and size time series can

provide distinct insights into species' responses to climate change and should be considered

separately in ecosystem conservation and stability assessments.

Keywords: Body size, Population abundance, Climate change, Ecosystem conservation,

Stability assessment

Introduction

Body size as one of the most crucial functional traits is typically closely linked to population abundance[1–3]. Much evidence shows that species with larger body sizes tend to result in lower abundances, following an inverse power law[1,2]. Essentially, this relationship is often attributed to differences in metabolic rates among organisms of varying sizes[1,2,4]. Larger species generally have lower metabolic rates and invest more energy in growth and maintenance[1,2]. This life-history tradeoff results in lower reproductive frequency, and thus longer generational intervals, which limits population size[5,6]. Larger producer species, such as phytoplankton, has lower nutrient conversion efficiency due to their surface-area-to-volume ratio, resulting in lower population abundance[7]. In the context of food webs, both theoretical and empirical work has well demonstrated that species at higher trophic levels tend to have larger body sizes and smaller population sizes[8–12] with limited dietary niche breaths and higher extinction risk[13]. However, the effect of body size on abundance has predominantly been documented across species, mostly based on snapshot samples. It is unclear whether a similar effect exists within species, across time series. In addition, while experiments have shown that body size changes over time can strongly influence population density over time within protist species[5,14–16], it remains uncertain whether the effect of size on abundance holds tightly in natural ecosystems.

The body size-abundance relationships could be *bidirectional* through time, and both directional effects may not be of equal strength[5,14]. Plastic or evolutionary shifts in body size can influence fitness and therefore population abundance, which in turn can affect size, creating feedback loops[5,17]. However, compared to the well-documented influence of size on abundance, the effect of abundance on body size time series are not well understood[5]. Such effect can in principle occur through eco-evolutionary feedback[17], phenotypic plasticity[5,14], and demographic processes[18,19]. For example, high population abundance can result in crowding and increased intraspecific competition[20,21]. Stronger competition can reduce resource availability, potentially resulting in growth limitation and smaller body sizes[22,23]. Indeed, changes in abundance affecting body size have been documented in indoor experiments[5,14]. However, this evidence has not yet been fully translated to natural ecosystems. Recent field studies suggest declines or increases in species abundance either had no consistent association with changes in body size[24] or showed complex associations[25,26]. However, these field studies focus on trends in body size across entire communities which thereby obscures the putative size-abundance relationships within species.

Therefore, it remains unclear whether abundance time series can influence body size time series.

Addressing this question is essential, as it determines whether species abundance time series

alone can reliably represent its body size dynamics in assessing species' responses to

environmental changes.

In nature, changes of abundance and body size often co-occur with environmental shifts

or community changes[5,27–29]. Factors such as warming[30–32], species diversity, and species traits

can potentially alter both body size and abundance. This raises the question if these factors will

change the established relationships between body size and population abundance. In this study,

we investigated whether body size and abundance were bidirectionally linked by analyzing data

from 41 natural ecosystems. The presence of bidirectional links (size → abundance, as well as

abundance → size) was evaluated using convergent cross mapping (CCM), a technique to detect

causal links (Fig. 1a). Then, we examined the proportion of bidirectional links in responses to

warming and biotic drivers. Our findings suggest that body size and abundance are only weakly

linked within species, and that warming further weakens this coupling. These results underscore

the importance of incorporating both species size and abundance into monitoring and

conservation efforts.

Results

Our results from CCM showed that bidirectional link between species' abundance and

size were not common within individual species (Fig 1b). Across 41 ecosystems, including

aquatic plankton and fish, and terrestrial species (primarily beetles)—a total of 1269 species

were analyzed. Of these, 21.5% ± 2.8% (mean ± s.e.) of the species had link from size to

abundance (size → abundance), while 14.0% ± 2.3% had link from abundance to size

(abundance → size) (Fig 1b). Only 7.7% of species had bidirectional causal links between size

and abundance. In addition, within plankton, 11.5% ± 3.6% of the species had size →

abundance, and 16.1 % ± 5.1% had abundance → size (Fig 1b). Within fish (and terrestrial

species), the proportions were 30.8% ± 7.4% (21.1% ± 3.1%) and 16.4% ± 6.5% (13.2% ±

1.8%), respectively (Fig 1b).

The link between size and abundance was system-dependent (Fig 1c-d). In four out of all

41 ecosystems, the proportion of link size → abundance exceeded 50% (Fig 1d; FigS1-S2).

Among the four systems, the three were aquatic fish ecosystems (i.e. Smokelumne River-

Georgiana Slough, Sacramento River-Koket, and American River), while the remaining one was

a terrestrial beetle ecosystem. In contrast, in just one system out of 41 systems, the proportion of

link abundance → size surpassed 50% (Fig 1c; FigS1-S2) and the one was again a fish

ecosystem (i.e. Smokelumne River-Georgiana Slough). These results indicate that species' body

size and abundance tend to be temporally decoupled rather than tightly linked. These conclusions

were not affected by including rare species—which are normally excluded from similar analyses

(Fig. S3-S5), and neither when only using the dominant species for analysis (Fig. S6-S8).

Across 41 systems, warming was negatively correlated with the proportion of both links

of abundance → size and size → abundance (Fig 2a, d). In contrast, mean species body mass was

only positively associated with the proportion of links of size → abundance (Fig 2c, f), whereas

species richness only had negative correlation with the link (Fig 2b, e). In addition, we found

mean temperature had no effect on either link of abundance → size or size → abundance (Table

S1). Within species group (plankton, fish or terrestrial species), the species richness and mean

body mass on proportion of link size → abundance were system dependent (Fig. 2). In most

species groups, species richness had negative effects on link of size → abundance (Fig. 2b).

Conversely, mean species body mass was positively correlated with link of size → abundance in

one out of three groups (Fig. 2c). Warming had negative effects on both links within each species

group (Fig. 2a, d).

Discussion

In the studied ecosystems, the bidirectional links between abundance and body size were

weak, with 14.0% of species exhibiting link of abundance → size and 21.5% exhibiting link of

size → abundance. Moreover, warming was associated with lower proportions of both links of

abundance → size and of size → abundance. These results suggest that species abundance and

size are not tightly linked over time, but tend to be independent under warming.

Body size responds rapidly to environmental changes[33] and can exert a significant

influence on population abundance[25]. Previous experimental studies have demonstrated that

temporal changes in species' body size are tightly associated with abundance in single-species

populations[5,14], a two-species competitive interactions[34], and a microbial food web[16]. However,

in natural ecosystems, we found the proportion of link size → abundance to be weak. The

weakening of the link was associated with warming and increased species richness (Fig 2).

Experimental evidence had documented weaker causal link of size → abundance at higher

temperatures[16]. This may be attributed to the different responses of size and abundance to

warming[26]. Higher temperatures can lead to a reduction in species body size[35], while it can have

no significant effect on population abundance[36]. Similarly, higher temperatures have shown to

accelerate growth, leading to an rapid increase in population abundance[16,37], while

simultaneously either leaving body size unaffected or slightly altered body size[26,38]. These

different responses among size and abundance can weaken the link between them[16]. Increased

species richness also reduced the proportion of the link of size → abundance. Enhanced species

richness may subject organisms to a broader spectrum of ecological interactions, such as

intensified interspecific competition and predation, which can disrupt the link of size →

abundance within species[34]. Indeed, evidence has documented that adding interspecific

competition can weaken the link strength of size → abundance[34]. We also found mean species

body mass increased the proportion of link size → abundance. To date, no direct empirical

evidence has been documented to validate this trend. However, evidence documented that

species with larger body mass exhibit longer generation times and slower population turnover[1,39],

allowing both body size and population abundance to respond more gradually to environmental

changes. This extended adaptation period may facilitate the temporal coupling relationship

between body size and population abundance. In addition, we found that the proportion of link

abundance→ size is also weak, and warming is associated with a lower proportion of the link.

This contrasts with previous experimental studies that demonstrated a strong relationship within

protist species[5]. Again, the weakening of the link driven by warming may be attributed to

different responses of size and abundance to warming[26,34].

The bidirectional feedback loop between body size and abundance dynamics can be

asymmetric. Indeed, previous experimental evidence has shown a stronger causal link of size →

abundance than the other way around[5,14]. Our results showed that the proportion of body size →

abundance is slightly higher than the reverse (21.5% vs. 14.0%). These findings suggest that

changes in body size may occur independently of changes in abundance rather than tightly

coupled. Indeed, body size and abundance have showed independent changes, particularly in

responses to environmental changes such as urbanization[33] and habitat fragmentation[40,41].

Species' size can even provide greater accuracy and serve as an earlier warning signals for

predicting population collapses than abundance, under environmental deterioration[42–44].

Furthermore, the proportion of link among size and abundance was species dependent (Fig. S9).

84.7% of species exhibited a proportion of links among size and abundance below 50%, while

only 4.1% had a proportion of links equal to 100%, including beetle species within the

Geotrupidae taxon (Fig. S9).

In summary, our findings reveal that species size and abundance time series are

decoupled over time and warming tends to accelerate this decoupling. The conventional

paradigm positing a robust cross-species association between size and abundance is not upheld

within species. Our results underscore the potential limitation of relying on either body size or

abundance time series alone to examine species' responses to warming and measure ecosystem

resilience.

Methods

Aquatic plankton Data. Species size and abundance data were compiled from a total of

10 long-term monitoring sites in aquatic planktonic ecosystems, including nine lakes and one

marine system, as detailed in Supplementary Table S2: (1) Wadden sea, EU, 2001-2010; (2) Lake

Mendota, USA, 1995-2005; (3) Lake Monona, USA, 1995-2005; (4) Lake Allequash, USA,

2009-2018; (5) Lake Big Muskellunge, USA, 1991-1999; (6) Lake Crystal Bog, USA, 2008-

2018; (7) Lake Crystal, USA, 1990-1999; (8) Lake Sparkling, USA, 1990-1999; (9) Trout Lake,

USA, 1990-1999; (10) Lake Trout Bog, USA, 2009-2018. For each aquatic planktonic

ecosystem, there were three types of variables: (1) species abundance; (2) species size (i.e. body

length, biovolume, or biomass); (3) water temperature. For species size (2), a single metric is

reported for each species and subsequently used for further analysis.

Next, for consistency, each species' time series were standardized to seasonal sampling

frequencies (if multiple samplings were conducted per season, e.g., monthly and bimonthly

sampling data sets, those were averaged). Here, the seasonal resolution (trimonthly) is the shared

consensus that can be applied to all 10 data sets, and the seasonal average here is also the most

representative measure across all data sets, because the equal sampling interval is necessary for

Convergent cross mapping (CCM) analysis. After the seasonal average, there were 0.31%

missing points across the whole planktonic dataset (41 out of 17032 data points). Those missing

data points were linearly interpolated using *na.approx* function in the package of zoo[45,46]. Then,

the species abundance across all data sets was scaled to the same unit (individuals per litre).

Species length, biovlume and biomass were standardized to the same unit (μm), (μm^3) and

(mg/m^3) respectively.

Last, we removed the long term linear and seasonal trend from each time series.

Specifically, we first removed the long-term linear trend from each time series by using the

residuals from a linear regression against time[47,48]. Next, we removed seasonality by scaling

against the mean and standard deviation of values occurring in the same year, $D_{season}(t) =$

$(O(t) - \mu_{season})/\sigma_{season}$, where $D_{season}(t)$ is the deseasoned time series, $O(t)$ is the original

time series, μ_{season} is the year mean, and the σ_{season} is the year standard deviation[47,48]. Then, we

selected species time series with number of zeros <60% because too many zero values may cause

statistical problems in time series analysis. Finally, each time series (both species abundance and

size) was scaled to zero mean and unit variance.

Fish Data. Fish size and abundance data were compiled from 11 monitoring sites

(Supplementary Table S3), including six rivers and five lakes, as detailed below: (1) American

River, USA, 1976-2023; (2) San Joaquin River (Elkhorn), USA, 1981-2023; (3) Sacramento

River (Koket), USA, 1976-2023; (4) Keller Beach, USA, 1980-2023; (5) Smokelumne River

(Georgiana Slough), USA, 1976-2023; (6) Old river (Franks Tract), USA, 1993-2023; (7) Lake

Allequash, USA, 1981-2018; (8) Lake Big Muskellunge, USA, 1981-2018; (9) Lake Mendota,

USA, 1981-2018; (10) Lake Sparkling, USA, 1981-2018; (11) Trout Lake, USA, 1981-2018.

Again, for each Fish ecosystem, there were three types of variables: (1) species abundance; (2)

species size (i.e. body length); (3) water temperature.

For consistency each species time series of abundance and size were standardized to the

yearly sampling frequency (if multiple samplings were conducted per year, e.g., quarterly

sampling data sets, those were averaged). Again, the yearly resolution is shared across all data

sets. After the yearly average, there were no missing points across the fish dataset. Then, the fish

abundance across all data sets was standardized to the same unit (caught per effort). Species

length was standardized to the same unit (mm). As all fish data were standardized to yearly

intervals, there was no need to remove long-term linear or seasonal trends from each time series.

Therefore, the original data were utilized for further analysis. Again, we selected species time

series with number of zeros <60%. Finally, each time series (both species abundance and size)

was scaled to zero mean and unit variance.

Terrestrial species. Terrestrial species size and abundance data were compiled from 20

monitoring sites (Supplementary Table S4). Of these, 18 sites are ground beetles, while one site

monitored lizard populations and another monitored rat species. Again, for each site, there were

three types of variables: (1) species abundance; (2) species size (i.e. biomass, mg/m^3); (3)

surface temperature. Each site reports both abundance and size. The daily temperature and

precipitation for each site were retrieved from NOAA's National Centers for Environmental

Information[49].

Next, to ensure consistency, each species' time series were standardized to seasonal

sampling frequencies (if multiple samplings were conducted per season, those were averaged).

After calculating seasonal averages, 5.4% of data points were missing in the dataset. These

missing values were subsequently linearly interpolated using the na.approx function from the zoo

package[45,46]. Then, long-term linear and seasonal trends were removed from each time series

using the same methodology applied to the plankton data. Again, species time series with less

than 60% zero values were retained for analysis. Finally, each time series, including both species

abundance and size, was standardized to have a mean of zero and a unit variance.

Quantifying the bidirectional causal link. For each species (plankton, fish or terrestrial

species), we identified the bidirectional causal link among species size and its abundance (i.e.

size →abundance, abundance →size) using convergent cross-mapping (CCM) (Fig 1). CCM is

based on Takens's theorem for dynamic ecosystems, which infers the causal relationship among

variables from their empirical time series[50,51] (see three brief videos for a summary introduction:

http://tinyurl.com/EDM-intro). If the causal variable X (e.g. species abundance) and affected

variable Y (e.g. species size) belongs to the same dynamical system, information on the causal

variable X is encoded in the affected variable Y. Thus, one can predict the states of causal

variable X using the affected variable Y. CCM infers causality by measuring the extent to which

the causal variable X has left signatures in the time series of the affect variable Y; a procedure

known as cross mapping (cross-prediction)[50]. In this study, the appropriate embedding

dimensions for cross-mapping E were determined by the optimized the hindcast cross-mapping

in which X(t) projected one-step backward to Y(t-1), examining values of E from 2 to square

root of n, where n is the length of the time series[46]. Thus, we computed n as the geometric mean

time series length across all data sets. E was finally examined from 2 to 7 across all data sets.

We used simplex projection to select the best E that gave the highest prediction skill[52].

To accommodate the fact that the causal variables (e.g., species abundance) can exhibit

time-delayed effects on the affected variables (e.g., species size), we carried out 0 to 1 year (i.e.

0~4 time point for plankton and terrestrial species but 1 time point for fish) lagged CCM

analyses, in which we retained the CCM with the time lag resulting in the highest cross mapping

skill ρ[53]. The criterion for testing causality is the convergence toward higher cross-mapping skill

with increasing library length (i.e., the number of points used for state space reconstruction)[50].

An increase in library length enhances the density of points in the reconstructed attractor,

facilitating more precise identification of nearest neighboring points for predictions. This, in turn,

leads to improved predictive accuracy[54]. Convergence can be assessed by evaluating whether

there is a significant monotonic increasing trend in cross-mapping skill ρ as library length

increases using Kendall's τ test[48], and whether the ρ at the largest library length is significantly

greater than that the ρ at the smallest library length using Fisher's Z test[48]. In this study, library

length was set from the minimum (E) to the maximum length (i.e., the length of the entire time

series). Throughout this study, an interaction link (e.g., X→Y) was considered causal if both

Kendall's τ test and Fisher's Z test were statistically significant ($P < 0.05$) for testing

convergence[48].

Finally, the warming rate, representing the long-term warming trend of water temperature

or land surface temperature in each ecosystem, was measured using the Theil–Sen median-based

trend estimator, implemented via 'sens.slope' from 'trend' package[48,55]. This method was chosen

due to its widespread use in quantifying warming rates[56,57], its robustness to episodic extreme

events, and its ability to provide stable estimates of long-term trends in the presence of outliers[55].

Species mean body mass was computed as average across whole time series, estimated from

biovolume, assuming that organisms have the same density as water, if unavailable, was obtained

from published sources[58,59].

Bidirectional links in responses to climate change and biotic drivers. We employed a

Zero-inflated Generalized Linear Mixed Model (glmmTMB function in glmmTMB package[60])

with a binomial error distributions to test the main effects of warming, mean temperature, species

richness, and mean species body mass on proportion of either link (size →abundance or

abundance →size). To ensure normality, mean temperature, species richness, and mean species

body mass were log-transformed using base 10. In the model, the proportion of each link was

used as the response variable, consisting of binary values: 1 indicating the presence of a causal

link and 0 indicating the absence of a link. Last, we set species groups (e.g. plankton) nested

within sample locations (e.g. Trout Lake) as random intercepts. Details of statistics see Table S1.

Sensitivity analysis. Given that low-frequency rare species or dominant species may

alter bidirectional link between species abundance and size, we extended the analysis to: 1)

include a large number of rare species (number of zeros 60%~80%) into analysis; these were

species excluded in analysis in main text, and 2) only use the very dominant species into analysis

(number of zeros <50%, that is species presence once per two samplings). The extended analysis

(1)-(2) did not change the conclusions (Fig. S3-S8). All analyses were done with R (ver. 4.1.2).

The CCM analysis was examined using the rEDM package (version 1.2.3).

Competing interests

The authors declare no competing interests.

Author Contributions

Q.H.Z., F.D.L. and Y.X.G.W designed the research. Q.H.Z. and Y.X.G.W performed the study

and analyzed the data. Q.H.Z and G.S performed the CCM analysis. Q.H.Z., F.D.L. and Y.X.G.W

led the writing, with contributions from C.X. and G.S.

Data and code availability

The data reproducing all the results in this study are available on GitHub

(https://github.com/QZhao16/link.abundance-size).

Acknowledgements

We thank all participants in each long-term monitoring site. Q.Z. acknowledges funding from the

concerted research action (ARC) from the special research fund from the University of Namur,

Belgium (ARC grant DIVERCE, Convention 18/23-095).

Figures

Figure 1. **Proportion of bidirectional link among species' abundance and size**. **a**, An

overview of the methods. The bidirectional link among abundance and body size can be

modulated by climate change and biotic drivers. In this study, we investigate the link within

individual species rather than across species. **b**, The average proportion of causal links for

aquatic plankton, fish, or terrestrial species across 41 ecosystems. The frequency and proportion

of causal links abundance → size (**c**) and size → abundance (**d**) across 41 ecosystems. The

details of proportion of causal link within each of the 41 ecosystems see Fig. S1–S2.

Figure 2. **The bidirectional link between abundance and size in response to warming and**
biotic drivers. Proportion of casual link from size to abundance (size →abundance) and from
abundance to size (abundance →size) in responses to warming (**a, d**), species richness (**b, e**), and
mean individual body mass (**c, f**). **a-f,** Causal links identified through CCM are coded as 1 and
non-causal link as 0. The bold black lines and error bands depict the significant best-fit trendline
and the 95% confidence interval in the logistic regressions (two-sided) across 41 ecosystems,
respectively, while the dashed bold black lines indicate nonsignificant fit across 41 ecosystems.
The non-bold dashed colored lines indicate the best-fit trendline in the logistic regressions (two-
sided) within each group (plankton, fish or terrestrial species). Colored points represent the
values of causal links between abundance and size within each species across 41 ecosystems
(total points in each figure, *n* = 1269). For statistical results see table S1.

327
328

 Supplementary Figures

 Figure S1. **The bidirectional causal link between species size and abundance in each of**

 aquatic ecosystems. Red points indicate causal link identified by CCM, while blue points

 indicate non-causal link. The numbers in parentheses above each violin plot represent the

 number of causal links (blue) and the total links (black). The percentage above each violin plot

 indicates the proportion of causal links to total links. The abbreviations of the sites see Table S2-

 S3.

Figure S2. **The bidirectional causal link between species size and abundance in each of**
terrestrial ecosystems. Red points indicate causal link identified by CCM, while blue points
indicate non-causal link. The numbers in parentheses above each violin plot represent the
number of causal links (blue) and the total links (black). The percentage above each violin plot
indicates the proportion of causal links to total links. The abbreviations of the sites see Table S4.

Figure S3. The average proportion of causal links for aquatic plankton, fish, and terrestrial

species across 41 ecosystems, by including rare species (number of zeros 60%~80% in

species time series) into analysis. Within each system, the proportion of the bidirectional causal

link see Figure S4-S5.

Figure S4. **The bidirectional causal link between species size and abundance in each of**
aquatic ecosystems, by including rare species (number of zeros 60%~80% in each species
time series) into analysis. Red points indicate causal link identified by CCM, while blue points
indicate non-causal link. The numbers in parentheses above each violin plot represent the
number of causal links (blue) and the total links (black). The percentage above each violin plot
indicates the proportion of causal links to total links. The abbreviations of the sites see Table S2-
S3.

357
358

Figure S5. **The bidirectional causal link between species size and abundance in each of**

terrestrial ecosystems, by including rare species (number of zeros 60%~80% in each

species time series) into analysis. Red points indicate causal link identified by CCM, while blue

points indicate non-causal link. The numbers in parentheses above each violin plot represent the

number of causal links in CCM (blue) and the total links (black). The percentage above each

violin plot indicates the proportion of causal links to total links. The abbreviations of the sites see

Table S4.

Figure S6. The average proportion of causal links for aquatic plankton, fish, and terrestrial

species across 41 ecosystems, by only using very dominant species (number of zeros <50%

in each species time series) into analysis. Within each system, the proportion of the

bidirectional causal link see Figure S7-S8.

Figure S7. **The bidirectional causal link between species size and abundance in each of**
aquatic ecosystems, by including rare species (number of zeros <50% in each species time
series) into analysis. Red points indicate causal link identified by CCM, while blue points
indicate non-causal link. The numbers in parentheses above each violin plot represent the
number of causal link (blue) and the total links (black). The percentage above each violin plot
indicates the proportion of causal links to total links. The abbreviations of the sites see Table S2-
S3.

382
383

Figure S8. **The bidirectional causal link between species size and abundance in each of**
terrestrial ecosystems, by including rare species (number of zeros 60%~80% in each
species time series) into analysis. Red points indicate causal link identified by CCM, while blue
points indicate non-causal link. The numbers in parentheses above each violin plot represent the
number of causal links (blue) and the total links (black). The percentage above each violin plot
indicates the proportion of causal links to total links. The abbreviations of the sites see Table S4.

390
391

392 **Figure S9**. **Species-dependent casual link across 41 ecosystems**. The absence of bars

393 corresponding to a specific taxon (e.g., Atherinidae) indicates a proportional value of zero for the

394 associated link.

     Supplementary Tables

Table S1. Generalized Linear Mixed Model (glmmTMB function in glmmTMB package)

with a binomial error distribution was employed to test the main effects of warming,

mean temperature, species richness, and mean species body mass on proportion of

either link (size →abundance or abundance →size). In the model, the proportion of

each link was used as the response variable, consisting of binary values: 1 indicating

the presence of a causal link and 0 indicating the absence of a link. This model set

species groups (e.g. plankton) nested within sample locations (e.g. Trout Lake) as

random intercepts. Mean temperature, species richness, and mean species body mass

were log-transformed using base 10 to ensure normality. For the model testing the

link proportion of size → abundance, model's marginal and conditional R^2 are 0.15

and 0.22 respectively. For the model testing the link proportion of abundance → size,

the R^2 is 0.14 and 0.24, respectively. ***P< 0.001, **P<0.01, *P<0.05.

	proportion of size → abundance			proportion of abundance → size		
	coefficient	Z	P value	coefficient	Z	P value
(Intercept)	1.84	0.88	0.38	-1.42	-0.72	0.47
Species richness	-1.61	-2.13	**0.03***	1.18	1.43	0.15
Mean body mass	0.10	2.17	**0.02***	0.03	0.79	0.43
Mean Temperature	0.61	0.60	0.55	0.76	0.80	0.43
Warming	-22.93	-2.88	**0.004****	-47.08	-5.21	**<0.0001*****

411 **Table S2**. Plankton data for the 10 ecosystems were averaged seasonally (trimonthly) for species

412 abundance and size using raw data, with the resulting values used in the main analysis.

	System	Latitude	Longitude	Abbreviation	Duration	Region	Data Source
1	Wadden sea	55.03	8.46	WS	2001-2010	Europe	[61]
2	Lake Mendota	43.11	-89.42	ME	1995-2005	USA	[62,63]
3	Lake Monona	43.07	-89.35	MO	1995-2005	USA	[64,65]
4	Lake Allequash	46.04	-89.62	AL	2009-2018	USA	[64,65]
5	Lake Big Muskellunge	46.02	-89.60	BM	1991-1999	USA	[64,65]
6	Lake Crystal Bog	46.01	-89.61	CB	2008-2018	USA	[64,65]
7	Lake Crystal	45.99	-89.61	CR	1990-1999	USA	[64,65]
8	Lake Sparkling	46.00	-89.70	SP	1990-1999	USA	[64,65]
9	Trout Lake	46.07	-89.66	TR	1990-1999	USA	[64,65]
10	Lake Trout Bog	46.04	-89.69	TB	2009-2018	USA	[64,65]

415   **Table S3**. Fish data sources for the 11 ecosystems. For each system, species abundance and size

416   were averaged on an annual basis using the corresponding raw data. These average values were

417   then used in the main text.

	System	Latitude	Longitude	Abbreviation	Duration	Region	Data Source
1	American River	38.62	-121.30	AR	1976-2023	USA	66,67
2	San Joaquin River	37.28	-120.82	SJ	1981-2023	USA	66,67
3	Sacramento River	38.24	-121.55	SR	1976-2023	USA	66,67
4	Keller Beach	37.92	-122.38	KB	1980-2023	USA	66,67
5	Smokelumne River	38.12	-121.57	SM	1976-2023	USA	66,67
6	Old River	37.92	-121.55	OR	1993-2023	USA	66,67
7	Lake Allequash	46.04	-89.62	AL	1991-2018	USA	68,69
8	Lake Big Muskellunge	46.02	-89.60	BM	1991-2018	USA	68,69
9	Lake Mendota	43.11	-89.42	ME	1991-2018	USA	68,69
10	Lake Sparkling	46.00	-89.70	SP	1991-2008	USA	68,69
11	Trout Lake	46.07	-89.66	TR	1991-2018	USA	68,69

419 **Table S4**. Terrestrial data sources across 20 ecosystems. Within each system, species abundance
420 and size were averaged on a seasonal basis using the corresponding raw data. These seasonally
421 averaged values were subsequently used for analysis in the main text.

	System	Latitude	Longitude	Duration	Species	Region	Data Source
1	WA	40.17	-79.26	2005-2014	Beetle	US	BioTIME[70]
2	NI	32.62	-106.74	2004-2014	Beetle	US	BioTIME[70]
3	HO	42.67	141.63	2004-2014	Beetle	Japan	BioTIME[70]
4	WK	34.07	135.53	2005-2014	Beetle	Japan	BioTIME[70]
5	MI	31.86	131.3	2004-2014	Lizard	Japan	BioTIME[70]
6	FU	36.94	140.59	2004-2014	Beetle	Japan	BioTIME[70]
7	NA	36.84	138.5	2005-2014	Beetle	Japan	BioTIME[70]
8	NJ	38.21	138.44	2004-2014	Beetle	Japan	BioTIME[70]
9	SA	35.94	138.8	2004-2014	Beetle	Japan	BioTIME[70]
10	HJ	42.68	141.61	2004-2014	Beetle	Japan	BioTIME[70]
11	TO	43.43	144.06	2004-2014	Beetle	Japan	BioTIME[70]
12	TH	41.81	140.46	2004-2014	Beetle	Japan	BioTIME[70]
13	TJ	43.32	141.49	2004-2014	Beetle	Japan	BioTIME[70]
14	TK	42.18	143.09	2004-2014	Beetle	Japan	BioTIME[70]
15	TA	40.92	141.31	2004-2014	Beetle	Japan	BioTIME[70]
16	AK	28.33	129.45	2005-2014	Beetle	Japan	BioTIME[70]
17	AM	32.05	131.19	2004-2014	Beetle	Japan	BioTIME[70]
18	SM	38.25	140.85	2005-2014	Rat	Japan	BioTIME[70]
19	YA	36.7	138.5	2005-2014	Beetle	Japan	BioTIME[70]
20	SH	33.15	132.92	2005-2014	Beetle	Japan	BioTIME[70]

References

1.   White, E., Ernest, S. K. M., Kerkhoff, A. & Enquist, B. Relationships between body size and abundance in ecology. *Trends in ecology & evolution* **22**, 323–30 (2007).

2.   Enquist, B. J., Brown, J. H. & West, G. B. Allometric scaling of plant energetics and population density. *Nature* **395**, 163–165 (1998).

3.   Cohen, J. E., Jonsson, T. & Carpenter, S. R. Ecological community description using the food web, species abundance, and body size. *Proceedings of the National Academy of Sciences* **100**, 1781–1786 (2003).

4.   Allen, A. P., Brown, J. H. & Gillooly, J. F. Global Biodiversity, Biochemical Kinetics, and the Energetic-Equivalence Rule. *Science* **297**, 1545–1548 (2002).

5.   Gibert, J. P., Han, Z.-Y., Wieczynski, D. J., Votzke, S. & Yammine, A. Feedbacks between size and density determine rapid eco-phenotypic dynamics. *Functional Ecology* **36**, 1668–1680 (2022).

6.   Gillooly, J. F. Effect of body size and temperature on generation time in zooplankton. *Journal of Plankton Research* **22**, 241–251 (2000).

7.   Marañón, E. *et al*. Unimodal size scaling of phytoplankton growth and the size dependence of nutrient uptake and use. *Ecology Letters* **16**, 371–379 (2013).

8.   Arim, M., Abades, S. R., Laufer, G., Loureiro, M. & Marquet, P. A. Food web structure and body size: trophic position and resource acquisition. *Oikos* **119**, 147–153 (2010).

9.   Jennings, S., Pinnegar, J., Polunin, N. & Warr, K. Linking size-based and trophic analyses of benthic community structure. *Mar. Ecol. Prog. Ser.* **226**, 77–85 (2002).

10.   Dalponti, G., Guariento, R. & Caliman, A. Hunting high or low: body size drives trophic position among and within marine predators. *Mar. Ecol. Prog. Ser.* **597**, 39–46 (2018).

11.   Romanuk, T. N., Hayward, A. & Hutchings, J. A. Trophic level scales positively with body size in fishes. *Global Ecology and Biogeography* **20**, 231–240 (2011).

12.   Potapov, A. M., Brose, U., Scheu, S. & Tiunov, A. V. Trophic Position of Consumers and Size Structure of Food Webs across Aquatic and Terrestrial Ecosystems. *The American Naturalist* **194**, 823–839 (2019).

13.   Robinson, J. G. & Redford, K. H. Body Size, Diet, and Population Density of Neotropical Forest Mammals. *The American Naturalist* **128**, 665–680 (1986).

14.   Gibert, J. P., Wieczynski, D. J., Han, Z.-Y. & Yammine, A. Rapid eco-phenotypic
feedback and the temperature response of biomass dynamics. *Ecology and Evolution* **13**,
e9685 (2023).

15.   Govaert, L. & Klauschies, T. Eco-phenotypic feedback loops differ in multistressor
environments. *Ecology* **n/a**, e4480.

16.   Han, Z.-Y., Wieczynski, D. J., Yammine, A. & Gibert, J. P. Temperature and nutrients
drive eco-phenotypic dynamics in a microbial food web. *Proceedings of the Royal Society
B: Biological Sciences* **290**, 20222263 (2023).

17.   De Meester, L. *et al.* Analysing eco-evolutionary dynamics—The challenging
complexity of the real world. *Functional Ecology* **33**, 43–59 (2019).

18.   Savage, V. M., Gillooly, J. F., Brown, J. H. & Charnov, E. L. Effects of body size and
temperature on population growth. *Am Nat* **163**, 429–441 (2004).

19.   DeLong, J. P. *et al.* The body size dependence of trophic cascades. *Am Nat* **185**, 354–
366 (2015).

20.   Gavina, M. K. A. *et al.* Multi-species coexistence in Lotka-Volterra competitive
systems with crowding effects. *Sci Rep* **8**, 1198 (2018).

21.   Barabás, G., Michalska-Smith, M. J. & Allesina, S. Self-regulation and the stability of
large ecological networks. *Nat Ecol Evol* **1**, 1870–1875 (2017).

22.   He, R.-C., Wang, L., Goodale, E., Li, K. & Quan, R.-C. The extremely small body size
of Williamson's mouse deer (Tragulus williamsoni) allows coexistence with sympatric larger
ungulates through temporal avoidance. *Front. Ecol. Evol.* **11**, (2023).

23.   Wellborn, G. A. Trade-Off between Competitive Ability and Antipredator Adaptation
in a Freshwater Amphipod Species Complex. *Ecology* **83**, 129–136 (2002).

24.   Terry, J. C. D., O'Sullivan, J. D. & Rossberg, A. G. No pervasive relationship between
species size and local abundance trends. *Nat Ecol Evol* **6**, 140–144 (2022).

25.   Martins, I. S. *et al.* 0. Widespread shifts in body size within populations and
assemblages. *Science* **381**, 1067–1071 (2023).

26.   Genner, M. J. *et al.* Body size-dependent responses of a marine fish assemblage to
climate change and fishing over a century-long scale. *Global Change Biology* **16**, 517–527
(2010).

27.   Åkesson, A. *et al.* The importance of species interactions in eco-evolutionary
community dynamics under climate change. *Nat Commun* **12**, 4759 (2021).

28.   Thrall, P. H. & Blanc, S. Unravelling eco-evolutionary dynamics: Understanding
adaptation to global change. *Ecology Letters* **26**, S3–S4 (2023).

29.   Alberti, M. *et al*. The Complexity of Urban Eco-evolutionary Dynamics. *BioScience*
70, 772–793 (2020).

30.   Kraemer, B. M. *et al*. Climate change drives widespread shifts in lake thermal
habitat. *Nat. Clim. Chang.* **11**, 521–529 (2021).

31.   Tong, Y. *et al*. Global lakes are warming slower than surface air temperature due to
accelerated evaporation. *Nat Water* **1**, 929–940 (2023).

32.   Woolway, R. I. *et al*. Lake heatwaves under climate change. *Nature* **589**, 402–407
(2021).

33.   Merckx, T. *et al*. Body-size shifts in aquatic and terrestrial urban communities.
Nature **558**, 113–116 (2018).

34.   Govaert, L. & Klauschies, T. Eco-phenotypic feedback loops differ in multistressor
environments. *Ecology* **e4480**, e4480 (2024).

35.   Audzijonyte, A. *et al*. Fish body sizes change with temperature but not all species
shrink with warming. *Nat Ecol Evol* **4**, 809–814 (2020).

36.   Sebastian, P., Stibor, H., Berger, S. & Diehl, S. Effects of water temperature and
mixed layer depth on zooplankton body size. *Mar Biol* **159**, 2431–2440 (2012).

37.   Brandenburg, K. M., Velthuis, M. & Van de Waal, D. B. Meta-analysis reveals
enhanced growth of marine harmful algae from temperate regions with warming and
elevated CO_2 levels. *Glob Chang Biol* **25**, 2607–2618 (2019).

38.   Crozier, L. G., Zabel, R. W., Hockersmith, E. E. & Achord, S. Interacting effects of
density and temperature on body size in multiple populations of Chinook salmon. *Journal*
of Animal Ecology **79**, 342–349 (2010).

39.   Gillooly, J. F. Effect of body size and temperature on generation time in zooplankton.
Journal of Plankton Research **22**, 241–251 (2000).

40.   Warzecha, D., Diekötter, T., Wolters, V. & Jauker, F. Intraspecific body size increases
with habitat fragmentation in wild bee pollinators. *Landscape Ecol* **31**, 1449–1455 (2016).

41.   Zastavniouk, C., Weir, L. K. & Fraser, D. J. The evolutionary consequences of habitat
fragmentation: Body morphology and coloration differentiation among brook trout
populations of varying size. *Ecology and Evolution* **7**, 6850–6862 (2017).

42. Clements, C. F., McCarthy, M. A. & Blanchard, J. L. Early warning signals of recovery
in complex systems. *Nat Commun* **10**, 1681 (2019).

43. Clements, C. F., Blanchard, J. L., Nash, K. L., Hindell, M. A. & Ozgul, A. Body size
shifts and early warning signals precede the historic collapse of whale stocks. *Nat Ecol*
Evol **1**, 1–6 (2017).

44. Clements, C. F. & Ozgul, A. Including trait-based early warning signals helps predict
population collapse. *Nat Commun* **7**, 10984 (2016).

45. Zhao, Q. *et al.* Relationships of temperature and biodiversity with stability of natural
aquatic food webs. *Nat Commun* **14**, 3507 (2023).

46. Karakoç, C., Clark, A. T. & Chatzinotas, A. Diversity and coexistence are influenced
by time-dependent species interactions in a predator–prey system. *Ecology Letters* **23**,
983–993 (2020).

47. Chang, C.-W. *et al.* Causal networks of phytoplankton diversity and biomass are
modulated by environmental context. *Nat Commun* **13**, 1140 (2022).

48. Chang, C.-W. *et al.* Long-term warming destabilizes aquatic ecosystems through
weakening biodiversity-mediated causal networks. *Global Change Biology* **26**, 6413–6423
(2020).

49. Global Historical Climatology Network - Daily (GHCN-Daily), Version 3. NOAA
National Centers for Environmental Information https://doi.org/10.7289/V5D21VHZ (2012).

50. Sugihara, G. *et al.* Detecting causality in complex ecosystems. *Science* **338**, 496–
500 (2012).

51. Deyle, E. R. & Sugihara, G. Generalized Theorems for Nonlinear State Space
Reconstruction. *PLOS ONE* **6**, e18295 (2011).

52. Hsieh, C., Glaser, S. M., Lucas, A. J. & Sugihara, G. Distinguishing random
environmental fluctuations from ecological catastrophes for the North Pacific Ocean.
Nature **435**, 336–340 (2005).

53. Ye, L. *et al.* Functional diversity promotes phytoplankton resource use efficiency.
Journal of Ecology **107**, 2353–2363 (2019).

54. Sugihara, G. & May, R. M. Nonlinear forecasting as a way of distinguishing chaos
from measurement error in time series. *Nature* **344**, 734–741 (1990).

55. Mohsin, T. & Gough, W. A. Trend analysis of long-term temperature time series in the
Greater Toronto Area (GTA). *Theor Appl Climatol* **101**, 311–327 (2010).

56.   Pinsky, M. L. *et al*. Warming and cooling catalyse widespread temporal turnover in
biodiversity. *Nature* 1–5 (2025) doi:10.1038/s41586-024-08456-z.

57.   Tong, Y. *et al*. Global lakes are warming slower than surface air temperature due to
accelerated evaporation. *Nat Water* **1**, 929–940 (2023).

58.   Cohen, R. E. & Lough, R. G. Length-Weight Relationships for Several Copepods
Dominant in the Georges Bank-Gulf of Maine Area. *J. Northw. Atl. Fish. Sci.* **2**, 47–52 (1981).

59.   Huntley, M. E. Temperature-Dependent Production of Marine Copepods: A Global
Synthesis. *The American Naturalist* **140**, 201–242 (1992).

60.   Brooks, M. *et al*. glmmTMB: Generalized Linear Mixed Models using Template Model
Builder. 1.1.9 https://doi.org/10.32614/CRAN.package.glmmTMB (2017).

61.   Felden, J., Möller, L., Schindler, U., Huber, R., Schumacher, S., Koppe, R.,
Diepenbroek, M., Glöckner, FO (2023): PANGAEA – Data Publisher for Earth &
Environmental Science. Scientific Data, 10(1), 347, https://doi.org/10.1038/s41597-023-
02269-x. https://www.pangaea.de/.

62.   Magnuson, J., Carpenter, S. & Stanley, E. North Temperate Lakes LTER: Zooplankton
- Madison Lakes Area 1997 - current. Environmental Data Initiative
https://doi.org/10.6073/PASTA/A03B42039E6DB8C45A9BB9E480A2BA6A (2024).

63.   Magnuson, J. J., Carpenter, S. R. & Stanley, E. H. North Temperate Lakes LTER:
Phytoplankton - Madison Lakes Area 1995 - current. Environmental Data Initiative
https://doi.org/10.6073/PASTA/BD03330E83FC52A5442BB39E131C95E9 (2022).

64.   Magnuson, J. J., Carpenter, S. R. & Stanley, E. H. North Temperate Lakes LTER:
Zooplankton - Trout Lake Area 1982 - current. Environmental Data Initiative
https://doi.org/10.6073/PASTA/04030C61782378DD44ABB005B29EA5D3 (2023).

65.   Magnuson, J., Carpenter, S. & Stanley, E. North Temperate Lakes LTER:
Phytoplankton - Trout Lake Area 1984 - 2006. Environmental Data Initiative
https://doi.org/10.6073/PASTA/E876543EBE841BF7608F800A6CFB357E (2023).

66.   CDFW. Interagency Ecological Monitoring Program. 2023. https://
iep.ca.gov/Science-Synthesis-Service/Monitoring-Programs/ EMP.

67.   CDFW. California Department of Fish and Wildlife San Francisco Bay study: Long-
term fish and water quality monitoring data. 2023.
https://filelib.wildlife.ca.gov/Public/BayStudy/.

68.   Magnuson, J. J., Carpenter, S. R. & Stanley, E. H. North Temperate Lakes LTER: Fish

Abundance 1981 - current. Environmental Data Initiative

https://doi.org/10.6073/PASTA/A47E2F761A5F4F99F31998AEF065E2D1 (2024).

69.   Magnuson, J. J., Carpenter, S. R. & Stanley, E. H. North Temperate Lakes LTER: Fish

Lengths and Weights 1981 - current. Environmental Data Initiative

https://doi.org/10.6073/PASTA/E97E5046660A7144F4B6AADE41C2125D (2024).

70.   Dornelas, M. *et al.* BioTIME: A database of biodiversity time series for the

Anthropocene. *Global Ecology and Biogeography* **27**, 760–786 (2018).

588