Molecular evidence for pre-chordate origins of ovarian cell types and

neuroendocrine control of reproduction

Periklis Paganos[1], Carsten Wolff[1], Danila Voronov[2], S. Zachary Swartz[1*]

[1] Marine Biological Laboratory, 7 MBL Street, Woods Hole, Massachusetts, 02543,

United States of America

[2] Max Planck Institute for Evolutionary Biology, August-Thienemann-Straße 2, 24306

Plön, Germany

*correspondence: zswartz@mbl.edu

Abstract

Sexual reproduction in animals requires the development of oocytes, or egg cells. This

process, termed oogenesis, requires complex interactions amongst germline and somatic

cell types in the ovary. How did these cell types and their signaling interactions evolve?

Here we use the sea star *Patiria miniata* as a non-chordate deuterostome representative

to define the ovarian cell type toolkit in echinoderms. Sea stars continuously produce

millions of new oocytes throughout their lifespan, making them a practical system to

understand the mechanisms that drive oogenesis from a biomedical and evolutionary

perspective. We performed scRNA-seq combined with high-resolution 3D-imaging to

reveal the ovarian cell types and their spatial organization. Our data support the presence

of actively dividing oogonial stem cells and granulosa-like and theca-like cells, which

display similarities and possible homology with their mammalian counterparts. Lastly, our

data support the existence of an endocrine signaling system between oogonial stem cells

and intrinsic ovarian neurons with striking similarities to the vertebrate hypothalamic-

pituitary-gonadal axis. Overall, this study provides molecular evidence supporting the

possible pre-chordate origins of conserved ovarian cell types, and the presence of an

intrinsic neuroendocrine system which potentially controls oogenesis and predates the

formation of the hypothalamic-pituitary-gonadal axis in vertebrates.

Keywords: sea star, single cell transcriptomics, oogenesis, oogonial stem cells, HPG-axis.

Introduction

Oogenesis is essential for the propagation of most animal species. This process takes place within the ovary, where somatic and germ cells interact to form a fundamental reproductive unit: the follicle (1). Diverse ovarian cell types underlie these interactions and define the reproductive potential of the organism. For instance, somatic follicle cells, also referred to as granulosa in mammals, are in direct contact with developing oocytes and provide signals and metabolic support to regulate growth and meiotic arrest. Mammalian theca cells in the outer layers of the follicles have a key role in steroid hormone production, nutrient delivery to the follicle through the vasculature, and regulating granulosa cell function (2). In vertebrates, the hypothalamic-pituitary gland-gonadal (HPG) axis provides hormonal cues from the central nervous system essential for proper germ cell growth, maturation, and ovulation (3).

Many germ line processes are broadly conserved, but different organisms have diverse reproductive potentials due to the presence or absence of specific cell types or states. For instance, the mammalian oocyte reserve is established during embryogenesis, during which germ cells are limited to either entering meiosis, generating oocytes that remain arrested until puberty, or undergoing apoptosis (4). This limited number of oocytes shows signs of developmental decline with age in both mice and humans (5). Other animals such as the fruit fly instead produce oocytes throughout their lifespans due to a well-defined germ line stem cell niche that persists through adulthood (6). However, the mechanisms controlling oogenesis and the establishment of reproductive potential in other metazoans remain less understood. Moreover, numerous evo-devo studies have identified conserved genes expressed in germ line cells (7), but much less is known about the evolutionary history of the other somatic cell types within ovaries.

Echinoderms are non-chordate deuterostomes (8), a phylogenetic position informative for both vertebrates and other invertebrates, thus making them a valuable experimental

system for addressing questions about the deep evolution of reproductive cell types (**Fig.**
1A). Echinoderms including sea stars continuously produce millions of oocytes that
develop into optically transparent embryos and larvae, and have documented lifespans
that can exceed a century (9). However, while echinoderms have been extensively used
to study fertilization, gene regulation, and embryogenesis due to the abundance of eggs
they provide, we still lack a mechanistic understanding for how they achieve this
remarkable reproductive output (10,11).
70
In this work, we asked how the reproductive potential of the sea star *Patiria miniata* is
controlled, and whether sea star ovarian cell types are conserved with distantly related
animals. Using high-resolution microscopy and single cell RNA sequencing (scRNA-seq),
we defined the cellular organization of the sea star ovary, determined its cell types, and
explored their evolutionary relationships with other animals. We found the ovary to consist
of epithelial cells, immune cells, neurons, oocytes, three diversified groups of follicle cells,
and putative oogonial stem cells organized in germ cell cysts. Our cross-species
comparison revealed the presence of potentially conserved ovarian cell types, including
a granulosa-like population. Last, we report conserved signaling pathways involved in
oogenesis, as well as an intrinsic hypothalamic-like and gonadotropic-like signaling
system operating between oogonial stem cells and ovarian neurons which we propose
may be analogous, or possibly homologous, to the vertebrate HPG axis.
83
Materials and Methods
Animal husbandry, ovary isolation, *in vitro* culture
Adult *Patiria miniata* were obtained from South Coast Bio (San Diego, CA, USA) and
Monterey Abalone Company (Monterey, CA, USA). The animals were maintained in
temperature controlled (15°C) circulating seawater aquariums at the Marine Resources
Center (MBL, MA, USA). Ovarian fragments were surgically isolated through the oral side
of the sea star's arm using a Stoelting™ Sterile Disposable Scalpel, No.11 and ovaries
were collected using forceps in Eppendorf tubes containing filter sea water (FSW). For
the single cell RNA sequencing experiments the ovaries were used immediately. For the

rest of the experiments of the manuscript, ovaries were kept in FSW 10 µg/ml

trimethoprim and 50 µg/ml sulfamethoxazole at 15°C as previously described (12).

Ovary dissociation

Dissociation of isolated sea star intact ovarian lobes into single cells was performed as

previously described with minor modifications (13). Briefly, ovarian fragments were

transferred with forceps to a petri dish containing Ca^{2+} Mg^{2+}-free artificial sea water

(CaMg-free ASW) and allowed to settle for 3 minutes (min). Then the ovary was

transferred to an Eppendorf tube containing the dissociation buffer (1M glycine, 0.02M

EDTA, in CaMg-free ASW) and incubated for 10 min. Dissociation was promoted through

gentle pipette aspiration every 2 min. Once dissociation was complete, cells were spun

down with a swing bucket centrifuge at 500g for 10 min and washed twice with CaMg-

free ASW. Propidium Iodide and Fluorescein diacetate were used to assess cell viability

and only cell suspensions with cell viability ≥ 90 % were further processed. Single cells

were passed through a 40 µm cell strainer to remove aggregates. The number of cells

was estimated using a hemocytometer and diluted according to the manufacturer's

protocol (10X Genomics). During the entire process, specimens were kept on ice, except

for the centrifugations in which they were kept at 4°C. The isolated cells were loaded on

the 10x Genomics Chromium Controller according to the manufacturer's instructions and

cDNA libraries were generated using the Chromium Single Cell 3' Reagent Kit (v3.1

Chemistry Dual Index). Specimens were sequenced by the Azenta Life Sciences

(Burlington, MA) sequencing service using the Illumina NextSeq 550 with a maximum

read length 2 x 150 bp at a resolution of 350M PE reads.

Single cell RNA sequencing mapping of reads and data analysis.

Single cells from ovaries originating from three different animals were processed using

the 10X Genomics Chromium scRNA-seq capturing system. The libraries generated were

biased towards early oogenesis since the 10X Genomics Chromium capturing system

allows for the capture of cells up to 40 µm, which means that in our single cell atlas the

maximum oogonium size represented is 40 µm. Fully grown oocytes in *Patiria miniata* are

~170 µm. *Patiria miniata* genome version 3.0 (14) was used for single cell RNA

sequencing read mapping. The corresponding annotation was modified to contain only

protein coding entries using AGAT v0.7.0 (15) and Linux command line tools GNU

coreutils v8.26. Cell Ranger Software Suite v7.1.0 (16) was used to create the index from

the genome FASTA file and the annotation GTF file as well as for mapping with --no-bam

and --nosecondary flags, with all other settings left as default so the number of cells was

estimated automatically by Cell Ranger. Cell Ranger output files were loaded in R Studio

and processed with the Seurat pipeline (17). Seurat objects were created by excluding

cells containing genes that are transcribed in less than three cells and cells that have less

than a minimum of 400 and a maximum of 5000 transcribed genes. Next the Seurat

objects were normalized, and variable genes were found through the variance stabilizing

transfer (VST) method with a maximum of 2000 variable features. The three different

objects were integrated through identification of gene anchors (FindIntegrationAnchors).

Following integration, scaling and principal component analysis (PCA) were performed.

Jackstraw was used to evaluate PCA significance. Uniform Manifold Approximate and

Projection (UMAP) was used to perform clustering dimensionality reduction. The final

object consisted of 35,877 cells. The FindAllMarkers command was used to identify

differentially expressed marker genes. The average expression of genes of interest were

visualized using the DotPlot and DoHeatmap commands included in the Seurat R

package. To compare the *Patiria miniata* ovary integrated single cell atlas with the publicly

available human (18), mouse (19), zebrafish (20), sea urchin (21) and fruit fly (22) ovarian

atlases as well as with the mouse pituitary gland atlas (23) and the sea urchin larva one

(13) SAMap v1.02 (24) was used as previously described (25). For the mouse, zebrafish,

fruit fly and sea urchin (ovary and larva) datasets the already available rds files were

used, while for the human fetal ovary and the mouse pituitary gland datasets clustering

analysis was performed similar to what was previously described (18,23). The average

score of genes of interest was performed using the AddModuleScore function

incorporated in the Seurat R package. The oogenesis trajectory was performed using the

R package Monocle3 (26). The package EnhancedVolcano was used to visualize the

differentially expressed genes between the germ cell cysts and oocytes along the

trajectory, while the R packages topGO and clusterProfiler were used for the GO

enrichment analysis.

Light sheet microscopy/Segmentation 3D analysis

Ovarian samples were mounted individually in Fluidflon FEP tubes (Proliquid 1.6x2.4mm) using 1% low gelling agarose (Sigma-Aldrich A0701, type VII-A) with their elongated morphology parallel to the tube. A Zeiss Lightsheet 7 was used to acquire multiple-view data. It was equipped with a 10x/0.5 W Plan-Apochromat detection objective and two 10x/0.2 air illumination objectives producing two light-sheets 5.8 µm thick at the waist and 12µm thick at the edges of a 1100 µm x 1100 µm field of view. Illumination mode was dual (mean fused) with active pivot scan. Samples were imaged from five views (72 degrees apart) and Zoom factor 0.8. Each z-stack has a pixel frame of 1920x1920 and a voxel size of 0.57 µm x 0.57 µm x 1.8 µm. Each optical slice was acquired with a combination of 405, 488, and 561 nm laser with following filter settings: 405nm: EF1 - BP 420-470, CAM BS - SBS LP 490; 488 nm: EF1 - BP 505-545, CAM BS - SBS LP 560, 561 nm: EF1 - BP 575-615, CAM BS - SBS LP 640. For registration of multi-view acquisitions and fusion plus deconvolution of the registered data, the Fiji plugin BigStitcher v2.3.4 (27) was used with standard settings. Image segmentation, 3D-reconstructions and data analysis were performed using the Amira software package (FEI Visualization Sciences Group Thermo Fisher Scientific), version 2024.1. Morphological features such as muscles and nuclei were segmented semi-automatically using various tools in Amira, such as "Thresholding" and "Magic Wand". Oocytes and germ cell cysts were primarily segmented manually. Volumes of oocytes and germ cell cysts were measured using the "Material Statistic" tool. Visualization of all segmented reconstructions was achieved using tools like "Volren", "Surface Rendering" and "Surface View" within the software package.

Immunohistochemistry (IHC) and fluorescent stainings

Ovarian fragments were fixed with 4% PFA in FSW for 15-20 min at room temperature (RT). Specimens were washed twice with FSW and five times with PBS supplemented with 0.1 % Tween (1x PBST). Next, samples were incubated in blocking solution [1 mg/ml Bovine Serum Albumin (BSA) and 4 % sheep serum (SS)] in PBST for one hour at RT. Blocking solution was removed and primary antibodies were added in the appropriate dilution in blocking solution and incubated overnight at 4°C. 1E11 (gift from Dr. Ina

Arnone) was used in a 1:20 dilution, anti-acetylated tubulin (Sigma) 1:100, anti-DM1α
(Developmental Studies Hybridoma bank) 1:100, anti-alpha tubulin (Sigma) 1:100, anti-
MHC (PRIMM) 1:50, anti-serotonin (Sigma) 1:500, anti-TH (Sigma) 1:100m and anti-
Myosin II (Developmental Studies Hybridoma bank) (1:50). Samples were washed five
times with 1x PBST (15 min each) and incubated for one hour at RT with secondary
antibodies (AlexaFluor) diluted 1:1000 in PBST. Samples were washed five times with 1x
PBST. In cases where phalloidin (Invitrogen) was used, it was diluted 1:20 (from a 40x
stock solution) in 1x PBST and allowed to incubate overnight at 4°C. For WGA staining
(Invitrogen) a working solution of 5 µg/mL was prepared, and the specimens were
incubated overnight at 4°C. Ovaries were washed five times with 1x PBST, nuclei were
stained with DAPI (1 µg/mL), and specimens were mounted for imaging. During the entire
procedure the interval of each wash was 15 min.
198
Whole mount Hybridization Chain Reaction (HCR)
HCR was performed according to previous protocols (28) with minor modifications.
Probes were designed either by Molecular Instruments or HCR 3.0 Probe Maker (29).
Ovarian fragments were fixed in 4% PFA in Fixative Buffer (0.1M MOPS, 0.5M NaCl,
2mM EGTA, 1mM $MgCl_2$ and 1x PBS) for 1 hour at room temperature (RT). Specimens
were washed three times with Fixative Buffer (without PFA) at RT. Samples were washed
twice with 100% methanol, transferred into fresh 100% methanol, and placed at -20°C for
long term storage. Specimens were rehydrated in 75%, 50%, 25% methanol in RNAase
free water on ice (each for 10 min). Next, ovaries were washed four times for 5 min each
in RNase free 1x PBST. Ovaries were treated with permeabilization buffer (1.0% SDS,
0.5% Tween, 50 mM Tris-HCl (pH 7.5), 1.0mM EDTA (pH 8.0), 150 mM NaCl) and
incubated for 1h at RT. Next, specimens were washed four times for 5 min each in 1x
PBST. Pre-heated (37°C) Probe Hybridization Buffer (PHB) was added to the samples
and incubated for 3 hours at 37°C. PHB was exchanged with PHB containing in total 0.04
µM of each probe and specimens were left to incubate for one day at 37°C. Following
hybridization, samples were washed 4 times with pre-heated Probe Wash Buffer (PWB)
for 5 min each, followed by two washes with PWB for 30 min each at 37°C. Samples were
then washed two times with 5x SSCT (5x Saline Sodium Citrate and 0.1% Tween) and

two times with 1x SSCT (1x Saline Sodium Citrate and 0.1% Tween) each for 5 min. Next,

specimens were incubated with Amplification Buffer for 30 min at RT, hairpins were added

to the specimens according to manufacturer's guidelines and incubated overnight at RT

in the dark. Lastly, samples were washed twice with 1x SSCT for 5 min, two more times

for 30 min and four times with 1x PBST. Nuclei were stained with DAPI (1 µg/mL), and

specimens were mounted for imaging.

Fluorescent *in situ* hybridization (FISH)

FISH and antisense RNA probes design was performed as previously described (30,31).

In brief, ovaries were fixed in 4% PFA in MOPS Buffer for 1 hour at RT. Samples were

washed five times with MOPS buffer, gradually dehydrated (20%, 50% ethanol) and

stored in 70% ethanol and kept at -20°C for long term storage. Specimens were gradually

rehydrated (50%, 20% ethanol) and washed five times with MOPS Buffer. At this step

ovaries were dissected to smaller pieces using micro scissors. Specimens were then pre-

hybridized for 3 hours at 65°C, followed by a 48 hour period of probe hybridization. Next,

specimens were washed five times with MOPS Buffer and incubated in blocking solution

(Akoya Biosciences) for 30 min at RT. Next, specimens were transferred to a solution

containing the anti-Digoxigenin (Roche) antibody diluted 1:1000 in blocking solution. The

antibody solution was removed, samples were washed with MOPS Buffer five times and

the signal was developed using fluorophore conjugated tyramide technology (TOCRIS).

Nuclei were stained with DAPI (1 µg/mL), and specimens were mounted for imaging.

EdU labeling

Click-It EdU Cell Proliferation Kit for Imaging Alexa Fluor 488 (Thermo Fisher Scientific)

was used and was combined with either HCR or IHC as previously described (13,32).

Isolated intact ovarian fragments were treated with EdU at a final concentration of 20 µM

in FSW with 10 µg/ml trimethoprim and 50 µg/ml sulfamethoxazole and allowed to

incubate for 2 days at 15°C. Depending on whether the EdU cell proliferation was

combined with HCR or IHC specimens were fixed accordingly as described in their

respective sections above. To develop the EdU signal, the Click-iT reaction mix was

prepared according to the manufacturer's guidelines. In cases where the EdU assay was

combined with HCR after the washes with PWB, specimens were washed twice with 3%

BSA in PBST for 15 min at RT. Next the Click-iT reaction mix was added and incubated

for 30 min in the dark at RT. After this step the HCR procedure was carried out as

described above. For combination of the EdU Cell Proliferation assay with

immunostaining the ovaries were fixed in 4 % PFA in FSW for 15 min at RT and washed

five times with PBST. PBST was removed and the reaction mix was added to the samples

for 30 min (RT). Specimens were washed five times with PBST, mounted for imaging.

Results

The structural organization of the *Patiria miniata* adult ovary

We first sought to define the structure of its ovary, which is multi-lobed and optically

translucent (**Fig. 1B**). We stained for F-actin with phalloidin as a cell boundary marker

and acetylated tubulin to visualize cilia (**Fig. 1C**). Each ovarian lobe is a hollow tube that

consists of a thin, tightly packed epithelial cell layer with long motile cilia (**Fig. 1B**). We

also identified cells organized in plexus-like structures resembling a nerve net underneath

the epithelial layer that are positive for synaptotagmin 1, a pan-neuronal marker in

echinoderms (**Fig. 1D, S1A, E-H**). We also identified ovarian neurons in the related sea

urchin *Strongylocentrotus purpuratus* (**Fig. S1I-J**) and the sea cucumber *Holothuria*

tubulosa (**Fig. S1K-L**), arguing that these are a conserved cell type across echinoderms.

We also identified extensive musculature below and in close proximity to the nerve net

(**Fig. 1E, S1B**), which is likely responsible for the contraction of the ovary during spawning

(33). We observed axonal projections in close proximity to the muscle layer, which may

regulate their contraction (**Fig. S1E-F, I-J**). Below the muscle layer and within the lumen,

we detected oocytes at different stages of maturation enwrapped by follicle cells, which

are filamentous in nature and ciliated (**Fig. 1F, Fig. S1C-D**).

We determined the stages of oocyte development, and the nature of oocyte-follicle

complexes. Early small oogonia are spatially organized in closely associated groups with

an average size of 8-16 μm (**Fig. 1G, Fig. S2A-B**). These groups were found in all ovaries

tested, although their number varied with different individuals. The chromatin morphology

of these cells was distinct, with condensed DNA concentrated towards the nuclear

periphery. This may be consistent with the leptotene to zygotene stage of meiosis, in
which chromosomes adopt a bouquet configuration (34). The cells within these groups
were compact with intense actin staining at the cell boundaries, and surrounded by small
somatic nuclei that do not intercalate within the cluster (**Fig. 1G**). This organization
resembles germ cell nests or cysts that are found in other species, including fruit flies and
mammalian fetal ovaries (35). Similar structures were observed by electron microscopy
in the ovaries of the phylogenetically distant sea star *Asterias rubens* (36). Co-staining
for F-actin and membrane lectins revealed cortical channel-like structures between
distinct oogonia within the germ cell groups (**Fig. 1H-J**). Such an organization strongly
resembles the intercellular bridges that physiologically connect the germ cells within the
germ cell cysts in mammals and protostomes, allowing for the exchange of the germ
plasm and suggests that germ cells are organized in cysts also in a non-chordate
deuterostome. Therefore, from here on we refer to these groups as "cysts".

To test the germline nature of these cysts, we performed HCR for the conserved germ
cell factor *nanos1,* and the yolk protein *vtg2*, which is expressed in sea star follicle cells
in other echinoderms (37–39). We found that *nanos1* is expressed specifically in oocytes
of all sizes as well as in the cysts (**Fig. 1K**). *vtg2* was expressed by the follicle cells that
surround the oocytes. We also found that the *vtg2+* follicle cells enwrap the cysts as a
whole, while none of the cells within the cysts are individually enwrapped (**Fig. 1K**). This
configuration further resembles the germ cell cysts observed in vertebrates, suggesting
that sea star adult oogenesis may share conserved features. Co-staining for *nanos1* and
vtg2 also enabled us to ask what is the smallest oocyte size necessary to form a primary
follicle. We found that the smallest *nanos1+* oocytes are located next to germ cell cysts
and by using the diameter of these oocytes as a proxy to calculate the size and volume
we found they are within the range of 17-42 µm (**Fig. S2C-D**).

We next asked how oogenesis is spatially organized within the ovarian lobe. Is there a
defined stem cell niche, or is oogenesis instead uniformly distributed? To test this, we
used large volume light sheet microscopy on entire ovarian lobes (**Fig. 1L-M, Fig. S2G**).
Segmentation and 3D reconstruction revealed that oogenesis is spatially polarized from

outside to inside. Germ cell cysts and smaller oocytes were localized entirely to the

periphery of the lobe interior, but were broadly distributed. Fully grown oocytes in

prophase I arrest, instead, were centrally located within the lumen (**Fig. 1L-M, Fig. S2G**).

Quantification of segmented cell volumes found that the diameter of the entire germ cell

cysts ranges from 27-56 μm, of the small oocytes from 17-106 μm and of the pre-GVBD

ones from 107-187 μm (**Fig. S2E-F**). Altogether, these results define the cell type

organization of the sea star ovary at a molecular level and the oocyte size distribution

during oogenesis (**Fig. 1N, Fig. S2A-F**). Moreover, our data provide molecular evidence

for the presence of germ cell cysts and show how *P. miniata* oogenesis is spatially

oriented within the ovary.

Figure 1. Molecular characterization of the *P. miniata* ovary. (**A**) Phylogenetic tree showing the evolutionary relationships of *P. miniata* with other metazoan experimental systems. (**B**) *P. miniata* ovarian explant. Scale bar is 1 mm. (**C**) Phalloidin and acetylated tubulin staining to label cell boundaries and cilia. Scale bar is 20 μm. (**D**) IHC for α-tubulin labeling the ovarian nerve plexus. Scale bar is 40 μm. (**E**) Phalloidin staining showing ovarian muscle fibers. Scale bar is 40 μm.(**F**) Acetylated tubulin labeling cilia on the follicle cells. Dotted circle indicates oocytes, while germ cell cysts are outlined with dotted lines. Scale bar is 20 μm. (**G**) Phalloidin staining labeling the cell boundaries of the germ cell cysts. Germ cell cysts are outlined with dotted lines. Scale bar is 20 μm. (**H-J**) Phalloidin and WGA lectin co-staining labeling the cell boundaries and membranes. Yellow arrowheads indicate intercellular bridge-like structures. Scale bar is 1 μm. (**K**) HCR for *nanos1* (magenta) and *vtg2* (yellow) labeling the entire germ lineage and follicle cells respectively. Germ cell cysts are outlined with dotted lines. Scale bar is 20 μm. Nuclei were labelled with DAPI (cyan). (**L-M**) 3D rendering of segmented germ cell cysts and oocytes. Scale bar is 100 μm. Ovary seen from the lateral (I) and top (J) views. (**N**) Schematic summary of the germ cell cyst, young oocyte and mature oocyte size distributions in the *P. miniata* ovary. fc, follicle cells; l, lobe; o, oocytes.

The *P. miniata* ovary consists of nine molecularly distinct cell type groups

Having defined the overall structures of the ovary, we next sought to determine the molecular identities of its constituent cell types with scRNA-seq (**Fig. 2A, Fig. S3A**). We used ovaries from three independent *Patiria miniata* females (**Fig. S3B-C**). One ovary originated from an individual in which spawning was induced 2 months prior to ovarian isolation, one was highly enriched in germ cell cysts and early oocytes, and one consisted mostly of fully grown oocytes. For each ovary used, the size distribution of their oogonia/oocytes was determined by dissociating the ovaries and manually measuring the oogonia/oocytes diameter to estimate their volume (**Fig. S3B-C**). Computational analysis and integration of the sequenced libraries yielded an atlas of 35,877 cells distributed in twenty-seven clusters or cell states (**Fig. 2A, Fig. S3A**).

We next inferred the cell type identities of the twenty-seven clusters using the average expression of differentially expressed marker genes**,** and gene homologs known to label ovarian cell types, in echinoderms and other organisms. (**Fig. 2B)** We curated 9 cell type groups expressing genes corresponding to endothelial and ciliated cells (*btub2/3, cfap77, dhc2, col1a, vwf*) (13,40,41), mammalian granulosa cells (*tgfb2, foxl2, inhbb, cyp26, foxo, gata4*) (1,42–45), echinoderm follicle cells (*vtg2, vtg5*) (39), mammalian theca cells (*mmp14, angpt1, star, cyp17a1, vegfa, fgf18*) (46–50), muscles (*mhc, mlck, tnn1*) (13),

immunocytes (*lyz, mpeg1, bpi, cd151*) (51–54), neurons (*syt1, smp, pplnp, oxy/vp, hdc*)

(13,55,56) and germ cells/oocytes (*nanos1, nanos3, boll, piwi1, msl3, sycp1, figla, ddx4,*

sox2, pum2) (37,38,57–64). We found that most cell states were equally distributed

amongst the three ovary samples (**Fig. S3D**) with the exception of clusters 5, 9, 10, 11,

12 and cluster 4, identified as follicle cells and epithelium, respectively, which

predominantly originated from the previously spawned ovaries of female 1. Germ cell cyst

clusters 13 and 14 were also enriched in sample 2, which was used for the 3D volume

analysis (**Fig. 1I-K**) and contained numerous germ cell cysts. Therefore, our single cell

atlas reflects the experimentally validated reproductive states of the ovaries used. With

this confidence in our dataset, we next sought to spatially map their location within intact

ovaries using HCR and FISH (**Fig. 2C-H. Fig. S3E-I**).

We first examined genes expressed in follicle cells, which enwrap and support oocytes

as they develop. Using HCR, we detected *vtg2* (a previously validated marker) and the

signaling molecule *tgfb2* in follicle cells, with the mammalian theca marker *tgfb2* (65)

being expressed in a subset of them (**Fig. 2C-D**). We also detected *foxl2*, a known

mammalian granulosa cell transcription factor, in follicle cells (**Fig. 2E**). Therefore, the

clusters that contain *foxl2* (clusters 5-8, 23-24), *tgfb2* (clusters 5, 7-8, 22, 24-25) and *vtg2*

(clusters 5-8, 9-12, 23) transcripts were confirmed to correspond to the partially non

overlapping follicle cell groups as identified by scRNA-seq.

We next turned to the putative germ line clusters. The transcription factor *figla*, which

functions in mammalian oogenesis, was detected exclusively in the germ cell lineage

including cysts and oocytes (**Fig. 2F**). Similar results were obtained for the germline

factors *piwi1* and *nanos3* (**Fig. 2G, Fig. S3D-D'**). Surprisingly, we did not detect the

conserved germ line RNA helicase *ddx4/vasa* in the germ cell cysts, though it was

detected in all other oocyte stages (**Fig. 2H**). This result suggests that *ddx4/vasa*

transcription is activated only once the primary follicle is formed. These expression

patterns are consistent with the single-cell predictions and suggest the presence of two

cell groups: the germ cell cysts (clusters 13 and 14) and early oocytes (clusters 15, 16,

17, 18 and 19). To further validate our clustering analysis, we examined *syt1* expression,

which was unexpectedly predicted to be present not only in the nervous system but also

in germ cell cysts and oocytes. HCR confirmed *syt1* in the nerve net as well as in the

germ cell cysts and oocytes (**Fig. S3F-G'**), while a positive control in *P. miniata* larvae

and post-metamorphic juveniles verified the probe specificity, showing *syt1* expression in

the nervous system as reported in other echinoderms (**Fig. S3H, I**). Taken together, our

spatial and transcriptomic analyses enabled us to identify and map nine cell type groups

distributed in twenty-seven clusters (**Fig. 2I**).

Figure 2. Cell type/state diversity of the *P. miniata* adult ovary. (A) Integrated UMAP of the three
different scRNA-seq libraries. Color code is indicative of different cell type groups. **(B)** Dotplot showing the
average expression of known echinoderm cell type markers as well as genes labeling ovarian and
associated cell types in other organisms. Color code is the same as in panel A. **(C-D)** HCRs for the follicle
cells markers *vtg2* (C) and *tgfb2* (D)*.* **(E-F)** FISH using antisense probes for the transcription factors *foxl2*
(E) and *figla* (F) labelling the follicle cells and germ lineage respectively. Dotted circle in panel E is indicative
of oocytes, while germ cell cysts in panel F are outlined with dotted lines. Scale bars for panels C and E
are 20 µm, and for panel D 60 µm. **(G-H)** HCRs for the germ line markers *piwi1* and *ddx4*. Germ cell cysts
in panels G and H are outlined with dotted lines. Scale bars are 20 µm. **(I)** Schematic representation of the
ovary depicting the cell type groups revealed by our single cell RNA sequencing atlas. Color code is the
same as in panel A. fc, follicle cells; o, oocytes.

Shared cell type toolkits between *P. miniata* and other deuterostomes

How similar are sea star ovarian cell types to those of animals that diverged millions of

years ago? Our gene candidate-based analysis identified similarities—for instance, sea

star follicle cells (clusters follicle I and follicle III) express gene orthologs also found in

mammalian granulosa (*foxl2, inhbb, tgfb2, gata4*) and theca (*tgfb2, mmp14, angpt1r, star,*

cyp17a1, vegfa, fgf18) cells. To more broadly test this similarity, we used SAMap to

compare our atlas with published datasets of adult fruit fly, sea urchin, zebrafish, mouse,

and fetal human ovaries (**Fig. 3, S4**). As expected, sea star cluster 22 (muscle-related)

aligned with muscle clusters in all animal ovaries tested. This similarity is not unique to

ovarian muscles, as alignment with a sea urchin larval scRNA-seq dataset also revealed

strong similarity with esophageal muscles (**Fig. 3, S4**). We also identified similarity

between the sea star follicle cell I (cluster 6) and their follicular counterparts in fruit fly and

sea urchin (**Fig. 3A, S4A**). Surprisingly, despite substantial evolutionary distance, this

follicle cell cluster also aligned with follicle cells in zebrafish, and granulosa cells in both

mouse and human, although in the latter with lower alignment scores (**Fig. 3B,C, S4B**).

In addition, the sea star oocyte cluster 19 aligns with germ and oogonial cell clusters in

fruit fly, sea urchin, zebrafish and human (**Fig. 3A-C, S4A**). Zebrafish and mouse theca

cells aligned with sea star epithelial clusters, while zebrafish vasculature aligned to one

sea star follicle cluster (**Fig. 3B**). Similarities become scarcer when we compare the

ovaries of sea star with the adult mouse, which does not contain oogonial stem cells (66)

(**Fig. S4B**). However, when we compared the sea star ovary with the human embryonic

ovary, in which oogenesis is still occurring, we found that one human germ cell cluster aligned with the sea star oocytes cluster 17 (**Fig. 3C**). We also detected similarities between sea star follicle cells (I) with human endothelium, sea star follicle cells (III) with human granulosa cells, as well as sea star epithelium with human stromal and theca cells (**Fig. 3C**). A summary of the main findings can be found in (**Fig. 3D**). In support of the specificity of the similarities we detected, comparison to the sea urchin larva, which serves as a null comparison, yielded no strong hits apart from muscle (**Fig. S4C**). Overall, our analysis has shown the existence of transcriptionally similar ovarian cell type programs across phylogenetically distant organisms, suggesting the possible pre-chordate origins of ovarian cell types, including follicle cells.

 Figure 3. Cross-species comparison of ovarian single cell atlases with SAMap. Sankey plot depicting the alignment scores between the adult sea star ovarian cell clusters with the adult sea urchin **(A)**, adult zebrafish **(B)** and fetal human ovarian clusters **(C)**. **(D)** Summary of the cell types shared between sea star and the taxa used for comparison. Alignment score is defined as the average number of mutual nearest cross-species neighbors of each cell relative to the maximum possible number of neighbors. Alignment scores equal or greater to 0.1 were considered significant.

Identification and characterization of mitotic oogonial stem cells

Having defined similar ovarian elements across species, we then asked what unique features of the sea star ovary support its extreme reproductive capability, such as adult oogonial stem cells (OSCs). To identify potential OSC populations, we computationally scored for the co-expression of cell cycle-associated and meiotic genes with germline marker genes (**Fig. 4A, S5A**). We found that the epidermal cell clusters 2 and 3, the germ cell cysts clusters 13-14 and oocytes (15-19) express genes associated with G1/S phase (including *cyclinD2, cyclinE, cdt1* and *pcna*), while the epidermal cluster 4, the follicle cell clusters 9-12, the germ cell cyst cluster 14 and oocytes (15-19) express genes operating during G2/M phase (including *cyclinA, cyclinB* and *bub3*) (**Fig. 4A**). Genes specific to meiosis, such as *spo11, mlh1, hormad, sycp1, sycp2* and *sycp3* were expressed in the germ cell cyst clusters 13-14 and oocyte clusters 15-19 (**Fig. 4A**). HCR for one of those genes, *mlh1,* confirmed the predicted expression in both the germ cell cysts and the oocytes (**Fig. 5SB**).

To directly test cell proliferation, we cultured ovarian lobes *in vitro* and performed EdU cell proliferation assays (**Fig. 5SC-E**). We found cells undergoing DNA replication in both somatic and germ cell populations. This included the ovarian epithelium, follicle cells, and within germ cell cysts (**Fig. 5SC-E**). To further define the germ line population, we paired EdU incorporation with HCR for *nanos1, piwi1* and *ddx4/vasa* (**Fig. 4B-E**). The EdU signal was exclusively detected in *nanos1* and *piwi1* positive germ cell cyst nuclei, but not in the *ddx4* positive growing oocytes. Altogether, these data revealed the presence of replicating cells within the sea star ovary (**Fig. 4F**) and support our hypothesis of a self-renewing oogonial stem cell pool in adult sea star ovaries.

Figure 4. Identification of self-renewing oogonial stem cells. (A) Dotplot showing the average score of

genes involved in G1/S, G2/M phases of the cell cycle and meiosis across the *P. miniata* ovarian cell type

groups. (**B-E**) HCR for the germline markers *piwi1* and *ddx4* paired with the EdU proliferation assay. Scale

bars are 20 μm. Nuclei were labelled with DAPI (cyan). Dotted lines outline the germ cell cyst in panels B-

D. **(F)** Summary of the ovarian cell types undergoing mitosis as revealed by the EdU nuclear incorporation.

o, oocytes.

We next asked how gene expression changes as oocytes develop. We used Monocle3

to identify a developmental trajectory linking the 7 germ line cell clusters (13, 14, 15, 16,

17, 18), from OSC to oocyte. Cluster 13 was chosen as the root because its molecular

signature and cellular morphology was most similar to typical germ cells, and because it

did not express *ddx4*, as experimentally validated by HCR (**Fig. 5C-E**). We found a

substantial number of genes that are upregulated in germ cell cysts and downregulated

in oocytes, and fewer that are more highly expressed in oocytes. (3378 genes

upregulated in cysts, 849 upregulated in small oocytes; **Fig. S6A**). Gene ontology (GO)

enrichment analysis shows that most of the cyst genes correspond to transcription and

translation factors, while many of the oocyte genes are related to splicing, protein

synthesis and transport. (**Fig. S6B-B', S7**). Among the genes also upregulated in the

germ cell cyst clusters are the transcriptional and epigenetic regulators *atrx1* and *cxxc1*.

Both genes are important for mammalian sexual differentiation and oogenesis

respectively (67–69). Plotting their average expression on the trajectory as well as HCR

for both genes confirmed their co-expression in the oocyte lineage and presence within

the germ cell cysts (**Fig. 5F-K, Fig. S6C-C'''**). Taken together our data highlight, for the

first time in echinoderms, distinct transcriptional differences occurring during the transition

from oogonia to early oocytes (**Fig. 5L**).

Figure 5. Reconstruction of the early oogenesis developmental trajectory. (A-B) HCR for the germline markers *piwi1* and *ddx4* highlights the absence of *ddx4* in the germ cell cysts. Dotted lines outline the germ cell cyst. Nuclei were labelled with DAPI (cyan). **(C-D)** Feature plot showing the expression levels of *piwi1* and *ddx4* along the germline clusters of the *P. miniata* single cell atlas. **(E)** UMAP depicting the developmental trajectory of the germline clusters reconstructed with Monocle3. The germ cell cluster 13 has been used as a root. **(F-I)** HCR for the transcriptional regulators *atrx1* and *cxxc1* paired with *piwi1* confirms their expression in germ cell cysts. Scale bars are 5 µm. **(J-K)** Feature plot showing the distribution

502 of *atrx1* and *cxxc1* transcripts along the trajectory confirms their differential expression in clusters
503 corresponding to germ cell cysts. Nuclei were labelled with DAPI (cyan). **(L)** Cartoon highlighting the
504 difference in gene expression of the oogonia in the germ cell cysts versus the oocyte stages. o, oocytes.
505

Signaling pathways across ovarian cell types
What are the major signaling pathways that could control oogenesis and folliculogenesis?
We first mined our atlas for the expression of gene groups corresponding to either ligands
or downstream signaling cascades of the Wnt, Tgfb, Delta/Notch, Hh, and RA signaling
pathways (**Fig. 6A, S8A**).We found that the clusters that most strongly express Wnt ligands
correspond to epithelium, oocytes and immune cells, while follicle I, germ, oocytes,
neurons and muscles are able to respond to the signal based on the expression of frizzled
receptors, *dvl3, gsk3, apc, tcf/Lef* and *β-catenin* (**Fig. 6A, S8A**). Similarly, Tgfβ ligands are
expressed by the epithelium, follicle I, follicle III and muscle clusters, while their receptors
and signaling cascade factors are expressed in germ, oocytes, muscles, follicle III and
immune cells (**Fig. 6A, S8A**). For Delta/Notch signaling, *delta* transcripts are found in
epithelial, follicle II, follicle III and oocytes cell type groups, and the signal is predicted to
be received by follicle I, oocytes and neurons *notch* receptors and gamma secretase
subunits (**Fig. 6A, S8A**). Hedgehog (*hh*) is exclusively expressed by neurons, and the
signal may be received by follicle I and germ cell type groups based on the expression of
patched and smoothened receptors (**Fig. 6A, S8A**). Lastly, retinoic acid synthesis
machinery gene orthologs, including *sdr7, aldh1, aldh3* and *aldh8* are differentially
expressed in the follicle II, germ, and oocyte cell groups, and the cells that express RA
receptors *rar, rxr* and *ppar* belong to the follicle I, III, germ, neural and muscle ones (**Fig.**
6A, S8A). Degradation of RA is performed by cells expressing *cyp26,* which includes the
follicle I group, while its transport is mediated by cells that express *fabpl3,* which includes
the follicle I and follicle II groups (**Fig. 6A, S8A**). Our results indicate the presence of a
complex signaling system involved in sea star oogenesis by which diverse signals are
produced and received both by the follicle cells, the cells of the germ lineage, and the
ovarian neurons.
531

We next focused on Tgfβ signaling, which has been implicated in primordial germ cell
development and follicular growth in mammals (1,70,71) (**Fig. 6C-F, S8B-D**). Our atlas

indicates that Tgfβ signaling may be active within the oocyte-follicle complex. Of the
genes tested, only *tgfb2* ligand was uniquely expressed in somatic tissues and primarily
in follicle cells enwrapping the oogonia and mature oocytes (**Fig. 6C**). We found Tgfβ
receptors *tgfbr3*, *alk2* and the intercellular messenger *smad5* expressed by both the germ
cell cysts and developing oocytes, validating the scRNA-seq predictions (**Fig. 6D-F**). In
addition, our atlas showed that several Bmp ligands are expressed within the germ
lineage, suggesting that Tgfβ signaling may also be active between growing oocytes, or
acting upon follicle cells, a dual role also reported in mammals (**Fig. 6A, S8A**). HCR for
bmp1 and *bmp4* confirmed these predicted expression patterns (**Fig. S8B-D**). In summary,
our results support the deep evolutionary conservation of Tgfβ signaling pathway
components within the ovarian follicle.

Figure 6. Putative signaling pathways involved in ovarian cell type crosstalk. (A) Dotplot showing the
average score of gene groups corresponding to ligands and signaling cascade genes in relation to Wnt,
Tgfβ, Delta/Notch, Hh and RA signaling pathways across the *P. miniata* ovarian cell type groups. (**B**)
Schematic representation of the predicted paracrine crosstalk between follicle cells, neurons and germ cell
cysts. (**C**) HCR for *tgfb2* labeling follicle cells. Scale bar is 40 μm. (**D-F**) HCRs for *tgfbr3*, *alk2* and *smad5*
show localization within the germ cell cysts and oocytes spatially confirming the scRNA-seq predictions.

Scale bar is 20 µm. Dotted lines outline the germ cell cysts. Nuclei were labelled with DAPI (cyan). fc, follicle
cells; o, oocytes.

Evidence for pre-chordate origins of the HPG axis

Vertebrate oogenesis is regulated by endocrine signaling between the hypothalamus, the

pituitary gland, and the ovary. Therefore, we asked how the ovarian nervous system might

serve as an intrinsic paracrine regulator of oogenesis. We found that most neuropeptide

synthesis genes are specifically expressed by neuronal cluster 20, including, tryptophan

hydroxylase (*tph*), tyrosine hydroxylase (*th*), choline O-acetyltransferase (*chat*), histidine

decarboxylase (*hdc*), and glutamate decarboxylase (*gad*) are expressed , indicating

serotonergic, dopaminergic, cholinergic, histaminergic and GABAergic neurons,

respectively (**Fig. S9A**) (72,73). Moreover, we found that sixteen different neuropeptides

are expressed in the same cell cluster (**Fig. S9A**). Immunofluorescence for TH and

serotonin (5HT) confirmed the presence of extensive dopaminergic and serotonergic

neuronal subepidermal networks respectively (**Fig. S9C-D'**). To our surprise, we also

detected serotonin within the germ cell cysts, (**Fig. 7A-B**) as well as genes encoding for

neuromodulators and neuropeptides including *tph, chat*, prolactin-releasing peptide

(*prrp*), luqin (*lqp*) and uncharacterized neuropeptide precursor (*np25*) (**Fig. S9A**). The

widespread presence of neuromodulators within the germ cell cysts prompted us to

further assess the link between ovarian neurons and germ cell cysts (**Fig. 7C**).

Importantly, most of the ligand/receptor pairs used for this analysis have been previously

identified and some even de-orphanized in sea stars and sea urchins (74–80). We found

that neuropeptides involved in the function and regulation of the mammalian HPG axis,

including gonadotropin-releasing hormone (*gnrh*), kisspeptin (*kpp*), *oxy/vp* (81) and the

echinoderm muscle relaxant neuropeptide F-Salmf (76) were specifically expressed in

the neuron cell type group and in both germ cell groups (**Fig. 7C**). Their receptors were

predominantly expressed in the germ cell cyst clusters, suggesting crosstalk between

neurons and germ cell cysts (**Fig. S7C**). Similarly, the receptors for serotonin, dopamine

and GABA were found expressed within the germ cell cyst clusters (**Fig. S9B**). On the

other hand, we found the ngffyamide (*ngffap*) neuropeptide to be expressed primarily in

the germ and oocytes cell type groups and not in the neurons, while the latter contained

transcripts for its receptor (**Fig. S7C**). Moreover, we also found neuropeptides including

prrp and *lqp* as well as their receptors expressed by both germ cell cysts and neurons

(**Fig. S7C**). Last, we report additional putative neuropeptidergic crosstalk of the ovarian

neurons with the muscles and the ovarian epithelium (**Fig. S7C**).

Figure 7. Neuropeptidergic profile of the *P. miniata* intrinsic ovarian nervous system. (A-B)
Immunofluorescence of 5HT and Myosin II. Dotted lines outline the germ cell cysts. Scale bars 20 µm. **(C)**
Dotplot showing the average expression of the reconstructed and/or deorphanized pairs of
neuropeptides/receptors. **(D-D')** HCR for *oxy/vp* and *nanos1* shows the relatively short distance between
the oxy/vp positive neurons and the germ cell cysts. Scale bars in panels are 20 µm. Dotted lines outline
the germ cell cysts. **(E)** Orthogonal views of the *oxy/vp* and *nanos1* HCR, shows the close proximity of the
oxy/vp neuron in respect to the *nanos1* positive germ cell cysts. Dotted lines outline the germ cell cysts.
Scale bar is 10 µm. **(F-F')** HCR for *gnrh* in subepithelial ovarian neurons. Scale bars in panels are 20 µm.
Dotted lines outline the germ cell cysts. **(G)** Orthogonal views of a *gnrh* HCR, highlights the spatial
organization of the *gnrh* positive neurons in close proximity to the germ cell cysts. Scale bar is 10 µm.
Dotted lines outline the germ cell cysts. **(H-H')** HCR for *gnrhr* confirms the single cell predictions of
expression in the germ lineage. Scale bars are 20 µm. Dotted lines outline the germ cell cysts. **(I)** Sankey
plot depicting the alignment scores between adult sea star ovarian cell atlas with the adult mouse pituitary
gland one. **(J)** Schematic representation of the components participating in the predicted crosstalk between
the intrinsic ovarian neurons and the germ cell cysts. Nuclei are labelled with DAPI (cyan). fc, follicle cells;
o, oocytes.
605

To test these predictions, we examined expression of *oxy/vp* as well as for the lowly
expressed *gnrh* and *gnrhr*. Using HCR, we found that oxy/vp are expressed exclusively
in rare subepidermal neurons (1-2 per ovarian lobe, **Fig. 7D-D', S9 E-E'**). These *oxy/vp*
positive neurons were in close proximity to the germ cell cysts, suggesting potential
crosstalk (**Fig. 7E**). Similarly, we detected *gnrh+* cells in proximity to germ cell cysts, while
gnrhr was expressed both in the cysts and oocytes (**Fig. 7F-H'**). Previous studies in the
adult sea star *A. rubens* reported that *gnrh* is expressed predominantly within the sea
star's nervous system (oral nerve ring and radial nerve cords). Consistent with this prior
work, and supporting the specificity of our HCR probes, *gnrh* and *gnrhr* were in
complementary regions of the oral nerve ring and the radial nerve cord (**Fig. S9F-F'**).
616

Taken together, our analysis found that many known hypothalamic neuromodulators
involved in the vertebrate HPG axis such as 5HT, dopamine, GABA, GNRH, KPP and
OXY/VP (81) as well as the pituitary gland secreted PrRP (82) are all present in the sea
star ovarian neurons. To further test if such a system is in place in sea star ovaries, we
assessed the similarity between sea star cell type families with the mammalian pituitary
gland using SAMap (**Fig. 7I**). Surprisingly, this analysis showed that the sea star neurons

exclusively align with pituitary gland gonadotropes, which upon stimulation by

hypothalamic *gnrh* produce the gonadotropins that control sex hormone production,

oogenesis, and ovulation in mammals (83). Our comparison showed alignments between

pituitary gland thyrotropes with the sea star follicle cells (follicle I) while the germ cell cysts

aligned with the pituitary gland lactotropes (**Fig. 7I**). Altogether, our results reveal a

diverse and putatively multifunctional intrinsic ovarian nervous system that may

communicate with the germ cell cysts (**Fig. 7J**). Lastly, our data suggest that the sea star

ovarian nervous system has a dual hypothalamic-like and gonadotropic-like state that

could act as a localized HPG-like center operating in parallel or independently from the

oral nerve ring and radial nerve cords to regulate sea star oogenesis, oocyte maturation

and spawning behavior.

Discussion

Sexual reproduction depends on the interactions between multiple cell types within the

ovary. Ovarian structure varies across animals, reflecting differences in reproductive

strategies and adaptations to different environments. For instance, many marine

invertebrates follow seasonal patterns and release large quantities of gametes into the

water column, whereas in mammals seasonal mating patterns result in the fertilization of

only a limited number of oocytes within the oviduct. Despite these differences, in most

animals studied, the ovary serves two critical roles as both an endocrine organ that

produces steroid hormones, and as the niche for oocyte development. Here we took

advantage of the easily accessible, optically transparent and highly fecund sea star ovary,

representing an extreme case of reproductive output, to explore the identity and the

evolutionary origins of the ovarian cell types in a non-chordate deuterostome. Our single

cell atlas identified nine cell type groups distributed across twenty-seven transcriptionally

distinct clusters which we then spatially mapped within the ovary. We show that the ovary

is made of seven somatic cell groups corresponding to ciliated epithelium, neurons,

muscles, immune cells and three different types of follicle cells, corroborating previous

morphological studies (36). We also found cell groups corresponding to young oocytes

as well as proliferating germ cells organized into cysts. Single cell trajectory analysis finds

connections between the cysts and the oocytes and shows how the gene expression

changes during early oogenesis. Collectively, this analysis provides a framework for
understanding echinoderm oogenesis at a new depth, and a valuable resource for future
investigation of reproductive biology and cell type evolution.

Sea star follicle cells are a heterotypic group

Follicle cells that enwrap oocytes serve important roles in oogenesis across animals (84).
In insects, different types of follicle cells produce yolk components or contribute to the
formation of egg protective layers (85). In mammals and fish, granulosa cells are involved
in the production of estrogen, nutrient transfer and regulation of oocyte maturation
(1,86,87). Theca cells instead populate the outer layers of the follicle and produce steroid
hormones that are taken up and converted to estrogen by granulosa cells. Similarly in
echinoderms, follicle cells produce maturation-inducing molecules, such as 1-
methyladenine in sea stars, upon stimulation by the gonad stimulating substance (GSS)
from the nervous system that triggers the resumption of meiosis in arrested oocytes
(88,89). However, the molecular signature of follicle cells has not been well defined. Here
we provide evidence for three diversified sea star follicle cell groups, one that appears to
be echinoderm specific (follicle II), involved primarily in vitellogenesis, and two that
express genes typically found in granulosa (follicle I) and theca cells (follicle III) in
mammals. Surprisingly, our results suggest possible pre-chordate evolutionary origins for
theca and granulosa cells.

In support of this hypothesis, our transcriptomic and spatial analysis discovered that
several genes associated with mammalian granulosa cells are also expressed by a
subset of sea star follicle cells. This includes the transcription factors *foxl2* and *gata4*, and
the protein hormone inhibin beta (*inhbb*) (1,44). The forkhead transcription factor *foxl2* is
a crucial member of the GRN controlling the granulosa cell proliferation and differentiation
in mammals (43), and in sea stars we find it expressed by a subset of follicle cells that
are in direct contact with the developing oocytes. In addition, we detected primary cilia on
the surface of the follicle cells enwrapping oocytes, which is a conserved feature of
estrogen-producing granulosa cells of antral follicles in mice (90). Our cross-species
analysis revealed that a granulosa-like cell cluster (follicle I) consistently aligned with
granulosa cell clusters in humans and mice, and with follicle cells in the rest of the species

analyzed. This raises the possibility that sea star granulosa cells influence follicle
development through similar mechanisms as in mammals. For instance, in mammals
granulosa cells secrete several Tgfβ family ligands that influence oocyte development
(91). Similarly, we find that such ligands are expressed by the granulosa-like (follicle)
cells. Last, our scRNA-seq data suggest that sea star granulosa-like cells produce and
respond to RA signaling, which prior work has been found to regulate granulosa cell
proliferation and steroidogenesis in mammals (92). Moreover, our data show that the
follicle III cell group expresses star and *cyp17a1*, genes essential for steroidogenesis in
mammalian theca cells. The sea star ovarian epithelium shares similarities with theca
cells in zebrafish, mouse, and fetal human ovaries, with two clusters expressing *cyp17a1*
and one expressing star, indicating a potential role in steroid hormone production similar
to what was previously reported in humans (93). Taken together our data suggest that
the sea star ovary harbors follicle cell types evolutionarily conserved to distinct follicle cell
populations in other deuterostomes as well as protostomes.

Molecular identification of putative oogonial stem cells

An evolutionarily conserved feature of oogenesis is the formation of germ cell cysts.
Clusters of cytoplasmically interconnected oogonia have been found in the ovaries of a
variety of animals including insects, amphibians, reptiles, birds, and mammals (35,94–
99). Here we define the germline identity of sea star germ cell nests and provide evidence
for intercellular bridges within them, thus suggesting that germ cell cysts are conserved
in non-chordate deuterostomes. Using 3D segmentation and reconstruction, we also
defined the spatial organization of oogenesis, in which germ cell cysts are localized to the
periphery of the ovarian lobe along with young oocytes organized in primary follicles. Cyst
expressed genes include the conserved germline markers *piwi1*, *nanos1*, *nanos3*, *boll*,
msl3, meiotic recombination components *sycp1* and *mlh1*, and transcriptional regulators
including *figla*, *atrx1* and *cxxc1* (59,100–102). We also detect cell proliferation within the
cysts, supporting them as the likely source of new oogonia and oocytes in the adult ovary
(103). The conserved germline RNA helicase *ddx4/vasa* is inactive in the cysts, but is
expressed in early follicles, similar to what was shown in marsupial fetal germ cells (104-

715 106). This repression may contribute to maintaining an undifferentiated state, as seen in

716 fruit flies where *ddx4* knockout led to arrested oocyte development (107). Additionally, we

717 hypothesize that the self-renewal-meiotic entry decision could be regulated by RA, Wnt,

718 Tgfβ and Hh pathways, whose respective receptors and intracellular components were

719 found expressed in the germ cell cysts. These results support an ancient role for these

720 pathways in oogenesis, as demonstrated in other species including mice and amphibians

721 (108–111).

723 Neuronal control of oogenesis predates the establishment of chordate HPG axis

724 Oogenesis and ovulation are regulated by neuroendocrine signals from the nervous

725 system in many animals. In vertebrates, this signaling occurs between the hypothalamus,

726 the pituitary gland, and the gonad, also known as the HPG axis. HPG signaling molecules

727 include neuropeptides *gnrh* and *oxy/vp* that are secreted by the hypothalamus and

728 received by gonadotropin cells within the pituitary gland. This induces the secretion of the

729 gonadotropin hormones follicle stimulating hormone (FSH) and luteinizing hormone (LH),

730 that in turn regulate gametogenesis within the gonad through activation of the

731 steroidogenic pathway. In mammals, the secretion of upstream hypothalamic

732 neuropeptides is regulated by several neuromodulators including 5HT, dopamine, GABA,

733 PrRP and KPP (112–116). How might oogenesis and ovulation be regulated in animals

734 like sea stars that lack a central nervous system?

735 Foundational work in sea stars found that soluble signals from radial nerves induce oocyte

736 maturation and spawning, indicating that oogenesis is under neural control even in

737 invertebrates (117). Our data further argue that the sea star ovary itself contains an

738 intrinsic neurochemical regulatory system. For instance, we found that the epithelium,

739 germ cells, and oocytes produce previously identified neuropeptides such as *rgpp* and

740 *trh*, both of which have been found to be produced by the radial nerve cords and regulate

741 oogenesis in echinoderms (118,119). We also found numerous other HPG-associated

742 signaling factors, including 5HT, dopamine, GABA, KPP, GNRH, PrRP and OXY/VP to

743 be expressed in the intrinsic ovarian nerve plexus and their respective receptors in the

744 germ cell cysts. Strikingly, our data suggest the presence of a novel echinoderm ovarian

cell type that features a dual molecular identity resembling chordate hypothalamic neurons and pituitary gonadotropic endocrine cells. In support of this is the similarity of the sea star neuronal cluster with the one corresponding to the mouse pituitary gland gonadotropes.

Taken together our analysis provides molecular evidence towards the presence of a hypothalamic-like and gonadotropic-like intrinsic ovarian neuroendocrine signaling system in non-chordate deuterostomes. We hypothesize that the echinoderm ovarian neurons are analogous to the mammalian HPG axis elements. We speculate that during animal evolution, oogenesis was initially regulated by intrinsic neuroendocrine systems within the ovary and as organisms became more complex the regulatory role was co-opted by distinct organs such as complex interconnected nervous systems and endocrine glands.

Acknowledgements

We thank Jamie MacKinnon, Akshay Kane, Beverly Naigles and Karen Velez, current and past members of the Swartz lab, for advice and fruitful discussions throughout the duration of this project. We thank Maria Ina Arnone (Stazione Zoologica Anton Dohrn) for helping by providing the 1e11 and Mhc antibodies, and *S. purpuratus* ovaries. We also thank Rossella Annunziata (Stazione Zoologica Anton Dohrn) for kindly providing us with *H. tubulosa* ovaries. We would also like to thank Louie Kerr, the Central Microscopy Facility at the Marine Biological Laboratory (Woods Hole, MA, USA) and Chris Bjornsson (ZEISS), for imaging assistance with confocal microscopy. We are grateful to Lisa Abbo and the Marine Resources Center staff for providing excellent animal care and husbandry of adult *P. miniata* used in this work.

Funding

This work was supported by the National Institutes of Health (R00HD099315 to S.Z.S.) and by a Faculty Recruitment Gift from the Hibbitt Foundation. P.P. was supported by a Postdoctoral fellowship awarded by the Global Consortium for Productive Health

(previously known as Global Consortium for Reproductive Longevity and Equality) part of

the Buck Center for Healthy Aging in Women.

Competing interests

The authors declare no competing interests.

Author contributions

Conceptualization: P.P., S.Z.S.; Methodology: P.P., S.Z.S.; Formal analysis: P.P., C.W.,

D.V.; Investigation: P.P., S.Z.S.; Writing - original draft: P.P.; Writing - review & editing:

P.P., C.W., D.V., S.Z.S.; Visualization: P.P., C.W.; Supervision: S.Z.S.; Project

administration: S.Z.S.; Funding acquisition: S.Z.S.

References

1.   Prasasya RD, Mayo KE. Regulation of follicle formation and development by
ovarian signaling pathways. In: The Ovary. Elsevier; 2019. p. 23–49.

2.   Orisaka M, Tajima K, Tsang BK, Kotsuji F. Oocyte-granulosa-theca cell interactions
during preantral follicular development. J Ovarian Res. 2009 Jul 9;2(1):9.

3.   Tsukamura H. Kobayashi Award 2019: The neuroendocrine regulation of the
mammalian reproduction. Gen Comp Endocrinol. 2022 Jan 1;315(113755):113755.

4.   Grive KJ, Freiman RN. The developmental origins of the mammalian ovarian
reserve. Development. 2015 Aug 1;142(15):2554–63.

5.   Packer C, Tatar M, Collins A. Reproductive cessation in female mammals. Nature.
1998 Apr 23;392(6678):807–11.

6.   Armstrong AR, Drummond-Barbosa D. Insulin signaling acts in adult adipocytes via
GSK-3β and independently of FOXO to control Drosophila female germline stem
cell numbers. Dev Biol. 2018 Aug 1;440(1):31–9.

7.   Juliano CE, Swartz SZ, Wessel GM. A conserved germline multipotency program.
Development. 2010 Dec;137(24):4113–26.

8.   Arnone MI, Byrne M, Martinez P. Echinodermata. In: Evolutionary Developmental
Biology of Invertebrates 6. Vienna: Springer Vienna; 2015. p. 1–58.

9.   Bodnar AG. Marine invertebrates as models for aging research. Exp Gerontol. 2009
Aug;44(8):477–84.

10.   Ernst SG. Offerings from an urchin. Dev Biol. 2011 Oct 15;358(2):285–94.

11.   Arnone MI, Andrikou C, Annunziata R. Echinoderm systems for gene regulatory
studies in evolution and development. Curr Opin Genet Dev. 2016 Aug;39:129–37.

12.   Swartz SZ, McKay LS, Su KC, Bury L, Padeganeh A, Maddox PS, et al. Quiescent
cells actively replenish CENP-A nucleosomes to maintain centromere identity and
proliferative potential. Dev Cell. 2019 Oct 7;51(1):35–48.e7.

13.   Paganos P, Voronov D, Musser JM, Arendt D, Arnone MI. Single-cell RNA
sequencing of the Strongylocentrotus purpuratus larva reveals the blueprint of
major cell types and nervous system of a non-chordate deuterostome. Elife. 2021
Nov 25;10.

14.   Magri MS, Voronov D, Foley S, Martínez-García PM, Franke M, Cary GA, et al.
Deep conservation of cis-regulatory elements and chromatin organization in
echinoderms uncover ancestral regulatory features of animal genomes. bioRxiv.
2024.

15.   Dainat J, Hereñú D, Murray KD, Davis E, Ugrin I, Crouch K, et al.
NBISweden/AGAT: AGAT-v1.4.1. Zenodo. 2024.

16.   Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al. Massively
parallel digital transcriptional profiling of single cells. Nat Commun. 2017 Jan
16;8(1):14049.

17.   Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, et al.
Dictionary learning for integrative, multimodal and scalable single-cell analysis. Nat
Biotechnol. 2024 Feb;42(2):293–304.

18.   Chitiashvili T, Dror I, Kim R, Hsu FM, Chaudhari R, Pandolfi E, et al. Female
human primordial germ cells display X-chromosome dosage compensation despite
the absence of X-inactivation. Nat Cell Biol. 2020 Dec;22(12):1436–46.

19.   Isola JVV, Ocañas SR, Hubbart CR, Ko S, Mondal SA, Hense JD, et al. A single-
cell atlas of the aging mouse ovary. Nat Aging. 2024 Jan;4(1):145–62.

20.   Liu Y, Kossack ME, McFaul ME, Christensen LN, Siebert S, Wyatt SR, et al.
Single-cell transcriptome reveals insights into the development and function of the
zebrafish ovary. Elife. 2022 May 19;11:e76014.

21.   Oulhen N, Morita S, Pieplow C, Onorato TM, Foster S, Wessel G. Conservation
and contrast in cell states of echinoderm ovaries. Mol Reprod Dev. 2024
Aug;91(8):e23721.

22. Rust K, Byrnes LE, Yu KS, Park JS, Sneddon JB, Tward AD, et al. A single-cell
840   atlas and lineage analysis of the adult Drosophila ovary. Nat Commun. 2020 Nov
841   6;11(1):5628.

842   23. Allensworth-James M, Banik J, Odle A, Hardy L, Lagasse A, Moreira ARS, et al.
843   Control of the anterior pituitary cell lineage regulator POU1F1 by the stem cell
844   determinant Musashi. Endocrinology. 2021 Mar 1;162(3):bqaa245.

845   24. Tarashansky AJ, Musser JM, Khariton M, Li P, Arendt D, Quake SR, et al. Mapping
846   single-cell atlases throughout Metazoa unravels cell type evolution. Elife. 2021 May
847   4;10:e66747.

848   25. Piovani L, Leite DJ, Yañez Guerra LA, Simpson F, Musser JM, Salvador-Martínez I,
849   et al. Single-cell atlases of two lophotrochozoan larvae highlight their complex
850   evolutionary histories. Sci Adv. 2023 Aug 2;9(31):eadg6034.

851   26. Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, et al. The single-cell
852   transcriptional landscape of mammalian organogenesis. Nature. 2019
853   Feb;566(7745):496–502.

854   27. Hörl D, Rojas Rusak F, Preusser F, Tillberg P, Randel N, Chhetri RK, et al.
855   BigStitcher: reconstructing high-resolution image datasets of cleared and expanded
856   samples. Nat Methods. 2019 Sep;16(9):870–4.

857   28. Formery L, Peluso P, Kohnle I, Malnick J, Thompson JR, Pitel M, et al. Molecular
858   evidence of anteroposterior patterning in adult echinoderms. Nature. 2023
859   Nov;623(7987):555–61.

860   29. Kuehn E, Clausen DS, Null RW, Metzger BM, Willis AD, Özpolat BD. Segment
861   number threshold determines juvenile onset of germline cluster expansion in
862   Platynereis dumerilii. J Exp Zool B Mol Dev Evol. 2022 Jun;338(4):225–40.

863   30. Paganos P, Caccavale F, La Vecchia C, D'Aniello E, D'Aniello S, Arnone MI. FISH
864   for all: A fast and efficient fluorescent in situ hybridization (FISH) protocol for
865   marine embryos and larvae. Front Physiol. 2022 Apr 19;13:878062.

866   31. Perillo M, Paganos P, Spurrell M, Arnone MI, Wessel GM. Methodology for whole
867   mount and fluorescent RNA in situ hybridization in echinoderms: Single, double,
868   and beyond. In: Methods in Molecular Biology. New York, NY: Springer US; 2021.
869   p. 195–216. (Methods in molecular biology (Clifton, N.J.)).

870   32. Ćorić A, Stockinger AW, Schaffer P, Rokvić D, Tessmar-Raible K, Raible F. A fast
871   and versatile method for simultaneous HCR, immunohistochemistry and Edu
872   labeling (SHInE). Integr Comp Biol. 2023 Aug 23;63(2):372–81.

873   33. Mita M, Osugi T, Takahashi T, Watanabe T, Satake H. Mechanism of gamete
874   shedding in starfish: Involvement of acetylcholine in extracellular Ca2+-dependent
875   contraction of gonadal walls. Gen Comp Endocrinol. 2020 May

1;290(113401):113401.

34. Zickler D, Kleckner N. The leptotene-zygotene transition of meiosis. Annu Rev
Genet. 1998;32:619–97.

35. Pepling ME. From primordial germ cell to primordial follicle: mammalian female
germ cell development. Genesis. 2006 Dec;44(12):622–32.

36. Schoenmakers HJ, Colenbrander PH, Peute J, van Oordt PG. Anatomy of the
ovaries of the starfish Asterias rubens (Echinodermata). A histological and
ultrastructural study. Cell Tissue Res. 1981;217(3):577–97.

37. Fresques T, Swartz SZ, Juliano C, Morino Y, Kikuchi M, Akasaka K, et al. The
diversity of nanos expression in echinoderm embryos supports different
mechanisms in germ cell specification. Evol Dev. 2016 Jul;18(4):267–78.

38. Reich A, Dunn C, Akasaka K, Wessel G. Phylogenomic analyses of Echinodermata
support the sister groups of Asterozoa and Echinozoa. PLoS One. 2015 Mar
20;10(3):e0119627.

39. Zazueta-Novoa V, Onorato TM, Reyes G, Oulhen N, Wessel GM. Complexity of
yolk proteins and their dynamics in the sea star Patiria miniata. Biol Bull. 2016
Jun;230(3):209–19.

40. Xia L, Yin GL, Long Y, Sun F, Wang BB, Zhu Y, et al. Essential role of CFAP77-
CCDC105-TEX43 subcomplex in connecting axonemal A and B tubules for
mammalian sperm motility. Cell Biology. bioRxiv. 2025.

41. Fan X, Bialecka M, Moustakas I, Lam E, Torrens-Juaneda V, Borggreven NV, et al.
Single-cell reconstruction of follicular remodeling in the human adult ovary. Nat
Commun. 2019 Jul 18;10(1):3164.

42. Juengel JL, Bibby AH, Reader KL, Lun S, Quirke LD, Haydon LJ, et al. The role of
transforming growth factor-beta (TGF-beta) during ovarian follicular development in
sheep. Reprod Biol Endocrinol. 2004 Nov 25;2(1):78.

43. Pisarska MD, Barlow G, Kuo FT. Minireview: roles of the forkhead transcription
factor FOXL2 in granulosa cell biology and pathology. Endocrinology. 2011
Apr;152(4):1199–208.

44. Bennett J, Baumgarten SC, Stocco C. GATA4 and GATA6 silencing in ovarian
granulosa cells affects levels of mRNAs involved in steroidogenesis, extracellular
structure organization, IGF-I activity, and apoptosis. Endocrinology. 2013
Dec;154(12):4845–58.

45. Liu Z, Castrillon DH, Zhou W, Richards JS. FOXO1/3 depletion in granulosa cells
alters follicle growth, death and regulation of pituitary FSH. Mol Endocrinol. 2013
Feb;27(2):238–52.

46. Vos MC, van der Wurff AA, Last JTJ, de Boed EAM, Smeenk JMJ, van Kuppevelt
TH, et al. Immunohistochemical expression of MMP-14 and MMP-2, and MMP-2
activity during human ovarian follicular development. Reprod Biol Endocrinol. 2014
Jan 31;12(1):12.

47. Abramovich D, Rodriguez Celin A, Hernandez F, Tesone M, Parborell F.
Spatiotemporal analysis of the protein expression of angiogenic factors and their
related receptors during folliculogenesis in rats with and without hormonal
treatment. J Reprod Fertil. 2009 Feb;137(2):309–20.

48. Kiriakidou M, McAllister JM, Sugawara T, Strauss JF 3rd. Expression of
steroidogenic acute regulatory protein (StAR) in the human ovary. J Clin Endocrinol
Metab. 1996 Nov;81(11):4122–8.

49. Yamamoto S, Konishi I, Tsuruta Y, Nanbu K, Mandai M, Kuroda H, et al.
Expression of vascular endothelial growth factor (VEGF) during folliculogenesis and
corpus luteum formation in the human ovary. Gynecol Endocrinol. 1997
Dec;11(6):371–81.

50. Estienne A, Relav L, Benkoura M, Monniaux D, Morin F, Fabre S, et al. Endothelial
cell-derived fibroblast growth factor-18 regulates ovarian function in sheep. J Cell
Physiol. 2022 May;237(5):2528–38.

51. Chai C, Liang L, Mikkelsen NS, Wang W, Zhao W, Sun C, et al. Single-cell
transcriptome analysis of epithelial, immune, and stromal signatures and
interactions in human ovarian cancer. Commun Biol. 2024 Jan 26;7(1):131.

52. Bayly-Jones C, Pang SS, Spicer BA, Whisstock JC, Dunstone MA. Ancient but not
forgotten: New insights into MPEG1, a macrophage perforin-like immune effector.
Front Immunol. 2020 Oct 15;11:581906.

53. Bülow S, Zeller L, Werner M, Toelge M, Holzinger J, Entzian C, et al.
Bactericidal/permeability-increasing protein is an enhancer of bacterial lipoprotein
recognition. Front Immunol. 2018 Dec 5;9:2768.

54. Baldwin LA, Hoff JT, Lefringhouse J, Zhang M, Jia C, Liu Z, et al. CD151-α3β1
integrin complexes suppress ovarian tumor growth by repressing slug-mediated
EMT and canonical Wnt signaling. Oncotarget. 2014 Dec 15;5(23):12203–17.

55. Wood NJ, Mattiello T, Rowe ML, Ward L, Perillo M, Arnone MI, et al.
Neuropeptidergic systems in Pluteus larvae of the sea urchin Strongylocentrotus
purpuratus: Neurochemical complexity in a "simple" nervous system. Front
Endocrinol (Lausanne). 2018 Oct 25;9:628.

56. Pagowski V. A description of the bat star nervous system throughout larval
ontogeny. Evol Dev. 2024 Jan;26(1):e12468.

57. Kee K, Angeles VT, Flores M, Nguyen HN, Reijo Pera RA. Human DAZL, DAZ and

BOULE genes modulate primordial germ-cell and haploid gamete formation.
Nature. 2009 Nov 12;462(7270):222–5.

58. Gonzalez LE, Tang X, Lin H. Maternal Piwi regulates primordial germ cell
development to ensure the fertility of female progeny in Drosophila. Genetics. 2021
Aug 26;219(1).

59. McCarthy A, Sarkar K, Martin ET, Upadhyay M, Jang S, Williams ND, et al. Msl3
promotes germline stem cell differentiation in female Drosophila. Development.
2022 Jan 1;149(1).

60. Sage J, Martin L, Meuwissen R, Heyting C, Cuzin F, Rassoulzadegan M. Temporal
and spatial control of the Sycp1 gene transcription in the mouse meiosis: regulatory
elements active in the male are not sufficient for expression in the female gonad.
Mech Dev. 1999 Jan;80(1):29–39.

61. Hu W, Gauthier L, Baibakov B, Jimenez-Movilla M, Dean J. FIGLA, a basic helix-
loop-helix transcription factor, balances sexually dimorphic gene expression in
postnatal oocytes. Mol Cell Biol. 2010 Jul;30(14):3661–71.

62. Raz E. The function and regulation of vasa-like genes in germ-cell development.
Genome Biol. 2000 Sep 1;1(3):REVIEWS1017.

63. Campolo F, Gori M, Favaro R, Nicolis S, Pellegrini M, Botti F, et al. Essential role of
Sox2 for the establishment and maintenance of the germ cell line. Stem Cells. 2013
Jul;31(7):1408–21.

64. Moore FL, Jaruzelska J, Fox MS, Urano J, Firpo MT, Turek PJ, et al. Human
Pumilio-2 is expressed in embryonic stem cells and germ cells and interacts with
DAZ (Deleted in AZoospermia) and DAZ-like proteins. Proc Natl Acad Sci U S A.
2003 Jan 21;100(2):538–43.

65. Nilsson EE, Doraiswamy V, Skinner MK. Transforming growth factor-beta isoform
expression during bovine ovarian antral follicle development. Mol Reprod Dev.
2003 Nov;66(3):237–46.

66. Zarate-Garcia L, Lane SIR, Merriman JA, Jones KT. FACS-sorted putative oogonial
stem cells from the ovary are neither DDX4-positive nor germ cells. Sci Rep. 2016
Jun 15;6(1):27991.

67. Huyhn K, Renfree MB, Graves JA, Pask AJ. ATRX has a critical and conserved
role in mammalian sexual differentiation. BMC Dev Biol. 2011 Jun 14;11(1):39.

68. Yu C, Fan X, Sha QQ, Wang HH, Li BT, Dai XX, et al. CFP1 regulates histone
H3K4 trimethylation and developmental potential in mouse oocytes. Cell Rep. 2017
Aug 1;20(5):1161–72.

69. Jiang Y, Zhang HY, Lin Z, Zhu YZ, Yu C, Sha QQ, et al. CXXC finger protein 1-

mediated histone H3 lysine-4 trimethylation is essential for proper meiotic
crossover formation in mice. Development. 2020 Mar 17;147(6):dev183764.

70. Moreno SG, Attali M, Allemand I, Messiaen S, Fouchet P, Coffigny H, et al.
TGFbeta signaling in male germ cells regulates gonocyte quiescence and fertility in
mice. Dev Biol. 2010 Jun 1;342(1):74–84.

71. Belli M, Shimasaki S. Molecular aspects and clinical relevance of GDF9 and
BMP15 in ovarian function. Vitam Horm. 2018;107:317–48.

72. D'Aniello E, Paganos P, Anishchenko E, D'Aniello S, Arnone MI. Comparative
Neurobiology of Biogenic Amines in Animal Models in Deuterostomes. Front Ecol
Evol. 2020 Sep 25;8.

73. Bhat US, Shahi N, Surendran S, Babu K. Neuropeptides and behaviors: How small
peptides regulate nervous system function and behavioral outputs. Front Mol
Neurosci. 2021 Dec 2;14:786471.

74. Tian S, Zandawala M, Beets I, Baytemur E, Slade SE, Scrivens JH, et al.
Urbilaterian origin of paralogous GnRH and corazonin neuropeptide signalling
pathways. Sci Rep. 2016 Jun 28;6:28788.

75. Yañez-Guerra LA, Zhong X, Moghul I, Butts T, Zampronio CG, Jones AM, et al.
Echinoderms provide missing link in the evolution of PrRP/sNPF-type neuropeptide
signalling. Elife. 2020 Jun 24;9.

76. Escudero Castelán N, Semmens DC, Guerra LAY, Zandawala M, Dos Reis M,
Slade SE, et al. Receptor deorphanization in an echinoderm reveals kisspeptin
evolution and relationship with SALMFamide neuropeptides. BMC Biol. 2022 Aug
24;20(1):187.

77. Yañez-Guerra LA, Delroisse J, Barreiro-Iglesias A, Slade SE, Scrivens JH, Elphick
MR. Discovery and functional characterisation of a luqin-type neuropeptide
signalling system in a deuterostome. Sci Rep. 2018 May 8;8(1):7220.

78. Odekunle EA, Semmens DC, Martynyuk N, Tinoco AB, Garewal AK, Patel RR, et
al. Ancient role of vasopressin/oxytocin-type neuropeptides as regulators of feeding
revealed in an echinoderm. BMC Biol. 2019 Jul 31;17(1):60.

79. Semmens DC, Dane RE, Pancholi MR, Slade SE, Scrivens JH, Elphick MR.
Discovery of a novel neurophysin-associated neuropeptide that triggers cardiac
stomach contraction and retraction in starfish. J Exp Biol. 2013 Nov 1;216(Pt
21):4047–53.

80. Perillo M, Arnone MI. Characterization of insulin-like peptides (ILPs) in the sea
urchin Strongylocentrotus purpuratus: insights on the evolution of the insulin family.
Gen Comp Endocrinol. 2014 Sep 1;205:68–79.

81. Acevedo-Rodriguez A, Kauffman AS, Cherrington BD, Borges CS, Roepke TA,
Laconi M. Emerging insights into hypothalamic-pituitary-gonadal axis regulation
and interaction with stress signalling. J Neuroendocrinol. 2018 Oct;30(10):e12590.

82. Szukiewicz D. Current insights in prolactin signaling and ovulatory function. Int J
Mol Sci. 2024 Feb 6;25(4):1976.

83. Clay CM, Cherrington BD, Navratil AM. Plasticity of anterior pituitary gonadotrope
cells facilitates the pre-ovulatory LH surge. Front Endocrinol (Lausanne).
2020;11:616053.

84. Albertini D, Bromfield J. Soma-Germline Interactions in the Ovary: An Evolutionary
Perspective. In: Oogenesis. Chichester, UK: John Wiley & Sons, Ltd; 2010. p. 103–
13.

85. Wu Z, Yang L, He Q, Zhou S. Regulatory mechanisms of vitellogenesis in insects.
Front Cell Dev Biol. 2020;8:593613.

86. Liu W, Xin Q, Wang X, Wang S, Wang H, Zhang W, et al. Estrogen receptors in
granulosa cells govern meiotic resumption of pre-ovulatory oocytes in mammals.
Cell Death Dis. 2017 Mar 9;8(3):e2662–e2662.

87. Planas JV, Athos J, Goetz FW, Swanson P. Regulation of ovarian steroidogenesis
in vitro by follicle-stimulating hormone and luteinizing hormone during sexual
maturation in Salmonid Fish1. Biol Reprod. 2000 May 1;62(5):1262–9.

88. Shirai H, Kanatani H, Taguchi S. 1-methyladenine biosynthesis in starfish ovary:
action of gonad-stimulating hormone in methylation. Science. 1972 Mar
24;175(4028):1366–8.

89. Kanatani H, Shirai H, Nakanishi K, Kurokawa T. Isolation and identification of
meiosis inducing substance in starfish Asterias amurensis. Nature. 1969
Jan;221(5177):273–4.

90. Johnson ET, Nicola T, Roarty K, Yoder BK, Haycraft CJ, Serra R. Role for primary
cilia in the regulation of mouse ovarian function. Dev Dyn. 2008 Aug;237(8):2053–
60.

91. Trombly DJ, Woodruff TK, Mayo KE. Roles for transforming growth factor beta
superfamily proteins in early folliculogenesis. Semin Reprod Med. 2009
Jan;27(1):14–23.

92. Kipp JL, Golebiowski A, Rodriguez G, Demczuk M, Kilen SM, Mayo KE. Gene
expression profiling reveals Cyp26b1 to be an activin regulated gene involved in
ovarian granulosa cell proliferation. Endocrinology. 2011 Jan;152(1):303–12.

93. Ivarsson K, Sundfeldt K, Brännström M, Janson PO. Production of steroids by
human ovarian surface epithelial cells in culture: possible role of progesterone as

growth inhibitor. Gynecol Oncol. 2001 Jul;82(1):116–21.

94. Kloc M, Tworzydło W, Szklarzewicz T. Germline and somatic cell syncytia in insects. Results Probl Cell Differ. 2024;71:47–63.

95. Tingen C, Kim A, Woodruff TK. The primordial pool of follicles and nest breakdown in mammalian ovaries. Mol Hum Reprod. 2009 Dec;15(12):795–803.

96. Roco ÁS, Ruiz-García A, Bullejos M. Testis development and differentiation in amphibians. Genes (Basel). 2021 Apr 16;12(4):578.

97. Smaga CR, Bock SL, Johnson JM, Parrott BB. Sex determination and ovarian development in reptiles and amphibians: From genetic pathways to environmental influences. Sex Dev. 2023;17(2-3):99–119.

98. Hall GB, Long JA, Wood BJ, Bedecarrats GY. Germ cell dynamics during nest breakdown and formation of the primordial follicle pool in the domestic turkey (Meleagris gallopavo). Poult Sci. 2020 May;99(5):2746–56.

99. Spradling AC, Niu W, Yin Q, Pathak M, Maurya B. Conservation of oocyte development in germline cysts from Drosophila to mouse. Elife. 2022 Nov 29;11.

100. Wang Z, Liu CY, Zhao Y, Dean J. FIGLA, LHX8 and SOHLH1 transcription factor networks regulate mouse oocyte growth and differentiation. Nucleic Acids Res. 2020 Apr 17;48(7):3525–41.

101. De La Fuente R, Viveiros MM, Wigglesworth K, Eppig JJ. ATRX, a member of the SNF2 family of helicase/ATPases, is required for chromosome alignment and meiotic spindle organization in metaphase II stage mouse oocytes. Dev Biol. 2004 Aug 1;272(1):1–14.

102. Lesch BJ, Page DC. Genetics of germ cell development. Nat Rev Genet. 2012 Nov;13(11):781–94.

103. Nicholls PK, Schorle H, Naqvi S, Hu YC, Fan Y, Carmell MA, et al. Mammalian germ cells are determined after PGC colonization of the nascent gonad. Proc Natl Acad Sci U S A. 2019 Dec 17;116(51):25677–87.

104. Hickford DE, Frankenberg S, Pask AJ, Shaw G, Renfree MB. DDX4 (VASA) is conserved in germ cell development in marsupials and monotremes. Biol Reprod. 2011 Oct;85(4):733–43.

105. Hartung O, Forbes MM, Marlow FL. Erratum: Zebrafish vasa is required for germ-cell differentiation and maintenance. Mol Reprod Dev. 2016 Dec;83(12):1128–30.

106. Chen Y, Lin X, Dai J, Bai Y, Liu F, Luo D. Deletion of ddx4 ovary-specific transcript causes dysfunction of meiosis and derepress of DNA transposons in zebrafish ovaries. Biology (Basel). 2024 Dec 16;13(12).

107. Liu N, Han H, Lasko P. Vasa promotes *Drosophila* germline stem cell differentiation
by activating *mei-P26* translation by directly interacting with a (U)-rich motif in its 3′
UTR. Genes Dev. 2009 Dec 1;23(23):2742–52.

108. Bristol-Gould SK, Kreeger PK, Selkirk CG, Kilen SM, Cook RW, Kipp JL, et al.
Postnatal regulation of germ cells by activin: the establishment of the initial follicle
pool. Dev Biol. 2006 Oct 1;298(1):132–48.

109. Iang Y. Hedgehog pathway inhibition causes primary follicle atresia and decreases
female germline stem cell proliferation capacity or stemness. Stem Cell Res Ther.
2019;10.

110. Le Rolle M, Massa F, Siggers P, Turchi L, Loubat A, Koo BK, et al. Arrest of
WNT/β-catenin signaling enables the transition from pluripotent to differentiated
germ cells in mouse ovaries. Proc Natl Acad Sci U S A. 2021 Jul
27;118(30):e2023376118.

111. Zhang X, Gunewardena S, Wang N. Nutrient restriction synergizes with retinoic
acid to induce mammalian meiotic initiation in vitro. Nat Commun. 2021 Mar
19;12(1):1758.

112. Moguilevsky JA, Wuttke W. Changes in the control of gonadotrophin secretion by
neurotransmitters during sexual development in rats. Exp Clin Endocrinol Diabetes.
2001;109(4):188–95.

113. Liu X, Herbison AE. Dopamine regulation of gonadotropin-releasing hormone
neuron excitability in male and female mice. Endocrinology. 2013 Jan;154(1):340–
50.

114. Camille Melón L, Maguire J. GABAergic regulation of the HPA and HPG axes and
the impact of stress on reproductive function. J Steroid Biochem Mol Biol. 2016
Jun;160:196–203.

115. Kaiser UB. Hyperprolactinemia and infertility: new insights. J Clin Invest. 2012
Oct;122(10):3467–8.

116. Xie Q, Kang Y, Zhang C, Xie Y, Wang C, Liu J, et al. The role of kisspeptin in the
control of the hypothalamic-pituitary-gonadal axis and reproduction. Front
Endocrinol (Lausanne). 2022 Jun 28;13:925206.

117. A.B. Chaet and R.A. McConnaughy. Physiologic Activity of Nerve Extracts. The
Biological Bulletin. 1959;117:407–8.

118. Feng Y, Piñon Gonzalez VM, Lin M, Egertová M, Mita M, Elphick MR. Localization
of relaxin-like gonad-stimulating peptide expression in starfish reveals the
gonoducts as a source for its role as a regulator of spawning. J Comp Neurol. 2023
Sep;531(13):1299–316.

119.Chaiyamoon A, Tinikul R, Nontunha N, Chaichotranunt S, Poomtong T, Sobhon P,
et al. Characterization of TRH/GnRH-like peptides in the sea cucumber, Holothuria
scabra, and their effects on oocyte maturation. Aquaculture. 2020
Mar;518(734814):734814.

Supplementary Figures

Figure S1. Immunohistochemical identification of echinoderm ovarian cell types. (A-D) IHC for Syt1 labeling neurons (A), for Mhc labeling muscles (B) and for myosin (C) and α-tubulin (D) labeling follicle cells. Scale bars for panels A and D are 40 µm; for panel B 60 µm; for C 20 µm. **(E-F)** IHC for Syt1 and Mhc to visualize the interconnection of ovarian neurons and muscles. Scale bars for panels are 20 µm. **(G-H)** IHC for Syt1 and α-tubulin co-labels the ovarian neurons. Scale bars for panels are 20 µm. **(I-J)** IHC for Syt1 and Mhc confirms the presence of the neurons and muscles respectively in the sea urchin *S. purpuratus* ovary. Scale bars are 25 µm. **(K-L)** IHC for α tubulin labels the ovarian neurons in the sea cucumber *H. tubulosa* ovary. Scale bars are 50 µm. Nuclei were labelled with DAPI (cyan). fc, follicle cells; o, oocytes.

Figure S2. Size and spatial distributions of oogonia and oocytes in the sea star ovary. (A-B) Violin

plot showing the experimentally measured diameter (A) and estimated volume (B) of each oogonium within

germ cell cysts. **(C-D)** Violin plot showing the measured diameter (C) and estimated volume (D) of the

smallest *nanos1* positive oocyte in close proximity to the germ cell cysts. **(E-F)** Violin plot of the

segmentation-based calculated volumes (E) and estimated diameter (F) of germ cell cysts, oocytes close

to the ovarian surface, and those within the lumen. **(G)** 3D rendering of segmented germ cell cysts and

oocytes. Scale bar is 100 μm. Ovary seen from a side view. Estimated volumes or diameters were

calculated based on experimentally measured diameters or volumes respectively, assuming the shape of

the oogonia and oocytes is spherical.

1153

Figure S3. *P. miniata* ovary cell atlas and cell contribution of the different libraries. (A) Integrated
UMAP of the three different scRNA-seq libraries, in which color code is indicative of different clusters **(B-**
C) Violin plot showing the experimentally measured diameter (B) and estimated volume (C) of the oogonia
and oocytes within the ovaries used for the reconstruction of the single cell atlas. Due to the capturing
system limitations only the oocytes with a size ≤ 40µm in diameter are included in the single cell atlas. **(D)**
Integrated UMAP of the three different scRNA-seq libraries, split by the cell contribution of each library. The
color code is indicative of different clusters. **(E-E')** HCR for the *nanos3* gene with (E) and without (E') the
nuclei channel (DAPI) confirmed its localization in the germ lineage. Dotted circles in panel E are indicative
of oocytes. Scale bars are 20 µm. **(F-G')** HCR for the *syt1* gene with (F-F') and without (G-G') the nuclei
channel (DAPI) shows expression in the neurons (F-G) as well as the germ cell cysts and oocytes (F'-G').
Dotted circles in F' are indicative of oocytes. Germ cell cysts in panels F' and G' are outlined with dotted
lines. Scale bars are 20 µm. **(H-I) HCR** for the *syt1* gene with (G) and without (G') the nuclei channel (DAPI)
shows expression of the gene only in the nervous system of the *P. miniata* larvae (H) and post-metamorphic
juveniles (I), confirming the specificity of the probe. Scale bars for panels H and I are 50 and 100 µm
respectively. cb, ciliary band; m, mouth, oocytes; oh, oral hood; onr, oral nerve ring; rnc, radial nerve cord.

Figure S4. Cross-species comparison of sea star ovarian cell types with protostome and deuterostome representatives with SAMap. Sankey plot depicting the alignment scores between adult

 sea star ovarian cell atlas with the adult fruit fly ovary **(A)**, adult mouse ovary **(B)** and with sea urchin larva
1173   **(C)**. Alignment scores equal or greater to 0.1 were considered significant.

1174

1175

Figure S5. Identification of mitotic and meiotic molecular signatures in the sea star ovary. (A) Dotplot showing the average expression of the genes used for the scoring analysis shown in Fig.4 A. **(B)** HCR for the meiotic marker *mlh1* labelling the germ cell cysts and oocytes of various sizes. **(C-E)** IHC for Myosin II labeling cell peripheries with the EdU proliferation assay, confirms that cells from both the somatic and germ lineages are actively proliferating in the adult sea star ovary. Nuclei are labelled with DAPI (cyan). Dotted lines outline the germ cell cyst in panels C-E. Scale bars are 20 μm. fc, follicle cells; o, oocytes.

1182

Figure S6. Differential expression patterns along the sea star oogenesis developmental trajectory.

(A) Volcano plot showing the differential expression of the germ cell cyst (clusters 13 and 14) genes versus

the oocyte reference group (clusters 15-19). **(B-B')** Dotplot showing the top 20 enriched gene ontology terms associated with the germ cell cyst (B) and the oocyte (B') upregulated genes. **(C-C''')** HCR for the transcriptional regulators *atrx1* and *cxxc1* paired with *piwi1* shows expression in oocytes. Scale bars are 10 μm.

 Figure S7. Distinct transcription factor profiles during early oogenesis. Heatmap of the top 50
1193 differentially expressed transcription factors across the germline clusters.

1194

1195

1196 **Figure S8. Expression patterns of evolutionarily conserved ovarian signaling pathways. (A)** Dotplot
1197 showing the average expression of the genes used for the scoring analysis shown in Fig. 6A. D,
1198 degradation; L, ligand; S, synthesis; SC, signaling cascade; T, transport. **(B-D)** HCR for the *bmp1* and *bmp4*
1199 ligands confirms their spatial expression in the germ cell cysts and oocytes. Nuclei are labelled with DAPI
1200 (cyan). Dotted lines outline the germ cell cyst. Scale bars are 20 µm. o, oocytes.

1201

Figure S9. Neuropeptides and neurotransmitters expressed in the intrinsic ovarian nervous system.

(A) Dotplot showing the average expression of known echinoderm neurotransmitter synthesizing enzymes

and neuropeptides. **(B)** Dotplot showing the average expression of sea star neurotransmitter receptors. **(C-**

C') Immunofluorescence of TH shows the presence of subepidermal ovarian dopaminergic neurons. Scale

bars are 40 µm. **(D-D')** Immunofluorescence of 5HT confirms the presence of subepidermal ovarian

serotonergic neurons. Scale bars are 40 µm. **(E-E')** HCR for *oxy/vp* neuropeptide marks two *oxy/vp* positive

neurons per ovarian lobe. Scale bars are 20 µm. **(F-F')** HCR for *gnrh* and *gnrhr* shows expression in the

post-metamorphic *P. miniata* nervous system confirming the specificity of the probes. Scale bars are 50

µm. onr, oral nerve ring; rnc, radial nerve cord.