Discovering Polyphosphate and Polyhydroxyalkanoate-Accumulating Organisms Across Ecosystems: Phenotype-targeted Genotyping via FACS-sequencing

Yuan Yan[a,□], Mathew Baldwin[a,□], Jangho Lee[a,b], Zijian Wang[d], Guangyu Li[a], Il Han[a,c], Peisheng He[a], Varun Shrinivasan[a,e], Liam Wickes-Do[f], Michael A. Gore[f] and April Z. Gu[a*]

[a]School of Civil and Environmental Engineering, Cornell University, Ithaca, NY 14850, United States

[b]Department of Chemistry, CUNY Hunter College, 695 Park Ave, New York, NY 10065, United States

[c] Metrohm Spectro, Inc., 313 Enterprise Drive, Plainsboro Township, NJ 08536, United States

[d]School of Biological and Environmental Engineering, Cornell University, Ithaca, NY 14850, United States

[e] Trinnex, Boston MA 02109, United States

[f]Plant Breeding and Genetics Section, School of Integrative Plant Science, Cornell University, Ithaca, NY 14853, United States

□These authors contributed equally to the manuscript.

[*] Corresponding author: April Z. Gu. aprilgu@cornell.edu

Abstract

Intracellular biopolymers serve versatile functions that allow microbes to adapt to fluctuating

environmental conditions. The metabolic interdependence of dual intracellular polymers, namely

polyphosphate (polyP) and polyhydroxyalkanoates (PHA), is a defining feature of functionally

polyphosphate-accumulating organisms (PAOs), the key agents enabling enhanced biological

phosphorus removal (EBPR) for wastewater treatment. However, beyond EBPR systems, the

presence and identities of PAOs that possess both polyP and PHA in natural environments such

as soil have never been examined due to a lack of available detection tools, despite their potential

roles in carbon and phosphorus cycling. This study presents a novel phenotype-targeted approach

integrating triple-stained fluorescence-activated cell sorting (FACS) with 16S rRNA gene

amplicon sequencing (termed TriFlow-Seq) to simultaneously detect, quantify, and

phylogenetically characterize PAOs accumulating both polyP and PHA (referred to as PHA-

PAOs). TriFlow-Seq was validated using polymer staining image analysis and Single-Cell

Raman micro-spectroscopy. Application to EBPR systems successfully enriched known PHA-

PAOs, including *Candidatus* Accumulibacter, *Tetrasphaera*, *Dechloromonas*, *Pseudomonas.* It

also revealed novel candidate PHA-PAOs, particularly within the Rhodobacteraceae family.

When applied to soil samples, TriFlow-Seq led to the first discovery of diverse PHA-PAOs

dominated by *Pseudomonas*, *Halomonas*, and *Nannocystis* in maize rhizosphere soils. These

predominant genera are known rhizosphere inhabitants of essential crops with key plant growth-

promoting functions including phosphate solubilization, biofilm formation, and phytohormone

production, yet simultaneous polyP and PHA accumulation has not been directly reported in the

maize rhizosphere. Our findings revealed unexpectedly high PHA-PAO prevalence and distinct

phylogenetic patterns associated with different maize genotypes, suggesting a potentially

overlooked role for dual polymer storage in microbial rhizosphere dynamics and function. This

study establishes a pioneering approach to investigate dual polyP and PHA-containing PAO

identities and their important roles in rhizosphere structure and plant health.

Key words:

Environmental PAOs, Phenotype-targeted sorting, Triple-staining FACS

Synopsis

Novel method reveals dual-polymer bacteria in activated sludge and crop rhizospheres,

suggesting new candidates supporting wastewater treatment and agricultural sustainability.

1 Introduction

A diverse range of microbes contains intracellular biopolymers, including polyphosphate (polyP), polyhydroxyalkanoates (PHA), glycogen, waxy esters, triacylglycerides (TAGs), trehalose, and cyanophycin.[1] These biopolymers serve versatile functions such as energy storage, nutrient cycling, ion balance, redox homeostasis, and stress resistance, allowing microbes to adapt to fluctuating environmental conditions.[1–3] Among them, polyP-accumulating organisms (PAOs) are key agents in achieving sustainable phosphorus (P) removal and recovery in enhanced biological phosphorus removal (EBPR) systems.[4,5] PAOs exhibit a unique metabolism: under anaerobic conditions, they utilize stored polyP as an energy source, releasing phosphate and concurrently accumulating PHA. Then, under subsequent oxic or anoxic conditions, they uptake phosphate and synthesize new polyP while using stored PHA to fuel growth.[6] This unique metabolism provides competitive advantages for PAOs under fluctuating anaerobic/oxic conditions.[7–9] Therefore, the metabolic interdependence of dual intracellular polymers (polyP and PHA) is a defining feature of functionally verified PAOs in EBPR systems, such as *Candidatus* Accumulibacter,[6] *Pseudomonas*,[10] and *Dechloromonas;*[7] although some, such as *Tetrasphaera,* have distinct sub-groups with varying capacities to accumulate PHA.[11–14]

Beyond EBPR engineering systems, there is increasing interest in exploring the functional roles of PAOs in natural environments such as soil. Emerging evidence points to their ubiquitous presence and versatile functions, where they serve as important facilitators of phosphorus (P) availability, carbon (C) cycling, biofilm formation, and contribute to both plant and microbial community stress resilience.[15–18] In the natural environments, PAOs are often broadly identified as polyP-containing organisms, with little knowledge of the potential co-occurrence and co-metabolisms with C biopolymer storage. To explore this, we performed a

preliminary quantification of polyP-containing organisms (PCOs) in agricultural soil with single-

cell Raman spectroscopy analysis, which indicated that up to 73.7% of PCOs also contain C

storage biopolymers, with a significant Pearson's correlation between PCOs and C-accumulating

PCOs (r=0.8, p-value <0.001) (Figure S1). While exploratory, this unexpected observation of a

high occurrence of PAOs simultaneously possessing both polyP and PHA suggests that dual

storage may be an ecologically relevant functional trait of PAOs in soil. It highlights complex

intra- and interdependent metabolic pathways that remain poorly understood. As the most

common C storage biopolymer, PHA (~1-4 μg C/g soil) responds positively to C availability, but

is suppressed by nitrogen (N) availability, suggesting its role in stoichiometric balancing and

stress response as an alternative carbon allocation strategy.[19,20] The co-storage of C biopolymers

and polyP likely enhances microbial adaptability to nutrient fluctuations and contributes to

broader nutrient cycling and ecosystem functioning. This dual-storage strategy distinguishes

specialized PAOs that exhibit resource-dependent surplus storage from organisms that

accumulate relatively low level of polyP primarily for stress responses or cation sequestration.[20–

22] Fungal PAOs have been more extensively studied in soil,[23] whereas little is known about

bacterial PAOs in managed environments. The inability of polyP-synthesizing fungal organisms,

such as arbuscular mycorrhizal fungi (AMF), to produce PHA provides a clear demarcation

between bacterial and fungal PAOs. Although there have been emerging yet limited

investigations of the presence and roles of PAOs in nature environments, the presence and

identities of bacterial PHA accumulating PAOs (PHA-PAOs) in natural environments have not

yet been elucidated. Particularly, we are interested in identifying bacterial microorganisms

exhibiting surplus storage, defined by their specific ability to accumulate substantial polyP and

PHA beyond immediate biological maintenance, distinguishing them from microbes that

maintain constitutively low levels of these biopolymers for basic cellular functions (e.g.,

Escherichia coli or *Bacillus subtilis*).[24,25]

Despite the ecological, biogeochemical, and engineering significance of PHA-PAOs,

identifying functionally active and phylogenetically novel PAOs exhibiting both intracellular

polyp and PHA storage in complex environmental matrices remains challenging.[15,26–28] Methods

for detecting PAOs that co-accumulate polyP and PHA in natural systems have yet to be reported.

Conventional approaches rely heavily on gene-centric profiling such as polyP-metabolism

biomarkers (i.e., *ppk*) or metagenomics to identify and quantify previously known PAOs across

aquatic and terrestrial systems,[29,30] yet these methods have limitations and cannot reflect actual

phenotypic traits such as active polyP and/or PHA accumulation and co-accumulation. [25–27]

Discovery of novel PAOs is often hindered by *ppk* gene-based detection which relies on prior

genetic information of the target PAOs. Additionally, these methods have low specificity for

detecting PAOs with unique polyP accumulation abilities, as the polyP-synthesizing gene *ppk* is

ubiquitous across microorganisms.[31,32] Moreover, polyP metabolism is not solely regulated by

the *ppk* gene, as alternative proteins, such as actin-like Arp proteins, have been proposed to

exhibit PPK-like functions in bacteria.[33] A few recent studies have employed fluorescence-

activated cell sorting (FACS) to phenotypically track the polyP accumulation traits and resolve

the phylogeny and taxonomy of PAOs from EBPR systems.[34,35] However, these studies used

stains only for polyP, without detecting the co-metabolism polymers like PHA.[34] In addition,

FACS-based PAO detection in soil samples has never been reported.

In this study, we propose a refined definition of functional PAOs in natural environments as

organisms that co-accumulate both polyP and intracellular carbon, predominantly PHA, to

surplus levels, where this co-metabolism holds important ecological roles as discussed above,

and here refer them as PHA-PAOs. We developed TriFlow, a workflow integrating triple-color

fluorescence staining with FACS to enable the simultaneous detection and isolation of

functionally active PAOs that co-accumulate both polyP and PHA. When coupled with 16S

ribosomal RNA (16S rRNA) amplicon sequencing for phylogenetic characterization, the

workflow becomes TriFlow-Seq. This extended method links phenotypic traits to taxonomic

identities. We validated TriFlow by comparing the relative abundances of PHA-PAOs detected

using TriFlow and single cell Raman spectroscopy (SCRS), an established method for the

simultaneous detection of intracellular polyP and PHA in PAOs within EBPR systems.[27,36–39] We

further demonstrated the applicability of TriFlow-Seq across diverse environmental samples,

including activated sludge from four full-scale and one pilot-scale EBPR facilities known to

contain PHA-PAOs, as well as rhizosphere soil from experimental maize fields in New York.

This new method enables us to detect and identify previously unknown bacterial PHA-PAOs and

provide targets for further targeted investigations of novel metabolisms and functional roles of

these unique and prevalent functional groups of organisms across diverse environmental systems.

2   Material and Methods

2.1   Sample collection

2.1.1   Activated sludge samples from EBPR systems

Biomass samples from the aerobic zone were collected from three full-scale and one pilot-scale

plant, all of which effectively remove P and contain active PHA-PAOs. Activated sludge

samples from full-scale EBPR systems were stored at 4°C, transported overnight to the

laboratory, and activated with a phosphorus release/uptake test (Text S1, Figure S2), while

sludge samples from the pilot-scale S2EBPR were directly fixed onsite as described in Section

2.3.

2.1.2 Soil samples

In 2023, two sets of maize genotypes—inbred and hybrid lines (Table S1)—were planted at Cornell University's Musgrave Research Farm in Aurora, NY. The set of inbred lines included five of the 26 parents of the U.S. nested association mapping population,[40] as well as both parents of the intermated B73 × Mo17 population,[41] with B73 serving as a parent in both populations. Collectively, the inbred set included popcorn, sweet corn, temperate, and tropical maize inbred lines. The hybrid set consisted of 22 temperate-adapted hybrids that represented the high-intensity phenotyping site plots provided by the Genomes to Fields (G2F) Initiative (https://www.genomes2fields.org/home/); however, only 10 of the 22 hybrids were sampled in this experiment.

The inbred set was planted on May 18 in a field (42.723288, -76.657356) characterized by a Lima silt loam (0–3% slope), while the hybrid set was planted on May 12 in a field (42.733297, -76.653955) characterized by both Honeoye silt loam (3–8% slopes) and Kendaia & Lyons silt loams (0–3% slope). Both genotype sets were planted in a randomized complete block design with two replicates, with experimental units consisting of two-row plots of varying lengths at each field location.

Plot dimensions and planting rates are described in Table S1. Rhizosphere samples were collected at tasseling (VT) in accordance with Hanway et al.[36] Briefly, within each two-row plot, one plant per row was selected at random and uprooted with bulk soil gently shaken off; the remaining soil adhering to the roots was collected as rhizosphere soil. However, due to greater planting density in hybrid plots, three plants per two-row plot (split between the two rows) were sampled. Samples were pooled by plot (n=2 per genotype), stored, and transported at 4□C for sample pre-processing within 24 hours.

2.2 Microbiota isolation

2.2.1 Activated sludge

Sludge samples were collected from the middle of the aerobic phase and centrifuged to remove the supernatant. The remaining pellets were washed three times with $1 \times$ PBS to remove any excess cations. All centrifuge steps for activated sludge samples were carried out at $3200 \times$ g for 10 minutes due to the easy settleability unless otherwise specified.

2.2.2 Rhizosphere microbiota

Extensive sample pre-treatment and a targeted sequential gating strategy were developed to separate, disaggregate, and isolate PHA-PAOs at the single-cell scale from the soil matrix. The key steps for isolating PHA-PAOs and ensuring single-cell recovery included: 1) detergent-based cell detachment and subsequent cell separation based on sedimentation rate and filtering, 2) disaggregation via chemical and enzymatic washing to destabilize soil microbial aggregates, 3) identification and sorting during gating (Text S2) based on particle size using forward scatter height and area, and 4) triple-fluorescence indicating simultaneous accumulation of DNA, PHA, and polyP.

2.2.2.1 Physical-chemical cell detachment

To isolate the microbial fraction from the soil matrix, 1 gram of fresh soil was suspended in 20 mL of a detachment solution containing 3 mM sodium pyrophosphate, 0.5% Tween 20, and 0.35% polyvinylpyrrolidone in $1 \times$ phosphate-buffered saline (PBS) (0.4 M Na☐HPO☐/NaH☐PO☐, 150 mM NaCl, pH 7.2).[43] The soil suspension was agitated for 30 minutes at room temperature in accordance with Deng et al. 2019,[44] at 200 rpm using an orbital shaker to facilitate detergent based cell detachment (Figure S3, step 2).

2.2.2.2   Cell separation

Following cell detachment on orbital shaker, to enhance cell recovery, the resulting soil slurry

was vortexed for 10 seconds and left to settle for one minute to sediment larger soil particles

(Figure S3, step 3). Due to previously reported wide range of sedimentation rates of soils, known

to vary based on soil texture, physicochemical properties, and particle size, settling time should

be adjusted for individual samples.[45] For these soil samples, the sedimentation time was

optimized as 1 minute based on preliminary time series analysis used to monitor the

sedimentation process (Figure S4). Following sedimentation, cell separation from the soil matrix

was further enhanced by decanting through a 40 μm cell strainer (Corning®, USA, Cat. No.

431750). Here we acknowledge we are unable to directly isolate and ensure cell purity in the

decanted supernatant following filtration due to the small size of many particles within soils; we

further address cell purity during sorting in our gating strategy in Text S2. Decanted and filtered

cell suspension was then pelleted via centrifugation at $6000 \times g$ for 10 minutes with remaining

supernatant decanted (Figure S3, step 3). All centrifuge steps for rhizosphere samples were

carried out at $6000 \times g$ for 10 minutes unless otherwise specified.

2.2.2.3   Cell Cleaning

To enhance microbial dispersion for downstream fluorescent staining and FACS, the detached,

filtered, and pelleted soil community was resuspended and washed once with 5 mL of 4 mM

ethylenediaminetetraacetic acid (EDTA) in $1 \times$ PBS (pH ~7.2) to remove excess cations which

facilitate aggregation of microbial cells in electrostatic interactions with: microbial cell walls,

extracellular polymeric substances (EPS).[46] This was followed by two additional washings with

5mL of $1 \times$ PBS to remove any residual EDTA and solubilized cations (Figure S3, step 4).

2.3 Cell fixation

After microbial isolation in step 2.2, for both soil communities and sludge, the remaining sample pellets were resuspended in 5 mL of fixation buffer, vortexed for 10 s, and stored at 4°C for 24 hours. The fixation buffer was prepared according to previously described protocols, containing 5 mM nickel chloride ($NiCl_2$), 5 mM barium chloride ($BaCl_2$), 10% sodium azide (NaN_3) in 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) buffer (25 mM, pH 7.2) (Figure S3, step 5).[35,47] Previous efforts have noted no significant difference in fixed microbial community patterns within 9 days.[47]

2.4 Triple-staining

Following 24 hours of fixation, both sludge and soil samples were pelleted by centrifugation and washed three times with 1× PBS to remove residual fixative.

Fixed and washed activated sludge samples need an additional homogenization step: sludge was mechanically disrupted with a 26-gauge needle and 1 mL syringe by repeatedly drawing the sludge into the syringe and out vigorously for 5 mins. Compared to soil, activated sludge has a simpler composition with fewer inorganic particles, making it more amenable to single-cell separation by shear. This approach, adapted from established SCRS protocols, effectively isolates individual cells for analysis.[36,37] After homogenization for 5 min, sludge samples were washed an additional three times with $1 \times$ PBS.

After washing steps, both soil and sludge samples were resuspended to 5 mL with $1 \times$ PBS and treated with 1.6 mL solution A (0.11 M citric acid/4.1 Mm Tween 20 in distilled water) for 20 min to permeabilize the cells, then washed and resuspended in 5 mL of $1 \times$ PBS, and filtered through a 40 μm cell strainer (Figure S3, step 6).

Deoxyribonucleic acid (DNA), polyP, and PHA of the cells were stained with SYBR

Green I (SG) (Ex 488 nm / Em 515 nm) (Invitrogen, USA), tetracycline hydrochloride (TC) (Ex

390 nm / Em 560 nm) (Sigma Chemical Co., USA), and nile red (NR) (Ex 532 nm / Em 610 nm)

(Sigma Chemical Co., USA), respectively–each validated in previous studies.[35,48,49] TC was first

applied at 2 mg/mL in Milli-Q water (final concentration 0.225 mM in cell suspensions) and

incubated for 10 minutes.[35] NR stock (10 mg/mL in DMSO) was added to 1 mL of cell

suspension at a final concentration of 0.05 mg/mL and incubated for 10 minutes.[50] Lastly, 100

× SG was added to the cells and incubated for 4 to 12 hours (Figure S3, step 7).[44] All incubation

steps were performed at room temperature in the dark.

2.5   Soil disaggregation

Significant challenges were faced when applying the FACS step of the TriFlow-Seq workflow to

soil samples, which differ markedly from activated sludge in their physicochemical properties

and tendency to reaggregate. Traditional soil pretreatments using detergents or sonication for

quantification have proven inadequate for sorting low-abundance populations,[51,52] especially

when high cell numbers are required for quality genomic reads. Key challenges include

reaggregation during extended suspension, which reduces sorting efficiency and increases false

positives,[53] and soil heterogeneity, which affects reaggregation rates due to variations in cation

exchange capacity, plant and microbial protein activity, soil organic matter content, and biofilm

formation.[54] This re-aggregation can reduce sorting efficiency and increase the risk of nozzle

clogging, potentially contaminating ongoing cell sorting.[52] To address this, a gentle enzymatic

digestion using Trypsin-EDTA was applied after staining to degrade residual and persistent

extracellular organic materials in microbial pellets, thereby further preventing re-aggregation

during FACS.[55] The protocol involved firstly pelleting the stained sample by centrifugation after

incubation with SG and decanting the supernatant. The stained pellets were then resuspended in

3 mL of $1 \times$ Trypsin-0.5 mM EDTA and incubated for 1 min at 37°C in a water bath, followed

by trypsin deactivation using $1 \times$ FBS in a 1:1 ratio. Prior to FACS, trypsin treated samples were

diluted in $1 \times$ PBS to achieve a proper threshold rate of 1500 events s^{-1} (Figure S3, step 8).

Notably, this additional pretreatment step achieved sorting efficiencies ranging from 77-92%

with an average of 85.5% (Table S2) across the tested soil series, representing a substantial

improvement of approximately 50% higher compared to the 35.75% average sorting efficiency

obtained using activated sludge sample pretreatment (n = 8 samples, Table S2).

2.6   FACS instrument settings

Cell sorting was carried out using the 100-micron nozzle on a Fusion Aria III cell sorter (BD,

U.S.) equipped with a blue laser (488 nm excitation) with a 530 nm band-pass (BP) filter for SG,

a violet laser (405 nm excitation) with a 525 BP filter for TC, and a yellow-green laser (561 nm

excitation) with a 610 BP filter for NR. The blue laser was also used to analyze forward scatter

(FSC) and side scatter (SSC), corresponding to cell size and cell granularity, respectively.

Manual gating was performed as summarized in the supplementary material (Text S2, Figure S3).

Bacterial cells were sorted in purity mode at a rate below 1,500 events/second. To obtain

sufficient DNA for downstream analyses, up to 10□ cells were sorted into sterile 1.5 mL

microtubes and stored at −80°C for further analysis.[56]

2.7   FACS gating validation

A post-sort purity check was conducted to confirm single-cell isolation and targeted sorting of

PHA-PAOs. A 100 µl aliquot of sorted sample was diluted in $1 \times$ PBS and re-analyzed.

Approximately 89% of events fell within the FSC-H/FSC-A gate, indicating successful recovery

of single cells, while around 57% remained within the PHA-PAO gate—consistent with

expectations due to photobleaching during laser exposure.

To ensure the reliability and accuracy of the proposed TriFlow-Seq method in identifying

and quantifying single cell PHA-PAOs in complex microbial communities, we employed two

independent validation methods. First, we verified PHA-PAO enrichment via fluorescent

confocal microscopy. Approximately 4,000 sorted cells were collected to validate that all sorted

cells were positive for SG, TC, and NR. Unsorted cells (as a control) and sorted cells were first

filtered through a black polycarbonate filter (0.22 µm pore size) to concentrate the cells. The

filters were placed onto glass slides with a drop of antifade reagent. Validation of the gating

methods was carried out using confocal microscopy (Carl Zeiss) at 800X magnification with

three excitation/filter sets: blue light (bandpass (BP) 450–490 nm excitation filter, dichroic

beamsplitter (FT) at 500 nm, and longpass (LP) 515 nm emission filter), green light (BP 510–

560 nm, FT 565 nm, LP 590 nm), and ultraviolet (BP 340–380 nm, FT 400 nm, BP 435–485).

Quantification of TC- and NR-stained cells relative to SG-stained cells was performed using

Daime (version 1.2).[57]

2.8   DNA extraction and amplicon sequencing

Genomic DNA from sorted soil and sludge cells was extracted using the InstaGene Matrix (Bio-

Rad Laboratories, USA). Genomic DNA from unsorted raw soil and sludge was extracted using

the DNeasy PowerSoil Pro Kit (Qiagen, Valencia, CA, USA) on the QIAcube® (QIAGEN)

platform. The V4 region of the 16S rRNA genes from both sorted and unsorted samples was

amplified and sequenced using primers 515F and 806R at the University of Connecticut-

Microbial Analysis, Resources, and Services (UCONN-MARS) facility. Details of the

sequencing and analysis processes are listed in Text S3.

2.9   Classification of Detected Genera by Polyphosphate and PHA Accumulation

Potential

For each detected genus, PAO-related functionality was assessed through literature review and

genomic analysis. A color-coding system was applied: red stars indicated genera with

experimentally confirmed anaerobic phosphorus release and aerobic phosphorus uptake; yellow

stars denoted genera with documented polyP and PHA accumulation but unclear PAO activity;

green stars represented genera with genetic potential for PHA-PAO metabolism based on

functional gene presence in representative genomes from NCBI and BioCyc databases; blue stars

were assigned when taxonomic resolution was limited to family or order level, with functional

inference based on constituent genera. This color code system was applied in Figure 3 and 4, and

Table S3 and S4. Detailed methodology is provided in Text S4.

2.10 Single cell Raman spectroscopy (SCRS)

As a second validation, we compared FACS results with SCRS analysis, an independent method

previously developed by our group to quantify polyP- and PHA-containing cells.[36,37] Details of

SCRS analysis for PHA-PAO identification, polyP and PHA detection can be referred to our

previous publications.[58] Key details are provided in Text S5.

2.11 Statistical analysis

Spearman and Pearson correlation coefficients and corresponding *p*-values were computed using

the R *stats* (4.1.3) package. Principal coordinates analysis (PCoA) was performed using the R

vegan (2.6-4) package to assess beta diversity based on Bray-Curtis dissimilarities of normalized

relative abundance data. The ordination plot was visualized using the R *ggplot2* (3.4.4) package.

For the differential abundance analysis, log□□-transformed fold changes were compared

between genera using two-tailed independent t-tests implemented in Python (version 3.12) with

SciPy (version 1.10.1). The resulting t-statistics were visualized as a heatmap using the Seaborn

package (version 0.13.2).

3   Results and discussion

3.1   Evaluation of TriFlow for sorting PHA-containing PAOs

The effectiveness of TriFlow for identifying and sorting PHA-containing PAOs was tested with

sludge samples from EBPR and S2EBPR systems, which are known to be enriched in PHA-

PAOs and served as positive environmental controls. The density of the triple-stained cells

before and after the TriFlow enrichment was visualized using confocal microscopy (Figure 1A).

A significant and evident enrichment of candidate PHA-PAO cells containing both polyP and

PHA was observed in the sorted samples (magenta) compared to the original sludge, a pattern

validated in all four tested EBPR systems. In addition, the relative abundances of PHA-positive

and polyP-positive cells to total cells (SG-stained cells) were semi-quantified using digital image

analysis (Figure 1B). NR-stained and TC-stained cells were clearly enriched in all FACS sorted

samples compared to the original sludge samples. These results demonstrate that our gating

strategy in the FACS analysis effectively enriched dual polyP- and PHA-accumulating PAOs as

expected, confirming the appropriateness of the gating strategy employed.

A)

Site	Unsorted cells	Sorted cells	Site	Unsorted cells	Sorted cells
A	a	b	C	e	f
B	c	d	D	g	h

ᵃGreen, DNA-positive cells; Red, PHA-positive cells; Blue, polyP-positive cells; Magenta, polyP and PHA-positive cells

B)

Sample	Unsorted cells (%)		Sorted cells (%)	
	NR+ (PHA-positive)	TC+ (polyP-positive)	NR+ (PHA-positive)	TC+ (polyP-positive)
A	50.3	35.6	99	98.3
B	45.7	53.7	97.3	87.7
C	68.1	29.1	79.8	80.9
D	19	6.8	96	83.6

Figure 1 A) Confocal microscopy images comparing FACS sorted stain-positive cells to unsorted cells in activated

sludge. Samples were taken from three EBPR systems (A, B, and C), and a S2EBPR system (D). B) Relative

abundances of NR (PHA)-positive and TC (polyP)-positive cells in sorted samples versus original unsorted samples,

as determined by semi-quantitative confocal microscopic image analysis.

3.2   Validation of PHA-PAOs quantification using TriFlow

To Further validate the TriFlow method, we compared the relative abundance of PHA-PAOs

identified through FACS with those determined by SCRS across various microbial samples,

including both EBPR systems biomass and the maize rhizosphere soil samples (Figure 2). SCRS,

which targets the molecular composition of microbial cells, has demonstrated exceptional

sensitivity and accuracy in detecting and quantifying intracellular polymers such as polyP, PHA,

and glycogen.[38] SCRS-based detection of PHA-PAOs containing both polyP and PHA has been

demonstrated in previous studies[27,36–39] Importantly, the percentage of PHA-PAOs determined by

FACS and SCRS fell within the same range. A significant positive monotonic relationship was

observed between the two techniques (Spearman's $\rho = 0.62, p = 0.009$), demonstrating

statistically robust concordance in detecting functional PHA-PAOs by these two independent

methods.

Figure 2 Spearman's rank correlation coefficients of PHA-PAOs detected by FACS and SCRS. PHA-PAOs are

defined as PHA-PAO cells containing both polyP and PHA. The strong correlation between the two independent

techniques validates the effectiveness of TriFlow in detecting and quantifying PHA-PAOs in complex microbial

communities. AS denotes activated sludge from EBPR and S2EBPR systems.

Although the percentages of PHA-PAOs determined by TriFlow were consistent and

correlated with those identified by SCRS, discrepancies occurred due to inherent differences in

detection principles and sample processing protocols. SCRS is a non-invasive, fixation-free

method that detects intracellular PHA and polyP based on their unique Raman spectral signatures

and has proven effective in detecting multiple intracellular polymers with relatively less

influence from varying polyP composition (i.e., counter metal ions).[59,60] In contrast, FACS

requires cell fixation and permeation steps to enhance dye penetration, which permanently alters

cell wall and membrane permeability.[47] This modification typically results in a 10-20% loss of

intracellular polymers, similar to observations with fluorescence *in situ* hybridization (FISH)

fixatives.[39] In addition, the accuracy of FACS largely relies on the specificity and sensitivity of

fluorescent dyes. Nile red, although widely employed for PHA detection in flow cytometry, can

also bind to other lipophilic inclusions such as trehalose and triglycerides.[61] Similarly,

tetracycline detects polyP by preferentially complexing with countercharged divalent cations

(e.g., Ca^{2+}, Mg^{2+}) in polyP granules.[35,62,63] These methodological differences suggest that

FACS may slightly overestimate PHA-containing PHA-PAOs relative to the SCRS method,

consequently reducing the slope of the linear curve. Furthermore, FACS employs size-based

filtration (40 μm cell strainer) to prevent flow cytometer nozzle clogging, while SCRS utilizes

density-gradient centrifugation for cell extraction.[6460] These procedural variations likely

influence the organisms being analyzed, contributing to the observed differences in PHA-PAO

abundance percentages. Despite these discrepancies, the consistent trend in PHA-PAO detection

by both methods demonstrates the viability of the TriFlow for identifying and quantifying PHA-

PAOs in complex microbial communities.

3.3 Application of TriFlow-Seq for the identification and quantification of PHA-PAOs in

EBPR systems

Combining TriFlow with downstream 16S rRNA gene amplicon sequencing (termed TriFlow-

Seq) enabled the enrichment, quantification, and identification of both known and candidate

PHA-PAOs in EBPR systems. The relative abundances of canonical PHA-PAOs in the original

sludge prior to cell sorting and after TriFlow enrichment in pilot- and full-scale EBPR plants are

depicted in Figure 3.

388
Figure 3 A) Relative abundances of known candidate PHA-PAOs before (unsort) and after (sort) TriFlow-seq in

three full-scale EBPR plants and one pilot-scale S2EBPR plant. Groups with relative abudances below 0.1% are not

shown; B) Differential abundance heatmap of bacterial genera before and after TriFlow-seq sorting. The heat map

represents t-statistics derived from two-tailed t-tests on log_{10}-transformed fold changes in relative abundance of

bacterial genera after TriFlow sorting in comprison to prior-sorting. Blue hues denote positive t-statistics

(enrichment via sorting), while red hues indicate negative t-statistics (depletion after sorting). Statistical significance

levels: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ The red pentagram indicates the genus has been experimentally

confirmed with PHA/polyP and PHA-PAO activities; the yellow pentagram indicates the presence of polyP and

PHA; the green pentagram indicates that the reference genome contains necessary genes for polyP and carbon

storage synthesis; the blue pentagram indicates that some genera within the Family have polyP accumulation ability.

Detailed information can be found in Table S3. Analysis is limited to the top 50 genera by abundance.

3.3.1 Enrichment of known PHA-PAOs in EBPR systems using TriFlow-Seq

Across all sampled wastewater treatment plants, TriFlow-Seq consistently detected well-

established PHA-PAO genera, including *Pseudomonas*, *Dechloromonas*, and *Ca.*

Accumulibacter (Figure 3A). These genera are recognized for their dual accumulation of polyP

and PHA in EBPR systems,[65–68] validating TriFlow-Seq's ability to target functionally active

PHA-PAOs.

To further characterize the sorted communities, we calculated enrichment ratios, defined

as the $\log_{10}$-transformed ratio of post-sorting to pre-sorting relative abundances for each genus.

Higher ratios indicate a greater enrichment of bacteria capable of accumulating both polyP and

intracellular C polymers, thus reflecting the proportion of putative PHA-PAOs within that

taxonomic group (Figure 3B). It should be noted that even within known PHA-PAOs, some sub-

genus populations were not actively performing polyP and PHA accumulation, resulting in a

negative enrichment factor, as observed for *Ca*. Accumulibacter. Despite being consistently

identified as the dominant PHA-PAO in raw sludge samples from the three full-scale plants

(except for the S2EBPR pilot plant, with 1.2~4.0%) (Figure 3A), *Ca*. Accumulibacter showed

varied enrichment ratios among the EBPR plants (Figure 3B). Its relative abundance even

decreased after sorting in sample C. Interestingly, *Tetrasphaera* emerged as the most

significantly enriched genus after sorting (Figure 3A and 3B). Although initially thought to lack

PHA synthesis capability, recent studies have revealed that anaerobic PHA synthesis depends on

clade composition of *Tetrasphaera*.[11–14] For instance, the *T. japonica* genome possesses PHA

synthase (*phaC*) and has been experimentally shown to synthesize PHA under anaerobic

conditions.[11] This evidence suggests that *T. japonica*, or potentially uncultured *Tetrasphaera*

with Accumulibacter-like PAO metabolism, resides in full-scale EBPR systems.[69] Overall, the

relative abundances of known PHA-PAOs accounted for only 2 to 12% of total PHA+polyP-

positive cells (Figure 3A), highlighting the greater phylogenetic diversity of PHA-PAOs than our

current understanding.

3.3.2   Discovery of candidate PHA-PAOs in EBPR systems

Based on TriFlow results, we proposed putative PHA-PAOs in EBPRs based on (1) consistent

detection across sorted communities via TripleFlow-seq; (2) genomic potential for polyP and

PHA accumulation within the detected genus. A diverse putative PHA-PAOs, including

unclassified Rhodobacteraceae, unclassified Rhizobiales, unclassified Caldilineaceae, *Lautropia*,

and others (Figure 3B). Among them, a substantial portion lacked genus-level classification,

suggesting the presence of uncharacterized genera that may possess PHA-PAO-like functions.

Although these families or orders have not traditionally been associated with PHA-PAO activity,

related members within the same taxonomic groups have been reported to accumulate

intracellular polyP and PHA based on genomics analysis. For instance, some bacteria isolated

from the order Rhizobiales were found to contain PHA and polyP.[70] An uncultured genus close

to the Rhodobacteraceae family is speculated to be sulfur PHA-PAOs.[71] These findings provide

preliminary evidence for novel candidate PHA-PAOs, though further studies are needed to

determine whether they contribute to phosphorus removal through classical or alternative EBPR

pathways.

These results revealed important insights into PHA-PAO diversity and functionality in EBPR

systems. Not all genetically identified *Ca*. Accumulibacter displayed metabolically active PAO

phenotypes, as evidenced by their varied enrichment ratios, highlighting the distinction between

genetic potential and phenotypic expression. *Tetrasphaera* showed unexpected and significant

enrichment with variable intracellular PHA content, revealing phenotypic diversity beyond the

current understanding of this genus. Additionally, the TriFlow-Seq method uncovered novel

candidate PHA-PAOs from previously unrecognized taxonomic groups, expanding our

understanding of PHA-PAO diversity in EBPR systems. Although further investigation and

confirmation of the roles of these candidate PHA-PAOs in the tested plants is beyond the focus

of this study, this application has demonstrated that TriFlow-Seq is a valuable tool for rapidly

enriching and identifying targeted bacterial populations that exhibit the phenotypic traits

characteristic of functionally relevant PHA-PAOs. This approach was further adapted and

optimized to be applicable to more heterogeneous sample matrices, such as bulk and rhizosphere

soil, enabling the identification and characterization of PHA-PAOs in these complex

environmental matrices.

3.4 Discovery of the identities of previously unknown rhizosphere PHA-PAOs
communities

3.4.1 Effectiveness in application of TriFlow-Seq in rhizosphere samples

Application of targeted single-cell FACS for PHA-PAO identification in soil samples has not

been previously reported. Our newly developed TriFlow-Seq pretreatment method and sequential

targeted gating successfully identified and isolated PHA-PAO populations from the rhizosphere

microbiome. Genomic DNA recovery ranged from 1 ng/µL to 16.3 ng/µL, with high-quality

(Phred >30) 16S rRNA gene amplicon reads obtained in all but one sequenced sorted sample (n

$= 192$), demonstrating the method's effectiveness across diverse soil types.

Successful application and recovery of the 16S rRNA gene amplicon enabled taxonomic

identification of the previously unknown PHA-PAOs community, revealed to include the

unanimous presence of canonical PHA-PAOs associated EBPR systems and performance.

Proteobacteria, Actinobacteria, and Acidobacteria were the most abundant phyla in both unsorted

and sorted communities (63.2-66.2% and 59.1-59.5% respectively) (Figure 4A), aligning with

the typically observed community compositions in both maize rhizosphere and BEPR

communities.[72] Notably, the phylum Myxococcota, known for complex social behaviors and

recently identified to harbor *ppk1* associated with polyP accumulation in activated sludge,[73,74]

was consistently detected only within the sorted communities (6.58-8.54%) (Figure 4A). At the

genus level, while typical PHA-PAOs such as *Tetrasphaera*, *Ca.* Accumulibacter, and

Dechloromonas were detected within the sorted community, their combined abundance was

below 1%, suggesting minimal contribution to the rhizosphere processes. Instead, *Pseudomonas*

(9.03-12.7%) and *Halomonas* (11-15.6%), both putative PHA-PAOs identified in EBPR systems,

were the predominant genera in the sorted communities (Figure 4B). Further, of the

conventionally known PHA-PAOs, including *Ca. Accumulibacter*, *Tetrasphaera*,

Dechloromonas, *Acinetobacter*, *Gemmatimonas*, and *Pseudomonas*. there was no significant loss

of unrarefied observed species richness within the *Tetrasphaera* and *Acinetobacter* genera

between the sorted and unsorted communities indicating the scope and accuracy of our TriFlow-

Seq method in surveying and conserving the active PHA-PAO diversity in rhizosphere samples;

capable of identifying functionally verified PHA-PAOs despite naturally low abundances (Figure

S7).

3.4.2   Discovery of novel PHA-PAOs in the maize rhizosphere

In addition to the discovery of canonical PHA-PAO activity in the rhizosphere, for the first time

we were able to capture a markedly different co-polymer accumulating community than those

reported in EBPR systems and propose novel candidate PHA-PAOs functionally active in

rhizosphere systems. The enrichment profiles of key sorted taxa across different maize genotypes

were examined to identify candidate functional PHA-PAOs in the maize rhizosphere. The

selection criteria used were comparable to those established for activated sludge systems in

Section 3.3.1 but adapted to consider the inherent variability of soil systems which leads to

differences in phenotypic profiles and diverse metabolic activities among potential PHA-PAOs.

The enrichment profiles at the sub-genus level revealed the selective dominance of

specific amplicon sequence variants (ASVs) within the predominate sorted genera, leading to the

identification of three candidate rhizosphere PHA-PAOs. Among these, two predominant

putative PHA-PAO ASVs exhibited positive enrichment across all maize lines: *Halomonas*

ASV_14 (21.1-22.1-fold) and *Pseudomonas* ASV_51 (20.7-22-fold) (Figure 4D). Additionally,

Nannocystis ASV_54, within the Myxococcota phylum, most closely related to *Nannocystis*

exedens (93.68% 16S rRNA sequence similarity, BLASTN)*,* exhibited positive enrichment

(1.14-4.46-fold) across 68.75% of samples, specifically in sampled inbreds (including B73) and

most maternal B73 hybrids (p-value <0.05) (Figure 4D). Analysis of the available annotated

genome of *N. exedens* (GenBank GCA_002343915.1) revealed the presence of *ppk1*, *ppx*, and

sterol and steroid biosynthetic pathways.[75] While this study does not aim to exhaustively

compare PHA-PAO populations across diverse maize genotypes, our proof-of-concept analysis

revealed distinct identities of novel PHA-PAO with distinct key C and P functions and their

unique associations with different maize genotypes, warranting further investigation.

3.4.3 Exploration of rhizosphere PHA-PAO community dynamics

In addition to revealing the PHA-PAO community's taxonomic identity, we first investigated

community dynamics across maize host genetic diversity, an emerging driver in rhizosphere

structure.[72,76] Principal coordinate analysis (PCoA) based on Bray-Curtis dissimilarity revealed

distinct clustering and significant separation between the unsorted and sorted community along

PCo1 (35.4% variance; PERMANOVA, p-value <0.001), indicating isolation of a distinct sub-

community (Figure 4C). Within both clusters, two significant sub-clusters emerged based on

maize genetic background (inbred versus hybrid) along PCo2 (11.1% variation, p-value <0.001,

PERMANOVA) (Figure 4C), suggesting that the sorted population retains its community

fingerprint from its original rhizosphere population. Alpha diversity was significantly reduced in

sorted samples across all metrics, including species richness, Shannon index, and Simpson's

index (p < 0.001; Figure S10), confirming that TriFlow-Seq enriched a specialized sub-

community dominated by a few taxa.

523
Figure 4 of candidate rhizosphere PHA-PAOs and characterization of the rhizosphere PHA-PAO community via

TriFlow-Seq. A) Phylum- and B) Genus-level community composition and relative abundance of unsorted raw

rhizosphere community and sorted PHA-PAO community, separated by field experiment, abundance filtered > 0.03;

B) Principal coordinate analysis (PCoA) based on Bray-Curtis dissimilarity of raw rhizosphere community and

FACS-sorted PHA-PAO community across maize inbred and hybrid lines; D) Differential abundance analysis of

candidate rhizosphere PHA-PAO taxa in inbred (e.g., B73) and hybrid (e.g., B73×Mo17) rhizospheres following

TriFlow-seq. The blue scale denotes positive enrichment in the PHA-PAO community, and the red scale denotes

negative enrichment. Significance is based on a two-tailed t-test with an unscaled $\log_{10}$-fold change: *$p < 0.05$, **p

< 0.01, ***$p < 0.001$. The red pentagram indicates the genus has been experimentally confirmed with PHA/polyP

and PHA-PAO activities; the yellow pentagram indicates the presence of polyP and PHA; the green pentagram

indicates that the reference genome contains necessary genes for polyP and carbon storage synthesis; the blue

pentagram indicates that some genera within the Family have polyP accumulation ability. Detailed information can

be found in Table S4. Analysis was limited to the top 50 amplicon sequence variants (ASVs) by abundance.

These results, for the first time, revealed the characteristics, diversity, composition, and

distribution pattern of the rhizosphere PHA-PAO community. Emerging evidence highlighted

the important roles of PHA-PAOs in rhizosphere communities and processes, including

halotolerance,   phytohormone   production,   siderophore   production,   beneficial

antagonism/antibiotic production, and nutrient cycling.[77–79] Our TriFlow-Seq method enabled the

discovery of new PHA-PAOs in agricultural soils, expanding our understanding of their identity

and diversity within agricultural systems.

4   TriFlow-Seq: Achievements, Applications, and Future Prospects

Linking phenotype-based detection and genotype-based characterization provides valuable

insights into the discovery of organisms of interest. In this study, we developed and validated the

TriFlow-Seq workflow and demonstrated two key applications: identifying functional PHA-

PAOs in wastewater treatment facilities and uncovering previously unrecognized PHA-PAOs

involved in P and C cycling within the maize rhizosphere. By modifying FACS preprocessing

protocols, we enabled the identification of candidate PHA-PAOs in modified environments, such

as soil matrix. Our findings revealed unexpectedly high PHA-PAO prevalence and distinct

phylogenetic patterns across different maize genotypes, establishing a pioneering approach to

exploring PHA-PAO identities and functions in rhizosphere systems.

Future applications of Reflow-Seq can be enhanced through several approaches: (1)

incorporating additional phenotyping targets as new staining materials become available,

potentially revealing greater PHA-PAO diversity; (2) integrating mini-metagenomics or other

omics techniques to obtain deeper functional and metabolic insights into candidate PHA-PAOs;

and (3) combining with FISH to increase specificity for targeting particular microbial groups.

During our investigation, tetracycline and DAPI were the only fluorescent dyes proven effective

for FACS-based PHA-PAO selection.[34,35] However, recent advancements have introduced more

specific dyes, such as JC-D7, which offer improved specificity for polyP detection in

environmental samples.[80] Incorporating such novel dyes into the TriFlow-Seq workflow could

enhance the accuracy and sensitivity of PHA-PAO identification in future research. Despite

current limitations, this technique demonstrates significant potential for identifying novel PHA-

PAOs within complex microbial consortia, advancing our understanding of microbial polyP

metabolism and phosphorus cycling across diverse ecosystems.

5   Acknowledgements

This study was partially supported by the Cornell Institute for Digital Agriculture (CIDA), and

the Center for Research on Programmable Plant System (CROPPS) funded by the National

Science Foundation (Grant No. DBI-2019674). We thank Dr. Lydia Tesfa, Jaclyn Mahoney, and

Michael Sledziona from the Flow Cytometry Facility of the Biotechnology Resource Center at

Cornell University (RRID:SCR_021740) for their technical assistance with FACS experiments.

The authors would like to thank the G2F consortium for providing the High-Intensity

Phenotyping Site (HIPS) experiment, which consisted of hybrids that were used in this study.

This consortium involves more than 30 researchers representing more than 20 research

institutions. Details about the initiative and publicly available resources can be found at

www.Genomes2Fields.org. We thank Dr. Maria Cinta Romay and Nicholas Lepak for their

generous time and effort in coordinating and planting the HIPS experiment.

References

(1)   Mason-Jones, K.; Robinson, S. L.; Veen, G. F. (Ciska); Manzoni, S.; van der Putten, W. H.

582   Microbial Storage and Its Implications for Soil Ecology. *ISME J* **2022**, *16* (3), 617–629.

583   https://doi.org/10.1038/s41396-021-01110-w.

584   (2)   Rao, N. N.; Gómez-García, M. R.; Kornberg, A. Inorganic Polyphosphate: Essential for

585   Growth and Survival. *Annual Review of Biochemistry* **2009**, *78* (Volume 78, 2009), 605–

586   647. https://doi.org/10.1146/annurev.biochem.77.083007.093039.

587   (3)   Kornberg, A.; Rao, N. N.; Ault-Riché, D. Inorganic Polyphosphate: A Molecule of Many

588   Functions.   *Annual   Review   of   Biochemistry*   **1999**,   *68*   (1),   89–125.

589   https://doi.org/10.1146/annurev.biochem.68.1.89.

590   (4)   Oehmen, A.; Lemos, P. C.; Carvalho, G.; Yuan, Z.; Keller, J.; Blackall, L. L.; Reis, M. A.

591   M. Advances in Enhanced Biological Phosphorus Removal: From Micro to Macro Scale.

592   *Water Research* **2007**, *41* (11), 2271–2300. https://doi.org/10.1016/j.watres.2007.02.030.

593   (5)   Seviour, R. J.; Mino, T.; Onuki, M. The Microbiology of Biological Phosphorus Removal

594   in Activated Sludge Systems. *FEMS Microbiology Reviews* **2003**, *27* (1), 99–127.

595   https://doi.org/10.1016/S0168-6445(03)00021-4.

596   (6)   Martín, H. G.; Ivanova, N.; Kunin, V.; Warnecke, F.; Barry, K. W.; McHardy, A. C.;

597   Yeates, C.; He, S.; Salamov, A. A.; Szeto, E.; Dalin, E.; Putnam, N. H.; Shapiro, H. J.;

598   Pangilinan, J. L.; Rigoutsos, I.; Kyrpides, N. C.; Blackall, L. L.; McMahon, K. D.;

599   Hugenholtz, P. Metagenomic Analysis of Two Enhanced Biological Phosphorus Removal

600   (EBPR)   Sludge   Communities.   *Nat   Biotechnol*   **2006**,   *24*   (10),   1263–1269.

601   https://doi.org/10/dct6rr.

(7) Petriglieri, F.; Singleton, C.; Peces, M.; Petersen, J. F.; Nierychlo, M.; Nielsen, P. H.

"Candidatus Dechloromonas Phosphoritropha" and "Ca. D. Phosphorivorans", Novel

Polyphosphate Accumulating Organisms Abundant in Wastewater Treatment Systems.

ISME J **2021**, *15* (12), 3605–3614. https://doi.org/10.1038/s41396-021-01029-2.

(8) Li, G.; Tooker, N. B.; Wang, D.; Srinivasan, V.; Barnard, J. L.; Russell, A.; Stinson, B.;

McQuarrie, J.; Schauer, P.; Menniti, A.; Varga, E.; Hauduc, H.; Takács, I.; Bott, C.;

Dobrowski, P.; Onnis-Hayden, A.; Gu, A. Z. Modeling Versatile and Dynamic Anaerobic

Metabolism for PAOs/GAOs Competition Using Agent-Based Model and Verification via

Single Cell Raman Micro-Spectroscopy. *Water Research* **2023**, *245*, 120540.

https://doi.org/10/gs5c3w.

(9) Zhao, W.; Bi, X.; Peng, Y.; Bai, M. Research Advances of the Phosphorus-Accumulating

Organisms of Candidatus Accumulibacter, Dechloromonas and Tetrasphaera: Metabolic

Mechanisms, Applications and Influencing Factors. *Chemosphere* **2022**, *307*, 135675.

https://doi.org/10/gq8m45.

(10) Tobin, K. M.; McGrath, J. W.; Mullan, A.; Quinn, J. P.; O'Connor, K. E. Polyphosphate

Accumulation by Pseudomonas Putida CA-3 and Other Medium-Chain-Length

Polyhydroxyalkanoate-Accumulating Bacteria under Aerobic Growth Conditions. *Applied*

and Environmental Microbiology **2007**, *73* (4), 1383–1387.

https://doi.org/10.1128/AEM.02007-06.

(11) Kristiansen, R.; Nguyen, H. T. T.; Saunders, A. M.; Nielsen, J. L.; Wimmer, R.; Le, V. Q.;

McIlroy, S. J.; Petrovski, S.; Seviour, R. J.; Calteau, A.; Nielsen, K. L.; Nielsen, P. H. A

Metabolic Model for Members of the Genus Tetrasphaera Involved in Enhanced Biological

Phosphorus Removal. *ISME J* **2013**, *7* (3), 543–554. https://doi.org/10.1038/ismej.2012.136.

(12) Close, K.; Marques, R.; Carvalho, V. C. F.; Freitas, E. B.; Reis, M. A. M.; Carvalho, G.;

Oehmen, A. The Storage Compounds Associated with Tetrasphaera PAO Metabolism and

the Relationship between Diversity and P Removal. *Water Research* **2021**, *204*, 117621.

https://doi.org/10/gsmsnj.

(13) Marques, R.; Santos, J.; Nguyen, H.; Carvalho, G.; Noronha, J. P.; Nielsen, P. H.; Reis, M.

A. M.; Oehmen, A. Metabolism and Ecological Niche of Tetrasphaera and Ca.

Accumulibacter in Enhanced Biological Phosphorus Removal. *Water Research* **2017**, *122*,

159–171. https://doi.org/10.1016/j.watres.2017.04.072.

(14) McIlroy, S. J.; Onetto, C. A.; McIlroy, B.; Herbst, F. A.; Dueholm, M. S.; Kirkegaard, R.

H.; Fernando, E.; Karst, S. M.; Nierychlo, M.; Kristensen, J. M.; Eales, K. L.; Grbin, P. R.;

Wimmer, R.; Nielsen, P. H. Genomic and in Situ Analyses Reveal the Micropruina Spp. as

Abundant Fermentative Glycogen Accumulating Organisms in Enhanced Biological

Phosphorus Removal Systems. *Frontiers in Microbiology* **2018**, *9* (MAY), 1–12.

https://doi.org/10.3389/fmicb.2018.01004.

(15) Akbari, A.; Wang, Z.; He, P.; Wang, D.; Lee, J.; Han, I.; Li, G.; Gu, A. Z. Unrevealed

Roles of Polyphosphate☐accumulating Microorganisms. *Microb Biotechnol* **2021**, *14* (1),

82–87. https://doi.org/10.1111/1751-7915.13730.

(16) Cordell, D.; Rosemarin, A.; Schröder, J. J.; Smit, A. L. Towards Global Phosphorus

Security: A Systems Framework for Phosphorus Recovery and Reuse Options.

Chemosphere **2011**, *84* (6), 747–758. https://doi.org/10/d4mvgw.

(17) Zabel, F.; Putzenlechner, B.; Mauser, W. Global Agricultural Land Resources–a High

Resolution Suitability Evaluation and Its Perspectives until 2100 under Climate Change

Conditions. *PloS one* **2014**, *9* (9), e107522.

(18) Saia, S. M.; Carrick, H. J.; Buda, A. R.; Regan, J. M.; Walter, M. T. Critical Review of

Polyphosphate and Polyphosphate Accumulating Organisms for Agricultural Water Quality

Management. *Environ. Sci. Technol.* **2021**, *55* (5), 2722–2742.

https://doi.org/10.1021/acs.est.0c03566.

(19) Mason-Jones, K.; Banfield, C. C.; Dippold, M. A. Compound-Specific 13C Stable Isotope

Probing Confirms Synthesis of Polyhydroxybutyrate by Soil Bacteria. *Rapid*

Communications in Mass Spectrometry **2019**, *33* (8), 795–802. https://doi.org/10/gr8ctq.

(20) Mason-Jones, K.; Breidenbach, A.; Dyckmans, J.; Banfield, C. C.; Dippold, M. A.

Intracellular Carbon Storage by Microorganisms Is an Overlooked Pathway of Biomass

Growth. *Nat Commun* **2023**, *14* (1), 2240. https://doi.org/10/gr745w.

(21) Butler, O. M.; Manzoni, S.; Warren, C. R. Community Composition and Physiological

Plasticity Control Microbial Carbon Storage across Natural and Experimental Soil Fertility

Gradients. *ISME J* **2023**, *17* (12), 2259–2269. https://doi.org/10.1038/s41396-023-01527-5.

(22) Ratcliff, W. C.; Kadam, S. V.; Denison, R. F. Poly-3-Hydroxybutyrate (PHB) Supports

Survival and Reproduction in Starving Rhizobia: PHB Increases Rhizobium Fitness. *FEMS*

Microbiology Ecology **2008**, *65* (3), 391–399. https://doi.org/10.1111/j.1574-

6941.2008.00544.x.

(23) Smith, S. E.; Smith, F. A. Roles of Arbuscular Mycorrhizas in Plant Nutrition and Growth:

New Paradigms from Cellular to Ecosystem Scales. *Annual Review of Plant Biology* **2011**,

62 (Volume 62, 2011), 227–250. https://doi.org/10.1146/annurev-arplant-042110-103846.

(24) Rao, N.; Kornberg, A. Inorganic Polyphosphate Regulates Responses of Escherichia Coli to

Nutritional Stringencies, Environmental Stresses and Survival in the Stationary Phase.

Inorganic polyphosphates: Biochemistry, biology, biotechnology **1999**, 183–195.

(25) Gray, M. J.; Jakob, U. Oxidative Stress Protection by Polyphosphate — New Roles for

an Old Player. *Current Opinion in Microbiology* **2015**, *24*, 1–6.

https://doi.org/10.1016/j.mib.2014.12.004.

(26) Alam, M. M.; Srinivasan, V.; Mueller, A. V.; Gu, A. Z. Status and Advances in

Technologies for Phosphorus Species Detection and Characterization in Natural

Environment- A Comprehensive Review. *Talanta* **2021**, *233*, 122458.

https://doi.org/10.1016/j.talanta.2021.122458.

(27) Li, Y.; Cope, H. A.; Rahman, S. M.; Li, G.; Nielsen, P. H.; Elfick, A.; Gu, A. Z. Toward

Better Understanding of EBPR Systems via Linking Raman-Based Phenotypic Profiling

with Phylogenetic Diversity. *Environmental Science and Technology* **2018**, *52* (15), 8596–

8606. https://doi.org/10.1021/acs.est.8b01388.

(28) Majed, N.; Li, Y.; Gu, A. Z. Advances in Techniques for Phosphorus Analysis in Biological

Sources. *Current Opinion in Biotechnology* **2012**, *23* (6), 852–859.

https://doi.org/10.1016/j.copbio.2012.06.002.

(29) He, Z.; Deng, Y.; Van Nostrand, J. D.; Tu, Q.; Xu, M.; Hemme, C. L.; Li, X.; Wu, L.;

Gentry, T. J.; Yin, Y.; Liebich, J.; Hazen, T. C.; Zhou, J. GeoChip 3.0 as a High-

Throughput Tool for Analyzing Microbial Community Composition, Structure and

Functional Activity. *The ISME Journal* **2010**, *4* (9), 1167–1179.

https://doi.org/10.1038/ismej.2010.46.

(30) Tu, Q.; Yu, H.; He, Z.; Deng, Y.; Wu, L.; Van Nostrand, J. D.; Zhou, A.; Voordeckers, J.;

Lee, Y.; Qin, Y. GeoChip 4: A Functional Gene□array□based High□throughput

Environmental Technology for Microbial Community Analysis. *Molecular ecology*

resources **2014**, *14* (5), 914–928.

(31) Kornberg, A. Inorganic Polyphosphate: Toward Making a Forgotten Polymer Unforgettable.

Journal of Bacteriology **1995**, *177* (3), 491–496. https://doi.org/10.1128/jb.177.3.491-

496.1995.

(32) Kulaev, I. S.; Vagabov, V.; Kulakovskaya, T. *The Biochemistry of Inorganic*

Polyphosphates, 2 nd.; John Wiley & Sons, 2005.

(33) Whitehead, M. P.; Eagles, L.; Hooley, P.; Brown, M. R. W. Most Bacteria Synthesize

Polyphosphate by Unknown Mechanisms. *Microbiology* **2014**, *160* (5), 829–831.

https://doi.org/10.1099/mic.0.075366-0.

(34) Terashima, M.; Kamagata, Y.; Kato, S. Rapid Enrichment and Isolation of Polyphosphate-

Accumulating Organisms Through 4'6-Diamidino-2-Phenylindole (DAPI) Staining With

Fluorescence-Activated Cell Sorting (FACS). *Front. Microbiol.* **2020**, *11*.

https://doi.org/10.3389/fmicb.2020.00793.

(35) Günther, S.; Trutnau, M.; Kleinsteuber, S.; Hause, G.; Bley, T.; Röske, I.; Harms, H.;

Müller, S.; Gunther, S.; Kleinsteuber, S.; Trutnau, M.; Bley, T.; Muller, S.; Hause, G.;

Roske, I.; Harms, H.; Gunther, S.; Bley, T.; Muller, S.; Hause, G.; Roske, I.; Harms, H.

Dynamics of Polyphosphate-Accumulating Bacteria in Wastewater Treatment Plant

Microbial Communities Detected via DAPI (4′,6′-Diamidino-2- Phenylindole) and

Tetracycline Labeling. *Applied and Environmental Microbiology* **2009**, *75* (7), 2111–2121.

https://doi.org/10.1128/AEM.01540-08.

(36) Wang, D.; He, P.; Wang, Z.; Li, G.; Majed, N.; Gu, A. Z. Advances in Single Cell Raman

Spectroscopy Technologies for Biological and Environmental Applications. *Current*

Opinion in Biotechnology **2020**, *64*, 218–229. https://doi.org/10.1016/j.copbio.2020.06.011.

(37) Majed, N.; Chernenko, T.; Diem, M.; Gu, A. Z. Identification of Functionally Relevant

Populations in Enhanced Biological Phosphorus Removal Processes Based on Intracellular

Polymers Profiles and Insights into the Metabolic Diversity and Heterogeneity.

Environmental Science and Technology **2012**, *46* (9), 5010–5017.

https://doi.org/10.1021/es300044h.

(38) Majed, N.; Matthäus, C.; Diem, M.; Gu, A. Z. Evaluation of Intracellular Polyphosphate

Dynamics in Enhanced Biological Phosphorus Removal Process Using Raman Microscopy.

Environmental Science and Technology **2009**, *43* (14), 5436–5442.

https://doi.org/10.1021/es900251n.

(39) Fernando, E. Y.; McIlroy, S. J.; Nierychlo, M.; Herbst, F.-A.; Petriglieri, F.; Schmid, M. C.;

Wagner, M.; Nielsen, J. L.; Nielsen, P. H. Resolving the Individual Contribution of Key

Microbial Populations to Enhanced Biological Phosphorus Removal with Raman–FISH.

ISME J **2019**, *13* (8), 1933–1946. https://doi.org/10/gn7s5g.

(40) McMullen, M. D.; Kresovich, S.; Villeda, H. S.; Bradbury, P.; Li, H.; Sun, Q.; Flint-Garcia,

S.; Thornsberry, J.; Acharya, C.; Bottoms, C.; Brown, P.; Browne, C.; Eller, M.; Guill, K.;

Harjes, C.; Kroon, D.; Lepak, N.; Mitchell, S. E.; Peterson, B.; Pressoir, G.; Romero, S.;

Rosas, M. O.; Salvo, S.; Yates, H.; Hanson, M.; Jones, E.; Smith, S.; Glaubitz, J. C.;

Goodman, M.; Ware, D.; Holland, J. B.; Buckler, E. S. Genetic Properties of the Maize

Nested Association Mapping Population. *Science* **2009**, *325* (5941), 737–740.

https://doi.org/10.1126/science.1174320.

(41) Lee, M.; Sharopova, N.; Beavis, W. D.; Grant, D.; Katt, M.; Blair, D.; Hallauer, A.

Expanding the Genetic Map of Maize with the Intermated B73 × Mo17 (IBM) Population.

Plant Mol Biol **2002**, *48* (5), 453–461. https://doi.org/10.1023/A:1014893521186.

(42) Hanway, J. J. Growth Stages of Corn (Zea Mays, L.)1. *Agronomy Journal* **1963**, *55* (5),

487–492. https://doi.org/10.2134/agronj1963.00021962005500050024x.

(43) Eichorst, S. A.; Strasser, F.; Woyke, T.; Schintlmeister, A.; Wagner, M.; Woebken, D.

Advancements in the Application of NanoSIMS and Raman Microspectroscopy to

Investigate the Activity of Microbial Cells in Soils. *FEMS Microbiology Ecology* **2015**, *91*

(10), fiv106. https://doi.org/10.1093/femsec/fiv106.

(44) Deng, L.; Fiskal, A.; Han, X.; Dubois, N.; Bernasconi, S. M.; Lever, M. A. Improving the

Accuracy of Flow Cytometric Quantification of Microbial Populations in Sediments:

Importance of Cell Staining Procedures. *Frontiers in Microbiology* **2019**, *10* (April), 1–13.

https://doi.org/10.3389/fmicb.2019.00720.

(45) Bakken, L. R.; Lindahl, V. Recovery of Bacterial Cells from Soil. In *Nucleic Acids in the*

Environment; Trevors, J. T., van Elsas, J. D., Eds.; Springer: Berlin, Heidelberg, 1995; pp

9–27. https://doi.org/10.1007/978-3-642-79050-8_2.

(46) Hu, X.; Zucherman, K. S. Removing Monolayer Cells from Culture Dishes by Incubation

with EDTA in Studies of Cell Surface Receptors. *Biotechniques* **1996**, *21* (5), 784, 786.

https://doi.org/10.2144/96215bm06.

(47) Günther, S.; Hübschmann, T.; Rudolf, M.; Eschenhagen, M.; Röske, I.; Harms, H.; Müller,

S. Fixation Procedures for Flow Cytometric Analysis of Environmental Bacteria. *Journal of*

Microbiological Methods **2008**, *75* (1), 127–134. https://doi.org/10/bq9dsz.

(48) Cao, J.-S.; Xu, R.-Z.; Luo, J.-Y.; Feng, Q.; Fang, F. Rapid Quantification of Intracellular

Polyhydroxyalkanoates via Fluorescence Techniques: A Critical Review. *Bioresource*

Technology **2022**, *350*, 126906. https://doi.org/10/gstqvm.

(49) Kawaharasaki, M.; Manome, A.; Kanagawa, T.; Nakamura, K. Flow Cytometric Sorting

and RFLP Analysis of Phosphate Accumulating Bacteria in an Enhanced Biological

Phosphorus Removal System. *Water Science and Technology* **2002**, *46* (1–2), 139–144.

https://doi.org/10.2166/wst.2002.0469.

(50) Zuriani, R.; Vigneswari, S.; Azizan, M. N. M.; Majid, M. I. A.; Amirul, A. A. A High

Throughput Nile Red Fluorescence Method for Rapid Quantification of Intracellular

Bacterial Polyhydroxyalkanoates. *Biotechnology and Bioprocess Engineering* **2013**, *18* (3),

472–478. https://doi.org/10.1007/s12257-012-0607-z.

(51) Bressan, M.; Trinsoutrot Gattin, I.; Desaire, S.; Castel, L.; Gangneux, C.; Laval, K. A Rapid

Flow Cytometry Method to Assess Bacterial Abundance in Agricultural Soil. *Applied Soil

Ecology* **2015**, *88*, 60–68. https://doi.org/10.1016/j.apsoil.2014.12.007.

(52) Frossard, A.; Hammes, F.; Gessner, M. O. Flow Cytometric Assessment of Bacterial

Abundance in Soils, Sediments and Sludge. *Front. Microbiol.* **2016**, *7*.

https://doi.org/10.3389/fmicb.2016.00903.

(53) Sutermaster, B. A.; Darling, E. M. Considerations for High-Yield, High-Throughput Cell

Enrichment: Fluorescence versus Magnetic Sorting. *Sci Rep* **2019**, *9* (1), 227.

https://doi.org/10.1038/s41598-018-36698-1.

(54) Ansari, F. A.; Jafri, H.; Ahmad, I.; Abulreesh, H. H. Factors Affecting Biofilm Formation

in in Vitro and in the Rhizosphere. In *Biofilms in Plant and Soil Health*; John Wiley & Sons,

Ltd, 2017; pp 275–290. https://doi.org/10.1002/9781119246329.ch15.

(55) Büks, F.; Kaupenjohann, M. Enzymatic Biofilm Digestion in Soil Aggregates Facilitates

the Release of Particulate Organic Matter by Sonication. *SOIL* **2016**, *2* (4), 499–509.

https://doi.org/10.5194/soil-2-499-2016.

(56) Metz, S.; Dos Santos, A. L.; Berman, M. C.; Bigeard, E.; Licursi, M.; Not, F.; Lara, E.;

Unrein, F.; Lopes dos Santos, A.; Berman, M. C.; Bigeard, E.; Dos Santos, A. L.; Berman,

M. C.; Bigeard, E.; Licursi, M.; Not, F.; Lara, E.; Unrein, F. Diversity of Photosynthetic

Picoeukaryotes in Eutrophic Shallow Lakes as Assessed by Combining Flow Cytometry

Cell-Sorting and High Throughput Sequencing. *FEMS Microbiology Ecology* **2019**, *95* (5),

1–12. https://doi.org/10.1093/femsec/fiz038.

(57) Daims, H. Use of Fluorescence in Situ Hybridization and the Daime Image Analysis

Program for the Cultivation-Independent Quantification of Microorganisms in

Environmental and Medical Samples. *Cold Spring Harbor Protocols* **2009**, *4* (7), 1–8.

https://doi.org/10.1101/pdb.prot5253.

(58) He, P.; Son, Y.; Berkowitz, J.; Li, G.; Lee, J.; Han, I.; Craft, E.; Piñeros, M.; Kao-Kniffin,

J.; Gu, A. Z. Recycled Phosphorus Bioamendments from Wastewater Impact

Rhizomicrobiome and Benefit Crop Growth: Sustainability Implications at Water-Food

Nexus. *Environ. Sci. Technol.* **2025**, *59* (4), 2131–2143.

https://doi.org/10.1021/acs.est.4c07901.

(59) Li, Y.; Rahman, S. M.; Li, G.; Fowle, W.; Nielsen, P. H.; Gu, A. Z. The Composition and

Implications of Polyphosphate-Metal in Enhanced Biological Phosphorus Removal Systems.

Environ. Sci. Technol. **2019**, *53* (3), 1536–1544. https://doi.org/10.1021/acs.est.8b06827.

(60) Wang, D.; Li, Y.; Cope, H. A.; Li, X.; He, P.; Liu, C.; Li, G.; Rahman, S. M.; Tooker, N. B.;

Bott, C. B.; Onnis-Hayden, A.; Singh, J.; Elfick, A.; Marques, R.; Jessen, H. J.; Oehmen, A.;

Gu, A. Z. Intracellular Polyphosphate Length Characterization in Polyphosphate

Accumulating Microorganisms (PAOs): Implications in PAO Phenotypic Diversity and

Enhanced Biological Phosphorus Removal Performance. *Water Research* **2021**, *206*,

117726. https://doi.org/10.1016/j.watres.2021.117726.

(61) Chavan, S.; Yadav, B.; Tyagi, R. D.; Drogui, P. A Review on Production of

Polyhydroxyalkanoate (PHA) Biopolyesters by Thermophilic Microbes Using Waste

Feedstocks. *Bioresource Technology* **2021**, *341*, 125900. https://doi.org/10/gr8kr7.

(62) Chang, C. F.; Shuman, H.; Somlyo, A. P. Electron Probe Analysis, X-Ray Mapping, and

Electron Energy-Loss Spectroscopy of Calcium, Magnesium, and Monovalent Ions in Log-

Phase and in Dividing Escherichia Coli B Cells. *J Bacteriol* **1986**, *167* (3), 935–939.

https://doi.org/10.1128/jb.167.3.935-939.1986.

(63) Smith, R. J. Calcium and Bacteria. *Adv Microb Physiol* **1995**, *37*, 83–133.

https://doi.org/10.1016/s0065-2911(08)60144-7.

(64) Barra Caracciolo, A.; Grenni, P.; Cupo, C.; Rossetti, S. In Situ Analysis of Native

Microbial Communities in Complex Samples with High Particulate Loads. *FEMS*

Microbiology Letters **2005**, *253* (1), 55–58. https://doi.org/10.1016/j.femsle.2005.09.018.

(65) Kamika, I.; Coetzee, M.; Mamba, B. B.; Msagati, T.; Momba, M. N. B. The Impact of

Microbial Ecology and Chemical Profile on the Enhanced Biological Phosphorus Removal

(EBPR) Process: A Case Study of Northern Wastewater Treatment Works, Johannesburg.

International Journal of Environmental Research and Public Health **2014**, *11* (3), 2876–

2898. https://doi.org/10.3390/ijerph110302876.

(66) Nielsen, P. H.; McIlroy, S. J.; Albertsen, M.; Nierychlo, M. Re-Evaluating the

Microbiology of the Enhanced Biological Phosphorus Removal Process. *Current Opinion*

in Biotechnology **2019**, *57* (Figure 1), 111–118.

https://doi.org/10.1016/j.copbio.2019.03.008.

(67) Stokholm-Bjerregaard, M.; McIlroy, S. J.; Nierychlo, M.; Karst, S. M.; Albertsen, M.;

Nielsen, P. H. A Critical Assessment of the Microorganisms Proposed to Be Important to

Enhanced Biological Phosphorus Removal in Full-Scale Wastewater Treatment Systems.

Frontiers in Microbiology **2017**, *8* (APR), 1–18. https://doi.org/10.3389/fmicb.2017.00718.

(68) Bai, Y.; Zhang, Y.; Quan, X.; Chen, S. Nutrient Removal Performance and Microbial

Characteristics of a Full-Scale IFAS-EBPR Process Treating Municipal Wastewater. *Water*

Science and Technology **2016**, *73* (6), 1261–1268. https://doi.org/10.2166/wst.2015.604.

(69) Singleton, C. M.; Petriglieri, F.; Wasmund, K.; Nierychlo, M.; Kondrotaite, Z.; Petersen, J.

F.; Peces, M.; Dueholm, M. S.; Wagner, M.; Nielsen, P. H. The Novel Genus, 'Candidatus

Phosphoribacter', Previously Identified as Tetrasphaera, Is the Dominant Polyphosphate

Accumulating Lineage in EBPR Wastewater Treatment Plants Worldwide. *ISME Journal*

2022, *16* (6), 1605–1616. https://doi.org/10.1038/s41396-022-01212-z.

(70) Pankratov, T. A.; Grouzdev, D. S.; Patutina, E. O.; Kolganova, T. V.; Suzina, N. E.;

Berestovskaya, J. J. Lichenibacterium Ramalinae Gen. Nov, Sp. Nov., Lichenibacterium

Minor Sp. Nov., the First Endophytic, Beta-Carotene Producing Bacterial Representatives

from Lichen Thalli and the Proposal of the New Family Lichenibacteriaceae within the

Order Rhizobiales. *Antonie van Leeuwenhoek* **2020**, *113* (4), 477–489.

https://doi.org/10/gstz8p.

(71) Wang, H.-G.; Huang, H.; Liu, R.-L.; Mao, Y.-P.; Biswal, B. K.; Chen, G.-H.; Wu, D.

Investigation on Polyphosphate Accumulation in the Sulfur Transformation-Centric EBPR

(SEBPR) Process for Treatment of High-Temperature Saline Wastewater. *Water Research*

2019, *167*, 115138. https://doi.org/10/gmngmh.

(72) Lima, D. C.; Aviles, A. C.; Alpers, R. T.; Perkins, A.; Schoemaker, D. L.; Costa, M.;

Michel, K. J.; Kaeppler, S.; Ertl, D.; Romay, M. C.; Gage, J. L.; Holland, J.; Beissinger, T.;

Bohn, M.; Buckler, E.; Edwards, J.; Flint-Garcia, S.; Gore, M. A.; Hirsch, C. N.; Knoll, J.

E.; McKay, J.; Minyo, R.; Murray, S. C.; Schnable, J.; Sekhon, R. S.; Singh, M. P.; Sparks,

E. E.; Thomison, P.; Thompson, A.; Tuinstra, M.; Wallace, J.; Washburn, J. D.;

Weldekidan, T.; Xu, W.; de Leon, N. 2020-2021 Field Seasons of Maize GxE Project

within the Genomes to Fields Initiative. *BMC Res Notes* **2023**, *16* (1), 219.

https://doi.org/10.1186/s13104-023-06430-y.

(73) Kurashita, H.; Hatamoto, M.; Tomita, S.; Yamaguchi, T.; Narihiro, T.; Kuroda, K.

Comparative Metagenomic Analysis Provides Novel Insights into the Ecophysiology of

Myxococcota in Activated Sludge Systems. bioRxiv April 28, 2024, p 2024.04.26.591422.

https://doi.org/10.1101/2024.04.26.591422.

(74) Zhou, T.; Xiang, Y.; Liu, S.; Ma, H.; Shao, Z.; He, Q.; Chai, H. Microbial Community

Dynamics and Metagenomics Reveal the Potential Role of Unconventional Functional

Microorganisms in Nitrogen and Phosphorus Removal Biofilm System. *Science of The*

Total Environment **2023**, *905*, 167194. https://doi.org/10.1016/j.scitotenv.2023.167194.

(75) Akone, S. H.; Hug, J. J.; Kaur, A.; Garcia, R.; Müller, R. Structure Elucidation and

Biosynthesis of Nannosterols A and B, Myxobacterial Sterols from Nannocystis Sp.

MNa10993. *Journal of Natural Products* **2023**, *86* (4), 915.

https://doi.org/10.1021/acs.jnatprod.2c01143.

(76) Peiffer, J. A.; Spor, A.; Koren, O.; Jin, Z.; Tringe, S. G.; Dangl, J. L.; Buckler, E. S.; Ley, R.

E. Diversity and Heritability of the Maize Rhizosphere Microbiome under Field Conditions.

Proceedings of the National Academy of Sciences **2013**, *110* (16), 6548–6553.

https://doi.org/10.1073/pnas.1302837110.

(77) Pérez-García, L.-A.; Sáenz-Mata, J.; Fortis-Hernández, M.; Navarro-Muñoz, C. E.; Palacio-

Rodríguez, R.; Preciado-Rangel, P. Plant-Growth-Promoting Rhizobacteria Improve

Germination and Bioactive Compounds in Cucumber Seedlings. *Agronomy* **2023**, *13* (2),

315. https://doi.org/10.3390/agronomy13020315.

(78) Kumar Arora, N.; Fatima, T.; Mishra, J.; Mishra, I.; Verma, S.; Verma, R.; Verma, M.;

Bhattacharya, A.; Verma, P.; Mishra, P.; Bharti, C. Halo-Tolerant Plant Growth Promoting

Rhizobacteria for Improving Productivity and Remediation of Saline Soils. *Journal of*

Advanced Research **2020**, *26*, 69–82. https://doi.org/10.1016/j.jare.2020.07.003.

(79) Dai, W.; Liu, M.; Wang, N.; Ye, X.; Liu, Y.; Yao, D.; Wang, L.; Cui, Z.; Yan, P.; Cheng,

C.; Huang, Z.; Wang, H. Positive Contribution of Predatory Bacterial Community to

Multiple Nutrient Cycling and Microbial Network Complexity in Arsenic-Contaminated

Soils. *Applied Soil Ecology* **2023**, *185*, 104792.

https://doi.org/10.1016/j.apsoil.2022.104792.

(80) Yang, X.; Gao, R.; Zhang, Q.; Yung, C. C. M.; Yin, H.; Li, J. Quantification of

Polyphosphate in Environmental Planktonic Samples Using a Novel Fluorescence Dye JC-

D7. *Environ. Sci. Technol.* **2024**, *58* (32), 14249–14259.

https://doi.org/10.1021/acs.est.4c04545.

For Table of Contents Only

895