Discovering Polyphosphate and Polyhydroxyalkanoate-Accumulating Organisms Across

Ecosystems: Phenotype-targeted Genotyping via FACS-sequencing

Yuan Yan[a], Jangho Lee[a,b], Mathew Baldwin[a], Guangyu Li[a], Il Han[a,c], Zijian Wang[d], Peisheng

He[a], Varun Shrinivasan[a,e], Liam Wicked-Do[f], Michael Gore[f] and April Z. Gu[a*]

[a]School of Civil and Environmental Engineering, Cornell University, Ithaca, NY 14850, United

States

[b]Department of Chemistry, CUNY Hunter College, 695 Park Ave, New York, NY 10065, United

States

[c] Metrohm Spectro, Inc., 313 Enterprise Drive, Plainsboro Township, NJ 08536, United States

[d]School of Biological and Environmental Engineering, Cornell University, Ithaca, NY 14850,

United States

[e] Trinnex, Boston MA 02109, United States

[f]Plant Breeding and Genetics Section, School of Integrative Plant Science, Cornell University,

Ithaca, NY 14850, United States

[*] Corresponding author: April Z. Gu. aprilgu@cornell.edu

Abstract

Polyphosphate-accumulating organisms (PAOs) play a central role in enhanced biological

phosphorus removal (EBPR) systems by coupling intracellular polyphosphate (polyP) cycling with

carbon storage through polyhydroxyalkanoates (PHA). However, in natural environments such as

soil and sediments, the term PAOs has been loosely applied to polyphosphate-containing

organisms (PCOs) without consideration of their carbon storage capacity or phylogeny. This

oversimplification overlooks a critical metabolic trait—surplus co-accumulation of polyP and

PHA—that defines functionally active PAOs in engineered systems. This study presents a novel

phenotype-targeted approach that integrates triple-stained fluorescence-activated cell sorting

(FACS) with 16S rRNA gene sequencing, TriFlow-Seq, enabling the simultaneous detection,

quantification, and phylogenetic characterization of functional PAOs capable of dual polyP and

PHA accumulation. TriFlow-Seq was applied to microbial communities from full-scale and pilot-

scale EBPR systems, as well as maize rhizosphere soils across multiple genotypes and fields. In

EBPR systems, the method successfully enriched known PAOs, including *Candidatus*

Accumulibacter, *Tetrasphaera*, *Dechloromonas*, and *Pseudomonas*, while also revealing novel

candidate PAOs, particularly within *Rhodobacteraceae*. In agricultural soils, TriFlow-Seq

uncovered a previously unrecognized and diverse PAO community, dominated by *Pseudomonas*,

Halomonas, and *Nannocystis*. Notably, PAO community composition in the maize rhizosphere

exhibited significant genotype-specific patterns, indicating plant genetic background influences

the recruitment and enrichment of functional PAOs. These findings highlight the importance of

exploring alternative PAOs and their specific metabolic preferences across environmental samples.

By providing insights into the detection and characterization of bacteria of interest, flow cytometry

combined with phylogenetic analysis offers a promising avenue for advancing our understanding

of PAOs and their roles in wastewater treatment and environmental processes.

1   Introduction

A diverse range of microbes contain intracellular biopolymers, including polyphosphate (polyP),

polyhydroxyalkanoates (PHA), glycogen, wax esters, triacylglyceride (TAGs), trehalose, and

cyanophycin.[1] These biopolymers serve versatile functions such as energy storage, nutrient

cycling , ion balance, redox homeostasis, and stress resistance, allowing microbes to adapt to

fluctuating environmental conditions.[1–3] Among them , polyP-accumulating organisms (PAOs)

have been widely recognized as a key agent to achieve sustainable phosphorus (P) removal and

recovery such as in enhanced biological phosphorus removal (EBPR) systems .[4,5] In EBPR systems,

PAOs exhibit a unique metabolism: under anaerobic conditions, they utilize stored polyP as an

energy source, releasing phosphate and concurrently accumulating PHA. Then under subsequent

oxic or anoxic conditions, they take up phosphate and synthesize new polyP while using stored

PHA to fuel growth.[6] The metabolic interdependence of dual intracellular polymers (polyP and

PHA) is a defining feature of functionally verified PAOs in EBPR systems, such as *Candidatus*

Accumulibacter,[6] *Pseudomonas*,[7] and *Dechloromonas*,[8] although some cannot accumulate PHA

such as *Tetrasphaera*.[9] This unique metabolism provides competitive advantages for PAOs under

fluctuating anaerobic/oxic conditions.[8,14,15]

Beyond EBPR engineering systems, there are increasing interests in exploring and

identifying functional roles of PAOs in natural environments, as emerging evidence points to their

ubiquitous presence and versatile functions.[10–13] However, PAOs in natural environments are often

identified broadly as polyphosphate-containing organisms (PCOs), with little knowledge of the

potential co-metabolisms with carbon (C) biopolymer storage. Here, we are specifically interested

and thus propose a refined definition of functional PAOs in natural environments as organisms

that co-accumulate both polyP and intracellular C, predominately PHA, to surplus levels, where

the co-metabolism holds important ecological roles in these systems for following reasons. Firstly,

as one of the dominant C storage biopolymers, PHA concentrations (~1-4µgC/g soil) are positively

responsive to C availability, but are suppressed by nitrogen availability, suggesting its role as an

alternate C allocation strategy for stoichiometric balancing and stress response.[16,17] The co-storage

of C biopolymers and polyP likely enhances microbial adaptability to nutrient fluctuations and

contributes to broader nutrient cycling and ecosystem functioning. This dual-storage strategy

distinguishes specialized PAOs exhibiting resource-dependent surplus storage from organisms that

accumulate polyP primarily for stress responses or cation sequestration.[17–19] Secondly, our

preliminary results indicate that up to 73.7% of polyP-accumulating cells in crop soil contain C

storage biopolymers with significant Pearson correlation between polyP-accumulating cells and

polyP-C-accumulating cells (R=0.8, P<0.001) (Figure S1). Lastly, the inability of polyP-

synthesizing fungal organisms, such as arbuscular mycorrhizal fungi, to produce PHA provides a

clear demarcation between bacterial and fungal PAOs. Fungal PAOs were more extensively

studied, whereases relatively little is known about bacterial PAOs in natural environments.[24]

Therefore, we focus on identifying bacterial microorganisms exhibiting surplus storage, defined

by their specific ability to accumulate "substantial" polyP and PHA beyond immediate biological

maintenance, distinguishing them from microbes that maintain constitutively low levels of these

biopolymers for basic cellular functions (e.g., *Escherichia coli* or *Bacillus subtilis*).[20,21]

Despite the ecological, biogeochemical, and engineering significance of PAOs, identifying

functionally active and phylogenetically novel PAOs in complex environmental matrices remains

challenging.[10,22–24] Conventional approaches rely heavily on gene-centric profiling such as polyP-

metabolism biomarkers (i.e. ppk) or metagenomics to identify and quantify previously known

PAOs across aquatic and terrestrial systems,[25,26] yet these methods have limitations and can not

reflect actual phenotypic traits such as PolyP accumulation. [25–27] Discovery of novel PAOs is often

prohibited with *ppk* gene-based detection that rely on prior genetic information of the targeted

PAOs and, it also has low specificity for detecting PAOs with unqiue polyP accumulation ability

as the polyP-synthesizing gene *ppk* is ubiquitous across microorganisms.[27,28] Moreover, polyP

metabolism is not solely regulated by *ppk* gene, as alternative proteins, such as actin-like Arp

proteins, have been proposed to exhibit PPK-like functions in bacteria.[29] Methods for detecting

PAOs that co-accumulate polyP and PHA in natural systems have never been reported. A few

recent studies have employed fluorescence-activated cell sorting (FACS) to phenotypically track

the polyP accumulation traits and resolve the phylogeny and taxonomy of PAOs from EBPR

systms,[30,31] although these studies stains only for polyP, without detecting the co-metabolism

polymers like PHA.[31] In addition, FACS-based PAO detection in soil samples have never been

reported.

In this study, we developed an integrated approach combining triple-staining fluorescence-

activated cell sorting (FACS) with genomic sequencing, termed here as TriFlow-Seq. Beyond

polyP-based PAO detection, TriFlow-Seq enables the simultaneous detection of functionally

active PAOs with both polyP and PHA, where sorted cells were subsequently subject to 16S rRNA

amplicon sequencing for phylogenetic identification. To validate this method, we applied an

independent phenotypic detection technique—Single-Cell Raman Spectroscopy (SCRS)—an

established approach for the simultaneous detection of intracellular polyP and PHA in EBPR

PAOs,[23,32–35] in parallel for comparison. We further demonstrated the successful application of

TriFlow-Seq across diverse environmental samples, including activated sludge from four full- and

pilot-scale EBPR facilities and rhizosphere soil from maize croplands in the U.S. By addressing

key limitations of conventional methods, this approach enhances our ability to resolve the

identities of bacterial PAOs beyond known taxa across diverse environmental systems, from soils

and sediments to water bodies.

2   Material and Methods

2.1   Sample collection.

2.1.1   Activated sludge samples from EBPR systems

Biomass samples from the aerobic zone were collected from three different full-scale EBPR plants

from Hampton Roads Sanitation District (HRSD) that have effective P removal via EBPR

activities thus containing active PAOs, namely in HRSD Nansemond Treatment Plant (Suffolk,

VA, USA), HRSD Virginia initiative Treatment Plant (Nortfolk, VA, USA), HRSD Atlantic

Treatment Plant (Virginia Beach, VA, USA), as well as from one pilot-scale side-stream EBPR

(S2EBPR) (VA, USA). The collected activated sludge samples from EBPR systems were stored

at 4 °C, transported overnight to the laboratory, and activated with phosphorus release/uptake test

(Text S1, Figure S2) while sample from pilot-scale S2EBPR was directly fixed onsite as described

in section 2.4.

2.1.2   Soil samples

Sixteen maize genotypes, six inbred and ten hybrid lines (Table S1), were each planted in a

randomized complete block design (n=2) across two fields respectively with standard planting

density (24 plants/plot), each characterized as Lima Silt Loam with a 0-3% slope at Musgrave

Farm in Aurora NY (42.733317, -76.652394). Rhizosphere samples were collected at tasseling

(VT) in accordance with Hanway et al.[36] Briefly, from each plot, 3 randomly selected plants were

uprooted with loosely attached bulk soil shaken off, root adhered soil was taken as rhizosphere

soil, pooled by plot and stored and transported at 4⁰C for sample pre-processing within 24 hours.

2.2   Microbiota isolation

2.2.1   Rhizosphere microbiota

The microbial fraction of the soil was isolated via detached from 1 g of soils using detachment

detergent (3 mM sodium pyrophosphate, 0.5% Tween20, 0.35% PVPP in 1xPBS) in a 1:20 ratio

shaken at 200 rpm on an orbital shaker for 30 mins at room temperature as previously described

in Deng et al., (2019). Following detachment, to prevent re-aggregation, detached soil community

was washed 1x with 0.4 mM EDTA dissolved in 1X PBS (pH ~7.2) for excess extracellular cation

removal followed by 2x washing with 1x PBS.

2.2.2   Activated sludge sample

Sludge samples were collected in the middle of the aerobic phase from each condition and were

centrifuged to remove supernatant. The remaining pellets were washed three times with 1X PBS

for excess cation removal.

2.3   Cell fixation

The fixation buffer was prepared according to previously described protocol,[38,39] containing 5 mM

Nickel Chloride ($NiCl_2$), 5 mM Barium Chloride ($BaCl_2$), 10 % Sodium azide (NaN_3) in HEPES

Buffer (25 mM), pH = 7.2. The remaining pellets were resuspended in a 5 ml fixation buffer,

vortexed for 10 s, and were stored at 4 °C for 1 day.

2.4   Triple-staining

Fixed sludge samples were washed three times with phosphate-buffered saline (PBS: 0.4 M

Na_2HPO_4/NaH_2PO_4, 150 mM NaCl, pH 7.2) with centrifugation at 3200 g for 10 min while fixed

soil samples were washed three times with 1X PBS with centrifugation at 6000g for 10 mins at

4°C. Fixed and washed sludge (1 ml) were treated with 0.584 ml solution A (0.11M citric

acid/4.1mMTween 20 in distilled water) for 20 min, washed and resuspended in 1ml of PBS buffer.

Fixed, washed, and resuspended (5 mL) soil pellets were similarly treated, with 1.6mL solution A

for 20 mins, washed, and resuspended in 5mL of PBS.

DNA, polyP and PHA of the cells were stained with SYBR Green I (SG) (Ex 488 nm / Em 515

nm), tetracycline hydrochloride (TC) (Ex 390 nm / Em 560 nm) (Sigma Chemical Co., USA), and

Nile red (NR) (Ex 532 nm / Em 610 nm) (Sigma Chemical Co., USA), respectively–each validated

in previous studies.[30,40–43] First, TC was applied as a 2 mg/ml stock solution in milli-Q water (0.225

mM in cell suspensions) and incubated for 10 min.[39] NR stock solution (10 mg/ml in DMSO) was

added to 1 ml of cells a as final concentration of 0.05 mg/ml, and then incubated for 10 min.[44]

Lastly, 100X SG was applied into the cells, and it was incubated for 4 hours.[37] All incubation steps

were done at room temperature in the dark.

2.5   Floc Disaggregation

Unlike activated sludge, rhizosphere samples are particularly prone to aggregation due to root

exudates, mucilage, and extracellular polymeric substances (EPS), which can reduce sorting

efficiency and risk nozzle clogging. A disaggregation process was therefore developed to further

prevent re-aggregation during FACS. The protocol involves resuspending the stained pellet in 3

mL of 1X Trypsin-0.5mM EDTA for 1 minute, followed by trypsin deactivation using 1X FBS

(1:1 ratio). Prior to FACS, trypsinized samples were diluted in 1X PBS to achieve a proper

threshold rate of 1500 events s^{-1}.

2.6   FACS settings and Validation of FACS gating strategy

Cell sorting was carried out using the 100-micron nozzle on Fusion Aria III cell sorter (BD, U.S.)

equipped with a blue laser (488nm excitation) with 530 nm band pass (BP) filter for SG, a violet

laser (405nm excitation) with 525 BP filter for TC, and a yellow-green laser (561nm excitation)

with a 610 BP filter for NR. The blue laser was also used to analyze forward scatter (FSC) which

is related to cell size and side scatter (SSC) which is related to cell granularity. Manual gating was

performed and summarized in supplementary material (Text S2, Figure S3). Bacterial cells were

sorted in purity mode at a rate less than 1500 cells per second. To obtain sufficient DNA for the

following DNA preparation and analyses, up to 10^5 cells were sorted.[45] The cells were sorted into

sterile 1.5 mL micro tubes and frozen for further analysis.

To ensure the reliability and accuracy of the proposed TriFlow-Seq in identifying and quantifying

PAOs in complex microbial communities, we employed two independent validation methods.

We first verified the PAO enrichment via fluorescent confocal microscopy. Approximately 4000

sorted cells were collected to validate that all sorted cells were positive with SG, TC and NR.

Unsorted cells as control and sorted cells were first filtered through black polycarbonate filter (0.22

um pore size) to concentrate cells. The filters were placed onto slide glass with a drop of antifade

reagent. The validation of the gating methods was carried out by observation through confocal

microscopy (Carl Zeiss) with 800X magnification and three excitation/filter sets: blue light (BP

450–490 nm, FT 500 nm, LP 515 nm), green light (BP 510–560 nm, FT 565 nm, LP 590 nm) and

ultraviolet (BP 340–380 nm, FT 400 nm, BP 435–485). Quantification of TC and NR-stained cells

relative to SG-stained cells was performed using Daime (version 1.2).[46]

Additional validation was conducted by comparing FACS results with SCRS analysis, an

independent method previously developed by our group quantifying polyP and PHA-containing

cells.[32,33] Samples were prepared on optically polished CaF_2 windows (Wilmad LabGlass, Buena,

NJ). Raman spectra were acquired using a Horiba LabRam Evolution (Horiba, Kyoto, Japan)

equipped with 50X objective (Olympus LMPLFLN 50x, Tokyo, Japan) in accordance with

established protocol.[23,47,48] Relative polymer content in each individual cell was evaluated based

on the signature peak intensity (CCD counts) of each polymers.

2.7   DNA extraction and Amplicon sequencing

Genomic DNA of sorted soil and sludge cells was extracted using InstaGene matrix (Bio-

Rad Laboratories, USA). Genomic DNA from unsorted raw soil and sludge was extracted via

DNeasy Power Soil Pro kit (Qiagen, Valencia, CA, USA) carried out on the QIAcube® (QIAGEN)

platform. The V4 region of the 16S ribosomal RNA (16S rRNA) genes from both sorted and

unsorted samples were amplified and sequenced using primers 515F and 806R at University of

Connecticut-MARS (Microbial Analysis, Resources, and Services) facility. Details of sequencing

were listed in Text S3.

2.8   Statistical Analysis

The Spearman correlation coefficients and corresponding *p*-value were computed using Python3

Scipy 1.10.1 package. Principal Coordinates Analysis (PCoA) was performed using *R vegan* (2.6-

4) to assess beta diversity based on Bray-Curtis dissimilarities of normalized relative abundance

data. The ordination plot was visualized using *ggplot2* (3.4.4).

3   Results and discussion

3.1   Development of FACS TriFlow method for sorting functional PAO with dual

accumulation of polyP and PHA

The effectiveness of the triple-stained FACS approach for identifying and sorting PHA-containing

PAOs were tested with EBPR systems that are known to contain PAOs (positive control

environmental samples). The density of the triple-stained cells before and after the FACS

enrichment were visualized using confocal microscopy (Figure 1A). A significant and evident

enrichment of candidate PAO cells that contain both polyP and PHA were shown in the sorted

samples (Magenta) compared to the original sludge, which was validated in all four tested EBPR

systems. In addition, relative abundances of PHA-positive and polyP-positive cells as of total cells

(SG-stained cells) were semi-quantified using digital image analysis (Figure 1B). NR-stained and

TC-stained cells were clearly enriched in all FACS sorted samples in comparison to the original

sludge samples. These results demonstrated that our gating strategy in the FCAS analysis

effectively led to dual-polyp-PHA-accumulating PAO enrichment as expected, and the gating

strategy employed was appropriate.

A)

*Green, DNA-postive cells; Red, PHA-positive cells; Blue, polyP-positive cells; Magenta, polyP and PHA-positive cells

B)

Sample	Unsorted cells (%)		Sorted cells (%)	
	NR+ (PHA-positive)	TC+ (polyP-positive)	NR+ (PHA-positive)	TC+ (polyP-positive)
ATL	50.3	35.6	99	98.3
NM	45.7	53.7	97.3	87.7
VIP	68.1	29.1	79.8	80.9
S2EBPR	19	6.8	96	83.6

NR+, Nile Red-stained cells; TC+, Tetracycline-stained cells.

233 **Figure 1 Images from confocal microscopy for comparison between FACS sorted stain-positive cells and**
234 **unsorted cells in activated sludge. Samples from Atlantic, Nansemond, VIP EBPR systems and a S2EBPR**
235 **system, respectively; B) Relative abundance of NR (PHA)-positive and TC (polyP)-positive- cells in FACS**
236 **sorted samples versus original unsorted samples by semi-quantitative confocal microscopic image analysis.**

3.2   Validation of polyP-PHA-acumulating PAO quantification through TriFlow

Prior to implementing the TriFlow in soil samples, we validated polyP-PHA dual staining by comparing the relative abundance of polyP-containing cells through FACS with those determined by SCRS. SCRS, which targets the molecular composition of microbial cells, has demonstrated exceptional sensitivity and accuracy in detecting and quantifying intracellular polymers such as PolyP, PHA, and glycogen.[49] SCRS-based detecton of PAOs containing both polyP and PHA were demonstrated in our previous studies (ref.) Detailed sampling information and the linear correlation between the two methods are provided in Text S4.

We compared the quantification of PAOs with polyP-detection irrespective of intracellular PHA containment, as well as the quantification of dual polyP-PHA-accumjuting PAOs, across various microbial samples from EBPR systems, from bulk soil and from maize rhizosphere. A positive correlation between the percentage of polyP-containing cells identified through FACS and SCRS was demonstrated (Pearson coefficient r^2 = 0.6974158, p-value = 0.000309; Figure 2A). Importantly, the percentage of PAOs determined by FACS and SCRS falls in the same range. This significant and strong correlation between the two methods for detecting and sorting polyP containing cells validated the proposed FACS protocol for sorting PAOs.

We further compared the performance of triple-stained FACS and SCRS in identifying the relative abundances of putative functional PAOs (containing both polyP and PHA) across diverse samples including soil, rhizosphere, EBPR activated sludge, and isolate PAO strains (*T. elongata* DSM 14184) (Figure 2B). They demonstrate a weak linear (Pearson's r = 0.488) yet still a significant positive monotonic relationship (Spearman's r = 0.62, p-value = 0.009).

Figure 2 Spearman's rank correlation coefficients of PAOs and polyP-containing cells detected by FACS and
SCRS. (A) polyP-containing cells, irrespective of PHA presence; (B) PAOs, defined as cells containing both
polyP and PHA. The strong correlation between the two independent techniques validates the effectiveness of
the triple-stained FACS in detecting and quantifying polyP-containing cells and PAOs in complex microbial
communities.

The percentages determined by FACS were consistently lower than those identified by SCRS,

likely stems from fixation-free characteristics of SCRS methods as discussed in the following

section.It is important to note that FACS and SCRS employ fundamentally different phenotyping

mechanisms, leading to distinct approaches in quantifying target cell populations. SCRS is a

fixation-free method that can detect intracellular PHA and polyP based on the unique Raman

spectral signature regardless of counterion type.[50] FACS, however, requires a fixation step that

enhances dye penetration but permanently modifies cell wall and membrane permeability.[51] This

modification typically results in a 10-20% loss of intracellular polymers, similar to observations

with FISH fixatives.[52] In addition, the accuracy of FACS largely rely on the specificity and

sensitivity of fluorescent dyes. Nile Red, although widely employed for PHA detection in flow

cytometry, can also capture other lipophilic inclusions such as trehalose and triglycerides.[53]

Similarly, tetracycline detects polyP through preferentially complexes with countercharge divalent

(e.g., Ca^{2+}, Mg^{2+}) in polyP granules.[54–56] These methodological differences suggest that FACS may

slightly overestimate PHA-containing PAOs relative to SCRS method, consequently reducing the

slope of the correlation curve. Furthermore, FACS employs size-based filtration (40μm cell

strainer) to prevent flow cytometer nozzle clogging, while SCRS utilizes density-gradient

centrifugation for cell extraction.[57] These procedural variations likely influence the organisms

being analyzed, contributing to the observed differences in PAO abundance percentages. Despite

these discrepancies, the consistent trend in PAO detection by both methods underscores the

viability of the triple-stained FACS approach for identifying and quantifying PAOs in complex

microbial communities.

3.3   Application of TriFlow-Seq in Identification and Quantification of PAOs in EBPRs

Combining triple-stained FACS with downstream 16S rRNA genes amplicon sequencing (termed

as TriFlow-Seq) led to enrichment, quantification, and identification of both known and candidate

PAOs in pilot and full scale EBPR plants. The relative abundances of known PAOs in the original

sludge prior to the FACS sorting, versus those after the FACS enrichment were depicted in Figure

3.

291
Figure 3 A) Relative abundances of known candidate PAOs before (left) and after (right) FACS sorting in three
full-cale EBPR plants and one pilot-scale S2EBPR plant. These samples were activated with P release and
uptake tests. The groups that had relative abudances below 0.1% were not shown; B) Differential abundance
analysis for idnetification of unknown candidate PAOs at the genus level following FACS sorting. Heat map
represents t-statistics derived from two-tailed t-tests on log10-transformed fold changes. Blue hues denote
positive t-statistics (enrichment), while red hues indicate negative t-statistics (depletion). Statistical significance
levels: *p < 0.05, **p < 0.01, ***p < 0.001. Red pentagram indicates the genus has been experimentally
confirmed with PHA/polyP and PAO activities; Yellow pentagram indicates the presence of polyP and PHA;
green pentagram indicates the the reference genome contains necessary genes for polyP and C storage synthesis;
blue pentagram indicates that some genus within the Family have polyP accumulation ability. Detailed
information can be found in Table S2. Analysis limited to the top 50 genera by abundance.

3.3.1 TriFlow-Seq enriched known PAOs in EBPR systems

TriFlow-Seq effectively identified previously known and expected PAO genera in EBPR systems,

including *Pseudomonas*, *Dechloromonas*, and *Ca*. Accumulibacter[58–61] across all plant samples

(Figure 3A), validating its capability to identify organisms known to contain both polyP and PHA

in EBPR systems.

The sorted communities were characterized by the enrichment ratios, calculated as the

relative abundance of each genus post-sorting divided by its pre-sorting abundance (log10 scale).

Higher ratios indicate a greater enrichment of bacteria capable of accumulating both polyP and

intracellular C polymers, thus reflecting the proportion of putative PAOs in that taxonomic group

(Figure 3B).

Ca. Accumulibacter, despite being consistently identified as the dominant PAO in unsorted

samples in three full-scale plants except S2EBPR pilot plant (1.2~4.0%) (Figure 3A), showed

varied enrichment ratios among the EBPR plants (Figure 3B). Its relative abundance even

decreased after sorting in the VIP samples. Interestingly, *Tetrasphaera* emerged as the most

significantly enriched genus after sorting (Figure 3A and 3B). Although initially thought to lack

PHA synthesis capability, recent studies have revealed that anaerobically PHA synthesis depends

on clade composition of *Tetrasphaera*.[62–65] For instance, the *T. japonica* genome possesses PHA

synthase (*phaC*) and was proved experimentally to synthesize PHA under anaerobic condition.[63]

This evidence indicates that *T. japonica* or potentially uncultured *Tetrasphaera* that possess

Accumulibacr-like PAO metabolism reside in full-scale EBPR.[66] Overall, the relative abundances

of known PAOs were only 2 to 12% out of total PHA+polyP positive cells (Figure 3A),

highlighting the larger phylogentic diversity of PAOs than our current understanding.

3.3.2   Discovery of candidate PAOs in EBPR

It should be noted that even within known PAOs, some sub-genus populations were not

actively performing polyP and PHA accumulation, resulting in a negative enrichment factor, for

example, Ca. Accumulibacter. Therefore, the selection of putative PAOs in sludge were based on

(1) consistent detection across sorted communities, and (2) prior evidence of or genomic potential

for polyP and PHA accumulation. A notable enrichment of putative PAOs was observed, including

unclassified *Rhodobacteraceae*, unclassified Rhizobiales, unclassified Caldilineaceae, *Lautropia*,

etc (Figure 3B). The identified OTUs associated with these bacterial taxa were not classified to

genus level, suggesting the potential presence of unknown genera, which could be candidate PAOs

with or without classical EBPR metabolisms. Bacteria isolated from the order Rhizobiales were

found to contain PHA and polyP.[67] An uncultured genus close to the *Rhodobacteraceae* family is

speculated to be sulfur PAOs.[68] these new insights provide initial novel candidate PAOs and

whether these putative PAOs can contribut to P rempval as those cronical PAOs require further

investigation.

These results led to the following findings: (1) not all genetically identified *Ca*.

Accumulibacter display metabolically active PAO phenotypes, as evidenced by their varied

enrichment ratios; (2) *Tetrasphaera* showed unexpected and significant enrichment with variable

intracellular PHA content, revealing phenotypic diversity beyond current understanding of this

genus; and (3) The TriFlow-Seq method uncovered novel candidate PAOs from previously

unrecognized taxonomic groups, expanding our understanding of PAO diversity in EBPR systems.

Although a comprehensive comparison of the PAO population in the tested EBPR plants is beyond

the focus of this study, this application has demonstrated that the TriFlow-Seq is a valuable tool

for rapidly enriching targeted bacterial populations that exhibit the phenotypic traits characteristic

of functionally relevant PAOs. This approach was further adapted and optimized to be applicable

to more heterogeneous sample types, such as bulk soil, rhizosphere soil, and sediments, enabling

the identification and characterization of PAOs in these complex environmental matrices.

3.4 Identification and Quantification of the Previously Unknown Rhizosphere PAO Community

3.4.1 Key Steps Modified for soil sample processing.

Significant challenges were faced in applying FACS to soil samples, which differ markedly from activated sludge in their physico-chemical properties and proneness to reaggregation. Traditional soil pretreatments using detergents or sonication quantifications prove inadequate for sorting low-abundance populations,[69,70] especially when high cell numbers are required for quality genomic reads. Key challenges include reaggregation during extended suspension, which reduces sorting efficiency and increases false positives,[71] and soil heterogeneity, which affects reaggregation rates due to variations in cation exchange capacity, plant and microbial protein activity, soil organic matter content and biofilm formation.[72] Our newly developed pretreatment method (Section 2.5) successfully overcomes these obstacles, demonstrating effectiveness across three diverse soil types: a silt loam from central NY, a soil fertilized with ammonium sulfate and high soil organic matter in Urbana Champaign Illinois, and a potting soil mixture. This method not only allows for effective sorting of low-abundance community members but also represents a significant step towards standardization in FACS preparation for varied soil types.

Notably, our soil pretreatment method achieved sorting efficiencies ranging from 77-92% across two soil series tested (n=8 samples), with an average of 85.5% (Table S3). This represents a substantial improvement, approximately 50% higher than the 35.75% average sorting efficiency obtained using standard WWT sample pretreatment (n=8 samples, Table S3). DNA extraction revealed the success of sorting with DNA quantification ranging from 1ng/ul to 16.3 ng/ul with downstream 16S rRNA amplicon sequencing quality control retrieving high quality reads (Phred >30) from all but one sequenced sorted sample (n=192).

    3.4.2   Effectiveness of TriFlow-Seq in rhizosphere samples

The identity of active prokaryotic PAOs in the rhizosphere of maize are mostly unknown. For the

first time, the bacterial PAO community in the maize rhizosphere of sampled Nested Association

Mapping (NAM) and a subset of phenotypically diverse hybrids from the High Intensity

Phenotyping Site (HIPS) panel were revealed through our triple stained FACS coupled to

downstream 16S rRNA sequencing.[73] Enrichment profiles of key sorted taxa across maize

genotypes were further investigated to identify candidate functional PAOs in maize rhizosphere.

381
Figure 4. Identification of candidate rhizosphere PAOs and characterization of rhizosphere PAO community
via coupled triple stained FACS and downstream 16S sequencing. A) Principal Coordinate Analysis (PCoA)
based on Bray-Curtis dissimilarity of unsorted raw and triple stained FACS sorted-16S sequenced communities
across NAM (inbred) and HIPS (hybrid) lines; B) Phylum and C) Genus level community composition and
relative abundance of unsorted raw rhizosphere community and FACS sorted-16S sequenced PAO community,
separated by field, abundance filtered > 0.03; D) Differential abundance analysis of candidate rhizosphere PAO
taxonomies in NAM/HIPS rhizosphere following FACS sorting-16S sequencing. Blue scale denotes positive
enrichment in PAO community and red denotes negative enrichment, significance based on two-tailed t-test

To assess the effectiveness of TriFlow-Seq in identifying rhizosphere PAOs, beta and alpha

diversity analysis were conducted to assess shifts in community structure post-sorting. Principal

Coordinate Analysis (PCoA) revealed distinct clustering and significant separation ($p<0.001$,

PERMANOVA) between the unsorted and sorted community along PCo1, which explained 35.4%

of the total variance. This separation indicates effective isolation of a distinct sub-community

(Figure 4A). Within both raw and sorted clusters, two significant sub-clusters emerged ($p<0.001$,

PERMANOVA) based on maize genotypes (inbred versus hybrid) along PCo2, accounting for an

additional 11.1% of variation (Figure 4A). This pattern indicates the sorted population retains its

community fingerprinted from its original rhizosphere population. Alpha diversity was then

conducted to understand impacts on community complexity. The sorted community exhibited

significantly lower diversity ($p<0.001$) across all metrics, including Species Richness, Shannon

Index, and Simpson Index (Figure S5), indicating that TriFlow-Seq successfully concentrated a

highly specialized sub-community characterized by a few dominant taxa from rhizosphere soil.

Proteobacteria, Actinobacteria, and Acidobacteria were the most abundant phyla in both

unsorted and sorted communities (63.2-66.2% and 59.1-59.5% respectively), aligning with

typically observed community compositions in both maize rhizosphere and activated sludge

communities (Figure 4B). Notably, the "highly social" Myxcococcota phyla, known for complex

sterol synthesis and recently revealed in activated sludge to widely contain PAO-like *ppk1*,[74,75]

was consistently detected only within the sorted communities (6.58-8.54%) (Figure 4B). While

putative PAOs such as *Tetrasphaera*, *Ca.* Accumulibacter, and *Dechloromas* were detected within

the sorted community, their combined prevalence <1% suggests minimal contribution to the

rhizosphere processes. Rather, *Pseudomonas* (9.03-12.7%) and *Halomonas* (11-15.6%), both

putative PAOs identified in EBPR, were the predominate genera detected in the sorted

communities. Additionally, *Nannocysitis* (3.25-7.56%), a *Myxcoccota* genus capable of polyP

accumulation,[76,77] was consistently detected across sorted communities regardless of soil types or

maize genotypes. The taxonomic composition of sorted communities further supports the efficacy

of our method in identifying putative PAOs as well as taxa with known genomic potential for

polyP and internal C accumulation.

3.4.3   Discovery of new PAOs in maize rhizosphere

Following confirmation of the efficacy of our TriFlow-Seq in identifying PAO or PAO-

like populations, candidate functional rhizosphere PAOs were identified using selection criteria

similar to those established for activated sludge systems. These criteria were adapted to address

the inherent heterogeneity of soil systems, which results in variable phenotypic profiles and diverse

metabolic activities of putative PAO. Enrichment profiles at the sub-genus level revealed selective

dominance of specific ASVs within the predominate sorted genera, leading to the identification of

three candidate rhizosphere PAOs. Among these, two predominant putative PAO ASVs exhibited

positive enrichment across all maize lines: *Halomonas* ASV_14 (21.1-22.1-fold) and

Pseudomonas ASV_51 (20.7-22-fold) (Figure 4D). Additionally, *Nannocystis* ASV_54, most

closely related to *Nannocystsis exedens* (93.68% 16S rRNA sequence simialrity, BLASTN)*,*

exhibited positive enrichment (1.14-4.46-fold) across 68.75% of samples localized to sampled

NAM lines (including B73), and most B73 parent hybrid lines (p<0.05) (Figure 4D). Analysis of

the available annotated genome (GenBank GCA_002343915.1) of *N. exedens* revealed the

presence of *ppk1*, *ppx*, sterol and steroid biosynthetic pathways.[78] While this study does not aim

to exhaustively compare PAO populations across maize genotypes, our proof-of-concept analysis

revealed distinct patterns of PAO association with different maize genetics, which intrigues further

investigation, .

These results, for the first time, revealed the characteristics, diversity, composition, and

distribution pattern of rhizosphere PAO community. The emerging evidence points to the

important roles of PAOs in rhizosphere communities and processes, including halotolerance,

phytohormone production, siderophore production, beneficial antagonism/antibiotic production,

and nutrient cycling.[79–81] Our proposed TriFlow-Seq method enables the discovery of new PAOs

in agricultural soil or other natural environments, which will expand our understanding of the

identity and diversity of PAOs in the natural environments.

4   TriFlow-Seq: Achievements, Applications, and Future Prospects

Linking phenotype-based detection and genotype-based characterization provides valuable

insights into the discovery of organisms of interest. This study developed and validated the

proposed TriFlow-Seq method and demonstrated two distinct applications: identification of

functional PAOs in wastewater treatment and resource recovery facilities; and elucidation of

unknown PAOs with previously unrecognized roles in P and C cycling within the maize

rhizosphere. By successfully modifying FACS preprocessing protocols for soil samples, we

enabled the identification of candidate PAOs in complex natural environments including soil and

sediment systems. Our findings revealed unexpectedly high PAO prevalence and distinct

phylogenetic patterns across different maize genotype lines, establishing a pioneering approach to

explore PAO identities and functions in rhizosphere systems. Future applications of TriFlow-Seq

could be enhanced through several approaches: (1) incorporation of additional phenotyping targets

as new staining materials become available, potentially revealing greater PAO diversity; (2)

integration with downstream mini-metagenomics or other omics techniques to obtain deeper

functional and metabolic insights into candidate PAOs; and (3) combination with FISH to increase method specificity for targeting particular microbial groups. Despite current limitations, this technique demonstrates significant potential for identifying novel PAOs within complex microbial consortia, advancing our understanding of microbial polyP metabolism and phosphorus cycling across diverse ecosystems.

5   Acknowledgement

This study was partially supported by the Cornell Institute for Digital Agriculture (CIDA), and Center for Research on Programmable Plant System (CROPPS) funded by the National Science Foundation (Grant No. DBI-2019674). We thank Dr. Lydia Tesfa, Jaclyn Mahoney, and Michael Sledziona from the Flow Cytometry Facility of the Biotechnology Resource Center at the Cornell University (RRID:SCR_021740) for their technical assistance with FACS experiments. Special thanks are given to the generous support of operators and staff of HRSD for their assistance with activated sludge sample collection for the Water Research Foundation project（Grant No. 4901). We thank Dr. Michael Gore and Liam Wickes-Do for their generous time and effort in curating, planting, tending, and growth of maize NAM and HIPS Panels.

478   References

479   (1)   Mason-Jones, K.; Robinson, S. L.; Veen, G. F. (Ciska); Manzoni, S.; van der Putten, W. H.

480   Microbial Storage and Its Implications for Soil Ecology. *ISME J* **2022**, *16* (3), 617–629.

481   https://doi.org/10.1038/s41396-021-01110-w.

482   (2)   Rao, N. N.; Gómez-García, M. R.; Kornberg, A. Inorganic Polyphosphate: Essential for

483   Growth and Survival. *Annual Review of Biochemistry* **2009**, *78* (Volume 78, 2009), 605–647.

484   https://doi.org/10.1146/annurev.biochem.77.083007.093039.

485   (3)   Kornberg, A.; Rao, N. N.; Ault-Riché, D. Inorganic Polyphosphate: A Molecule of Many

486   Functions.   *Annual   Review   of   Biochemistry*   **1999**,   *68*   (1),   89–125.

487   https://doi.org/10.1146/annurev.biochem.68.1.89.

488   (4)   Oehmen, A.; Lemos, P. C.; Carvalho, G.; Yuan, Z.; Keller, J.; Blackall, L. L.; Reis, M. A. M.

489   Advances in Enhanced Biological Phosphorus Removal: From Micro to Macro Scale. *Water*

490   *Research* **2007**, *41* (11), 2271–2300. https://doi.org/10.1016/j.watres.2007.02.030.

491   (5)   Seviour, R. J.; Mino, T.; Onuki, M. The Microbiology of Biological Phosphorus Removal in

492   Activated   Sludge   Systems.   *FEMS   Microbiology   Reviews*   **2003**,   *27*   (1),   99–127.

493   https://doi.org/10.1016/S0168-6445(03)00021-4.

494   (6)   Martín, H. G.; Ivanova, N.; Kunin, V.; Warnecke, F.; Barry, K. W.; McHardy, A. C.; Yeates,

495   C.; He, S.; Salamov, A. A.; Szeto, E.; Dalin, E.; Putnam, N. H.; Shapiro, H. J.; Pangilinan, J.

496   L.; Rigoutsos, I.; Kyrpides, N. C.; Blackall, L. L.; McMahon, K. D.; Hugenholtz, P.

497   Metagenomic Analysis of Two Enhanced Biological Phosphorus Removal (EBPR) Sludge

498   Communities. *Nat Biotechnol* **2006**, *24* (10), 1263–1269. https://doi.org/10/dct6rr.

499   (7)   Tobin, K. M.; McGrath, J. W.; Mullan, A.; Quinn, J. P.; O'Connor, K. E. Polyphosphate

500   Accumulation   by   Pseudomonas   Putida   CA-3   and   Other   Medium-Chain-Length

Polyhydroxyalkanoate-Accumulating Bacteria under Aerobic Growth Conditions. *Applied and Environmental Microbiology* **2007**, *73* (4), 1383–1387. https://doi.org/10.1128/AEM.02007-06.

(8) Petriglieri, F.; Singleton, C.; Peces, M.; Petersen, J. F.; Nierychlo, M.; Nielsen, P. H. "Candidatus Dechloromonas Phosphoritropha" and "Ca. D. Phosphorivorans", Novel Polyphosphate Accumulating Organisms Abundant in Wastewater Treatment Systems. *ISME J* **2021**, *15* (12), 3605–3614. https://doi.org/10.1038/s41396-021-01029-2.

(9) Kristiansen, R.; Nguyen, H. T. T.; Saunders, A. M.; Nielsen, J. L.; Wimmer, R.; Le, V. Q.; McIlroy, S. J.; Petrovski, S.; Seviour, R. J.; Calteau, A.; Nielsen, K. L.; Nielsen, P. H. A Metabolic Model for Members of the Genus Tetrasphaera Involved in Enhanced Biological Phosphorus Removal. *ISME J* **2013**, *7* (3), 543–554. https://doi.org/10.1038/ismej.2012.136.

(10) Akbari, A.; Wang, Z.; He, P.; Wang, D.; Lee, J.; Han, I.; Li, G.; Gu, A. Z. Unrevealed Roles of Polyphosphate-accumulating Microorganisms. *Microb Biotechnol* **2021**, *14* (1), 82–87. https://doi.org/10.1111/1751-7915.13730.

(11) Cordell, D.; Rosemarin, A.; Schröder, J. J.; Smit, A. L. Towards Global Phosphorus Security: A Systems Framework for Phosphorus Recovery and Reuse Options. *Chemosphere* **2011**, *84* (6), 747–758. https://doi.org/10/d4mvgw.

(12) Zabel, F.; Putzenlechner, B.; Mauser, W. Global Agricultural Land Resources–a High Resolution Suitability Evaluation and Its Perspectives until 2100 under Climate Change Conditions. *PloS one* **2014**, *9* (9), e107522.

(13) Saia, S. M.; Carrick, H. J.; Buda, A. R.; Regan, J. M.; Walter, M. T. Critical Review of Polyphosphate and Polyphosphate Accumulating Organisms for Agricultural Water Quality

Management. *Environ. Sci. Technol.* **2021**, *55* (5), 2722–2742.

https://doi.org/10.1021/acs.est.0c03566.

(14) Li, G.; Tooker, N. B.; Wang, D.; Srinivasan, V.; Barnard, J. L.; Russell, A.; Stinson, B.;

McQuarrie, J.; Schauer, P.; Menniti, A.; Varga, E.; Hauduc, H.; Takács, I.; Bott, C.;

Dobrowski, P.; Onnis-Hayden, A.; Gu, A. Z. Modeling Versatile and Dynamic Anaerobic

Metabolism for PAOs/GAOs Competition Using Agent-Based Model and Verification via

Single Cell Raman Micro-Spectroscopy. *Water Research* **2023**, *245*, 120540.

https://doi.org/10/gs5c3w.

(15) Zhao, W.; Bi, X.; Peng, Y.; Bai, M. Research Advances of the Phosphorus-Accumulating

Organisms of Candidatus Accumulibacter, Dechloromonas and Tetrasphaera: Metabolic

Mechanisms, Applications and Influencing Factors. *Chemosphere* **2022**, *307*, 135675.

https://doi.org/10/gq8m45.

(16) Mason-Jones, K.; Banfield, C. C.; Dippold, M. A. Compound-Specific 13C Stable Isotope

Probing Confirms Synthesis of Polyhydroxybutyrate by Soil Bacteria. *Rapid*

Communications in Mass Spectrometry **2019**, *33* (8), 795–802. https://doi.org/10/gr8ctq.

(17) Mason-Jones, K.; Breidenbach, A.; Dyckmans, J.; Banfield, C. C.; Dippold, M. A.

Intracellular Carbon Storage by Microorganisms Is an Overlooked Pathway of Biomass

Growth. *Nat Commun* **2023**, *14* (1), 2240. https://doi.org/10.1038/s41467-023-37713-4.

(18) Butler, O. M.; Manzoni, S.; Warren, C. R. Community Composition and Physiological

Plasticity Control Microbial Carbon Storage across Natural and Experimental Soil Fertility

Gradients. *ISME J* **2023**, *17* (12), 2259–2269. https://doi.org/10.1038/s41396-023-01527-5.

(19) Ratcliff, W. C.; Kadam, S. V.; Denison, R. F. Poly-3-Hydroxybutyrate (PHB) Supports

Survival and Reproduction in Starving Rhizobia: PHB Increases Rhizobium Fitness. *FEMS*

Microbiology Ecology **2008**, *65* (3), 391–399. https://doi.org/10.1111/j.1574-
6941.2008.00544.x.

(20) Rao, N.; Kornberg, A. Inorganic Polyphosphate Regulates Responses of Escherichia Coli to
Nutritional Stringencies, Environmental Stresses and Survival in the Stationary Phase.
Inorganic polyphosphates: Biochemistry, biology, biotechnology **1999**, 183–195.

(21) Gray, M. J.; Jakob, U. Oxidative Stress Protection by Polyphosphate — New Roles for an
Old Player. *Current Opinion in Microbiology* **2015**, *24*, 1–6.
https://doi.org/10.1016/j.mib.2014.12.004.

(22) Alam, M. M.; Srinivasan, V.; Mueller, A. V.; Gu, A. Z. Status and Advances in Technologies
for Phosphorus Species Detection and Characterization in Natural Environment- A
Comprehensive Review. *Talanta* **2021**, *233*, 122458.
https://doi.org/10.1016/j.talanta.2021.122458.

(23) Li, Y.; Cope, H. A.; Rahman, S. M.; Li, G.; Nielsen, P. H.; Elfick, A.; Gu, A. Z. Toward
Better Understanding of EBPR Systems via Linking Raman-Based Phenotypic Profiling with
Phylogenetic Diversity. *Environmental Science and Technology* **2018**, *52* (15), 8596–8606.
https://doi.org/10.1021/acs.est.8b01388.

(24) Majed, N.; Li, Y.; Gu, A. Z. Advances in Techniques for Phosphorus Analysis in Biological
Sources. *Current Opinion in Biotechnology* **2012**, *23* (6), 852–859.
https://doi.org/10.1016/j.copbio.2012.06.002.

(25) He, Z.; Deng, Y.; Van Nostrand, J. D.; Tu, Q.; Xu, M.; Hemme, C. L.; Li, X.; Wu, L.; Gentry,
T. J.; Yin, Y.; Liebich, J.; Hazen, T. C.; Zhou, J. GeoChip 3.0 as a High-Throughput Tool for
Analyzing Microbial Community Composition, Structure and Functional Activity. *The ISME*
Journal **2010**, *4* (9), 1167–1179. https://doi.org/10.1038/ismej.2010.46.

(26) Tu, Q.; Yu, H.; He, Z.; Deng, Y.; Wu, L.; Van Nostrand, J. D.; Zhou, A.; Voordeckers, J.;

Lee, Y.; Qin, Y. GeoChip 4: A Functional Gene-array-based High-throughput Environmental

Technology for Microbial Community Analysis. *Molecular ecology resources* **2014**, *14* (5),

914–928.

(27) Kornberg, A. Inorganic Polyphosphate: Toward Making a Forgotten Polymer Unforgettable.

Journal of Bacteriology **1995**, *177* (3), 491–496. https://doi.org/10.1128/jb.177.3.491-

496.1995.

(28) Kulaev, I. S.; Vagabov, V.; Kulakovskaya, T. *The Biochemistry of Inorganic Polyphosphates*,

2 nd.; John Wiley & Sons, 2005.

(29) Whitehead, M. P.; Eagles, L.; Hooley, P.; Brown, M. R. W. Most Bacteria Synthesize

Polyphosphate by Unknown Mechanisms. *Microbiology* **2014**, *160* (5), 829–831.

https://doi.org/10.1099/mic.0.075366-0.

(30) Günther, S.; Trutnau, M.; Kleinsteuber, S.; Hause, G.; Bley, T.; Röske, I.; Harms, H.; Müller,

S. Dynamics of Polyphosphate-Accumulating Bacteria in Wastewater Treatment Plant

Microbial Communities Detected via DAPI (4′,6′-Diamidino-2-Phenylindole) and

Tetracycline Labeling. *Applied and Environmental Microbiology* **2009**, *75* (7), 2111–2121.

https://doi.org/10/c6h3z8.

(31) Terashima, M.; Kamagata, Y.; Kato, S. Rapid Enrichment and Isolation of Polyphosphate-

Accumulating Organisms Through 4'6-Diamidino-2-Phenylindole (DAPI) Staining With

Fluorescence-Activated Cell Sorting (FACS). *Front. Microbiol.* **2020**, *11*.

https://doi.org/10.3389/fmicb.2020.00793.

(32) Wang, D.; He, P.; Wang, Z.; Li, G.; Majed, N.; Gu, A. Z. Advances in Single Cell Raman

Spectroscopy Technologies for Biological and Environmental Applications. *Current Opinion*

in Biotechnology **2020**, *64*, 218–229. https://doi.org/10.1016/j.copbio.2020.06.011.

(33) Majed, N.; Chernenko, T.; Diem, M.; Gu, A. Z. Identification of Functionally Relevant

Populations in Enhanced Biological Phosphorus Removal Processes Based on Intracellular

Polymers Profiles and Insights into the Metabolic Diversity and Heterogeneity.

Environmental Science and Technology **2012**, *46* (9), 5010–5017.

https://doi.org/10.1021/es300044h.

(34) Majed, N.; Matthäus, C.; Diem, M.; Gu, A. Z. Evaluation of Intracellular Polyphosphate

Dynamics in Enhanced Biological Phosphorus Removal Process Using Raman Microscopy.

Environmental Science and Technology **2009**, *43* (14), 5436–5442.

https://doi.org/10.1021/es900251n.

(35) Fernando, E. Y.; McIlroy, S. J.; Nierychlo, M.; Herbst, F.-A.; Petriglieri, F.; Schmid, M. C.;

Wagner, M.; Nielsen, J. L.; Nielsen, P. H. Resolving the Individual Contribution of Key

Microbial Populations to Enhanced Biological Phosphorus Removal with Raman–FISH.

ISME J **2019**, *13* (8), 1933–1946. https://doi.org/10/gn7s5g.

(36) Hanway, J. J. Growth Stages of Corn (Zea Mays, L.)1. *Agronomy Journal* **1963**, *55* (5), 487–

492. https://doi.org/10.2134/agronj1963.00021962005500050024x.

(37) Deng, L.; Fiskal, A.; Han, X.; Dubois, N.; Bernasconi, S. M.; Lever, M. A. Improving the

Accuracy of Flow Cytometric Quantification of Microbial Populations in Sediments:

Importance of Cell Staining Procedures. *Frontiers in Microbiology* **2019**, *10* (April), 1–13.

https://doi.org/10.3389/fmicb.2019.00720.

(38) Eschenhagen, M.; Müller, S.; Günther, S.; Harms, H.; Hübschmann, T.; Röske, I.; Rudolf, M.

Fixation Procedures for Flow Cytometric Analysis of Environmental Bacteria. *Journal of*

Microbiological Methods **2008**, *75* (1), 127–134.

https://doi.org/10.1016/j.mimet.2008.05.017.

(39) Günther, S.; Trutnau, M.; Kleinsteuber, S.; Hause, G.; Bley, T.; Röske, I.; Harms, H.; Müller,

S.; Gunther, S.; Kleinsteuber, S.; Trutnau, M.; Bley, T.; Muller, S.; Hause, G.; Roske, I.;

Harms, H.; Gunther, S.; Bley, T.; Muller, S.; Hause, G.; Roske, I.; Harms, H. Dynamics of

Polyphosphate-Accumulating Bacteria in Wastewater Treatment Plant Microbial

Communities Detected via DAPI (4′,6′-Diamidino-2- Phenylindole) and Tetracycline

Labeling. *Applied and Environmental Microbiology* **2009**, *75* (7), 2111–2121.

https://doi.org/10.1128/AEM.01540-08.

(40) Cao, J.-S.; Xu, R.-Z.; Luo, J.-Y.; Feng, Q.; Fang, F. Rapid Quantification of Intracellular

Polyhydroxyalkanoates via Fluorescence Techniques: A Critical Review. *Bioresource*

Technology **2022**, *350*, 126906. https://doi.org/10/gstqvm.

(41) Kawaharasaki, M.; Manome, A.; Kanagawa, T.; Nakamura, K. Flow Cytometric Sorting and

RFLP Analysis of Phosphate Accumulating Bacteria in an Enhanced Biological Phosphorus

Removal System. *Water Science and Technology* **2002**, *46* (1–2), 139–144.

https://doi.org/10.2166/wst.2002.0469.

(42) Vidal-Mas, J.; Resina-Pelfort, O.; Haba, E.; Comas, J.; Manresa, A.; Vives-Rego, J. Rapid

Flow Cytometry - Nile Red Assessment of PHA Cellular Content and Heterogeneity in

Cultures of Pseudomonas Aeruginosa 47T2 (NCIB 40044) Grown in Waste Frying Oil.

Antonie van Leeuwenhoek, International Journal of General and Molecular Microbiology

2001, *80* (1), 57–63. https://doi.org/10.1023/A:1012208225286.

(43) Zuriani, R.; Vigneswari, S.; Azizan, M. N. M.; Majid, M. I. A.; Amirul, A. A. A High

Throughput Nile Red Fluorescence Method for Rapid Quantification of Intracellular

Bacterial Polyhydroxyalkanoates. *Biotechnology and Bioprocess Engineering* **2013**, *18* (3),

472–478. https://doi.org/10/gpdfn2.

(44) Zuriani, R.; Vigneswari, S.; Azizan, M. N. M.; Majid, M. I. A.; Amirul, A. A. A High

Throughput Nile Red Fluorescence Method for Rapid Quantification of Intracellular

Bacterial Polyhydroxyalkanoates. *Biotechnology and Bioprocess Engineering* **2013**, *18* (3),

472–478. https://doi.org/10.1007/s12257-012-0607-z.

(45) Metz, S.; Dos Santos, A. L.; Berman, M. C.; Bigeard, E.; Licursi, M.; Not, F.; Lara, E.; Unrein,

F.; Lopes dos Santos, A.; Berman, M. C.; Bigeard, E.; Dos Santos, A. L.; Berman, M. C.;

Bigeard, E.; Licursi, M.; Not, F.; Lara, E.; Unrein, F. Diversity of Photosynthetic

Picoeukaryotes in Eutrophic Shallow Lakes as Assessed by Combining Flow Cytometry Cell-

Sorting and High Throughput Sequencing. *FEMS Microbiology Ecology* **2019**, *95* (5), 1–12.

https://doi.org/10.1093/femsec/fiz038.

(46) Daims, H. Use of Fluorescence in Situ Hybridization and the Daime Image Analysis Program

for the Cultivation-Independent Quantification of Microorganisms in Environmental and

Medical Samples. *Cold Spring Harbor Protocols* **2009**, *4* (7), 1–8.

https://doi.org/10.1101/pdb.prot5253.

(47) Li, G.; Wu, C.; Wang, D.; Srinivasan, V.; Dy, J. G.; Gu, A. Z. Machine Learning-Based

Determination of Sampling Depth for Complex Environmental Systems: Case Study with

Single-Cell Raman Spectroscopy Data in EBPR Systems. *Environ. Sci. Technol.* **2022**, *56*

(18), 13473–13484. https://doi.org/10.1021/acs.est.1c08768.

(48) Wang, D.; Li, Y.; Cope, H. A.; Li, X.; He, P.; Liu, C.; Li, G.; Rahman, S. M.; Tooker, N. B.;

Bott, C. B.; Onnis-Hayden, A.; Singh, J.; Elfick, A.; Marques, R.; Jessen, H. J.; Oehmen, A.;

Gu, A. Z. Intracellular Polyphosphate Length Characterization in Polyphosphate

Accumulating Microorganisms (PAOs): Implications in PAO Phenotypic Diversity and

Enhanced Biological Phosphorus Removal Performance. *Water Research* **2021**, *206*, 117726.

https://doi.org/10.1016/j.watres.2021.117726.

(49) Majed, N.; Matthäus, C.; Diem, M.; Gu, A. Z. Evaluation of Intracellular Polyphosphate

Dynamics in Enhanced Biological Phosphorus Removal Process Using Raman Microscopy.

Environmental science & technology **2009**, *43* (14), 5436–5442.

(50) Li, Y.; Rahman, S. M.; Li, G.; Fowle, W.; Nielsen, P. H.; Gu, A. Z. The Composition and

Implications of Polyphosphate-Metal in Enhanced Biological Phosphorus Removal Systems.

Environ. Sci. Technol. **2019**, *53* (3), 1536–1544. https://doi.org/10.1021/acs.est.8b06827.

(51) Günther, S.; Hübschmann, T.; Rudolf, M.; Eschenhagen, M.; Röske, I.; Harms, H.; Müller,

S. Fixation Procedures for Flow Cytometric Analysis of Environmental Bacteria. *Journal of*

Microbiological Methods **2008**, *75* (1), 127–134. https://doi.org/10/bq9dsz.

(52) Fernando, E. Y.; McIlroy, S. J.; Nierychlo, M.; Herbst, F.-A.; Petriglieri, F.; Schmid, M. C.;

Wagner, M.; Nielsen, J. L.; Nielsen, P. H. Resolving the Individual Contribution of Key

Microbial Populations to Enhanced Biological Phosphorus Removal with Raman–FISH. *The*

ISME Journal **2019**, *13* (8), 1933–1946. https://doi.org/10/gn7s5g.

(53) Chavan, S.; Yadav, B.; Tyagi, R. D.; Drogui, P. A Review on Production of

Polyhydroxyalkanoate (PHA) Biopolyesters by Thermophilic Microbes Using Waste

Feedstocks. *Bioresource Technology* **2021**, *341*, 125900. https://doi.org/10/gr8kr7.

(54) Chang, C. F.; Shuman, H.; Somlyo, A. P. Electron Probe Analysis, X-Ray Mapping, and

Electron Energy-Loss Spectroscopy of Calcium, Magnesium, and Monovalent Ions in Log-

Phase and in Dividing Escherichia Coli B Cells. *J Bacteriol* **1986**, *167* (3), 935–939.

https://doi.org/10.1128/jb.167.3.935-939.1986.

(55) Smith, R. J. Calcium and Bacteria. *Adv Microb Physiol* **1995**, *37*, 83–133.

https://doi.org/10.1016/s0065-2911(08)60144-7.

(56) Günther, S.; Trutnau, M.; Kleinsteuber, S.; Hause, G.; Bley, T.; Röske, I.; Harms, H.; Müller,

S. Dynamics of Polyphosphate-Accumulating Bacteria in Wastewater Treatment Plant

Microbial Communities Detected via DAPI (4′,6′-Diamidino-2-Phenylindole) and

Tetracycline Labeling. *Applied and Environmental Microbiology* **2009**, *75* (7), 2111–2121.

https://doi.org/10.1128/AEM.01540-08.

(57) Barra Caracciolo, A.; Grenni, P.; Cupo, C.; Rossetti, S. In Situ Analysis of Native Microbial

Communities in Complex Samples with High Particulate Loads. *FEMS Microbiology Letters*

2005, *253* (1), 55–58. https://doi.org/10.1016/j.femsle.2005.09.018.

(58) Kamika, I.; Coetzee, M.; Mamba, B. B.; Msagati, T.; Momba, M. N. B. The Impact of

Microbial Ecology and Chemical Profile on the Enhanced Biological Phosphorus Removal

(EBPR) Process: A Case Study of Northern Wastewater Treatment Works, Johannesburg.

International Journal of Environmental Research and Public Health **2014**, *11* (3), 2876–

2898. https://doi.org/10.3390/ijerph110302876.

(59) Nielsen, P. H.; McIlroy, S. J.; Albertsen, M.; Nierychlo, M. Re-Evaluating the Microbiology

of the Enhanced Biological Phosphorus Removal Process. *Current Opinion in Biotechnology*

2019, *57* (Figure 1), 111–118. https://doi.org/10.1016/j.copbio.2019.03.008.

(60) Stokholm-Bjerregaard, M.; McIlroy, S. J.; Nierychlo, M.; Karst, S. M.; Albertsen, M.;

Nielsen, P. H. A Critical Assessment of the Microorganisms Proposed to Be Important to

Enhanced Biological Phosphorus Removal in Full-Scale Wastewater Treatment Systems.

Frontiers in Microbiology **2017**, *8* (APR), 1–18. https://doi.org/10.3389/fmicb.2017.00718.

(61) Bai, Y.; Zhang, Y.; Quan, X.; Chen, S. Nutrient Removal Performance and Microbial

Characteristics of a Full-Scale IFAS-EBPR Process Treating Municipal Wastewater. *Water*

Science and Technology **2016**, *73* (6), 1261–1268. https://doi.org/10.2166/wst.2015.604.

(62) Close, K.; Marques, R.; Carvalho, V. C. F.; Freitas, E. B.; Reis, M. A. M.; Carvalho, G.;

Oehmen, A. The Storage Compounds Associated with Tetrasphaera PAO Metabolism and

the Relationship between Diversity and P Removal. *Water Research* **2021**, *204*, 117621.

https://doi.org/10/gsmsnj.

(63) Kristiansen, R.; Nguyen, H. T. T.; Saunders, A. M.; Nielsen, J. L.; Wimmer, R.; Le, V. Q.;

McIlroy, S. J.; Petrovski, S.; Seviour, R. J.; Calteau, A.; Nielsen, K. L.; Nielsen, P. H. A

Metabolic Model for Members of the Genus Tetrasphaera Involved in Enhanced Biological

Phosphorus   Removal.   *ISME   Journal*   **2013**,   *7*   (3),   543–554.

https://doi.org/10.1038/ismej.2012.136.

(64) Marques, R.; Santos, J.; Nguyen, H.; Carvalho, G.; Noronha, J. P.; Nielsen, P. H.; Reis, M.

A. M.; Oehmen, A. Metabolism and Ecological Niche of Tetrasphaera and Ca.

Accumulibacter in Enhanced Biological Phosphorus Removal. *Water Research* **2017**, *122*,

159–171. https://doi.org/10.1016/j.watres.2017.04.072.

(65) McIlroy, S. J.; Onetto, C. A.; McIlroy, B.; Herbst, F. A.; Dueholm, M. S.; Kirkegaard, R. H.;

Fernando, E.; Karst, S. M.; Nierychlo, M.; Kristensen, J. M.; Eales, K. L.; Grbin, P. R.;

Wimmer, R.; Nielsen, P. H. Genomic and in Situ Analyses Reveal the Micropruina Spp. as

Abundant Fermentative Glycogen Accumulating Organisms in Enhanced Biological

Phosphorus Removal Systems. *Frontiers in Microbiology* **2018**, *9* (MAY), 1–12.

https://doi.org/10.3389/fmicb.2018.01004.

(66) Singleton, C. M.; Petriglieri, F.; Wasmund, K.; Nierychlo, M.; Kondrotaite, Z.; Petersen, J.

F.; Peces, M.; Dueholm, M. S.; Wagner, M.; Nielsen, P. H. The Novel Genus, 'Candidatus

Phosphoribacter', Previously Identified as Tetrasphaera, Is the Dominant Polyphosphate

Accumulating Lineage in EBPR Wastewater Treatment Plants Worldwide. *ISME Journal*

2022, *16* (6), 1605–1616. https://doi.org/10.1038/s41396-022-01212-z.

(67) Pankratov, T. A.; Grouzdev, D. S.; Patutina, E. O.; Kolganova, T. V.; Suzina, N. E.;

Berestovskaya, J. J. Lichenibacterium Ramalinae Gen. Nov, Sp. Nov., Lichenibacterium

Minor Sp. Nov., the First Endophytic, Beta-Carotene Producing Bacterial Representatives

from Lichen Thalli and the Proposal of the New Family Lichenibacteriaceae within the Order

Rhizobiales. *Antonie van Leeuwenhoek* **2020**, *113* (4), 477–489. https://doi.org/10/gstz8p.

(68) Wang, H.-G.; Huang, H.; Liu, R.-L.; Mao, Y.-P.; Biswal, B. K.; Chen, G.-H.; Wu, D.

Investigation on Polyphosphate Accumulation in the Sulfur Transformation-Centric EBPR

(SEBPR) Process for Treatment of High-Temperature Saline Wastewater. *Water Research*

2019, *167*, 115138. https://doi.org/10/gmngmh.

(69) Bressan, M.; Trinsoutrot Gattin, I.; Desaire, S.; Castel, L.; Gangneux, C.; Laval, K. A Rapid

Flow Cytometry Method to Assess Bacterial Abundance in Agricultural Soil. *Applied Soil*

Ecology **2015**, *88*, 60–68. https://doi.org/10.1016/j.apsoil.2014.12.007.

(70) Frossard, A.; Hammes, F.; Gessner, M. O. Flow Cytometric Assessment of Bacterial

Abundance in Soils, Sediments and Sludge. *Front. Microbiol.* **2016**, *7*.

https://doi.org/10.3389/fmicb.2016.00903.

(71) Sutermaster, B. A.; Darling, E. M. Considerations for High-Yield, High-Throughput Cell Enrichment: Fluorescence versus Magnetic Sorting. *Sci Rep* **2019**, *9* (1), 227. https://doi.org/10.1038/s41598-018-36698-1.

(72) Ansari, F. A.; Jafri, H.; Ahmad, I.; Abulreesh, H. H. Factors Affecting Biofilm Formation in in Vitro and in the Rhizosphere. In *Biofilms in Plant and Soil Health*; John Wiley & Sons, Ltd, 2017; pp 275–290. https://doi.org/10.1002/9781119246329.ch15.

(73) Lima, D. C.; Aviles, A. C.; Alpers, R. T.; Perkins, A.; Schoemaker, D. L.; Costa, M.; Michel, K. J.; Kaeppler, S.; Ertl, D.; Romay, M. C.; Gage, J. L.; Holland, J.; Beissinger, T.; Bohn, M.; Buckler, E.; Edwards, J.; Flint-Garcia, S.; Gore, M. A.; Hirsch, C. N.; Knoll, J. E.; McKay, J.; Minyo, R.; Murray, S. C.; Schnable, J.; Sekhon, R. S.; Singh, M. P.; Sparks, E. E.; Thomison, P.; Thompson, A.; Tuinstra, M.; Wallace, J.; Washburn, J. D.; Weldekidan, T.; Xu, W.; de Leon, N. 2020-2021 Field Seasons of Maize GxE Project within the Genomes to Fields Initiative. *BMC Res Notes* **2023**, *16* (1), 219. https://doi.org/10.1186/s13104-023-06430-y.

(74) Kurashita, H.; Hatamoto, M.; Tomita, S.; Yamaguchi, T.; Narihiro, T.; Kuroda, K. Comparative Metagenomic Analysis Provides Novel Insights into the Ecophysiology of Myxococcota in Activated Sludge Systems. bioRxiv April 28, 2024, p 2024.04.26.591422. https://doi.org/10.1101/2024.04.26.591422.

(75) Zhou, T.; Xiang, Y.; Liu, S.; Ma, H.; Shao, Z.; He, Q.; Chai, H. Microbial Community Dynamics and Metagenomics Reveal the Potential Role of Unconventional Functional Microorganisms in Nitrogen and Phosphorus Removal Biofilm System. *Science of The Total Environment* **2023**, *905*, 167194. https://doi.org/10.1016/j.scitotenv.2023.167194.

(76) Nguyen, H. T. T.; Nielsen, J. L.; Nielsen, P. H. 'Candidatus Halomonas Phosphatis', a Novel

Polyphosphate-Accumulating Organism in Full-Scale Enhanced Biological Phosphorus

Removal Plants. *Environmental Microbiology* **2012**, *14* (10), 2826–2837.

https://doi.org/10.1111/j.1462-2920.2012.02826.x.

(77) Stokholm-Bjerregaard, M.; McIlroy, S. J.; Nierychlo, M.; Karst, S. M.; Albertsen, M.;

Nielsen, P. H. A Critical Assessment of the Microorganisms Proposed to Be Important to

Enhanced Biological Phosphorus Removal in Full-Scale Wastewater Treatment Systems.

Frontiers in Microbiology **2017**, *8*. https://doi.org/10.3389/fmicb.2017.00718.

(78) Akone, S. H.; Hug, J. J.; Kaur, A.; Garcia, R.; Müller, R. Structure Elucidation and

Biosynthesis of Nannosterols A and B, Myxobacterial Sterols from Nannocystis Sp.

MNa10993. *Journal of Natural Products* **2023**, *86* (4), 915.

https://doi.org/10.1021/acs.jnatprod.2c01143.

(79) Pérez-García, L.-A.; Sáenz-Mata, J.; Fortis-Hernández, M.; Navarro-Muñoz, C. E.; Palacio-

Rodríguez, R.; Preciado-Rangel, P. Plant-Growth-Promoting Rhizobacteria Improve

Germination and Bioactive Compounds in Cucumber Seedlings. *Agronomy* **2023**, *13* (2), 315.

https://doi.org/10.3390/agronomy13020315.

(80) Kumar Arora, N.; Fatima, T.; Mishra, J.; Mishra, I.; Verma, S.; Verma, R.; Verma, M.;

Bhattacharya, A.; Verma, P.; Mishra, P.; Bharti, C. Halo-Tolerant Plant Growth Promoting

Rhizobacteria for Improving Productivity and Remediation of Saline Soils. *Journal of*

Advanced Research **2020**, *26*, 69–82. https://doi.org/10.1016/j.jare.2020.07.003.

(81) Dai, W.; Liu, M.; Wang, N.; Ye, X.; Liu, Y.; Yao, D.; Wang, L.; Cui, Z.; Yan, P.; Cheng, C.;

Huang, Z.; Wang, H. Positive Contribution of Predatory Bacterial Community to Multiple

Nutrient Cycling and Microbial Network Complexity in Arsenic-Contaminated Soils.

Applied Soil Ecology **2023**, *185*, 104792. https://doi.org/10.1016/j.apsoil.2022.104792.

793