Genetic underpinnings of chills from art and music

Giacomo Bignardi[1,2*], Danielle Admiraal[1], Else Eising[1†], & Simon E. Fisher[1,3†]

[1] *Language and Genetics Department, Max Planck Institute for Psycholinguistics, Nijmegen, 6525, XD, the Netherlands;*
[2] *Max Planck School of Cognition, Leipzig, Germany;*
[3] *Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, 6500, HB, the Netherlands;*

† These authors jointly supervised this work.
* Corresponding author
Email: giacomo.bignardi@mpi.nl;

Author Contributions: G.B. conceptualised the work; E.E. & S.E.F. supervised the research; G.B. performed primary analyses; E.E. and D.A. performed secondary analyses and quality control; G.B. created (supporting) figures; G.B. drafted the manuscript; E.E., D.A. & S.E.F. contributed to revising the manuscript.

Competing Interest Statement: The authors declare no competing interest.

Keywords: Aesthetics, Genetics, Chills, Art, Music, Heritability.

Abstract

Art can evoke strong emotional responses in humans. Here, we examine genetic contributions to chills, a marker of such responses. We gather self-reports from a genotyped sample of thousands of partly related individuals from the Netherlands (N =15,606). Using genomic relationships based on common single-nucleotide polymorphism (SNP) data, we find that up to 29% of the variation in proneness to aesthetic (visual art and poetry) and music chills can be explained by familial relatedness effects, one-fourth of which is attributed to SNP variation. Furthermore, we reveal a moderate genetic correlation of .58 between aesthetic and music chills, pointing to shared genetic variation affecting susceptibility to strong emotional responses across different art forms. Finally, we find that a polygenic index (PGI) capturing openness to experiences, including propensities towards art (N = 234,880), is associated with susceptibilities both to aesthetic and music chills. Our results show that additive genetic variation, but also familial relatedness beyond shared common SNPs, contributes to proneness to chills from artistic, poetic, and musical expressions. These results open up a promising path towards studying the human attitude towards art, via both state-of-the-art genomics and intergenerational models of transmission.

Author Summary

Many people experience chills when listening to music, reading poetry, or viewing art. Yet not everyone feels these reactions in the same way. These differences provide a window into how our brains and bodies respond to art, revealing individual variation in emotional experiences. To investigate what drives these differences, we analysed data from over 15,000 participants with available genetic information, examining whether DNA variation could help explain why some people are more prone to these intense responses. We estimated that roughly 30% of the variation in chills is linked to family-related factors, of which about one fourth was attributable to common DNA variants. Some genetic influences appear to be shared across music, poetry, and art, and are associated with individual openness to experience and general artistic interests, while others may be unique to each domain. These results suggest that genetics contributes to how strongly people respond to cultural experiences and pave the way for future studies on the genetics of sensitivity to art and music experiences.

Introduction

Darwin experienced them while hearing the anthem in King's College Chapel, which gave him "intense pleasure so that [his] backbone would sometimes shiver"(1). Nabokov elevated them as "the telltale tingle" needed to bask in the book of genius(2). When people encounter art, they may experience intense physiological responses that coincide with peak subjective pleasure and emotions, described as "chills"(3). Since chills are clear, measurable events(4) that link subjective responses with physiological manifestations(4–7), they serve as a model for studying responses to art(3,6–11). Investigations of chills have revealed that the pleasure evoked by music and poetry recruits neural systems similar to those involved in the evaluation of other biologically relevant non-abstract stimuli(3,7), and have linked stable subjective differences to inter-individual variability at both physiological and neurobiological levels(8,10,12).

Here, we investigate the genetics behind individual differences in proneness to chills, which we suggest can offer an additional approach for empirically studying how humans respond to art. As with many other complex human traits(13), proneness to chills is partly heritable(14). For example, across diverse cultures, proneness to chills from visual art and poetry (referred to as aesthetic chills) has moderate heritability (h^2). Genetic effects have been estimated to account for 36 to 43% of the variance in large twin-based samples—including more than 10,000 individuals from the Netherlands(14), and thousands from Germany, Denmark, North America, Australia, and Japan(15)—with the remaining variance attributed to residual effects, such as environmental influences and measurement error, with no evidence for environmental effects shared by family members(14). However, key questions about the genetics of inter-individual differences in proneness to chills remain unanswered. First, to date, h^2 estimates have been derived from family studies only, mainly from those relying on the Classical Twin Design (CTD). As such, it is unclear whether previous results reflect molecular genetic effects or biases introduced by unmet assumptions inherent to the CTD. Second, there is a complete lack of evidence regarding whether genetic effects on aesthetic chills may extend to other artistic domains, such as music, since the only estimates derived so far are specific to chills from visual art and poetry. This gap raises additional questions of whether sensitivity to different art forms is shaped by entirely different processes or is tied to similar underlying biological mechanisms. Finally, it is still unclear whether the genetics of proneness to chills is indeed an index of broader predispositions towards art, or if it reflects an independent phenomenon.

To resolve these uncertainties, we used a large sample of more than fifteen thousand genotyped individuals from Lifelines, a large multi-generational cohort study of individuals living in the North of the Netherlands(16), from whom we gathered self-reports on their proneness to aesthetic and music chills. We applied a molecular approach to quantify h^2 that contrasts patterns of measured genotypic resemblance with phenotypic resemblance in unrelated and related individuals(17,18). The strength of this approach relies on the possibility of partitioning family-based (also called pedigree-based) estimates for h^2 (h_{ped}^2) into two types of h^2. The first, obtained from unrelated and distantly related individuals and referred to as the Single-Nucleotide Polymorphism heritability (h_{SNP}^2), estimates the additive effect of genetic variants common in the population under study, captured using DNA genotyping chips.The second, obtained instead from related individuals, assesses additional additive effects of common genetic variants untagged by genotyping chips, as well as biases due to, for instance, non-additive genetic and environmental effects shared between members of families(19,20). The gap between such h_{SNP}^2 and h_{ped}^2 estimates is sometimes referred to as the (still) missing heritability (21). Moreover, by collecting information on people's proneness to chills evoked by different art modalities, we could investigate similarities and differences between responses to diverse forms of artistic, poetic and musical expression. We accomplished this by estimating genetic correlations (r_g) between aesthetic and music chills. Finally, we investigated the degree to which proneness to chills might reflect a broader genetic predisposition towards art. To do so, we leveraged the largest published genome-wide association study (GWAS) on personality to date(22) to construct a openness to experience polygenic

105   index (PGI), mostly related to artistic interests and active imagination, testing their association with
106   proneness to chills in our target sample.
107
108   **Results**
109
Working with Lifelines(16), we gathered self-report data on proneness to chills in response to diverse
art forms from 35,114 adults (21,249 women). Some 15,615 of these individuals (9,907 women) had
genotype data available, and their ages covered the entire adult lifespan, ranging from 18 to 96 years
old. Proneness to aesthetic and music chills was assessed using the self-reported items "Sometimes
when I am reading poetry or looking at a work of art, I feel a chill or wave of excitement"(23) and "I
sometimes feel chills when I hear a melody that I like"(24), respectively. Participants were asked to
report how much they agreed with these statements by choosing one out of five response options
ranging from "strongly disagree" to "strongly agree". Within-trait descriptive statistics and between-trait
phenotypic similarities and differences are reported in Supporting Information.
119
To assess the extent to which genetic influences account for variation in proneness to aesthetic and
music chills, we applied a linear mixed model with a Genome-based Restricted Maximum Likelihood
(GREML) estimator(18). This method estimates the amount of the observed phenotypic variance that
can be explained by genomic relatedness between individuals, as indexed by SNP-derived Genetic
Relatedness Matrices (GRM). We focused our analysis on a total sample of 15,606 individuals,
removing a total of 9 individuals form pairs with estimated genomic relatedness (π) larger then .90 (i.e.,
identical twins). Taking advantage of the family relatedness structure of the participants included in the
Lifelines cohort, we constructed two complementary SNP-derived GRM, one representative of the
pairwise genomic relatedness (π) across all 15,606 individuals and another for which π below a set
threshold (i.e., $\pi < .05$, the recommended genetic relatedness threshold for GREML-based analysis in
family data(19)) was adjusted to zero(17). Using this approach, also known as threshold GREML(25),
we could partition h_{ped}^2 into h_{SNP}^2 and $h_{\pi \geq .05}^2$, the latter assessing effects beyond common SNP variation
tagged by genotyping array. As shown in Table 1, GREML-based analysis of adjusted two-stage rank
normalised and residualised data for age, sex, genotyping array, and ten genomic principal components
(PC) indicates significant h_{SNP}^2 for aesthetic ($\chi^2(1) = 6.30$, $p = .006$) and music ($\chi^2(1) = 9.25$, $p = .001$)
chills, with h_{ped}^2 explaining up to 29% of the total variance in proneness to chills (see Table 1). These
findings indicate that the proneness to chills from art, poetry and music is influenced by genetic
variation. Results obtained from raw, transformed data, and by using a different threshold ($\pi < .02$), did
not differ (see Supplementary Table S1 and Table S2).
139
Since both aesthetic and music chills displayed significant h_{SNP}^2, we assessed evidence for shared
genetic contributions to variability in the two traits. First, we estimated the phenotypic correlation
between proneness to aesthetic and music chills. To obtain a more precise estimate, we utilised the full
Lifelines sample of 35,114 individuals with available proneness to chills data. Proneness to aesthetic
and music chills (raw scores) correlated modestly, with a r of .43 (95% Confidence Interval (CI) [.42,
.44]). Then, we applied a bivariate extension of GREML to estimate the genetic correlation between the
proneness to aesthetic and music chills in the remaining genotyped sample of 15,606 individuals. We
found a moderate genetic correlation, with an r_g of .58 (95% CI [.20, .95], $\chi^2(1) = 4.61$, $p = .02$; which
was also significantly lower than 1, $\chi^2(1) = 4.17$, $p = .02$, both one-tailed test). These results suggest
that a substantial proportion of the genetic variation associated with proneness to aesthetic chills is also
associated with music chills. Yet, they also reveal some degree of specificity in genetic contributions to
the different traits.
152
153   Finally, we assessed whether genetic proneness to chills could in part be reflected by broader genetic
154   predispositions towards art. We derived individual PGI from the largest published genome-wide
155   association study of openness to experience, which was previously conducted in a fully independent
156   sample of 234,880 individuals(22). To construct the PGI, we used SNP associations with openness to
157   experience, assessing, in particular, artistic interests and active imagination (see Methods). We found

that PGI were significantly associated with proneness to aesthetic ($\beta = .05$, $p = 4.79 \times 10^{-10}$) and music chills ($\beta = .03$, $p = 1.97 \times 10^{-6}$), explaining 0.3% (r_{PGI}^2 95% CI [0.01%,0.4%]) and 0.1% (r_{PGI}^2 95% CI [0.0%, 0.3%]) of the overall variance, respectively. Similar results were obtained in a subset of 10,703 distantly related and unrelated individuals ($\pi < .05$; See Supporting Information). Further analyses provided no evidence for heterogeneous PGI associations with either aesthetic or music chills ($Q_{SNP} = 1.95$, $p = .10$), suggesting that PGI explained about 0.5% (95% CI [0.2%, 0.8%]) of the overall covariance between the two traits (see Methods and Supporting Information). Thus, genetic variation associated with individual differences in art interest contributes to proneness to chills across art domains. Table 1 provides a comprehensive overview of the results of these different investigations.

We note that assortative and cross-assortative mating can lead to bias when estimating h_{SNP}^2 and r_g. Assortative and cross-assortative mating refer to phenomena where two individuals are more likely to mate when they resemble each other in one or more (e.g., cross) traits. These phenomena have been shown to confound h_{SNP}^2 and r_g by upwardly biasing estimates[26,27]. To provide evidence of minimal upward bias on h_{SNP}^2, and help solidify the interpretation of r_g as an estimate for widespread pleiotropy unconfounded by both types of assortment, we estimated trait and cross-trait partner correlations in proneness to chills. To do so, we studied 3424 partners for whom phenotypic data were available. We observed small yet significant partner correlations for aesthetic chills, $r = .09$ (95% CI [.06,.12]), and music chills, $r = .10$ (95% CI [.07,.13]). There was no evidence of cross-trait partner correlation, with a non-significant correlation of only .02 (95% CI [-.03,.04]). These findings indicate that assortative and cross-assortative mating do not largely bias our estimates for h_{SNP}^2 and r_g (for details, see Supporting Information and reference[26]).

Table 1. *Overview of estimates.*

Component of interest	Parameter	Trait	Estimate	95% CI
Trait variance explained by tagged common DNA variation (with additive effect)	h_{SNP}^2	Aesthetic chills	.06	[.01, .10]
		Music chills	.07	[.03, .11]
Gap between effects of common DNA variation and familial relatedness	$h_{\pi \geq .05}^2$	Aesthetic chills	.18	[.10, .27]
		Music chills	.23	[.14, .31]
Trait variance explained by overall effects of familial relatedness	h_{ped}^2	Aesthetic chills	.24	[.17, .32]
		Music chills	.29	[.22, .36]
Phenotypic correlation	r	Aesthetic-music chills	.43	[.42, .44]
Genetic correlation	r_g	Aesthetic-music chills	.58	[.20, .95]
Percentage of variance explained by PGI for openness to experience	r_{PGI}^2	Aesthetic chills	0.3%	[0.1%, 0.4%]
		Music chills	0.1%	[0.0%, 0.3%]
Partner phenotypic correlation	r_{AM}	Aesthetic chills	.09	[.05, .12]
		Music chills	.09	[.06, .13]
Cross-trait partner phenotypic correlation	r_{xAM}	Aesthetic-music chills	.01	[-.03, .04]

Note. h_{SNP}^2: GREML-SNP-based heritability; $h_{\pi \geq .05}$: GREML-based excess heritability in related individuals; h_{ped}^2: GREML pedigree-based heritability; r: Pearson-correlation; r_g: GREML-SNP-based additive genetic correlation; PGI: Polygenic Index; r_{AM} within-trait Pearson-correlation between partners; r_{xAM} cross-trait Pearson-correlation between partners. All results shown here were obtained from fully adjusted two-stage rank normalisation procedure (rank transformed) on the phenotype data, with

exception of phenotypic correlations, which used non-adjusted rank transformed scores. 95%
Confidence Intervals (CI) for heritability and genetic correlation estimates are derived from the Standard
Errors (i.e. 95% CI = estimate ± 1.96*SE).

Discussion

Our results provide converging evidence for the existence of a genetic predisposition to the ability to
experience chills from cultural products, such as artistic, poetic, and musical expression. By
investigating the similarity between genotypic and phenotypic resemblance in related and unrelated
individuals, we find significant SNP-based heritability for aesthetic and music chills, aligning with the
claim that proneness to chills may represent a "vector of biological variation"(28). Consistent with other
studies(14,15), the substantial proportion of variance attributable to residual effects—including
environmental influences, measurement error, but also stochastic developmental processes unique to
each individual—indicates that factors beyond genetic variation also contribute to individual differences
in proneness to chills. Furthermore, we reveal a substantial genetic correlation between the traits, which
indicates that proneness to chills from different art modalities is partly tied to shared genetic variation.
In line with these findings, we also find that genetic propensities for openness to experience, including
artistic interests, are significantly associated with proneness to aesthetic and music chills. At the same
time, genetic correlations between propensity to aesthetic and music chills are significantly lower than
1, indicating less than full overlap in the underlying genetic architecture for responses to different forms
of artistic expression.

206
Notwithstanding the significant SNP-based heritability, this explained only around a fourth of the total
pedigree heritability and a fifth of the previously reported twin-based heritability for aesthetic chills
(obtained from another sample of the Dutch population(14)). While in line with well-established concepts
of "missing heritability"(29), as SNP-based estimates tend to explain only one-third to one-half of family-
based estimates of complex human traits(20,21,30), the gap observed here for proneness to chills
seems relatively large. (Note, however, that this is still smaller than the gap observed for some other
behavioural characteristics, for which h_{SNP}^2 is only about one-tenth of the h_{twin}^2; see reference(30) for
an example). The observed gap between SNP- and family-based heritability estimates likely involves a
complex mixture of factors. These include contributions of imperfect tagging of causal variants from
genotyping arrays, such as rare variants, non-additive genetic effects, such as dominance deviations
or higher-order epistasis (gene-by-gene interaction at different genetic loci), and common
environmental effects shared across families(17). Additional contributors may include unequal
environmental sharing among relatives and interactions between genes and shared environmental
factors(31).

221
It is worth noting here that while h_{SNP}^2 estimates can be considered the upper bound of the additive
genetic influences that genome-wide association studies (GWAS) could capture on the same
genotyping platform, they are generally expected to underestimate the true heritability of the trait in the
population under study. This discrepancy arises because the GRM imperfectly captures the contribution
of ungenotyped variants among unrelated or distantly related individuals. By contrast, when the same
GRM is obtained from pairs of individuals above a certain threshold of relatedness (e.g., $\pi > .05$), the
estimates tend to be closer to the true heritability. This is because, at higher levels of relatedness (larger
values of π), the GRM better approximate the genetic covariance across the genome. Therefore, a gap
between h_{SNP}^2 and h_{ped}^2 is expected(17). At the same time, the h_{ped}^2 estimates derived from related
individuals can be upwardly biased if shared environmental effects, non-additive genetic effects, or
effects of rare variants are present. Previous CTD studies on proneness to aesthetic chills found no
evidence of shared environmental effects(14,15), with correlations among monozygotic twins exceeding
twice those of dizygotic twins(14). These findings, under the assumptions of identical shared
environmental influences in monozygotic and dizygotic twins and no gene-by-shared environment
interaction, suggest that non-additive genetic effects, rather than shared environmental effects, may
partly contribute to the gap between h_{SNP}^2 and h_{ped}^2 as found in this study. However, an alternative

explanation, which cannot be ruled out in the current study, is also possible: gene-by-shared
environment interactions. If gene-by-shared environment interactions were to influence phenotypic
variation, then both the current h_{ped}^2 and the previously reported h_{twin}^2 would yield upwardly biased
estimates of h^2. Furthermore, although we noted that the observed weak correlation between partners
was not sufficient to largely bias h_{SNP}^2 estimates in this study, it is still possible that assortment may
have previously upwardly biased h_{twin}^2 estimates (as these were obtained from a constrained ADE
model, which is expected to provide inflated h_{twin}^2 estimates under assortative mating[33]). Therefore,
assortative mating may still play a role in the gap observed between the estimates provided in this study
and the ones reported from previous CTD studies. Future studies with different research paradigms will
help to better understand this gap. In particular, more sophisticated twin-based approaches, such as
extended family twin designs (in which parents, partners, and siblings of twins are included[33,34]),
may be particularly suited to partly explain the gap observed in this study.

In addition, our study reveals the association between proneness to chills and polygenic indices for
openness to experience—mainly assessed as individuals' interest towards the arts and self-reported
active imagination (see below and Methods). We note that the percentage of variance in proneness to
aesthetic and music chills explained by the PGI is less than 1%. However, the modest proportion of
variance accounted for here is in line with theory-driven expectations on PGI-based predictions. In
particular, it is known that the expected percentage of phenotypic variance that can be explained by a
PGI is a function of the heritability of the target phenotype being predicted, the one from which the PGI
is derived, and their genetic correlation, as well as the sample size of the original GWAS, along with the
effective number of SNPs included[35,36]. Considering the observed h_{SNP}^2 of proneness to chills and
the previously reported heritability of openness to experience[22] ($h_{SNP}^2 = .05$), then the maximum
expected percentage of variance that we could have explained is ~1% (see Supporting information) for
aesthetic and music chills, respectively, even assuming an unlikely perfect genetic correlation of 1.
These expectations are not far from our observed r^2, thus contextualising our estimates for the absolute
percentage of variance explained by the PGI as being relatively close to theoretical expectations.

This study comes with a few limitations. First, our PGI analyses made use of summary statistics from
meta-GWAS efforts conducted with self-reports that included art interest and other broader
characteristics defining openness to experience[22]. As such, individual PGI may assess more complex
genetic predispositions than art interest alone. However, we note that the largest cohort included in the
meta-analysis, including 220,015 individuals from the Million Veteran Program (comprising 94% of the
combined samples), was conducted on the sum score of only two self-report items, one assessing art
interests and the other evaluating active imagination. As such, we expect the genetic signal used to
construct the PGI to reflect individual traits that revolve around art interest and imagination over other
aspects of openness to experience. Second, the results in the present study were obtained from a
population representative of the Netherlands and focused on individuals of European descent. Further
studies should assess the generalisability of our findings beyond ancestries of European descent and
culture. Indeed, the cross-cultural prevalence of experiencing chills[37] and proneness towards
these[23], and the observation that such proneness may rely less on cultural bounds and influences
from Western humanistic traditions than other traits[23,28], leads to intriguing questions for future
research. In particular, it will be interesting to ask whether proneness to chills may be used to investigate
the genetics of human attitudes towards art across cultures. Moderating effects introduced by cultural
or generational differences in genetic influences could be an especially promising avenue of study[38].
Third, our results do not provide insights into how molecular associations with proneness to chills may
extend to biological mechanisms underlying related physiological processes, such as piloerection (i.e.,
hairs standing on end). Previous research suggests that piloerection tends to co-occur with the
experience of chills[39], although this is not always the case[40]. At the same time, the findings of
genetic associations with proneness to chills may allow future research to better disambiguate between
these two seemingly tied phenomena. Fourth, sampling bias in the Lifelines cohort may affect heritability
and genetic correlation estimates. While Lifelines represents the northern Netherlands well, women,

middle-aged individuals, and individuals with higher educational attainment tend to be slightly overrepresented(41). Since proneness to chills is linked to both demographics and education(42), this may introduce bias in genetic estimates. However, evidence from other biobanks suggests that heritability is typically only modestly underestimated under such bias(43), with downstream analyses being more strongly affected(44). Therefore, genome-wide association studies may be more susceptible to this type of bias than the present study. Future work should address this issue, for example, by applying weighting to genetic associations based on census data on sex and educational attainment, to better align the sample estimates to the effects in the overall population(45). Fifth, the items used to assess proneness to chills are likely subject to measurement error. For instance, the test-retest over a 5-year timespan for aesthetic chills in adults has been previously estimated to range between .58 and .61, in a comparable but not overlapping Dutch sample(14). As such, our h^2 estimates are likely downwardly biased by intra-individual variance (including measurement error), and therefore should be considered conservative lower bounds for the additive effects of SNP variation.

In conclusion, our research highlights shared molecular heritability for proneness to chills from art, poetry, and music. Furthermore, it reveals that proneness to chills is associated with a PGI for openness to experience, partly indexing genetic propensities towards art. Yet it also reveals larger estimates for genetic effects on proneness to chills in related individuals, hinting at deviations from pure additive genetic transmission of sensitivity to art and underscoring the need for extended intergenerational designs to relax classic modelling assumptions. We anticipate that larger-scale GWAS of proneness to chills, combined with extended family designs, could further illuminate the origins and biological bases of affective responses to art. Such studies would enable the exploration of both molecular and cultural processes and may even clarify whether the etiology of these subjective experiences differs from that of other sensory experiences(46).

Materials and Methods

Ethics
The general Lifelines protocol has been approved by the University Medical Center Groningen medical ethical committee (2007/152), and the "Speech, Language and Musicality" study was reviewed by the medical ethical committee of the Erasmus Medical Center (MEC-2022-0313). All participants gave written informed consent prior to taking part in the survey. More information on how informed consent was obtained can be found at https://wiki.lifelines.nl/doku.php?id=informed_consent.

Sample
Individuals included here participated in the Lifelines study "Speech, Language and Musicality". Lifelines is a multi-disciplinary prospective population-based cohort study examining in a unique three-generation design the health and health-related behaviours of 167,729 persons living in the North of the Netherlands. It employs a broad range of investigative procedures in assessing the biomedical, socio-demographic, behavioural, physical, and psychological factors which contribute to the health and disease of the general population, with a special focus on multi-morbidity and complex genetics(16). The study population included in Lifeline is broadly representative of the adult population of the north of the Netherlands. Participants in Lifelines were invited by their General Practitioner (GP) to be part of the original study. Family members of individual participants, including partners, were invited by the participants themselves to take part in the study. Family relations data in Lifelines come from municipal registries. The UMCG Genetics Lifelines Initiative (UGLI) consortium refined these using anonymised surname data and family composition questionnaires, then optimised them with genetic data. Couples living together without children were also registered as partners within the same family. Documentation of the reconstruction and quality control process is available at https://wiki-lifelines.web.rug.nl/doku.php?id=family_relations. Partner information was based on baseline data (i.e., between 2007 and 2013). Lifelines had no specific selection criteria for exclusion, but GPs were asked

to decide which individuals to invite based on a list of possible criteria provided by Lifelines. The list can
be found at https://wiki-lifelines.web.rug.nl/doku.php?id=cohort.

Between September and October 2022, approximately 115,000 adult Lifelines participants were asked
to participate online in the "Speech, Language and Musicality" study, of whom 35,179 completed the
questionnaire. The response rate observed here (~30%) is consistent with prior musicality research in
national registries. For instance, studies of music, art, and cultural engagement in the Swedish Twin
Registry reported comparable rates of participation of 29 to 36%[47]. It is also worth noting that previous
work found no evidence of sampling bias for music reward sensitivity, a construct that encompasses
proneness to chills.

Phenotyping
Aesthetic and music chills were measured using validated items in the revised NEO personality
inventory (NEO PI-R[23]) and the Barcelona Music Reward Questionnaire (BMRQ[8]). The
questionnaires, which were administered online, were collected in Dutch. The sample of participants for
which we had data available on both items was 35,114 individuals. This included a subsample of 15,615
individuals for whom we had genome-wide SNP genotype data available. From this subset, we
additionally removed 9 individuals from pairs with $\pi > .90$, resulting in a final sample of 15,606
individuals. The sample stratified by π can be found in Supplementary Table 3.

Genotyping
Genotyping was carried out by the using the Infinium Global Screening Array for 9251 individuals
(release 1, version 2) and the FinnGen Thermo Fisher Axiom for 6364 individuals (release 2, version
2). Information about quality control of genotyped data and samples, genetic imputation, and genomic
PC calculations can be found at https://wiki.lifelines.nl/doku.php?id=ugli. For the individual genotypes
obtained using the FinnGen Thermo Fisher Axiom array (release 2, version 2), filtered non-imputed
genotyping data was pruned and then used to calculate principal components using PLINK (version
1.9). Further information can be found in the Supporting Information file.

Covariates
Following De Hoyos et al.[48], data were transformed before model fitting using a fully adjusted two-
stage rank normalisation procedure[49]. Age, sex, genotyping array, and the first ten genomic PC were
the covariates of interest. Results from untransformed raw data are included in Supplementary Table
S1.

Phenotypic analysis
Given that the full sample comprised both related and unrelated individuals, the standard errors (SE)
and, by extension, the 95% confidence intervals (CI) surrounding the estimated correlation coefficient
are expected to be downwardly biased[50]. To address this limitation and obtain robust estimates of
uncertainty for the point estimate, individuals were clustered within families using the cluster argument
in the sem() function of the lavaan[51] R package (i.e., sem(..., cluster = "FAM_ID")).
More information on how Lifelines assigns individuals' family IDs can be found at https://wiki-
lifelines.web.rug.nl/doku.php?id=family_relations.

Genome-based Restricted Maximum Likelihood
Estimates for h_{SNP}^2, $h_{\pi \geq .05}^2$, and h_{ped}^2 of aesthetic and music chills were obtained by using a threshold
GREML approach[17,52] as implemented in the GCTA software
(https://cnsgenomics.com/software/gcta/), version 1.93.2beta[18]. The threshold for the GRM was set
based on the recommended value of .05 for double threshold GREML approach[19]. Estimates
obtained using a more stringent threshold of $\pi < .02$, following[20], are provided in Supplementary
Table 2. Similarly, a bivariate extension of GREML[53] was used to obtain estimates of the covariances
between the variance components of the two phenotypes, which were then used to obtain estimates

for r_g. To test whether h_{SNP}^2 and r_g estimates significantly differ from 0 and 1, GCTA applies a Likelihood
Ratio Test comparing the model in which h_{SNP}^2 and r_g are freely estimated to a model in which h_{SNP}^2
and r_g are fixed to 0 and 1, respectively. To control for two comparisons, we used a more stringent α of
.025. Schematic representations of the GREML approaches are given in Supplementary Figure S1. The
extended method section can be found in the Supporting Information file. Robustness analyses with
meta-analytic estimates obtained from subsamples corresponding to the two UGLI genotyping releases
are provided in Supplementary Table S4. We note that, despite $h_{\pi \geq .05}^2$ being equal to $h_{ped}^2 - h_{SNP}^2$, $h_{\pi \geq .05}^2$
and h_{ped}^2 are sometimes used interchangeably in the literature to describe pedigree-based heritability.
The latter nomenclature is appropriate when $h_{\pi \geq .05}^2$ and h_{SNP}^2 are not estimated simultaneously, as in
reference[20].
403
Polygenic Index analyses
We computed individuals' Polygenic Index (PGI) for openness to experience using meta-analysis
summary statistics from a GWAS meta-analysis that totalled 234,880 individuals. The largest GWAS
cohort in this meta-analysis included 220,015 individuals, for which sum scores were derived from only
two self-reports: "I see myself as someone who has few artistic interests" (reverse coded) and "I see
myself as someone who has an active imagination" (five response options, "disagree strongly" to "agree
strongly"). The remaining GWAS cohorts that contributed to the meta-analyses included summary
statistics from openness to experience scores derived from 12 to 48 items(54). The meta-analytic
summary statistics were filtered, removing ambiguous, duplicated and rare (MAF < 0.01) SNPs. PGI
were calculated using PRS-CS (version Jan 4 2021)(55) in PLINK, using a European linkage
disequilibrium panel from the 1000 Genomes. Genotyping data of individuals with proneness for chills
data remained for PGI analyses. We then regressed proneness to aesthetic or music chills on
standardised PGI. The r_{PGI}^2 estimates were derived from a PGI regression with robust standard error,
as in reference(56). Supplementary analyses with retained unrelated or distantly related individuals,
corresponding to $\pi < .05$ are included in the Supporting Information. The bivariate PGI heterogeneity
statistic, Q_{SNP} (i.e., $\chi^2(1)$), as well as the amount of covariance explained by the PGI, was obtained via
an adapted bivariate PGI Common Pathway Model(57,58). More details are provided in the Supporting
Information file.

References

1. Darwin C, Barlow N, Darwin C. The autobiography of Charles Darwin, 1809 - 1882: with original omissions restored. New York: Norton; 1993. 253 p.

2. Nabokov V. Lectures on Literature. Boston: Houghton Mifflin Harcourt; 2002. 1 p.

3. Blood AJ, Zatorre RJ. Intensely pleasurable responses to music correlate with activity in brain regions implicated in reward and emotion. Proc Natl Acad Sci. 2001 Sep 25;98(20):11818–23.

4. Salimpoor VN, Benovoy M, Longo G, Cooperstock JR, Zatorre RJ. The Rewarding Aspects of Music Listening Are Related to Degree of Emotional Arousal. PLOS ONE. 2009 Oct 16;4(10):e7487.

5. Grewe O, Nagel F, Kopiez R, Altenmüller E. Emotions over time: Synchronicity and development of subjective, physiological, and facial affective reactions to music. Emotion. 2007;7(4):774–88.

6. Salimpoor VN, Benovoy M, Larcher K, Dagher A, Zatorre RJ. Anatomically distinct dopamine release during anticipation and experience of peak emotion to music. Nat Neurosci. 2011 Feb;14(2):257–62.

7. Wassiliwizky E, Koelsch S, Wagner V, Jacobsen T, Menninghaus W. The emotional power of poetry: neural circuitry, psychophysiology and compositional principles. Soc Cogn Affect Neurosci. 2017 01;12(8):1229–40.

8. Mas-Herrero E, Zatorre RJ, Rodriguez-Fornells A, Marco-Pallarés J. Dissociation between Musical and Monetary Reward Responses in Specific Musical Anhedonia. Curr Biol. 2014 Mar 17;24(6):699–704.

9. Zatorre RJ, Salimpoor VN. From perception to pleasure: Music and its neural substrates. Proc Natl Acad Sci. 2013 Jun 18;110(supplement_2):10430–7.

10. Martínez-Molina N, Mas-Herrero E, Rodríguez-Fornells A, Zatorre RJ, Marco-Pallarés J. Neural correlates of specific musical anhedonia. Proc Natl Acad Sci. 2016 Nov 15;113(46):E7337–45.

11. Wassiliwizky E, Menninghaus W. Why and How Should Cognitive Science Care about Aesthetics? Trends Cogn Sci. 2021 Jun 1;25(6):437–49.

12. Williams PG, Johnson KT, Curtis BJ, King JB, Anderson JS. Individual differences in aesthetic engagement are reflected in resting-state fMRI connectivity: Implications for stress resilience. NeuroImage. 2018 Oct;179:156–65.

13. Polderman TJC, Benyamin B, de Leeuw CA, Sullivan PF, van Bochoven A, Visscher PM, et al. Meta-analysis of the heritability of human traits based on fifty years of twin studies. Nat Genet. 2015 Jul;47(7):702–9.

14. Bignardi G, Chamberlain R, Kevenaar ST, Tamimy Z, Boomsma DI. On the etiology of aesthetic chills: a behavioral genetic study. Sci Rep. 2022 Feb 28;12(1):3247.

15. Mõttus R, Sinick J, Terracciano A, Hřebíčková M, Kandler C, Ando J, et al. Personality
characteristics below facets: A replication and meta-analysis of cross-rater agreement, rank-
order stability, heritability, and utility of personality nuances. J Pers Soc Psychol. 2019
Oct;117(4):e35–50.

16. Sijtsma A, Rienks J, van der Harst P, Navis G, Rosmalen JGM, Dotinga A. Cohort Profile
Update: Lifelines, a three-generation cohort study and biobank. Int J Epidemiol. 2022 Oct
1;51(5):e295–302.

17. Zaitlen N, Kraft P, Patterson N, Pasaniuc B, Bhatia G, Pollack S, et al. Using Extended
Genealogy to Estimate Components of Heritability for 23 Quantitative and Dichotomous
Traits. PLOS Genet. 2013 May 30;9(5):e1003520.

18. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: A Tool for Genome-wide Complex Trait
Analysis. Am J Hum Genet. 2011 Jan 7;88(1):76–82.

19. Yang J, Zeng J, Goddard ME, Wray NR, Visscher PM. Concepts, estimation and
interpretation of SNP-based heritability. Nat Genet. 2017 Sep;49(9):1304–10.

20. Kemper KE, Yengo L, Zheng Z, Abdellaoui A, Keller MC, Goddard ME, et al. Phenotypic
covariance across the entire spectrum of relatedness for 86 billion pairs of individuals. Nat
Commun. 2021 Feb 16;12(1):1050.

21. Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, et al. Finding the
missing heritability of complex diseases. Nature. 2009 Oct;461(7265):747–53.

22. Gupta P, Galimberti M, Liu Y, Beck S, Wingo A, Wingo T, et al. A genome-wide investigation
into the underlying genetic architecture of personality traits and overlap with
psychopathology. Nat Hum Behav. 2024 Nov;8(11):2235–49.

23. McCrae RR. Aesthetic Chills as a Universal Marker of Openness to Experience. Motiv Emot.
2007 Mar 1;31(1):5–11.

24. Mas-Herrero E, Marco-Pallares J, Lorenzo-Seva U, Zatorre RJ, Rodriguez-Fornells A.
Individual Differences in Music Reward Experiences. Music Percept. 2013 Dec 1;31(2):118–
38.

25. Hagenbeek FA, Hirzinger JS, Breunig S, Bruins S, Kuznetsov DV, Schut K, et al. Maximizing
the value of twin studies in health and behaviour. Nat Hum Behav. 2023 May 15;1–12.

26. Border R, O'Rourke S, de Candia T, Goddard ME, Visscher PM, Yengo L, et al. Assortative
mating biases marker-based heritability estimators. Nat Commun. 2022 Feb 3;13(1):660.

27. Border R, Athanasiadis G, Buil A, Schork AJ, Cai N, Young AI, et al. Cross-trait assortative
mating is widespread and inflates genetic correlation estimates. Science. 2022 Nov
18;378(6621):754–61.

28. Schwaba T, Thalmayer AG. Openness/intellect: The weirdest trait in the Big Five world?
[Internet]. OSF; 2024 [cited 2024 Dec 26]. Available from: https://osf.io/wnzy4

29. Zuk O, Hechter E, Sunyaev SR, Lander ES. The mystery of missing heritability: Genetic
interactions create phantom heritability. Proc Natl Acad Sci. 2012 Jan 24;109(4):1193–8.

30. Cheesman R, Selzam S, Ronald A, Dale PS, McAdams TA, Eley TC, et al. Childhood
behaviour problems show the greatest gap between DNA-based and twin heritability. Transl
Psychiatry. 2017 Dec 12;7(12):1–9.

31. Keller MC, Coventry WL. Quantifying and addressing parameter indeterminacy in the
classical twin design. Twin Res Hum Genet Off J Int Soc Twin Stud. 2005 Jun;8(3):201–13.

32. Wainschtein P, Jain D, Zheng Z, Cupples LA, Shadyab AH, McKnight B, et al. Assessing the
contribution of rare variants to complex trait heritability from whole-genome sequence data.
Nat Genet. 2022 Mar;54(3):263–73.

33. Keller MC, Medland SE, Duncan LE, Hatemi PK, Neale MC, Maes HHM, et al. Modeling
Extended Twin Family Data I: Description of the Cascade Model. Twin Res Hum Genet.
2009 Feb;12(1):8–18.

34. Keller MC, Medland SE, Duncan LE. Are Extended Twin Family Designs Worth the Trouble?
A Comparison of the Bias, Precision, and Accuracy of Parameters Estimated in Four Twin
Family Models. Behav Genet. 2010 May 1;40(3):377–93.

35. Vlaming R de, Okbay A, Rietveld CA, Johannesson M, Magnusson PKE, Uitterlinden AG, et
al. Meta-GWAS Accuracy and Power (MetaGAP) Calculator Shows that Hiding Heritability Is
Partially Due to Imperfect Genetic Correlations across Studies. PLOS Genet. 2017 Jan
17;13(1):e1006495.

36. Okbay A, Wu Y, Wang N, Jayashankar H, Bennett M, Nehzati SM, et al. Polygenic prediction
of educational attainment within and between families from genome-wide association
analyses in 3 million individuals. Nat Genet. 2022 Apr;54(4):437–49.

37. Beier EJ, Janata P, Hulbert JC, Ferreira F. Do you chill when I chill? A cross-cultural study of
strong emotional responses to music. Psychol Aesthet Creat Arts. 2022;16(1):74–96.

38. Uchiyama R, Spicer R, Muthukrishna M. Cultural evolution of genetic heritability. Behav
Brain Sci. 2022 Jan;45:e152.

39. de Fleurian R, Pearce MT. Chills in music: A systematic review. Psychol Bull.
2021;147(9):890–920.

40. McPhetres J, Zickfeld JH. The physiological study of emotional piloerection: A systematic
review and guide for future research. Int J Psychophysiol. 2022 Sep 1;179:6–20.

41. Klijs B, Scholtens S, Mandemakers JJ, Snieder H, Stolk RP, Smidt N. Representativeness of
the LifeLines Cohort Study. PLOS ONE. 2015 Sep 2;10(9):e0137203.

42. Schoeller F, Christov-Moore L, Lynch C, Diot T, Reggente N. Predicting individual
differences in peak emotional response. PNAS Nexus. 2024 Mar;3(3):pgae066.

43. Song S, Benonisdottir S, Liu JS, Kong A. Participation bias in the estimation of heritability
and genetic correlation. Proc Natl Acad Sci. 2025 Jun 24;122(25):e2425530122.

44. Schoeler T, Speed D, Porcu E, Pirastu N, Pingault JB, Kutalik Z. Participation bias in the UK
Biobank distorts genetic associations and downstream analyses. Nat Hum Behav. 2023
Jul;7(7):1216–27.

45. van Alten S, Domingue BW, Faul J, Galama T, Marees AT. Correcting for volunteer bias in
GWAS increases SNP effect sizes and heritability estimates. Nat Commun. 2025 Apr
15;16(1):3578.

46. Nadal M, Skov M. The sensory valuation account of aesthetic experience. Nat Rev Psychol.
2024 Nov 28;1–15.

47. Bignardi G, Wesseldijk LW, Mas-Herrero E, Zatorre RJ, Ullén F, Fisher SE, et al. Twin
modelling reveals partly distinct genetic pathways to music enjoyment. Nat Commun. 2025
Mar 25;16(1):2904.

48. de Hoyos L, Barendse MT, Schlag F, van Donkelaar MMJ, Verhoef E, Shapland CY, et al.
Structural models of genome-wide covariance identify multiple common dimensions in
autism. Nat Commun. 2024 Feb 27;15(1):1770.

49. Sofer T, Zheng X, Gogarten SM, Laurie CA, Grinde K, Shaffer JR, et al. A fully adjusted two-
stage procedure for rank-normalization in genetic association studies. Genet Epidemiol.
2019 Apr;43(3):263–75.

50. Rebollo I, de Moor MHM, Dolan CV, Boomsma DI. Phenotypic factor analysis of family data:
correction of the bias due to dependency. Twin Res Hum Genet Off J Int Soc Twin Stud.
2006 Jun;9(3):367–76.

51. Rosseel Y. lavaan: An R Package for Structural Equation Modeling. J Stat Softw. 2012 May
24;48(1):1–36.

52. Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, et al. Common SNPs
explain a large proportion of the heritability for human height. Nat Genet. 2010
Jul;42(7):565–9.

53. Deary IJ, Yang J, Davies G, Harris SE, Tenesa A, Liewald D, et al. Genetic contributions to
stability and change in intelligence from childhood to old age. Nature. 2012
Feb;482(7384):212–5.

54. de Moor MHM, Costa PT, Terracciano A, Krueger RF, de Geus EJC, Toshiko T, et al. Meta-
analysis of genome-wide association studies for personality. Mol Psychiatry. 2012
Mar;17(3):337–49.

55. Ge T, Chen CY, Ni Y, Feng YCA, Smoller JW. Polygenic prediction via Bayesian regression
and continuous shrinkage priors. Nat Commun. 2019 Apr 16;10(1):1776.

56. Gustavson DE, Borriello GA, Karhadkar MA, Rhee SH, Corley RP, Rhea SA, et al. Stability
of general cognitive ability from infancy to adulthood: A combined twin and genomic
investigation. Proc Natl Acad Sci. 2025 May 27;122(21):e2426531122.

57. Grotzinger AD, Mallard TT, Akingbuwa WA, Ip HF, Adams MJ, Lewis CM, et al. Genetic
architecture of 11 major psychiatric disorders at biobehavioral, functional genomic and
molecular genetic levels of analysis. Nat Genet. 2022 May;54(5):548–59.

58. Alagöz G, Eising E, Mekki Y, Bignardi G, Fontanillas P, Nivard MG, et al. The shared genetic
architecture and evolution of human language and musical rhythm. Nat Hum Behav. 2025
Feb;9(2):376–90.

Acknowledgements

The Lifelines initiative has been made possible by subsidy from the Dutch Ministry of Health, Welfare and Sport, the Dutch Ministry of Economic Affairs, the University Medical Center Groningen (UMCG), Groningen University and the Provinces in the North of the Netherlands (Drenthe, Friesland, Groningen). The authors wish to acknowledge the services of the Lifelines Cohort Study, the contributing research centres delivering data to Lifelines, and all the study participants. We further wish to thank MacKenzie D. Trupp for feedback on an earlier version of the manuscript, Lucia De Hoyos for sharing the R script to apply the fully adjusted two-stage rank normalisation procedure, and Henkjan Honing for assisting with the Dutch translation of the music chill item. G.B. was supported by the German Federal Ministry of Education and Research (BMBF) and the Max Planck School of Cognition; G.B., D.A., E.E. and S.E.F. were supported by the Max Planck Society. E.E. is also supported by a Veni grant from the Dutch Research Council (NWO; VI.Veni.202.072).