Neural dynamics of induced throat vibrations during vocal emotion recognition

Damien Benis, PhD (a,b)*

Garance Selosse MSc (a, b)*

Jennifer Greilsamer, MSc (c)

Ben Meuleman, PhD (b)

Didier Grandjean, PhD[†] (a, b)

Leonardo Ceravolo, PhD[†] (a, b)

[*] Shared first authors

[†] Shared last authors

(a)   *Department of Psychology, University of Geneva, Geneva, Switzerland,*

(b)   *Swiss Center for Affective Sciences, University of Geneva, Geneva, Switzerland,*

(c)   *Department of Medicine, University of Fribourg, Fribourg, Switzerland*

Corresponding author:

Leonardo Ceravolo, PhD
University of Geneva
Boulevard Pont-d'Arve 40
CH-1205 Switzerland
email: leonardo.ceravolo@unige.ch

Abstract

Despite a large corpus of literature in psychological and brain mechanisms on emotional prosody perception, the perspective of embodied cognition in these mechanisms have been largely neglected. Here we investigated the influence of induced bodily vibrations on the categorization of ambiguous emotional vocalizations using event-related potentials (ERPs). Emotional voices were morphed between a fearful expression with the speaker's identity-matching angry expression, creating blends of emotions in each voice. Emotional congruent and incongruent vibrations were delivered on the skin close to the vocal cords. Congruent with our hypotheses, behavioural results revealed that induced vibrations skewed the participants' emotional ratings by biasing responses towards the vibration's emotion. ERPs indicated that N100 and P200 components subtending the early processing of emotional prosody were significantly modulated by induced vibrations in the congruent setting, considered as a facilitation effect for emotion recognition at early stages of processing. A modulation of the late positive component was also observed in the incongruent setting, suggesting an error processing mechanism. Source reconstruction highlighted effects of vibration types in prefrontal, motor, somatosensory, and insular cortices. Our results suggest that voice-associated vibrations may play a significant role in vocal emotion processing and recognition through an embodied mechanism.

Keywords: *voice prosody, emotion, bodily vibrations, feedback, embodiment, electroencephalography*

Word count (excluding abstract, references, tables, legends): 4996

Introduction

Verbal and non-verbal communication is essential for humans. Emotional prosody represents the temporal fluctuations of the pitch, amplitude and spectrum (Grandjean et al., 2006; Laukka et al., 2005; Risberg & Lubker, 1978; Scherer & Oshinsky, 1977). Processing emotional prosody variations allows listeners to subjectively attribute the emotional state of speakers, influenced by context (Brunswik, 1956; Grandjean et al., 2006; Scherer, 2003). One key mechanism is the embodied 'simulation' of what we hear (Hawk et al., 2012), and emotional 'embodiment' is defined as the motor, sensory and perceptual reexperiencing or simulating of the emotion perceived in a peer in the framework of the Simulation of Smiles (SIMS) model (Niedenthal, 2007; Niedenthal et al., 2010). This process has primarily been described in the visual modality, with the perception of facial expressions in others generating facial mimicry in the observer—stronger for emotional vs. non-emotional faces (Chartrand & Bargh, 1999; Dimberg, 1982; Korb et al., 2010; Künecke et al., 2017; Moody & McIntosh, 2011; Moody et al., 2007). Recent studies have supported the idea that facial mimicry represents context-dependent sensorimotor simulation of an observed emotion rather than a mere muscular reproduction of an observed facial expression (Borgomaneri et al., 2020; Drimalla et al., 2019; Hess & Blairy, 2001; Hess & Fischer, 2014; Philip et al., 2018). Importantly, some studies suggest that this process may impact the accurate decoding of facial emotional information especially in women (Hyniewska & Sato, 2015; Korb et al., 2010; Schneider et al., 2013; Stel & van Knippenberg, 2008), although not exclusively (Blairy et al., 1999; Hess & Blairy, 2001; Holland et al., 2021).

While studies on the contribution of embodiment in the field of emotional prosody decoding are scarcer than in the visual domain, some point to a contribution of this mechanism in the auditory modality. Indeed, several studies showed that passive listening of speech signals elicits activations in motor regions of the brain similar to those involved during speech production (Fadiga et al., 2002; Watkins et al., 2003; Wilson et al., 2004). Facilitated language comprehension when the primary motor areas controlling the lips/tongue were stimulated transcranially was also observed (D'Ausilio et al., 2009). One mechanism underlying emotion embodiment in voice signals may be body resonances or vibrations. In fact, vibrations—especially close to the vocal folds—may represent a reliable interoceptive and/or exteroceptive feedback during vocal emotion expression. Indeed, during voice expression, vibrations from the vocal cords are transmitted through the speaker's skin and body tissues (Munger & Thomson, 2008; Švec et al., 2005), with a frequency corresponding

to the fundamental frequency of the produced auditory tone (Sundberg, 1992). These bodily

vibrations would constitute, for the speaker, a dynamic feedback on several characteristics of

his/her own voice (Sundberg, 1992). Lylykangas and colleagues (2009) highlighted the

potential to successfully use vibrotactile feedback as relevant information for speed regulation

messages. Similarly, Tuuri and colleagues (2010) compared audio and tactile feedback from

the same speech stimuli as speed regulation cues and found that the information was delivered

in a similar manner across modalities. However, there is still a paucity of research in this field

and we intend to fill this gap. To investigate precise dynamic neural correlates of the

embodiment of vocal emotion perception concurrent to induced throat vibrations, we used

surface electroencephalography (EEG) using event-related potentials (ERPs) as the

underlying dependent measure.

ERP correlates of emotional prosody decoding have been extensively studied, with

evidence for a three-stage processing (Schirmer & Kotz, 2006). In line with this model, a

wealth of ERP studies has shown the implication of several early and late ERP components in

the processing of emotional vocalizations. Early components underlie auditory-sensory

processing, attention and early recognition of meaning around 100ms after stimulus onset

(N100, frontocentral area). It might reflect early low-level prosody analysis and emotion

discrimination (Gädeke et al., 2013; Paquette et al., 2020; Tarai & Srinivasan, 2019). Basic

emotions are also distinguished in the frontal/parietal P200 component (Duville et al., 2022;

Gädeke et al., 2013; Maltezou-Papastylianou et al., 2022; Paulmann et al., 2013; Paulmann &

Uskul, 2017). The P200 is a positive deflection around 200 milliseconds reflecting increased

attention to emotional stimuli for facilitated processing (Paulmann & Kotz, 2008; Thomas et

al., 2007). Later components are commonly associated with higher-level mechanisms of

emotional prosody processing such as appraisal, contextual relevance and mental

representations (Duville et al., 2022; Pell & Kotz, 2021). This includes ~300-400 milliseconds

deflections (P300/N400) in frontal/frontocentral areas during emotional prosody processing

(Carminati et al., 2018; Proverbio et al., 2020; Tarai & Srinivasan, 2019), underlying deviant

auditory stimuli detection (Lindín et al., 2004; Tsolaki et al., 2015). Finally, the Late Positive

Component (LPC) is a positive deflection observed at a later stage in frontal/frontocentral and

parietal areas, associated with higher-order cognitive processes such as arousal information

(Paulmann et al., 2013), expectancy violation in prosody (Astésano et al., 2004; Brattico et al.,

2006; Paulmann et al., 2012) and emotional 'meaning' (Wei et al., 2022; Zora et al., 2020).

The LPC may reflect attentional processes rather than emotional face processing *per se*

(Ashley et al., 2004), consistent with the attentional capture effect associated with violation or
incongruence (Vachon et al., 2012). Mismatching prosody stimuli also elicits an LPC,
potentially indicating an integration process of the different sources of information (Brattico
et al., 2006; Paulmann et al., 2012).

Concerning functional MRI studies, processing emotional prosody activates
subcortical and cortical networks (Ceravolo et al., 2021; Grandjean, 2021; Peron et al., 2016;
Witteman et al., 2012), including the bilateral superior temporal, inferior frontal (Ceravolo et
al., 2026) and orbitofrontal cortex, cerebellum, etc. Additionally, the motor cortex is also
involved during speech perception (Fadiga et al., 2002; Watkins et al., 2003) while
interoceptive signals trigger insular activity, a region responsible for the integration of internal
and external physiological signals and emotional processes (Adolfi et al., 2017; Casals-
Gutierrez & Abbey, 2020). The somatosensory cortex or the supplementary motor area
(SMA) also play a critical role in interoceptive and proprioceptive processing (Engelen et al.,
2023; Gibson, 2019; Khalsa et al., 2018).

Our study aims to explore the potential interoceptive and/or exteroceptive feedbacks
represented by bodily resonances—induced external throat vibrations, and their impact on
emotional prosody recognition. Revealing such a link between vocal emotion perception and
induced vibrations would constitute convincing evidence for a crucial role of vibrotactile
feedback as a physiological mechanism of vocal emotion embodiment. We recently
demonstrated the neural correlates of measured throat vibrations during emotional voice
expression (Selosse et al., 2025) and our results are coherent with above mentioned
literature—although pertaining to throat vibrations specifically. Associations between
interoception/proprioception and emotional processing were previously studied but never
using vibrations concurrent to expressed voice signals (Burleson & Quigley, 2021; Couto et
al., 2015; Critchley & Garfinkel, 2017). Other related work includes: interoceptive accuracy
(Mendoza-Medialdea & Ruiz-Padial, 2021), ERPs of interoception in emotion processing
(P300 component; Herbert et al., 2007; Pollatos et al., 2005), a somatosensory-specific P300
component (parietal regions) involved in somatosensory perception (Al et al., 2020;
Auksztulewicz & Blankenburg, 2013; Has et al., 2021), interoceptive priming on faces
(Salamone et al., 2021). In addition, other studies further support the idea that emotionally
relevant stimuli can modulate attention and early perceptual processing across modalities
(Brosch, Grandjean, et al., 2008; Brosch, Sander, et al., 2008)—in the case of the present
study, auditory and vibrotactile modalities.

No study has specifically investigated the contribution of throat vibrations to the auditory

perception of emotions using EEG. The present study aims to further explore this mechanism.

The main mechanistic rationale is that vocal cords vibrations, seen as a 'mechanical'

consequence of the production of vocalized signals, would represent a feedback that could

impact the auditory perception and recognition of vocal emotions. Namely, according to an

embodied perception perspective: as a valid feedback, vibratory signals would influence voice

production for the speaker and should therefore also have an impact on perception if induced

in the listener. At the behavioral level, we first postulated that: i) vibrations would affect the

discriminative perceptual processes of emotional vocalizations (more frequent "anger"

response when the vibrations also express anger and likewise for fear) or when compared to

the 'no-vibration' condition; ii) these effects would occur especially for ambiguous stimuli

(i.e., voices containing 40-60% of each emotion), as these would trigger sensory information

seeking for clarification (Niedenthal, 2007; Niedenthal et al., 2010), here induced throat

vibrations. At the neural level, we postulated: i) vibration-specific ERPs between 200 and

300ms modulated by interoceptive/proprioceptive processes (Herbert et al., 2007; Mendoza-

Medialdea & Ruiz-Padial, 2021; Pollatos et al., 2005) with source modulations in the insular,

somatosensory, prefrontal, and cingulate cortices as well as in the SMA/pre-SMA (Adolfi et

al., 2017; Casals-Gutierrez & Abbey, 2020; Engelen et al., 2023; Gibson, 2019; Khalsa et al.,

2018), for anger or fear vibrations compared to no-vibration; ii) incongruence between

emotional voice and vibrations–angry voice with fear vibration and *vice versa*, would trigger

early (100ms and 200-300ms) and late (400-700ms) ERPs with source modulations in the

above-mentioned brain areas, related to error prediction and/or integration of vibratory

feedbacks in the context of emotional prosody information processing.

Method

Participants

Sample size was decided a priori based on repeated measures MANOVA in 'G*Power

3.1' software, with: effect size=.25, power=.8, alpha=.05, number of groups=2 (fixed, no

other option here), number of measures=9 and correlation among repeated measures=0.5

(Faul et al., 2007). This approximation—the closest possible to mixed-effects modeling—

resulted in a required sample size of N=24. We therefore recruited twenty-four healthy

participants (18 females and 6 males; Age=18-41; M_{Age}=23.67; SD_{Age}=5.11). Participants had

normal hearing, no neurologic or psychiatric history and gave informed written consent. The
study was approved by state-wise ethics committee (nr. 2020-02262) and conducted
according to the declaration of Helsinki.

Material

Voices ('Aah', Montreal affective voices database; Belin et al., 2008) were presented
through headphones (HD25, Sennheiser, DE) and morphed on the emotion dimension (anger,
fear) using Matlab (The Mathworks Inc., Natick, MA, USA) and the PsychToolbox (Brainard,
1997; Kleiner et al., 2007; Pelli, 1997). Morphing fearful expression with the speaker's
identity-matching angry expression resulted in five conditions of 10 voices each [fear%-
anger%: 10-90 (A_{90}), 40-60 (A_{60}), 50-50 (A_{50}), 60-40 (A_{40}), and 90-10 (A_{10})], using the
STRAIGHT toolbox (Kawahara & Matsui, 2003). Voice stimuli and vibrations were
acoustically assessed (amplitude, pitch, loudness, duration, spectral centroid and slope, F1
frequency; Supplementary Table 1 & 2, respectively). Morphing conditions and actors were
randomly distributed across trials. The choice of using anger and fear was due to the fact that
these emotions are among the most biologically-relevant for survival.

Throat vibrations were induced using a small vibrator (BC-10, Ortofon, DK;
dimensions: 13.5*29.5*18.0mm, weight: 16.5gr, sensitivity: 118dB, harmonic distortion:
1.5%, impedance: 15Ω, sensibility: 100-1000Hz, sampling rate: 1000Hz). These expressed
pure anger or pure fear in $2/3^{rd}$ of the trials (unmorphed 'Aah'; anger, fear) while no
vibrations were induced in the remaining trials (control, $1/3^{rd}$). The vibrator was placed close
to the vocal cords (Figure 1), taped on the left side to the laryngeal prominence with
kinesiology tape. This location fits our general hypothesis that throat vibrations may be used
as a vibrotactile feedback. It also matches relevant locations in this context (Munger &
Thomson, 2008; Nolan et al., 2009; Sundberg, 1992). The same speaker was used for both
vibrator and voice stimuli for the sake of consistency and to avoid intonation issues. There
were 20 stimuli for the vibrations (two emotions, ten actors).

Task Procedure

The experiment comprised six runs (two fear-induced, two anger-induced vibrations,
two no-vibration) of N=113 randomly presented trials each (Figure 1). Each trial started with
a fixation cross (500ms) concurrent to a voice stimulus (1000ms), then explicit emotion
categorization for 3000ms (buttons: anger, fear; key labels counterbalanced between).
Participants were instructed to answer as fast and as accurately as possible. Vibrations were

induced simultaneously to the voices and were emotionally congruent or incongruent with the

voice stimulus. The final factorial design is a 5 EmotionalVoice (A_{90}, A_{60}, A_{50}, A_{40}, A_{10}) by 3

Vibration (No Vibrations, Anger, Fear) within-subjects design.

219
Figure 1. **Experimental design.** Participants were instructed to focus on a black central
fixation cross displayed on a grey screen for five hundred milliseconds. Emotional voice
stimuli were presented through headphones for one second. Vibrations were induced at the
same time in two thirds of the trials and expressed pure anger or pure fear. After each voice
stimulus, the cross turned white to indicate that the participant had to categorize the emotion
expressed auditorily using the response keys, as fast as accurately as possible and within the
3-seconds response window.

229   **Statistical Analysis**

230   *Trial-level behavioral analysis*

231   Responses were categorical: anger=0, fear=1. A Generalized Linear Mixed Model

(GLMM) was performed in R through Rstudio (Venables & Smith, 2003) to analyze the

responses as a function of Vibration and EmotionalVoice. An LMM was chosen to analyze

reaction times with the same factors (no significant effects and not part of our hypotheses;

Supplementary Table 3). Both models were computed using the 'lme4' package (Bates et al.,

2015). For both models, fixed effects included the interaction between Vibration and

Emotional voice. Random effects included the random slope of Vibration and

EmotionalVoice as a function of participant identity (uncorrelated) and the random intercept

of stimulus identity, participant and speaker gender, runs. The formula was the following

(glmer for response, lmer for reaction times):

Model.response / rt <- glmer / lmer

(response/rt~Vibration*EmotionalVoice+ParticipantGender+SpeakerGender+(1+Vib

ration+EmotionalVoice||ParticipantID)+(1|Run)+(1|StimulusID))

The effect size of each model was computed and labelled according to the thresholds

defined by Cohen (1988). Planned contrasts according to our hypotheses use Bonferroni

correction for multiple testing (Bland & Altman, 1995) and estimated marginal means

package for odds ratios (Lenth et al., 2019).

EEG preprocessing

Surface EEG was acquired with 64 amplified electrodes (Biosemi ActiveThree system,

BioSemi B.V., Netherlands), preprocessing was performed using Brainvision analyser 2.2

(Brain Products, Munich, Germany) and Fieldtrip (Oostenveld et al., 2011). Spline

interpolation was applied to artifacted channels (5.34±4.74% of channels) and an average

reference was applied. Blink and saccade artifacts were rejected using independent

component analysis. Raw data were split into two-seconds epochs (-1s to +1s) centered on

voice onset. Peak-to-peak amplitude was calculated for all epochs across all channels, and a

trial rejection threshold was calculated as the median +1.96 standard deviation of the obtained

peak-to-peak amplitude distribution (~114μv). For each channel, epochs exceeding the

threshold were rejected (average rejection: 9.44±11.95% trials). For each participant/channel,

ERPs were obtained by averaging epochs for each emotional condition and each vibration

type, and their conjunction (anger90-fear10 with anger vibration for example). Grand-average

ERPs were then obtained by averaging the resulting ERPs across participants for each channel

(final average number of trials per participant/condition: 40.95±6.08; Supplementary Table 7).

Topographical clustering methods are reported in the supplementary material

('Supplementary EEG methods', *I.*).

ERP statistical analysis

ERP for each channel, condition, trial was averaged into twenty 50ms bins from stimulus

onset to 1s post-stimulus-onset. A database was created with average power at each of the 20

timebins, 3 Vibration and 5 EmotionalVoice conditions, for each EEG channel for the 4

clusters and electrode placement (left, center, right) as factors. As power followed a linear

distribution, a Bayesian general LMM approach was used to validate the result of

topographical clustering. For each timebin, data was entered into a GLMM to test the

Cluster*EmotionalVoice(A_{90}, A_{60}, A_{50}, A_{40}, A_{10})*Vibration(Anger, Fear, No vibrations)

interaction. Random factors included participant identity and EEG channels.

Cluster significance calculations and source reconstruction methods are reported in the

supplementary material ('Supplementary EEG methods', *II. & III.*).

Results

278   In this study, the participants had to categorize morphed voice emotions concomitant

279   to induced throat vibrations in 2/3rd of the trials. We analyzed the influence of these vibrations

280   on voice recognition responses, ERPs and source modulations in the brain.

Behavioral responses

The main effect of Vibration ($\chi^2(2)=8.25$, $p<.05$, Figure 2**A**), EmotionalVoice

($\chi^2(4)=17.81$, $p<.01$, Supplementary Figure 1) and their interaction ($\chi^2(8)=23.92$, $p< .01$,

Figure 2**B**) were significant. Model effect size was small-to-medium (Fixed effects $R^2=.14$;

Fixed and random effects $R^2=.36$).

The impact of Vibration on responses (Fear vibrations: *m*=.57, *sd*=.50, Anger

vibrations: *m*=.53, *sd*=.50, No vibrations: *m*=.55, *sd*=.50) was tested using hypothesis-driven

planned contrasts: [Vibration Anger>Fear], ($\chi^2(1)=6.86$, $p<.01$, *OR*=0.83); [Vibration

Fear>None], ($\chi^2(1)=4.87$, $p<.05$, *OR*=1.12; [Vibration Anger>None], ($\chi^2(1)=1.68$, $p=.19$,

OR=0.93; Figure 2**A**). These results already partly confirm our first hypothesis (more 'anger'

response for an anger vibration, and similarly for fear).

Vibration*Emotional Voice interaction on responses revealed significant planned

contrasts: [Vibration Anger>Fear]*[Ambiguous (A_{60}, A_{50}, A_{40}) voice stimuli], ($\chi^2(1)=7.99$,

$p<.01$, *OR*=0.51, [Vibration Anger>None]*[Ambiguous (A_{60}, A_{50}, A_{40}) voice stimuli],

($\chi^2(1)=4.21$, $p<.05$, *OR*=0.67, [Vibration Fear>None]*[Ambiguous (A_{60}, A_{50}, A_{40}) voice

stimuli], ($\chi^2(1)=4.20$, $p<.14$, *OR*=1.31; Figure 2**B**). More specific contrasts on the most

ambiguous condition of Emotional Voice were computed: [Vibration

Anger>Fear]*[Ambiguous (A_{50}) voice stimuli], ($\chi2(1)=3.94$, $p<.05$, *OR*=0.81), [Vibration

Anger>None]*[Ambiguous (A_{50}) voice stimuli], ($\chi2(1)=7.08$, $p<.01$, *OR*=0.77), [Vibration

Fear>None]*[Ambiguous (A_{50}) voice stimuli], ($\chi2(1)=0.28$, $p=.59$, *OR*=0.95; Figure 2**B**).

These results also partly confirm our second hypothesis, especially the influence of angry

vibrations on ambiguous voice stimuli. Results for non-ambiguous voices are reported in

Supplementary Table 8, with only an effect of incongruence for fear vibrations on A_{90} voices.

Figure 2. **A. Main effect of Vibration on the probability of fear responses.** On the X axis,
each point represents a Vibration condition (i.e., None = no vibrations, Anger, Fear). **B.**
Interaction between Emotional Voice and Vibration on the probability of fear responses. On
the X axis, each point represents an Emotional Voice condition (i.e., A_{90}, A_{60}, A_{50}, A_{40}, A_{10})
and each coloured curve represents a Vibration condition (i.e., None = no vibrations; Anger,
Fear). Error bars indicate the standard error of the mean. *$p < .05$.

EEG results

Event-Related Potentials

Influence of induced vibrations during vocal emotion processing on brain dynamics

revealed differences between our clusters considering conditions response with a three-way

Cluster*EmotionalVoice*Vibration interaction along trial length (Supplementary Table 6,

$F(24)=[3.04-23.36$ min-max]; $p<.001$ FDR corrected). Three clusters contained at least one

50ms timebin presenting an EmotionalVoice*Vibration interaction. In the paragraph below,

we describe the results from the functionally best-defined cluster, i.e., the frontocentral cluster

(cluster C4, Figure 3**G** & 4**K**). Results from the two other ERP clusters (occipital and

temporal) are described in the supplementary material, together with posthoc ERP results not

pertaining to our hypotheses ('Supplementary EEG results', *I.* & *II*, and Supplementary

Figures 2-4) and for source reconstruction (Supplementary Figures 4-5).

Effect of Emotional voice X Vibration: frontocentral cluster C4

The frontocentral cluster C4 (Figure 3**G** & 4**K**) presented an EmotionalVoice*

Vibration interaction at 50-150ms, 300-400ms, 450-550ms, 600-650ms, 750-850ms, and 900-

950ms time intervals (Supplementary Table 6; Figure 3 & 4). Three clearly defined peaks

exceeded baseline threshold: a positive deflection (100-150ms), another positive deflection

(200-350ms) and a negative one (from 500ms post-stimulus onset). For the sake of clarity, a

complete list of all statistical contrasts is available in the supplementary material

('Supplementary EEG results', *II.)*.

Low emotional voice ambiguity ERPs

At 125ms, A_{90} voices with anger vibrations elicited a weaker positive deflection

compared to both A_{50} and A_{10} voices. At the same time point, we observed that in A_{90} voices,

anger vibration presented a weaker deflection compared to both fear and no-vibration

conditions. No effects were observed for the P200 peak amplitude. Around 625ms, A_{10} voices

showed a weaker negative deflection in the fear vibration condition compared to both A_{50} and

A_{90} voices. Full results are detailed in the supplementary material (Figure S2, Supplementary

Table 5).

High emotional voice ambiguity ERPs

For ambiguous voices, no expected results were observed around 100ms. In the descending

phase of the P200, at 325ms, A_{40} voices differed from A_{50} voices (fear vibration). At 475ms,

A_{60} voices differed from A_{50} voices (anger vibration). Around 625ms, A_{60} voices differed

from A_{40} voices (anger vibration) and A_{40} voices showed a weaker negative deflection than

both A_{50} and A_{60} voices (fear vibration). As for vibration contrasts, fear vibration elicited a

greater negative deflection than both other conditions for A_{60} and A_{50} voices. At 825ms, all

vibration conditions differed for A_{40} and A_{50} voices, while anger vibration led to a greater

negative deflection than both other conditions for A_{60} voices. Finally, at 925ms, fear vibration

showed a weaker negative deflection than both other conditions for A_{50} and A_{60} voices. Full

results in Figure S2 and Supplementary Table 5. These results already support our EEG

hypotheses, namely the presence of early (100-300ms) and later-stage (400-700ms) ERPs

specific to the vibration conditions, both when these were mostly congruent or incongruent

with the emotion of the voices presented auditorily.

Source reconstruction

Source reconstruction analysis revealed correlates at the brain level of the vibration ERP-dissociations observed at the sensor level (Figure 3**B**-**D**,**F** and 4**B**-**E**,**G**-**J**).

Low emotional voice ambiguity brain sources

We observed in the 300-350ms time interval a fear vs anger vibrations source-ERP contrast in the superior posterior lobe (Figure 3**B**,**C**) but also for incongruent fear vs no-vibrations in the medial orbitofrontal cortex (OFC), insula and primary somatosensory cortex (550-600ms, Figure S4) followed by the SMA (750-800ms) and primary motor cortex (900-950ms; Figure 3**D**). For fearful voices, congruent fear vs no-vibration contrasts were observed in the SMA (700-750ms) and superior parietal lobe (700-750ms; 850-900ms; Figure 3**F**).

Figure 3. **Contrasts between vibrations for Low Ambiguity vocalizations.** Top panel: Contrasts between vibrations for low ambiguity angry voices (A_{90}) for **A:** Event-related potentials and **B-C-D**: Brain activations from source reconstruction. Bottom panel: Contrasts between vibrations for low ambiguity fearful voices (F_{90}) for **E**: Event-related potentials and **F**: Brain activations from source reconstruction. **G**: Graphical representation of the topographical location of the analyzed cluster. For ERPs graphs, line colors for each condition represent the vibration applied (anger in red, fear in blue, no-vibration control in black). Upper horizontal lines represent contrast with substantial proof of difference (BF>3, see Methods) between conditions. Black lower horizontal line represents the period where the two-way Condition X Vibration interaction was important. Blue dashed horizontal line represents the 3 X baseline standard deviation interval. All statistics reported here have Bayes factors > 3. M1: primary motor cortex, SMA: supplementary motor area, SPL: superior parietal lobule.

High emotional voice ambiguity brain sources

384   At the 350-400ms peak, source-ERP fear vs no vibration contrasts were observed in
385 the anterior cingulate cortex and insula for fearful voices (Figure 4**G**-**H**). A significant anger
386 vs fear vibration contrast was also observed at the same timing in the superior parietal cortex
387 for fearful voices (Figure 4**G**). At the 450-500ms peak, a congruent (anger vibration vs. no-
388 vibration) contrast was observed for angry voices in the primary sensory, primary motor, and
389 premotor areas, OFC, dorsolateral prefrontal cortex (DLPFC) as well as in the intraparietal
390 sulcus (IPS; Figure 4**B**). In the late period, a congruent anger vibration vs no-vibration
391 contrast was observed for angry voices in the precuneus (700-750ms, Figure 4**D**), as well as
392 superior parietal lobule and IPS (900-950ms, Figure 4**E**). A congruent fear vibration vs no-
393 vibration contrast was also observed for fearful voices in the primary motor cortex, IFG and
394 OFC (650-700ms) while a fear and anger (global) vibration vs no-vibration contrast was
395 observed in the primary motor area at the same timing (Figure 4**I**). Finally, an incongruent
396 anger vibration vs no-vibration contrast was observed for fearful voices in primary motor
397 cortex and SMA (750-800ms; Figure 4**J**).

398   Source reconstruction results therefore also fit well with our EEG hypotheses both for
399 vibration-specific locations and for error-prediction sites. Taken together, ERP and source
400 reconstruction results mostly confirm our hypotheses and paint a clearer picture of the
401 dynamic neural underpinnings of induced throat vibrations on the recognition of both
402 emotionally 'pure' and ambiguous voices.

Figure 4. **Contrasts between vibrations for High Ambiguity vocalizations.** Top panel:
Contrasts between vibrations for high ambiguity angry voices (A_{10}) for **A**: Event-related
potentials and **B-C-D-E**: Brain activations from source reconstruction. Bottom panel:
Contrasts between vibrations for high ambiguity fearful voices (F_{10}) for **F:** Event-related
potentials and **G-H**: Brain activations from source reconstruction. **K:** Graphical representation

of the topographical location of the analyzed cluster. For ERPs graphs, line colors for each
condition represent the vibration applied (anger in red, fear in blue, no-vibration control in
black). Upper horizontal lines represent contrast with substantial proof of difference (BF>3,
see Methods) between conditions. Black lower horizontal line represents the period where the
two-way Condition X Vibration interaction was significant. Blue dashed horizontal line
represents the 3 X baseline standard deviation interval. All statistics reported here have Bayes
factors > 3. ACC: anterior cingular cortex, DLPFC: dorsolateral prefrontal cortex, IFG:
inferior frontal gyrus, IPS: intraparietal sulcus, M1: primary motor cortex, OFC: orbitofrontal
cortex, op: pars opercularis, Prec: precuneus, PreM1: premotor cortex, S1: primary
somatosensory cortex, SMA: supplementary motor area, SPL: superior parietal lobule.

Discussion

Research on the impact of embodied cognition on vocal emotion processing is very

scarce. Our study investigated whether vocal tract vibrations could represent useable

vibrotactile feedback, and whether this feedback could influence auditory emotion perception,

especially for emotionally ambiguous voices. Behavioural and EEG results revealed effects

that support our assumptions: induced throat vibrations modulated behavioral responses to

emotional voices as well as early and late ERP components, within relevant and expected

brain modulators.

As mentioned, our study highlighted an impact of induced vibrations on the

behavioural responses to morphed emotional voices, which is in line with our main

hypothesis. Although not all the expected contrasts were observed, it was clear that our

participants categorized emotional vocalizations as 'anger' significantly more often when a

concomitant anger vibration was induced than when it was a fear vibration. This result is

consistent with language comprehension facilitation effect of by the transcranial stimulation

of the primary motor areas controlling the lips/tongue (D'Ausilio et al., 2009). We therefore

interpret the results of our study as an emotion-specific stimulation of throat

mechanoreceptors that would facilitate vocal emotion recognition. This is also coherent with

studies on congruence-specific facilitation effect in multimodal processing of emotion (Lin &

Ding, 2019; Niedenthal et al., 2001). Following a similar reasoning, participants categorized

ambiguous emotional vocalizations (A_{60}, $A_{50,}$ and A_{40}) as 'anger' more often when anger (vs.

fear) vibrations were induced—or when there were no vibrations were induced. This latter

contrast is consistent with the SIMS model (Niedenthal et al., 2010): interoceptive feedback

would be integrated especially for non-prototypical emotional stimuli—here, ambiguous
voices, when available emotional information is not clear enough.

At the neural level, the frontocentral cluster C4 was the locus of an interaction
between voice ambiguity and induced throat vibrations—both for early and late time-
windows. Our results showed that the induction of both congruent fear and anger vibrations
led to an increase in P100 amplitude—for low, high and highest (A_{50}) emotional ambiguity. In
the literature, the P100 was linked to attention-driven interoceptive accuracy effects—
auditory tone appearing in synchrony with heartbeat or with a delay (Mendoza-Medialdea &
Ruiz-Padial, 2021). In cross-modal studies, multimodal integration of emotion recognition
occurred very early, around 110ms post-stimulus onset (Pourtois et al., 2000). Furthermore,
studies investigating interoception and somatosensory stimulation observed P300
modulations—slightly later that in our data, particularly in the parietal area and therefore in
line with our results (Al et al., 2020; Auksztulewicz & Blankenburg, 2013; Has et al., 2021;
Herbert et al., 2007; Pollatos et al., 2005).

We also observed the presence of a P200 component in this frontocentral area,
consistent with emotional voice processing (Duville et al., 2022; Gädeke et al., 2013;
Maltezou-Papastylianou et al., 2022; Paulmann et al., 2013; Paulmann & Uskul, 2017). Our
results showed anger vibrations lead to P200 amplitude *increase* (low-ambiguity), while fear
vibrations lead to P200 *decrease* (ambiguous voices), and for incongruent high-ambiguity
angry voices. These results are in line with above-mentioned findings on multimodal
integration and interoception and occur at a slightly later stage. These data therefore support
the impact of vibrotactile throat feedback on behavior through P100 and P200 modulations.
They also align with interoception and somatosensory stimulation instigating a P300
component—especially in the parietal region (Al et al., 2020; Auksztulewicz & Blankenburg,
2013; Has et al., 2021; Herbert et al., 2007; Pollatos et al., 2005). Moreover, the effect of fear
vibrations for ambiguous voice categorization only is also supported by the SIMS model
(Niedenthal, 2007; Niedenthal et al., 2010).

Concerning later ERP components, we observed modulations by induced vibrations
most notably between 500 and 800ms post-stimulus onset. These modulations involve both
congruent and incongruent cases, especially for low-ambiguity voices—and both emotions.
For instance, at 500ms, anger vibrations produce an incongruent negative deflection for low-
ambiguity fearful voices. These results recall the N400 component observed in frontal,
parietal and frontocentral areas during emotional voice processing (Föcker & Röder, 2019;

Maltezou-Papastylianou et al., 2022; Tarai & Srinivasan, 2019). At 800ms, power increase—
although still negative—for both emotional vibrations compared to no-vibration for
ambiguous voices compares to the Late Positive Component (LPC). The LPC is usually
associated with higher-order cognitive processes: arousal information (Paulmann et al., 2013),
prosody expectancy violation (Astésano et al., 2004; Brattico et al., 2006; Paulmann et al.,
2012), emotional 'meaning' (Wei et al., 2022; Zora et al., 2020). Nevertheless, this echoes
once again with the SIMS model—embodied simulation is engaged when the emotional
stimulus is non-prototypical (Niedenthal et al., 2010). Here, the vibrations were integrated
specifically for ambiguous voices, as prototypical emotional voice processing would not be
enough to identify the emotion. At the interoception and proprioception levels, these results
globally show that vibrations are integrated during emotional processing at multiple stages,
dynamically and throughout the time course of an event, similarly to the multimodal
integration shown in audiovisual studies (Föcker & Röder, 2019; Klasen et al., 2012).

ERP source reconstruction revealed source modulations in several key regions. Some
are involved in vibratory processing—the DLPFC, vibrotactile working memory (Burton et
al., 2010) or in vibratory processing, crucially in vibration rate or pattern discrimination
(Christensen et al., 2007; Coghill et al., 1994; Golaszewski et al., 2006; Hernández et al.,
2002; Lenoir et al., 2017; Li Hegner et al., 2010; Macar et al., 2002; Romo et al., 2004;
Woolgar & Zopf, 2017; Wu et al., 2018). Concerning more internal feedback or interoception,
we observed source modulations in the insula, known to be crucially involved in interoceptive
processing (Adolfi et al., 2017; Casals-Gutierrez & Abbey, 2020) and precuneus
(interoceptive awareness; Herbert et al., 2007; Longarzo et al., 2021; Wilson-Mendenhall et
al., 2019). The modulation of the primary somatosensory cortex might be related to
proprioceptive information processing while modulations in the OFC were also reported in
interoception studies (Singer & Klimecki, 2014; Stern et al., 2017).

Regarding emotional prosody perception, we observed source modulations in the
DLPFC (prosodic working memory; Mitchell, 2007), superior parietal lobule (emotional
prosody judgement; Bach et al., 2008), as well as in other regions also involved in emotional
prosody processing (IPS, SMA; Alba-Ferrara et al., 2011; Belyk & Brown, 2014; Kotz et al.,
2003). OFC modulations fit well with higher-level processing of emotional prosody
(Wildgruber et al., 2006)—especially when ambiguous (Grandjean, 2021), or for accurately
categorizing human voice (Ceravolo et al., 2023). Source modulations in the insula may
contribute to external emotional stimuli integration (vibrotactile feedback) and emotional

experiences with interoceptive signals (Adolfi et al., 2017; Nguyen et al., 2016; Zaki et al.,
2012). Finally, above-mentioned modulations—superior parietal lobule, IPS, premotor,
primary somatosensory cortex—are in line with multimodal integration for auditory, visual,
sensorimotor and tactile stimuli (Baker et al., 2018; Culham & Valyear, 2006; de Borst & de
Gelder, 2017; Gentile et al., 2011; Kreifelts et al., 2007; Parker et al., 2020). Source
modulation locations for the Vibration and Emotion factors are fully in line with our previous
functional MRI results on neural correlates of emitted, measured vocal tract vibrations during
emotional expression (Selosse et al., 2025). Combined with the present data, they help draw a
clearer picture of emotion embodiment in voice signals.

Taken together, these results suggest the implication of a dynamic, wide-spread
network of brain areas in the processing of throat induced vibrations and their integration with
emotional voices. This is especially true for highly ambiguous voice signals, for which such
information is most relevant for improved decision making. This interpretation relies on
previous literature on interoception and multisensory integration, and suggests that induced
throat vibrations are a type of feedback critical enough to impact auditory perception and
decision.

Limitations and perspectives
Our sample consists mainly of female participants, despite gender differences in emotional
prosody recognition (Imaizumi et al., 2004; Lambrecht et al., 2014) and facial mimicry (Korb
et al., 2015; Niedenthal et al., 2012). Second, it would be necessary to replicate the observed
effects we with different emotions to generalize our results, although morphing positive and
negative emotions may be problematic. Third, induced vibrations should be those of the
participants for improved ecological validity—and better consideration of interindividual
differences. Fourth, vibrations were induced on the skin, therefore more subtle internal tissue
vibrations encoded by different mechano-receptors may not be triggered. Further research is
needed to clarify the relationships between actual vibrations created by the vocal cords and
those induced on the skin level. Future studies would benefit from the inclusion of measures
of bodily self-consciousness or interoceptive awareness. In fact, interindividual variability in
this dimension probably impacts subsequent data as the feedback mechanism highlighted in
the present work might highly depend on such sensibility.

Conflict of interest

The authors declare no conflict of interest whatsoever.

Data availability statement

Data and codes will be made available upon request to the corresponding author.

Funding

This work as well as Open Access publication charges were funded by the Swiss National Science Foundation (grant number 10531C_189397/DG-LC).

References

Adolfi, F., Couto, B., Richter, F., Decety, J., Lopez, J., Sigman, M., & Ibáñez, A. (2017).

Convergence of interoception, emotion, and social cognition: a twofold fMRI meta-

analysis and lesion approach. *Cortex*, *88*, 124--142.

Al, E., Iliopoulos, F., Forschack, N., Nierhaus, T., Grund, M., Motyka, P., & Villringer, A.

(2020). Heart–brain interactions shape somatosensory perception and evoked

potentials. *Proceedings of the National Academy of Sciences*, *117*(19), 10575--10584.

Alba-Ferrara, L., Hausmann, M., Mitchell, R. L., & Weis, S. (2011). The Neural Correlates of

Emotional Prosody Comprehension: Disentangling Simple from Complex Emotion.

PLoS ONE, *6*(12), 28701. https://doi.org/10.1371/journal.pone.0028701

Ashley, V., Vuilleumier, P., & Swick, D. (2004). Time course and specificity of event-related

potentials to emotional expressions. *Neuroreport*, *15*(1), 211--216.

Astésano, C., Besson, M., & Alter, K. (2004). Brain potentials during semantic and prosodic

processing in French. *Cognitive Brain Research*, *18*(2), 172--184.

Auksztulewicz, R., & Blankenburg, F. (2013). Subjective rating of weak tactile stimuli is

parametrically encoded in event-related potentials. *Journal of Neuroscience*, *33*(29),

11878--11887.

Bach, D. R., Grandjean, D., Sander, D., Herdener, M., Strik, W. K., & Seifritz, E. (2008). The

effect of appraisal level on processing of emotional prosody in meaningless speech.

NeuroImage, *42*(2), 919--927. https://doi.org/10.1016/j.neuroimage.2008.05.034

Baker, C. M., Burks, J. D., Briggs, R. G., Conner, A. K., Glenn, C. A., Taylor, K. N., Sali, G.,

McCoy, T. M., Battiste, J. D., O'Donoghue, D. L., & Sughrue, M. E. (2018). A

Connectomic Atlas of the Human Cerebrum—Chapter 7: The Lateral Parietal Lobe.

Operative Neurosurgery, *15*(suppl_1), 295--349. https://doi.org/10.1093/ons/opy261

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., Dai, B.,

Grothendieck, G., Green, P., & Bolker, M. B. (2015). Package 'lme4'. *convergence*,

12(1), 2.

Belin, P., Fillion-Bilodeau, S., & Gosselin, F. (2008). The Montreal Affective Voices: A

validated set of nonverbal affect bursts for research on auditory affective processing.

Behavior research methods, *40*(2), 531--539.

Belyk, M., & Brown, S. (2014). Perception of affective and linguistic prosody: An ALE meta-

analysis of neuroimaging studies. *Social Cognitive and Affective Neuroscience*, *9*(9),

1395--1403. https://doi.org/10.1093/scan/nst124

Blairy, S., Herrera, P., & Hess, U. (1999). Mimicry and the judgment of emotional facial

expressions. *Journal of Nonverbal Behavior*, *23*, 5--41.

Bland, J. M., & Altman, D. G. (1995). Multiple significance tests: the Bonferroni method.

BMJ, *310*(6973), 170.

Borgomaneri, S., Bolloni, C., Sessa, P., & Avenanti, A. (2020). Blocking facial mimicry

affects recognition of facial and body expressions. *PLoS ONE*, *15*(2), e0229364.

Brainard, D. H. (1997). The psychophysics toolbox. *Spatial Vision*, *10*(4), 433--436.

https://doi.org/10.1163/156856897x00357

Brattico, E., Tervaniemi, M., Näätänen, R., & Peretz, I. (2006). Musical scale properties are

automatically processed in the human auditory cortex. *Brain Research*, *1117*(1), 162--

174.

Brosch, T., Grandjean, D., Sander, D., & Scherer, K. R. (2008). Behold the voice of wrath:

Cross-modal modulation of visual attention by anger prosody. *Cognition*, *106*(3),

1497--1503.

Brosch, T., Sander, D., Pourtois, G., & Scherer, K. R. (2008). Beyond fear: Rapid spatial

orienting toward positive emotional stimuli. *Psychological science*, *19*(4), 362--370.

Brunswik, E. (1956). *Perception and the representative design of psychological experiments*.
University of California Press.

Burleson, M. H., & Quigley, K. S. (2021). Social interoception and social allostasis through
touch: legacy of the somatovisceral afference model of emotion. *Social neuroscience*,
16(1), 92--102.

Burton, H., Sinclair, R. J., & Dixit, S. (2010). Working memory for vibrotactile frequencies:
Comparison of cortical activity in blind and sighted individuals. *Human Brain
Mapping*, *31*(11), 1686--1701. https://doi.org/10.1002/hbm.20966

Carminati, M., Fiori-Duharcourt, N., & Isel, F. (2018). Neurophysiological differentiation
between preattentive and attentive processing of emotional expressions on French
vowels. *Biological psychology*, *132*, 55--63.

Casals-Gutierrez, S., & Abbey, H. (2020). Interoception, mindfulness and touch: A meta-
review of functional MRI studies. *International Journal of Osteopathic Medicine*, *35*,
22--33.

Ceravolo, L., Debracque, C., Pool, E., Gruber, T., & Grandjean, D. (2023). Frontal
mechanisms underlying primate calls recognition by humans. *Cerebral Cortex
Communications*, *4*(4), tgad019.

Ceravolo, L., Frühholz, S., Pierce, J., Grandjean, D., & Péron, J. (2021). Basal ganglia and
cerebellum contributions to vocal emotion processing as revealed by high-resolution
fMRI. *Scientific reports*, *11*(1), 10645.

Ceravolo, L., Moisa, M., Grandjean, D., Ruff, C., & Frühholz, S. (2026). Behavioral and
neural aftermath of right inferior frontal cortex disruption on ambiguous vocal
emotion decisional processes. *Scientific reports*. https://doi.org/10.1038/s41598-026-
39668-0

Chartrand, T. L., & Bargh, J. A. (1999). The chameleon effect: The perception–behavior link
and social interaction. *Journal of personality and social psychology*, *76*(6), 893.

Christensen, M. S., Lundbye-Jensen, J., Geertsen, S. S., Petersen, T. H., Paulson, O. B., &
Nielsen, J. B. (2007). Premotor cortex modulates somatosensory cortex during
voluntary movements without proprioceptive feedback. *Nature Neuroscience*, *10*(4),
417--419. https://doi.org/10.1038/nn1873

Coghill, R., Talbot, J., Evans, A., Meyer, E., Gjedde, A., Bushnell, M., & Duncan, G. (1994).
Distributed processing of pain and vibration by the human brain. *The Journal of
Neuroscience*, *14*(7), 4095--4108. https://doi.org/10.1523/jneurosci.14-07-04095.1994

Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*. Hillsdale.

Couto, B., Adolfi, F., Velasquez, M., Mesow, M., Feinstein, J., Canales-Johnson, A., &
Ibanez, A. (2015). Heart evoked potential triggers brain responses to natural affective
scenes: a preliminary study. *Autonomic Neuroscience*, *193*, 132--137.

Critchley, H. D., & Garfinkel, S. N. (2017). Interoception and emotion. *Current opinion in
psychology*, *17*, 7--14.

Culham, J. C., & Valyear, K. F. (2006). Human parietal cortex in action. *Current Opinion in
Neurobiology*, *16*(2), 205--212. https://doi.org/10.1016/j.conb.2006.03.005

D'Ausilio, A., Pulvermüller, F., Salmas, P., Bufalari, I., Begliomini, C., & Fadiga, L. (2009).
The motor somatotopy of speech perception. *Current Biology*, *19*(5), 381--385.

de Borst, A. W., & de Gelder, B. (2017). fMRI-based multivariate pattern analyses reveal
imagery modality and imagery content specific representations in primary
somatosensory, motor and auditory cortices. *Cerebral Cortex*, *27*(8), 3994--4009.
https://doi.org/10.1093/cercor/bhw211

Dimberg, U. (1982). Facial reactions to facial expressions. *Psychophysiology*, *19*(6), 643--
647.

Drimalla, H., Landwehr, N., Hess, U., & Dziobek, I. (2019). From face to face: the
contribution of facial mimicry to cognitive and emotional empathy. *Cognition and*
Emotion.

Duville, M. M., Alonso-Valerdi, L. M., & Ibarra-Zarate, D. I. (2022). Neuronal and
behavioral affective perceptions of human and naturalness-reduced emotional
prosodies. *Frontiers in Computational Neuroscience*, *16*, 1022787.

Engelen, T., Solcà, M., & Tallon-Baudry, C. (2023). Interoceptive rhythms in the brain.
Nature Neuroscience, *26*(10), 1670--1684.

Fadiga, L., Craighero, L., Buccino, G., & Rizzolatti, G. (2002). Speech listening specifically
modulates the excitability of tongue muscles: a TMS study. *European Journal of*
Neuroscience, *15*(2), 399--402.

Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G* Power 3: A flexible statistical
power analysis program for the social, behavioral, and biomedical sciences. *Behavior*
research methods, *39*(2), 175--191.

Föcker, J., & Röder, B. (2019). Event-related potentials reveal evidence for late integration of
emotional prosody and facial expression in dynamic stimuli: An ERP study.
Multisensory Research, *32*(6), 473--497.

Gädeke, J. C., Föcker, J., & Röder, B. (2013). Is the processing of affective prosody
influenced by spatial attention? An ERP study. *BMC neuroscience*, *14*(1), 1--15.

Gentile, G., Petkova, V. I., & Ehrsson, H. H. (2011). Integration of Visual and Tactile Signals
From the Hand in the Human Brain: An fMRI Study. *Journal of Neurophysiology*,
105(2), 910--922. https://doi.org/10.1152/jn.00840.2010

Gibson, J. (2019). Mindfulness, interoception, and the body: A contemporary perspective.
Frontiers in Psychology, *10*.

Golaszewski, S. M., Siedentopf, C. M., Koppelstaetter, F., Fend, M., Ischebeck, A., Gonzalez-
Felipe, V., Haala, I., Struhal, W., Mottaghy, F. M., Gallasch, E., Felber, S. R., &
Gerstenbrand, F. (2006). Human brain structures related to plantar vibrotactile
stimulation: A functional magnetic resonance imaging study. *NeuroImage*, *29*(3), 923-
-929. https://doi.org/10.1016/j.neuroimage.2005.08.052

Grandjean, D. (2021). Brain networks of emotional prosody processing. *Emotion Review*,
13(1), 34--43.

Grandjean, D., Bänziger, T., & Scherer, K. R. (2006). Intonation as an interface between
language and affect. *Progress in brain research*, *156*, 235--247.

Has, N. A., Jamaludin, M. N., Rejab, S. M., Ahmad, Z., Zainuddin, N. F., Salam, A. H. A., &
As'ari, M. A. (2021). P300 Somatosensory Validation of Vibrotactile Haptic Feedback
for Upper Limb Prosthesis. 2021 IEEE National Biomedical Engineering Conference
(NBEC),

Hawk, S. T., Fischer, A. H., & van Kleef, G. A. (2012). Face the noise: embodied responses
to nonverbal vocalizations of discrete emotions. *Journal of personality and social*
psychology, *102*(4), 796.

Herbert, B. M., Pollatos, O., & Schandry, R. (2007). Interoceptive sensitivity and emotion
processing: an EEG study. *International Journal of Psychophysiology*, *65*(3), 214--
227.

Hernández, A., Zainos, A., & Romo, R. (2002). Temporal Evolution of a Decision-Making
Process in Medial Premotor Cortex. *Neuron*, *33*(6), 959--972.
https://doi.org/10.1016/s0896-6273(02)00613-x

Hess, U., & Blairy, S. (2001). Facial mimicry and emotional contagion to dynamic emotional
facial expressions and their influence on decoding accuracy. *International Journal of*
Psychophysiology, *40*(2), 129--141.

Hess, U., & Fischer, A. (2014). Emotional mimicry: Why and when we mimic emotions.
Social and personality psychology compass, *8*(2), 45--57.

Holland, A. C., O'Connell, G., & Dziobek, I. (2021). Facial mimicry, empathy, and emotion
recognition: a meta-analysis of correlations. *Cognition and Emotion*, *35*(1), 150--168.

Hyniewska, S., & Sato, W. (2015). Facial feedback affects valence judgments of dynamic and
static emotional expressions. *Frontiers in Psychology*, *6*, 291.

Imaizumi, S., Homma, M., Ozawa, Y., Maruishi, M., & Muranaka, H. (2004). Gender
differences in emotional prosody processing—An fMRI study. *Psychologia*, *47*(2),
113--124.

Kawahara, H., & Matsui, H. (2003). Auditory morphing based on an elastic perceptual
distance metric in an interference-free time-frequency representation. 2003 IEEE
International Conference on Acoustics, Speech, and Signal Processing,

Khalsa, S. S., Adolphs, R., Cameron, O. G., Critchley, H. D., Davenport, P. W., Feinstein, J.
S., & Zucker, N. (2018). Interoception and mental health: a roadmap. *Biological*
psychiatry: cognitive neuroscience and neuroimaging, *3*(6), 501--513.

Klasen, M., Chen, Y. H., & Mathiak, K. (2012). Multisensory emotions: perception,
combination and underlying neural processes. *Reviews in the Neurosciences*, *23*(4),
381--392.

Kleiner, M., Brainard, D., & Pelli, D. (2007). What's new in Psychtoolbox-3. In.

Korb, S., Grandjean, D., & Scherer, K. R. (2010). Timing and voluntary suppression of facial
mimicry to smiling faces in a Go/NoGo task—An EMG study. *Biological psychology*,
85(2), 347--349.

Korb, S., Malsert, J., Rochas, V., Rihs, T. A., Rieger, S. W., Schwab, S., & Grandjean, D.
(2015). Gender differences in the neural network of facial mimicry of smiles–An
rTMS study. *Cortex*, *70*, 101--114.

Kotz, S. A., Meyer, M., Alter, K., Besson, M., von Cramon, D. Y., & Friederici, A. D. (2003).

On the lateralization of emotional prosody: An event-related functional MR

investigation. *Brain and Language*, *86*(3), 366--376. https://doi.org/10.1016/s0093-

934x(02)00532-1

Kreifelts, B., Ethofer, T., Grodd, W., Erb, M., & Wildgruber, D. (2007). Audiovisual

integration of emotional signals in voice and face: An event-related fMRI study.

NeuroImage, *37*(4), 1445--1456. https://doi.org/10.1016/j.neuroimage.2007.06.020

Künecke, J., Wilhelm, O., & Sommer, W. (2017). Emotion recognition in nonverbal face-to-

face communication. *Journal of Nonverbal Behavior*, *41*, 221--238.

Lambrecht, L., Kreifelts, B., & Wildgruber, D. (2014). Gender differences in emotion

recognition: Impact of sensory modality and emotional category. *Cognition &*

Emotion, *28*(3), 452--469.

Laukka, P., Juslin, P., & Bresin, R. (2005). A dimensional approach to vocal expression of

emotion. *Cognition & Emotion*, *19*(5), 633--653.

Lenoir, C., Huang, G., Vandermeeren, Y., Hatem, S. M., & Mouraux, A. (2017). Human

primary somatosensory cortex is differentially involved in vibrotaction and

nociception. *Journal of Neurophysiology*, *118*(1), 317--330.

https://doi.org/10.1152/jn.00615.2016

Lenth, R., Singmann, H., Love, J., Buerkner, P., & Herve, M. (2019). Package 'emmeans'. *R*

package version, *1*(3.2).

Li Hegner, Y., Lee, Y., Grodd, W., & Braun, C. (2010). Comparing Tactile Pattern and

Vibrotactile Frequency Discrimination: A Human fMRI Study. *Journal of*

Neurophysiology, *103*(6), 3115--3122. https://doi.org/10.1152/jn.00940.2009

Lin, Y., & Ding, H. (2019). Multisensory integration of emotions in a face-prosody-semantics

Stroop task. In *NeuroManagement and Intelligent Computing Method on Multimodal*

Interaction (pp. 1--5).

Lindín, M., Zurrón, M., & Díaz, F. (2004). Changes in P300 amplitude during an active

standard auditory oddball task. *Biological psychology*, *66*(2), 153--167.

Longarzo, M., Mele, G., Alfano, V., Salvatore, M., & Cavaliere, C. (2021). Gender Brain

Structural Differences and Interoception. *Frontiers in Neuroscience*, *14*, 586860.

https://doi.org/10.3389/fnins.2020.586860

Lylykangas, J., Surakka, V., Rantala, J., Raisamo, J., Raisamo, R., & Tuulari, E. (2009).

Vibrotactile information for intuitive speed regulation. In *People and Computers XXIII*

Celebrating People and Technology (pp. 112--119).

Macar, F., Lejeune, H., Bonnet, M., Ferrara, A., Pouthas, V., Vidal, F., & Maquet, P. (2002).

Activation of the supplementary motor area and of attentional networks during

temporal processing. *Experimental Brain Research*, *142*(4), 475--485.

Maltezou-Papastylianou, C., Russo, R., Wallace, D., Harmsworth, C., & Paulmann, S. (2022).

Different stages of emotional prosody processing in healthy ageing–evidence from

behavioural responses, ERPs, tDCS, and tRNS. *PLoS ONE*, *17*(7), e0270934.

Mendoza-Medialdea, M. T., & Ruiz-Padial, E. (2021). Understanding the capture of

exogenous attention by disgusting and fearful stimuli: the role of interoceptive

accuracy. *International Journal of Psychophysiology*, *161*, 53--63.

Mitchell, R. L. C. (2007). fMRI delineation of working memory for emotional prosody in the

brain: Commonalities with the lexico-semantic emotion network. *NeuroImage*, *36*(3),

1015--1025. https://doi.org/10.1016/j.neuroimage.2007.03.016

Moody, E. J., & McIntosh, D. N. (2011). Mimicry of dynamic emotional and motor-only

stimuli. *Social Psychological and Personality Science*, *2*(6), 679--686.

Moody, E. J., McIntosh, D. N., Mann, L. J., & Weisser, K. R. (2007). More than mere

mimicry? The influence of emotion on rapid facial reactions to faces. *Emotion*, *7*(2),

447.

Munger, J. B., & Thomson, S. L. (2008). Frequency response of the skin on the head and neck

during production of selected speech sounds. *The Journal of the Acoustical Society of*

America, *124*(6), 4001--4012.

Nguyen, V. T., Breakspear, M., Hu, X., & Guo, C. C. (2016). The integration of the internal

and external milieu in the insula during dynamic emotional experiences. *NeuroImage*,

124, 455--463. https://doi.org/10.1016/j.neuroimage.2015.08.078

Niedenthal, P. M. (2007). Embodying Emotion. *Science*, *316*(5827), 1002--1005.

Niedenthal, P. M., Augustinova, M., Rychlowska, M., Droit-Volet, S., Zinner, L., Knafo, A.,

& Brauer, M. (2012). Negative relations between pacifier use and emotional

competence. *Basic and Applied Social Psychology*, *34*(5), 387--394.

Niedenthal, P. M., Brauer, M., Halberstadt, J. B., & Innes-Ker, Å. H. (2001). When did her

smile drop? Facial mimicry and the influences of emotional state on the detection of

change in emotional expression. *Cognition & Emotion*, *15*(6), 853--864.

Niedenthal, P. M., Mermillod, M., Maringer, M., & Hess, U. (2010). The Simulation of

Smiles (SIMS) model: Embodied simulation and the meaning of facial expression.

Behavioral and brain sciences, *33*(6), 417.

Nolan, M., Madden, B., & Burke, E. (2009). Accelerometer based measurement for the

mapping of neck surface vibrations during vocalized speech. Annual International

Conference of the IEEE Engineering in Medicine and Biology Society,

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: open source

software for advanced analysis of MEG, EEG, and invasive electrophysiological data.

Computational intelligence and neuroscience, *2011*(1), 156869.

Paquette, S., Rigoulot, S., Grunewald, K., & Lehmann, A. (2020). Temporal decoding of

vocal and musical emotions: Same code, different timecourse? *Brain Research*,

146887.

Parker, A. G., Briggs, R. G., Conner, A. K., O'Neal, C. M., Bonney, P. A., Maxwell, B. D.,

Baker, C. M., Burks, J. D., Sali, G., & Sughrue, M. E. (2020). Parcellation-based

tractographic modeling of the ventral attention network. *Journal of the Neurological*

Sciences, *408*, 116548.

Paulmann, S., Bleichner, M., & Kotz, S. A. (2013). Valence, arousal, and task effects in

emotional prosody processing. *Frontiers in Psychology*, *4*, 345.

Paulmann, S., Jessen, S., & Kotz, S. A. (2012). It's special the way you say it: An ERP

investigation on the temporal dynamics of two types of prosody. *Neuropsychologia*,

50(7), 1609--1620.

Paulmann, S., & Kotz, S. A. (2008). An ERP investigation on the temporal dynamics of

emotional prosody and emotional semantics in pseudo-and lexical-sentence context.

Brain and Language, *105*(1), 59--69.

Paulmann, S., & Uskul, A. K. (2017). Early and late brain signatures of emotional prosody

among individuals with high versus low power. *Psychophysiology*, *54*(4), 555--565.

Pell, M. D., & Kotz, S. A. (2021). Comment: The next frontier: Prosody research gets

interpersonal. *Emotion Review*, *13*(1), 51--56.

Pelli, D. G. (1997). The videotoolbox software for visual psychophysics: Transforming

numbers into movies. *Spatial Vision*, *10*(4), 437--442.

Peron, J., Frühholz, S., Ceravolo, L., & Grandjean, D. (2016). Structural and functional

connectivity of the subthalamic nucleus during vocal emotion decoding. *Social*

Cognitive and Affective Neuroscience, *11*(2), 349-356.

Philip, L., Martin, J. C., & Clavel, C. (2018). Rapid facial reactions in response to facial

expressions of emotion displayed by real versus virtual faces. *i-Perception*, *9*(4),

2041669518786527.

Pollatos, O., Kirsch, W., & Schandry, R. (2005). On the relationship between interoceptive

awareness, emotional experience, and brain processes. *Cognitive Brain Research*,

25(3), 948--962.

Pourtois, G., de Gelder, B., Vroomen, J., Rossion, B., & Crommelinck, M. (2000). The time-

course of intermodal binding between seeing and hearing affective information.

Neuroreport, *11*(6), 1329--1333.

Proverbio, A. M., Benedetto, F., & Guazzone, M. (2020). Shared neural mechanisms for

processing emotions in music and vocalizations. *European Journal of Neuroscience*,

51(9), 1987--2007.

Risberg, A., & Lubker, J. (1978). Prosody and speechreading. *Speech Transmission*

Laboratory Quarterly Progress Report and Status Report, *4*, 1--16.

Romo, R., Hernández, A., & Zainos, A. (2004). Neuronal Correlates of a Perceptual Decision

in Ventral Premotor Cortex. *Neuron*, *41*(1), 165--173. https://doi.org/10.1016/s0896-

6273(03)00817-1

Salamone, P. C., Legaz, A., Sedeño, L., Moguilner, S., Fraile-Vazquez, M., Campo, C. G., &

Ibañez, A. (2021). Interoception primes emotional processing: multimodal evidence

from neurodegeneration. *Journal of Neuroscience*, *41*(19), 4276--4292.

Scherer, K. R. (2003). Vocal communication of emotion: A review of research paradigms.

Speech communication, *40*(1-2), 227--256.

Scherer, K. R., & Oshinsky, J. S. (1977). Cue utilization in emotion attribution from auditory

stimuli. *Motivation and Emotion*, *1*(4), 331--346.

Schirmer, A., & Kotz, S. A. (2006). Beyond the right hemisphere: brain mechanisms
mediating vocal emotional processing. *Trends in cognitive sciences*, *10*(1), 24--30.

Schneider, K. G., Hempel, R. J., & Lynch, T. R. (2013). That "poker face" just might lose you
the game! The impact of expressive suppression and mimicry on sensitivity to facial
expressions of emotion. *Emotion*, *13*(5), 852.

Selosse, G., Grandjean, D., & Ceravolo, L. (2025). Neural correlates of embodied and
vibratory mechanisms associated with emotional prosody production. *Social Cognitive*
and Affective Neuroscience, *20*(1), nsaf084.

Singer, T., & Klimecki, O. M. (2014). Empathy and compassion. *Current Biology*, *24*(18),
875--878.

Stel, M., & van Knippenberg, A. (2008). The role of facial mimicry in the recognition of
affect. *Psychological science*, *19*(10), 984.

Stern, E. R., Grimaldi, S. J., Muratore, A., Murrough, J., Leibu, E., Fleysher, L., & Burdick,
K. E. (2017). Neural correlates of interoception: Effects of interoceptive focus and
relationship to dimensional measures of body awareness. *Human Brain Mapping*,
38(12), 6068--6082.

Sundberg, J. (1992). Phonatory vibrations in singers: A critical review. *Music Perception: An*
Interdisciplinary Journal, *9*(3), 361--381.

Švec, J. G., Titze, I. R., & Popolo, P. S. (2005). Estimation of sound pressure levels of voiced
speech from skin vibration of the neck. *The Journal of the Acoustical Society of*
America, *117*(3), 1386--1394.

Tarai, S., & Srinivasan, N. (2019). Emotional prosody Stroop effect in Hindi: an event related
potential study. In *Progress in brain research* (Vol. 247, pp. 193--217).

Thomas, S. J., Johnstone, S. J., & Gonsalvez, C. J. (2007). Event-related potentials during an
emotional Stroop task. *International Journal of Psychophysiology*, *63*(3), 221--231.

Tsolaki, A., Kosmidou, V., Hadjileontiadis, L., Kompatsiaris, I. Y., & Tsolaki, M. (2015).

Brain source localization of MMN, P300 and N400: Aging and gender differences.

Brain Research, 32--49.

Tuuri, K., Eerola, T., & Pirhonen, A. (2010). Leaping across modalities: Speed regulation

messages in audio and tactile domains. International Workshop on Haptic and Audio

Interaction Design,

Vachon, F., Hughes, R. W., & Jones, D. M. (2012). Broken expectations: violation of

expectancies, not novelty, captures auditory attention. *Journal of Experimental*

Psychology: Learning, Memory, and Cognition, *38*(1), 164.

Venables, W. N., & Smith, D. M. (2003). The R development core team. *An Introduction to*

R, Version, *1*(0).

Watkins, K. E., Strafella, A. P., & Paus, T. (2003). Seeing and hearing speech excites the

motor system involved in speech production. *Neuropsychologia*, *41*(8), 989--994.

Wei, H., He, Y., Kauschke, C., Scharinger, M., & Domahs, U. (2022). An EEG-study on L2

categorization of emotional prosody in German. Speech Prosody,

Wildgruber, D., Ackermann, H., Kreifelts, B., & Ethofer, T. (2006). Cerebral processing of

linguistic and emotional prosody: fMRI studies. In *Progress in brain research* (Vol.

156, pp. 249--268). https://doi.org/10.1016/s0079-6123(06)56013-3

Wilson-Mendenhall, C. D., Henriques, A., Barsalou, L. W., & Barrett, L. F. (2019). Primary

Interoceptive Cortex Activity during Simulated Experiences of the Body. *Journal of*

Cognitive Neuroscience, *31*(2), 221--235. https://doi.org/10.1162/jocn_a_01346

Wilson, S. M., Saygin, A. P., Sereno, M. I., & Iacoboni, M. (2004). Listening to speech

activates motor areas involved in speech production. *Nature Neuroscience*, *7*(7), 701--

702.

Witteman, J., van Heuven, V. J., & Schiller, N. O. (2012). Hearing feelings: a quantitative

meta-analysis on the neuroimaging literature of emotional prosody perception.

Neuropsychologia, *50*(12), 2752--2763.

Woolgar, A., & Zopf, R. (2017). Multisensory coding in the multiple-demand regions:

Vibrotactile task information is coded in frontoparietal cortex. *Journal of*

Neurophysiology, *118*(2), 703--716. https://doi.org/10.1152/jn.00559.2016

Wu, Y., Uluç, I., Schmidt, T. T., Tertel, K., Kirilina, E., & Blankenburg, F. (2018).

Overlapping frontoparietal networks for tactile and visual parametric working memory

representations. *NeuroImage*, *166*, 325--334.

Zaki, J., Davis, J. I., & Ochsner, K. N. (2012). Overlapping activity in anterior insula during

interoception and emotional experience. *NeuroImage*, *62*(1), 493--499.

Zora, H., Rudner, M., & Magnusson, A. K. M. (2020). Concurrent affective and linguistic

prosody with the same emotional valence elicits a late positive ERP response.

European Journal of Neuroscience, *51*(11), 2236--2249.

907