Evolutionary responses and genomic consequences of polyploidization in natural

populations of *Orychophragmus*

Qiang Lai[1], Zeng Wang[1], Changfu Jia[1], Xiner Qumu[1], Rui Wang[1], Zhipeng Zhao[1], Yao

Liu[1], Yukang Hou[1], Jianquan Liu[1], Pär K. Ingvarsson[2], Jing Wang[1*]

[1] Key Laboratory for Bio-Resources and Eco-Environment of Ministry of Education,

College of Life Sciences, Sichuan University, Chengdu, China

[2] Linnean Centre for Plant Biology, Department of Plant Biology, Uppsala BioCenter,

Swedish University of Agricultural Sciences, Uppsala, Sweden

*Corresponding authors: wangjing2019@scu.edu.cn

Abstract

Polyploidization has occurred throughout the tree of life and is particularly common in

plants. Despite its ubiquity, our understanding of the short- and long-term effects and

consequences of genome doubling in natural populations remains incomplete. In this

study, we identified a novel ploidy-variable species system within the ornamental and

industrial oilseed genus *Orychophragmus* (Brassicaceae), which comprises six species,

including diploid and tetraploid cytotypes of *O. taibaiensis*. By integrating population-

scale genomic and transcriptomic datasets across the species in this genus, we

constructed a robust phylogenetic framework and investigated the divergence and

demographic history of *O. taibaiensis* in comparison to its relatives. Specifically, we

characterized the geographical distribution patterns of diploids and tetraploids in

natural populations of *O. taibaiensis*, confirmed the autopolyploid origin of tetraploids,

and inferred their origin time relative to diploid counterparts. Our findings further

revealed that, following genome doubling, tetraploids accumulated a higher genetic

load of deleterious mutations, likely due to relaxed purifying selection facilitated by

allelic redundancy. Additionally, genome doubling was associated with pronounced

changes in gene expression patterns, with differentially expressed genes evolving under

relaxed selective constraints. These results highlight that the initial masking of

deleterious mutations, changes in expression regulation, and divergent efficacy of

selection likely all contribute to shaping the establishment and evolutionary potential

of polyploids.

Key words: demographic history, gene expression, genetic load, *Orychophragmus*,

polyploidization, selection efficacy

Introduction

Polyploidization, resulting from whole-genome duplication (WGD), has long been regarded as a key driver of plant speciation, adaptation to novel and extreme environments, and genetic innovation (Otto and Whitton 2000; Van de Peer, et al. 2017). While allopolyploidy has received considerable attention, autopolyploidy remains markedly less studied, despite its prevalence and suitability for exploring the direct impacts of immediate genome doubling without the complications associated with hybridity (Parisod, et al. 2010; Barker, et al. 2016). Autopolyploidy has been repeatedly shown to arise and establish from diploid populations, often leading to the coexistence of tetraploid and diploid populations within a single species (Monnahan, et al. 2019; Morgan, et al. 2020). However, our understanding of the evolutionary dynamics and genomic consequences of genome doubling in these populations remains limited (Kolář, et al. 2017; Mortier, et al. 2024). For example, it is unclear to what extent the additional chromosomal copies resulting from WGD can mask deleterious mutations, thereby influencing the genetic load (Griswold 2021). Furthermore, the degree to which genome duplication alters the efficiency of selection processes in tetraploid populations compared to their diploid progenitors remains largely unexplored (Spoelhof, et al. 2017). Polyploidy is also frequently associated with morphological and ecological changes, but the extent to which autopolyploidy reshapes gene expression patterns and the functional roles of differentially expressed genes remains an open and intriguing question (Osborn, et al. 2003).

Species diversification within the Brassicaceae family is intricately linked to repeated cycles of WGDs, with polyploidy being a common feature across many lineages (Marhold and Lihová 2006; Franzke, et al. 2011). Notably, the ancestors of the *Orychophragmus* genus underwent a unique WGD event (Zhang, et al. 2023), followed

by rediploidization and speciation, resulting in a genome size of approximately 1.3

Gb—larger than that of other Brassicaceae species (Johnston, et al. 2005; Lysak, et al.

2008). Recently, *Orychophragmus* species have garnered attention as early-flowering

ornamental plants and as promising industrial oilseed crops, owing to their high

dihydroxy fatty acids content. This unique trait provides superior lubrication properties

and ensures broad adaptability to diverse environmental conditions (Li, et al. 2018;

Wang, et al. 2021; Huang, et al. 2023; Jia, et al. 2024). Moreover, their close

evolutionary relationship with *Brassica* species—renowned for their ease of

hybridization (Li and Heneen 1999; Li and Ge 2007)—makes *Orychophragmus* species

valuable germplasm resources for *Brassica* genetics and breeding.

Despite their potential importance, the phylogenetic relationships within the

Orychophragmus genus remain poorly understood. Currently, six species are

recognized in this genus: *O. violaceus*, *O. longisiliqus*, *O. zhongtiaoshanus*, *O.*

taibaiensis, *O. hupehensis*, and *O. diffusus* (Hu, et al. 2015; Hu, et al. 2016; Hu, et al.

2018). These species display significant morphological variation and inhabit diverse

environments across East Asia (Fig. 1). However, previous analyses have relied on a

limited number of nuclear and chloroplast genes, resulting in ongoing debates about the

genus's evolutionary history and relationships. To address these uncertainties, it is

essential to incorporate broader genomic datasets and advanced methodologies to

construct a more robust phylogeny, with particular attention to the potential effects of

incomplete lineage sorting and introgression.

Notably, the *Orychophragmus* genus includes both widespread and regionally

distributed species. *O. violaceus*, the most widely distributed species in the genus, is

commonly cultivated as an ornamental plant and is known for its small purple flowers

that typically bloom in early spring. In contrast, *O. taibaiensis* is endemic to the

mountainous regions of the Taibai Mountains in northwest China. This endemic species

is an ideal candidate for exploring its evolutionary and demographic histories,

particularly its remarkable adaptation to the local alpine environment, in contrast to the

more widespread and closely related *O. violaceus*. Additionally, *O. taibaiensis* has been

reported to consist of both diploid ($2n = 2x = 24$) and tetraploid ($2n = 4x = 48$) plants

(Zhou, et al. 2009), although the mode of origin of the polyploids remains unclear. As

such, it also serves as an excellent model for studying the immediate evolutionary and

genomic consequences of polyploidization. A direct comparison between tetraploid

populations and their diploid progenitors could provide valuable insights into how

polyploidization influences both the short-term adaptive responses and long-term

evolutionary potential of these populations (Comai 2005; Mayrose, et al. 2011).

In this study, we integrated whole-genome sequencing and transcriptomic datasets

to explore the phylogenetic relationships and divergence order of the six species within

the genus *Orychophragmus*. We then examined the demographic and divergence

history of the widespread species *O. violaceus* and the alpine endemic *O. taibaiensis*,

comparing selection efficacy and the deleterious mutation load between the two species.

Given that *O. taibaiensis* includes both diploid and tetraploid forms, we characterized

the geographical distribution patterns of these cytotypes through extensive field

investigations and karyotype analyses, while also investigating the origin of the

tetraploid forms. Finally, we explored the genetic and gene expression changes

associated with polyploidization in *O. taibaiensis* populations to gain deeper insights

into the evolutionary consequences of whole-genome duplication and its potential

short- and long-term selective effects on the ecology and evolution of these populations.

2 Materials and Methods

2.1 Taxon sampling, genomic sequencing and genetic data collection

After integrating newly sequenced genomic datasets from this study with transcriptomic datasets from previous research (Zhong, et al. 2019), we collected genomic and/or transcriptomic data for 52 individuals, representing all six *Orychophragmus* taxa: 7 *O. longisiliqus*, 13 *O. zhongtiaoshanus*, 10 *O. taibaiensis*, 1 *O. hupehensis*, 6 *O. diffusus*, and 14 *O. violaceus*. Additionally, we included one species of *Sinalliaria limprichtiana* as the outgroup. Detailed information about the data is provided in the supplemental information (Table S1). For the newly sequenced dataset, genomic DNA was extracted from leaf samples with the Qiagen DNeasy plant kit. Whole-genome paired-ends reads were generated using the T7 platform at BGI (PE150).

2.2 Phylogenomic construction

To construct the phylogenetic relationships among six *Orychophragmus* species and its relative species, we first downloaded protein sequences from six other species within the Brassicaceae family: *Aethionema arabicum*, *Capsella rubella*, *Arabidopsis lyrata*, *Arabidopsis thaliana*, *Raphanus sativus* and *Brassica rapa*. We then performed de novo transcriptome assembly using Trinity v.2.8.5 (Grabherr, et al. 2011) with default parameters applied to the filtered RNA-Seq reads processed by fastp v.0.20.1 (Chen, et al. 2018) for an individual from each species. We retained only the longest isoform for each gene and discarded redundant contigs using CD-HIT v4.8.1 (Fu, et al. 2012). Then, we utilized TransDecoder v.5.26.3 (Haas, et al. 2013) to identify protein-coding regions and subsequently used OrthoFinder v.2.5.4 (Emms and Kelly 2019) to recover single-copy genes. These single-copy genes were then used as references to extract the corresponding reads from each genome-resequenced individual and

assemble each gene for each individual using aTRAM v.2.4.0 (Allen, et al. 2017).

Finally, we used 514 single-copy genes longer than 300 bp for phylogenetic analysis.

We aligned the amino acid sequences using MAFFT v7.475 (Katoh and Standley 2013)

and then converted these alignments to codon alignments using PAL2NAL (Suyama, et

al. 2006). After removing spurious sequences and poorly aligned regions with trimAl

v1.4.rev22 (Capella-Gutiérrez, et al. 2009), the remaining alignments were combined

into a supergene matrix for phylogenetic construction using RAxML v8.2.8(Stamatakis

2014) under the GTRCAT model. To estimate divergence times between species or

clades, we employed the MCMCTree program implemented in the PAML package

v4.10.0 (Yang 2007; dos Reis, et al. 2012), utilizing fossil data sourced from TimeTree

(Kumar, et al. 2017). Furthermore, we performed coalescent-based species tree

inference using ASTRAL v.5.15.5 (Yin, et al. 2019) and visualized gene trees and

species tree discordances using DensiTree.v2.2.7 (Bouckaert 2010).

We further used population genomic data to cofirm the interspecific relationship

within *Orychophragmus.* Whole-genome resequencing data and transcriptome data

were combined to extract variant sites. For raw genomic resequencing and

transcriptome sequencing reads, we used fastp v.0.20.1 (Chen, et al. 2018) for adapter

removal and to trim bases from the start or end of reads if the base quality was <20.

Reads shorter than 36 bases after trimming were further discarded. After quality control,

all high-quality genomic resequencing reads were mapped to our new de novo

assembled *O. violaceus* (Jia, et al. 2023) genome using the BWA-MEM algorithm of

bwa v.0.7.17 (Li 2013) with default parameters, whereas the filtered transcriptome

sequencing high-quality reads were mapped using HISAT2 v. 2.2.1 (Kim, et al. 2019).

All generated bam files were subsequently sorted with SAMtools v.1.9 (Li, et al. 2009),

and PCR duplicates were marked using the MarkDuplicates tool from Picard v.2.18.21

(http://broadinstitute.github.io/picard/, last accessed March 16, 2022). Genetic variants

(SNP calling) were identified using the Genome Analysis Toolkit (GATK) v.4.2.5.0

(DePristo, et al. 2011) through its subcomponents HaplotypeCaller, CombineGVCFs,

and GenotypeGVCFs. A merged VCF file including "all sites" (both variant and non-

variant sites) were generated using the 'EMIT_ALL_SITES' flag. After combining the

GVCF files from the resequencing and transcriptome datasets, the GenotypeGVCFs

tool was applied to genotype the variants, which was followed by applying a strict set

of   filtering   criteria: 1)   Heng   Li's   SNPable   tool

(http://lh3lh3.users.sourceforge.net/snpable.shtml) was used to mask genomic regions

where reads were not uniquely mapped. To achieve this, the reference genome was

divided into overlapping 100-mers, which were then aligned back to the genome using

"bwa aln -R 1,000,000 -O 3 -E 3". Only the sites (712,157,434 out of 1,285,736,775

sites) in which all 100-mers mapped uniquely and without mismatches were retained

for downstream analyses; 2) SNPs with more than two alleles (>2), read depth (DP) <

5, or read depth greater than three times the average sequencing depth were removed;

3) SNPs with quality by depth (QD) < 2.0, Fisher Strand value (FS) >60, mapping

quality (MQ) < 20, MQRankSum < -12.5 and ReadPosRankSum < -8.0 were filtered;

4) SNPs with a missing rate >20% were excluded. Finally, we used the Degeneracy tool

to   identify   four-fold   and   zero-fold   degenerate   sites

(https://github.com/tvkent/Degeneracy). Only four-fold degenerate sites were retained

to create a robust intersect dataset after merging resequencing and transcriptome

datasets with publicly available transcriptomic data from *Sinalliaria limprichtiana*

(SRA ID: SRR6441722), which was used as an outgroup, and a total of 93,969 SNPs

were used for population genomic phylogenetic constructions.

Neighbor-joining (NJ) phylogenetic trees were first constructed using PLINK

v.1.90 (Purcell, et al. 2007) with the parameter distance 1-ibs to calculate the pairwise

identify-by-state (IBS) genetic distance matrix, followed by MEGAZ (Kumar, et al.

2018) for tree construction. And maximum likelihood (ML) phylogenetic trees were

constructed using IQ-TREE v.2.2.0.3 (Nguyen, et al. 2015). For the ML tree,

ModelFinder (Kalyaanamoorthy, et al. 2017) was used to select the best-fitting

substitution model (-B 1000 -m MFP) with 1000 ultrafast bootstraps. All trees were

visualized using FigTree v1.4.4 (https://tree.bio.ed.ac.uk/software/Fig.tree).

Additionally, to evaluate and test for the possibility of introgression among these

Orychophragmus species, the ABBA-BABA tests were constructed using the Dtrios

program in Dsuite (Malinsky, et al. 2021) to calculate the Patterson's D-statistics and

corresponding test values (Z-scores) (Patterson, et al. 2012). The D-statistics (D) were

useful to detect introgression, while the Z-scores (Z) were used to assess statistical

significance, with an absolute value of $Z > 3$ often considered as a critical threshold for

gene flow (Excoffier, et al. 2013; Martin, et al. 2015).

2.3 Demographic history analyses of *O. violaceus* and *O. taibaiensis*

To reconstruct the historical demography of the widespread species *O. violaceus*

and the locally endemic species *O. taibaiensis*, species distribution models (SDMs)

(Elith and Leathwick 2009) were developed using MAXENT v.3.4.4 (Phillips and

Dudík 2008; Merow, et al. 2013). These models were applied to evaluate shifts in

suitable habitat ranges across different time periods for both species. To achieve this,

current native distribution records for *O. violaceus* (445 records) and *O. taibaiensis* (91

records) were collected from the Chinese Virtual Herbarium (CVH,

https://www.cvh.ac.cn/) and field investigations. Environmental layers for 19

biologically significant climate variables (BIO1–BIO19) at a 2.5-arc-minute resolution

were obtained from the WorldClim database (http://www.worldclim.org) (Hijmans, et

al. 2005) for three time periods: the present (1970–2000), the Mid Holocene (~6,000

years ago), and the Last Glacial Maximum (~20,000 years ago). To minimize

overfitting, only climate variables with Pearson correlation coefficients below 0.8 ($r <$

0.8) were included in the SDM analysis. The accuracy of the SDMs was evaluated using

the area under the curve (AUC) of the receiver operating characteristic (ROC) plot

generated by Maxent (Fawcett 2006), which is a threshold-independent and prevalence-

insensitive metric.

To further investigate the demographic history of the two species, we initially

applied the Pairwise Sequentially Markovian Coalescent (PSMC) method (Li and

Durbin 2011) to infer the historical effective population size (N_e) dynamics. The

analysis was conducted using the parameters 'N25 -t15 -r5 -p "4+25×2+4+6"' for both

species, with 100 bootstrap replicates performed. Assuming a mutation rate of $8.22 \times$

10^{-9} mutations per site per generation (Beilstein, et al. 2010) and a generation time of 1

year, we converted the coalescent-scaled time into absolute time measured in years. To

gain deeper insights into recent population dynamics, particularly over the past 10,000

years, we employed the Multiple Sequentially Markovian Coalescent approach

(MSMC2) (Schiffels and Durbin 2014). This analysis was performed on phased whole-

genome sequences that were generated by Beagle v.4.1 (Browning and Browning 2009)

from four individuals (representing eight haplotypes) for each species. For both species,

MSMC2 was run on all possible individual configurations, and the medians and

standard deviations of N_e changes were subsequently estimated.

Subsequently, we inferred the divergence history between *O. violaceus* and *O.*

taibaiensis using a coalescent simulation-based approach implemented in *fastsimcoal*2

v.27 (Excoffier, et al. 2013; Excoffier, et al. 2021). From the whole-genome

resequencing dataset of the two species, we utilized only fourfold degenerate sites to

construct the two-dimensional site frequency spectrum (SFS) via easySFS

(https://github.com/isaacovercast/easySFS). We evaluated twelve demographic models,

each representing different scenarios, including divergence with or without gene flow,

and constant or variable population sizes following divergence (Fig. S6). For each

model, we performed 50 independent runs, with each run consisting of 40 cycles and

100,000 coalescent simulations. Model comparisons were based on the maximum

likelihood values from the 50 runs, further assessed using Akaike weights and

Δlikelihood. The model with the highest Akaike weight was selected as the optimal

demographic scenario. To estimate confidence intervals, we conducted parametric

bootstrapping with 100 bootstrap replicates, each including 50 independent runs. As in

the PSMC and MSMC2 analyses, we assumed a mutation rate of 8.22×10^{-9} mutations

per site per generation (Beilstein, et al. 2010) and a generation time of 1 year.

2.4 Assessment of selection efficiency and divergent selection signatures between

O. violaceus* and *O. taibaiensis

Given the differences in demographic history and effective population sizes

between *O. violaceus* and *O. taibaiensis*, we compared the efficacy of natural selection

and the deleterious genetic load between the two species. First, we calculated and

compared the genome-wide ratio of 0-fold nonsynonymous to 4-fold synonymous

nucleotide diversity ($\pi_{0\text{-fold}}/\pi_{4\text{-fold}}$) using pixy v1.0.4 (Korunes and Samuk 2021) across

100-kb nonoverlapping window. Next, we used the folded site frequency spectrum

(SFS) of 0-fold nonsynonymous and 4-fold synonymous sites to estimate the

distribution of fitness effects (DFE) for new nonsynonymous mutations using DFE-

alpha v2.16 (Keightley and Eyre-Walker 2007). A demographic model with a stepwise

population size change was fitted to the neutral SFS, and the estimated parameters were

incorporated to infer the fitness effects of new deleterious mutations and the strength

of purifying selection ($N_e s$) for each species. Confidence intervals (95% CIs) for all

estimates were generated using 200 bootstrap replicates, resampling across all sites

within each class and excluding the top and bottom 2.5% of replicates.

Furthermore, to investigate the biological significance of genomic regions

potentially under divergent selection during the divergence of *O. violaceus* and *O.*

taibaiensis, we identified candidate divergent selection regions using the XP-CLR

method (Chen, et al. 2010). Specifically, we calculated cross-population composite

likelihood ratio (XP-CLR) scores between the two species within 5-kb non-overlapping

windows. Genomic regions with scores in the top 1% (2397 windows) were identified

as candidate selective regions. To validate these regions, we calculated independent

estimates of inter-species genetic divergence (F_{ST}) and intra-species Tajima's *D* values,

confirming signatures of divergent selection. Finally, we extracted the genes (340)

located within the candidate regions and performed Gene Ontology (GO) enrichment

analysis using the topGO R package (https://bioconductor.org/packages/topGO/) to

investigate the biological significance of these regions.

2.5 Determination of diploid and tetraploid distribution across *O. taibaiensis*

populations

To thoroughly investigate the geographical distribution of diploid and tetraploid

cytotypes across the natural populations of *O. taibaiensis*, seeds were collected from

94 individuals across seven populations, covering the entire known distribution of *O.*

taibaiensis identified in the field. The seeds were germinated on wet filter paper in Petri

dishes at 25°C. Root meristems were carefully excised from germinating seeds and pre-

treated with a 0.1% colchicine solution at 25°C for 3 hours. The samples were then

washed and fixed in a 3:1 solution of ethyl alcohol and acetic acid at 4°C for 3 hours.

Subsequently, the samples were rinsed in distilled water and immersed in 1 M HCl at

37°C for 45 minutes to facilitate hydrolysis. After hydrolysis, the root tips were washed

in distilled water and stained with a modified carbol-fuchsin solution at room

temperature for more than 3 hours. Finally, the meristems were dissected and squashed

onto glass slides, and chromosomal counts were analyzed using a microscope under oil

immersion (×1000 magnification) and photographed.

To further investigate and differentiate the origin of polyploidy (autopolyploidy or

allopolyploidy) in the identified diploids and tetraploids of *O. taibaiensis*, we selected

five diploids and five tetraploids for high-depth whole-genome resequencing (average

depth ~52.48×). First, we utilized nQuire (Weiß, et al. 2018) to confirm the ploidy

levels of the sequenced individuals. The "denoise" subcommand was applied to reduce

baseline noise, and sequencing reads were mapped back to genome assemblies to

evaluate read depths and allele frequencies. The density distribution of the three allele

types (reference, alternative, and both) was calculated to validate ploidy levels. Next,

we integrated GenomeScope and Smudgeplots (Ranallo-Benavidez, et al. 2020) to

distinguish between autotetraploids and allotetraploids. This was achieved by analyzing

patterns of nucleotide heterozygosity for polyploid samples, following the construction

of *K-mer* frequency distributions using Jellyfish v.2.2.9 (Marçais and Kingsford 2011)

from the resequencing data. Finally, we re-performed variant calling for the five

resequenced individuals of tetraploid *O. taibaiensis* using GATK, setting the parameter

"--ploidy 4". The relative proportions of different genotypes (e.g., Aaaa, AAaa, aaaA)

were calculated to further verify whether the tetraploids were of autotetraploid or

allotetraploid origin.

2.6 Population genomic analysis and assessment of genetic load for diploid and

tetraploid individuals of *O. taibaiensis*

To construct the phylogenetic relationships of *O. taibaiensis* at different ploidy

levels, we re-performed variant calling and filtering, following the method described in

section 2.2, for the ten high-depth resequenced individuals. First, we used PLINK

v.1.90 (Purcell, et al. 2007) with the parameter distance 1-ibs to calculate the pairwise

identify-by-state (IBS) genetic distance matrix, quantifying the relatedness among the

five diploids and five tetraploids. Based on the resulting distance matrix, we employed

MEGAZ (Kumar, et al. 2018) to construct an unrooted NJ tree among individuals with

different ploidy levels. Next, we utilized *fastsimcoal2* (Excoffier, et al. 2021) to infer

the divergence history of the two cytotypes based on a two-dimensional joint SFS of

fourfold degenerate sites. Whole-genome resequencing data from *O. longisiliqus* and

O. zhongtiaoshanus were included as outgroups to infer the derived states of alleles

using the est-sfs software (Keightley and Jackson 2018). The analysis was performed

under two models: one accounting for gene flow and one assuming no gene flow (see

Fig. S14a, S14b). The detailed procedures followed those outlined in section 2.3.

To test for the potential effects of polyploidization on the efficacy of purifying

selection and the accumulation of deleterious genetic load, we estimated and compared

the ratio of 0-fold nonsynonymous to 4-fold synonymous nucleotide diversity ($\pi_{0\text{-fold}}/\pi_{4\text{-fold}}$)

between diploid and tetraploid individuals of *O. taibaiensis*. Additionally, we

examined and compared the distribution of fitness effects (DFE) of new mutations

across both ploidy levels. The intensity of purifying selection ($N_e s$) at 0-fold

nonsynonymous sites was estimated using DFE-alpha v2.16 (Keightley and Eyre-

Walker 2007) by fitting a stepwise population size change model to account for

differences in the SFS of 0-fold nonsynonymous sites and putatively neutral 4-fold

synonymous sites. To evaluate the potential effects of the diploid assumption in these

analyses, we re-performed variant calling for tetraploid individuals using GATK,

setting the parameter "--ploidy 4". For each tetraploid individual, we randomly

subsampled two alleles from the four alleles per site to generate six independent

subsampling datasets. These datasets were then used for associated analyses, including

fastsimcoal2 modeling, genetic load estimation, and DFE analysis. This approach

ensured robust comparison between diploid and tetraploid cytotypes while accounting

for the polyploid nature of the tetraploid individuals.

2.7 Differential gene expression analysis between the two cytotypes of *O.*
338
339   *taibaiensis*

To investigate how polyploidy influences transcriptomic changes, seeds of diploid

and tetraploid *O. taibaiensis* collected from two geographic regions (Fig. 3a red

triangles) were grown under identical conditions in a growth chamber. The environment

was controlled with a 16/8 h light/dark photoperiod, a constant temperature of 25°C,

and a light intensity of $100\,\mu\mathrm{mol\,m^{-2}\,s^{-1}}$. After six months of growth, leaf and root

tissues were collected from plants with similar growth characteristics for both cytotypes.

Three biological replicates were collected per cytotype. Harvested materials were

frozen in liquid nitrogen and stored at -80°C until RNA extraction. Total RNA was

extracted using the TIANGEN RNA Kit following the manufacturer's protocol.

Sequencing libraries were constructed and sequenced on the DNBSEQ-T7 platform

using paired-end sequencing strategy.

After filtering and trimming RNA reads using fastp v.0.20.1 (Chen, et al. 2018)

with default parameters, the clean reads were mapped to the *O. violaceus* genome (Jia,

et al. 2023) using HISAT2 v.2.2.1 (Kim, et al. 2019). Gene expression levels were

quantified with StringTie v1.3.6 (Pertea, et al. 2015; Kovaka, et al. 2019), and

normalized expression levels were calculated as transcripts per million (TPM). Out of

52,812 annotated genes in the reference genome (Jia, et al. 2023), a total of 37,607

genes (TPM > 0 for each sample) were expressed across both leaf and root tissues in

diploid and tetraploid samples. For differential expression analysis, the R package

DESeq2 (Love, et al. 2014) was used, with thresholds of |log2 fold change (FC)| ≥ 2

and adjusted *p*-value < 0.001 to identify significant differentially expressed genes

(DEGs). Genes with |log2 fold change (FC)| < 1 or adjusted *p*-value > 0.05 were

classified as non-differentially expressed genes (NDEs). The DEGs were further

clustered and visualized using the ClusterGVis package

(https://github.com/junjunlab/ClusterGVis), which employs the "mfuzz" clustering

method to group genes based on their expression patterns. To identify the biological

functions enriched among the DEGs associated with polyploidy in *O. taibaiensis*, GO

enrichment analysis was performed using the topGO R package

(https://bioconductor.org/packages/topGO/).

2.8 Selection on genes exhibiting differential expression between cytotypes

To examine and compare differences in the strength and direction of natural

selection on DEGs and NDEs associated with polyploidization in diploids and

tetraploids of *O. taibaiensis*, we first estimated pairwise relative genetic divergence

(F_{ST}), and absolute genetic divergence (d_{xy}) within and between resequenced

individuals of different cytotypes for the DEGs and NDEs gene sets using PIXY v1.0.4

(Korunes and Samuk 2021). Next, we calculated the ratio of zero-fold nonsynonymous

to four-fold synonymous nucleotide diversity to compare deleterious mutation loads

between the two gene sets. Finally, we assessed the strength of selective constraint on

the two gene sets by modeling the distribution of fitness effects (DFE). As described

earlier, the strength of purifying selection (N_es) on zero-fold nonsynonymous sites was

estimated for each gene set in both diploids and tetraploids, using four-fold

synonymous sites as a neutral reference, with DFE-alpha v2.16 (Keightley and Eyre-

Walker 2007). For tetraploid individuals, six additional independent datasets, generated

by randomly sampling two alleles from the four alleles, were applied in all analyses in

this section.

3 Results

3.1 Phylogenetic and evolutionary relationships among species in the genus

Orychophragmus

To investigate the phylogenetic relationships within the genus *Orychophragmus*,

we incorporated seven additional species from the Brassicaceae family and utilized 514

single-copy genes to construct phylogenetic trees using both concatenated and

coalescent approaches.   Phylogenetic analysis based on single-copy genes revealed the

monophyly of *Orychophragmus*, with its closest relative being the genus *Sinalliaria*

(Fig. 1a). According to MCMCTree, the divergence between *Orychophragmus* and its

closest relative *Sinalliaria* was estimated to be 15.43 million years ago (MYA, 95%

highest posterior density: 10.23-21.69 MYA). Within the clade *Orychophragmus*, the

concatenated analysis produced a strongly supported topology of (outgroups, (*O.*

longisiliqus, (*O. zhongtiaoshanus*, (*O. taibaiensis*, (*O. violaceus*, (*O. diffusus, O.*

hupehensis)))))) (Fig. 1a). This topology was also supported by the ASTRAL coalescent

approach (Fig. S1a), although some branches exhibited weak support values. To further

explore this, we used DensiTree as a visualization tool to display and quantify the

concordance and discordance between individual gene trees and the species tree, which

revealed significant inconsistencies between many individual gene trees and the species

tree within the *Orychophragmus* clade (Fig. S1b).

To further validate the phylogenetic relationship within the genus *Orychophragmus*,

we integrated population-level resequencing and transcriptomic datasets, using

Sinalliaria as the outgroup to construct phylogenies. The phylogenetic relationships

within *Orychophragmus*, inferred using both maximum likelihood (ML) and neighbor-

joining (NJ) methods (Fig. S2), were consistent with those derived from single-copy

gene analyses (Fig. 1a, Fig. S1). Given the observed inconsistencies in individual gene

trees within *Orychophragmus*, we used ABBA-BABA analyses to assess potential gene

flow between these species. Specifically, we systematically tested for signatures of

introgression by calculating Patterson's *D*-statistics and corresponding test values (*Z*-

scores) (Patterson, et al. 2012). The results provide strong evidence of pervasive

historical introgression among the *Orychophragmus* species (Fig. S3), which could

partially explain the observed inconsistencies in the individual gene trees of

Orychophragmus.

3.2 Demographic histories and divergent selection between *O. violaceus* and *O.*

taibaiensis

Given the unique features of *O. taibaiensis*—the only species in the genus reported

to consist of both diploids and tetraploids as well as being the endemic species found

at the highest altitudes within the genus (Fig. S4)—we specifically explored and

compared the demographic histories of this species with the closely related widespread

species *O. violaceus*. To achieve this, we first applied species distribution models

(SDMs) to reconstruct the suitable habitats of *O. violaceus* and *O. taibaiensis* across

three evolutionary periods: the present, the Middle Holocene (MH, ~6 ka), and the Last

Glacial Maximum (LGM, ~21–18 ka). Our analysis revealed that the suitable habitat

of *O. violaceus* expanded significantly from the LGM and MH to the present. In

contrast, the suitable habitat of *O. taibaiensis* remained relatively stable, being

primarily confined to the Qinling-Daba mountain regions (Fig. 2a). Furthermore, over

time, its range became increasingly restricted to the Taibai Mountain region, from the

LGM and MH to the present.

To assess the long-term effective population size (N_e) dynamics of these two

species, we applied the pairwise sequential Markovian coalescent (PSMC) method (Li

and Durbin 2011). The results revealed that both species experienced a reduction in N_e

from the Riss glaciation to the LGM (Fig. 2b). Consistent with the broader distribution

range of *O. violaceus* predicted by the SDMs, the N_e of *O. violaceus* was generally

much larger than that of *O. taibaiensis* (Fig. 2b). Since PSMC can only estimate

population dynamics up to around 10,000 years ago, we also employed MSMC2 to

focus specifically on the population histories of these two species over the latest 10,000

years. The results indicated that the *O. violaceus* population underwent a more

significant recovery following LGM when compared to *O. taibaiensis* (Fig. 2b; Fig.

S5).

To further infer the divergence history of the two species, we employed a

coalescent simulation-based approach using *fastsimcoal*2 (Excoffier, et al. 2013;

Excoffier, et al. 2021). Twelve models were evaluated (Fig. S6), differing in the

presence or absence of post-divergence gene flow and changes in population size

following species divergence. The best-fitting model (Model 12 in Fig. S6; Table S2)

suggested that *O. violaceus* and *O. taibaiensis* diverged approximately 5.25 MYA (95%

confidence interval (CI) = 1.40–7.33 MYA) (Fig. 2c; Table S3), consistent with the

divergence time estimated from phylogenetic analysis (Fig. 1a). Furthermore, the

model revealed divergent patterns of changes in effective population sizes and

asymmetric gene flow between the species following their divergence, although no

gene flow has occurred between the two species within the recent 5.22 thousand years

ago (KYA) (95% CI = 5.02–5.79KYA) (Fig. 2c, Table S2). Additionally, the model

inferred that *O. taibaiensis* experienced a recent population reduction ($N_e = 1.59 \times 10^5$,

$[1.45 \times 10^5 – 1.99 \times 10^5]$), in contrast to *O. violaceus*, which underwent population

expansion in both effective population size ($N_e = 1.08 \times 10^6$ $[8.45 \times 10^5 – 1.29 \times 10^6]$) and

distribution range (Fig. 2a,c).

To further compare the likely influence of different demographic histories on the

efficacy of selection and the accumulation of mutational load between the two species,

we estimated the ratio of nucleotide diversity at 0- to 4-fold degenerate sites. Consistent

with the expectation that species with smaller population sizes have a reduced efficacy

of selection to purge deleterious mutations at 0-fold sites (Lynch, et al. 1995), we

observed that the ratio of 0- to 4-fold nucleotide diversity was significantly elevated in

O. taibaiensis compared to *O. violaceus* (Fig. 2d). Similarly, the distribution of fitness

effects (DFE) analysis revealed that *O. taibaiensis* exhibited weaker purifying selection

against strongly deleterious nonsynonymous mutations ($N_e s > 100$) compared to *O.*

violaceus (Fig. 2e). Moreover, given that *O. taibaiensis* contains both diploid and

tetraploid cytotypes, we re-analyzed and compared the population history divergence,

genetic load, and DFE solely between *O. violaceus* and diploid *O. taibaiensis*. These

results were highly consistent with the broader analysis, further supporting the

conclusion that *O. taibaiensis* exhibits reduced selection efficacy and a higher

mutational load compared to *O. violaceus* (Fig. S7b).

Lastly, we applied and calculated XP-CLR statistics to identify potential divergent

selection regions between the two species. In total, we detected 2,397 outlier windows

in the top 1% of XP-CLR scores (Fig. 2f). Further gene ontology (GO) enrichment

analyses of genes within the candidate selective regions revealed significant enrichment

in GO terms such as "cellular macromolecule metabolic process" and "response to

reactive oxygen species" (Fig. S8a; Table S4), suggesting their association with

divergent adaptation to different altitudinal environments between the two species.

Many genes known to be involved in stress and defense responses were identified

within these regions. For example, the *Arabidopsis* orthologous gene *PAD4*, which

plays a critical role in salicylic acid signaling and resistance gene-mediated plant

disease resistance(Zeng, et al. 2023; Yu, et al. 2024), was found within these regions.

In addition to the significant signals of XP-CLR, we observed substantially increased

inter-species genetic divergence (F_{ST}) and reduced intra-species Tajima's *D* values (Fig.

S8b), providing strong evidence of divergent selection within these genic regions.

Similarly, the *Arabidopsis* orthologous gene *PEC2*, which is involved in responding to

environmental stimuli such as jasmonic acid, light, and wounding (Völkner, et al. 2021),

was also identified, highlighting its role in stress responses and adaptive signaling

pathways. We found substantially increased F_{ST} and reduced Tajima's *D* values in both

species around these genic regions (Fig. S8b). Similar patterns were observed for other

genes related to fundamental processes crucial to plant growth, development, and

epigenetic regulation, such as genes *DME* (Khouider, et al. 2021) and *XTH7*

(Volyanskaya, et al. 2023) (Fig. S8d,e)

3.3 Cytotype diversity and distribution in natural populations of *O. taibaiensis*

Although *O. taibaiensis* was previously reported to have both diploid and tetraploid

cytotypes locally distributed in the Taibai Mountains of central China (Tan, et al. 1998;

Zhou, et al. 2008), the spatial overlap and distribution patterns of populations with

different ploidy levels remain unclear. To address this, we determined the ploidy levels

of *O. taibaiensis* in a relatively large sample set of 94 individuals from various locations.

Chromosome number determination confirmed the findings of earlier studies (Zhou, et

al. 2008) and revealed variation in chromosome size. Specifically, we identified two

ploidy levels, with chromosome counts of 2n = 24 for diploids (Fig. 3b) and 2n = 48

for tetraploids (Fig. 3c). Cytotype distribution analysis indicated that individuals with

different ploidy levels were primarily separated by local mountain barriers within

relatively limited geographical ranges (Fig. 3a). To further investigate whether there

was environmental differentiation among cytotypes, we performed a principal

component analysis (PCA) using five uncorrelated environmental variables derived

from 19 bioclimatic factors (bio1-bio19) and elevation (Fig. S9a,c, Table S5, Table S6).

Our results revealed little to no evidence of niche differentiation between cytotypes,

with only subtle variation observed along environmental and climatic gradients (Fig.

S9).

Next, we selected five diploid and five tetraploid individuals of *O. taibaiensis* for

high-depth whole-genome resequencing. We first employed nQuire (Weiß, et al. 2018),

a statistical method designed to determine the most plausible ploidy model based on

the distribution of base frequencies in the sequencing data. The results from nQuire

confirmed the diploid and tetraploid forms of *O. taibaiensis* that we had previously

identified (Fig. S10a, b). Specifically, the read depth density distribution of the three

alleles (alternative, reference and both) in diploid individuals was close to 1:1:2 (Fig.

S10c), while in tetraploid individuals, it was approximately 1:3:4 (Fig. S10d). To

differentiate between autopolyploidization and allopolyploidization, we utilized

Smudgeplots (Ranallo-Benavidez, et al. 2020) to visualize the expected allele ratio

patterns (Fig. S11). We found that the AAAB pattern was more prominent than AABB

for tetraploids, consistent with the expectation of an autopolyploid origin. Additionally,

genome-wide heterozygosity analysis using GenomeScope v.2.0 revealed a distribution

of aaab (3.52%) > aabb (1.96%) (Fig. S12a), further supporting the notion of

autotetraploids. Lastly, we recalled SNPs based on the tetraploid model for each of the

five resequenced tetraploids and calculated the proportions of various genotypes. The

results indicated that the AAAa genotype was significantly more prevalent than AAaa

across the entire genome (Fig. 3d; Fig. S12b-f), reinforcing the hypothesis of

autotetraploidy in the polyploid *O. taibaiensis*.

3.4 Origin history of autotetraploids and the evolutionary consequences of short-

term polyploidization in *O. taibaiensis*

To investigate the origin and genetic differentiation between diploid and

autotetraploid *O. taibaiensis*, we constructed an unrooted tree using ten highly deep-

sequenced individuals from both ploidy levels. The analysis revealed that the individuals formed two distinct clusters, albeit with short genetic distances and low divergence (Fig. 3e). Using *fastsimcoal2*, we identified the gene flow model as the best fit for the divergence history and estimated that diploids and tetraploids diverged approximately 328 KYA (95% CI: 326–468 KYA). The contemporary N_e of diploids was estimated to be $2.53×10^5$ (95% CI: $2.40×10^5$–$3.61×10^5$), which is slightly larger than that of autotetraploids, estimated at $1.52×10^5$ (95% CI: $1.45×10^5$–$2.06×10^5$) (Fig. 3f, Table S7).

To further explore the effects of ploidy on the efficacy of purifying selection, we evaluated and compared the DFE between diploids and tetraploids. The results showed that diploids exhibited a significantly higher proportion of loci under strong purifying selection (N_es > 100) compared to autotetraploids. This finding suggests relaxed purifying selection acting on nonsynonymous sites in the tetraploid population compared to the diploid population (Fig. 3g). These results remained robust when we randomly subsampled two out of four alleles per site from the genotypes of tetraploid population to account for potential biases introduced by genotype calling (Fig. S13). However, it is important to note that such biases cannot be fully resolved, as a diploid model was assumed for DFE estimation. Additionally, we calculated the ratio of overall 0- to 4-fold nucleotide diversity (Fig. S14c-S14i) and found that the ratio was significantly higher in autotetraploids compared to diploids. These results further suggest that the tetraploid population has accumulated a higher proportion of deleterious mutations, which aligns with the findings from *fastsimcoal2* and DFE analysis, showing that the tetraploid population, with higher genetic load, exhibits a smaller effective population size and lower purifying selection efficiency.

3.5 Gene expression changes and associated signatures of selection upon

polyploidization in *O. taibaiensis*

To explore transcriptional changes and responses to WGD, we examined gene

expression differences using RNA-seq data from diploids and autotetraploids of *O.*

taibaiensis in both leaf and root tissues. Principal component analysis (PCA) clearly

distinguished samples based on tissue type and cytotype (Fig. S15a). To specifically

investigate gene expression changes associated with polyploidization, we identified

differentially expressed genes (DEGs) between diploids and autotetraploids. While the

majority of genes showed no significant change in expression (NDEs), we identified

3,254 and 3,422 DEGs in leaf and root tissues, respectively (Fig. S15b), with 1,910

DEGs shared between the two tissues.

GO enrichment analysis of these DEGs revealed that genes differentially

expressed between diploids and tetraploids are highly enriched in terms related to signal

transduction, cell communication, defense response, regulation of gene expression, and

circadian rhythm (Fig. 4a). For instance, genes downregulated in tetraploids, such as

SNI1 and *CERK1*, are involved in gene transcription, DNA recombination, and defense

signaling (Fig. 4b). Notably, *SNI1* in *Arabidopsis* plays a crucial role in preventing

errors during meiotic recombination in plants (Liu, et al. 2018; Zhu, et al. 2021).

Similarly, *α-DOX1* and *ADC* are key players in metabolic processes, with *α-DOX1*

participating in fatty acid alpha-oxidation and *ADC* being involved in polyamine

biosynthesis, both of which are essential for maintaining cellular integrity and stress

responses (Vicente, et al. 2012; Rossi, et al. 2015) (Fig. 4b). The significantly reduced

expression of these genes in tetraploids may be attributed to genomic instability and the

regulatory complexity caused by gene dosage effects in tetraploids. Conversely, in

autotetraploid plants, the upregulated genes are primarily involved in functions critical

for maintaining genomic stability and responding to environmental stresses. For

example, *EXT3* is essential for plant cell wall organization, while *RECA2* is involved

in DNA repair (Cannon, et al. 2008; Saha, et al. 2013; Odahara, et al. 2017).

Additionally, *LecRK* and *LNK3* have been reported to play roles in cellular responses

to salicylic acid, defense against pathogens, and the regulation of circadian rhythms

(Bouwmeester and Govers 2009; Kidokoro, et al. 2023). These findings may reflect a

broader theme of adaptation and stress response, highlighting the ability of tetraploids

to cope with environmental challenges (Fig. 4a, 4b).

To further investigate the association between expression divergence and sequence

divergence following genome doubling in *O. taibaiensis*, we compared both the relative

(F_{ST}) and absolute (d_{xy}) genetic divergence between the two cytotypes for DEGs and

NDEs. We observed that DEGs exhibited significantly higher d_{xy} compared to NDEs,

while no significant differences were observed for F_{ST}. These findings suggest that

genes differentially expressed between diploids and tetraploids have accumulated a

significantly higher average number of nucleotide differences between ploidies, while

nucleotide diversity within each ploidy type remained relatively stable (Fig. 5a,b).

Additionally, we compared the ratio of nucleotide diversity at 0-fold to 4-fold

degenerate sites (π_{0fold}/π_{4fold}) between DEGs and NDEs and found a substantially

increased π_{0fold}/π_{4fold} ratio within DEGs. This result aligns with expectations of a higher

mutation load and reduced purifying selection efficiency for these genes (Fig. 5c). DFE

analysis further reinforced these observations, revealing that DEGs showed a

significantly lower proportion of novel mutations with likely highly deleterious effects

compared to NDEs, implying relaxed purifying selection and reduced functional

constraints on these genes (Fig. 5d). Finally, these results were robust to random

subsampling of two out of four alleles per site from the genotypes of autotetraploids.

In the resampling datasets, F_{ST} values for DEGs were found to be significantly higher

than those for NDEs, although the overall trends remained consistent (Fig. S16-S18).

4 Discussion

Polyploidy resulting from WGD is widespread and has repeatedly occurred throughout the evolution of eukaryotes (Masterson 1994; Wood, et al. 2009; Van de Peer, et al. 2017). However, our understanding of its effects on shaping patterns of genomic variation and influencing the selection process remains limited. In this study, we examined the non-model plant genus *Orychophragmus*, which includes both widespread and endemic species, as well as species with variable ploidy levels. By integrating population genomics and transcriptomic datasets, we explored the evolutionary consequences of diverse population demographic histories and the inherent effects of genome doubling on the long-term evolutionary potential of populations.

Using single-copy nuclear genes and population-level SNP datasets, our phylogenomic analyses constructed a more robust species-level phylogeny for this genus compared to previous studies (Hu, et al. 2016; Zhong, et al. 2019). However, we also observed inconsistencies among individual gene trees, likely caused by introgressive hybridization between species (Morales-Briones, et al. 2020; Morales-Cruz, et al. 2021), which merits further investigation in future studies. Notably, *O. taibaiensis* is a locally endemic mountainous species that occupies the highest elevations within the genus and is the only known species to exhibit a mixed-ploidy system. Despite this, little is known about the spatial distribution of different cytotypes across landscapes or the establishment history of polyploids (Barker, et al. 2016). To address this, we first examined the divergence history of *O. taibaiensis* and the closely related, predominantly widespread species *O. violaceus* within the genus. Ecological modeling and population genomic analyses revealed that the two species have experienced divergent demographic histories following their divergence. Specifically,

O. taibaiensis underwent a more drastic population contraction and a mild recent

population size recovery, whereas *O. violaceus* maintained a more stable demographic

history. These divergent demographic trajectories have resulted in differences in

selection efficacy between the two species. The lower effective population size of *O.*

taibaiensis has likely led to the accumulation of a greater deleterious mutation load due

to reduced selection efficacy in this species (Harris and Nielsen 2016; Liu, et al. 2022).

Additionally, we explored signals of divergent selection between the two species and

identified genomic regions enriched with genes involved in responses to oxidative

stress. These genes are likely crucial for *O. taibaiensis* to survive and thrive in high

mountain regions, highlighting the need for further in-depth studies to uncover the

adaptive mechanisms of *O. taibaiensis* in such harsh environments.

The diploid-tetraploid variation in *O. taibaiensis* provides a unique study system

to investigate the origin of polyploids, the niche similarity and differentiation of

ploidies across spatial scales, and the genomic and evolutionary consequences of

genome doubling (Parisod, et al. 2010; Padilla-García, et al. 2023). For the first time,

by integrating extensive field surveys, karyotype analysis, and bioinformatics

approaches, our results provide strong evidence supporting the autotetraploid origin

model of polyploids. Additionally, we clearly delineate the geographic distribution of

the two cytotypes across spatial scales. Despite being generally separated by a mountain,

we found little environmental differentiation between the cytotypes at local spatial

scales across the distribution range of *O. taibaiensis*. Population demographic analyses

further revealed that the two cytotypes diverged approximately 328 KYA, with close

phylogenetic relationships among individuals of both cytotypes further supporting the

relatively recent autopolyploid origin of *O. taibaiensis*.

To better understand how autotetraploids evolve and are affected by selection

processes, we examined and compared the distribution of fitness effects of new

mutations and the genetic load, estimated as the ratio of 0-fold to 4-fold diversity. Our

findings suggest that polysomic masking in autotetraploids, compared to diploids,

likely reduces the efficacy of purifying selection. As a result, this relaxed purifying

selection leads to a higher accumulation of deleterious mutations in tetraploid

populations (Cheng, et al. 2018; Monnahan, et al. 2019). In the short term, the masking

of deleterious mutations and the relaxed purifying selection that follow genome

doubling may facilitate the rapid establishment and niche expansion of autotetraploid

populations (Van de Peer, et al. 2017; Baduel, et al. 2019). However, it remains

uncertain how the accumulation of deleterious mutations will balance against the

beneficial effects of masking in the long run, and how this dynamic interplay will shape

the persistence and coexistence of the two cytotypes.

Moreover, nuclear volume changes arising from polyploidization can increase

genome complexity and influence gene expression (Burns, et al. 2021; Yu, et al. 2023).

By analyzing gene expression patterns in leaf and root tissues of the two cytotypes of

O. taibaiensis, we identified a set of differentially expressed genes (DEGs) between

diploids and autotetraploids, while the majority of genes were not differentially

expressed (NDEs). Consistent with the expected nuclear volume changes, DEGs were

significantly enriched in functions related to extracellular structure organization, cell

wall assembly, and cell communication (Westermann 2021; Zhu, et al. 2024).

Additionally, genes involved in signal transduction, response to stimuli, defense

responses, regulation of primary and secondary metabolic processes, and circadian

rhythm were also enriched among DEGs. These findings provide a likely explanation

for the frequent association of polyploidization with enhanced stress resistance and

local environmental adaptation following divergence from diploid relatives (Parisod, et

al. 2010; McDaniel 2024). Strikingly, we found that genes with expression changes

between ploidy cytotypes evolve under relaxed selective constraints and accumulate

more sequence divergence compared to NDEs. This highlights a strong association

between expression and sequence changes in response to genome doubling for these

genes. Future studies are needed to uncover the potential evolutionary forces driving

this association and to explore how selection and transcriptional mechanisms jointly

respond to genome doubling, shaping the subsequent evolution of autopolyploids

(Bhaskara, et al. 2023; Louder, et al. 2024).

Importantly, there are a few caveats that must be acknowledged. First, biases may

arise because many populations genetic analyses assume a diploid model of allele

frequencies at mutation-selection-drift balance, which could affect and bias the

estimates for autotetraploids (Monnahan, et al. 2019; Yu, et al. 2023). In this study, we

called variants in both diploid and polyploid modes for the autotetraploids and also

performed random subsampling of two alleles per site to facilitate comparisons across

various datasets. While we found consistent results, we must acknowledge that we

cannot completely rule out the possibility of bias. Second, our transcriptome analyses

compared the relative expression levels of genes to the total transcriptome between the

two cytotypes. However, we did not estimate the true transcriptome sizes due to the

lack of normalization of transcripts for cell number and biomass content changes in

autotetraploids (Coate 2023; Srikant, et al. 2024). This means we cannot entirely

exclude the possibility that an overall change in the total number of transcripts occurred

following genome doubling. Nevertheless, given that the majority of genes showed

balanced expression levels between cytotypes, we believe the likelihood of substantial

changes in cell number and transcriptome sizes in autotetraploids is low.

Altogether, our findings provide a novel empirical study system to explore the

genomic consequences of genome doubling and the potential evolutionary drivers

underlying the successful establishment of newly formed autotetraploid lineages in the

local mountainous endemic species *O. taibaiensis*. Future work could focus on

sampling and sequencing a broader range of diploids and autotetraploids to better

understand the factors influencing the evolutionary potential and establishment of

polyploids, as well as their long-term coexistence with their diploid ancestors.

Acknowledgments

722   This work was supported by the National Natural Science Foundation of China

723   (32000265) and Fundamental Research Funds for the Central Universities

724   (2023SCUNL105) to J.W.

Author contributions

727   J.W. conceived and supervised the study. Q.L., C.J., R.W., Y.L., Y.H. handled the

728   sampling, material collection and performed experiments. Q.L., Z.W., X. Q., Z. Z.

729   analyzed the data, Q.L. and J.W. wrote the manuscript with the input from P.K.I and

730   J.L.. All authors approved the final version of the manuscript.

Competing interests

733   The authors declare no competing interests.

Data availability

All data needed to evaluate the conclusions in this study are present in the paper and/or

the Supplementary information. The newly generated whole-genome resequencing data

and transcriptome data of the samples produced in this study have been deposited in

the National Genomics Data Center (https://ngdc.cncb.ac.cn) under the accession

number PRJCA035915.

Reference:

Allen JM, Boyd B, Nguyen N-P, Vachaspati P, Warnow T, Huang DI, Grady PGS, Bell KC, Cronk
QCB, Mugisha L, et al. 2017. Phylogenomics from whole genome sequences using aTRAM.
Systematic Biology **66**: 786-798.

Baduel P, Quadrana L, Hunter B, Bomblies K, Colot V. 2019. Relaxed purifying selection in
autopolyploids drives transposable element over-accumulation which provides variants for local
adaptation. *Nature Communications* **10**: 5818.

Barker MS, Arrigo N, Baniaga AE, Li Z, Levin DA. 2016. On the relative abundance of autopolyploids
and allopolyploids. *New phytologist* **210**: 391-398.

Beilstein MA, Nagalingum NS, Clements MD, Manchester SR, Mathews S. 2010. Dated molecular
phylogenies indicate a Miocene origin for *Arabidopsis thaliana*. *Proceedings of the National Academy*
of Sciences **107**: 18724-18728.

Bhaskara GB, Haque T, Bonnette JE, Napier JD, Bauer D, Schmutz J, Juenger TE. 2023.
Evolutionary analyses of gene expression divergence in *Panicum hallii*: exploring constitutive and
plastic responses using reciprocal transplants. *Molecular Biology and Evolution* **40**: msad210.

Bouckaert RR. 2010. DensiTree: making sense of sets of phylogenetic trees. *Bioinformatics* **26**: 1372-
1373.

Bouwmeester K, Govers F. 2009. *Arabidopsis* L-type lectin receptor kinases: phylogeny, classification,
and expression profiles. *Journal of Experimental Botany* **60**: 4383-4396.

Browning BL, Browning SR. 2009. A unified approach to genotype imputation and haplotype-phase
inference for large data sets of trios and unrelated individuals. *The American Journal of Human*
Genetics **84**: 210-223.

Burns R, Mandáková T, Gunis J, Soto-Jiménez LM, Liu C, Lysak MA, Novikova PY, Nordborg
M. 2021. Gradual evolution of allopolyploidy in *Arabidopsis suecica*. *Nature Ecology and Evolution*.

Cannon MC, Terneus K, Hall Q, Tan L, Wang Y, Wegenhart BL, Chen L, Lamport DTA, Chen Y,
Kieliszewski MJ. 2008. Self-assembly of the plant cell wall requires an extensin scaffold. *Proceedings*
of the National Academy of Sciences **105**: 2226-2231.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**: 1972-1973.

Chen H, Patterson N, Reich D. 2010. Population differentiation as a test for selective sweeps. *Genome*

Research **20**: 393-402.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**: i884-i890.

Cheng F, Wu J, Cai X, Liang J, Freeling M, Wang X. 2018. Gene retention, fractionation and
subgenome differences in polyploid plants. *Nature Plants* **4**: 258-268.

Coate JE 2023. Beyond transcript concentrations: quantifying polyploid expression responses per
biomass, per genome, and per cell with RNA-Seq. In: Van de Peer Y ed. *Polyploidy: Methods and*
Protocols. New York, NY: Springer US, 227-250.

Comai L. 2005. The advantages and disadvantages of being polyploid. *Nature Reviews Genetics* **6**: 836-
846.

DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, del Angel
G, Rivas MA, Hanna M, et al. 2011. A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nature Genetics* **43**: 491-498.

dos Reis M, Inoue J, Hasegawa M, Asher RJ, Donoghue PCJ, Yang Z. 2012. Phylogenomic datasets
provide both precision and accuracy in estimating the timescale of placental mammal phylogeny.
Proceedings of the Royal Society B: Biological Sciences **279**: 3491-3500.

Elith J, Leathwick JR. 2009. Species Distribution Models: Ecological explanation and prediction across
space and time. *Annual Review of Ecology, Evolution, and Systematics* **40**: 677-697.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics.
Genome Biology **20**: 238.

Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC, Foll M. 2013. Robust demographic
inference from genomic and SNP data. *PLoS Genetics* **9**: e1003905.

Excoffier L, Marchi N, Marques DA, Matthey-Doret R, Gouy A, Sousa VC. 2021. *fastsimcoal*2:
demographic inference under complex evolutionary scenarios. *Bioinformatics* **37**: 4882-4885.

Fawcett T. 2006. An introduction to ROC analysis. *Pattern Recognition Letters* **27**: 861-874.

Franzke A, Lysak MA, Al-Shehbaz IA, Koch MA, Mummenhoff K. 2011. Cabbage family affairs:
the evolutionary history of Brassicaceae. *Trends in Plant Science* **16**: 108-116.

Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-generation
sequencing data. *Bioinformatics* **28**: 3150-3152.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
Raychowdhury R, Zeng Q, et al. 2011. Full-length transcriptome assembly from RNA-Seq data
without a reference genome. *Nature Biotechnology* **29**: 644-652.

Griswold CK. 2021. The effects of migration load, selfing, inbreeding depression, and the genetics of
adaptation on autotetraploid versus diploid establishment in peripheral habitats. *Evolution* **75**: 39-55.

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D,
Li B, Lieber M, et al. 2013. De novo transcript sequence reconstruction from RNA-seq using the
Trinity platform for reference generation and analysis. *Nature Protocols* **8**: 1494-1512.

Harris K, Nielsen R. 2016. The genetic cost of Neanderthal introgression. *Genetics* **203**: 881-891.

Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005. Very high resolution interpolated
climate surfaces for global land areas. *International Journal of Climatology* **25**: 1965-1978.

Hu H, Al-Shehbaz IA, Sun Y, Hao G, Wang Q, Liu J. 2015. Species delimitation in *Orychophragmus*
(Brassicaceae) based on chloroplast and nuclear DNA barcodes. *Taxon* **64**: 714-726.

Hu H, Hu Q, Al-Shehbaz IA, Luo X, Zeng T, Guo X, Liu J. 2016. Species delimitation and
interspecific relationships of the genus *Orychophragmus* (Brassicaceae) inferred from whole
chloroplast genomes. *Frontiers in Plant Science* **7**: 1826.

Hu H, Zeng T, Wang Z, Al-Shehbaz IA, Liu J. 2018. Species delimitation in the *Orychophragmus*
violaceus species complex (Brassicaceae) based on morphological distinction and reproductive
isolation. *Botanical Journal of the Linnean Society* **188**: 257-268.

Huang F, Chen P, Tang X, Zhong T, Yang T, Nwafor CC, Yang C, Ge X, An H, Li Z, et al. 2023.
Genome assembly of the Brassicaceae diploid *Orychophragmus violaceus* reveals complex whole-
genome duplication and evolution of dihydroxy fatty acid metabolism. *Plant Communications* **4**:
100432.

Jia C, Hou Y, Lai Q, Zhang Y, Wang R, Liu J, Wang J. 2023. A reference genome and its epigenetic
landscape of potential *Orychophragmus violaceus*, an industrial crop species. *bioRxiv*: doi:
10.1101/2023.1109.1121.558835.

Jia C, Lai Q, Zhu Y, Feng J, Dan X, Zhang Y, Long Z, Wu J, Wang Z, Qumu X, et al. 2024.
Intergrative metabolomic and transcriptomic analyses reveal the potential regulatory mechanism of
unique dihydroxy fatty acid biosynthesis in the seeds of an industrial oilseed crop *Orychophragmus*
violaceus. *BMC Genomics* **25**: 29.

Johnston JS, Pepper AE, Hall AE, Chen ZJ, Hodnett G, Drabek J, Lopez R, Price HJ. 2005.
Evolution of genome size in Brassicaceae. *Annals of Botany* **95**: 229-235.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast
model selection for accurate phylogenetic estimates. *Nature Methods* **14**: 587-589.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements
in performance and usability. *Molecular Biology and Evolution* **30**: 772-780.

Keightley PD, Eyre-Walker A. 2007. Joint inference of the distribution of fitness effects of deleterious
mutations and population demography based on nucleotide polymorphism frequencies. *Genetics* **177**:
2251-2261.

Keightley PD, Jackson BC. 2018. Inferring the probability of the derived vs. the ancestral allelic state
at a polymorphic site. *Genetics* **209**: 897-906.

Khouider S, Borges F, LeBlanc C, Ungru A, Schnittger A, Martienssen R, Colot V, Bouyer D. 2021.
Male fertility in *Arabidopsis* requires active DNA demethylation of genes that control pollen tube
function. *Nature Communications* **12**: 410.

Kidokoro S, Konoura I, Soma F, Suzuki T, Miyakawa T, Tanokura M, Shinozaki K, Yamaguchi-
Shinozaki K. 2023. Clock-regulated coactivators selectively control gene expression in response to
different temperature stress conditions in *Arabidopsis*. *Proceedings of the National Academy of*
Sciences **120**: e2216183120.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* **37**: 907-915.

Kolář F, Čertner M, Suda J, Schönswetter P, Husband BC. 2017. Mixed-ploidy species: progress and

opportunities in polyploid research. *Trends in plant science* **22**: 1041-1055.

Korunes KL, Samuk K. 2021. pixy: Unbiased estimation of nucleotide diversity and divergence in the

presence of missing data. *Molecular Ecology Resources* **21**: 1359-1368.

Kovaka S, Zimin AV, Pertea GM, Razaghi R, Salzberg SL, Pertea M. 2019. Transcriptome assembly

from long-read RNA-seq alignments with StringTie2. *Genome Biology* **20**: 278.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: molecular evolutionary genetics

analysis across computing platforms. *Molecular Biology and Evolution* **35**: 1547-1549.

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: A resource for timelines, timetrees, and

divergence times. *Molecular Biology and Evolution* **34**: 1812-1819.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv:*

Genomics: doi: 10.6084/M6089.FIGSHARE.963153.V963151.

Li H, Durbin R. 2011. Inference of human population history from individual whole-genome sequences.

Nature **475**: 493-496.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,

Subgroup GPDP. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics* **25**:

2078-2079.

Li X, Teitgen AM, Shirani A, Ling J, Busta L, Cahoon RE, Zhang W, Li Z, Chapman KD, Berman

D, et al. 2018. Discontinuous fatty acid elongation yields hydroxylated seed oil with improved

function. *Nature Plants* **4**: 711-720.

Li Z-y, Ge X-h. 2007. Unique chromosome behavior and genetic control in *Brassica×Orychophragmus*

wide hybrids: a review. *Plant Cell Reports* **26**: 701-710.

Li Z, Heneen WK. 1999. Production and cytogenetics of intergeneric hybrids between the three

cultivated *Brassica* diploids and *Orychophragmus violaceus*. *Theoretical and Applied Genetics* **99**:

694-704.

Liu J, Liu B, Chen S, Gong B-Q, Chen L, Zhou Q, Xiong F, Wang M, Feng D, Li J-F, et al. 2018. A

tyrosine phosphorylation cycle regulates fungal activation of a plant receptor Ser/Thr kinase. *Cell Host*

and Microbe **23**: 241-253.

Liu S, Zhang L, Sang Y, Lai Q, Zhang X, Jia C, Long Z, Wu J, Ma T, Mao K, et al. 2022.

Demographic history and natural selection shape patterns of deleterious mutation load and barriers to

introgression across *Populus* genome. *Molecular Biology and Evolution* **39**: msac008.

Louder MIM, Justen H, Kimmitt AA, Lawley KS, Turner LM, Dickman JD, Delmore KE. 2024.

Gene regulation and speciation in a migratory divide between songbirds. *Nature Communications* **15**:

98.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. *Genome Biology* **15**: 550.

Lynch M, Conery J, Burger R. 1995. Mutation accumulation and the extinction of small populations.

The American Naturalist **146**: 489-518.

Lysak MA, Koch MA, Beaulieu JM, Meister A, Leitch IJ. 2008. The dynamic ups and downs of

genome size evolution in Brassicaceae. *Molecular Biology and Evolution* **26**: 85-98.

Malinsky M, Matschiner M, Svardal H. 2021. Dsuite - Fast D-statistics and related admixture evidence

from VCF files. *Molecular Ecology Resources* **21**: 584-595.

Marçais G, Kingsford C. 2011. A fast, lock-free approach for efficient parallel counting of occurrences
of k-mers. *Bioinformatics* **27**: 764-770.

Marhold K, Lihová J. 2006. Polyploidy, hybridization and reticulate evolution: lessons from the
Brassicaceae. *Plant Systematics and Evolution* **259**: 143-174.

Martin SH, Davey JW, Jiggins CD. 2015. Evaluating the use of ABBA–BABA statistics to locate
introgressed loci. *Molecular Biology and Evolution* **32**: 244-257.

Masterson J. 1994. Stomatal size in fossil plants: evidence for polyploidy in majority of angiosperms.
Science **264**: 421-424.

Mayrose I, Zhan SH, Rothfels CJ, Magnuson-Ford K, Barker MS, Rieseberg LH, Otto SP. 2011.
Recently formed polyploid plants diversify at lower rates. *Science* **333**: 1257-1257.

McDaniel SF. 2024. Local adaptation, recombination, and the fate of neopolyploids. *New Phytologist*
244: 32-38.

Merow C, Smith MJ, Silander Jr JA. 2013. A practical guide to MaxEnt for modeling species'
distributions: what it does, and why inputs and settings matter. *Ecography* **36**: 1058-1069.

Monnahan P, Kolář F, Baduel P, Sailer C, Koch J, Horvath R, Laenen B, Schmickl R, Paajanen P,
Šrámková G, et al. 2019. Pervasive population genomic consequences of genome duplication in
Arabidopsis arenosa. *Nature Ecology and Evolution* **3**: 457-468.

Morales-Briones DF, Kadereit G, Tefarikis DT, Moore MJ, Smith SA, Brockington SF, Timoneda
A, Yim WC, Cushman JC, Yang Y. 2020. Disentangling sources of gene tree discordance in
phylogenomic data sets: testing ancient hybridizations in Amaranthaceae s.l. *Systematic Biology* **70**:
219-235.

Morales-Cruz A, Aguirre-Liguori JA, Zhou Y, Minio A, Riaz S, Walker AM, Cantu D, Gaut BS.
2021. Introgression among North American wild grapes (*Vitis*) fuels biotic and abiotic adaptation.
Genome Biology **22**: 254.

Morgan EJ, Čertner M, Lučanová M, Kubíková K, Marhold K, Kolář F. 2020. Niche similarity in
diploid-autotetraploid contact zones of *Arabidopsis arenosa* across spatial scales. *American Journal*
of Botany **107**: 1375-1388.

Mortier F, Bafort Q, Milosavljevic S, Kauai F, Prost Boxoen L, Van de Peer Y, Bonte D. 2024.
Understanding polyploid establishment: temporary persistence or stable coexistence? *Oikos* **2024**:
e09929.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic
algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* **32**: 268-
274.

Odahara M, Kishita Y, Sekine Y. 2017. MSH1 maintains organelle genome stability and genetically
interacts with *RECA* and *RECG* in the moss *Physcomitrella patens*. *The Plant Journal* **91**: 455-465.

Osborn TC, Pires JC, Birchler JA, Auger DL, Chen ZJ, Lee H-S, Comai L, Madlung A, Doerge R,
Colot V. 2003. Understanding mechanisms of novel gene expression in polyploids. *Trends in genetics*
19: 141-147.

Otto SP, Whitton J. 2000. Polyploid incidence and evolution. *Annual review of genetics* **34**: 401-437.

Padilla-García N, Šrámková G, Záveská E, Šlenker M, Clo J, Zeisek V, Lučanová M, Rurane I,
Kolář F, Marhold K. 2023. The importance of considering the evolutionary history of polyploids
when assessing climatic niche evolution. *Journal of Biogeography* **50**: 86-100.

Parisod C, Holderegger R, Brochmann C. 2010. Evolutionary consequences of autopolyploidy. *New*
Phytologist **186**: 5-17.

Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, Genschoreck T, Webster T, Reich
D. 2012. Ancient admixture in human history. *Genetics* **192**: 1065-1093.

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. 2015. StringTie enables
improved reconstruction of a transcriptome from RNA-seq reads. *Nature Biotechnology* **33**: 290-295.

Phillips SJ, Dudík M. 2008. Modeling of species distributions with Maxent: new extensions and a
comprehensive evaluation. *Ecography* **31**: 161-175.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de
Bakker PIW, Daly MJ, et al. 2007. PLINK: A tool set for whole-genome association and population-
based linkage analyses. *The American Journal of Human Genetics* **81**: 559-575.

Ranallo-Benavidez TR, Jaron KS, Schatz MC. 2020. GenomeScope 2.0 and Smudgeplot for
reference-free profiling of polyploid genomes. *Nature Communications* **11**: 1432.

Rossi FR, Marina M, Pieckenstain FL. 2015. Role of arginine decarboxylase (ADC) in *Arabidopsis*
thaliana defence against the pathogenic bacterium *Pseudomonas viridiflava*. *Plant Biology* **17**: 831-
839.

Saha P, Ray T, Tang Y, Dutta I, Evangelous NR, Kieliszewski MJ, Chen Y, Cannon MC. 2013. Self-
rescue of an EXTENSIN mutant reveals alternative gene expression programs and candidate proteins
for new cell wall assembly in *Arabidopsis*. *The Plant Journal* **75**: 104-116.

Schiffels S, Durbin R. 2014. Inferring human population size and separation history from multiple
genome sequences. *Nature Genetics* **46**: 919-925.

Spoelhof JP, Soltis PS, Soltis DE. 2017. Pure polyploidy: closing the gaps in autopolyploid research.
Journal of Systematics and Evolution **55**: 340-352.

Srikant T, Gonzalo A, Bomblies K. 2024. Chromatin accessibility and gene expression vary between a
new and evolved autopolyploid of *Arabidopsis arenosa*. *Molecular Biology and Evolution* **41**:
msae213.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* **30**: 1312-1313.

Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence alignments
into the corresponding codon alignments. *Nucleic Acids Research* **34**: W609-W612.

Tan Z, Xu J, Zhao B, Zhang X. 1998. New taxa of *Orychophragmus* (Cruciferae) from China. *Acta*
Phytotaxonomica Sinica **36**: 544-548.

Van de Peer Y, Mizrachi E, Marchal K. 2017. The evolutionary significance of polyploidy. *Nature*
Reviews Genetics **18**: 411-424.

Vicente J, Cascón T, Vicedo B, García-Agustín P, Hamberg M, Castresana C. 2012. Role of 9-
lipoxygenase and α-dioxygenase oxylipin pathways as modulators of local and systemic defense.
Molecular Plant **5**: 914-928.

Völkner C, Holzner LJ, Day PM, Ashok AD, Vries Jd, Bölter B, Kunz H-H. 2021. Two plastid
POLLUX ion channel-like proteins are required for stress-triggered stromal Ca^{2+}release. *Plant*
Physiology **187**: 2110-2125.

Volyanskaya AR, Antropova EA, Zubairova US, Demenkov PS, Venzel AS, Orlov YL, Makarova
AA, Ivanisenko TV, Gorshkova TA, Aglyamova AR, et al. 2023. Reconstruction and analysis of the
gene regulatory network for cell wall function in *Arabidopsis thaliana* L. leaves in response to water
deficit. *Vavilovskii Zhurnal Genet Selektsii* **27**: 1031-1041.

Wang Z, Zhai L, Xiong S, Li X, Han Y, Wang G, Feng L, Fan Z, Lei Y, Yang B, et al. 2021. February
orchid cover crop improves sustainability of cotton production systems in the Yellow River basin.
Agronomy for Sustainable Development **41**: 67.

Weiß CL, Pais M, Cano LM, Kamoun S, Burbano HA. 2018. nQuire: a statistical framework for
ploidy estimation using next generation sequencing. *BMC Bioinformatics* **19**: 122.

Westermann J. 2021. Two is company, but four Is a party-Challenges of tetraploidization for cell wall
dynamics and efficient tip-growth in pollen. *Plants* **10**: 2382.

Wood TE, Takebayashi N, Barker MS, Mayrose I, Greenspoon PB, Rieseberg LH. 2009. The
frequency of polyploid speciation in vascular plants. *Proceedings of the National Academy of Sciences*
of the United States of America **106**: 13875-13879.

Yang Z. 2007. PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular Biology and*
Evolution **24**: 1586-1591.

Yin J, Zhang C, Mirarab S. 2019. ASTRAL-MP: scaling ASTRAL to very large datasets using
randomization and parallelization. *Bioinformatics* **35**: 3961-3969.

Yu H, Xu W, Chen S, Wu X, Rao W, Liu X, Xu X, Chen J, Nishimura MT, Zhang Y, et al. 2024.
Activation of a helper NLR by plant and bacterial TIR immune signaling. *Science* **386**: 1413-1420.

Yu R-M, Zhang N, Zhang B-W, Liang Y, Pang X-X, Cao L, Chen Y-D, Zhang W-P, Yang Y, Zhang
D-Y, et al. 2023. Genomic insights into biased allele loss and increased gene numbers after genome
duplication in autotetraploid *Cyclocarya paliurus*. *BMC Biology* **21**: 168.

Zeng Y, Zheng Z, Hessler G, Zou K, Leng J, Bautor J, Stuttmann J, Xue L, Parker JE, Cui H. 2023.
Arabidopsis PHYTOALEXIN DEFICIENT 4 promotes the maturation and nuclear accumulation of
immune-related cysteine protease RD19. *Journal of Experimental Botany* **75**: 1530-1546.

Zhang K, Yang Y, Zhang X, Zhang L, Fu Y, Guo Z, Chen S, Wu J, Schnable JC, Yi K, et al. 2023.
The genome of *Orychophragmus violaceus* provides genomic insights into the evolution of
Brassicaceae polyploidization and its distinct traits. *Plant Communications* **4**: 100431.

Zhong L, Liu H, Ru D, Hu H, Hu Q. 2019. Population genomic evidence for radiative divergence of
four *Orychophragmus* (Brassicaceae) species in eastern Asia. *Botanical Journal of the Linnean Society*
191: 18-29.

Zhou L, Liu Z, Wu J, Wang J, Yang Y, Li X. 2008. Karyotype variation and evolution in populations
of the Chinese endemic *Orychophragmus violaceus* complex (Brassicaceae). *Nordic Journal of*
Botany **26**: 375-383.

Zhou L, Yu Y, Song R, Hen X, Jiang Y, Li X, Yang Y. 2009. Phylogenetic relationships within the
Orychophragmus violaceus complex (Brassicaceae) endemic to China. *Acta Botanica Yunnanica* **31**:

127-137.

Zhu L, Fernández-Jiménez N, Szymanska-Lejman M, Pelé A, Underwood CJ, Serra H, Lambing

C, Dluzewska J, Bieluszewski T, Pradillo M, et al. 2021. Natural variation identifies SNI1, the

SMC5/6 component, as a modifier of meiotic crossover in *Arabidopsis*. *Proceedings of the National*

Academy of Sciences **118**: e2021970118.

Zhu Y, Wang X, He Y, Liu Y, Wang R, Liu Y, Wang S. 2024. Chromosome doubling increases *PECTIN*

METHYLESTERASE 2 expression, biomass, and osmotic stress tolerance in kiwifruit. *Plant*

Physiology **196**: 2841-2855.

1019

1020 **Figure legend**

1021

Fig. 1. Phylogenetic relationships and geographical distribution of
Orychophragmus. **(a)** Phylogenetic inference and divergence time estimation for six
Orychophragmus species alongside other cruciferous species. Blue numbers indicate
estimated divergence times (95% highest posterior density, in million years ago [MYA]),
while red numbers represent bootstrap support values. The black star denotes the
Orychophragmus-specific whole-genome duplication event, the black triangle marks
the Brassica-specific triplication event, and the black tetragon represents the ancestral
whole-genome duplication within the Brassicaceae family. **(b)** Geographical
distribution and morphological characteristics of the six species within the genus
Orychophragmus.

1032

Fig. 2. Demographic histories and divergent selection between *O. violaceus* and *O. taibaiensis*. (a) Species distribution modeling under the Last Glacial Maximum (LGM), Mid-Holocene (MH), and present climate conditions. **(b)** Demographic history inferred using the PSMC (outer) and MSMC (inner, see Fig. S5 for details) model. Gray vertical bars indicate the LGM, Riss glaciation, and Pre-Pastonian glaciation periods. Bold lines represent dynamic changes in effective population size (N_e), while faint lines show 100 bootstrap replicates, ensuring robustness. **(c)** The best-fit demographic model inferred using *fastsimcoal2*. Each block represents a current or ancestral population, with arrows indicating gene flow after divergence (per-generation migration rates). The timing of historical events is shown in thousands of years ago (KYA). **(d)** Ratio of nucleotide diversity at 0-fold sites relative to 4-fold sites. **(e)** Distribution of fitness effects (DFE)

in bins of N_es for new 0-fold nonsynonymous mutations for *O. violaceus* and *O.*

taibaiensis. Error bars indicate 95% confidence intervals based on 200 bootstrap

replicates. **(f)** Selective sweep analysis based on XP-CLR scores along chromosomes.

The top 1% of scores, above the red dashed horizontal line, are considered candidate

selective regions. Red circles highlight representative candidate genes located within

these regions, with black arrows indicating their names. Asterisks indicate statistical

significance from the Wilcoxon test (two-tailed) (NS. > 0.05, *$P < 0.05$, **$P < 0.01$,

***$P < 0.001$).

1053

Fig. 3 Comparison of the geographic distribution, divergence history, genetic load, and selection efficiency of diploid and tetraploid *O. taibaiensis*. (a) Geographic distribution of diploid and tetraploid *O. taibaiensis*, with stars representing the sequenced individuals mentioned in the text. Red triangles indicate the seed collection sites used for gene expression analysis across ploidy levels. (b) Karyotype of diploid *O. taibaiensis*. (c) Karyotype of tetraploid *O. taibaiensis*. (d) Relative proportions of various genotypes within the five whole-genome resequenced tetraploid *O. taibaiensis* individuals. (e) Phylogenetic relationships among the five resequenced diploid and tetraploid *O. taibaiensis*. (f) Population historical dynamics and divergence time

estimation between diploid and tetraploid *O. taibaiensi*s, based on *Fastsimcoal2*. (g)

Distribution of fitness effects (DFE) in bins of N_es for new 0-fold nonsynonymous

mutations for diploid and tetraploid *O. taibaiensis*. Error bars represent 95% confidence

intervals based on 200 bootstrap replicates. Asterisks indicate the level of significance

in the Wilcoxon test (two-tailed) (NS: not significant, *$P < 0.05$, **$P < 0.01$, ***$P <$

0.001).

Fig. 4 Differential expression analysis between diploid and tetraploid *O. taibaiensis*. (a) Hierarchical clustering of gene expression in diploid and autotetraploid *O. taibaiensis* for leaves (yellow, L) and roots (blue, R), highlighting key representative Gene Ontology (GO) terms enriched among differentially expressed genes (DEGs) within each cluster. Multiple representative DEGs from each cluster are shown. (b) Examples of expression level comparisons between diploid and tetraploid *O. taibaiensis* for selected representative DEGs identified in (a).

Fig. 5 Evolutionary and selection consequences of gene expression changes following autopolyploidization in *O. taibaiensis*. Comparison of genetic divergence of F_{ST} (a), d_{xy} (b) between differential expressed genes (DEGs, blue) and non-differentially expressed genes (NDEs, yellow) across samples with different ploidy levels. (c) Comparison of the ratio of 0-fold to 4-fold genetic diversity, used as a measure of selection efficiency, between DEGs and NDEs in diploid (light color) and tetraploid (dark color) populations in *O. taibaiensis*. (d) Distribution of fitness effects (DFE) for DEGs and NDEs in diploid (light color) and tetraploid (dark color) populations in *O. taibaiensis*. Errors bars represent 95% confidence interval based on 200 bootstrap replicates. Asterisks denote significance levels in the Wilcoxon test (two-tailed) (NS. > 0.05, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$).