Rapid Dopaminergic Signatures in Movement: Reach Vigor Reflects Reward Prediction Error
and Learned Expectation

Colin C. Korbisch[1], Alaa A. Ahmed[1,2]

1. Department of Mechanical Engineering, University of Colorado Boulder
2. Biomedical Engineering Program, University of Colorado Boulder

Running head: Reaching vigor tracks learned prediction error

Address for Correspondence:
Alaa Ahmed
Neuromechanics Laboratory
Department of Mechanical Engineering
University of Colorado Boulder
Boulder, CO 80309-0354
email: alaa@colorado.edu

KEYWORDS: reaching; reward; value; decision making; movement vigor

GRANTS: Work is supported by grants from the National Institutes of Health (1R01NS096083) and
the National Science Foundation (CAREER award 1352632) to AAA.

DISCLOSURES: The authors report no potential conflicts of interest.

AUTHOR CONTRIBUTIONS: CCK and AAA conceived and designed experiment. CCK performed
experiments and analyzed data. CCK and AAA interpreted results of experiments. CCK prepared
figures and drafted manuscript. CCK and AAA edited and revised manuscript. CCK and AAA
approved final manuscript.

ORCiD Numbers: CK:0000-0002-6566-2860; AA:0000-0002-1596-342X

Institutional ID: http://ror.org/02ttsq026

ABSTRACT

Movement vigor across multiple modalities increases with reward, suggesting that the neural circuits that represent value influence the control of movement. Dopaminergic neuron (DAN) activity has been suggested as the potential mediator of this response. If DAN activity is the bridge between value and vigor, then vigor should track canonical mediators of DAN activity, namely learning signals in the form of reward expectation and reward prediction error. Here we ask if a similar time-locked response is present in vigor of reaching movements. We explore this link by leveraging the known phasic dopaminergic response to stochastic rewards, where activity is modulated by both reward expectation at cue and the reward prediction error at feedback. We used probabilistic rewards to create a reaching task rich in reward expectation, reward prediction error, and learning. In one experiment, target reward probabilities were explicitly stated, and in the other, were left unknown and to be learned by the participants. We included two stochastic rewards (probabilities 33% and 66%) and two deterministic ones (probabilities 100% and 0%). In both experiments, outgoing peak velocity increased with increasing reward expectation. Furthermore, we observed a short-latency response in the vigor of the ongoing movement, that tracked reward prediction error: either invigorating or enervating velocity consistent with the sign and magnitude of the error. Reaching kinematics also revealed the value-update process in a trial-to-trial fashion, similar to the effect of prediction error signals typical in dopamine-mediated striatal phasic activity. Lastly, reach vigor increased with reward history over trials, mirroring the motivational effects often linked to fluctuating dopamine levels. Taken together, our results highlight the link between known short-latency dopaminergic learning signals and the invigoration of movement, not only at the time of cue presentation and movement initiation, but during an ongoing movement immediately after feedback is provided.

NEW & NOTEWORTHY

Previous research has demonstrated the invigorating effects of reward on movement. Growing evidence suggests this is causally explained by midbrain dopamine transients. Here, we demonstrate that reach vigor tracks canonical variables of learning and motivation across time scales ranging from milliseconds to minutes. Velocity was modulated by reward expectation, reward prediction error and reward rate, key variables that have also been associated with striatal dopaminergic fluctuations. These results point to a potential neural mechanism by which dopamine can influence both decision making and movement control and support the proposition that reward-based invigoration of movement is in part influenced by dopaminergic circuits.

INTRODUCTION

Imagine sitting down in front of a series of slot machines and being instructed to pull their levers, one after another. You find that these machines, frequently referred to as one-armed bandits for their ability to extricate money from patrons, have different rates of payouts, with some better than others. After learning these differences, you will likely prefer to pull the levers that offer a greater payout. But would you also choose to pull those levers faster or with more force?

Previous inquiry has shown that individuals will move faster towards goals or targets associated with greater value[1–10]. People are willing to produce greater muscle forces, or expend greater effort, to reach a cued target in less time. A potential explanation for how this may be represented in the brain lies in the neurotransmitter dopamine (DA), which is implicated in both the learning and representation of value as well as the control of movement[11]. Basal ganglia activity, already known to be influenced by dopaminergic inputs, has been found to invigorate not just movement vigor[12–15], but the decision-making process as well[16,17]. If vigor is indeed a reflection of value and DA the mediator, then vigor may also reflect the machinery that underlies the learning process, due to the phasic dopamine release coincident with learning and reward prediction error[18]. Seminal work has shown that dopaminergic neuron (DAN) responses in a learned environment scale with reward prediction error at the time of both stimulus presentation and feedback presentation[19]. DAN activity is greater in response to cues associated with greater expectation of reward (greater positive reward prediction error), and lower upon feedback of that reward (lower positive reward prediction error). If a similar time-locked response is present in movement vigor at cue and feedback presentation, this would suggest that DA-related activity contributes to reward-driven increases in vigor.

While short-term phasic DAN response is implicated in prediction error and learning, tonic dopamine levels have been found to be sensitive to the history of reward reception[1,20–22]. Particularly, midbrain DA levels come to match average reward rate and are implicated in motivation and invigoration. Hamid et. al. found a significant relationship between history of reward and relative DA in the nucleus accumbens compared to other potential analytes. Greater tonic levels of DA can also be interpreted as greater motivation or *drive*, resulting in behavioral invigoration[23,24]. Recent reward history has been extensively recorded influencing movement vigor[20,25–28]. Returning to our hypothetical gambler, receiving more rewards over the past, regardless of which levers were pulled, would result in greater vigor in subsequent lever pulls.

A growing body of work has found correlates between phasic dopaminergic activity and movement vigor[29–37]. Cue-evoked phasic dopamine response within rat nucleus accumbens core (NAc core) was significantly correlated with trial movement vigor[30]. Similarly, for rat substantia nigra pars compacta (SNc) neurons sensitive to free movement initiation, relative activity levels were correlated with ensuing vigor[34]. Optogenetic activation of these neurons when not moving increased the probability of movement initiation, but did not produce any measurable changes in ongoing movement vigor. However, a separate study found that ongoing movement vigor was reduced with inhibition of striatal activity[33]. These and other works provide evidence that not only do longer timescale tonic dopamine levels potentially modulate animal movement vigor, so too may short latency, phasic responses.

In this study, we sought to investigate whether human kinematic response to probabilistic rewards would mirror the characteristic reward prediction error response observed in mesencephalic

dopaminergic neurons (DAN) on a sub-second timescale. We also asked if kinematic response would reflect tonic DA response correlated with reward history. Given the short-latency and tuned response of DANs to reward value, we hypothesized vigor response in human arm reaching to be modulated by both the expectation of reward (probabilistic expectation) as well as the prediction error (difference between binary reward outcome and expectation). In effect, we expect outcomes congruent with two of the axioms for RPE-based models: 1) vigor responses to various reward expectations should demonstrate consistent lottery ordering, and 2) should show evidence of the "no surprise equivalence," i.e., fully deterministic reward outcomes show no relative difference . We also hypothesized that greater reward amounts received over the recent past would lead to greater relative invigoration, as a result of increased striatal DA levels. To test these hypotheses, we performed two experiments, each of which involved human subjects performing out-and-back reaching movements to probabilistically rewarding targets. Our primary findings reveal that movement vigor is modulated online in alignment with reward prediction error on a sub-second time-scale, while also reflecting the learned value update on a trial-to-trial basis as well as average reward rate. These results highlight how motor planning, execution, and feedback control are all likely influenced by DA response, reflecting DA's tripartite role in learning, motivation, and movement control.

RESULTS

In a series of two experiments, we asked whether movement vigor tracked canonical determinants of dopamine-related learning and performance: learned value, reward prediction error, and reward history. In both experiments, human subjects directed arm reaching movements to targets associated with a probability of receiving a reward (Figure 1a-c; see Methods). Each of four target locations was associated with a unique probability of receiving the reward (0%, 33.$\overline{3}$%, 66.$\overline{6}$%, 100%) that was changed each block (Figure 1d) of 180 trials. Feedback of reward reception or reward denial was provided upon target acquisition. Assuming perfect representation of task instruction, participants could experience five reward prediction errors (RPE): -0.33 and +0.66 for the 66% target, -0.66 and 0.33 for the 33% target, and 0 RPE for the 0% and 100% targets (Figure 1e). This paradigm allowed us to probe the effect of reward expectation on the vigor of the outgoing movement. Additionally, we could investigate the effect of RPE at time of feedback on the vigor of the return movement. Lastly, we could test the effect of reward history on reach vigor across trials, independent of the expected value of the current target. We predicted that with increasing expectation, outgoing vigor would increase, reflected in an increase in peak velocity (Figure 1g), that vigor would be modulated on the return portion of the reach, scaled by the reward prediction error, and that vigor across trials would increase with reward history.

Figure 1. Experimental protocol. A) Participants made out-and-back reaching movements towards prompted, virtual targets. B) Targets were located at four different positions, each 10 cm from a central home location. C) Trial sequence. After the target was cued, participants were to make an out-and-back reach towards said target. Reward feedback was dependent on stated target reward frequency (in the first experiment), with reward feedback consisting of a yellow flash and high-pitched tone. Omission of feedback coincided with the target turning gray. D) At the beginning of each 180-trial block, in the first experiment, participants were told the reward frequencies of each target, either 0%, 33%, 66%, or 100%. Shown here is one possible experiment progression order. Relative order of reward-target associations was randomized across participants. Each block's rewards were pseudorandomized such that, in the single-target trials, the target reward frequencies were exact to the underlying frequency. Each block consisted of 5 sets of 36 trials, wherein each target was presented 9 times, and amount of reward per target was precisely equal to the stated expectation (i.e., rewards were given 0, 3, 6, or 9 times within each set of 36 trials depending on the target). Target presentation order within each set of 36 trials was fully randomized. E) Reward prediction error (RPE) was defined as the difference between the stated (and actual) reward frequency per target and the binary outcome of the trial. G) *Left* Kinematic measures of interest included the time between target presentation (cue) and movement onset (reaction time), peak velocity of the outgoing portion of the movement, time between movement onset and target hit, peak velocity of the return portion, and difference between the absolute value of the two peak velocities. Example positional trace of trial reach trajectory given on *Right*, with spatiotemporal points of interest highlighted.

Experiment 1

In the first experiment, we sought to focus solely on the vigor response to a known expectation of reward (E[R]). We did so by informing the participants of the reward probability of the targets at the beginning of each block.

Figure 2. Vigor tracks reward expectation. A) *Upper* Radial velocity trace for first experiment, aligned to time of outgoing peak velocity. Highlighted region shows differences in peak excursion velocity for differences in reward expectations. *Lower* To calculate the difference traces, each participant's average signed radial velocity, at each sampled time point for the 0% expected reward condition, was subtracted from all other trials. Traces shown are the grand average of these difference traces, +/- standard error. B) *Left* Outgoing peak velocity increased with increasing expected reward. Data were normalized by subtracting the per-participant average for 0% expected reward condition. *Right* Similarly, the time to target hit, defined as the time between movement onset and reaching out to a 10cm radial distance, decreased as reward expectation increased. C) Return peak velocities were residualized by first fitting a reduced model including target direction and outgoing peak velocity. These residuals were significantly correlated with trial reward prediction error, increasing with greater RPE.

Peak velocity to target tracks reward expectation

As the cued target's E[R] increased, peak velocity of the outgoing movement increased as well (GLMM, $\beta_{E[R]}$=0.0159±0.00629, p=0.0152, Figure 2a,b). Likewise, time to target, defined as the time between movement onset and reaching a 10 cm radial distance, decreased with increasing reward expectation ($\beta_{E[R]}$=-0.0125±0.00527, p=0.0226). Reaction times decreased with greater E[R] (Supplemental Figure 1a-b).

Return velocity is influenced by reward prediction error

When the cursor hits the target, participants can either receive reward feedback or not. Does this feedback, or lack thereof, influence subsequent movement kinematics? Does the degree of surprise also modulate movement kinematics? In other words, does the reward prediction error influence the speed of the ensuing movement?

To answer these questions, we analyzed participants' return movements and whether receiving (or not receiving) reward feedback on a given trial produced significant influences. Specifically, we questioned whether the sign and magnitude of the reward prediction error, *RPE,* influenced the velocity:

$$RPE = R - E[R]$$

with R a binary coding for presence or absence of reward feedback (0,1), and $E[R]$ the instructed expectation of reward feedback (see Methods).

Participants exhibited diminished peak velocity on return movements compared to the outgoing movements (paired mean difference = 0.0601 m/s [0.0445,0.0757 95% CI], t_{41}=7.7789, p=1.34e-9). After controlling for the average relationship between outward and return peak velocities (β_{OutPV}=0.580±0.0192, p<2e-16), we observed an effect of reward feedback and reward expectation, i.e., RPE (β_{RPE}=0.00865±0.00260, p=0.00186). This indicated that the return portion of the reach was not merely driven by feedforward mechanisms, but also feedback. No significant main effect of reward was found (β_{Reward}=0.000876±0.00159, p=0.586), nor interaction with RPE ($\beta_{RPE×Reward}$=-0.00545±0.00392, p=0.171), signifying a slope of response across prediction error values without discontinuities. Additional control analyses confirmed return movements were modulated by reward feedback and not other potential confounds (Supplementary Figures 2-4).

Figure 3. Vigor response reflects reward prediction error. A) For a within-trial measure of relative invigoration, instantaneous velocity was normalized by dividing by the trial outgoing peak. Difference traces were then calculated in a manner similar to figure 2A. From SPM analysis, a significant effect of RPE on percent peak velocity was found beginning at 212 ms after reward feedback was received. B) The within-trial difference in absolute peak velocities, *Left* return divided by outgoing *Right* return minus outgoing, significantly varied with reward prediction error. As RPE increased, the return peak more closely matched the outgoing peak.

To better account for variance in the outgoing portion of the movement, we normalized instantaneous velocity and focused on within-trial effects. First, instantaneous radial velocity was divided by the within-trial outgoing peak velocity. Next, the difference traces of these %-outgoing velocity values were calculated in a similar manner as in figure 2A, though instead of taking the difference to the 0% reward trials, the differences were taken relative to the per-participant averaged normalized velocity trace for all 0% and 100% reward trials, i.e. RPE=0 trials. After calculating the grand average of these difference traces, we see a clear striation in relative normalized return velocity corresponding with reward prediction error (Figure 3a).

We next performed a hierarchical random effects analysis (Supplementary Figure 5; see Methods) to determine the population-average effect of reward prediction error on instantaneous, normalized

velocity. At 212 ms after target feedback, we found a significant negative effect of RPE, indicative of
greater invigoration in the return movement with greater RPE (Figure 3a).

Aside from the effect on instantaneous velocity, the within-trial difference in velocities, i.e., the
difference between excursion and return peak velocity, also exhibited a significant effect of RPE
(Figure 3b); with the slope of response consistent across positive and negative RPE conditions ($\beta_{RPE \times}$
$_{Reward}$ =-0.00176±0.00502, p=0.727). For both percent relative to outgoing peak (3b *Left*) and velocity
difference (3b *Right*) the slope of response was significant across RPE conditions
(β_{RPE}=0.02788±0.00788, p=1.03e-3; β_{RPE}=0.0117±0.00325, p=9.00e-4, respectively). When RPE was
most positive (+0.66) the difference between excursion and return peak velocity was at its smallest
compared to RPE at its most negative (-0.66). There was no separate effect of reward feedback in
either metric (β_{Reward}=-0.00269±0.00463, p=0.5643; β_{Reward}=-0.00355±0.00208, p=0.0978).

Figure 4. Experiment 2 choice behavior. A) In the second experiment, reward probabilities were left unstated to the participants. Instead, they were incentivized to learn the underlying rewards during the single-target period, as additional monetary compensation was dependent on performance during later choice trials (see Methods). B) Per-participant average accuracy rate over four blocks. Participant choices were highly variable across the population. Mean accuracy was 71%. C) Participant choices were found to be sensitive to relative expectation, choosing the more rewarding option as difference increased. D) Response time decreased with increasing difference of reward expectations of the presented targets. E) Target direction was effected choice selection. When targets 1 or 3 were the more rewarding option, i.e. associated with lesser biomechanical effort, they were selected at higher rates compared to when targets 2 or 4 were the more rewarding option, consistent with a potential influence of biomechanical effort on choice selection. F) When grouping choices by whether the greater reward has greater relative effective mass, significant differences in choice accuracies were found. Individual participant responses are shown in color. G) Baseline selection rates were shifted depending on relative effort. No interaction between effort and reward difference was found.

Overall, we found a significant effect of reward feedback on the return movement that was
dependent on the reward expectation, i.e., the reward prediction error. With greater RPE, relative
return velocity was greater compared to when the prediction error was negative. This difference
emerged 212 ms after feedback was received and could not be accounted for by variation in
maximum excursion, nor preemptive differences in the outgoing portion of the movement (potential
correlations between outgoing peak velocity and future reward reception). Participants employed
neither set return velocity, nor set return velocity difference strategies. In the first case, outgoing peak
velocity would be unrelated to the return peak, and in the second case, return velocity difference
would be invariant to the experienced prediction error.

Experiment 2

We next asked whether vigor was modulated by learned value, rather than instructed value. To probe
this we conducted a second experiment where participants were left uninformed of the targets'
reward frequencies. Instead, participants learned through experience over the course of a block,
affording us a measure of trial-to-trial response of changes in reward estimation. Within each block,
after a sequence of 144 single-target trials similar to the first experiment, there were 36 two-
alternative forced choice trials (2-AFC) to assess degree of learning (Figure 4a). Individuals were
incentivized to choose greater expected rewards on choice trials as additional monetary
compensation was dependent on performance (see Methods). Critically, no feedback was given on
target hit during these choice trials to limit learning to the single target trials only.

Choices reflected both expectation of reward and effort

Choice accuracy, defined as the selection of the greater reward frequency of the two options, varied
across the participants, ranging from 42.7% to 91.7% (Figure 4b). The population-average accuracy
rate was 71%. Underlying reward difference, hidden to participants, was found to significantly
predict rate of choosing one option over another ($\beta_{\Delta E[R]}$=1.233±0.476, p=0.0096; Figure 4c). Response
time of the decision was found to significantly decrease with increasing difference of reward
expectation (Figure 4d; $\beta_{|\Delta E[R]|}$=-0.202±0.053, p=1.47e-4), providing further evidence that underlying
expectations were learned. Interestingly, response time and outgoing peak velocity response differed
during choice trials in their sensitivity to reward difference and choice accuracy (Supplementary
Figure 6). Whereas outgoing peak velocity only varied with the selected option's reward expectation,
response time varied with both relative and selected expectation.

Target direction was also found to influence decision making. We first categorized all choices by
whether a given target direction was associated with the greater of the two potential rewards
presented on a given choice trial. We then calculated the choice accuracy rate for each of these
subsets. If direction had no influence, we should expect no variation in selection rates between these
four categories, however this was not the case. Instead, significant differences were found (Figure
4e). Selection rate biases matched the arm's inertial axes[38,39] (Figure 4e *Right*). Using this measure of
effort, we categorized choice trials by whether the more rewarding option was associated with more
or less effort compared to the alternative choice (Figure 4f). We found significant differences in
accuracy rates, with idiosyncratic responses to effort readily apparent. Augmenting our logistic
regression analysis from before by including a relative effort term, we found a significant effect of
said relative effort, with average rates of choosing the more rewarding option increasing when this
option was associated with less effort (β_e=0.472±0.188, p=0.012; Figure 4g). Thus, participants'

choices revealed they had largely learned the target reward contingencies and demonstrated that
these value-based choices were influenced by the effort of the arm reach.
231

Figure 5. Experiment 2: Vigor tracked reward expectation and reward prediction error. A) *Left* Outgoing peak velocity increased with increasing reward expectation. Data were normalized by taking the difference to per-participant average for 0% reward. Points and error bars represent grand means, ± standard error. The slope of peak velocity response to reward expectation changed over the course of a block. In the initial 36 trials (start of block), response was essentially flat, but by the final 36 trials (end of block), outgoing velocity significantly increased with respect to reward frequency. *Right* Time to target, taken as the duration between movement onset and reaching out to a 10 cm radius (target hit) changed with respect to underlying expectation over the course of the single-target period. B) End of block response to reward (i.e. per-participant slope of response to E[R]) was significantly correlated with block choice accuracy (rate of choosing the more rewarding option, Figure 4B). C) Normalized velocity difference traces for experiment 2. As in figure 3C, velocity data were first normalized by dividing by the within-trial peak outgoing velocity. Difference traces were calculated by subtracting from each trial the average instantaneous normalized velocity for trials where the underlying reward expectation was 0% or 100% (RPE = 0). Data presented is the grand average of these difference traces. Differences in relative velocity emerged after reward feedback that scaled with prediction error. D) As RPE increased, the relative reduction in peak velocities between the return and outgoing portions of the reaching movement decreased (i.e. the difference was less negative).

Velocity in single-target trials tracked reward expectation and reflected learning

Given that participants' choices revealed a learned reward estimation, we asked whether vigor on
single target trials would reveal this estimation as learning progressed. Over the course of the single-
target period, we found that average slope of response of outgoing peak velocity relative to reward
expectation increased (GLMM; $\beta_{Trial \times E[R]}$ = 0.0392±0.01447, p=0.00682), demonstrating a dynamic
response to the probabilistic reward (Figure 5a). Likewise, the effect of expectation on the time to
target significantly varied over this period ($\beta_{Trial \times E[R]}$ = -0.0356±0.0142, p=0.0125). At the beginning of
a block, there was no significant effect of reward expectation ($\beta_{E[R]}$= -0.0108±0.0073, p=0.142). The
change in response of instantaneous velocity over the course of the single-target period is further
evident in the velocity difference traces (Supplementary Figure 7). However, no such interaction
effect between trial and hidden reward expectation was found when modeling reaction times
(Supplementary Figure 1).

The strength of peak velocity-reward expectation response (i.e. the slope of velocity relative to
reward) could predict within-block choice accuracy (the rate of choosing the more rewarding of the
two options). Across the participant pool, we found a significant correlation between these measures
(Kendall's rank correlation, τ = 0.156, p=0.0384; Figure 5b). The stronger the slope of response in an
individual at the end of the single target period, the greater the rate in selecting the more rewarding
of the two options presented during choice trials.

Even when participants were not told of the expected reward frequencies at the outset of a block,
both outgoing peak velocity and time to target responded to increasing expectation. Learning of the
different reward frequencies is shown clearly by the differential response of velocity at the beginning
and end of the single-target period. Initially, kinematic response did not differ between the four
targets, but by the end, average peak velocity increased with increasing expectation of reward
feedback. Within-block performance could be predicted based on participants' slope of response to
reward at the end of the single-target period, suggesting that the change in kinematics over the
course of a block was related to the learning of relative reward frequencies.

Change in return velocity tracks reward prediction error

Turning to within-trial measures of relative vigor, we performed the same hierarchical random effect
SPM analysis on normalized velocity as in the first experiment and found a population-level effect of
RPE (Figure 5c). Additional supporting detail for this analysis is provided in supplementary
documentation (Supplementary Figure 8). At 214 ms after reward feedback, relative velocity varied
significantly with reward prediction error. Likewise, the within-trial velocity difference (return minus
outgoing) also significantly varied with RPE (β_{RPE}=8.22e-3±2.61e-3, p=0.00166; Figure 5d). With
increasing positive prediction error, the return peak velocity more closely matched the outgoing peak
velocity. Reward feedback had no significant interaction effect on this difference ($\beta_{RPE \times Reward}$=-2.156e-
3±3.682e-3, p=0.558), However, there was a significant main effect of reward (β_{Reward} =-4.414e-
3±1.825e-3, p=0.0217). As such, for both 0% and 100% rewarding trials, the differences in outgoing
and return peak velocities were statistically distinct. The slope of response to reward prediction error
was not found to change over the course of a block ($\beta_{RPE \times Trial}$ =-4.746e-3±4.770e-3, p=0.320). As in the
previous experiment, we performed additional control analysis to isolate the effect of reward
feedback on the return movement (Supplementary Figure 9).

To reiterate, even when expected reward must be experienced and learned, individuals change their
behavior on the return movement in a manner consistent with RPE sign and magnitude.

Biomechanical effort slowed outgoing peak velocity

Considering the effect of target direction (i.e. relative effort), on choice bias, we were also concerned
whether there would be differences in average peak velocities towards the different directions.
Investigating this potential effect (Figure 6a *Left*), we found that each direction was significantly
different in average peak outgoing velocity from the other (Tukey-HSD comparison testing;
Supplementary Table 2). In addition to the directional effect, we found that average outgoing peak
velocity increased over the course of the experiment from block-to-block (β_{Block}=0.158±0.051,
p=0.0054).

We also evaluated whether this directional influence on velocity was conserved in the previous
experiment (Figure 6a *Right*). Once again, differences in average outgoing velocity across the
directions were consistent with the effective mass of the arm. Peak velocity increased over the
course of the experiment (β_{Block}=0.2352±0.0058, p=1.25e-9) for all targets, and post-hoc multiple
comparison testing (Tukey HSD) revealed significant differences across all four targets
(Supplementary Table 1).

In short, average velocity to the different target directions in both experiments reflected the relative
inertia, or effective mass, of the arm when making this reaching movement. Outgoing peak velocity
was greatest towards the 45° target, and slowest towards the 135° target.

Subjective vigor response in single-target trials predicted choice

Summarizing our results thus far for the second experiment, we have shown how participant choices
reflected subjective value, and kinematic response during single-target trials reflected the learning
of relative reward frequencies. To further substantiate the relationship between vigor and preference,
we examined the degree to which reach velocity during the single-target trial period could predict
choices across our participant pool.

To start, we analyzed whether the subjective effort bias evidenced in participant choices would also
be present in the kinematic response. Comparing the per-participant relative velocities and the target
selection rates, we found a significant correlation (Kendall's rank correlation, τ = 0.194, p=0.00846;
Figure 6b *Left*). With MANOVA analysis, after averaging across participants, we found significant
differences between the relative velocities and selection rates across the four target directions
(Figure 6b *Right;* $WTS_{(6)}$=43.165, p<0.001). Next, we questioned whether kinematic behavior during
the end of the single target period (the final 36 single-target trials) could predict later choice
selections directly. To do so, we analyzed the correlation between the relative peak velocity per-
target, per-participant and that target's within-block selection rate (quantified as a fraction of the
total 18 potential selections within the choice period). A significant correlation was found (Kendall's
rank correlation, τ = 0.104, p=0.00496; Figure 6c *Left*). The faster someone reached towards a target
at the end of the single-target period, the more frequently they chose that option during choice trials.

As an additional piece of evidence for the relationship between outgoing peak velocity and relative
selection rate, we directly compared the difference in mean velocity between two targets at the end
of the single target period and the relative rate of selection within that block. Via logistic regression

analysis, we found a significant relationship between the velocity difference and rate of selection ($\beta_{\Delta PV}$ = 9.56±2.55, p=1.75e-4; Figure 6c *Right*). Roughly, a 0.2 m/s difference in peak reach velocity translates to a 3:1 preferred rate of selection.

Figure 6. (Top) Relation between velocity, subjective effort, and learned value. A) *Left* In experiment 2, arm reaches towards target 1 were found to have a greater peak velocity on average, with targets 2 and 4 associated with comparatively lesser averages. *Right* In the first experiment, the same ordering of relative peak velocities was observed. B) *Upper* Participant-relative velocity, in single-target trials, were correlated with relative rates of selection of the different target directions. *Lower* Averaged over all participants per target, there was evidence of a separation in relative average velocities and selection rates. C) *Left* Relative velocity to a given target at the end of the single-target period could predict preference. *Right* Difference in average velocity between two targets, within the final 36 single-target trials, was continuous with relative rates of selection for those two targets. The greater the difference in velocity, the more frequently one option was chosen over the other in the proceeding six choice trials where the two targets were presented together. (Bottom) Performance of Learned Value Model D) Individuals were sensitive to learned value differences during decisions, selecting targets with greater relative values. E) Choice prediction accuracy improvement was participant specific. Average improvement was 4.42%, and some participants saw choice prediction improvements up to 20% (roughly 29 choices during the experiment). F) Choice trial response time significantly decreased with difference of the presented targets' learned values. G) Normalizing value relative to per-participant minimum and maximum estimate (min-max norm.), we found that outgoing peak velocity during the single-target trials significantly increased with increasing learned value. H) On trials where the same target was prompted in repetition, the previous trial's reward prediction error was significantly correlated with the trial-to-trial change in outgoing peak velocity. I) Value prediction error, taken as the difference between feedback and expected value, was also found to predict the relative difference in peak velocity between outgoing and return.

Taken together, we found that participant decisions varied widely across the population but could be
predicted by each participant's relative outgoing velocities on single target trials: the greater the
relative peak velocity, the more frequently that option was chosen.

Value Estimation

Reward and effort (in the form of target direction) had significant influences on outgoing velocity. Both
reward and effort also affected choice preference and relative selection rates. We posit that the
target characteristics of reward expectation and direction could be combined into a singular
subjective value or decision variable that would describe both participant-specific reach velocity and
choice preference.

Learned value better explained single-target trial reach vigor

To examine whether this was the case and estimate this subjective decision variable, we fit a
Bayesian hierarchical delta-rule learning model, similar to a Rescorla-Wagner model[40], to choices at
the end of each block to determine the per-trial subjective value for each participant. After each trial,
the updated value estimate of a target V'^T is calculated:

331
$$V'^T_i = V^T_i + \eta_s\left(R_i + e^T_s - V^T_i\right)$$

Here, R_i is the reward received on the current trial, η_s is learning rate, and e^T_s, the relative effort of the
target, dependent on direction (major or minor inertial axis). Participant-specific parameters, η_s and
e^T_s, were selected and sampled from an estimated population distribution in a hierarchical fashion.

A separate logistic regression confirmed that value differences predicted choice behavior (Figure 6d).
As relative value increased, individuals were more likely to choose that option. Additionally, the
model-derived learned value was found to better predict choices as compared to reward
expectation, with an aggregate accuracy rate of 75.4% compared to 70.98%. This was corroborated
with a significant difference in ROC curves between the two metrics (Bootstrap test for difference in
AUC: D=8.054, n=2000, p=8.014e-16). However, prediction improvements were highly idiosyncratic
in the participant pool (Figure 6e). Lastly, choice trial response time was significantly influenced by
value, decreasing with increasing value difference ($\beta_{|\Delta V|}$=-0.1373±0.0464, p=0.0094; Figure 6f).
Outgoing reach during choice trials was also influenced by the value estimate (Supplementary Figure
10).

We next used this learned value estimate to predict reach peak velocity on single target trials. There
was found to be a significant relationship between value and outgoing peak velocity (GLMM;
β_{Value}=0.0223±0.00895, p=0.0126; Figure 6g). The regression model including value improved in
performance relative to one using E[R] (AICc: -13479.4 versus -13369.3 respectively). Likewise, time
to target decreased with learned value as well, and better matched the data compared to reward
expectation (β_{Value}=-0.0265±0.007148, p=0.0013; Figure S6B; AICc: -12225.6 versus -12157.7).

Interestingly, even though subjective directional efforts were incorporated into the target value, there
were still significant effects of direction. Post-hoc Tukey HSD testing revealed significant differences
between the target-specific regression factors (Supplementary Table 3). This implies that the
difference in vigor between directions does not fully capture the resultant difference in value as
shown by later choices.

We predicted, as an additional piece of evidence relating learned value and vigor, that an individual's
change in value (the value prediction error) would be correlated with the trial-to-trial change in peak
velocity after a target is repeated (Figure 6h). Calculating this correlation, we found that, as
hypothesized, a greater value update leads to greater velocity changes, and the relationship was
significant (Pearson's correlation, $t_{(2801)}$=3.083, r=0.058, p=2.07e-3). Testing to ensure this effect was
target-specific, we calculated this correlation on trials where relative effort, but not direction, was
repeated, and found no significance ($t_{(2732)}$=-0.624, r=-0.012, p=0.533).

The relative difference in peak velocities between the outgoing and return portions of the movement
decreased with increasing value prediction error (VPE) (β_{VPE}=0.008414±0.00370, p=0.0297; Figure 6i);
the comparative velocity of the return portion of the reach was greater with more positive error.
Performance in modeling the within-trial velocity difference also improved when using VPE
compared to RPE (AICc: -37118 versus -37098.2). Unlike the RPE regression model that showed a
significant effect of reward reception on average difference in return velocity (see above, Figure 5d),
the value prediction error model did not (β_{Reward}=-0.00441±0.00296, p=0.153). Additionally, no
significant interaction was found ($\beta_{VPE \times Reward}$ = -0.00510±0.00298, p=0.0873). In effect, the slope of
response of velocity difference to value prediction error remained constant with or without reward
feedback. Taken together, relative return velocities varied continuously with value prediction error
with no additional significant effect of reward reception.

Recent history of reward led to faster movements

Having shown how target-specific learned value influenced the reach kinematics on the outgoing
and return portions of the movement, we sought to investigate the potential effects of target-agnostic
reward reception, independent of value estimation. We investigated the potential effect in both
experiments, testing the influence of previous trial reward reception and an integrated reward history,
making use of the "leaky integrator" model[41] (see Methods). Based on previous findings, we
hypothesized that reward history would invigorate the outgoing movement, increasing peak velocity,
and decrease response times.

In our first experiment, we found an effect of previous reward reception on outgoing peak velocity.
Peak velocity was greater on trials following reward compared to trials following no reward
(β_{PriorR}=0.0152±0.003, p=5.93e-6; Figure 7a). For our reward history model, after fitting to reaction
times, the smoothing factor, α, was found to equal 0.647. Incorporating reward history into our
previous regression model for outgoing peak velocity, we found that velocity significantly increased
with increasing $\bar{R}$ ($\beta_{History}$=0.0209±0.00319, p=7.15e-8; Figure 7b). Model performance also relatively
improved (AICc: -85753.7 versus -85728.6). This model-derived reward history value was significant
even when accounting for current trial reward expectation (Figure 7c, Supplementary Figure 11).

In our second experiment, unlike the first, there was no significant response caused by reward
reception on the previous trial (β_{PriorR}=-0.0033±0.006, p=0.592; Figure 7d). With the same reaction
time dependent likelihood maximization, we found a smoothing factor, α, equal to 0.0521. Outgoing
peak velocity again significantly increased with reward history (β_{Hist}=0.0884±0.0151, p=4.66e-9;
Figure 7e, Supplementary Figure 11e). Model performance when predicting velocity was also
improved (AICc: E[R]+History -36774.7, E[R] -36742.4).

Figure 7. Peak velocity increased with reward history. Experiment 1: A) Outgoing velocity significantly increased with prior reward reception B) Reward history was estimated based on an exponentially smoothed moving average fit to reaction time (see Methods). With greater reward history, outgoing peak velocity increased. C) Even when accounting for movements to the same reward expectation, reward history has a significant effect on outgoing peak velocity. Experiment 2: D) Prior reward reception did not influence the subsequent trial velocity. E) The effect of reward history was determined in the same manner as in experiment 1. Outgoing peak velocity significantly increased with the amount of reward received in recent trial history. F) Vigor of reaching movements towards targets of similar learned value, were influenced by recent trial reward history.

398 Lastly, including both learned value and reward history in a single generalized regression model
399 significantly improved model likelihood, compared to an alternative model that excluded the reward
400 history term (LRT: χ_1^2 =30.156, p=3.987e-8). This result strengthens our argument that both recent,

401   target-agnostic rewards and effort-conscious valuation of specific targets influence the motivation
402   of movement.

DISCUSSION

Here, we demonstrated that reach vigor tracks neural correlates of learning and motivation across
time scales ranging from milliseconds to minutes. Velocity was modulated by reward expectation,
reward prediction error and reward rate, key variables that have also been associated with striatal
dopaminergic fluctuations. These results point to a potential neural mechanism by which dopamine
can provide the bridge between decision making and movement control.

Reward prediction error is typically examined with respect to recent reward history and current target
expectation. Saccadic reaction times are faster or slower following a positive or negative reward
prediction error[42]. In this study, we are the first to show RPE can have a modulatory effect of an
ongoing movement, leading to either acceleration or deceleration that aligns with the sign of the
prediction error. Moreover, we find that this effect scales with the magnitude of the prediction error.
The rapid reflection of RPEs in vigor may be related to vigor's already known response to average
reward rate of the environment[27,43], which is tracked by tonic dopamine levels[21,26]. Striatal dopamine
levels rapidly rise and fall as a consequence of the temporal integration of phasic prediction error
signals[32,44]. It could be through this mechanism that we see rapid sub-second changes in vigor.
However, the exact relationship between spiking activity and striatal dopamine levels is
multifaceted[29,45] and the impact we observe on vigor is small and fleeting. This may explain why
targeted optogenetic manipulations were limited or temporally restricted in their effects on vigor[34].

A substantial body of evidence has detailed the relationship between midbrain DA transients and
action initiation[46]. In both freely-moving, and head-fixed mice, phasic activity accompanied
movement initiation[34,47], with its correlation to the ensuing movement's vigor being context
dependent. Likewise, many studies found significant relationships between phasic DAN activity and
reward-seeking motor behaviors[20,35,48].

It may seem unlikely that the absence or presence of the reward feedback visual stimulus would
influence ongoing velocity at sub-second speeds, approximately 215 ms after stimuli presentation.
However, neurons in the superior colliculus, that then project to dopaminergic neurons within the
VTA or SNc, have a response latency between 40 and 60 ms[49]. Another study found median response
latencies of cue-elicited DAN responses within the SNc were on the order of 112 ms[50]. Others still
have found that modulation of rapid motor responses, on the order of 150 ms, is dependent on task
specific visuospatial features[51]. Reward prediction error modulation in saccade latencies were found
to be affected on the order of ~150 ms[42]. A more recent study has found that changes in direction of
movement, specifically when a reach target suddenly decreases in expected value, occur on average
~248 ms after cue[52]. This body of literature detailing the short-latency rapid motor response suggests
that the rapid response in relative velocity that we see can be feasibly attributed to tuned
sensorimotor reward-based predictions.

To generate these prediction errors, we provided individuals with stochastic reward feedback
independent of motor performance. Previous research has focused on either the effect of varying
reward levels[53–55], or the difference between predictable and unpredictable reward[56,57] on the

invigoration of movement. One other work has found a continuous relationship between movement
response tempo (i.e. execution time) and reward-contingency belief[58]. Our work in this study details
how the reward-vigor relationship in human reaching is continuous with varying degrees of
probabilistic expectation as would be predicted based on the utility-maximization theoretic basis[38].
In both explicit contexts where the reward probabilities are stated, and within an implicit context
where individuals must learn and experience the differences in expectations, greater reward
probabilities were associated with greater average movement vigor.

Beyond the effects of within-trial reward and prediction error, we investigated the relationship
between biomechanical effort, subjective choice preference, and movement vigor. Individual
willingness to expend effort has been found to be idiosyncratic[59]. We build upon this finding in our
current study, showing that individuals differed in their willingness to expend biomechanical effort to
accrue monetary reward, and that these differences were demonstrated both in their choices and
kinematic behavior during the preceding single-target trials. Specifically, the magnitude of effect of
target-direction on choice preference was correlated with the magnitude of the effect on an
individual's movement vigor (Figure 6B).

Aside from reward-mediated invigorative effects of DA, both in movement and deliberation, other
research has found the neurotransmitter instrumental in overcoming costs and providing motivation
in the face of effortful action[60–64], which has been shown to influence motor behavior and deliberation
[27,39,65–67]. However, phasic midbrain DAN activity may not necessarily code for upcoming effort
value[68]. In monkey (Macaca mulatta) SNc, only a minority 13% of spiking activity in a reaching task
was correlated with net utility (rewards minus cost) compared to reward-alone. One hypothesis is
that the phasic DA response incorporates effort costs only if reward rate, or discounted value of
future action, were meaningfully affected[64]. Other neurotransmitters and pathways may be
implicated in utility prediction. Specific serotonin receptor activation in mice was found to increase
motivation and vigor alongside an increase in dorsomedial striatum extracellular DA[69]. Thus, the
effect of variable effort on the behavioral and dopaminergic response to reward prediction error
remains an open question.

Here we use a delta-rule learning model to estimate and quantify the trial-to-trial, subjective value
of a given target, motivated by its success in literature modelling prediction error responses in human
dopaminergic systems[19,70–73]. We found that movement vigor reflected the learned value, updated
from trial to trial[43,58]. A critical assumption in our model is that a value prediction error that integrates
reward and effort updates learned value, rather than solely a reward prediction error. Effort too has
been shown to influence neural activity in rat ventral striatum and dorsal anterior cingulate cortex
(ACCd)[74,75]. Likewise, human ACCd activity was found to reflect the interaction between both
expected reward and expected effort[76]. We also modeled effort as additively discounting
experienced reward, rather than multiplicatively as in previous work[77]. Additive discounting accounts
for the invigorating effect of reward and aligns with the conceptual framework that the brain
modulates movement vigor so as to maximize reward rate[38,78]. Our approach is validated in our data
where the value prediction error, which integrates learned reward and effort costs, has a significant
effect on both the within-trial change in velocity and trial-to-trial changes (Figure 6H,I).

Nonetheless, an alternative approach is one where the prediction error does not incorporate the
experienced target effort, which would be modeled as follows:

$$483 \qquad \hat{R}'^T_i = \hat{R}^T_i + \eta_s\left(R_i - \hat{R}^T_i\right)$$

$$484 \qquad V^T_i = \hat{R}^T_i + e^T_s$$

Here, learning is driven by reward prediction alone, and behavior is a result of both learned reward
and already known effort. Both models were fit to choice data and compared. The first model, with
learning driven from a value prediction error, had greater conditional R^2, 0.686, compared to the
alternative (0.543) when predicting participant choice. However, AICc scores show a preference
towards model 2: 2890.61 compared to 2917.76. From our experimental design, the kinematic data
would be equally well described with either model, as effort values were constant. Ultimately, the
evidence is ambiguous to state conclusively which of the feasible delta-rule models for learning
target value best describes the data. Our experiment did not originally set out to test these
distinctions, and so additional follow-up with the requisite design considerations would be needed.

The results highlight the differences between an experienced stochasticity and described one[79],
although our first experiment may be better described as a hybrid protocol, as trial-to-trial feedback
is still provided to reinforce the previously stated reward expectations. The use of kinematic
responses, specifically peak velocity on outgoing and return portions of the movement, are useful
tools in understanding not just these differences[80], but also the similarities between the two
environments. The effect of reward prediction error was similar, occurring within the same time frame
and showing comparable magnitudes of response. The most apparent distinction was the influence
of recent reward history. In our second experiment, the found relationship between prior reward and
velocity can be said to have a longer view of the past, integrating incrementally over many trials to
arrive in near proximity to the reward average (0.5). The first experiment could be said to elicit more
impulsive or rapid response, with the influence of trials further back in time quickly diminishing. This
behavior may speak to the difference in environmental uncertainties, where average reward
expectation is immediately known to the participant compared to when it must be experienced and
learned over time. Previous research has found significant effects of environmental uncertainty in
mice and monkey reward learning rates[81], with greater uncertainty resulting in reducing update rates,
which would explain differences in reward history α-coefficients between the two experiments.

Conclusion: Across two experiments, we demonstrate the sensitivity of movement vigor to
dopaminergic correlates of learning and motivation. As the likelihood of reward feedback increased,
so too did vigor. Movement vigor was also rapidly updated during an ongoing movement, in
proportional response to reward prediction error. Trial-specific learned value, modelled via a delta-
rule update formulation, could incorporate both rewards and subjective efforts to predict individual
reaching vigor. Lastly, target-agnostic reward history significantly influenced vigor, even when the
value expectation was matched. Taken together, our results highlight the link between known short-
latency dopaminergic learning signals and the invigoration of movement, not only at the time of cue
presentation and movement initiation, but during an ongoing movement immediately after feedback
is provided.

MATERIALS AND METHODS

Participants: The study consisted of two experiments, each with an independent population (first:
n=42, f=16, age=22.3±0.76; second: n=22, f=16, age=23.5±0.9). All individuals were either
ambidextrous or right-handed as determined by the Edinburgh Handedness Inventory survey, as well
as free of upper extremity injury or self-reported neurological condition. In the first experiment,
participants were compensated at a fixed rate of $10 per hour with no difference depending on
performance. For the second experiment, the $10/hr rate remained with the potential of an additional
$5 depending on performance (average compensation = $14.02±0.13). Participants provided written
and informed consent, and procedures were approved of by the University of Colorado Boulder
institutional review board.

Experimental Design: Participants performed a bandit-like task using the KINARM end-point robotic
arm (BKIN Technologies, ON, CA) to control a white cursor presented onscreen. From a home circle
(r=1cm) at the center of a ring of radius 10 cm, individuals were instructed to make an out-and-back
reaching motion to move a cursor (r=0.5cm) a cued target (r=1cm), displayed at 45, 135, 225, or 315
degrees relative to the home circle. Target cues were displayed at a variable time interval uniformly
ranging from 800 ms to 1000 ms after trial initiation. On reaching to the cued target, participants need
not hit the prompted target exactly. If absolute angular error at the 10 cm radial distance was less
than 22°, the target was counted as "hit." If individuals failed to hit the prompted target, either within
the ±22° accuracy constraint or within 4 seconds, a large red "X" appeared onscreen with an
accompanying tone (400 Hz, 100 ms duration) indicating a trial failure.

Individuals were instructed that each target had a unique probability (100%, 66%, 33%, or 0%) of
providing rewarding feedback at the moment of target hit. Reward feedback consisted of a brief, high-
pitched tone (880 Hz, 100 ms duration) being played, the target instantly doubling in size, changing
color to yellow, and blinking intermittently (2 blinks, on-off duration at 50 ms). Sample trial
progression is shown in figure 1c. Reward probability per target changed between blocks, with a total
of 4 blocks each consisting of 180 trials. Trial order and rewards were pseudorandomized such that
for each set of 36 trials, each target was cued 9 times, and reward feedback per target was given 9,
6, 3 or 0 times for the reward expectations of 100%, 66%, 33%, and 0% respectively. Individuals were
not told the total number of trials to be experienced, only that the total time in the experiment would
require approximately 1 hour.

Reward prediction error was defined as the difference between the received feedback and the
presented target's reward expectation, i.e.:

$$R_i \in \{0,1\}$$

$$RPE_i = R_i - E[R_i^T]$$

Thus, the four target reward probabilities led to five different reward prediction errors varying in both
magnitude and sign. The probabilistic targets (33% and 66%) led to both smaller and larger, positive
and negative reward prediction errors. The deterministic targets (0% and 100%) led to 0 RPE, but also
allowed us to compare the effects of reward feedback per se, independent of reward prediction error.

In experiment 1, participants were explicitly told which targets were to have what expectation of
reward within a block. In experiment 2, the rewards were identical to experiment 1 except that
participants were not told of the target reward probabilities but rather had to learn from experience

when reaching to the target when prompted. To quantify the degree to which they learned the reward
probabilities, the final 36 trials of each block were 'choice trials' where two targets were cued rather
than a single target. No reward feedback was provided within these choice trials after a target was
hit on the reach. Individuals were familiarized with the nature of these choice trials beforehand and
were informed that the underlying reward frequencies remained constant within a block. Each
unique choice pair (n=6) was presented six times during these choice trial periods. For choice trials,
we analyzed the choices as a function of expected value and expected reward. We also measured
the response time, the time between target presentation and movement onset. Participants were
instructed in the second experiment that additional compensation was contingent on the potential
rewards received during these choice trials and thus were incentivized to reach towards targets with
previously experienced greater expectation for reward feedback. Total bonus compensation was
calculated as the sum of chosen expected reward, divided by maximum potential expected chosen
reward (18.66), times five dollars:

575
$$\frac{\sum_i E[R]_{choice,i}}{18.66} \times \$5$$

In the first experiment, individuals were familiarized with the reaching task by completing 16 out-and-
back reaches, four each to the four different targets. In the second experiment, an additional 8 choice
trials were provided as familiarization. Reward feedback was withheld on all familiarization trials.

Kinematic Metrics: KINARM encoders sampled hand position and velocity (x,y cartesian coordinates)
at 1000 Hz. Raw kinematics data were filtered via third-order double pass filter with cutoff of 10 Hz.
Radial velocity was computed from numerical differentiation of the radial position relative to the
home circle using a second order centered finite difference. The primary metrics on each trial (both
single target trials and choice trials) were movement initiation time (reaction time), peak outgoing
and return radial velocities, maximum excursion radial distance, and movement duration from onset
to target hit. Reaction time, defined as the time between cue presentation and detected movement
onset, was calculated by use of the MACC-based onset detection method[82].

Trial Exclusion Criteria: For the first experiment, 1.83±0.35% of trials per participant on average were
excluded from kinematic analysis either due to failure to complete the trial, failing to reach the outer
target ring during the initial outward reach (a "double-peak" movement), or missing the target
completely (angular error was ≥22.5°). In the second experiment 7±2.37% of single-target trials per
participant on average were excluded from further kinematic analysis. These trials were included,
however, for the purposes of calculating reward history and recorded as providing no reward
feedback.

Linear Regression Models: To determine the relationship between velocity, target reward, and reward
prediction error, we used generalized Gamma linear mixed models with a log link function to estimate
the relative effects of factors of interest and to account for between-participant variability. The
Gamma distribution was selected after post-hoc residual analysis of a gaussian-residual
assumption revealed significant heteroskedasticity and skewness. The log-link was ultimately
selected after model comparisons to different potential link functions (inverse and identity). A similar
analysis was used to probe the relationship between reaction time and target reward expectation.

In the first experiment, our regression models for peak velocity and reaction time was as follows:

$$\log(\mu_y) = (1 + \text{Target} * \text{Repeat} + \text{Block} * \text{Trial} + \text{E}[\text{R}] + \text{Prior}_{\text{RWD}}) + (1 + \text{E}[\text{R}] + \text{Target}$$
$$+ \text{Block}|Subj)$$

With μ_y representing the mean of the outcome of interest, i.e., velocity or reaction time.

And for the second experiment, an additional interaction term with trial was added:
$$\log(\mu_y) = (1 + \text{Target} * \text{Repeat} + \text{Block} * \text{Trial} + \text{E}[\text{R}] + \text{Prior}_{\text{RWD}} + \text{E}[\text{R}]{:}\text{Trial}) + (1 + \text{E}[\text{R}]$$
$$+ \text{Target} + \text{Block}|Subj)$$

Target is treated as factored variable, and all others as continuous variables.
The regression model for return peak velocity was similar, but with added terms for reward reception
and prediction error while controlling for outward peak velocity:
$$\log(\mu_y) = (1 + \text{OutPV} * \text{Target} + \text{Repeat} + \text{Trial} + \text{RPE} * \text{Reward}) + (1 + \text{RPE} * \text{Reward} + \text{OutPV}$$
$$+ \text{Trial}|Subj)$$
For velocity difference models, a gaussian distribution was found to better approximate the
residuals, so a typical LMER with an identity link function was used instead of the generalized
Gamma model. All mixed regression models were fitted to restricted maximum likelihood (REML)
with the *lme4* and *lmerTest* packages for the *R* language[83,84], except for when model comparisons
were performed with the *performance* R package[85]. P-values for mixed model regression coefficients
are derived from Satterthwaite approximations. Hypothesis testing for linear combinations (LHT) of
regression coefficients and calculation of asymptotic Chi-squared test statistics was performed with
the linearHypothesis function in the *car* R package[86].

One-Dimensional SPM Analysis: Continuous time analysis of radial velocity was performed by one-
dimensional statistical parametric mapping[87]. Biomechanical curves are well-suited to analysis by
this methodology, owing to their smoothness and discrete bounds. We measured the potential effect
of reward feedback on different expectation contexts, either deterministic or stochastic, with either
single-sample t-tests or 2-factor ANOVAs respectively. This was done to both test for potential
significant effects as well as when such effect may occur on instantaneous radial velocity. We
focused on the time window from 150 ms prior to hitting the target (i.e. feedback reception) and 300
ms afterward.

Population-level inference on the effect of reward prediction error on within-trial normalized velocity
(taken as instantaneous velocity divided by the within-trial peak outgoing radial velocity) was
conducted in a manner following previous work[88]. First, regression analyzes were fitted per
participant, then this collection of 1-D beta values were submitted to a second-level single-sample
t-test.

Learning Model Design: To model the trial-to-trial update process in experiment 2, we developed a
Bayesian hierarchical Rescorla-Wagner learning model. In each of a block's single-target trials, the
presented target's estimated value was updated based on this rule:

$$V'^T_i = V^T_i + \eta_s(R_i + e^T_s - V^T_i)$$

With *T* the presented target at 45, 135, 225, or 315 degrees, *R* the reward feedback [0,1], *e* the effort
cost, *s*: participant, *i*: trial, and η: learning rate. Participant-specific learning rates (η_s) and effort costs

(e_s) were found via posterior maximum likelihood estimation based on choices. The full model (with
priors) is characterized as follows:

642
$$(Choice_i = A) \sim \text{Bernoulli} \left(\frac{1}{1 + \exp\left(-\tau_s\left(V^T_{A\,i} - V^T_{B\,i}\right)\right)} \right)$$

643
$$\bar{\zeta} \sim N(0,10); \ \bar{e} \sim N(0,5); \ \bar{\tau} \sim N(0,5)$$

644
$$\sigma_\zeta \sim \Gamma(2,0.01); \ \sigma_e \sim \Gamma(2,0.01); \ \sigma_\tau \sim \Gamma(2,0.01)$$

645
$$\zeta_s \sim N\left(\bar{\zeta},\sigma_\zeta\right); \ e_s \sim N(\bar{e},\sigma_e); \ \tau_s \sim N(\bar{\tau},\sigma_\tau)$$

646
$$\eta_s = \frac{1}{1 + \exp(-\zeta_s)}$$

647 In each choice trial, targets were randomly labelled as either A or B. Value prediction error was
648 defined as the sum of reward feedback and experienced effort minus learned value:

649
$$VPE_i = R_i + e^T_s - V^T_i$$

650 Sampling for Bayesian models was performed by NUTS Hamiltonian MCMC algorithm via use of the
651 *RStan* package[89].

652 *Reward History Model*: To investigate the effect of prior reward on excursion peak velocity, we applied
653 an exponential moving average to reward reception history[41]. For each single-target trial, average
654 reward was updated with this rule:
655
$$\bar{R}_{i+1} = \alpha R_i + (1 - \alpha)\bar{R}_i$$
656 The smoothing factor, α, was fit to maximize the likelihood of a reaction time random intercept
657 regression[20]:
658
$$\log(\mu_{RT}) \sim (1 + \text{Block} + \bar{R} + \text{Repeat} + \text{Trial} * E[R] + \text{Target}) + (1|Subj)$$
659 The effects of block number (Block), direction repetition (Repeat), trial number (Trial), reward
660 expectation (E[R]) and factored target direction (Target) were included as control variables. We then
661 used this model-derived reward history term to independently predict outgoing peak velocity in
662 subsequent linear models.

REFERENCES

1. Beierholm, U., Guitart-Masip, M., Economides, M., Chowdhury, R., Düzel, E., Dolan, R., and
Dayan, P. (2013). Dopamine Modulates Reward-Related Vigor. Neuropsychopharmacology *38*,
1495–1503. https://doi.org/10.1038/npp.2013.48.

2. Choi, J.E.S., Vaswani, P.A., and Shadmehr, R. (2014). Vigor of Movements and the Cost of Time
in Decision Making. J Neurosci *34*, 1212–1223. https://doi.org/10.1523/JNEUROSCI.2798-
13.2014.

3. Opris, I., Lebedev, M., and Nelson, R.J. (2011). Motor Planning under Unpredictable Reward:
Modulations of Movement Vigor and Primate Striatum Activity. Front Neurosci *5*.
https://doi.org/10.3389/fnins.2011.00061.

4. Reppert, T.R., Lempert, K.M., Glimcher, P.W., and Shadmehr, R. (2015). Modulation of Saccade
Vigor during Value-Based Decision Making. Journal of Neuroscience *35*, 15369–15378.
https://doi.org/10.1523/JNEUROSCI.2621-15.2015.

5. Shadmehr, R., and Ahmed, A.A. (2020). Vigor: Neuroeconomics of movement control (MIT
Press).

6. Summerside, E.M., Shadmehr, R., and Ahmed, A.A. (2018). Vigor of reaching movements:
reward discounts the cost of effort. Journal of Neurophysiology *119*, 2347–2357.
https://doi.org/10.1152/jn.00872.2017.

7. Smoulder, A.L., Marino, P.J., Oby, E.R., Snyder, S.E., Miyata, H., Pavlovsky, N.P., Bishop, W.E.,
Yu, B.M., Chase, S.M., and Batista, A.P. (2024). A neural basis of choking under pressure.
Neuron *112*, 3424-3433.e8. https://doi.org/10.1016/j.neuron.2024.08.012.

8. Korbisch, C.C., Apuan, D.R., Shadmehr, R., and Ahmed, A.A. (2022). Saccade vigor reflects the
rise of decision variables during deliberation. Current Biology *32*, 5374-5381.e4.
https://doi.org/10.1016/j.cub.2022.10.053.

9. Berret, B., and Jean, F. (2016). Why Don't We Move Slower? The Value of Time in the Neural
Control of Action. J. Neurosci. *36*, 1056–1070. https://doi.org/10.1523/JNEUROSCI.1921-
15.2016.

10. Marbaker, R.M., Schmad, R.C., Al-Ghamdi, R.A., Sukumar, S., and Ahmed, A.A. (2025). Reward
invigorates isometric gripping actions. Journal of Neurophysiology *133*, 1282–1294.
https://doi.org/10.1152/jn.00384.2024.

11. Ungerstedt, U. (1971). Adipsia and Aphagia after 6-Hydroxydopamine Induced Degeneration of
the Nigro-striatal Dopamine System. Acta Physiologica Scandinavica *82*, 95–122.
https://doi.org/10.1111/j.1365-201X.1971.tb11001.x.

12. Klaus, A., Alves da Silva, J., and Costa, R.M. (2019). What, If, and When to Move: Basal Ganglia
Circuits and Self-Paced Action Initiation. Annu. Rev. Neurosci. *42*, 459–483.
https://doi.org/10.1146/annurev-neuro-072116-031033.

13. Larry, N., Zur, G., and Joshua, M. (2024). Organization of reward and movement signals in the
basal ganglia and cerebellum. Nat Commun *15*, 2119. https://doi.org/10.1038/s41467-024-
45921-9.

14. Dudman, J.T., and Krakauer, J.W. (2016). The basal ganglia: from motor commands to the
control of vigor. Current Opinion in Neurobiology *37*, 158–166.
https://doi.org/10.1016/j.conb.2016.02.005.

15. Jurado-Parras, M.-T., Safaie, M., Sarno, S., Louis, J., Karoutchi, C., Berret, B., and Robbe, D.
(2020). The Dorsal Striatum Energizes Motor Routines. Current Biology *30*, 4362-4372.e6.
https://doi.org/10.1016/j.cub.2020.08.049.

16. Thura, D., and Cisek, P. (2017). The Basal Ganglia Do Not Select Reach Targets but Control the
Urgency of Commitment. Neuron *95*, 1160-1170.e5.
https://doi.org/10.1016/j.neuron.2017.07.039.

17. Herz, D.M., Bange, M., Gonzalez-Escamilla, G., Auer, M., Ashkan, K., Fischer, P., Tan, H.,
Bogacz, R., Muthuraman, M., Groppa, S., et al. (2022). Dynamic control of decision and
movement speed in the human basal ganglia. Nat Commun *13*, 7530.
https://doi.org/10.1038/s41467-022-35121-8.

18. Schultz, W. (1998). Predictive Reward Signal of Dopamine Neurons. Journal of Neurophysiology
80, 1–27. https://doi.org/10.1152/jn.1998.80.1.1.

19. Fiorillo, C.D., Tobler, P.N., and Schultz, W. (2003). Discrete coding of reward probability and
uncertainty by dopamine neurons. Science *299*, 1898–1902.
https://doi.org/10.1126/science.1077349.

20. Hamid, A.A., Pettibone, J.R., Mabrouk, O.S., Hetrick, V.L., Schmidt, R., Vander Weele, C.M.,
Kennedy, R.T., Aragona, B.J., and Berke, J.D. (2016). Mesolimbic dopamine signals the value of
work. Nat Neurosci *19*, 117–126. https://doi.org/10.1038/nn.4173.

21. Niv, Y., Daw, N.D., Joel, D., and Dayan, P. (2007). Tonic dopamine: opportunity costs and the
control of response vigor. Psychopharmacology (Berl.) *191*, 507–520.
https://doi.org/10.1007/s00213-006-0502-4.

22. Wang, Y., Toyoshima, O., Kunimatsu, J., Yamada, H., and Matsumoto, M. (2021). Tonic firing
mode of midbrain dopamine neurons continuously tracks reward values changing moment-by-
moment. eLife *10*, e63166. https://doi.org/10.7554/eLife.63166.

23. Ikemoto, S., and Panksepp, J. (1999). The role of nucleus accumbens dopamine in motivated
behavior: a unifying interpretation with special reference to reward-seeking. Brain Res Brain
Res Rev *31*, 6–41. https://doi.org/10.1016/s0165-0173(99)00023-5.

24. Salamone, J.D., and Correa, M. (2002). Motivational views of reinforcement: implications for
understanding the behavioral functions of nucleus accumbens dopamine. Behav Brain Res
137, 3–25. https://doi.org/10.1016/s0166-4328(02)00282-6.

25. Mah, A., Schiereck, S.S., Bossio, V., and Constantinople, C.M. (2023). Distinct value
computations support rapid sequential decisions. Nat Commun *14*, 1–14.
https://doi.org/10.1038/s41467-023-43250-x.

26. Rigoli, F., Chew, B., Dayan, P., and Dolan, R.J. (2016). The Dopaminergic Midbrain Mediates an
Effect of Average Reward on Pavlovian Vigor. Journal of Cognitive Neuroscience *28*, 1303–1317.
https://doi.org/10.1162/jocn_a_00972.

27. Sukumar, S., Shadmehr, R., and Ahmed, A.A. (2024). Effects of reward and effort history on
decision making and movement vigor during foraging. Journal of Neurophysiology *131*, 638–
651. https://doi.org/10.1152/jn.00092.2023.

28. Yoon, T., Geary, R.B., Ahmed, A.A., and Shadmehr, R. (2018). Control of movement vigor and
decision making during foraging. PNAS *115*, E10476–E10485.
https://doi.org/10.1073/pnas.1812979115.

29. Cai, X., Liu, C., Tsutsui-Kimura, I., Lee, J.-H., Guo, C., Banerjee, A., Lee, J., Amo, R., Xie, Y.,
Patriarchi, T., et al. (2024). Dopamine dynamics are dispensable for movement but promote
reward responses. Nature, 1–9. https://doi.org/10.1038/s41586-024-08038-z.

30. Engel, L., Wolff, A.R., Blake, M., Collins, V.L., Sinha, S., and Saunders, B.T. (2024). Dopamine
neurons drive spatiotemporally heterogeneous striatal dopamine signals during learning.
Current Biology *34*, 3086-3101.e4. https://doi.org/10.1016/j.cub.2024.05.069.

31. Howe, M.W., and Dombeck, D.A. (2016). Rapid signalling in distinct dopaminergic axons during
locomotion and reward. Nature *535*, 505–510. https://doi.org/10.1038/nature18942.

32. Mohebi, A., Wei, W., Pelattini, L., Kim, K., and Berke, J.D. (2024). Dopamine transients follow a
striatal gradient of reward time horizons. Nat Neurosci, 1–10. https://doi.org/10.1038/s41593-
023-01566-3.

33. Panigrahi, B., Martin, K.A., Li, Y., Graves, A.R., Vollmer, A., Olson, L., Mensh, B.D., Karpova, A.Y.,
and Dudman, J.T. (2015). Dopamine Is Required for the Neural Representation and Control of
Movement Vigor. Cell *162*, 1418–1430. https://doi.org/10.1016/j.cell.2015.08.014.

34. da Silva, J.A., Tecuapetla, F., Paixão, V., and Costa, R.M. (2018). Dopamine neuron activity
before action initiation gates and invigorates future movements. Nature *554*, 244–248.
https://doi.org/10.1038/nature25457.

35. Syed, E.C.J., Grima, L.L., Magill, P.J., Bogacz, R., Brown, P., and Walton, M.E. (2016). Action
initiation shapes mesolimbic dopamine encoding of future rewards. Nat Neurosci *19*, 34–36.
https://doi.org/10.1038/nn.4187.

36. Schall, T.A., Li, K.-L., Qi, X., Lee, B.T., Wright, W.J., Alpaugh, E.E., Zhao, R.J., Liu, J., Li, Q., Zeng,
B., et al. (2024). Temporal dynamics of nucleus accumbens neurons in male mice during
reward seeking. Nat Commun *15*, 9285. https://doi.org/10.1038/s41467-024-53690-8.

37. Jang, H.J., Ward, R.M., Golden, C.E.M., and Constantinople, C.M. (2024). Acetylcholine
demixes heterogeneous dopamine signals for learning and moving. Preprint at bioRxiv,
https://doi.org/10.1101/2024.05.03.592444 https://doi.org/10.1101/2024.05.03.592444.

38. Shadmehr, R., Huang, H.J., and Ahmed, A.A. (2016). A Representation of Effort in Decision-
Making and Motor Control. Current Biology *26*, 1929–1934.
https://doi.org/10.1016/j.cub.2016.05.065.

39. Pierrieau, E., Lepage, J.-F., and Bernier, P.-M. (2021). Action Costs Rapidly and Automatically
Interfere with Reward-Based Decision-Making in a Reaching Task. eNeuro *8*.
https://doi.org/10.1523/ENEURO.0247-21.2021.

40. Wagner, A.R., and Rescorla, R.A. (1972). Inhibition in Pavlovian conditioning: Application of a
theory. Inhibition and learning, 301–336.

41. Daw, N.D., Kakade, S., and Dayan, P. (2002). Opponent interactions between serotonin and
dopamine. Neural Networks *15*, 603–616. https://doi.org/10.1016/S0893-6080(02)00052-7.

42. Sedaghat-Nejad, E., Herzfeld, D.J., and Shadmehr, R. (2019). Reward Prediction Error
Modulates Saccade Vigor. J. Neurosci. *39*, 5010–5017.
https://doi.org/10.1523/JNEUROSCI.0432-19.2019.

43. Guitart-Masip, M., Beierholm, U.R., Dolan, R., Duzel, E., and Dayan, P. (2011). Vigor in the Face
of Fluctuating Rates of Reward: An Experimental Examination. Journal of Cognitive
Neuroscience *23*, 3933–3938. https://doi.org/10.1162/jocn_a_00090.

44. Mohebi, A., Pettibone, J.R., Hamid, A.A., Wong, J.-M.T., Vinson, L.T., Patriarchi, T., Tian, L.,
Kennedy, R.T., and Berke, J.D. (2019). Dissociable dopamine dynamics for learning and
motivation. Nature *570*, 65–70. https://doi.org/10.1038/s41586-019-1235-y.

45. Sippy, T., and Tritsch, N.X. (2023). Unraveling the dynamics of dopamine release and its actions
on target cells. Trends in Neurosciences *46*, 228–239.
https://doi.org/10.1016/j.tins.2022.12.005.

46. Coddington, L.T., and Dudman, J.T. (2019). Learning from Action: Reconsidering Movement
Signaling in Midbrain Dopamine Neuron Activity. Neuron *104*, 63–77.
https://doi.org/10.1016/j.neuron.2019.08.036.

47. Dodson, P.D., Dreyer, J.K., Jennings, K.A., Syed, E.C.J., Wade-Martins, R., Cragg, S.J., Bolam, J.P.,
and Magill, P.J. (2016). Representation of spontaneous movement by dopaminergic neurons is
cell-type selective and disrupted in parkinsonism. Proceedings of the National Academy of
Sciences *113*, E2180–E2188. https://doi.org/10.1073/pnas.1515941113.

48. Coddington, L.T., and Dudman, J.T. (2018). The timing of action determines reward prediction
signals in identified midbrain dopamine neurons. Nat Neurosci *21*, 1563–1573.
https://doi.org/10.1038/s41593-018-0245-7.

49. Dommett, E., Coizet, V., Blaha, C.D., Martindale, J., Lefebvre, V., Walton, N., Mayhew, J.E.W.,
Overton, P.G., and Redgrave, P. (2005). How Visual Stimuli Activate Dopaminergic Neurons at
Short Latency. Science *307*, 1476–1479. https://doi.org/10.1126/science.1107026.

50. Ravel, S., and Richmond, B.J. (2006). Dopamine neuronal responses in monkeys performing
visually cued reward schedules. European Journal of Neuroscience *24*, 277–290.
https://doi.org/10.1111/j.1460-9568.2006.04905.x.

51. Pruszynski, J.A., King, G.L., Boisse, L., Scott, S.H., Flanagan, J.R., and Munoz, D.P. (2010).
Stimulus-locked responses on human arm muscles reveal a rapid neural pathway linking visual
input to arm motor output. European Journal of Neuroscience *32*, 1049–1057.
https://doi.org/10.1111/j.1460-9568.2010.07380.x.

52. Marti-Marca, A., Deco, G., and Cos, I. (2020). Visual-reward driven changes of movement during
action execution. Sci Rep *10*, 15527. https://doi.org/10.1038/s41598-020-72220-2.

53. Codol, O., Holland, P.J., Manohar, S.G., and Galea, J.M. (2020). Reward-Based Improvements in
Motor Control Are Driven by Multiple Error-Reducing Mechanisms. J. Neurosci. *40*, 3604–3620.
https://doi.org/10.1523/JNEUROSCI.2646-19.2020.

54. Sporn, S., Chen, X., and Galea, J.M. (2022). The dissociable effects of reward on sequential
motor behavior. Journal of Neurophysiology *128*, 86–104.
https://doi.org/10.1152/jn.00467.2021.

55. Summerside, E.M., Shadmehr, R., and Ahmed, A.A. (2018). Vigor of reaching movements:
reward discounts the cost of effort. Journal of Neurophysiology *119*, 2347–2357.
https://doi.org/10.1152/jn.00872.2017.

56. Glaser, J.I., Wood, D.K., Lawlor, P.N., Ramkumar, P., Kording, K.P., and Segraves, M.A. (2016).
Role of expected reward in frontal eye field during natural scene search. Journal of
Neurophysiology *116*, 645–657. https://doi.org/10.1152/jn.00119.2016.

57. Opris, I., Lebedev, M., and Nelson, R.J. (2011). Motor Planning under Unpredictable Reward:
Modulations of Movement Vigor and Primate Striatum Activity. Front. Neurosci. *5*.
https://doi.org/10.3389/fnins.2011.00061.

58. Tecilla, M., Großbach, M., Gentile, G., Holland, P., Sporn, S., Antonini, A., and Ruiz, M.H. (2023).
Modulation of Motor Vigor by Expectation of Reward Probability Trial-by-Trial Is Preserved in
Healthy Ageing and Parkinson's Disease Patients. J. Neurosci. *43*, 1757–1777.
https://doi.org/10.1523/JNEUROSCI.1583-22.2022.

59. Reppert, T.R., Rigas, I., Herzfeld, D.J., Sedaghat-Nejad, E., Komogortsev, O., and Shadmehr, R.
(2018). Movement vigor as a traitlike attribute of individuality. Journal of Neurophysiology *120*,
741–757. https://doi.org/10.1152/jn.00033.2018.

60. Collins, A.G.E., and Frank, M.J. (2016). Surprise! Dopamine signals mix action, value and error.
Nat Neurosci *19*, 3–5. https://doi.org/10.1038/nn.4207.

61. Hagura, N., Haggard, P., and Diedrichsen, J. (2017). Perceptual decisions are biased by the cost
to act. eLife *6*, e18422. https://doi.org/10.7554/eLife.18422.

62. Kurniawan, I., Guitart-Masip, M., and Dolan, R. (2011). Dopamine and Effort-Based Decision
Making. Frontiers in Neuroscience *5*.

63. Salamone, J., Correa, M., Farrar, A., Nunes, E., and Pardo, M. (2009). Dopamine, behavioral
economics, and effort. Frontiers in Behavioral Neuroscience *3*.

64. Walton, M.E., and Bouret, S. (2019). What Is the Relationship between Dopamine and Effort?
Trends in Neurosciences *42*, 79–91. https://doi.org/10.1016/j.tins.2018.10.001.

65. Courter, R.J., Alvarez, E., Enoka, R.M., and Ahmed, A.A. (2023). Metabolic costs of walking and
arm reaching in persons with mild multiple sclerosis. Journal of Neurophysiology *129*, 819–832.
https://doi.org/10.1152/jn.00373.2022.

66. Summerside, E.M., and Ahmed, A.A. (2021). Using metabolic energy to quantify the subjective
value of physical effort. Journal of The Royal Society Interface *18*, 20210387.
https://doi.org/10.1098/rsif.2021.0387.

67. Schweighofer, N., Xiao, Y., Kim, S., Yoshioka, T., Gordon, J., and Osu, R. (2015). Effort, success,
and nonuse determine arm choice. Journal of Neurophysiology *114*, 551–559.
https://doi.org/10.1152/jn.00593.2014.

68. Pasquereau, B., and Turner, R.S. (2013). Limited encoding of effort by dopamine neurons in a
cost-benefit trade-off task. J Neurosci *33*, 8288–8300.
https://doi.org/10.1523/JNEUROSCI.4619-12.2013.

69. Bailey, M.R., Goldman, O., Bello, E.P., Chohan, M.O., Jeong, N., Winiger, V., Chun, E., Schipani,
E., Kalmbach, A., Cheer, J.F., et al. (2018). An Interaction between Serotonin Receptor Signaling
and Dopamine Enhances Goal-Directed Vigor and Persistence in Mice. J. Neurosci. *38*, 2149–
2162. https://doi.org/10.1523/JNEUROSCI.2088-17.2018.

70. Daw, N.D., O'Doherty, J.P., Dayan, P., Seymour, B., and Dolan, R.J. (2006). Cortical substrates
for exploratory decisions in humans. Nature *441*, 876–879.
https://doi.org/10.1038/nature04766.

71. Maes, E.J.P., Sharpe, M.J., Usypchuk, A.A., Lozzi, M., Chang, C.Y., Gardner, M.P.H.,
Schoenbaum, G., and Iordanova, M.D. (2020). Causal evidence supporting the proposal that
dopamine transients function as temporal difference prediction errors. Nat Neurosci *23*, 176–
178. https://doi.org/10.1038/s41593-019-0574-1.

72. Starkweather, C.K., and Uchida, N. (2021). Dopamine signals as temporal difference errors:
recent advances. Current Opinion in Neurobiology *67*, 95–105.
https://doi.org/10.1016/j.conb.2020.08.014.

73. Steinberg, E.E., Keiflin, R., Boivin, J.R., Witten, I.B., Deisseroth, K., and Janak, P.H. (2013). A
causal link between prediction errors, dopamine neurons and learning. Nat Neurosci *16*, 966–
973. https://doi.org/10.1038/nn.3413.

74. Schweimer, J., and Hauber, W. (2005). Involvement of the rat anterior cingulate cortex in control
of instrumental responses guided by reward expectancy. Learn. Mem. *12*, 334–342.
https://doi.org/10.1101/lm.90605.

75. Walton, M.E., Bannerman, D.M., Alterescu, K., and Rushworth, M.F.S. (2003). Functional
Specialization within Medial Frontal Cortex of the Anterior Cingulate for Evaluating Effort-
Related Decisions. J. Neurosci. *23*, 6475–6479. https://doi.org/10.1523/JNEUROSCI.23-16-
06475.2003.

76. Croxson, P.L., Walton, M.E., O'Reilly, J.X., Behrens, T.E.J., and Rushworth, M.F.S. (2009). Effort-
Based Cost–Benefit Valuation and the Human Brain. J. Neurosci. *29*, 4531–4541.
https://doi.org/10.1523/JNEUROSCI.4515-08.2009.

77. Klein-Flügge, M.C., Kennerley, S.W., Saraiva, A.C., Penny, W.D., and Bestmann, S. (2015).
Behavioral Modeling of Human Choices Reveals Dissociable Effects of Physical Effort and
Temporal Delay on Reward Devaluation. PLOS Computational Biology *11*, e1004116.
https://doi.org/10.1371/journal.pcbi.1004116.

78. Emlen, J.M. (1966). The Role of Time and Energy in Food Preference. The American Naturalist
100, 611–617. https://doi.org/10.1086/282455.

79. Garcia, B., Cerrotti, F., and Palminteri, S. (2021). The description–experience gap: a challenge
for the neuroeconomics of decision-making under uncertainty. Philosophical Transactions of
the Royal Society B: Biological Sciences *376*, 20190665.
https://doi.org/10.1098/rstb.2019.0665.

80. Fitzgerald, T.H.B., Seymour, B., Bach, D.R., and Dolan, R.J. (2010). Differentiable neural
substrates for learned and described value and risk. Curr Biol *20*, 1823–1829.
https://doi.org/10.1016/j.cub.2010.08.048.

81. Woo, J.H., Aguirre, C.G., Bari, B.A., Tsutsui, K.-I., Grabenhorst, F., Cohen, J.Y., Schultz, W.,
Izquierdo, A., and Soltani, A. (2023). Mechanisms of adjustments to different types of
uncertainty in the reward environment across mice and monkeys. Cogn Affect Behav Neurosci
23, 600–619. https://doi.org/10.3758/s13415-022-01059-z.

82. Botzer, L., and Karniel, A. (2009). A simple and accurate onset detection method for a
measured bell-shaped speed profile. Frontiers in Neuroscience *3*.

83. Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting Linear Mixed-Effects Models
Using lme4. Journal of Statistical Software *67*, 1–48. https://doi.org/10.18637/jss.v067.i01.

84. Kuznetsova, A., Brockhoff, P.B., and Christensen, R.H.B. (2017). lmerTest Package: Tests in
Linear Mixed Effects Models. Journal of Statistical Software *82*, 1–26.
https://doi.org/10.18637/jss.v082.i13.

85. Lüdecke, D., Ben-Shachar, M.S., Patil, I., Waggoner, P., and Makowski, D. (2021). performance:
An R Package for Assessment, Comparison and Testing of Statistical Models. Journal of Open
Source Software *6*, 3139. https://doi.org/10.21105/joss.03139.

86. Fox, J., and Weisberg, S. (2019). An R Companion to Applied Regression Third. (Sage).

87. Pataky, T.C. (2010). Generalized n-dimensional biomechanical field analysis using statistical
parametric mapping. Journal of Biomechanics *43*, 1976–1982.
https://doi.org/10.1016/j.jbiomech.2010.03.008.

88. Beckmann, C.F., Jenkinson, M., and Smith, S.M. (2003). General multilevel linear modeling for
group analysis in FMRI. NeuroImage *20*, 1052–1063. https://doi.org/10.1016/S1053-
8119(03)00435-X.

89. Stan Development Team (2024). RStan: the R interface to Stan.