Metaproteome analysis of short-term thermal stress in three sympatric coral species reveals divergent host responses.

Authors

Shrinivas Nandi[1,†], Timothy G. Stephens[1,†], Erin E. Chille[1], Samantha Goyen[2], Line K. Bay[2,*], Debashish Bhattacharya[1,*]

Affiliations

[1]Department of Biochemistry and Microbiology, Rutgers University, New Brunswick, New Jersey, 08901, USA

[2]Australian Institute of Marine Science, Townsville, Queensland, Australia

[†]Equal contribution; co-first authors.

[*]Corresponding authors: Debashish Bhattacharya, dbhattac@rutgers.edu; Line K. Bay, L.Bay@aims.gov.au

Abstract

The accelerating loss of coral reefs worldwide due to anthropogenic climate change has led to a myriad of studies aimed at understanding the basis of coral resilience to support reef conservation. Here, we integrate physiological measurements with proteomic and metabolomic profiles to examine species-specific responses to increased temperature in three sympatric reef-building corals from the Great Barrier Reef: *Acropora hyacinthus*, *Porites lobata*, and *Stylophora pistillata*. We find species-specific stress response strategies with *A. hyacinthus*, a thermally sensitive species, exhibiting rapid decline in endosymbiont physiology, coupled with a one-third reduction in protein abundance. In contrast, *P. lobata* displayed a delayed physiological response to stress and a muted proteome response, suggesting greater resilience. *S. pistillata* initially showed minor shifts in the proteome followed by colony "bail-out". Overall, we observed markedly different responses in most biochemical pathways in the three coral species. Nonetheless, some known biomarkers of stress, including heat-shock proteins, showed conserved responses to thermal stress with differences in temporal abundance reflecting bleaching resistance. Our results underscore the species-specific nature of coral responses to thermal stress and highlight proteomic signatures associated with symbiosis breakdown, offering mechanistic insights into coral bleaching susceptibility and resilience.

Keywords: Coral bleaching, Great Barrier Reef, holobiont, metaproteomics, protein abundance, stony corals, thermal stress.

Introduction

Stony corals (Scleractinia) thrive in oligotrophic waters due to their ability to form a symbiotic association with dinoflagellate algae in the family Symbiodiniaceae (Gordon and Leggat, 2010). These photosynthetic endosymbionts sustain the host coral through the transfer of photosynthates (Muscatine and Porter, 1977). In exchange, corals provide Symbiodiniaceae with a stable, sheltered environment, inorganic carbon, and other essential nutrients (Frankowiak et al., 2016). Algal symbionts are contained within host gastrodermal cells in symbiosome compartments which provide an intracellular microenvironment optimized for photosynthesis (Barott et al., 2022, 2015; Bertucci et al., 2013; Thies et al., 2022). Under stressful conditions (e.g., high nutrient enrichment, cold- and heat-stress), coral hosts may lose their endosymbiotic algae, which is known as coral bleaching. During bleaching, some corals can sustain themselves for extended periods of time through heterotrophic feeding (Grottoli et al., 2006). However, bleached corals are more susceptible to disease and may succumb to starvation if the symbiosis is not reestablished (Bourne et al., 2016; Muller et al., 2008). The increase in mean global oceanic temperature, a result of anthropogenic climate change, has led to increased bleaching episodes and a significant decline in global coral populations (Hughes et al., 2017).

Although coral bleaching is a conserved stress response across Scleractinia, bleaching risk and the degree of bleaching for a colony depends on several factors. Coral bleaching responses are shaped by a complex interplay of ecological, physiological, and evolutionary factors, leading to species-specific differences in susceptibility (Loya et al., 2001; Woesik et al., 2011). Bleaching outcomes under different levels of thermal stress have been extensively studied in the Great Barrier Reef, Australia (Burn et al., 2023; Hughes et al., 2019, 2018, 2017; Pratchett et al., 2020). For instance, massive and slow-growing corals such as *Porites* often exhibit greater thermal tolerance when compared to branching, fast-growing species such as *Acropora*, which tend to bleach more readily (Burn et al., 2023; Fitt et al., 2009; Marshall and Baird, 2000; Pratchett et al., 2020), likely reflecting different strategies for energy allocation (i.e., resilience vs. rapid growth) of these long-lived sessile organisms under dyanamic environmental conditions. Variations in skeletal morphology (Swain et al., 2018, 2016), tissue thickness (Barkley et al., 2018; Qin et al., 2020), and metabolic strategies, such as heterotrophic plasticity (Grottoli et al., 2006; Martinez et al., 2024; Sangmanee et al., 2020), can contribute to differential tolerance to thermal and other environmental stressors. Variations in algal symbiont composition have been extensively studied, with some algal lineages conferring higher thermal tolerance to the coral colony (Bhattacharya et al., 2024; Buerger et al., 2020; Putnam et al., 2017; Rowan, 2004; Wang et al., 2023; Ziegler et al., 2018). The prokaryotic microbiome may also play a major role in coral holobiont resilience (Peixoto et al., 2017; Reshef et al., 2006; Rosado et al., 2019; Ziegler et al., 2017). In addition, local environmental conditions including water flow, light intensity, nutrient levels, and previous exposure to stress modulate bleaching susceptibility. Corals in high-flow or turbid environments may experience stress buffering, whereas those inhabiting sheltered, low-flow conditions are more likely to experience thermal extremes (Morgan et al., 2017; Risk and Edinger, 2011). Prior exposure to sublethal stress, or high frequency thermal variability can induce acclimatization or select for stress-tolerant genotypes, resulting in site- and species-specific patterns in bleaching severity (Barshis et al., 2013; Kenkel et al., 2013; Oliver and Palumbi, 2011; Safaie et al., 2018).

Developing reliable molecular biomarkers to detect early stress responses in corals is a major goal in reef monitoring and conservation. These tools will enable rapid, pre-bleaching health

assessment which can guide targeted interventions (Chille et al., 2025; Downs et al., 2005).
Transcriptomic datasets have been extensively used to identify putative biomarkers of
thermal stress (Chille et al., 2025; Cowen and Putnam, 2022; Cziesielski et al., 2019).
However, a central challenge is that stress produces high variation in gene expression levels
in different species (Da-Anoy et al., 2024; Ip et al., 2022; Rosic et al., 2020), and even among
genotypes within a species (Chille et al., 2024; Kenkel and Matz, 2016). Nonetheless,
transcriptomic studies have identified some marker genes of thermal stress, including those
encoding heat-shock proteins (e.g., HSP70, HSP90) (Nakamura et al., 2011; Rosic et al.,
2011; Seveso et al., 2016), ubiquitin-related proteins (DeSalvo et al., 2008; Rodriguez-
Lanetty et al., 2010, 2009; Seneca and Palumbi, 2015), and antioxidant enzymes (Downs et
al., 2002; Krueger et al., 2015) among others. A recent review of 307 gene and protein
expression papers studying stress response in cnidarians found that most genes have
inconsistent responses across different experiments, with only 14 having shared responses
across at least two species tested and NF-κB having shared responses across four species,
thereby suggesting a high degree of species specificity vis-à-vis the coral stress response
(Molinari et al., 2025).

The use of proteomics in coral research is relatively under-developed and may enhance
biomarker identification by providing a more direct measure of the functional molecules that
drive cellular processes, capturing post-transcriptional regulation and protein modification
that transcriptomics alone cannot address (Greenbaum et al., 2003). In addition, protein
biomarkers offer an advantage to mRNA because they are more readily integrated into
existing point-of care diagnostic tools (Chille et al., 2025). Research across diverse
eukaryotic systems, including the coral *Montipora capitata*, demonstrates that transcriptomic
data often poorly represent the proteome, with weak correlations observed between changes
in transcript abundance and accumulation of the corresponding proteins (Williams et al.,
2023). Thus, proteomic data can provide a rich resource not only for biomarker development,
but also for further elucidation of coral thermal stress responses (Chille et al., 2025).

Metabolomics provides complementary insights to proteomic data by capturing the end
products of cellular activity, i.e., metabolites that directly reflect the real-time physiological
state of the organism (Fiehn, 2002; Roessner and Bowne, 2009). Previous metabolomic
studies in corals have reported changes in antioxidant-related metabolites during thermal
stress (Williams et al., 2021b) and shifts in amino acid pools (Chiles et al., 2022; Martinez et
al., 2020, 2022a). Notably, recent findings have demonstrated the accumulation of specific
dipeptides as reliable indicators of pre-bleaching thermal stress. These include lysine-
glutamine (KQ), arginine-glutamine (RQ), arginine-alanine (RA), and arginine-valine (RV)
(Huffmyer et al., 2024; Williams et al., 2021a).

In this study, we used proteomic data to investigate the coral holobiont stress response in
three sympatric species from the Great Barrier Reef (GBR), Australia: *Acropora hyacinthus*,
Porites lobata, and *Stylophora pistillata* [**Figure 1**]. These species were selected due to their
known differences in thermal tolerance: i.e., sensitive, resilient, and low-to-moderate
sensitivity, respectively (Burn et al., 2023; Fitt et al., 2009; Manalili et al., 2025; Marshall
and Baird, 2000; Pratchett et al., 2020). We examined key metabolic pathways in each
species to better understand their responses to thermal stress and searched for conserved
proteins across species that can potentially be used as biomarkers. Lastly, we investigated the
metabolomic data to validate previously reported shifts in metabolite abundance, including
dipeptides, amino acids, and antioxidants that play key roles in the coral thermal stress
response.

Results

Host bailout and mortality
No mortality was observed in *A. hyacinthus* and *P. lobata*. In contrast, *S. pistillata* exhibited
significant tissue loss (i.e. "bail-out") and loss of color (becoming paler) based on visual
assessment at TP1 [**Supplemental Figure 1**] and complete tissue loss was observed by TP2,
in most fragments.

Endosymbiont community composition and physiology
Because coral bleaching is associated with reduced algal symbiont abundance in host tissues
and reduced photosynthetic efficiency, cell counts [**Figure 2A**] and Pulse Amplitude
Modulation (PAM) fluorometer measurements [**Figure 2B**] were taken for each of the
samples collected at each time point to characterize the effect of thermal stress on
endosymbiont physiology in the three coral species [**Supplemental Table S1a**]. Symbiont
cell densities, calculated as a factor of coral tissue area; hereinafter, endosymbiont
abundance, were used to assess bleaching severity and to normalize the proteomic data
generated from the symbionts in each of the sampled coral fragments. The latter was done to
account for reduced symbiont protein abundance values because of symbiont cell loss and not
reduced accumulation of the protein within the symbiont cells. Under ambient conditions,
endosymbiont abundances remained stable across the experiment [**Figure 2A**]. Using
ANOVA, we observed significant shifts in endosymbiont abundance upon exposure to
thermal stress in *A. hyacinthus* (df = 5, F = 6.032, p-value = 0.0051), but not in *P. lobata* (df
= 5, F = 1.015, p-value = 0.450) or *S. pistillata* (df = 3, F = 3.285, p-value = 0.079). In *A.*
hyacinthus, when comparing endosymbiont abundance between ambient conditions (TP1:
20.50 ± 0.32, TP2: 20.24 ± 0.52) and thermal stress samples (TP1: 19.23 ± 0.70, TP2: 18.36
± 0.31), Tukey-HSD post hoc analysis indicated a significant difference only at TP2 (adj-p-
value = 0.014).

Same as endosymbiont abundance, photosynthetic efficiency (F_v/F_m) values remained stable
under ambient conditions in all three species. However, under thermal stress, we observed
significant declines in endosymbiont photosynthetic efficiency in all three coral species
[**Figure 2B**]. ANOVA analysis showed significant photosynthetic efficiency shifts in *A.*
hyacinthus (df = 5, F = 8.552, p-value = 0.0011). Tukey-HSD post hoc analyses revealed a
significant decrease in thermal stress samples at TP2 (adj-p-value = 0.0007, Ambient = 0.740
± 0.037, Hot = 0.59 ± 0.027). Similarly, in *P. lobata* (ANOVA: df = 5.0, F = 3.067, p-value =
0.051), a significant decline was observed only at TP2 (adj-p-value = 0.020, Ambient = 0.712
± 0.018, Hot = 0.657 ± 0.015). Lastly, in *S. pistillata* (df = 5, F = 23.89, p-value = 0.00024),
we observed a significant decline at TP1 (adj-p-value = 0.0015, Ambient = 0.668 ± 0.032,
Hot = 0.529 ± 0.046).

We also characterized the endosymbiont community profiles using ITS2 sequencing [**Figure**
2C] to determine whether differences in host performance could be attributed to differences
in resident endosymbiont composition. ITS2-based *Symbiodiniaceae* profiling of the three
coral species using SymPortal showed that all samples associated exclusively with the genus
Cladocopium (except for one *P. lobata* replicate, which had <1% relative abundance of
Gerakladium) [**Figure 2C**]. The three coral species exhibited distinct major ITS2 type
profiles, which were C15-C15l-C15n-C15bb-C15.8 for all samples of *P. lobata*, C8/C1-
C42.2 for all samples of *S. pistillata*, and C3k/C3-C50a-C29-C21ab-C3b, C3/C3k-C50a-

C3.10-C3gj-C3b, and C3k-C50a-C3-C3ba-C50q-C3dq-C50f-C3a for all samples of *A.*
hyacinthus [**Supplemental Table 1b**].

Overall host proteome analysis

Prior to filtering, we obtained 3,793 protein groups (hereinafter proteins) from *A. hyacinthus*
[**Supplemental Table 2**], 4,098 proteins from *P. lobata* [**Supplemental Table 3**], and 2,479
proteins from *S. pistillata* [**Supplemental Table 4**]. After filtering for unique and razor
peptides and removing proteins with low counts, 2,255 (*A. hyacinthus*), 2,351 (*P. lobata*), and
1,335 (*S. pistillata*) proteins remained for downstream analysis. We assessed differential
abundance for all proteins using pairwise comparisons. A protein was considered to have
increased abundance if it had a $\log_2$ fold-change (FC) > 0.5 and *p*-value < 0.05, and decreased
abundance if it had a FC < –0.5 with *p*-value < 0.05.

Differentially abundant proteins were detected at TP0 in all three species, suggesting the
presence of tank effects. In *A. hyacinthus*, 72 proteins showed increased abundance and 5
showed decreased abundance [**Supplemental Figure 2**]. In *P. lobata*, 46 proteins had
increased abundance and 82 had decreased abundance. In *S. pistillata*, 5 proteins exhibited
increased abundance and 148 exhibited decreased abundance. Although these results
indicated tank effects at TP0, we observed a markedly higher number of differentially
abundant proteins at later timepoints under thermal stress. In *A. hyacinthus* at TP1, 23
proteins showed increased abundance and 161 showed decreased abundance; at TP2, 75
proteins showed increased abundance and 101 showed decreased abundance. In *P. lobata* at
TP1, 23 proteins showed increased abundance and 96 showed decreased abundance; at TP2,
86 proteins showed increased abundance and 235 showed decreased abundance
[**Supplemental Figure 2**]. Lastly, in *S. pistillata*, 87 proteins showed increased abundance
and 57 showed decreased abundance at TP1. Samples were collected only for two timepoints,
due to complete bailout by TP2 (**see Discussion**).

Pathway analysis of coral hosts

Enrichment analysis at the "B-level" (KEGG Orthology) was done to compare pathway-level
differences in coral stress responses. Pathways were considered upregulated if ≥5% of the
total proteins in a pathway showed increased abundance under stress, and downregulated if
≥5% showed a decrease. Mixed responses were assigned when both patterns were observed.
After filtering for pathways with more 10 proteins in each species, 25 pathways were retained
for analysis [**Figure 3A**]. At TP1, *A. hyacinthus* showed downregulation in 16 pathways; *P.*
lobata showed downregulation in 10 pathways and a mixed response in one; *S. pistillata*
showed upregulation in seven pathways, downregulation in three, and a mixed response in
three. By TP2, *A. hyacinthus* displayed a shift, with 5 pathways downregulated, six
upregulated, and one mixed. In contrast, *P. lobata* exhibited stronger downregulation at TP2,
with 16 pathways downregulated, five mixed, and one upregulated. Several pathways
downregulated in *P. lobata* at TP2 mirrored those observed in *A. hyacinthus* at TP1 (e.g.,
energy metabolism).

To identify shared vs. species-specific responses, we examined conserved patterns, defined as
a consistent direction of regulation across all species, and divergent responses (those not
conserved). Pathways were considered conserved even if their trends emerged at different
timepoints. For example, energy metabolism was downregulated at TP1 in both *S. pistillata*
and *A. hyacinthus*, but in *P. lobata*, this downregulation occurred only at TP2. This approach
accounts for differences in bleaching rates among the species, as evident from physiological

measurements. Only four pathways showed conserved response wherein carbohydrate
metabolism, energy metabolism, lipid metabolism were downregulated, and
folding/sorting/degradation was upregulated [**Figure 3A, red stars**]. In contrast, 21 pathways
displayed divergent responses (for example: cell growth and death) or species-specific
responses (for example: downregulation of "digestive system" in *P. lobata* only).

We prioritized conserved pathways for further functional analysis. Although "amino acid
metabolism" showed divergent trends, it was included due to its known role in thermal stress
in corals. Furthermore, two custom sets of proteins, symbiosome maintenance (lysosome,
phagosome proteins, carbonic anhydrases, and Vtype ATPases) and reactive oxygen species
(ROS) detoxification enzymes were assessed due to their known role in coral bleaching
(**Supplemental Table 5, See Supplemental Text**). Some proteins may be part of multiple
pathways but are presented only once in the results. **Figure 3B** summarizes the results
presented below.

244
Shared proteome responses in coral hosts.
Carbohydrate metabolism
All three coral species exhibited downregulation of carbohydrate metabolism under thermal
stress, with 39 proteins differentially abundant. However, only two proteins were shared
across species, although they were differentially abundant at different timepoints. Malonate-
semialdehyde dehydrogenase (ALDH6A1, K00140), involved in propionate metabolism,
showed decreased abundance at TP1 in both *A. hyacinthus* (FC = –0.52, *p*-value = 0.0086)
and *P. lobata* (FC = –0.72, *p*-value = 0.011), and at TP2 in *P. lobata* (FC = –0.77, *p*-value =
0.025). Aconitate hydratase (ACO, K01681) exhibited a divergent pattern, with increased
abundance in *S. pistillata* at TP1 (FC = 1.09, *p*-value = 0.032) and decreased abundance in *P.*
lobata at TP2 (FC = –0.58, *p*-value = 0.044). The remaining 37 proteins were species-specific
[Figure 3B].

Energy metabolism
Downregulation of energy metabolism was observed in all three species. Key proteins in
oxidative phosphorylation pathways, including ATPases, NADHs, and cytochrome proteins,
were differentially abundant. In *A. hyacinthus* at TP1, ATPeF1A (K02132, FC = –0.50, *p*-
value = 0.037) and ATPeF1B (K02133, FC = –0.53, *p*-value = 0.035) both showed decreased
abundance. Additionally, the V-type ATPase (VHA) genes ATPeV1G (K02152, FC = –1.57,
p-value = 0.027) and ATPeV1F (K02151, FC = –1.44, *p*-value = 0.031) also showed
decreased abundance at TP1. None of these proteins were statistically significant at TP2.
Different proteins related to energy metabolism were differentially abundant in *P. lobata*. At
TP1, NADH dehydrogenase subunit NDUFA5 (K03949, FC = –0.60, *p*-value = 0.035)
showed decreased abundance. A stronger trend was observed in the same pathway at TP2,
wherein ATPeFG (K02140, FC = –0.79, *p*-value = 0.042), NDUFS4 (K03937, FC = –0.67, *p*-
value = 0.010), and ATPeV1F (FC = –1.49, *p*-value = 0.041) all showed decreased
abundance in *P. lobata*. No significant shifts in oxidative phosphorylation proteins were
observed in *S. pistillata*.

Amino acid metabolism
Amino acids play an essential role in the coral-algal symbiosis and are catabolized by the
host under thermal stress. Glutamine synthetase (GLNA; K01915) showed decreased
abundance in *A. hyacinthus* at both TP1 (FC = –0.98, *p*-value = 0.028) and TP2 (FC = –1.09,
p-value = 0.020). In *S. pistillata*, two copies of GLNA were detected, both of which showed

decreased abundance at TP1 (FC = –0.64, *p*-value = 0.0043 and FC = –0.74, *p*-value =
0.00032). No shifts in GLNA abundance were observed in *P. lobata* at TP1, however, a
significant decrease was observed at TP2 (FC = –2.35, *p*-value = 0.0051). We also assessed
the abundance of glutamate dehydrogenase (GDH). One GDH protein (K00262) was detected
in *S. pistillata*, which showed a significant decrease at TP1 (FC = –0.50, *p*-value = 0.0029).
In *A. hyacinthus*, a different GDH protein (K00261) was identified that showed increased
abundance at TP2 (FC = 0.80, *p*-value = 0.016). In *P. lobata*, no significant shifts in GDH
abundance were detected. Lastly, we assessed 4-hydroxyphenylpyruvate dioxygenase
(HPPD), which showed a significant increase only in *P. lobata* at TP2 (K00457, FC = 0.68,
p-value = 0.028).

Protein folding, sorting, and degradation
Elevation of heat shock protein (HSP) abundance is one of most conserved responses in
corals under thermal stress (Molinari et al., 2025) and showed significant responses in all
three species. At TP1, three HSPs showed a significant increase in *S. pistillata* only: HSPA5
(K09490, FC = 1.16, *p*-value = 0.0013), HSP90B (K09487, FC = 0.96, *p*-value = 0.00057),
and HSP90A (K04079, FC = 0.78, *p*-value = 0.007). At TP2, significant increases were
observed in both *A. hyacinthus* and *P. lobata* for HSPA5 (FC = 0.60, *p*-value = 0.016; FC =
0.57, *p*-value = 0.028, respectively) and HSP90B (FC = 1.01, *p*-value = 0.00086; FC = 1.26,
p-value = 0.0022). Additionally, DNAJ proteins (HSP40 family) showed similar patterns: at
TP1, two proteins increased in *S. pistillata*, DNAJC3 (K09523, FC = 1.22, *p*-value =
0.00083) and DNAJA2 (K09503, FC = 2.05, *p*-value = 0.010). At TP2, *A. hyacinthus* also
showed increased abundance of DNAJC3 (FC = 1.07, *p*-value = 0.016). Another chaperone
protein, Calreticulin (CALR, K08057), showed a staggered yet consistent response. At TP1,
it showed increased abundance in *P. lobata* (FC = 0.71, *p*-value = 0.026) and *S. pistillata* (FC
= 0.89, *p*-value = 0.0065). At TP2, we observed a significant increase in both *A. hyacinthus*
(FC = 0.70, *p*-value = 0.0058) and *P. lobata* (FC = 1.02, *p*-value = 0.0081). We also observed
significant shifts in proteasome subunit proteins involved in degradation **[See Supplemental**
Text].

Lipid metabolism
Shifts in lipid content are known markers of coral thermal stress. We observed two key fatty
acid biosynthesis genes that showed conserved and significant responses to thermal stress:
long-chain-fatty-acid--CoA ligase (ACSBG, K15013) and long-chain acyl-CoA synthetase
(ACSL, K01897). At TP1, decreased abundance was observed in two ACSBG proteins in *A.*
hyacinthus (FC = –0.60, *p*-value = 0.014; FC = –0.86, *p*-value = 0.043). At TP1, *P. lobata*
also showed decreased abundance of ACSL (FC = –1.18, *p*-value = 0.019). At TP2, we
observed a similar pattern in *A. hyacinthus*, with two ACSBG proteins showing decreased
abundance (FC = –0.61, *p*-value = 0.033; FC = –1.02, *p*-value = 0.046), and ACSL also
decreasing (FC = –0.93, *p*-value = 0.029). In contrast, at TP2, *P. lobata* showed increased
abundance of ACSL (FC = 0.67, *p*-value = 0.000048).

Symbiosome maintenance proteins
We observed shifts in several key proteins potentially involved in symbiosomal maintenance.
Two cathepsin proteins: CTSL (K01365) and CTSB (K01363), associated with the lysosome,
were differentially abundant. At TP1, CTSB showed increased abundance in *A. hyacinthus*
(FC = 0.79, *p*-value = 0.043) and *S. pistillata* (FC = 1.02, *p*-value = 0.015). CTSL also
showed increased abundance in *A. hyacinthus* (FC = 1.59, *p*-value = 0.046) and *S. pistillata*
(FC = 0.58, *p*-value = 0.025). No significant response was observed in *P. lobata* at TP1. At
TP2, CTSB showed increased abundance in both *A. hyacinthus* (FC = 0.93, *p*-value = 0.028)

and *P. lobata* (FC = 1.51, *p*-value = 0.028). CTSL showed a mixed response in *P. lobata*,
with one copy increasing (FC = 1.50, *p*-value = 0.0015) and another decreasing (FC = –1.97,
p-value = 0.022).

Niemann-Pick C2 (NPC2, K13443), a lysosome-associated protein, showed decreased
abundance in *A. hyacinthus* at TP2 (FC = –2.04, *p*-value = 0.0069). Neutrophil cytosolic
factor 2 (NCF2), involved in phagosomes and ROS production, showed decreased abundance
at TP2 in both *A. hyacinthus* (FC = –0.88, *p*-value = 0.045) and *P. lobata* (FC = –1.63, *p*-
value = 0.012). Carbonic anhydrase 2 (CA2, K18245) showed decreased abundance in *A.
hyacinthus* at TP1 (FC = –2.26, *p*-value = 0.025) and TP2 (FC = –3.61, *p*-value = 0.0018).
Additionally, the transcription factor NF-κB (K02580) has been implicated to play a role in
maintaining symbiosis (Mansfield et al., 2017), which showed increased abundance in *A.
hyacinthus* at both TP1 (FC = 0.63, *p*-value = 0.025) and TP2 (FC = 1.43, *p*-value = 0.025).
No significant shifts were observed in the other species [for additional signaling proteins see
Supplemental Text]

Tank effects in the endosymbiont proteome
After quality filtering of endosymbiont protein groups, we obtained 332, 275, and 184
proteins from the endosymbionts of *A. hyacinthus*, *P. lobata*, and *S. pistillata*, respectively.
Protein abundances were subsequently normalized to account for changes in endosymbiont
abundance across samples, thereby minimizing artifacts caused by bleaching-induced cell
loss. PCA analysis, along with assessments of median protein intensities **[Supplemental File
2]**, revealed strong tank effects that led to proteomic shifts even among ambient samples,
independent of thermal stress. These confounding effects limited our ability to interpret
biological signal; thus, we did not pursue further analysis of the endosymbiont proteome.

Differences in metabolomic biomarkers across species
We investigated metabolomic data to validate previous findings regarding dipeptides, amino
acids, and antioxidants. The dipeptides KQ, RQ, RA and RV accumulate under thermal stress
(Williams et al., 2021a). In *A. hyacinthus* at TP1, KQ (FC = 3.40, adj-*p*-value = 0.0049), RQ
(FC = 2.30, adj-*p*-value = 0.026), and RV (FC = 1.86, adj-*p*-value = 0.0049) all significantly
increased in abundance. At TP2, the same dipeptides remained elevated. In *S. pistillata* at
TP1, only RV showed a significant increase (FC = 6.73, adj-*p*-value = 0.029), while the other
dipeptides also had increased abundances, they were not statistically significant. In *P. lobata*,
no dipeptides were significantly differentially abundant at either TP1 or TP2, though
increased fold-changes were observed at TP2 **[Figure 4B, Table S6a]**.

We assessed the abundance of well-characterized antioxidant metabolites (Williams et al.,
2021b). Each coral holobiont showed a distinct profile of stress metabolite accumulation
[Table S6b]. In *A. hyacinthus*, methionine had an increased abundance at both TP1 (FC =
0.74, adj-*p*-value = 0.046) and TP2 (FC = 2.34, adj-*p*-value = 0.018). Hypotaurine also had
an increased abundance at TP2 in *A. hyacinthus* (FC = 11.23, adj-*p*-value = 0.035). CDP-
choline also had an increased abundance in *A. hyacinthus* at TP1 (FC = 2.10, adj-*p*-value =
0.00030). However, these shifts were exclusive to *A. hyacinthus* only. Whereas
phosphocholine was the only antioxidant metabolite that had an increased abundance at TP1
in *S. pistillata* (FC = 1.93, adj-*p*-value = 0.054) and a significant increase *P. lobata* (FC =
1.03, adj-*p*-value = 0.027)

In addition, all three coral species showed increased abundance of amino acids **[Figure 4A,
Table S6c]**. In *A. hyacinthus*, nine amino acids increased at TP1 (seven were statistically

significant, adj-*p*-value < 0.05), and nine remained increased at TP2 (four were significant).
In *S. pistillata*, 12 amino acids had an increased abundance (seven were significant). In *P.*
lobata, seven amino acids had an increased abundance (four were significant). Of note only
two amino acids (arginine and alanine) had an increased abundance in all three species,
though this was temporally varied. Arginine had an increased abundance at TP1 in *S.*
pistillata (FC = 3.01, adj-*p*-value = 0.035) and subsequently at TP2 in *A. hyacinthus* (FC =
1.12, adj-*p*-value = 0.036) and *P. lobata* (FC = 2.78, adj-*p*-value = 0.028). Alanine had an
increased abundance at TP1 in *A. hyacinthus* (FC = 1.00, adj-*p*-value = 0.027) and *S.*
pistillata (FC = 2.81, adj-*p*-value = 0.035), and at TP2 in *P. lobata* (FC = 1.69, adj-*p*-value =
0.017).

379
Discussion

Overall trends in coral physiology

The three sympatric coral species investigated in this study (*Acropora hyacinthus*, *Porites*
lobata, and *Stylophora pistillata*) have well-documented differences in thermal stress
response. *A. hyacinthus* is thermally sensitive, *P. lobata* is thermally resilient, and *S. pistillata*
displays a low to moderate thermal stress tolerance (Burn et al., 2023; Fitt et al., 2009;
Manalili et al., 2025; Meziere et al., 2022; Pratchett et al., 2020). In the experimental tanks,
all three corals exhibited expected physiological responses to heat stress, including reduction
in endosymbiont abundance (Glynn, 1991; Jokiel and Coles, 1977) and photosynthetic
efficiency (F_v/F_m) (Warner et al., 1999, 1996). Consistent with the expected resilience levels,
A. hyacinthus bleached first, wherein we observed a reduced endosymbiont abundance upon
initial (TP1) stress and subsequently a more significant loss under prolonged (TP2) stress.
Decline in photosynthetic efficiency was also observed. In contrast, *P. lobata* had only a
slight decline (non-significant) in endosymbiont abundance, however, a significant decline in
photosynthetic efficiency occurred under prolonged stress. Lastly, in *S. pistillata* colonies
exhibited polyp bail-out, a rapid tissue loss response that is a severe stress reaction in this
species (Chuang et al., 2021; Sammarco, 1982; Schweinsberg et al., 2021). Reduced
photosynthetic efficiency was observed at initial stress. All three species hosted
endosymbiotic algae from the genus *Cladocopium*, however the strains in each species were
different. The three corals were collected from the same reef in the GBR, and likely had
similar historical environmental stress profiles, helping to minimize variability due to
differences in past stress exposure (Barshis et al., 2013; Kenkel and Matz, 2016; Morgan et
al., 2017; Oliver and Palumbi, 2011; Risk and Edinger, 2011; Rogers et al., 2016; Safaie et
al., 2018).

We assessed both the endosymbiont and host proteomes and observed tank effects in each. In
the coral hosts, these effects were minimal, with only a few proteins showing differential
abundance at TP0 (**Supplemental File 1**). In contrast, the endosymbiont proteome exhibited
more pronounced variability. Endosymbiont protein abundances were normalized to account
for bleaching-induced endosymbiont cell loss, however, this adjustment revealed exacerbated
tank effects. Specifically, we observed a decline in protein abundance in ambient samples
over time (**Supplemental File 2**), despite the absence of thermal stress. As a result, we
excluded the endosymbiont proteome from further analysis and focused instead on host-
specific responses to heat stress. Physiological indicators, including reductions in
endosymbiont abundance and photosynthetic efficiency, supported the occurrence of
bleaching. Together, these findings suggest a decline in photosynthate transfer from

endosymbionts to the host under thermal stress, a response observed in previous coral
bleaching studies (Davy et al., 2012; Grottoli et al., 2006).

417
Pathway and protein level responses are consistent with diverged ecological outcomes to
thermal stress.
Corals depend on their algal symbionts to meet metabolic needs, thus, reduction in symbiont-
derived photosynthate can severely impact holobiont fitness (Gordon and Leggat, 2010;
Tremblay et al., 2012). Our data reveal that different coral species exhibit distinct proteomic
responses to thermal stress and the associated decline in photosynthate transfer (**Figure 5**).
However, the timing of these protein shifts was different between the species, likely due to
differences in bleaching sensitivity, i.e., susceptible versus resilient. *A. hyacinthus*
demonstrated early and extensive proteomic shifts, consistent with its high thermal
sensitivity. At initial thermal stress, this species showed dramatic decreases in protein
abundance across a substantial portion of its proteome (34.6% of proteins based on fold-
change; **Supplemental Figure S3**). Upon prolonged thermal stress, there were signs of
proteomic recovery. *P. lobata* displayed a more gradual and delayed response, wherein under
initial stress a moderate response was observed. However, with prolonged stress, the response
became more prominent, resembling the pattern seen in *A. hyacinthus*, but occurring later.
Notably, many proteins in *P. lobata* showed mixed responses across different copies,
suggesting complex pathway regulation. *S. pistillata* exhibited a unique response pattern,
showing both increases and decreases in protein abundances reflecting a mixed but active
stress-response strategy, before ultimately undergoing polyp bail-out

437
Consistent with their known ecological differences in thermal tolerance, we observed
divergent responses at the pathway level among the three coral species. We evaluated the
differential abundance of proteins across 25 KEGG pathways, 21 of which exhibited species-
specific patterns. Among these, the "cell growth and death" pathway, which includes
apoptosis and has been previously linked to coral thermal stress responses (Ainsworth et al.,
2011), showed the most pronounced divergence and thus serves as a prime example [**Figure**
3A]. In *A. hyacinthus*, this pathway was downregulated during initial stress but upregulated
under prolonged stress. In contrast, *P. lobata* exhibited a mixed initial response that shifted to
downregulation under longer exposure to stress. *S. pistillata* also showed a mixed response at
initial stress. These contrasting trajectories highlight the distinct stress-response strategies in
these species and align with their known phenotypes. Such divergent responses underscore
the challenges in identifying universal biomarkers of coral stress [**see below**].

Consistent with the known thermal sensitivity of species in the genus *Acropora* (Sakai et al.,
2019; Singh et al., 2019), we observed that in *A. hyacinthus*, 16 pathways were
downregulated under initial thermal stress, indicative of a broad metabolic collapse [**Figure**
5]. The host appeared to enter an energy-depleted state earlier than the other two coral
species, putatively suggesting a higher reliance on endosymbiont derived photosynthates. We
hypothesize that in response to reduced photosynthate transfer, the host shifts energy away
from endosymbiont maintenance, reflected by decreased abundance of carbon supply proteins
(e.g., CA2) (Bertucci et al., 2013; Zoccola et al., 2015) and increased abundance of proteins
associated with symbiont expulsion (e.g., NF-κB) (Mansfield et al., 2017). These trends were
reflected at the physiological level, wherein *A. hyacinthus* was the first of the three species to
exhibit substantial endosymbiont loss. Under prolonged stress, there was an apparent
recovery in the proteome, suggesting a compensatory shift towards mobilizing endogenous

carbohydrate, lipid, and amino acid reserves to sustain metabolic function following
bleaching (Rodrigues and Grottoli, 2007).

In *P. lobata*, a species known for its relative resilience to thermal stress (Huang et al., 2025;
Rivera et al., 2022), few pathways were affected during initial stress. This finding was
consistent with physiology, whereby no significant loss of endosymbiont cells was observed,
and reduced photosynthetic efficiency was noted only under prolonged stress. Furthermore,
under prolonged stress at an individual protein level, *P. lobata* had numerous instances of
mixed responses (i.e., one copy of a protein had increased abundance and another showed a
decrease). These mixed responses may indicate a compensatory mechanism in *P. lobata*
which may be responsible for the higher thermal tolerance in this genus. Species in the genus
Porites have larger nuclear genomes and subsequently encode more genes (30,000-60,000
genes) (Stephens et al., 2022), whereas other genera and species have smaller genomes and
gene inventories, such as *Acropora* (~20,000 genes) (Shinzato et al., 2021) and *Stylophora*
pistillata (24,833 genes) (Voolstra et al., 2017). We speculate that the larger genomic
repertoire in *Porites* species may enable a more complex and flexible stress response,
potentially underpinning their thermal resilience phenotype.

Physiologically, *S. pistillata* exhibited a bail-out response under thermal stress (Schweinsberg
et al., 2021). Unsurprisingly, at the molecular level we observed distinctive responses,
compared to the other two species. We observed many pathways upregulated at initial stress
(7 of the 25), which included transcription and translation, and were species-specific
responses in *S. pistillata*. A study that performed salinity induced bail-out in the coral
Pocillopora acuta found key enriched responses in signaling pathways (Chuang and Mitarai,
2020). Proteins in signaling pathways were relatively stable, however this may be due to the
measurements being made prior to bail-out and not during active bail-out or may simply
reflect the smaller number of proteins obtained from *S. pistillata* when compared to the other
two coral species. Shared with *P. acuta*, we found an increased abundance in proteosome
associated proteins that may be involved in proteolysis (PSMA3, PSMA4, and PSMD1)
(Chuang and Mitarai, 2020) [**see Supplemental Text**].

Diverged protein responses even in pathways with a conserved response
Only four of the 25 pathways exhibited a conserved response across all three coral species
[**Figure 3A**]. Within these, two key trends emerged. First, although the direction of change
was consistent across species (e.g., downregulation), the timing varied. For example, the
energy metabolism pathway was downregulated in *A. hyacinthus* and *S. pistillata* during
initial stress, but not until prolonged stress in *P. lobata* [**Figure 3A**]. These conserved, yet
temporally staggered, responses align with known bleaching susceptibility, and subsequently
endosymbiont photosynthate limitation. Declines in carbohydrate and energy metabolism
have been widely documented, although the magnitude and timing differ across species and
experimental conditions (Haydon et al., 2023; Petrou et al., 2021;). Lipid metabolism, a
commonly used biomarker of coral bleaching, was also consistently downregulated,
consistent with prior studies (Ermolenko and Sikorskaya, 2021; Grottoli et al., 2004; Imbs
and Dembitsky, 2023). In contrast, protein folding, sorting, and degradation pathways were
upregulated, which consist of heat-shock proteins, potentially reflecting increased demand for
the turnover and clearance of misfolded or damaged proteins under stress (Petrou et al., 2021;
Williams et al., 2021b).

The second trend we observed was that even within pathways exhibiting a conserved overall
response, though temporally varied, the specific proteins driving that response differed across

species. For instance, carbohydrate metabolism, which showed a consistent decrease across
all three corals only 2 of the 39 differentially abundant proteins were shared between species.
ALDH6A1 was differentially abundant in both *A. hyacinthus* and *P. lobata*, whereas ACO
was affected in *S. pistillata* and *P. lobata*, but showed divergent expression patterns. The
remaining 37 proteins were differentially abundant in only one species [**Figure 3B**].
513
Shared protein and metabolite responses under thermal stress
Conserved protein responses
In this section, we examined the individual proteins that exhibited a conserved response in
the three coral species. We first focused on canonical stress response proteins, commonly
observed in corals and other metazoans. Heat shock proteins (HSPs) are well-established
markers of thermal stress (Rosic et al., 2011; Sharp et al., 1997). These proteins have been
widely used as biomarkers and play a central role in promoting survival under elevated
temperatures in corals (Cleves et al., 2020; Louis et al., 2017). In our study, HSPA5 (also
known as HSP70) and HSP90B showed conserved upregulation across all species. Notably,
this response was temporally staggered wherein, *S. pistillata* exhibited increased abundance
at initial stress, whereas *A. hyacinthus* and *P. lobata* showed increases later in the stress
period. We also observed a conserved increase in calreticulin (CALR), a calcium-binding
endoplasmic reticulum chaperone involved in protein folding and quality control (Fucikova et
al., 2021). CALR has been proposed to act in concert with HSPs during thermal stress in
corals (Ishibashi et al., 2023; Ruiz-Jones and Palumbi, 2017), supporting its role as a
component of the conserved stress response machinery.
We also noted conserved proteins that are not directly involved in thermal stress responses
but have been implicated in previous studies. One such protein is glutamine synthetase
(GLNA), which is involved in ammonium assimilation. GLNA abundance decreased across
all three corals; GLNA has been previously observed to have a decreased abundance at the
transcriptomic level in *S. pistillata* from the Red Sea (Rädecker et al., 2021), suggesting a
conserved response across species and geographic locations. Lastly, we observed a conserved
response in two lysosomal cysteine proteases, cathepsin proteins CTSB and CTSL. Increased
expression of CTSB under copper and thermal stress has been observed in corals and is
suggested to play a role in the ROS response (Louis et al., 2017; Schwarz et al., 2013).
Increased CTSL expression has been previously associated with pathogenic infections and
thermal stress in corals (Louis et al., 2017; Skutnik et al., 2020; Wright et al., 2015).
541
Conserved metabolomic responses
In this study, we employed polar metabolomics to identify and validate previously proposed
metabolite biomarkers of thermal stress. Consistent with the physiological and proteomic
responses, a notable temporal variation was observed in the metabolomic data. Within the
coral holobiont, the host and endosymbiont share the same amino acid pool. Under thermal
stress, amino acids serve two key functions: first, to maintain symbiosis, corals that transfer
amino acids to their endosymbionts bleach more slowly than those that do not (Martinez et
al., 2022b, 2022a, 2020); and second, to support host metabolism upon symbiont loss, as
amino acid catabolism can compensate for reduced photosynthate supply (Rädecker et al.,
2021). In this study, we observed an increase in free amino acid abundance in all three coral
species, though the timing was species-specific. Only two amino acids, arginine and alanine,
were consistently elevated across all species. Whereas the role of the increased amino acid
pool remains unclear, species-specific physiological patterns suggest divergent strategies. In

A. hyacinthus, increased amino acids coincided with early endosymbiont loss, suggesting
they may supplement host metabolism under stress. Likewise, in *S. pistillata*, the increased
amino acid pool, may be used to serve as a source of nutrition post bail-out. In contrast, *P.*
lobata maintained stable endosymbiont abundances even under prolonged stress, implying
that the elevated amino acid pool may be transferred to the symbiont to sustain the symbiosis.

Dipeptides are markers of stress in other model systems (Williams et al., 2021a) and show
great promise in corals. Little is known however about the pathways involved in their
synthesis and downstream functions. Stable isotope analysis demonstrates large differences in
the turnover rates of dipeptides and free amino acids, suggesting that these have distinct
biosynthetic mechanisms in corals (Chiles et al., 2022). We observed a conserved increase in
dipeptide abundance across species, with all four dipeptides exhibited positive fold changes
under thermal stress. Although only KQ, RQ, and RV were significantly increased in *A.*
hyacinthus, and RV was significant in both *S. pistillata* and *P. lobata*. Like the patterns
observed in the proteome, dipeptide responses were temporally staggered, reflecting species-
specific dynamics.
570
Study Limitations
Despite careful experimental design, tank effects were clearly observed in the coral and algal
endosymbiont proteomes, which is a persistent issue in coral research (McLachlan et al.,
2020; Molinari et al., 2025). After normalization of the endosymbiont proteome using
changes in cell abundance, the magnitude of this effect became even more apparent
(**Supplemental File 1**), which disallowed use of the algal data. Notably, colonies maintained
under ambient, non-stress conditions still exhibited substantial shifts in endosymbiont
proteomes across all three coral species, as shown by changes in median protein abundance.
In contrast, tank effects on the coral host proteome were minimal. Whereas it is plausible that
alterations in the endosymbiont proteome would influence the coral host, PCA and median
protein abundance analyses suggest these downstream effects were not substantial
(**Supplemental File 2**). In addition, we obtained different coverage levels in the proteome,
wherein significantly fewer proteins were obtained in *S. pistillata* relative to the other
species.
585
Conclusion
In this study we subjected three sympatric coral species *A. hyacinthus* (sensitive), *P. lobata*
(resilient), and *S. pistillata* (bail-out) with known ecological divergent responses to thermal
stress. Consistent with the literature, we observed the expected physiological response for
each species (i.e., reduced endosymbiont loss and decreased photosynthetic efficiency) in all
three corals. However, we observe a clear diverged response in the proteome with only six
pathways having a consistent response in all three corals. Even within the six pathways, we
note that only a handful of proteins had a consistent response across species. We note that
dipeptides consistently show an increased accumulation in all three species upon exposure to
thermal stress, however, some species showed a stronger response than others, further
supporting the species-specific nature of these responses. Overall, in *A. hyacinthus* we
observe a sharp initial response to thermal stress followed by proteomic recovery, whereas in

P. lobata we observe a mixed response only after prolonged stress, and lastly, in *S. pistillata*
we observed an initial stress response which was followed by host bail-out.

600
601

Methods

Test species
A single large colony of *Acropora hyacinthus* (neat morphotype), *Stylophora pistillata*, and
Porites lobata were sampled from Davies Reef (–18°49′30″S, 147°38′42″E) on the central
GBR in February 2023 (Permit Number: G19:43148) (**Figure 1**). Only one genotype was
used to avoid the possibility of stochastic inter-colony variation in the stress proteome and
metabolome (Chille et al., 2024). These species were selected to represent different
morphologies and heat stress sensitivities: *Acropora hyacinthus* - sensitive, tabulate,
branching; *Stylophora pistillata* – low/moderate tolerance, branching; *Porites lobata* -
tolerant, massive.

Experimental design
Each colony was subdivided into 36 fragments (~7cm^2), mounted on experimental racks, and
allowed to recover for 3 weeks prior to the start of the experiment at the National Sea
Simulator (SeaSim) at the Australian Institute of Marine Science (AIMS) in outdoor flow
through tanks replicating Davies Reef conditions (28.0°C, ca. ~250 µmol photons m^{-2} s^{-1}).
Prior to heat stress treatments, fragments were randomly assigned across six 50 L
experimental flow through tanks (n = 18 per tank, 6 fragments per species), three for each of
the two experimental treatments and acclimated for seven days before initiating heat stress.
Fragments were exposed to two treatments: a temperature treatment of 32.4°C and a control
maintained at 28.38^{o}C (Maximum Monthly Mean at Davies Reef). The MMM is defined as
the maximum sea surface temperature (SST) of the 12 monthly mean values for the years
1985-1990 and 1993 (Heron et al., 2014) temperature ramping increased from 28.38°C to
32.4°C, over 5☐days (0.8°C/day) for treatment tanks. The heat stress experiment ran for a
further 4 days concluding after 9 days. Heat exchangers and precision mixers were used to
control tank water temperature within 0.1°C and programmed and monitored using the
Supervisory Control and Data Acquisition system (SCADA) at SeaSim. Experimental tanks
were immersed within temperature controlled exterior insulating jackets of seawater kept at
treatment temperatures (one jacket per treatment) and each tank fitted with a TC Direct PT-
100 temperature probe to maintain ramp rate and programmed temperatures. Filtered
seawater was continuously supplied to the experimental system at an inflow rate of
0.8L/minute and each tank was fitted with a circulator (Turbelle nanostream 6015 3.5 W,
Tunze Aquarientechnik Gmbh, Germany) for water movement. Light was on a 24 hour 12:12
light/dark cycle with a 2-hour sunrise/sunset ramp time (half-sine ramp) and maximum
photosynthetically active radiation (PAR) at 280 µmol/m^2/s supplied from SS Gen 1
Customised multichip LEDs (325 W, The National Sea Simulator, Townsville, Australia) and
measured continuously in each tank using a Li-192 sensor (UWQ8687, Li-COR, USA) and
Li-250Am meter (Li-COR, USA). Water temperature and PAR was stable at target levels
throughout the experiment.

642
Sampling plan
Corals were sampled at three time points (**Figure 1**): TP0, day 0, at the start of the
experiment (end of coral acclimation described above) but prior to temperature ramp-up;
TP1, day 5, at the end of temperature ramp-up; and TP2, day 9, four days after the end of

temperature ramp-up. At each timepoint, sampling included photographs of each fragment
[**Supplemental Figure 1**] with one fragment per species from each tank (n =
3/treatment/species) immediately snap frozen and kept at -80°C until processing for
metabolomics, proteomics, and algal endosymbiont cell counts and typing (see below). At
each timepoint, and additionally, on days 7 and 8, the maximal quantum yield (F_v/F_m) of
chlorophyll a fluorescence of all fragments was measured using a Diving-PAM (Waltz,
Germany; Fiberoptics outer diameter 8 mm; settings: measuring light intensity = 6, saturation
pulse intensity = 10, saturating width = 0.8s, damping = 2, gain = 4). Measurements were
made before dawn therefore, colonies were low light acclimated (Suggett et al., 2022). Each
fragment was measured three times, at separate parts of the fragment, and the average values
were recorded. Analysis of variance (ANOVA) was utilized to assess the difference in F_v/F_m
for each coral species. Additionally, Tukey-HSD test was performed to do a pairwise
comparison between ambient and thermal stress at each timepoint.
660
Endosymbiont counts
A small section of tissue (~1 cm^2) from each frozen fragment collected at each timepoint
(TP0, TP1, and TP2), was air-brushed in PBS buffer, separated (as above), and washed three
times in filtered seawater, these cells were used to calculate endosymbiont abundance. Cell
quantification was determined visually *via* a haemocytometer (Neubauer Haemocytometer,
Fisher Scientific, Loughborough, UK) with eight replicate quadrats counted for each sample.
Surface area measurements of each air-brushed coral skeleton were conducted using the
single wax dipping method (Veal et al., 2010). Eight cylindrical calibration standards were
measured using digital callipers with a precision of 0.001 mm to determine their geometric
surface areas. These were used to generate a standard curve relating wax mass to surface area.
Coral skeletons were weighed and then skeletons and calibration standards were immersed in
paraffin wax at 65 °C for 2 seconds, then removed and gently rotated to promote uniform
coating and initial air drying. Samples were subsequently left to dry for an additional 15
minutes at ambient temperature before being reweighed. The surface area of each skeleton
was then calculated based on the difference in mass before and after wax coating, using the
calibration curve.
677
Endosymbiont cell densities were subsequently normalized using surface area, i.e., cell
density / surface area (cells/ml/cm^2), hereinafter cell abundance. Normalized cell abundance
were log$_2$ transformed for visualization. Analysis of variance (ANOVA) was utilized to assess
the difference in normalized cell density for each coral species. Additionally, Tukey-HSD test
was performed to do a pairwise comparison between ambient and thermal stress at each
timepoint.
684
Endosymbiont typing
A small section of tissue (~2 cm^2) from each frozen fragment collected at TP0 was air-
brushed in PBS buffer for endosymbiont typing using amplicon sequencing of the ribosomal
Internal Transcribed Spacer 2 (ITS-2) region. The tissue slurry was centrifuged at 3000g for
10 minutes to separate the coral host and algal symbiont fractions. DNA was extracted from
the algal pellet using the Qiagen DNeasy Blood and Tissue kit (Qiagen, Germany) following
the "protocol: purification of total DNA from animal tissues" including a RNase A-based
RNA digestion procedure. Genomic DNA was sent to the Ramaciotti Centre for Genomics
(University of New South Wales, Sydney, Australia) and ITS-2 amplicons sequenced using
the sym_var_5.8s2/sym_var_rev primers on the NextSeq 1000 P1 platform with 2x300bp
paired-end sequencing (including PhiX spike-in) (Hume et al., 2018). ITS2 sequences were

submitted to the SymPortal analytical framework for quality control and ITS2 type profile
analysis (https://symportal.org) as described by Hume and coauthors (Hume et al., 2019).
698
Protein extraction
Proteins were extracted using a RIPA lysis buffer, composed of Tris-HCl (50mM), NaCl
(150mM), SDS (1%), and DI water. The buffer was chilled on ice until ready for use and one
cOmplete mini EDTA-free tablet was dissolved in 10 ml of lysis buffer immediately before
extraction. A total of 500μl of ice-cold RIPA lysis buffer was added to a tube filled with
0.5mm silica beads. Approximately 0.2-0.4 g of coral sample was clipped into each bead
tube. The samples were vortexed for at least five minutes, or until the skeleton was stripped
from the tissue. The samples were then incubated on ice for 30 minutes, before centrifuging
at 10,000 rcf for 10 minutes and extracting the supernatant. Protein quality was assessed
using a Qubit protein assay. Samples were stored in a freezer at -80˚C.
709
Proteomics analysis
Proteomic analysis was performed at the University of South Florida Proteomics Core. The
proteins were normalized before digestion using 25μg for each sample. After digestion, an
equal amount of sample (1000ng) was injected for each run. LC-MS analysis peptides were
characterized using a Thermo Q-exactive-HF-X mass spectrometer coupled to a Thermo Easy
nLC 1200. Samples separated at 300nl/min on an Acclaim PEPMAP 100 trap (75uM, 2CM,
c18 3um, 100A) and an Thermo easy spray column (75um, 25cm, c18, 100A) using a 180-
minute gradient with an initial starting condition of 2% B buffer (0.1% formic acid in 90%
Acetonitrile) and 98% A buffer (0.1% formic acid in water). Buffer B was increased to 28%
over 140 minutes, then up to 40% in an additional 10 minutes. High B (90%) was run for 15
minutes afterwards. The mass spectrometer was outfitted with a Thermo nanospray easy
source with the following parameters: Spray voltage: 2.00V, Capillary temperature: 300dC,
Funnel RF level=40. Parameters for data acquisition were as follows: for MS data the
resolution was 60,000 with an AGC target of $3e^6$ and a max IT time of 50 ms, the range was
set to 400-1600 m/z. MS/MS data was acquired with a resolution of 15,000, an AGC of $1e^4$,
max IT of 50 ms, and the top 30 peaks were picked with an isolation window of 1.6m/z with
a dynamic execution of 25s.
727
For each of the three species, a comprehensive metaproteomic database was assembled based
on the ITS2 results (**Figure 2C**) to aid protein identification. A proteomic database for
Porites lobata was assembled from: (1) predicted genes from the *P. lobata* transcriptome
assembly (v1) (Bhattacharya et al., 2016); (2) predicted genes from the *P. lobata* (KU572435)
mitochondrial genome assembly (Tisthammer et al., 2016); (3) predicted genes from the
Cladocopium goreaui nuclear genome assembly (Chen et al., 2022), (4) predicted genes from
the *Breviolum minutum* mitochondrial genome assembly (Shoguchi et al., 2015); and (5)
predicted genes from the *Cladocopium* C3 plastid genome assembly (Barbrook et al., 2014).
A proteomic database for *Acropora hyacinthus* was assembled from: (1) predicted genes from
the *A. hyacinthus* nuclear genome assembly (v1) (Singh et al., 2019); (2) predicted genes
from the *A. hyacinthus* (OP311657) mitochondrial genome assembly (Wang et al., 2022); (3)
predicted from the *Cladocopium goreaui* nuclear genome assembly; (4) predicted
genes from the *Breviolum minutum* mitochondrial genome assembly; and (5) predicted genes
from the *Cladocopium* C3 plastid genome assembly. Finally, a proteomic database for
Stylophora pistillata was assembled from: (1) predicted genes from the *S. pistillata* nuclear
genome assembly (v1) (Voolstra et al., 2017); (2) predicted genes from the *S. pistillata*
(EU400214) mitochondrial genome assembly; (3) predicted genes from the *Cladocopium*
goreaui nuclear genome assembly; (4) predicted genes from the *Breviolum minutum*

mitochondrial genome assembly; and (5) predicted genes from the *Cladocopium* C3 plastid
genome assembly.
748
Protein clean-up and normalization
Raw proteomic data was generated using Max quant 2.0.3.1 (Cox and Mann, 2008; Tyanova
et al., 2016). Proteomic data cleaning was conducted in RStudio v2024.04.2+764 using R
v4.4.1 and the protti R package v0.9.0 (Quast et al., 2022). This package was used to clean,
normalize, and generate pre- and post-cleaning reports for each of the three host species.
Assessments of peak intensity distribution, data completeness, coefficient of variation across
sites, and principal component analysis (PCA) were performed before and after cleaning. To
create a high-quality protein group library, LC-MS results were filtered to exclude symbiont
proteins, proteins with greater than one unique peptide and razor peptide for each protein
group (Unique peptides > 1 & Razor + unique peptides > 1) and a protein group Q-value less
than 0.01. Proteins belonging to a set of potential contaminants provided by the LC-MS
facility were also removed. The data matrix was then filtered to exclude observations with
intensity values below 5, which likely represent false assignments. Protein intensities were
log2-transformed using the dplyr *mutate* and base R *log2* functions. Outlier samples were
identified by PCA and data completeness using the protti *qc_pca* and *qc_data_completeness*
functions. Based on these analyses, two samples with low completeness and that were PCA
outliers in their respective datasets were removed, Acropora_638 and Stylophora_2312.
Protein groups missing from more than 80% of samples were also removed. This threshold
allowed proteins present in at least 80% of samples from a species to remain in the dataset,
therefore it had to have been missing in >3 samples to be removed. For the remaining high-
confidence observations, missing values were imputed using the *missForest* function from the
missForest package with default parameters (maxiter = 10, ntree = 100) and a set seed of 124.
Finally, log2-transformed peak intensities were median-normalized per treatment_timepoint
group using the protti *normalise* function, resulting in the final cleaned protein abundance
matrix.
774
Downstream proteomic analysis
GhostKOALA (v2.0) (Kanehisa et al., 2016) was used to annotate proteins with KEGG
Orthology (KO) numbers. We subsequently calculated the $\log_2$ Fold-Change (hereinafter, FC)
for each protein between the ambient and high temperature samples for TP0, TP1, and TP2
using Python (v.3.9.12) (additional packages pandas v1.4.2 and numpy v1.26.4), and using
SciPy (v1.12.0) to assess statistical significance with a student t-test ($n = 3$). A protein was
considered to have an increased abundance if it had a FC > 0.5 and p-value < 0.05, likewise,
a protein was considered to have a decreased abundance if it had a FC < -0.5 and p-value <
0.05. However, upon correcting for multiple testing (Benjamini Hochberg corrections), no
adjusted p – value was < 0.05, potentially due to the high variation in the proteomic data. We
therefore proceed with only p-value based analyses. We also performed a pathway level
analysis, using the "B-level Description" annotation generated from the KEGG database. A
pathway was upregulated if 5% of the total proteins in a pathway had an increased abundance
and was downregulated if 5% of the total proteins had a decreased abundance. A pathway was
considered to have a mixed response if both upregulation and downregulation was observed.
790
791 *Metabolomic analysis*
792 In addition to physiological and proteomic analyses, we generated untargeted polar
793 metabolomics data to assess metabolite abundance (notably dipeptides) previously linked to
794 thermal stress responses in corals (Williams et al., 2021a). Metabolites were extracted from
795 preserved coral tissue using a 40:40:20 methanol:acetonitrile:water buffer with 0.1 M formic

acid, followed by homogenization, centrifugation, acid neutralization with ammonium bicarbonate, and transfer to autosampler vials for LC-MS analysis. LC-MS analysis was performed at the Metabolomics Core Facility at the Cancer Institute of New Jersey. Prior to sample runs, LC-MS performance was verified using commercial and in-house standards to ensure mass accuracy and signal consistency. Samples were randomized to minimize batch effects, and method blanks were included to assess background signals. Metabolite separation was performed using HILIC on a Vanquish UHPLC (Thermo Fisher Scientific, Waltham, MA) and the subsequent full scan mass spectrometry was performed on the Thermo Q Exactive PLUS mass spectrometer operating in both positive and negative modes. Data were processed using MAVEN (Melamud et al., 2010) with identification based on accurate mass and retention time matching to an in-house library, and low-abundance metabolites were filtered from downstream analyses. For an in-depth description of the extraction protocols and mass-spectrometry settings **see supplemental text.**

We subsequently calculated the $\log_2FC$ for each of the metabolites across the species, comparing the ambient and thermally stressed treatments at each timepoint. We also assessed the statistical significance for each of the metabolites using a Welch t-test ($n = 6$) with multiple testing correction using the Benjamini-Hochberg procedure, using SciPy (v1.12.0).

Acknowledgments
We acknowledge the Bindal and Wulgurukaba Peoples as the Traditional Custodians of the land and sea Countries where the Great Barrier Reef part of this research took place. We wish to acknowledge Elders past, present, and emerging, and their continuing spiritual connection to sea Country. Thank you to the staff of the AIMS vessel and the National Sea Simulator for field and laboratory support. We also thank Elizabeth Wood at the University of South Florida, Proteomics Core for support with proteomic data collection and analysis. We thank Eric Chiles and Dr. Xiaoyang Su at the Rutgers Metabolomic Core. Finally, we wish to thank Eric Hochberg from the Bermuda Institute of Ocean Sciences for his contributions to the design of the study and assistance with collection of the samples.

Funding
National Science Foundation award titled "Collaborative Research: Edge CMT: Polygenic traits of heat stress phenome in coral "dark genes" from genome to functional applications" (#2128073; 12/01/2021 to 11/30/2025) awarded to DB, USDA National Institute of Food and Agriculture Hatch Formula award titled "Using multi-omics and experimental evolution to understand the response of eukaryotes to changing environmental conditions" (#NJ01180; 01/17/20 to 09/30/24) awarded to DB, Revive & Restore award titled "Toward predictive coral phenomics" (#2022-039; 01/10/2022 to 09/30/2025) awarded to LKB, SG, and DB, and the Australian Government Reef Trust and Great Barrier Reef Foundation titled "Reef Restoration and Adaptation Program (RRAP)" awarded to LKB.

Author contributions:
Conceptualization: SN, TGS, EEC, SG, LKB, DB
Methodology: SN, TGS, EEC, SG, LKB, DB
Investigation: SN, TGS, EEC, SG, LKB, DB
Formal Analysis: SN, TGS, EEC
Supervision: LKB, DB
Writing (original draft): SN, TGS
Writing (review & editing): SN, TGS, EEC, SKG, LB, DB

846 **Competing interests:** Authors declare that they have no competing interests.
847

Data and materials availability: All data are available in the main text or the supplementary
materials. The GBR coral mass spectrometry proteomics data have been deposited to the
ProteomeXchange Consortium *via* the PRIDE partner repository with the dataset identifier
PXD052478 (https://www.ebi.ac.uk/pride/). The metabolomic mass spectrometry data are
available from the MassIVE database with the dataset identifier MSV000094832
(https://massive.ucsd.edu/). The ITS2 sequence read files are available from the Sequencing
Read Archive (SRA) with the dataset identifier PRJNA1119670
(https://www.ncbi.nlm.nih.gov/sra/).

Figures

Figure 1. Study sites and experimental designs. Location of samples ($n = 48$) collected and
experimental design (right panel) are shown. The coral images were generated by Samantha
Goyen, map created in R using the Environmental Systems Research Institute (ESRI)
shapefiles and Mapdata package, experimental setup image made in Canva. TP refers to the
timepoint at which corals were sampled. Tank set up is displayed on the left with the mean
temperatures of each tank shown. Graph showing the ramp up protocol used for the tanks,
red, thermal stress tanks and blue, control tanks.

**Figure 2. Physiological measurements of GBR corals under ambient and heat stress
treatments.** (**A**) Endosymbiont cell abundance ($\log_2$ transformed) over the duration of the
experiment. Blue boxes represent ambient cell density at each timepoint (x-axis) and the
orange boxes represent cell density upon exposure to thermal stress. The three graphs are the
different species used in this study. The y-axis shows the $\log_2$ normalized cell density
(cells/ml/cm^2). (**B**) The y-axis shows the F_v/F_m ratio, which is a proxy for photosynthetic
efficiency, and the x-axis represents the experimental time points. Blue boxes are the values
for the ambient samples and the orange for the stress samples. * Represents statistical
significance (p-value < 0.05) between the ambient and heat stress treatments at that
timepoint. (**C**) Variation in major SymPortal ITS2 Symbiodiniaceae communities within
Acropora hyacinthus, *Porites lobata*, and *Stylophora pistillata* colonies from the Great
Barrier Reef. The y-axis is the relative abundance. One replicate of *P. lobata* consisted of
<1% relative abundance of *Gerakladium* and has not been included in the image.

**Figure 3. Proteomic responses at the pathway level and individual protein abundances
in conserved pathways.** (**A**) Heatmap showing the pathway level response (y-axis) for each
coral species (x-axis) at each timepoint. The blue boxes are downregulated pathways in a
species at that timepoint, the orange boxes show upregulated pathways, green boxes show a
mixed response, and grey boxes indicate no change (see figure legend). The red stars indicate
conserved pathways that are extensively discussed in the main text. (**B**) Heatmap showing
significant protein responses ($|FC| > 0.5$ and p-value < 0.05), with proteins on the y-axis
(Name and KEGG ID) and the species on the x-axis. The graph is divided in two timepoints,
TP1 left) and TP2 (right). Some proteins may be listed twice; this is due to the presence of
multiple copies in a species that are significant.

Figure 4. Differential abundance of polar metabolites. (**A**) Heatmap showing the various
targeted metabolites in the study (y-axis) and the coral species and timepoints (x-axis). Red
boxes indicate increased abundance, whereas blue boxes indicate decreased abundance. The

asterisks denote statistical significance (* adj-p-value < 0.05; ** adj-p-value < 0.01; *** adj-p-value < 0.001). (**B**) Box plots for the four dipeptides assessed in the metabolomic dataset. Abundances (y-axis) are plotted for each dipeptide at each timepoint (x-axis). The thermally stressed samples are shown in orange, whereas ambient samples are shown in blue. Each column shows a different coral species and each row, a different dipeptide.

Figure 5. Divergent outcomes of thermal stress in three GBR coral species. (**A**) Diagram depicting a healthy symbiosis between the host and algal endosymbiont, involving transfer of nutrients from the host to the endosymbiont and subsequent transfer of photosynthates from the endosymbiont to the host. (**B**) A less efficient endosymbiont upon exposure to stress, limiting photosynthate transfer to host. The blue cross represents the lack of transfer of photosynthates from the endosymbiont to the host. (**C**) The divergent host metabolic responses of the three studied GBR coral species. In each panel, blue arrows represent pathways with proteins that decrease in abundance in response to thermal stress. The combination of a blue and red arrow represents a mixed protein response: i.e., a similar number of proteins in a pathway had increasing (red) and decreasing (blue) abundance.

Supplemental Information

Table S1a.
Endosymbiont cell density and photosynthetic efficiency for samples from all three GBR coral species.
Table S1b.
SymPortal ITS2 Symbiodiniaceae communities within *Acropora hyacinthus, Porites lobata and Stylophora pistillata* from the Great Barrier Reef. ITS2 sequence data and major ITS2-type profiles.
Table S2.
Proteomic data for the *Acropora hyacinthus* host across timepoints, with differential accumulation statistics and KEGG annotations shown for each protein.
Table S3.
Proteomic data for the *Porites lobata* host across timepoints, with differential accumulation statistics and KEGG annotations shown for each protein.
Table S4.
Proteomic data for the *Stylophora pistillata* host across timepoints, with differential accumulation statistics and KEGG annotations shown for each protein.
Table S5.
Summary table of all GBR coral host proteins detected for each species across the stress experiment for specific pathways. Each protein is counted only once here for reference. Multiple copies of a given protein are included in these counts.
Table S6a.
Dipeptide metabolite abundance variation between control and treatment conditions at each time point from the GBR corals.
Table S6b.
Annotated antioxidant metabolite abundance variation between control and treatment conditions at each time point from the GBR corals.
Table S6c.
Amino acid metabolite abundance variation between control and treatment conditions at each time point from the GBR corals.

Supplemental File 1: Endosymbiont proteome distribution after cell density
normalization and principal component analysis.
Box and whisker plots of log2 protein intensities across treatments and timepoints for *A.*
hyacinthus (**A**), *P. lobata* (**C**), and *S. pistillata* (**E**) symbiont proteins. Protein intensities were
group-median normalized by treatment_timepoint for each species independently to examine
protein-wide up- or down- regulation in response to treatment. PCA plots of the first two
principal components based on group-normalized log2 protein intensities for *A.*
hyacinthus (**B**), *P. lobata* (**D**), and *S. pistillata* (**F**). Legends describing the colors and shapes
are shown to the right of each figure.
953
Supplemental File 2: Host protein intensity distribution and principal component
analysis.
Box and whisker plots of log2 protein intensities across treatments and timepoints for *A.*
hyacinthus (**A**), *P. lobata* (**C**), and *S. pistillata* (**E**) host proteins. Protein intensities were
group-median normalized by treatment_timepoint for each species independently to examine
protein-wide up- or down- regulation in response to treatment. PCA plots of the first two
principal components based on group-normalized log2 protein intensities for *A.*
hyacinthus (**B**), *P. lobata* (**D**), and *S. pistillata* (**F**). Legends describing the colors and shapes
are shown to the right of each figure.
963
Supplemental Figure 1: Coral fragment images over the course of the experiment.
This figure shows coral fragments from all three species at various timepoints during the
experiment. Timepoints and treatment conditions are indicated on the left, and fragment IDs
are shown below each image. In *Stylophora pistillata*, distinct paling of the host colonies is
visible as early as TP1 under thermal stress. Similarly, *Acropora hyacinthus* displays clear
signs of bleaching by TP2 based on visual assessment. In *Porites lobata*, mild paling is also
evident by TP2.
971
Supplemental Figure 2: Host total proteome distribution. Volcano plot of the proteins
identified in the host proteome data from the three sympatric species. The plots are arranged
horizontally by species and vertically by time point. Data is not available for *S. pistillata* TP2
due to the bail out phenotype, and as such, the respective position are filled with "NA". The
dots in each plot represent individual proteins, with the blue dots representing proteins with a
statistically significant decreased abundance (FC < -0.5, *p*-value < 0.05) and red dots
representing proteins with a statistically significant increased abundance (FC > 0.5, *p*-value <
0.05) at the given time point. A legend describing the colors is shown at the bottom of the
figure.
981
Supplemental Figure 3: Fold-change based pathway level assessment. (A) Stacked mirror
plot of differentially abundant proteins at TP1 in each of the three colonies. "B-Description"
(KEGG classification) is presented on the x-axis and the number of proteins that are
differentially abundant on the y-axis. Note only pathways with more than 5 proteins
differentially abundant (across all species) have been included in the plot. The different
species have been shown in different colors. Bars extending up are the number of proteins
that have an increased abundance and down show decreased abundance. **(B)** Similar to **(A)**
but shows differentially abundant proteins at TP2. Data is only available for *A. hyacinthus*
and *P. lobata*.
991
992 # References

Barbrook, A.C., Voolstra, C.R., Howe, C.J., 2014. The Chloroplast Genome of a Symbiodinium sp. Clade C3 Isolate. Protist 165, 1–13. https://doi.org/10.1016/j.protis.2013.09.006

Barkley, H.C., Cohen, A.L., Mollica, N.R., Brainard, R.E., Rivera, H.E., DeCarlo, T.M., Lohmann, G.P., Drenkard, E.J., Alpert, A.E., Young, C.W., Vargas-Ángel, B., Lino, K.C., Oliver, T.A., Pietro, K.R., Luu, V.H., 2018. Repeat bleaching of a central Pacific coral reef over the past six decades (1960–2016). Commun Biol 1, 177. https://doi.org/10.1038/s42003-018-0183-7

Barott, K.L., Thies, A.B., Tresguerres, M., 2022. V-type H^+-ATPase in the symbiosome membrane is a conserved mechanism for host control of photosynthesis in anthozoan photosymbioses. R. Soc. open sci. 9, 211449. https://doi.org/10.1098/rsos.211449

Barott, K.L., Venn, A.A., Perez, S.O., Tambutté, S., Tresguerres, M., 2015. Coral host cells acidify symbiotic algal microenvironment to promote photosynthesis. Proc. Natl. Acad. Sci. U.S.A. 112, 607–612. https://doi.org/10.1073/pnas.1413483112

Barshis, D.J., Ladner, J.T., Oliver, T.A., Seneca, F.O., Traylor-Knowles, N., Palumbi, S.R., 2013. Genomic basis for coral resilience to climate change. PNAS 110, 1387–1392. https://doi.org/10.1073/pnas.1210224110

Bertucci, A., Moya, A., Tambutté, S., Allemand, D., Supuran, C.T., Zoccola, D., 2013. Carbonic anhydrases in anthozoan corals—A review. Bioorganic & Medicinal Chemistry 21, 1437–1450. https://doi.org/10.1016/j.bmc.2012.10.024

Bhattacharya, D., Agrawal, S., Aranda, M., Baumgarten, S., Belcaid, M., Drake, J.L., Erwin, D., Foret, S., Gates, R.D., Gruber, D.F., Kamel, B., Lesser, M.P., Levy, O., Liew, Y.J., MacManes, M., Mass, T., Medina, M., Mehr, S., Meyer, E., Price, D.C., Putnam, H.M., Qiu, H., Shinzato, C., Shoguchi, E., Stokes, A.J., Tambutté, S., Tchernov, D., Voolstra, C.R., Wagner, N., Walker, C.W., Weber, A.P., Weis, V., Zelzion, E., Zoccola, D., Falkowski, P.G., 2016. Comparative genomics explains the evolutionary success of reef-forming corals. eLife 5, e13288. https://doi.org/10.7554/eLife.13288

Bhattacharya, D., Stephens, T.G., Chille, E.E., Benites, L.F., Chan, C.X., 2024. Facultative lifestyle drives diversity of coral algal symbionts. Trends in Ecology & Evolution 39, 239–247. https://doi.org/10.1016/j.tree.2023.10.005

Bourne, D.G., Morrow, K.M., Webster, N.S., 2016. Insights into the Coral Microbiome: Underpinning the Health and Resilience of Reef Ecosystems. Annual Review of Microbiology 70, 317–340. https://doi.org/10.1146/annurev-micro-102215-095440

Buerger, P., Alvarez-Roa, C., Coppin, C.W., Pearce, S.L., Chakravarti, L.J., Oakeshott, J.G., Edwards, O.R., van Oppen, M.J.H., 2020. Heat-evolved microalgal symbionts increase coral bleaching tolerance. Science Advances 6, eaba2498. https://doi.org/10.1126/sciadv.aba2498

Burn, D., Hoey, A.S., Matthews, S., Harrison, H.B., Pratchett, M.S., 2023. Differential bleaching susceptibility among coral taxa and colony sizes, relative to bleaching severity across Australia's Great Barrier Reef and Coral Sea Marine Parks. Marine Pollution Bulletin 191, 114907. https://doi.org/10.1016/j.marpolbul.2023.114907

Chen, Y., Shah, S., Dougan, K.E., van Oppen, M.J.H., Bhattacharya, D., Chan, C.X., 2022. Improved Cladocopium goreaui Genome Assembly Reveals Features of a Facultative Coral Symbiont and the Complex Evolutionary History of Dinoflagellate Genes. Microorganisms 10, 1662. https://doi.org/10.3390/microorganisms10081662

Chiles, E.N., Huffmyer, A.S., Drury, C., Putnam, H.M., Bhattacharya, D., Su, X., 2022. Stable isotope tracing reveals compartmentalized nitrogen assimilation in scleractinian corals. Frontiers in Marine Science 9.

Chille, E.E., Stephens, T.G., Misri, D., Strand, E.L., Putnam, H.M., Bhattacharya, D., 2024. Gene expression response under thermal stress in two Hawaiian corals is dominated by ploidy and genotype. Ecology and Evolution 14, e70037. https://doi.org/10.1002/ece3.70037

Chille, E.E., Stephens, T.G., Nandi, S., Jiang, H., Gerdes, M.J., Williamson, O.M., Neufeld, A., Montoya-Maya, P., Bhattacharya, D., 2025. Coral Restoration in the Omics Era: Development of Point-of-Care Tools for Monitoring Disease, Reproduction, and Thermal Stress. BioEssays 47, e70007. https://doi.org/10.1002/bies.70007

Chuang, P.-S., Ishikawa, K., Mitarai, S., 2021. Morphological and Genetic Recovery of Coral Polyps After Bail-Out. Front. Mar. Sci. 8. https://doi.org/10.3389/fmars.2021.609287

Chuang, P.-S., Mitarai, S., 2020. Signaling pathways in the coral polyp bail-out response. Coral Reefs 39, 1535–1548. https://doi.org/10.1007/s00338-020-01983-x

Cleves, P.A., Tinoco, A.I., Bradford, J., Perrin, D., Bay, L.K., Pringle, J.R., 2020. Reduced thermal tolerance in a coral carrying CRISPR-induced mutations in the gene for a heat-shock transcription factor. Proceedings of the National Academy of Sciences 117, 28899–28905. https://doi.org/10.1073/pnas.1920779117

Cowen, L.J., Putnam, H.M., 2022. Bioinformatics of Corals: Investigating Heterogeneous Omics Data from Coral Holobionts for Insight into Reef Health and Resilience. Annual Review of Biomedical Data Science 5, 205–231. https://doi.org/10.1146/annurev-biodatasci-122120-030732

Cox, J., Mann, M., 2008. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol 26, 1367–1372. https://doi.org/10.1038/nbt.1511

Cziesielski, M.J., Schmidt□Roach, S., Aranda, M., 2019. The past, present, and future of coral heat stress studies. Ecol Evol 9, 10055–10066. https://doi.org/10.1002/ece3.5576

Da-Anoy, J., Posadas, N., Conaco, C., 2024. Interspecies differences in the transcriptome response of corals to acute heat stress. PeerJ 12, e18627. https://doi.org/10.7717/peerj.18627

Davy, S.K., Allemand, D., Weis, V.M., 2012. Cell Biology of Cnidarian-Dinoflagellate Symbiosis. Microbiology and Molecular Biology Reviews 76, 229–261. https://doi.org/10.1128/mmbr.05014-11

DeSalvo, M.K., Voolstra, C.R., Sunagawa, S., Schwarz, J.A., Stillman, J.H., Coffroth, M.A., Szmant, A.M., Medina, M., 2008. Differential gene expression during thermal stress and bleaching in the Caribbean coral Montastraea faveolata. Mol Ecol 17, 3952–3971. https://doi.org/10.1111/j.1365-294X.2008.03879.x

Downs, C., Fauth, J., Halas, J., Dustan, P., Bemiss, J., Woodley, C., 2002. Oxidative stress and seasonal coral bleaching. Free radical biology & medicine 33, 533–43. https://doi.org/10.1016/S0891-5849(02)00907-3

Downs, C.A., Fauth, J.E., Robinson, C.E., Curry, R., Lanzendorf, B., Halas, J.C., Halas, J., Woodley, C.M., 2005. Cellular diagnostics and coral health: Declining coral health in the Florida Keys. Marine Pollution Bulletin, Coral Reef Ecotoxicology and Health 51, 558–569. https://doi.org/10.1016/j.marpolbul.2005.04.017

Ermolenko, E.V., Sikorskaya, T.V., 2021. Lipidome of the reef-building coral *Acropora cerealis:* Changes under thermal stress. Biochemical Systematics and Ecology 97, 104276. https://doi.org/10.1016/j.bse.2021.104276

Fiehn, O., 2002. Metabolomics--the link between genotypes and phenotypes. Plant Mol Biol 48, 155–171.

Fitt, W.K., Gates, R.D., Hoegh-Guldberg, O., Bythell, J.C., Jatkar, A., Grottoli, A.G., Gomez, M., Fisher, P., Lajuenesse, T.C., Pantos, O., Iglesias-Prieto, R., Franklin, D.J., Rodrigues, L.J., Torregiani, J.M., van Woesik, R., Lesser, M.P., 2009. Response of two species of Indo-Pacific corals, Porites cylindrica and Stylophora pistillata, to short-term thermal stress: The host does matter in determining the tolerance of corals to bleaching. Journal of Experimental Marine Biology and Ecology 373, 102–110. https://doi.org/10.1016/j.jembe.2009.03.011

Frankowiak, K., Wang, X.T., Sigman, D.M., Gothmann, A.M., Kitahara, M.V., Mazur, M., Meibom, A., Stolarski, J., 2016. Photosymbiosis and the expansion of shallow-water corals. Science Advances 2, e1601122. https://doi.org/10.1126/sciadv.1601122

Fucikova, J., Spisek, R., Kroemer, G., Galluzzi, L., 2021. Calreticulin and cancer. Cell Res 31, 5–16. https://doi.org/10.1038/s41422-020-
0383-9

Glynn, P.W., 1991. Coral reef bleaching in the 1980s and possible connections with global warming. Trends in Ecology & Evolution 6, 175–
179. https://doi.org/10.1016/0169-5347(91)90208-F

Gordon, B.R., Leggat, W., 2010. Symbiodinium—Invertebrate Symbioses and the Role of Metabolomics. Marine Drugs 8, 2546–2568.
https://doi.org/10.3390/md8102546

Greenbaum, D., Colangelo, C., Williams, K., Gerstein, M., 2003. Comparing protein abundance and mRNA expression levels on a genomic
scale. Genome Biology 4, 117. https://doi.org/10.1186/gb-2003-4-9-117

Grottoli, A.G., Rodrigues, L.J., Juarez, C., 2004. Lipids and stable carbon isotopes in two species of Hawaiian corals, Porites compressa and
Montipora verrucosa, following a bleaching event. Marine Biology 145, 621–631. https://doi.org/10.1007/s00227-004-1337-3

Grottoli, A.G., Rodrigues, L.J., Palardy, J.E., 2006. Heterotrophic plasticity and resilience in bleached corals. Nature 440, 1186–1189.
https://doi.org/10.1038/nature04565

Haydon, T.D., Matthews, J.L., Seymour, J.R., Raina, J.-B., Seymour, J.E., Chartrand, K., Camp, E.F., Suggett, D.J., 2023. Metabolomic
signatures of corals thriving across extreme reef habitats reveal strategies of heat stress tolerance. Proc. R. Soc. B. 290,
20221877. https://doi.org/10.1098/rspb.2022.1877

Heron, S.F., Liu, G., Eakin, C.M. (Carlon M., Skirving, W.J. (William J., Muller-Karger, F.E., Vega-Rodriguez, M., De La Cour, J.L.,
Burgess, T.F.R., Strong, A.E. (Alan E., Geiger, E.F., Guild, L.S., Lynds, S., 2014. Climatology development for NOAA Coral
Reef Watch's 5-km product suite. https://doi.org/10.7289/V59C6VBS

Huang, W., Chen, J., Yang, E., Meng, L., Feng, Y., Chen, Y., Huang, Z., Tan, R., Xiao, Z., Zhou, Y., Xu, M., Yu, K., 2025. Heat-tolerant
subtropical *Porites lutea* may be better adapted to future climate change than tropical one in the South China Sea. Science of The
Total Environment 962, 178381. https://doi.org/10.1016/j.scitotenv.2025.178381

Huffmyer, A.S., Ashey, J., Strand, E., Chiles, E.N., Su, X., Putnam, H.M., 2024. Coral larvae increase nitrogen assimilation to stabilize algal
symbiosis and combat bleaching under increased temperature. PLOS Biology 22, e3002875.
https://doi.org/10.1371/journal.pbio.3002875

Hughes, T.P., Anderson, K.D., Connolly, S.R., Heron, S.F., Kerry, J.T., Lough, J.M., Baird, A.H., Baum, J.K., Berumen, M.L., Bridge, T.C.,
Claar, D.C., Eakin, C.M., Gilmour, J.P., Graham, N.A.J., Harrison, H., Hobbs, J.-P.A., Hoey, A.S., Hoogenboom, M., Lowe, R.J.,
McCulloch, M.T., Pandolfi, J.M., Pratchett, M., Schoepf, V., Torda, G., Wilson, S.K., 2018. Spatial and temporal patterns of mass
bleaching of corals in the Anthropocene. Science 359, 80–83. https://doi.org/10.1126/science.aan8048

Hughes, T.P., Kerry, J.T., Álvarez-Noriega, M., Álvarez-Romero, J.G., Anderson, K.D., Baird, A.H., Babcock, R.C., Beger, M., Bellwood,
D.R., Berkelmans, R., Bridge, T.C., Butler, I.R., Byrne, M., Cantin, N.E., Comeau, S., Connolly, S.R., Cumming, G.S., Dalton,
S.J., Diaz-Pulido, G., Eakin, C.M., Figueira, W.F., Gilmour, J.P., Harrison, H.B., Heron, S.F., Hoey, A.S., Hobbs, J.-P.A.,
Hoogenboom, M.O., Kennedy, E.V., Kuo, C., Lough, J.M., Lowe, R.J., Liu, G., McCulloch, M.T., Malcolm, H.A., McWilliam,
M.J., Pandolfi, J.M., Pears, R.J., Pratchett, M.S., Schoepf, V., Simpson, T., Skirving, W.J., Sommer, B., Torda, G., Wachenfeld,
D.R., Willis, B.L., Wilson, S.K., 2017. Global warming and recurrent mass bleaching of corals. Nature 543, 373–377.
https://doi.org/10.1038/nature21707

Hughes, T.P., Kerry, J.T., Connolly, S.R., Baird, A.H., Eakin, C.M., Heron, S.F., Hoey, A.S., Hoogenboom, M.O., Jacobson, M., Liu, G.,
Pratchett, M.S., Skirving, W., Torda, G., 2019. Ecological memory modifies the cumulative impact of recurrent climate extremes.
Nature Clim Change 9, 40–43. https://doi.org/10.1038/s41558-018-0351-2

Hume, B.C.C., Smith, E.G., Ziegler, M., Warrington, H.J.M., Burt, J.A., LaJeunesse, T.C., Wiedenmann, J., Voolstra, C.R., 2019. SymPortal:
A novel analytical framework and platform for coral algal symbiont next-generation sequencing ITS2 profiling. Molecular
Ecology Resources 19, 1063–1080. https://doi.org/10.1111/1755-0998.13004

Hume, B.C.C., Ziegler, M., Poulain, J., Pochon, X., Romac, S., Boissin, E., de Vargas, C., Planes, S., Wincker, P., Voolstra, C.R., 2018. An
improved primer set and amplification protocol with increased specificity and sensitivity targeting the Symbiodinium ITS2
region. PeerJ 6, e4816. https://doi.org/10.7717/peerj.4816

Imbs, A.B., Dembitsky, V.M., 2023. Coral Lipids. Mar Drugs 21, 539. https://doi.org/10.3390/md21100539

Ip, J.C.-H., Zhang, Y., Xie, J.Y., Yeung, Y.H., Qiu, J.-W., 2022. Comparative transcriptomics of two coral holobionts collected during the
2017 El Niño heat wave reveal differential stress response mechanisms. Marine Pollution Bulletin 182, 114017.
https://doi.org/10.1016/j.marpolbul.2022.114017

Ishibashi, H., Minamide, S., Takeuchi, I., 2023. Expression analyses of stress-responsive genes in the hermatypic coral Acropora tenuis and
its symbiotic dinoflagellates after exposure to the herbicide Diuron. Mar Life Sci Technol 5, 289–299.
https://doi.org/10.1007/s42995-023-00183-0

Jokiel, P.L., Coles, S.L., 1977. Effects of temperature on the mortality and growth of Hawaiian reef corals. Mar. Biol. 43, 201–208.
https://doi.org/10.1007/BF00402312

Kanehisa, M., Sato, Y., Morishima, K., 2016. BlastKOALA and GhostKOALA: KEGG Tools for Functional Characterization of Genome
and Metagenome Sequences. J Mol Biol 428, 726–731. https://doi.org/10.1016/j.jmb.2015.11.006

Kenkel, C.D., Matz, M.V., 2016. Gene expression plasticity as a mechanism of coral adaptation to a variable environment. Nat Ecol Evol 1,
0014. https://doi.org/10.1038/s41559-016-0014

Kenkel, C.D., Meyer, E., Matz, M.V., 2013. Gene expression under chronic heat stress in populations of the mustard hill coral (Porites
astreoides) from different thermal environments. Molecular Ecology 22, 4322–4334. https://doi.org/10.1111/mec.12390

Krueger, T., Hawkins, T.D., Becker, S., Pontasch, S., Dove, S., Hoegh-Guldberg, O., Leggat, W., Fisher, P.L., Davy, S.K., 2015. Differential
coral bleaching—Contrasting the activity and response of enzymatic antioxidants in symbiotic partners under thermal stress.
Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology 190, 15–25.
https://doi.org/10.1016/j.cbpa.2015.08.012

Louis, Y.D., Bhagooli, R., Kenkel, C.D., Baker, A.C., Dyall, S.D., 2017. Gene expression biomarkers of heat stress in scleractinian corals:
Promises and limitations. Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology 191, 63–77.
https://doi.org/10.1016/j.cbpc.2016.08.007

Loya, Y., Sakai, K., Yamazato, K., Nakano, Y., Sambali, H., van Woesik, R., 2001. Coral bleaching: the winners and the losers. Ecology
Letters 4, 122–131. https://doi.org/10.1046/j.1461-0248.2001.00203.x

Manalili, S.E.N., Bataan, D.A.U., Ulanova, D., Sakurai, T., Sekida, S., Keshavmurthy, S., Mezaki, T., Kubota, S., 2025. Short-term recovery
responses in Acropora hyacinthus exposed to moderate-term thermal stress. Mar Biol 172, 76. https://doi.org/10.1007/s00227-
025-04636-1

Mansfield, K.M., Carter, N.M., Nguyen, L., Cleves, P.A., Alshanbayeva, A., Williams, L.M., Crowder, C., Penvose, A.R., Finnerty, J.R.,
Weis, V.M., Siggers, T.W., Gilmore, T.D., 2017. Transcription factor NF-κB is modulated by symbiotic status in a sea anemone
model of cnidarian bleaching. Sci Rep 7, 16025. https://doi.org/10.1038/s41598-017-16168-w

Marshall, P.A., Baird, A.H., 2000. Bleaching of corals on the Great Barrier Reef: differential susceptibilities among taxa. Coral Reefs 19, 155–163. https://doi.org/10.1007/s003380000086

Martinez, S., Grover, R., Baker, D.M., Ferrier-Pagès, C., 2022a. Symbiodiniaceae Are the First Site of Heterotrophic Nitrogen Assimilation in Reef-Building Corals. mBio 13, e01601-22. https://doi.org/10.1128/mbio.01601-22

Martinez, S., Grover, R., Ferrier-Pagès, C., 2024. Unveiling the importance of heterotrophy for coral symbiosis under heat stress. mBio 15, e01966-24. https://doi.org/10.1128/mbio.01966-24

Martinez, S., Grover, R., Ferrier-Pagès, C., 2022b. Symbiont starvation affects the stability of the coral–Symbiodiniaceae symbiosis. Front. Mar. Sci. 9, 979563. https://doi.org/10.3389/fmars.2022.979563

Martinez, S., Kolodny, Y., Shemesh, E., Scucchia, F., Nevo, R., Levin-Zaidman, S., Paltiel, Y., Keren, N., Tchernov, D., Mass, T., 2020. Energy Sources of the Depth-Generalist Mixotrophic Coral Stylophora pistillata. Front. Mar. Sci. 7, 566663. https://doi.org/10.3389/fmars.2020.566663

McLachlan, R.H., Price, J.T., Solomon, S.L., Grottoli, A.G., 2020. Thirty years of coral heat-stress experiments: a review of methods. Coral Reefs 39, 885–902. https://doi.org/10.1007/s00338-020-01931-9

Melamud, E., Vastag, L., Rabinowitz, J.D., 2010. Metabolomic Analysis and Visualization Engine for LC–MS Data. Anal. Chem. 82, 9818–9826. https://doi.org/10.1021/ac1021166

Meziere, Z., Rich, W.A., Carvalho, S., Benzoni, F., Morán, X.A.G., Berumen, M.L., 2022. *Stylophora* under stress: A review of research trends and impacts of stressors on a model coral species. Science of The Total Environment 816, 151639. https://doi.org/10.1016/j.scitotenv.2021.151639

Molinari, C.G., McDougall, C., Pitt, K.A., 2025. Understanding Dynamic Molecular Responses Is Key to Designing Environmental Stress Experiments: A Review of Gene and Protein Expression in Cnidaria Under Stress. Molecular Ecology 34, e17753. https://doi.org/10.1111/mec.17753

Morgan, K.M., Perry, C.T., Johnson, J.A., Smithers, S.G., 2017. Nearshore Turbid-Zone Corals Exhibit High Bleaching Tolerance on the Great Barrier Reef Following the 2016 Ocean Warming Event. Front. Mar. Sci. 4. https://doi.org/10.3389/fmars.2017.00224

Muller, E.M., Rogers, C.S., Spitzack, A.S., van Woesik, R., 2008. Bleaching increases likelihood of disease on Acropora palmata (Lamarck) in Hawksnest Bay, St John, US Virgin Islands. Coral Reefs 27, 191–195. https://doi.org/10.1007/s00338-007-0310-2

Muscatine, L., Porter, J.W., 1977. Reef Corals: Mutualistic Symbioses Adapted to Nutrient-Poor Environments. BioScience 27, 454–460. https://doi.org/10.2307/1297526

Nakamura, M., Morita, M., Kurihara, H., Mitarai, S., 2011. Expression of hsp70, hsp90 and hsf1 in the reef coral Acropora digitifera under prospective acidified conditions over the next several decades. Biology Open 1, 75–81. https://doi.org/10.1242/bio.2011036

Oliver, T.A., Palumbi, S.R., 2011. Do fluctuating temperature environments elevate coral thermal tolerance? Coral Reefs 30, 429–440. https://doi.org/10.1007/s00338-011-0721-y

Peixoto, R.S., Rosado, P.M., Leite, D.C. de A., Rosado, A.S., Bourne, D.G., 2017. Beneficial Microorganisms for Corals (BMC): Proposed Mechanisms for Coral Health and Resilience. Frontiers in Microbiology 8, 341. https://doi.org/10.3389/fmicb.2017.00341

Petrou, K., Nunn, B.L., Padula, M.P., Miller, D.J., Nielsen, D.A., 2021. Broad scale proteomic analysis of heat-destabilised symbiosis in the hard coral Acropora millepora. Sci Rep 11, 19061. https://doi.org/10.1038/s41598-021-98548-x

Pratchett, M.S., Caballes, C.F., Newman, S.J., Wilson, S.K., Messmer, V., Pratchett, D.J., 2020. Bleaching susceptibility of aquarium corals collected across northern Australia. Coral Reefs 39, 663–673. https://doi.org/10.1007/s00338-020-01939-1

Putnam, H.M., Barott, K.L., Ainsworth, T.D., Gates, R.D., 2017. The Vulnerability and Resilience of Reef-Building Corals. Current Biology 27, R528–R540. https://doi.org/10.1016/j.cub.2017.04.047

Qin, Z., Yu, K., Liang, Y., Chen, B., Huang, X., 2020. Latitudinal variation in reef coral tissue thickness in the South China Sea: Potential linkage with coral tolerance to environmental stress. Science of The Total Environment 711, 134610. https://doi.org/10.1016/j.scitotenv.2019.134610

Quast, J.-P., Schuster, D., Picotti, P., 2022. protti: an R package for comprehensive data analysis of peptide- and protein-centric bottom-up proteomics data. Bioinformatics Advances 2, vbab041. https://doi.org/10.1093/bioadv/vbab041

Rädecker, N., Pogoreutz, C., Gegner, H.M., Cárdenas, A., Roth, F., Bougoure, J., Guagliardo, P., Wild, C., Pernice, M., Raina, J., Meibom, A., Voolstra, C.R., 2021. Heat stress destabilizes symbiotic nutrient cycling in corals. *Proceedings of the National Academy of Sciences* 118, e2022653118. https://doi.org/10.1073/pnas.2022653118

Reshef, L., Koren, O., Loya, Y., Zilber-Rosenberg, I., Rosenberg, E., 2006. The Coral Probiotic Hypothesis. Environmental Microbiology 8, 2068–2073. https://doi.org/10.1111/j.1462-2920.2006.01148.x

Risk, M.J., Edinger, E., 2011. Impacts of Sediment on Coral Reefs, in: Encyclopedia of Modern Coral Reefs. Springer, Dordrecht, pp. 575–586. https://doi.org/10.1007/978-90-481-2639-2_25

Rivera, H.E., Cohen, A.L., Thompson, J.R., Baums, I.B., Fox, M.D., Meyer-Kaiser, K.S., 2022. Palau's warmest reefs harbor thermally tolerant corals that thrive across different habitats. Commun Biol 5, 1394. https://doi.org/10.1038/s42003-022-04315-7

Rodrigues, L.J., Grottoli, A.G., 2007. Energy reserves and metabolism as indicators of coral recovery from bleaching. Limnology and Oceanography 52, 1874–1882. https://doi.org/10.4319/lo.2007.52.5.1874

Rodriguez-Lanetty, M., Harii, S., Hoegh-Guldberg, O., 2010. Corrigendum. Molecular Ecology 19, 1071–1071. https://doi.org/10.1111/j.1365-294X.2010.04545.x

Rodriguez-Lanetty, M., Harii, S., Hoegh-Guldberg, O., 2009. Early molecular responses of coral larvae to hyperthermal stress. Molecular Ecology 18, 5101–5114. https://doi.org/10.1111/j.1365-294X.2009.04419.x

Roessner, U., Bowne, J., 2009. What is metabolomics all about? BioTechniques 46, 363–365. https://doi.org/10.2144/000113133

Rogers, J.S., Monismith, S.G., Koweek, D.A., Torres, W.I., Dunbar, R.B., 2016. Thermodynamics and hydrodynamics in an atoll reef system and their influence on coral cover. Limnology and Oceanography 61, 2191–2206. https://doi.org/10.1002/lno.10365

Rosado, P.M., Leite, D.C.A., Duarte, G.A.S., Chaloub, R.M., Jospin, G., Nunes da Rocha, U., P. Saraiva, J., Dini-Andreote, F., Eisen, J.A., Bourne, D.G., Peixoto, R.S., 2019. Marine probiotics: increasing coral resistance to bleaching through microbiome manipulation. ISME J 13, 921–936. https://doi.org/10.1038/s41396-018-0323-6

Rosic, N., Rémond, C., Mello-Athayde, M.A., 2020. Differential impact of heat stress on reef-building corals under different light conditions. Marine Environmental Research 158, 104947. https://doi.org/10.1016/j.marenvres.2020.104947

Rosic, N.N., Pernice, M., Dove, S., Dunn, S., Hoegh-Guldberg, O., 2011. Gene expression profiles of cytosolic heat shock proteins Hsp70 and Hsp90 from symbiotic dinoflagellates in response to thermal stress: possible implications for coral bleaching. Cell Stress and Chaperones 16, 69–80. https://doi.org/10.1007/s12192-010-0222-x

Rowan, R., 2004. Thermal adaptation in reef coral symbionts. Nature 430, 742–742. https://doi.org/10.1038/430742a

Ruiz-Jones, L.J., Palumbi, S.R., 2017. Tidal heat pulses on a reef trigger a fine-tuned transcriptional response in corals to maintain homeostasis. Sci Adv 3, e1601298. https://doi.org/10.1126/sciadv.1601298

Safaie, A., Silbiger, N.J., McClanahan, T.R., Pawlak, G., Barshis, D.J., Hench, J.L., Rogers, J.S., Williams, G.J., Davis, K.A., 2018. High frequency temperature variability reduces the risk of coral bleaching. Nat Commun 9, 1671. https://doi.org/10.1038/s41467-018-04074-2

Sakai, K., Singh, T., Iguchi, A., 2019. Bleaching and post-bleaching mortality of Acropora corals on a heat-susceptible reef in 2016. PeerJ 7, e8138. https://doi.org/10.7717/peerj.8138

Sammarco, P.W., 1982. Polyp Bail-Out: An Escape Response to Environmental Stress and a New Means of Reproduction in Corals. Marine Ecology Progress Series 10, 57–65.

Sangmanee, K., Casareto, B.E., Nguyen, T.D., Sangsawang, L., Toyoda, K., Suzuki, T., Suzuki, Y., 2020. Influence of thermal stress and bleaching on heterotrophic feeding of two scleractinian corals on pico-nanoplankton. Marine Pollution Bulletin 158, 111405. https://doi.org/10.1016/j.marpolbul.2020.111405

Schwarz, J.A., Mitchelmore, C.L., Jones, R., O'Dea, A., Seymour, S., 2013. Exposure to copper induces oxidative and stress responses and DNA damage in the coral *Montastraea franksi*. Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology 157, 272–279. https://doi.org/10.1016/j.cbpc.2012.12.003

Schweinsberg, M., Gösser, F., Tollrian, R., 2021. The history, biological relevance, and potential applications for polyp bailout in corals. Ecology and Evolution 11, 8424–8440. https://doi.org/10.1002/ece3.7740

Seneca, F.O., Palumbi, S.R., 2015. The role of transcriptome resilience in resistance of corals to bleaching. Molecular Ecology 24, 1467–1484. https://doi.org/10.1111/mec.13125

Seveso, D., Montano, S., Strona, G., Orlandi, I., Galli, P., Vai, M., 2016. Hsp60 expression profiles in the reef-building coral *Seriatopora caliendrum* subjected to heat and cold shock regimes. Marine Environmental Research 119, 1–11. https://doi.org/10.1016/j.marenvres.2016.05.007

Sharp, V.A., Brown, B.E., Miller, D., 1997. Heat shock protein (hsp 70) expression in the tropical reef coral Goniopora djiboutiensis. Journal of Thermal Biology 22, 1–19. https://doi.org/10.1016/S0306-4565(96)00029-0

Shinzato, C., Khalturin, K., Inoue, J., Zayasu, Y., Kanda, M., Kawamitsu, M., Yoshioka, Y., Yamashita, H., Suzuki, G., Satoh, N., 2021. Eighteen Coral Genomes Reveal the Evolutionary Origin of *Acropora* Strategies to Accommodate Environmental Changes. Molecular Biology and Evolution 38, 16–30. https://doi.org/10.1093/molbev/msaa216

Shoguchi, E., Shinzato, C., Hisata, K., Satoh, N., Mungpakdee, S., 2015. The Large Mitochondrial Genome of Symbiodinium minutum Reveals Conserved Noncoding Sequences between Dinoflagellates and Apicomplexans. Genome Biology and Evolution 7, 2237–2244. https://doi.org/10.1093/gbe/evv137

Singh, T., Iijima, M., Yasumoto, K., Sakai, K., 2019. Effects of moderate thermal anomalies on Acropora corals around Sesoko Island, Okinawa. PLoS ONE 14, e0210795. https://doi.org/10.1371/journal.pone.0210795

Skutnik, J.E., Otieno, S., Khoo, S.K., Strychar, K.B., 2020. Examining the Effect of Heat Stress on Montastraea cavernosa (Linnaeus 1767) from a Mesophotic Coral Ecosystem (MCE). Water 12, 1303. https://doi.org/10.3390/w12051303

Stephens, T.G., Lee, J., Jeong, Y., Yoon, H.S., Putnam, H.M., Majerová, E., Bhattacharya, D., 2022. High-quality genome assemblies from key Hawaiian coral species. GigaScience 11, giac098. https://doi.org/10.1093/gigascience/giac098

Suggett, D.J., Nitschke, M.R., Hughes, D.J., Bartels, N., Camp, E.F., Dilernia, N., Edmondson, J., Fitzgerald, S., Grima, A., Sage, A., Warner, M.E., 2022. Toward bio□optical phenotyping of reef□forming corals using Light□Induced FluorescenceTransient□FastRepetition Rate fluorometry. Limnology and Oceanography: Methods 20, 172–191. https://doi.org/10.1002/lom3.10479

Swain, T.D., DuBois, E., Gomes, A., Stoyneva, V.P., Radosevich, A.J., Henss, J., Wagner, M.E., Derbas, J., Grooms, H.W., Velazquez, E.M., Traub, J., Kennedy, B.J., Grigorescu, A.A., Westneat, M.W., Sanborn, K., Levine, S., Schick, M., Parsons, G., Biggs, B.C., Rogers, J.D., Backman, V., Marcelino, L.A., 2016. Skeletal light-scattering accelerates bleaching response in reef-building corals. BMC Ecology 16, 10. https://doi.org/10.1186/s12898-016-0061-4

Swain, T.D., Lax, S., Lake, N., Grooms, H., Backman, V., Marcelino, L.A., 2018. Relating Coral Skeletal Structures at Different Length Scales to Growth, Light Availability to Symbiodinium, and Thermal Bleaching. Front. Mar. Sci. 5. https://doi.org/10.3389/fmars.2018.00450

Ainsworth, T.D., Wasmund, K., Ukani, L., Seneca, F., Yellowlees, D., Miller, D., Leggat, W., 2011. Defining the tipping point. A complex cellular life/death balance in corals in response to stress. Sci Rep 1, 160. https://doi.org/10.1038/srep00160

Thies, A.B., Quijada-Rodriguez, A.R., Zhouyao, H., Weihrauch, D., Tresguerres, M., 2022. A Rhesus channel in the coral symbiosome membrane suggests a novel mechanism to regulate NH$_3$ and CO$_2$ delivery to algal symbionts. Sci. Adv. 8, eabm0303. https://doi.org/10.1126/sciadv.abm0303

Tisthammer, K.H., Forsman, Z.H., Sindorf, V.L., Massey, T.L., Bielecki, C.R., Toonen, R.J., 2016. The complete mitochondrial genome of the lobe coral *Porites lobata* (Anthozoa: Scleractinia) sequenced using ezRAD. Mitochondrial DNA Part B 1, 247–249. https://doi.org/10.1080/23802359.2016.1157770

Tremblay, P., Grover, R., Maguer, J.F., Legendre, L., Ferrier-Pagès, C., 2012. Autotrophic carbon budget in coral tissue: a new 13C-based model of photosynthate translocation. J Exp Biol 215, 1384–1393. https://doi.org/10.1242/jeb.065201

Tyanova, S., Temu, T., Cox, J., 2016. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. Nat Protoc 11, 2301–2319. https://doi.org/10.1038/nprot.2016.136

Veal, C.J., Holmes, G., Nunez, M., Hoegh-Guldberg, O., Osborn, J., 2010. A comparative study of methods for surface area and three-dimensional shape measurement of coral skeletons. Limnology and Oceanography: Methods 8, 241–253. https://doi.org/10.4319/lom.2010.8.241

Voolstra, C.R., Li, Y., Liew, Y.J., Baumgarten, S., Zoccola, D., Flot, J.-F., Tambutté, S., Allemand, D., Aranda, M., 2017. Comparative analysis of the genomes of Stylophora pistillata and Acropora digitifera provides evidence for extensive differences between species of corals. Sci Rep 7, 17583. https://doi.org/10.1038/s41598-017-17484-x

Wang, C., Zheng, X., Kvitt, H., Sheng, H., Sun, D., Niu, G., Tchernov, D., Shi, T., 2023. Lineage-specific symbionts mediate differential coral responses to thermal stress. Microbiome 11, 211. https://doi.org/10.1186/s40168-023-01653-4

Wang, W., Cao, B., Xu, Z., Jia, Z., Yu, S., Tian, P., Niu, W., Xiao, J., 2022. The complete mitochondrial genome of Montipora vietnamensis (Scleractinia, Acroporidae). BDJ 10, e91531. https://doi.org/10.3897/BDJ.10.e91531

Warner, M.E., Fitt, W.K., Schmidt, G.W., 1999. Damage to photosystem II in symbiotic dinoflagellates: A determinant of coral bleaching. Proc. Natl. Acad. Sci. U.S.A. 96, 8007–8012. https://doi.org/10.1073/pnas.96.14.8007

Warner, M.E., Fitt, W.K., Schmidt, G.W., 1996. The effects of elevated temperature on the photosynthetic efficiency of zooxanthellae in hospite from four different species of reef coral: a novel approach. Plant, Cell & Environment 19, 291–299. https://doi.org/10.1111/j.1365-3040.1996.tb00251.x

Williams, A., Chiles, E.N., Conetta, D., Pathmanathan, J.S., Cleves, P.A., Putnam, H.M., Su, X., Bhattacharya, D., 2021a. Metabolomic shifts associated with heat stress in coral holobionts. Sci. Adv. 7, eabd4210. https://doi.org/10.1126/sciadv.abd4210

Williams, A., Pathmanathan, J.S., Stephens, T.G., Su, X., Chiles, E.N., Conetta, D., Putnam, H.M., Bhattacharya, D., 2021b. Multi-omic
characterization of the thermal stress phenome in the stony coral *Montipora capitata*. PeerJ 9, e12335.
https://doi.org/10.7717/peerj.12335
Williams, A., Stephens, T.G., Shumaker, A., Bhattacharya, D., 2023. Peeling back the layers of coral holobiont multi-omics data. iScience.
Woesik, R. van, Sakai, K., Ganase, A., Loya, Y., 2011. Revisiting the winners and the losers a decade after coral bleaching. Marine Ecology
Progress Series 434, 67–76. https://doi.org/10.3354/meps09203
Wright, R.M., Aglyamova, G.V., Meyer, E., Matz, M.V., 2015. Gene expression associated with white syndromes in a reef building coral,
Acropora hyacinthus. BMC Genomics 16, 371. https://doi.org/10.1186/s12864-015-1540-2
Ziegler, M., Eguíluz, V.M., Duarte, C.M., Voolstra, C.R., 2018. Rare symbionts may contribute to the resilience of coral–algal assemblages.
ISME J 12, 161–172. https://doi.org/10.1038/ismej.2017.151
Ziegler, M., Seneca, F.O., Yum, L.K., Palumbi, S.R., Voolstra, C.R., 2017. Bacterial community dynamics are linked to patterns of coral heat
tolerance. Nat Commun 8, 14213. https://doi.org/10.1038/ncomms14213
Zoccola, D., Ganot, P., Bertucci, A., Caminiti-Segonds, N., Techer, N., Voolstra, C.R., Aranda, M., Tambutté, E., Allemand, D., Casey, J.R.,
Tambutté, S., 2015. Bicarbonate transporters in corals point towards a key step in the evolution of cnidarian calcification. Sci
Rep 5, 9983. https://doi.org/10.1038/srep09983

1304

Experimental Design

Great Barrier Reef

Davies Reef

MMM = 28.4°C — Control Tanks (n=3)

High = 32.4°C — Treatment Tanks (n=3)

Tank   Tank   Tank

0.8°C/day

28.4°C
32.4°C

Days

0   5   9
TP0   TP1   TP2

(n=48)

A. hyacinthus   *P. lobata*   *S. pistillata*

Pathway Level Summary

Protein Level Summary

A Ambient Symbiosis

B Thermal Stress

Divergent host responses

C Metabolic Crash

Acropora hyacinthus

Bail out

Stylophora pistillata

Resilient phenotype

Porites lobata