Warming up to a new coat: moulting king penguins exhibit hyperthermia and increased

peripheral heat loss

Juan D Zuluaga[1,2$], Emmanuel Pretti[3], Aude Leynaert[3], Elsa Marçon[4], Antoine Stier[4,5*], Agnès Lewden[3*$]

[1] Department of Biology and Marine Biology, University of North Carolina Wilmington, Wilmington, NC 28403, USA

[2] Department of Ecology and Evolutionary Biology, University of Tennessee Knoxville, Knoxville, TN 37996, USA

[3] Université de Brest - UMR 6539 CNRS/UBO/IRD/Ifremer, Laboratoire des sciences de l'environnement marin -

IUEM - Rue Dumont D'Urville - 29280 - Plouzané

[4] Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

[5] Department of Biology, University of Turku, Turku, Finland

[$]: corresponding authors: jzuluaga@utk.edu & agnes.lewden@univ-brest.fr

[*]: joint senior authorship

Corresponding Author contact information:

Email: jzuluaga@utk.edu

Phone number: (+1) 865-850-3020

Mailing Address: Mossman 217, 1311 Cumberland Ave, Knoxville, TN 37996

Keywords: Heat stress, moult, hyperthermia, thermography, ecophysiology

Abstract

Penguins are among the most specialized thermoregulators on the planet, however,

the same adaptations that maximize heat retention underwater likely hinder heat dissipation on

land, possibly creating dangerous thermoregulatory challenges when encountering warming

terrestrial habitats. Penguins are subject to strictly terrestrial phases, such as moulting, when

metabolic heat production, insulation, and energetic constraints are heightened. We assessed

thermoregulation in moulting captive king penguins (*Aptenodytes patagonicus*) using

simultaneous measurements of core and surface temperatures to test two hypotheses. Under

the *thermal challenge* hypothesis, an initial rise in heat dissipation effort (i.e., increased

peripheral vasomotion) followed by a rise in core temperature would indicate failure to prevent

hyperthermia. Under the *warm-up* hypothesis, an initial rise of core temperature concomitant

or followed by an increase of peripheral vasomotion would indicate regulated hyperthermia,

possibly to accelerate feather development. Core and surface temperatures increased

drastically but concomitantly during moult, providing tentative support for the *warm-up*

hypothesis. Moulting penguins did not pant, suggesting that peripheral heat dissipation was

sufficient to regulate moulting-induced hyperthermia. Core and subcutaneous temperatures in

wild individuals resembled patterns measured in captivity, despite lower heat load and

additional options for behavioural thermoregulation. These results indicate that hyperthermia is

prevalent in moulting king penguins, and documenting the timing of temperature changes

provides novel insights for the moulting physiology of penguins. Because moulting-induced

hyperthermia may contribute to heat load, we caution that moulting may increase the

susceptibility of wild penguins to heat stress, especially as regions near the poles warm at a

disproportionately rapid rate.

Introduction

Maintaining thermal insulation is essential for endotherms to thrive in their natural environments (1). Fur and feathers contribute to thermal insulation in most mammals and birds, but these keratinaceous structures degrade through wear and breakage as they age. Thus, the maintenance of proper insulation from fur or feathers through time requires their replacement through shedding: a protracted, year-round replacement, or moulting: a contracted, punctuated replacement (1). Feathers are unique among morphological traits due to the variety of functions they serve, with major roles in crypsis, mate choice, species signalling, thermoregulation, waterproofing, flight, and protection (2). Moulting is thus considered a key life-history stage because it conditions the ability of individuals to survive and reproduce in their environment (3).

Moult has been well characterized in various avian models from a metabolic perspective. Moult requires a massive amount of protein because feathers are composed of approximately 90% protein and can represent or even exceed 10% of body mass and 25% of total body protein content (4). Synthesizing new feathers also requires an increase in energy expenditure, with moulting birds generally exhibiting a metabolic rate *ca.* 9-111% higher than non-moulting individuals (4). From a thermoregulatory perspective, moult has been less well characterized to the best of our knowledge. The few studies available in birds suggest that an increase of *ca.* 20-60% of thermal conductance occurs during moult (5, 6), which may be explained by a decrease in insulation and/or an increase in evaporative heat transfer due to increased body-water turnover (4). Increases in thermal conductance have been shown to raise the lower critical temperature in long-eared owls (*Asio otus*) from approximately 18°C to 24°C (7) and in emperor penguins (*Aptenodytes forsteri*) from approximately -10°C to 0°C (8). Thus, moult has mostly been considered thermally challenging under cold environmental conditions. Alternatively, it has been hypothesized that the increased metabolism of moulting birds could more than compensate for the decrease in insulation, and thus that the increased heat generation may pose a challenge for moulting birds (4, 9, 10).

Penguins have specialized vascular arrangements that provide countercurrent heat
exchange in the legs (11), head (12), and wings (13). Penguins also boast efficient insulation
through a layer of subcutaneous fat (14), and specialized plumage comprised of an insulating
underlayer and a waterproof overlayer (8, 15). Despite penguins' adaptations for heat retention
underwater, they must breed and moult on land, where dissipating heat is more difficult than in
water (thermal conductance is 2.2-4.8 times higher in water than in air; (16)). Penguins undergo
a catastrophic moult once a year, replacing their entire plumage extremely quickly (between 13
and 34 days; (17)), which involves an increase of metabolic rate (and thus of heat production) of
ca. 30-50% (18, 19). Catastrophic moult involves the temporary overlap of new and old feather
layers (20), which increases thermal insulation (10). Penguins seem relatively sensitive to heat
stress while on land (21–23), and climate change is pronounced in polar and sub-polar regions
(24, 25), where many penguin species moult on land during the warmest season (26). Thus, if
heat dissipation is indeed a challenge during moult (4, 9, 10), the increasing temperatures
penguins are facing while moulting on land may challenge these iconic species.

Captive Gentoo penguins held within their thermoneutral zone had increased thermal
insulation and heat dissipation effort during the overlap of old and new feather layers (10).
Therefore, moulting seems to represent a challenge for heat dissipation in Gentoo penguins, but
the efficiency or inefficiency of their thermoregulatory response remains unknown because
core body temperature was not measured. We define the *thermal challenge* hypothesis as the
scenario where core body temperature rises because of failure to dissipate heat. Alternatively, it
has been suggested that birds may elevate core body temperature while moulting,
hypothetically under physiological control, to favour feather growth (3), an idea we are referring
to as the *warm-up* hypothesis. If heat dissipation effort increases but core temperature does
not, this would indicate that heat dissipation capacity is sufficient to offset the increased
metabolism (4) and insulation (10) that are known to occur during moult. If heat dissipation
effort increases and body temperature increases afterwards, this would indicate a failure to
regulate body temperature, thus supporting the *thermal challenge* hypothesis.

To test these hypotheses, we simultaneously recorded panting behaviour, heat
dissipation effort (*i.e.* surface temperature of thermal windows such as flippers, bill and feet),

insulation (*i.e.* surface temperature of the trunk) and core body temperature in captive king

penguins before, during and after the moult. To address ecological relevance of the results

obtained in captivity, we also include behavioural and physiological data of wild king penguins,

although limited in sample size. Under the thermal challenge hypothesis, we predict that wild

king penguins will be less likely to exhibit hyperthermia because environmental conditions are

colder in the field and because wild king penguins have additional methods to cool themselves

(*e.g.,* immersing feet in cold water). If core body temperature increases before heat dissipation

effort, or if both increase simultaneously, this would indicate a regulated increase in core body

temperature, thus supporting the *warm-up* hypothesis. Under the *warm-up* hypothesis, we

predict that both captive and wild king penguins will exhibit hyperthermia despite

environmental and behavioural variation.

Materials and Methods

Captive king penguins

We studied 8 captive adult king penguins (*Aptenodytes patagonicus*) in the spring of

2024 at Océanopolis aquarium, Brest, France. The birds were individually identified by their

coloured plastic band, and six individuals moulted during the study period. Individuals were

maintained indoors at 8.12 ± 0.07°C (mean ± SE), well within their estimated thermoneutral

zone (*i.e.* ⎯-5 to⎯+22 °C; (27)). They had permanent access to free water to swim/dive, but no

possibility to only immerse their feet. Researchers were the first to enter the enclosure after a

full night with no disturbance to collect thermal images shortly before 8:00 AM, and keepers

entered afterwards throughout the day for regular enclosure cleanings, feeding non-moulting

birds (*i.e.* moulting birds are naturally fasting and stay on land even in captivity). The lighting

program used in the enclosure includes a monthly variation in artificial light, with exposure

varying between 11 and 13 h of light per day during the study. Following the 7-stage moult

categorization (10), we assigned moult stage each day for each individual from uniform old

plumage (M1) to uniform new plumage (M7) before surface temperature data extraction. To

investigate the continuous changes that occur within the visibly distinct moult stages, we

tracked the moulting process numerically by assigning a moult day for each bird, which began at

day 0 when we visually detected changes in the plumage structure (*i.e.* the start of M2, (10)).

Moult actually starts before any visual changes occur (*i.e.* feather growth below the skin), which

occurs at sea in wild penguins and has been estimated to last *ca.* 10 days (20).

Individuals were fed with an ingestible core temperature (T_{core}) data logger (Bodycap,

Anipill, France, accuracy ± 0.2°C between 25 and 45°C, resolution ± 0.01°C, sampling rate =

5min) hidden in their normal diet (*i.e.* without handling) when they began exhibiting

hyperphagia, which precedes the fasting period during catastrophic moult (28). Individuals were

equipped at this stage because retention time in the stomach is quite variable (*ca.* 7 to 80 days

in wild king penguins, *pers. obs.*). We expect that stomach temperature provides a close

approximation of core body temperature during the moult because temperature measurements

in fasting penguins are not influenced by food intake. To avoid collecting measurements of core

temperature that may be influenced by food intake in non-moulting individuals, we collected

temperature data of all birds in the mornings shortly before 8:00 AM after a full fasting night

(*i.e.* 30 minutes after lights turned on, and before any keepers had entered the enclosure to

feed the non-moulting birds). Five moulting individuals were successfully equipped before the

onset of visible plumage change, and one at the time of early plumage change (*i.e.* early M2

phase, (10)). Only one (non-moulting) individual lost the logger before the completion of the

study, 32 days after ingestion. Moulting individuals were equipped on average 16.5 ± 4.8 days

before visible changes in plumage occurred. Data were downloaded remotely with an

Anilogger® monitor placed inside the enclosure throughout the study.

Upon entering the enclosure to collect data, the presence/absence of panting behaviour

was recorded for each individual (22), and we used a thermal imaging camera (three different

cameras were used due to technical issues: E96, E8, or T640; resolutions provided in Electronic

Supplementary Materials, *ESM*) to collect images of each individual from a distance of 1 m (10).

We collected images of birds in the profile orientation (29). Air temperature (T_a) and relative

humidity (RH) inside the enclosure were measured using a weather station (Kestrel 5400;

Kestrel® instrument, United Kingdom). Thermal imaging collection started on average 23.3 ± 5.4

days before visible changes in plumage occurred. Pictures of individuals being wet at the time of

thermal imaging were excluded from the analysis (*i.e.* n = 36 images from non-moulting

individuals, or moulting ones with moult day > 20). We report the average core body

temperature from 7:30 to 8:00 AM. This approach resulted in a sample size of 291 datapoints

with body surface and core temperatures measured simultaneously.

After collecting thermal images, we extracted surface temperatures using FLIR Thermal

Studio Pro. We set emissivity to 0.98 (10) and we specified T_a and RH for each image using

measurements collected during each thermal imaging session. We then measured the average

surface temperatures of the bill, flipper, and foot (T_{bill}, $T_{flipper}$, and T_{foot}, respectively) by drawing

separate region of interest (ROI) polygons for each area. We measured the maximum surface

temperature of the periorbital region (henceforth T_{eye}; (30)) using an ellipse. We measured

ground temperature (T_{ground}) below the bird using a line segment the length of the foot. To

determine a representative trunk surface temperature (T_{trunk}) despite irregular feather loss, we

measured the temperature of both the old and newly grown plumage (T_{old} and T_{new}) using a

square with side length equal to ~one third the length of the flipper. We used visual landmarks

to measure T_{ground}, T_{old}, and T_{new} because image resolution varied between the three cameras.

After standardizing between the cameras (see *ESM*), we used values of T_{old} and T_{new} to calculate

T_{trunk} according to (10). The moulting pattern of king penguins is sufficiently patchy for squares

to be consistently drawn in a section of completely old or new plumage with no overlap.

Data analysis was conducted in R Version 4.3.3 (31). We first calculated wet-bulb

temperature (T_w) in R package HeatStress (32) using the measurements of T_a and RH collected

while sampling. Following guidance from two external (*i.e.,* non-author) experts in the field of

thermography, we standardized measurements made with the three cameras using two

approaches (*i.e.* see ESM S0 for details). We then modelled T_{core}, T_{eye}, T_{bill}, $T_{flipper}$, T_{foot}, and T_{trunk}

as a function of moult day. To capture the non-linear changes of body temperatures over the

course of the moult (in moulting birds only), we created segmented models based on calculated

changes in slope (*i.e.,* breakpoints; (33)). We determined the presence/timing of breakpoints

and the associated 95% intervals using the R package segmented (34). After identifying

breakpoints, we fitted separate linear mixed models ((lmer()) in R package lme4 (35)) for each

time period as determined by the breakpoints for moulting birds. Although thermal conditions

were held relatively constant (Fig. S1), we included T_w as a fixed effect in all global models to

account for any subtle variation in thermal conditions; we used T_{ground} instead of T_w for the

models of foot temperature according to (10). Non-moulting birds were analysed in separate

mixed models and were assigned an arbitrary moulting day for graphical purpose only (Fig. 1).

We included the bird's assigned identification number (bird ID) as a random intercept in all

models to account for repeated measures. Model support was determined using AICc (36) using

the function dredge() in R package MuMIn (37). We report the top-ranked model with

parameter estimates for moult day and T_w or T_{ground} (Table S1). We assessed the fit of the top

models by calculating the conditional Nakagawa's R^2 using function r.squaredGLMM() in R

package MuMIn. We visually represented findings from our models by plotting model

predictions obtained from package ggeffects (38) along with raw data for each response

variable using package ggplot2 (39). Finally, predicted values and the associated 95% confidence

intervals were extracted from key stages based on breakpoint (*i.e.* pre-moult, peak-moult, post-

moult) as well as non-moulting individuals using semi-parametric bootstrapping (n = 1000

simulations) with confint() and bootMer() functions of R package lme4 (35)).

Wild king penguins

Wild king penguins were studied in 2023-24 (as part of a long-term observatory) in the

'Baie du Marin' colony on Possession Island (40), Crozet Archipelago, in the Southern Indian

Ocean (46°26' S, 51°52' E) from courtship in November 2023 to the 17[th] of October 2024, when

all scientific activities were shut down because of an outbreak of highly pathogenic avian

influenza (HPAI) at the study site (41). At breeding initiation (Nov-Dec 2023), 48 breeding pairs

were temporarily marked with hair dye on the breast feathers (*i.e.* lasting until the post-

breeding moult) and permanently marked with thermosensitive pit-tags (BioTherm13 tag,

resolution ± 0.1°C, accuracy ± 0.5°C, between 33°C and 43°C, Biomark, USA) in the inter-scapular

region. These thermosensitive pit-tags enable long-term individual identification and

subcutaneous temperature measurements (22). Subcutaneous temperature measurements

were conducted during incubation (3[rd] day of an incubation shift on land) as part of other long-

term monitoring procedures, in less than 3 minutes after capture to avoid handling-stress

induced changes in body temperature (40). A subsample of birds was equipped with core

temperature ingestible loggers (Bodycap, Anipill, France, sampling rate = 5 or 15 min) during

incubation. At the time of post-breeding moult (Sep-Oct 2024), the colony was searched daily to

identify marked penguins being back on land to moult (N = 43 penguins observed 1 to 10 times

during moult), scoring their moult according to ((10); M1 to M7 stages), noting the

presence/absence of panting behaviour (22) and the usage of the river as a potentially cooling

substrate (*i.e.* feet in the river: yes/no). Approximately on their 3^{rd} day on land, 3 moulting

individuals (*i.e.* stage M1 for two individuals and stage M2 for another one) were recaptured,

had their subcutaneous temperature recorded in less than 3 minutes after capture, and were

equipped with a core temperature ingestible logger. At moult completion (stage M7), individuals

were re-captured to download core body temperature data, which was unfortunately only

possible for one individual before field site closure linked to HPAI outbreak. Considering the

limited sample size, no statistical tests were performed and individual data points are

presented. To quantify the environmental heat load faced by wild *vs.* captive individuals (*i.e.*

including solar radiation and wind speed in addition to air temperature and humidity used for

T_w in captivity), Wet Bulb Globe Temperature (WBGT) values were computed using the R

package HeatStress function wbgt.Liljegren(). Environmental heat load in the wild was lower

during moult than breeding, and captive penguins experienced an environmental heat load

being intermediate between moulting and breeding individuals in the wild (Fig. S1).

Results

Captive penguins

Moulting and non-moulting birds were never observed panting during the study. The visible part of moult lasted on average 24.3 ± 0.9 days in captive individuals.

Changes in body temperatures during moult followed a non-linear pattern (Fig. 1). While T_{core} was best fitted with 2 breakpoints (*i.e.* an increase followed by a decrease during moult and then stabilisation post-moult, Fig 1a, Table S1), all surface temperatures were best fitted with at least 3 breakpoints that corresponded to 3 main phases: pre-moult, moult (with 2 segments surrounding peak-moult: early-moult and late-moult) and post-moult (Fig. 1b to 1f). While surface temperatures (T_{eye}, T_{bill}, $T_{flipper}$, T_{foot} and T_{trunk}) during pre- and post-moult periods were not always constant, the magnitude of changes during these periods were limited (Fig. 1 and Table S1). During moult, T_{eye}, T_{bill}, $T_{flipper}$ and T_{foot} initially increased during early-moult (although non-significantly for T_{foot}) and then decreased during late-moult, so post-moult levels were comparable to pre-moult and non-moulting levels (Fig. 1b to 1e; Fig 2b to 2e; Table S1). Conversely, T_{trunk} (*i.e.* an inverse proxy of insulation provided by the plumage) initially decreased and then increased back to levels comparable to pre-moult and non-moulting levels (Fig. 1f and 2f, Table S1).

Despite birds being equipped with a T_{core} logger on average 16.5 ± 4.8 days before visible changes in plumage occurred (*i.e.* moult day 0), we did not detect a pre-moulting stable period for T_{core} (Fig. 1a). Thus, we do not report a predicted T_{core} for the pre-moult stage (Fig. 2a), and we instead draw inferences by comparing T_{core} values between peak-moult and post-moult. T_{core} was *ca.* 1.3°C higher during peak moult than post-moult or than in non-moulting individuals. Body temperatures were overall mostly indistinguishable between pre-moult, post-moult and non-moulting individuals (Fig. 2). The changes in surface temperatures during moult were drastic, with an increase at peak moult of *ca.* +4.0°C for T_{eye} (Fig. 2b), +11.5°C for T_{bill} (Fig. 2c), +10.2°C for $T_{flipper}$ (Fig. 2d), +5.0°C for T_{foot} although 95% CI overlapped with most other stages (Fig. 2e), and a decrease of *ca.* -5.0°C for T_{trunk} (Fig. 2f).

For body surface temperatures, the initial rise (*i.e.* moult start) in T_{eye}, T_{bill}, $T_{flipper}$ and

T_{foot} occurred simultaneously around 10 days before moult day 0 (Fig. 3). The initial drop in T_{trunk}

occurred slightly later (*ca.* moult day -6), but not significantly so considering 95% CI overlapped

(Fig. 3). The peak in $T_{flipper}$ was reached approximately 10 days earlier than the peaks in T_{core},

T_{eye}, T_{bill}, and T_{foot}, while the peak in plumage insulation (*i.e.* lowest T_{trunk}) occurred

approximately 5 days later (Fig. 3, non-overlapping 95% CI). The peak in T_{core} occurred

approximately 3 days before the peaks In T_{eye}, T_{bill} and T_{foot}, but not significantly so considering

95% CI overlapped (Fig. 3). T_{foot} was the first to return to baseline levels at *ca.* moult day 14,

followed by T_{trunk}, T_{bill}, T_{eye} (*ca.* moult day 27), then T_{core} (*ca.* moult day 34) and then only by $T_{flipper}$

(*ca.* moult day 44). While 95% CI overlapped between T_{core} and T_{bill}, T_{eye}, $T_{flipper}$, this was not the

case between T_{core} and T_{trunk} (Fig. 3).

A glimpse at the natural scenario

Out of 87 observations from 43 wild individuals (mostly during early moulting stages),

none were observed panting, whereas breeding individuals have been reported to pant during

ca. 20% of behavioural observations in the same population (22). Moulting birds immersed their

feet in the river during 75.8% of our observations. The visible part of moult lasted 13 days in the

only individual that completed its moult before the HPAI-related closure of the field site.

T_{subcut} was approximately 1.4°C higher in early-moulting birds compared to the same

individuals measured while breeding (Fig. 4a). Average daily T_{core} was overall 1.2°C higher during

moulting compared to breeding in one wild individual (Fig. 4b), despite environmental heat load

(*i.e.* WBGT) being lower during moult (Fig. 4c) and this individual being consistently observed

with its feet immersed in the river. This wild individual had T_{core} within the range of captive non-

moulting individuals during breeding (except the first day), and within the range of captive

peak-moult between approximately 4 and 11 days of its moulting fast on land (Fig. 4b). T_{core} of

this moulting individual did not return to breeding or captive non-moulting values, even once

moult was visibly completed (Fig. 4b).

Discussion

Warming up to a new coat

Our results provide evidence that moulting elicits thermovascular responses that increase heat loss in king penguins, as previously described in gentoo penguins (10). We now also provide evidence that core body temperature in king penguins markedly increases during moult (*i.e.* up to +1.4°C and +1.2°C in captive and wild birds respectively), suggesting either a failure to dissipate excess heat generated by the up-regulation of metabolism (*thermal challenge hypothesis*), or a regulated increase of body temperature during moult (*warm-up hypothesis*). By assessing the timing of changes in surface and core body temperature, we found that surface temperatures increased rapidly, and likely after the rise in core temperature occurred, thereby providing support for the *warm-up* hypothesis. Coupled with the lack of panting observed in the study period, these findings suggest that increasing peripheral heat dissipation was likely enough to maintain core temperature below a maximum threshold. The limited evidence we gathered from wild king penguins (*i.e.* +1.4°C in subcutaneous and +1.2°C in core body temperature) suggests that the hyperthermic pattern we observed in captivity also occurs under natural conditions, where behavioural heat dissipation was less constrained because individuals had access to cold water, and used it for partial immersion.

King penguins must accomplish moulting efficiently to avoid reproductive failure because food availability decreases rapidly after the moulting period (42). It is known that king penguins that arrive from the sea later in the season initiate feather loss more quickly and undergo a contracted moult period relative to individuals that arrive earlier (43). Hyperthermia may benefit moulting penguins by accelerating the process of feather growth, thus enabling them to better match high food availability. Assuming that king penguins in our study had a Q10 equal to 2.3 (Roussel et al., *pers. Com.*; 44) the observed increase in body temperature of ~1.4°C would yield a 12.4% increase in metabolic rate, which may shorten moult by *ca.* 2.5 days, equal to a 9.3% reduction in the total duration of moult.

Positive correlations between feather growth and body temperature have been observed in chickens, where individuals with an early feathering genotype display higher body

temperatures and faster feather growth during the first six weeks post-hatching (45). A similar
pattern has been observed indirectly in turkeys, where individuals held at higher ambient
temperature had longer feathers at 14 days post-hatching (46), although this may partially be
explained by metabolic costs being diverted to thermoregulation instead of feather growth at
the lower temperature treatment used in that study. Further supporting the association
between high body temperature and accelerated feather growth, the majority of feather growth
occurs during the day when body temperature and metabolic rate are highest (47). In penguins,
moulting-induced hyperthermia occurs alongside increases in blood perfusion at the surface
((10); this study), which likely helps to deliver nutrients to peripheral tissues that is necessary
for feather growth (48).

Thermoregulation in captivity and in the wild during catastrophic moult

Core and surface temperatures in captive non-moulting birds remained stable during the
study period relative to moulting birds, which exhibited distinct changes over time. Core
temperature was the first to increase under captive conditions as we were not able to detect a
pre-moulting stable period. During the gradual rise in core temperature, we observed increases
in surface temperature in thermal windows (flippers, bill, and feet), indicating increased effort
to dissipate heat. Interestingly, the flippers were the only thermal window to reach the
maximum surface temperature before core body temperature reached its maximum, suggesting
that the flippers may function as a first line of defense for peripheral heat dissipation in
penguins, likely due to their large surface area and vascularization (49). On the other hand, the
surface temperatures of the bill and feet closely tracked fluctuations in core temperature, as
indicated by the similarities in the timing and direction of changes in these temperatures. That
heat dissipation via thermal windows increased after core temperature had already been rising
indicates that hyperthermia was likely regulated and maintained below a maximum threshold.
This interpretation is further supported by the lack of panting observed throughout the study
period. We found that the surface temperature of the trunk declined in captive moulting birds,
indicating increased insulation due to the overlap of new and old layers of plumage (10, 50).

Despite the increased insulation during the moult, core temperature had already begun
declining by the time insulation reached its maximum (i.e., the minimum of T_{trunk}). This
observation suggests that the rise in core temperature was primarily driven by increased
metabolic heat generation, not by increased insulation as hypothesized in (10).

Conducting this research in captivity enabled us to eliminate variation that would be
introduced by more dynamic thermal environments in the wild, thus isolating the effects of
moulting on thermoregulation. On one hand, captivity prevented heating from solar radiation;
on the other hand, captivity prevented cooling from wind chill and precipitation, and it limited
behavioural thermoregulation in the form of habitat selection (i.e., partial immersion in water).
However, behavioural thermoregulation and environmental conditions, such as solar radiation,
wind, and humidity remain essential considerations for wild animal thermoregulation (22, 51–
53). Therefore, to provide insights about the relevance of our finding in captivity, we measured
thermoregulatory behaviour, subcutaneous temperature, and core temperature in a limited
number of wild king penguins. During 75.8% of our observations, free-living moulting king
penguins had their feet immersed in the river, confirming that wild birds likely take advantage of
additional means to dissipate heat. Despite this behaviour and greater potential for wind chill,
core temperature measured in one wild moulting king penguin showed a similar hyperthermic
pattern (+1.2°C compared to incubation) as the one found in captive moulting king penguins
(+1.4°C). Although this result was obtained in only one individual, it is supported by the similar
pattern observed in the subcutaneous temperature of three wild king penguins measured
during both incubation and early-moult (+1.4°C). It is interesting to note that the core body
temperature of captive individuals was similar to the wild scenario, both during ($T_{core} \simeq 38.7$°C)
and outside ($T_{core} \simeq 37.3$°C) the moulting period, thereby suggesting that captivity does not
markedly affect thermoregulation in king penguins.

Seeking baselines: Operating points of body and surface temperatures

The data we report conform well to a negative feedback-based model of
thermoregulation with multiple autonomous thermo-effectors (54). The elevation of metabolic

heat generation while moulting represents a positive (i.e., heat) thermal challenge that ultimately triggers an increase in heat dissipation via the periphery. The rises in core, and subsequently surface, temperatures represent the changes in multiple operating points that would be necessary to balance heat loss with elevated heat generation. The thermal challenge introduced by moulting was large enough to elicit changes in peripheral heat loss, but not panting, suggesting that the thermoregulatory response observed lies within an inter-threshold zone (55–58). We interpret the difference in timing and magnitude of change in flipper temperature relative to other thermal windows to represent varying thresholds for response across body regions, a pattern supported by the specialization of regional heterothermy in birds (59), and indeed by the widespread presence of regional heterothermy in endotherms (60).

Previous studies have suggested that moulting poses a negative (i.e., cold) thermoregulatory challenge due to reduced insulation during feather loss, which is compensated by an increase in heat production (17–19, 61). If moulting poses a negative thermal challenge, one would expect a reduction in body temperature followed by upregulation of metabolic heat generation until the thermal challenge passes (i.e., hypothermia, not hyperthermia (54)). However, moulting in penguins [(62, 63), present study] and other birds (5, 64) can elicit elevations in body temperature (i.e., hyperthermia), the opposite of the pattern predicted if moulting was a negative thermal challenge. Moulting-induced hyperthermia may be less pronounced in species where heat loss occurs too efficiently for body temperature to be affected by upregulated metabolism (e.g., small passerines; (65)), or if the sampling period does not align with the onset of hyperthermia. In the wild, penguins arrive onshore with new feathers developed to between 18% to 42% of their final length (18, 66), and it is possible that studies in wild individuals may not have been able to detect early changes in core temperature occurring during the pre-moult or early-moult stages. Pre-moult elevation of core temperature is not mutually exclusive with increased energy cost of thermoregulation in later stages of the moult when insulation decreases quickly, as observed in (19).

Conducting this research in captivity enabled us to measure moulting and non-moulting individuals simultaneously throughout the entire moulting process. Despite starting to measure core temperature more than two weeks before the first visible change in plumage, our sampling

period was too late to capture a stable pre-moult phase of core temperature. In gentoo
penguins, lower surface temperatures were observed at the end of the study, possibly because
the sampling period may have begun after body temperatures had already begun increasing
(10). This and previous studies highlight the difficulty of establishing a baseline for
normothermia, and these differences in sampling period may explain some of the differences
between studies. We advise care when planning sampling periods for physiological studies of
moulting because physiological changes in preparation for moulting that underly visible change
in plumage may have an earlier start than what was previously appreciated in the literature.

We also advise care external and internal factors when measuring thermoregulatory
traits, such as core temperature, surface temperatures, and thermoregulatory behaviors,
because these can be influenced by multiple. For instance, acute stress can cause temperature
changes in regions throughout the body (29, 33, 40, 67) and body temperatures also exhibit
circadian patterns (68). Acute stress and circadian changes in body temperature could affect
behaviours that vary with heat load, such as panting, a common thermoregulatory behavior in
birds, including penguins (69, 70). In our system, we identified the daily visits of keepers as
potential sources of error because of variation in timing, tasks, and levels of disturbance
(underwater cleaning being less stressful than above-water cleaning). To avoid both stress-
related and circadian influences on body temperatures and thermoregulation, we collected data
at the same time each morning before keepers entered the enclosure. Yet, such potential issues
are way more difficult to deal with in wild settings, where timing of measurement is not
necessarily chosen by the experimenter, and acute stressors (*e.g.* aggressive interaction with
neighbours, presence of predator) cannot be controlled and may not be noticed by the
experimenter.

Warming up in captivity – heating up in the wild?

We found that the increase in core temperature in moulting king penguins was likely
regulated, as evidenced by the lack of influence from plumage insulation, the absence of
panting behavior, and the delay in vascular perfusion. These patterns suggest that hyperthermia

was likely employed as a regulated thermoregulatory strategy (71). Air temperature remained
stable and far below body temperature during our study, and the warming effect of solar
radiation was absent in captivity. Since environmental conditions are unlikely to be as forgiving
in the wild, the current study raises concerns regarding the vulnerability of moulting penguins in
the context of global climate change. High mortality rates have already been reported in
multiple penguin species immediately after moulting due to the energetic challenge of this
fasting period (18, 72–74). More recently, in 2019, a mortality event was recorded during early
moult in little penguins following a heatwave on land (*Eudyptula minor*; (75)). Based on our
finding that moulting poses a positive (i.e., heat) thermal challenge, we caution that moulting
may increase the susceptibility of wild penguins to heat stress, especially as regions near the
poles warm at a disproportionately rapid rate (24). Further investigation of thermoregulatory
responses of penguins during moulting in the wild is thus deeply needed.

Data availability and Electronic Supplementary Materials

446 The datasets, R code, and Electronic Supplementary Materials (Supplemental Figs. S0
447 and S1; Supplemental Tables S0-A, S0-B, and Table S1) used in this manuscript are available on
448 FigShare:

449 https://figshare.com/articles/dataset/Data_from_Warming_up_to_a_new_coat_moul_ng_king_p
450 enguins_exhibit_hyperthermia_and_increased_peripheral_heat_loss_/28629305

Acknowledgments

We are grateful to Oceanopolis© and Dominique Barthelemy for their logistical support during
the experiment. This work would not have been possible without the help of the animal care team:
Christine Dumas, Alexiane Corcuff, Maxence Leroy, Mélanie Robert and Agathe Lefranc. We thank the
members of polar program #119 (ECONERGY) JP Robin, VA Viblanc, P Bize, as well as to the French Polar
Institute (IPEV) and the Terres Australes et Antarctiques Françaises for providing financial and logistical
support for the study of wild king penguins. We are grateful to Marine Montblanc, Norith Eckbo, Samuel
Laporte, Zohria-Lys Guillerm, Camille De Pasquale and the entire Alfred Faure field station for their help

in the field. We thank Raymond M. Danner for providing financial and logistical support to JDZ, and for

his helpful comments on the proposal. We thank the following students for their assistance in extracting

data from thermal images: Abby E. Lehman, Olivia P. Wanex, Roberto Bonifacio-Dominguez, Tyler W.

Vanwinkle. We thank Glenn J. Tattersall and Dominic J. McCafferty for providing feedback regarding

standardization of the thermal camera measurements, as well as D. Roussel for providing unpublished

Q10 values of king penguins. The wild part of this study was part of the long-term Studies in Ecology and

Evolution (SEE-Life) program of the CNRS. AL was supported by ISblue project, Interdisciplinary graduate

school for the blue planet (ANR-17-EURE-0015) and co-funded by a grant from the French government

under the program "Investissements d'Avenir" embedded in France 2030. **This project received financial**

support from the "Région Bretagne". The project benefitted from an IdEx grant from the Université de

Strasbourg (*HotPenguin*) granted to AS. EM was funded by the French Polar Institute. We thank the

Company of Biologists and the Journal of Experimental Biology for providing a Traveling Fellowship to

JDZ.

Author's contributions

475   Study design: ALew, AS, JZ. Funding acquisition: ALew, AS, ALey, JZ. Data collection in captivity:

476   JZ, EP, ALew. Data collection in the field: EM, AS. Data analysis: JZ, EP, AS. Writing original draft: JZ, ALew

477   and AS. Writing review and editing: ALey, EP, EM.

Institutional Review Board Statement (IRB)

480   The study in captive penguins did not include invasive procedures. Scientific procedures for wild

481   king penguins were approved by the French Ethical Committee (*APAFIS#31268-2021042117037897 v3*)

482   and the Terres Australes et Antarctiques Françaises (Arrêtés TAAF A-2023-88).

New and Noteworthy

Penguins may experience heat stress while moulting, which causes increased metabolic heat

generation and insulation. We assessed thermoregulation in moulting captive king penguins

(*Aptenodytes patagonicus*) using simultaneous measurements of core and surface temperatures. By

measuring temperature throughout the entirety of the moult, we found that hyperthermia and

increased peripheral heat dissipation are prevalent in moulting king penguins. We caution that moulting-

induced hyperthermia may contribute to the susceptibility of penguins to heat stress in the wild.

References

1.   **Beltran RS**, **Burns JM**, **Breed GA**. Convergence of biannual moulting strategies across
birds and mammals. *Proc R Soc B* 285: 20180318, 2018. doi: 10.1098/rspb.2018.0318.

2.   **Terrill RS**, **Shultz AJ**. Feather function and the evolution of birds. *Biological Reviews* 98:
540–566, 2023. doi: 10.1111/brv.12918.

3.   **Jenni L**, **Winkler R**. *The Biology of Moult in Birds*. London: HELM, 2020.

4.   **Carey C**, editor. Energetics and Nutrition of Molt. In: *Avian Energetics and Nutritional*
Ecology. Springer US, p. 158–198.

5.   **Lustick S**. Energy Requirements of Molt in Cowbirds. *The Auk* 87: 742–746, 1970. doi:
10.2307/4083708.

6.   **Dietz MW**, **Daan S**, **Masman D**. Energy Requirements for Molt in the Kestrel *Falco*
tinnunculus. *Physiological Zoology* 65: 1217–1235, 1992. doi:
10.1086/physzool.65.6.30158276.

7.   **Wijnandts H**. Ecological Energetics of the Long-Eared Owl (Asio Otus). *Ardea* 38–90: 1–
92, 2002. doi: 10.5253/arde.v72.p1.

8.   **Le Maho Y**, **Delclitte P**, **Chatonnet J**. Thermoregulation in fasting emperor penguins
under natural conditions. *American Journal of Physiology-Legacy Content* 231: 913–922,
1976. doi: 10.1152/ajplegacy.1976.231.3.913.

9.   **Gavrilov V**. Metabolism of molting birds. *Zoologicheskii Zhurnal* 53: 1363–1375, 1974.

10. **Lewden A**, **Halna Du Fretay T**, **Stier A**. Changes in body surface temperature reveal the
thermal challenge associated with catastrophic moult in captive gentoo penguins. *Journal*
of Experimental Biology 227: jeb247332, 2024. doi: 10.1242/jeb.247332.

11. **Kazas S**, **Benelly M**, **Golan S**. The Humboldt Penguin (Spheniscus humboldti) Rete
Tibiotarsale – A supreme biological heat exchanger. *Journal of Thermal Biology* 67: 67–78,
2017. doi: 10.1016/j.jtherbio.2017.04.011.

12. **Frost PGH**, **Siegfried WR**, **Greenwood PJ**. Arterio-venous heat exchange systems in the
Jackass penguin *Spheniscus demersus*. *Journal of Zoology* 175: 231–241, 1975. doi:
10.1111/j.1469-7998.1975.tb01398.x.

13. **Thomas DB**, **Fordyce RE**. Biological Plasticity in Penguin Heat-Retention Structures. *The*
Anatomical Record 295: 249–256, 2012. doi: 10.1002/ar.21538.

14. **Enstipp MR**, **Bost C-A**, **Le Bohec C**, **Bost C**, **Le Maho Y**, **Weimerskirch H**, **Handrich Y**.
Apparent changes in body insulation of juvenile king penguins suggest an energetic
challenge during their early life at sea. *Journal of Experimental Biology* 220: 2666–2678,
2017. doi: 10.1242/jeb.160143.

15. **Ainley DG**, **Wilson RP**. Hot Penguins: Cold Water. In: *The Aquatic World of Penguins*.
Springer International Publishing, p. 217–256.

16. **De Vries J**, **Van Eerden MR**. Thermal Conductance in Aquatic Birds in Relation to the
Degree of Water Contact, Body Mass, and Body Fat: Energetic Implications of Living in a
Strong Cooling Environment. *Physiological Zoology* 68: 1143–1163, 1995. doi:
10.1086/physzool.68.6.30163797.

17. **Adams NJ**, **Brown CR**. Energetics of Molt in Penguins. In: *Penguin Biology*. Elsevier, p.
297–315.

18. **Cherel Y**, **Charrassin JB**, **Challet E**. Energy and protein requirements for molt in the king
penguin Aptenodytes patagonicus. *American Journal of Physiology-Regulatory, Integrative*
and Comparative Physiology 266: R1182–R1188, 1994. doi:
10.1152/ajpregu.1994.266.4.R1182.

19. **Baudinette R**, **Gill P**, **O'driscoll M**. Energetics of the Little Penguin, Eudyptula Minor:
Temperature Regulation, the Calorigenic Effect of Food, and Moulting. *Aust J Zool* 34: 35,
1986. doi: 10.1071/ZO9860035.

20. **Groscolas R**, **Cherel Y**. How to Molt While Fasting in the Cold: The Metabolic and
Hormonal Adaptations of Emperor and King Penguins. *Ornis Scandinavica* 23: 328, 1992.
doi: 10.2307/3676657.

21. **Stonehouse B**. The General Biology and Thermal Balances of Penguins. In: *Advances in*
Ecological Research. Elsevier, p. 131–196.

22. **Noiret A**, **Lewden A**, **Lemonnier C**, **Bocquet C**, **Montblanc M**, **Bertile F**, **Hoareau M**,
Marcon E, **Robin J-P**, **Viblanc VA**, **Stier A**. Mind the polar sun: Solar radiations trigger
frequent heat stress in breeding king penguins, despite relatively cool air temperatures.
Evolutionary Biology: 2024.

23. **Welman S**, **Pichegru L**. Nest microclimate and heat stress in African Penguins
Spheniscus demersus breeding on Bird Island, South Africa. *Bird Conservation*
International 33: e34, 2023. doi: 10.1017/S0959270922000351.

24. **Nel W**, **Hedding DW**, **Rudolph EM**. The sub-Antarctic islands are increasingly warming in
the 21st century. *Antarctic Science* 35: 124–126, 2023. doi: 10.1017/S0954102023000056.

25. **Gorodetskaya IV**, **Durán-Alarcón C**, **González-Herrero S**, **Clem KR**, **Zou X**, **Rowe P**,
Rodriguez Imazio P, **Campos D**, **Leroy-Dos Santos C**, **Dutrievoz N**, **Wille JD**,
Chyhareva A, **Favier V**, **Blanchet J**, **Pohl B**, **Cordero RR**, **Park S-J**, **Colwell S**, **Lazzara**
MA, **Carrasco J**, **Gulisano AM**, **Krakovska S**, **Ralph FM**, **Dethinne T**, **Picard G**. Record-
high Antarctic Peninsula temperatures and surface melt in February 2022: a compound
event with an intense atmospheric river. *npj Clim Atmos Sci* 6: 202, 2023. doi:
10.1038/s41612-023-00529-6.

26. **Davis LS**, **Darby JT**. *Penguin biology*. San Diego New York Boston [etc.]: Academic press,
1990.

27. **Froget G**, **Handrich Y**, **Maho YL**, **Rouanet J-L**, **Woakes AJ**, **Butler PJ**. The heart
rate/oxygen consumption relationship during cold exposure of the king penguin: a
comparison with that during exercise. *Journal of Experimental Biology* 205: 2511–2517,
2002. doi: 10.1242/jeb.205.16.2511.

28.   **Cherel Y**, **Charrassin J-B**, **Handrich Y**. Comparison of Body Reserve Buildup in
Prefasting Chicks and Adults of King Penguins (Aptenodytes patagonicus). *Physiological*
Zoology 66: 750–770, 1993. doi: 10.1086/physzool.66.5.30163822.

29.   **Tabh JKR**, **Burness G**, **Wearing OH**, **Tattersall GJ**, **Mastromonaco GF**. Infrared
thermography as a technique to measure physiological stress in birds: Body region and
image angle matter. *Physiol Rep* 9, 2021. doi: 10.14814/phy2.14865.

30.   **Jerem P**, **Jenni-Eiermann S**, **Herborn K**, **McKeegan D**, **McCafferty DJ**, **Nager RG**. Eye
region surface temperature reflects both energy reserves and circulating glucocorticoids in
a wild bird. *Sci Rep* 8: 1907, 2018. doi: 10.1038/s41598-018-20240-4.

31.   **R Core Team**. R: A Language and Environment for Statistical Computing [Online]. R
Foundation for Statistical Computing: 2025. https://www.R-project.org/.

32.   **Casanueva A**. HeatStress. Zenodo: 2019.

33.   **Zuluaga JD**, **Danner RM**. Acute stress and restricted diet reduce bill-mediated heat
dissipation in the song sparrow (*Melospiza melodia*): implications for optimal
thermoregulation. *Journal of Experimental Biology* 226: jeb245316, 2023. doi:
10.1242/jeb.245316.

34.   **Muggeo VMR**. segmented: Regression Models with Break-Points / Change-Points
Estimation (with Possibly Random Effects). 2.1-4, 2003.

35.   **Bates D**, **Mächler M**, **Bolker B**, **Walker S**. Fitting Linear Mixed-Effects Models Using
lme4. *J Stat Soft* 67, 2015. doi: 10.18637/jss.v067.i01.

36.   **Burnham KP**, **Anderson DR**, editors. Model Selection and Multimodel Inference. Springer
New York.

37.   **Bartoń K**. MuMIn: Multi-Model Inference. 1.48.11, 2010.

38.   **Lüdecke D**. ggeffects: Tidy Data Frames of Marginal Effects from Regression Models.
JOSS 3: 772, 2018. doi: 10.21105/joss.00772.

39.   **Wickham H**. ggplot2: Elegant Graphics for Data Analysis. Springer New York.

40.   **Lewden A**, **Ward C**, **Noiret A**, **Avril S**, **Abolivier L**, **Gérard C**, **Hammer TL**, **Raymond É**,
Robin J-P, **Viblanc VA**, **Bize P**, **Stier A**. Surface temperatures are influenced by handling
stress independently of corticosterone levels in wild king penguins (Aptenodytes
patagonicus). *Journal of Thermal Biology* 121: 103850, 2024. doi:
10.1016/j.jtherbio.2024.103850.

41.   **Clessin A**, **Briand F-X**, **Tornos J**, **Lejeune M**, **Pasquale CD**, **Fischer R**, **Souchaud F**,
Hirchaud E, **Bralet T**, **Guinet C**, **McMahon CR**, **Grasland B**, **Baele G**, **Boulinier T**. Mass
mortality events in the sub-Antarctic Indian Ocean caused by long-distance circumpolar
spread of highly pathogenic avian influenza H5N1 clade 2.3.4.4b. Microbiology: 2025.

42.   **Olsson O**. Seasonal Effects of Timing and Reproduction in the King Penguin: A Unique
Breeding Cycle. *Journal of Avian Biology* 27: 7, 1996. doi: 10.2307/3676955.

43. **Gauthier-Clerc M**, **Le Maho Y**, **Gendner J-P**, **Handrich Y**. Moulting fast and time constraint for reproduction in the king penguin. *Polar Biol* 25: 288–295, 2002. doi: 10.1007/s00300-001-0342-y.

44. **Monternier P-A**, **Marmillot V**, **Rouanet J-L**, **Roussel D**. Mitochondrial phenotypic flexibility enhances energy savings during winter fast in king penguin chicks. *J Exp Biol* 217: 2691–2697, 2014. doi: doi:10.1242/jeb.104505.

45. **Noubandiguim M**, **Erensoy K**, **Sarica M**. Feather growth, bodyweight and body temperature in broiler lines with different feathering rates. *SA J An Sci* 51, 2021. doi: 10.4314/sajas.v51i1.10.

46. **Wylie LM**, **Robertson GW**, **MacLeod MG**, **Hocking PM**. Effects of ambient temperature and restricted feeding on the growth of feathers in growing turkeys. *British Poultry Science* 42: 449–455, 2001. doi: 10.1080/00071660120070631.

47. **Lillie FR**, **Wang H**. Physiology of Development of the Feather: IV. The Diurnal Curve of Growth in Brown Leghorn Fowl. *Proc Natl Acad Sci USA* 26: 67–85, 1940. doi: 10.1073/pnas.26.1.67.

48. **Lillie FR**. Physiology of Development of the Feather. III Growth of the Mesodermal Constituents and Blood Circulation in the Pulp. *Physiological Zoology* 13: 143–176, 1940. doi: 10.1086/physzool.13.2.30151536.

49. **Lewden A**, **Nord A**, **Bonnet B**, **Chauvet F**, **Ancel A**, **McCafferty DJ**. Body surface rewarming in fully and partially hypothermic king penguins. *J Comp Physiol B* 190: 597–609, 2020. doi: 10.1007/s00360-020-01294-1.

50. **Le Maho Y**. The Emperor Penguin: A Strategy to Live and Breed in the Cold: Morphology, physiology, ecology, and behavior distinguish the polar emperor penguin from other penguin species, particularly from its close relative, the king penguin. *American Scientist* 65: 680–693, 1977.

51. **Freeman MT**, **Coulson B**, **Short JC**, **Ngcamphalala CA**, **Makola MO**, **McKechnie AE**. Evolution of avian heat tolerance: The role of atmospheric humidity. *Ecology* 105: e4279, 2024. doi: 10.1002/ecy.4279.

52. **Wolf BO**, **Walsberg GE**. Thermal Effects of Radiation and Wind on a Small Bird and Implications for Microsite Selection. *Ecology* 77: 2228–2236, 1996. doi: 10.2307/2265716.

53. **Cunningham SJ**, **Gardner JL**, **Martin RO**. Opportunity costs and the response of birds and mammals to climate warming. *Frontiers in Ecology and the Environment* 19: 300–307, 2021.

54. **Mitchell D**, **Fuller A**, **Snelling EP**, **Tattersall GJ**, **Hetem RS**, **Maloney SK**. Revisiting concepts of thermal physiology: understanding negative feedback and set-point in mammals, birds, and lizards. *Biological Reviews* 100: 1317–1346, 2025. doi: 10.1111/brv.70002.

55. **Bligh J**. The thermosensitivity of the hypothalamus and thermoregulation in mammals. *Biological Reviews* 41: 317–365, 1966. doi: 10.1111/j.1469-185X.1966.tb01496.x.

56. **Notley SR**, **Mitchell D**, **Taylor NAS**. A century of exercise physiology: concepts that
ignited the study of human thermoregulation. Part 3: Heat and cold tolerance during
exercise. *Eur J Appl Physiol* 124: 1–145, 2024. doi: 10.1007/s00421-023-05276-3.

57. **Taylor NAS**, **Gordon CJ**. The origin, significance and plasticity of the thermoeffector
thresholds: Extrapolation between humans and laboratory rodents. *Journal of Thermal
Biology* 85: 102397, 2019. doi: 10.1016/j.jtherbio.2019.08.003.

58. **Mitchell D**, **Snelling EP**, **Hetem RS**, **Maloney SK**, **Strauss WM**, **Fuller A**. Revisiting
concepts of thermal physiology: Predicting responses of mammals to climate change.
Journal of Animal Ecology 87: 956–973, 2018. doi: 10.1111/1365-2656.12818.

59. **McQueen A**, **Barnaby R**, **Symonds MRE**, **Tattersall GJ**. Birds are better at regulating
heat loss through their legs than their bills: implications for body shape evolution in
response to climate. *Biol Lett* 19: 20230373, 2023. doi: 10.1098/rsbl.2023.0373.

60. **Angilletta M**, **Cooper B**, **Schuler M**, **Boyles J**. The evolution of thermal physiology in
endotherms. *Front Biosci* E2: 861–881, 2010. doi: 10.2741/e148.

61. **Green JA**, **Butler PJ**, **Woakes AJ**, **Boyd IL**. Energetics of the moult fast in female
macaroni penguins *Eudyptes chrysolophus*. *Journal of Avian Biology* 35: 153–161, 2004.
doi: 10.1111/j.0908-8857.2004.03138.x.

62. **Farner DS**. Incubation and Body Temperatures in the Yellow-eyed Penguin. *The Auk* 75:
249–262, 1958.

63. **Groscolas R**. Study of molt pasting followed by an experimental forced fasting in the
emperor penguin Aptenodytes forsteri: Relationship between feather growth, body weight
loss, body temperature and plasma fuel levels. *Comparative Biochemistry and Physiology
Part A: Physiology* 61: 287–295, 1978. doi: 10.1016/0300-9629(78)90111-1.

64. **Newton I**. The Temperatures, Weights, and Body Composition of Molting Bullfinches. *The
Condor* 70: 323–332, 1968. doi: 10.2307/1365926.

65. **Chilgren J**. Dynamics and bioenergetics of postnuptial molt in captive White-crowned
Sparrows (Zonotrichia leucophrys gambelii). Washington State University: 1975.

66. **Brown CR**. Feather growth, mass loss and duration of moult in macaroni and rock-hopper
penguins. *Ostrich* 57: 180–184, 1986. doi: 10.1080/00306525.1986.9633647.

67. **Herborn KA**, **Jerem P**, **Nager RG**, **McKeegan DEF**, **McCafferty DJ**. Surface temperature
elevated by chronic and intermittent stress. *Physiology & Behavior* 191: 47–55, 2018. doi:
10.1016/j.physbeh.2018.04.004.

68. **Prinzinger R**, **Preßmar A**, **Schleucher E**. Body temperature in birds. *Comparative
Biochemistry and Physiology Part A: Physiology* 99: 499–506, 1991. doi: 10.1016/0300-
9629(91)90122-S.

69. **Murrish DE**. Acid-Base Balance in Three Species of Antarctic Penguins Exposed to
Thermal Stress. *Physiological Zoology* 55: 137–143, 1982.

70. **Welman S**, **Green JA**, **Ryan PG**, **Parsons NJ**, **Pichegru L**. Body temperature and
thermoregulatory behaviour in the Endangered African Penguin *Spheniscus demersus*.
Bird Conservation International 34: e29, 2024. doi: 10.1017/S095927092400025X.

71. **Gerson AR**, **McKechnie AE**, **Smit B**, **Whitfield MC**, **Smith EK**, **Talbot WA**, **McWhorter
TJ**, **Wolf BO**. The functional significance of facultative hyperthermia varies with body size
and phylogeny in birds. *Functional Ecology* 33: 597–607, 2019. doi: 10.1111/1365-
2435.13274.

72. **Heezik YV**, **Davis L**. Effects of food variability on growth rates, fledging sizes and
reproductive success in the Yellow-eyed Penguin *Megadyptes antipodes*. *Ibis* 132: 354–
365, 1990. doi: 10.1111/j.1474-919X.1990.tb01055.x.

73. **Boersma PD**. Breeding Patterns of Galápagos Penguins as an Indicator of Oceanographic
Conditions. *Science* 200: 1481–1483, 1978. doi: 10.1126/science.200.4349.1481.

74. **Dann P**, **Cullen JM**, **Thoday R**, **Jessop R**. Movements and Patterns of Mortality at Sea of
Little Penguins *Eudyptula minor* from Phillip Island, Victoria. *Emu - Austral Ornithology* 91:
278–286, 1991. doi: 10.1071/MU9910278.

75. **Tworkowski L**, **Dann P**, **Ellenburg U**, **Robert K**. Impacts of terrestrial heat waves on
survival of little penguins during moult. 2021.

 Figure legend:

 Figure 1. Variation in core (a) and surface temperatures (b: eye, c: bill, d: flipper, e: foot and f: trunk) of

 captive king penguins in relation to moult dynamics. The dashed line at day 0 indicates the first visible

 signs of moult, and the one at *ca.* day 24 indicates the completion of plumage renewal. Coloured circles

 and solid lines represent moulting birds, whereas white circles and dashed lines represent non-moulting

 birds. Lines are predictions from the top ranked models (Table S1). n = 6 moulting and n = 2 non-

 moulting individuals.

 Figure 2. Predicted core (a) and surface body temperatures (b: eye, c: bill, d: flipper, e: foot and f:

 trunk) of captive king penguins at key stages during moult (*i.e.* pre-moult, peak-moult and post-moult

 based on breakpoint analyses) and in non-moulting individuals. Error bars represent 95% CI; n = 2-6

 moulting and n = 2 non-moulting individuals.

 Figure 3. Temporal dynamics of changes in body temperatures during moult in captive king penguins

 based on breakpoint analyses. The dashed line at day 0 indicates the first visible signs of moult, and the

 one at *ca.* day 24 indicates the average time to visual completion of plumage renewal. Error bars

 represent 95% CI, N = 6 captive individuals.

 Figure 4. Subcutaneous (a) and core (b) body temperatures of moulting *vs.* breeding wild king

 penguin(s), along with (c) corresponding environmental heat load measured as Wet Bulb Globe

 Temperature. The same individual(s) were measured during incubation and moulting within the same

 breeding cycle (2023-24). (a) N = 3 individuals measured during incubation and early moult (*ca.* 3[rd] day of

 fasting on land), with individual values presented in colour with grey lines, and mean ± SE in colour with

 black dashed line. (b) N = 1 individual measured during incubation (blue) and moulting (red) within the

 same breeding cycle (2023-24), circles and error bars represent daily average core body temperature ±

 SE, and the black arrow represents the visible part of moult (*i.e.* from new feathers emergence to fully

 new plumage). For comparison to results from captive individuals, the shaded areas represent the 95%

 CI of core body temperature for captive non-moulters (blue, n = 2) and during peak moult (red, n = 6). (c)

 Circles and error bars represent daily average Wet Bulb Globe Temperature ± SE during incubation (blue)

731   and moulting (red), and the shaded purple area represent the 95% CI of Wet Bulb Globe Temperature for

732   captive individuals for comparison.

(a)

(b) visible part of moult

(c) visible part of moult

Moult Breeding

Captive T$_{core}$ peak moult (95% C.I.)

Captive T$_{core}$ non-moulting (95% C.I.)

Captive wet bulb globe temperature (95% C.I.)