A Sensitive Period for the Development of Temporally Coherent

Visual Cortical Processing in Humans: Evidence from Sight

Recovery Following Congenital Cataracts

Suddha Sourav [a, *], Max Emanuel Feucht [a], Ramesh Kekunnaya [b], and Brigitte Röder [a, b]

[a] Biological Psychology and Neuropsychology, University of Hamburg, 20146 Hamburg, Germany

[b] Jasti V Ramanamma Children's Eye Care Center, Child Sight Institute, L V Prasad Eye Institute,

500034 Hyderabad, India

Author Information

Suddha Sourav: (iD) https://orcid.org/0000-0002-4172-7711; Contributions: Designed research,

performed research, contributed unpublished analytic tools, analyzed data, wrote the paper.

* Corresponding author. E-Mail: suddha.sourav@uni-hamburg.de

Max Emanuel Feucht: (iD) https://orcid.org/0009-0009-5929-9258; Contributions: Performed

research, analyzed data, wrote the paper.

Ramesh Kekunnaya: (iD) https://orcid.org/0000-0001-5789-2300; Contributions: Designed research,

performed research, wrote the paper.

Brigitte Röder: (iD) https://orcid.org/0000-0003-3088-8023; Contributions: Designed research,

performed research, wrote the paper.

The authors have no conflicts to declare.

20 Abstract

Typical cortical development requires early sensory experience. In humans who had suffered

from a transient period of blindness due to dense bilateral congenital cataracts (CC), multiple

investigations of cortical activation *strength* have suggested that extrastriate visual regions are

especially vulnerable to aberrant early postnatal visual input. Yet, to date, how early experience

might sculpt the temporal stability of human visual cortical processing has not been

investigated. In CC individuals, we previously found a persistent and replicable reduction of

the earliest event-related marker of extrastriate processing, the *P1 wave*. Here, we report that

this robust reduction of the P1 in CC individuals results from an attenuated temporal coherence

of cortical oscillations across trials, rather than from a generally reduced extrastriate cortical

activation strength. In two experiments, compared to matched normally sighted controls, CC

individuals (n = 12 and 13, respectively) exhibited diminished oscillatory phase coherence (i.e.,

higher temporal variability), rather than lower oscillatory amplitudes during visual processing.

Furthermore, phase coherence information, but not activation strength, allowed classifying

sight recovery individuals with congenital vs. developmental etiologies. Finally, exchanging the

signal phase and activation strength information between CC and matched control individuals

confirmed that group differences in P1 amplitude were largely driven by oscillatory timing.

These group differences, specific to the CC group, were not found in groups of sight-recovery

individuals who had suffered from developmental cataracts. Our results indicate that the

intricate temporal orchestration of visual processing in human extrastriate cortex requires

visual experience in an early postnatal sensitive period.

Significance Statement

Rare individuals who were born with pattern visual blindness, but later recovered sight, provide

a unique opportunity for probing how early visual experience shapes brain development. A few

existing studies have reported more impaired extrastriate rather than striate cortical activation

strength in sight recovery individuals' visual cortex. Here, we demonstrate that this extrastriate

processing difference can be largely explained by a marked increase of the temporal variability

of oscillatory brain activity during visual processing, rather than by a general impairment of

visual cortical excitation. These results suggest that precisely temporally coordinated visual

processing critically depends on early visual experience, and in turn, indicate that the temporal

tuning of neural circuits might be of utmost importance for functional brain development.

Introduction

Sight in early childhood is essential for the typical development of the visual system (Wiesel

and Hubel, 1965a; Röder and Kekunnaya, 2021; Mitchell and Maurer, 2022). In humans who

had suffered from a period of congenital pattern visual blindness due to the presence of total,

bilateral congenital cataracts (CC), persistent visual deficits have been reported following sight

restoration, even after extended recovery periods. These deficits include a permanently

reduced visual acuity and stereoscopic ability (Tytla et al., 1993; Birch et al., 2009), atypical

mid- to high-level visual feature processing (Putzar et al., 2007; McKyton et al., 2015), and

altered face processing (Röder et al., 2013; De Heering and Maurer, 2014). In contrast, a

substantially higher recovery of visual functions has been observed if the onset of visual loss

was later (instead of congenital), e.g., through developmental cataracts (DC, Röder et al.,

2013; Bottari et al., 2018; Sourav et al., 2020). These results indicate the presence of *sensitive*

periods, during which the effect of experience is especially strong in the development of many

visual cortical circuits and the associated functions (Maurer and Lewis, 2013; Röder et al.,

2021).

Previous research in kittens and macaques have suggested a higher vulnerability of

extrastriate visual cortical areas compared to the primary visual cortex (Wiesel and Hubel,

1965b; Hyvärinen et al., 1981) following a period of bilateral congenital visual deprivation.

Corroborating these classic non-human animal results, in a rare, well characterized group of

sight recovered humans, we have previously reported a marked attenuation of the *P1 wave* of

the visual event-related potential (ERP) derived from electroencephalographic (EEG) data

(Sourav et al., 2020). The P1 wave peaks approximately 145 ms after visual stimulation and

has been indicated to have an extrastriate cortical origin (Di Russo et al., 2002; Miller et al.,

2015). The P1 reduction observed in CC individuals had a very large effect size and was

consistent across participants, allowing excellent classification of congenital vs. developmental

etiology in sight recovery individuals (Sourav et al., 2020). However, the precise neural

mechanism behind this persistent P1 wave reduction following a period of transient congenital

visual deprivation remains unexplained.

ERPs are derived by the averaging of scalp-recorded electrophysiological activity across many

trials. On one extreme, a reduction of the P1 wave in CC individuals might result from a

generally reduced cortical activation across trials; on the other extreme, a similar P1 wave

reduction could arise from a lack of temporal coherence across individual trials, with

unimpaired activation strength across the trials. At the level of the neural generators, these two

hypothetical extremes represent fundamentally different effects of a period of transient visual

deprivation, i.e., a generally suppressed extrastriate cortical activity vs. compromised temporal

stability of extrastriate visual cortical processing. To uncover the mechanism behind the P1

wave reduction, thus, is of essential importance to understand basic mechanisms of

experience-dependent brain development.

Here, we investigated the relative contribution of a reduced extrastriate cortical activation

strength vs. impaired temporal coherence to the persistent P1 wave reduction in CC

individuals. In the first part of a three-pronged approach, we employed time-frequency

decomposition using wavelet transforms to calculate the activation magnitude (operationalized

by the post-stimulus relative power change of cortical oscillations) and the oscillatory inter-trial

phase coherence. These two markers were subsequently compared between groups.

Secondly, we attempted to classify CC individuals based on relative power changes (compared

to pre-stimulus baseline), as well as inter-trial phase coherence information in the time-

frequency representations. Finally, we derived what we term *"mutual magnitude-phase*

transplants", a novel method that we here introduce, which allowed us to exchange the per-

trial activation strength and timing information between a cataract reversal individual and their

matched control participant. Applying this method, we investigated the relative contribution of

amplitude vs. phase information to the persistent P1 wave reduction after a period of transient

congenital visual deprivation. Finally, minimum-norm source estimation was performed to

localize the cortical sources of the P1 wave.

Materials and Methods

Participants

Datasets originating from two different experiments, termed experiment 1 and experiment 2, were utilized. Both experiments included a group of congenital cataract reversal individuals (CC) as well as a group of developmental cataract reversal individuals (DC), who had undergone the same surgical procedures. In addition, normally sighted control participants with a typical developmental history were recruited, and each normally sighted control participant was matched to a cataract reversal individual for age, sex, and handedness.

Thirteen CC individuals took part in experiment 1, of whom the data of 1 participant were not analyzed here due to a very short visual deprivation period (1 month). The remaining CC participants, whose data entered further analysis (n = 12, see Table 1 for participant details), had a mean age of 18.67 years (range = 10 – 37 years), 2 were female (and 10 were male), and 1 was left-handed. In addition, 16 DC individuals took part in experiment 1, of whom the data of 3 individuals were not analyzed respectively due to a diagnosis of seizure and developmental delay, the participant's unwillingness to continue the experiment shortly after the start and the inability of the participant to look straight ahead. The data of 13 DC individuals (Table 2) entered further analysis (mean age = 16 years, range = 11 – 24 years, 5 females and 8 males, all right-handed). The CC individuals had received surgical treatment after a mean of 41.33 months of blindness (range 4 – 213 months). The average age at treatment for DC individuals was 117.38 months (range 24 – 208 months). The visual acuity (VA) of the CC individuals (geometric mean, decimal = 0.23, range = 0.05 – 0.5) was significantly lower compared to that of the DC individuals (geometric mean, decimal = 0.74, range = 0.44 – 1.00), as revealed by a one-tailed t-test after converting the VAs to LogMAR units, t(14.531) = 5.642, p < .001. The mean age of the typically sighted control individuals matched to the CC individuals (MCC group) was 18.08 years (range = 10 – 34 years, 2 female and 10 male, 1 left-handed), and 16.85 years for the typically sighted control individuals matched to the DC individuals (MDC group, range = 10 – 25 years, 5 female and 8 male, all right-handed).

In experiment 2, 15 participants with a history of bilateral, dense congenital cataracts took part.

The data of 2 of these participants were rejected because in one case, the participant was born

with incomplete cataract and thus did not suffer from complete pattern visual blindness since

birth, and in another case, because of a very short visual deprivation period (1 month). Thus,

the data of 13 CC individuals entered further analysis (Table 3, mean age = 17.69 years, range

6 – 39 years, 3 female and 10 male, 1 left-handed). In addition, 16 DC individuals took part in

experiment 2. We did not analyze the data of 1 participant due to a history of seizures and

developmental delay. The data of 15 DC individuals entered further analysis (Table 4, mean

age = 14.47 years, range 9 – 24 years, 4 female and 11 male, 1 left-handed).

The CC individuals in experiment 2 whose data were analyzed (n = 13) had been treated for

cataract after a mean of 45.31 months of blindness (range 4 – 213 mo.), and the DC individuals

whose data were analyzed had been treated for cataract with a mean age of 88.33 months

(range 23 – 173 mo.). Once again, the CC individuals exhibited a generally lower mean

geometric visual acuity (0.21, range = 0.051 – 0.5) than the DC individuals (0.618, range =

0.23 – 1.00; one-tailed t-test, $t(21.532) = 5.297$, p < .001). The MCC group in experiment 2 had

a mean age of 17 years (range = 7 – 38 years, 3 female and 10 male, 1 lefthanded), and the

MDC group had a mean age of 14.93 years (range = 11 – 26 years, 4 female and 11 male, 1

lefthanded).

Ten CC individuals, 8 DC individuals and 5 normally sighted control individuals took part in

both experiment 1 and experiment 2; 35.7% of cataract reversal individuals were therefore

unique to experiment 2 (i.e., 3 CC and 7 DC individuals). All cataract reversal individuals from

both experiments were recruited and tested at the L V Prasad Eye Institute, Hyderabad, India.

Control individuals were tested at the University of Hamburg, Hamburg, Germany. Adult

participants received a small monetary compensation for taking part in the study, whereas

minors received a small gift. Costs for travel and accommodation were reimbursed. The study

obtained approval from the local ethics commission of the faculty of Psychology and Human

Movement at the University of Hamburg and the institutional ethics review board of LVPEI, and

adhered to the ethical principles outlined in the Declaration of Helsinki (World Medical

Association, 2013) .

Experimental Design

In both experiments, participants were presented monochromatic square wave grating stimuli

in front of a grey background, one at a time for 150 ms in one of the four visual field quadrants

with a mean interstimulus interval of 1.85 s (1.5 s – 2.2 s). The stimuli were presented with a

fixed eccentricity of 4 degrees from the center and subtended a visual angle of 2.5° as depicted

in Figure 2 in Sourav et al. (2020). The stimuli were presented at a polar angle of 25° in the

upper visual field and at a polar angle of −45° in the lower visual field. The gratings were either

horizontally (standard stimuli, P = .80) or vertically oriented (target stimuli, P = .20), with a

spatial frequency of 2 cycles/degree. Participants were instructed to attend to a central fixation

cross and to indicate when a rare target stimulus appeared in any of the locations by pressing

a foot pedal in experiment 1 and by clicking a computer mouse in experiment 2. Sight restored

cataract individuals were tested using a Dell IN2030 monitor at LVPEI, Hyderabad, and

typically sighted control individuals were tested on a Samsung P2370 monitor at the University

of Hamburg. Both monitors had a refresh rate of 60 Hz and a nominal luminance of 250 cdm⁻

². Participants sat at a distance of 45 cm to the monitor in experiment 1, and at a distance of

65 cm in experiment 2. The angular measurements described above were identical in both

experiments. Participants were presented additional stimuli in both experiments which are not

reported here.

For stimulus presentation we used the the PsychoPy framework (v1.83, Peirce, 2009).

Table 1. Participant characteristics for the sight recovery individuals with a history of dense bilateral congenital cataracts and subsequent surgery
(CC) in experiment 1. The table is based on Sourav et al. (2020).

Participant ID	Group	Age (yrs.)	Blindness Duration (mo.)	Time Since Surgery (mo.)	Sex	Handedness	Visual Acuity (VA), decimal	VA, LogMAR	Pre-surgical VA (Better Eye)	Nystagmus	Strabismus	Family History
CC – 001	CC	26	5	308	M	Right	0.180	0.745	NA	Yes	Esotropia	No
CC – 002	CC	37	24	428	M	Right	0.400	0.398	NA	Yes	Esotropia	Yes
CC – 003	CC	11	7	124	M	Right	0.430	0.367	Fixates and follows light	Yes	Exotropia	No
CC – 004	CC	13	4	158	M	Right	0.130	0.886	Fixates and follows light	Yes	Exotropia	No
CC – 005	CC	33	72	330	M	Right	0.051	1.292	NA	Yes	No	Yes
CC – 006	CC	23	4	279	M	Left	0.130	0.886	NA	Yes	Esotropia	Yes
CC – 007	CC	13	15	150	M	Right	0.500	0.301	Fixates and follows light	Yes	Exotropia	Yes
CC – 009	CC	10	11	116	M	Right	0.240	0.620	Fixates and follows light	Yes	Esotropia	No
CC – 010	CC	18	25	198	F	Right	0.310	0.509	Fixates and follows light	Yes	Esotropia	Yes
CC – 011	CC	19	213	25	M	Right	0.200	0.699	Counting fingers at 0.5 m	Yes	Esotropia	Yes
CC – 012	CC	11	42	89	F	Right	0.450	0.347	Fixates and follows light	Yes	Exotropia	Yes
CC – 013	CC	10	74	48	M	Right	0.250	0.602	0.017	Yes	No	Yes

182 **Table 2.** Participant characteristics for the sight recovery individuals with a history of bilateral developmental cataracts and subsequent surgery (DC)
183 in experiment 1. The table is based on Sourav et al. (2020).

Participant ID	Group	Age (yrs.)	Age at Surgery (yrs.)	Time Since Surgery (mo.)	Sex	Handedness	Visual Acuity (VA), decimal	VA, LogMAR	Pre-surgical VA (Better Eye)	Nystagmus	Strabismus	Family History
DC – 001	DC	16	12	53	M	Right	1.000	0.000	Counting fingers at 0.5 m	No	No	No
DC – 002	DC	19	14	55	M	Right	1.000	0.000	0.200	No	No	No
DC – 003	DC	12	6	63	F	Right	1.000	0.000	0.500	No	No	No
DC – 004	DC	11	8	39	F	Right	0.530	0.276	0.159	No	No	No
DC – 005	DC	13	8	68	M	Right	0.710	0.149	0.100	No	Exotropia	No
DC – 006	DC	12	8	51	M	Right	0.600	0.222	0.400	No	No	No
DC – 007	DC	12	6	77	M	Right	1.000	0.000	0.250	No	No	Yes
DC – 008	DC	18	12	74	F	Right	0.790	0.102	Sees hand movement	No	No	No
DC – 009	DC	16	10	67	M	Right	0.740	0.131	0.400	No	No	No
DC – 010	DC	24	2	264	F	Right	0.440	0.357	Fixates and follows light	No	Exotropia	No
DC – 011	DC	16	7	110	F	Right	0.530	0.276	Counting fingers @ 1 m	No	Exotropia	Yes
DC – 012	DC	20	12	97	M	Right	0.880	0.056	0.400	No	No	No
DC – 013	DC	19	17	17	M	Right	0.730	0.137	0.200	No	No	No

185 **Table 3.** Participant characteristics for the sight recovery individuals with a history of dense bilateral congenital cataracts and subsequent surgery
186 (CC) in experiment 2 (Please note that age of the same participant might vary from Table 1 due to the experiments being run at different time points).
187 The table is based on Sourav et al. (2020).

Participant ID	Group	Age (yrs.)	Blindness Duration (mo.)	Time Since Surgery (mo.)	Sex	Handedness	Visual Acuity (VA), decimal	VA, LogMAR	Pre-surgical VA (Better Eye)	Nystagmus	Strabismus	Family History
CC – 001	CC	26	5	308	M	Right	0.180	0.745	NA	Yes	Esotropia	No
CC – 005	CC	33	72	330	M	Right	0.051	1.292	NA	Yes	No	Yes
CC – 014	CC	6	5	70	M	Right	0.240	0.620	Not fixating/following light	Yes	No	No
CC – 002	CC	39	24	444	M	Right	0.400	0.398	NA	Yes	Esotropia	Yes
CC – 004	CC	14	4	175	M	Right	0.130	0.886	Fixates and follows light	Yes	Exotropia	No
CC – 006	CC	24	4	294	M	Left	0.130	0.886	NA	Yes	Esotropia	Yes
CC – 007	CC	15	15	166	M	Right	0.500	0.301	Fixates and follows light	Yes	Exotropia	Yes
CC – 009	CC	11	11	129	M	Right	0.240	0.620	Fixates and follows light	Yes	Esotropia	No
CC – 015	CC	8	48	48	F	Right	0.150	0.824	Perception of light	Yes	No	Yes
CC – 013	CC	11	74	61	M	Right	0.250	0.602	0.017	Yes	No	Yes
CC – 012	CC	11	42	90	F	Right	0.450	0.347	Fixates and follows light	Yes	Exotropia	Yes
CC – 011	CC	21	213	40	M	Right	0.200	0.699	Counting fingers at 0.5 m	Yes	Esotropia	Yes
CC – 016	CC	11	72	68	F	Right	0.260	0.585	Fixates and follows light	Yes	No	No

189 **Table 4.** Participant characteristics for the sight recovery individuals with a history of bilateral developmental cataracts and subsequent surgery (DC)
190 in experiment 2 (Please note that age of the same participant might vary from Table 2 due to the experiments being run at different time points). The
191 table is based on Sourav et al. (2020).

Participant ID	Group	Age (yrs.)	Age at Surgery (yrs.)	Time Since Surgery (mo.)	Sex	Handedness	Visual Acuity (VA), decimal	VA, LogMAR	Pre-surgical VA (Better Eye)	Nystagmus	Strabismus	Family History
DC – 002	DC	19	14	55	M	Right	1.000	0.000	0.200	No	No	No
DC – 011	DC	14	7	80	F	Right	0.530	0.276	Finger counting at 1 m	No	Exotropia	Yes
DC – 003	DC	13	6	74	F	Right	1.000	0.000	0.500	No	No	No
DC – 005	DC	14	8	80	M	Right	0.710	0.149	0.100	No	Exotropia	No
DC – 007	DC	12	6	77	M	Right	1.000	0.000	0.250	No	No	Yes
DC – 014	DC	12	7	66	M	Right	0.230	0.638	0.125	No	No	No
DC – 015	DC	12	7	67	M	Right	0.460	0.337	0.100	No	No	No
DC – 009	DC	16	10	67	M	Right	0.740	0.131	0.400	No	No	No
DC – 010	DC	24	2	264	F	Right	0.440	0.357	Fixates and follows light	No	Exotropia	No
DC – 008	DC	18	12	74	F	Right	0.790	0.102	Perceives hand movement	No	No	No
DC – 016	DC	17	12	60	M	Right	0.310	0.509	0.160	No	No	Yes
DC – 017	DC	9	2	86	M	Right	0.930	0.032	0.154	No	No	No
DC – 018	DC	16	6	113	M	Right	0.520	0.284	Finger counting at 1 m	No	No	No
DC – 019	DC	10	3	76	M	Left	0.730	0.137	0.065	No	No	Yes
DC – 020	DC	11	1	115	M	Right	0.667	0.176	0.154	No	No	No

EEG: Data Acquisition and Preprocessing

EEG data were acquired continuously during the whole experiment using 32 Ag/AgCl

electrodes with a left earlobe online reference. Brain activity was recorded at the 10-10

locations FP1, FP2, F7, F3, Fz, F4, F8, FC5, FC1, FCz, FC2, FC6, T7, C3, Cz, C4, T8, TP9,

CP5, CP1, CP2, CP6, TP10, P7, P3, Pz, P4, P8, O1, O2, F9, and F10, with an online high-

pass filter with a cutoff frequency of 0.016 Hz, a low-pass filter with a cutoff frequency of 250

Hz, and a sampling rate of 1 kHz. To eliminate electrical line noise, the data were notch-filtered

offline at 50 Hz and, if apparent in the frequency spectrum, at the corresponding harmonics.

Subsequently, the voltages were average referenced and biological artifacts such as blinks,

eye movements, muscle activity or heartbeats were removed from the signal with an

independent component analysis (ICA), using EEGLAB for MATLAB (EEGLAB, version

11.5.4b, Delorme and Makeig, 2004; MATLAB version 2012b; MathWorks, Natick, MA). To

prevent confounds due to eye movements or blinks during stimulus presentation, we excluded

stimulus epochs that contained blinks or eye movements within the time window of −25 ms to

175 ms relative to the stimulus, as previously reported (Sourav et al., 2020, 2024). The data

were then low pass filtered at 40 Hz and EEG epochs were generated from -1.7s before to 1.7

s after the stimulus onset. For increasing the statistical power, data from trials with a left

hemifield stimulation were remapped; at each electrode, the data were swapped with the data

of the electrode in its mirrored location with respect to the nasion-inion axis, ignoring the

midline electrodes (see Sourav et al., 2020). Additionally, to prevent artifacts related to motor

activation, epochs containing a motor response within ±500 ms of the stimulus onset were

rejected.

Only EEG trials/ERPs generated by the frequent, standard stimuli were used for further

analysis. Moreover, as P1 amplitudes elicited by upper visual field stimulation are larger than

those elicited by lower visual field stimulations in the present paradigm, we only used visual

ERPs for upper visual field stimulation for the subsequent analyses, as in Sourav et al. (2020).

Data Reuse

The dataset in the current study has been derived from Sourav et al. (2020). The only difference is the exclusion of one participant with a very short duration of visual deprivation prior to the commencement of data analysis (1 month in participant CC – 008; see supplemental materials for Sourav et al., 2020). Parts of the dataset have been included in two other publications (Sourav et al., 2018, 2024).

Time-Frequency Analysis

Time-frequency representations of the EEG data were derived by wavelet transformation of individual trials at each electrode using the *FieldTrip* software in MATLAB (Oostenveld et al., 2011). Signals at each electrode were convolved with complex Morlet wavelets which have a Gaussian shape over time (with standard deviation σ_t) as well as over frequency (with standard deviation σ_f), centered around a time t_0 and a frequency f_0 (Tallon-Baudry et al., 1999). Non-constant wavelets were applied with a ratio $f_0/\sigma_f = 4$ for the lowest frequency and increasing linearly up to 8.5 for the highest, identical to Bottari et al. (2016, 2018). Thereafter, the relative power change compared to baseline (ΔP_{rel}), and the inter-trial phase coherence (ITPC) across trials were computed at each frequency from 2 to 40 Hz with a step size of 1 Hz, and from 700 ms before to 700 ms after stimulus onset in steps of 20 ms. ΔP_{rel} was derived by normalizing the wavelet power with respect to a baseline interval from -700 ms to -300 ms, i.e., $\Delta P(t,f)_{rel} = [P(t,f) - P_{baseline}(f)]/P_{baseline}(f)$, see Bottari et al. (2016, 2018). The ITPC is a measure of oscillatory phase synchronization across trials at a given timepoint and frequency (Delorme & Makeig, 2004) and was derived by calculating the complex average of the normalized phase directions. The ITPC is a measure that can take on values in the interval [0, 1], where ITPC = 1 indicates perfect synchronization across trials and ITPC = 0 indicates complete desynchronization. Importantly, the ITPC is a biased but consistent statistical measure, e.g., for uniformly random, desynchronized phase values with a finite number of trials, the ITPC is expected to be positive (i.e., is overestimated) but approaches 0 as the number of trials is increased (Aydore et al., 2013). To unbias the ITPC estimates for group comparisons, and to derive mutual magnitude-phase transplants as described in a following section, we used an

247   equal number of EEG epochs for each cataract reversal participant and their control

248   participant.

Cluster Based Permutation Tests

The ΔP_{rel} and ITPC values were compared between each cataract reversal group and their

sighted controls (i.e., CC vs. MCC; DC vs. MDC) using two-tailed cluster-based permutation

tests. The tests were performed with the *FieldTrip* framework (Oostenveld et al., 2011) running

on MATLAB (R2022a, Natick, MA), over the post-stimulus time-frequency plane (*t*: 0 – 700 ms;

f: 2- 40 Hz) and over the 13 electrodes posterior to the midline, where the P1 wave is most

prominent. The cluster-based permutation tests were based on 10,000 randomizations and

used weighted cluster mass as the test metric.

After visual stimulus presentation, a task-related decrease of ΔP_{rel} ("desynchronization") is

typically observed in the alpha (8 – 12 Hz) frequency range (Klimesch, 2012). Based on a

previous report by Bottari et al. (2016) that this task-related post-stimulus decrease of ΔP_{rel} in

the alpha frequency range (8 – 12 Hz) might differ in the CC group, we additionally compared

ΔP_{rel} in each sight recovery group to their control group (i.e., CC vs. MCC; DC vs. MDC) with

cluster-based permutation tests, with the same parameters as above, in the "alpha

desynchronization" time-frequency window (time: 0.3 – 0.7 s, frequency: 8 – 12 Hz) . Here, we

underscore that the term *desynchronization* refers to microscopic, spatial, inter-generator

desynchronization, operationalized as a decreased scalp-recorded oscillatory power (and not

as a mesoscopic, temporal, inter-trial phase coherence).

Classification Based on Time-Frequency Features

In addition to statistical tests for group differences in time-frequency representations, we tested

group discriminability of CC vs. Non-CC (i.e., DC and MCC) and CC vs. DC individuals in a

classification paradigm based on the time-frequency features, separately for ΔP_{rel} and ITPC.

Classification was performed based on oscillatory activity from 2 to 40 Hz in steps of 1 Hz (i.e.,

in 39 frequency bins) and from 100 ms before stimulus onset to 500 ms after stimulus onset,

with 20 ms step size (i.e., 31 temporal bins) at all 13 posterior electrodes, yielding 15717

classification features for both ΔP_{rel} and ITPC. To deal with the high dimensionality of the data,

we employed principal component analysis (PCA) prior to classification (Buzzell et al., 2022).

Using the principal components' loadings, participants were then classified using linear support

vector machines (SVM). Data from experiment 1 were used for deriving the principal

components, for tuning the SVM hyperparameters, and for subsequent SVM training. In

contrast, data from experiment 2 were utilized to validate the previously trained SVMs. The

analysis was performed in R, version 4.0.3 (R Core Team, 2016).

Dimension Reduction of the Time-Frequency Features

The ΔP_{rel} and ITPC values at each frequency and timepoint at the 13 posterior electrodes were

used to derive principal components for ΔP_{rel} and ITPC separately. PCA was performed across

all participants of experiment 1 (including the normally sighted control individuals), without

centering or scaling the ΔP_{rel} and ITPC features, respectively. We chose the first three principal

components for classification which explained 84.84% of the variance of ITPC and 49.21% of

the variance of ΔP_{rel}. Importantly, the PCA components do not necessarily reflect components

with the highest *intergroup* variance, rather they reflect the highest *interindividual* variance

regardless of the group membership. According to the suggestion by Buzzell et al. (2022),

varimax rotation was first applied to the selected components before computing the single

participants' loadings at each electrode, yielding 39 ΔP_{rel} and 39 ITPC features per participant

(13 electrodes × 3 principal component loadings). The varimax-rotated components are

displayed in Fig. 1.

To assess the robustness of potential group differences, classification was additionally

performed based on loadings of each of the first three components averaged over all posterior

electrodes, yielding 3 classification features for both ΔP_{rel} and ITPC. The principal components

were solely derived from data of experiment 1 and the same components were used for

classification of the data from experiment 2.

Figure 1. Principal component loadings of relative power change and of ITPC.

a – c. The first three principal components of the relative power change compared to baseline (ΔP_{rel}), summed across all posterior electrodes for each time-frequency feature (unitless). **d – e**. The first three principal components of inter-trial phase coherence (ITPC), summed across all posterior electrodes for each time-frequency feature (unitless). Each pixel in each plot represents one time-frequency feature. Summation over all electrodes is performed to highlight overall positive and negative loading clusters for each component.

Support Vector Machine Configuration

Using the features derived from the PCA-based dimensionality reduction, the regularizing hyperparameter c of the SVM was tuned in a repeated cross-validation (CV) procedure, with a split of $k = 5$ (CC vs. non-CC classification) and $k = 3$ (CC vs. DC classification). The CV procedure was repeated 500 times, and the optimal c that was chosen most often of all repetitions was used to fit a final SVM to all data from experiment 1. For each feature class (i.e., ΔP_{rel} and ITPC), each classification type (CC vs. non-CC, and CC vs. DC), an SVM was trained once using all 39 component loadings from experiment 1 and the most often chosen optimal value of c, summing to 4 training procedures (2 feature classes, × 2 classification types). This procedure was repeated for the component loadings averaged over the posterior electrodes, yielding 4 additional training procedures each only using 3 classification features. For each SVM, a receiver operating characteristic (ROC) analysis was conducted to extract the optimal classification threshold by maximizing the *Youden's J* parameter. To validate each trained classifier, the decision boundary derived from experiment 1 was used to classify the participants in experiment 2. ROC analyses were conducted for each classifier and the

corresponding areas under the ROC curves (AUCs) were extracted. Subsequently, the

significance of the AUCs was determined using the Mann-Whitney U-Statistic (Mason and

Graham, 2002), and the resulting *p*-values were adjusted using the Benjamini and Hochberg

procedure (Benjamini and Hochberg, 1995). The sensitivity and specificity of each

classification procedure was obtained through the optimal thresholds derived from experiment

1.

Mutual Magnitude-Phase Transplants

The discrete Fourier transform decomposes a signal (e.g., scalp-recorded voltage at an

electrode) into a set ("spectrum") of complex sinusoidal oscillations, each possessing a

different frequency. A constituent oscillation has a magnitude, indicating the oscillation

strength, as well as a phase, indicating its relative position in time or space (Grami, 2016). The

magnitude and phase spectra of a signal generally carry distinct information about the signal.

For example, in an image, most of the global color information is held in the magnitude

spectrum of the color channels, whereas the phase spectrum contains most of the form

information (Weinhaus, 2011). It is well known that swapping the magnitude and phase

information between two images creates chimeric images that generally inherit the color

palette from the magnitude spectrum and the form information from the phase spectrum (Fig.

2a). To probe the relative contribution of signal strength vs. coherent timing to the P1 wave

reduction observed in CC individuals, we adapted this signal processing method for event-

related potentials at the level of individual trials. Here we briefly describe this method, which

we term *mutual magnitude-phase transplants* (MuTx) in the present context.

Event-related potentials (ERP) are derived by averaging across numerous trials (Fig. 2 b – c).

For each CC individual and their matched normally sighted control participant (MCC), we

derived the MuTx waves from a set balanced number of trials, as in all the other analyses

reported in this article. First, for both individuals, we applied a Hann window to each individual

trial and calculated the discrete Fourier transform with the FFT method in Matlab (R2022a,

Natick, MA). Then, the magnitude and the phase spectra of the FFT was calculated, e.g.,

348
$$x_i(t) \xrightarrow{FFT} X_i(\omega) := |X_i(\omega)|e^{j \cdot \angle X_i(\omega)}$$

349
$$y_i(t) \xrightarrow{FFT} Y_i(\omega) := |Y_i(\omega)|e^{j \cdot \angle Y_i(\omega)}$$

350 where $|X_i(\omega)|$ and $\angle X_i(\omega)$ are the magnitude and phase spectra of the i^{th} trial of the CC

351 individual, $|Y_i(\omega)|$ and $\angle Y_i(\omega)$ are the magnitude and phase spectra of the i^{th} trial of the MCC

352 individual, and j is the imaginary unit, i.e. $j^2 = -1$.

353 After this step, the phase of the CC and the MCC individuals were exchanged, and an inverse

354 fast Fourier transform (IFFT) yielded two MuTx trials (Fig. 2 d – e):

355
$$|X_i(\omega)|e^{j \cdot \angle Y_i(\omega)} \xrightarrow{IFFT} |x_i| \angle y_i(t)$$

356
$$|Y_i(\omega)|e^{j \cdot \angle X_i(\omega)} \xrightarrow{IFFT} |y_i| \angle x_i(t)$$

The first of the two MuTx trials, $|x_i| \angle y_i(t)$, had the magnitude information from the CC

individual's trial, but the phase information of the MCC individual's trial. Conversely, the second

MuTx trial, $|y_i| \angle x_i(t)$, had the magnitude of the MCC individual's trial but the phase of the CC

individual's trial. Averaging across these trials provided two MuTx ERPs, which we term

$|CC|\angle MCC$ and $|MCC|\angle CC$. Here, $|CC|\angle MCC$ is the ERP derived from the trials with the

magnitude of CC individuals and the phase of MCC individuals. Conversely, $|MCC|\angle CC$ had

the magnitude of MCC individuals and the phase of CC individuals. If a generally lower

activation magnitude across trials would be the main driver of the P1 reduction in CC

individuals, we would expect $|CC|\angle MCC$ to have lower P1 values than $|MCC|\angle CC$. In contrast,

if the P1 wave reduction would be driven mainly by a reduced phase alignment (i.e.,

coherence) in the CC individuals, then $|MCC|\angle CC$ would exhibit a reduced P1 wave compared

to the $|CC|\angle MCC$.

The role of magnitude and phase information in a constellation of sight recovery group and

their control participants (i.e., CC vs. MCC or DC vs. MCC) was investigated using the mean

of the standardized P1 wave activation (i.e., *z*-score) over the 13 posterior electrodes (120 –

170 ms post-stimulus at electrodes TP9/TP10, CP1/CP2, CP5/CP6, P7/P8, P3/P4, Pz, and

O1/O2) as the dependent variable (Sourav et al., 2020).

Figure 2. Derivation of mutual magnitude-phase transplants (MuTx).

a An image processing example motivating the MuTx approach. Two-dimensional fast Fourier transform (FFT) of
signals, here public-domain images of a house (Weber, 2018) and an imaginary animal (Roy, 1946), decomposes
each image into a set of sinusoidal spatial oscillations (here, in two spatial direction *u* and *v*). Each oscillation (i.e.
each member of the set of sinusoidal oscillations derived by FFT) have a magnitude, indicating the strength of
oscillation, as well as a phase, indicating its relative position. As an experiment, the magnitude and phases of the
two images can be exchanged. After this interchange, inverse fast Fourier transforms (IFFT) yield the mutual
magnitude-phase transplants (MuTx). It can be appreciated that for images, most of the shape information seems
to be contained in the signal phase, whereas color information seems to rely more on the signal magnitude. **b – c**
Derivation of magnitude and phase information via one-dimensional FFT of the individual trials of the EEG epochs
respectively in a CC individual and their matched normally sighted control participant (MCC), with the mean across
trials (unaltered event related potentials, ERPs) displayed. **d – e** After exchanging the magnitude and phase spectra,
the inverse FFT yields two MuTx waves with the magnitude of the CC individual but the phase of the MCC individuals
($|CC|\angle MCC$) and vice versa ($|MCC|\angle CC$). It can be appreciated that phase information seems to drive most of the
P1 magnitude difference (grey bar, $120 - 170$ ms post-stimulus) between the CC and the MCC participant.

The *m*ean *p*osterior *P*1 activation, termed MPP1, was investigated in Bayesian linear models

of the form `MPP1 ~ magnitude × phase`. The factors *magnitude* and *phase* had two levels

each, which indicated the source group for the magnitude or phase information, namely *sight*

recovery and *control*. Separate models were run for the CC vs. MCC and the DC vs. MDC

comparisons.

In addition to this model, we used a reparametrized model with custom contrasts (Schad et al.,

2020) to test for three differences of interest. For the CC vs. MCC comparison, the first of these

differences tested whether the CC group exhibited a P1 wave reduction (contrast: MCC – CC).

The other two contrasts tested the effect of "transplanted" phase on the P1 wave amplitude

(contrasts: |CC|∠MCC – CC, and |MCC|∠CC – MCC). Analogous models were run for the DC

vs. MDC comparison. In all cases, weak priors were set with the auto_prior() function of the

sjstat library (Lüdecke, 2018).

P1 Wave Source Analysis

Average-referenced ERP voltage topographies were converted to a 15,002-point,

unconstrained cortical source space using the sLORETA method for EEG inverse modeling

(Pascual-Marqui, 2002), implemented in the Brainstorm software package (Tadel et al., 2011).

For calculating the forward model used in the sLORETA modeling, we started with the *New*

York Head Model with six tissue types: scalp, skull, cerebrospinal fluid, gray matter, white

matter, and air cavities (Huang et al., 2016). The forward model was then calculated with the

DUNEuro partial differential equation solver package using realistic conductivity values for the

tissues and with the electrode locations in the present study (Piastra et al., 2021; Schrader et

al., 2021). An identity matrix was used as the noise covariance.

Similar to the P1 wave analysis at the sensor level, group comparisons (CC vs. MCC, DC vs.

MDC) of source activity were performed in a posterior cortical region of interest that included

the occipital, temporal and parietal regions, excluding the postcentral sulcus and the limbic

lobe. To this end, the mean sLORETA activation across the posterior region was extracted,

with automatic sign inversion implemented in the Brainstorm package for oppositely oriented

sLORETA solutions. The vector norm, i.e. magnitude of the mean sLORETA activation, served

as the dependent variable for a Bayesian lognormal model of the form `norm_of_mean ~`

`Group`; weakly informative priors were set for the intercept and groups, respectively $N(0, 10)$

and N(0, 2.5). For evidence synthesis across experiments, we added a random intercept of

participants to take participation in both experiments into account, i.e., `norm_of_mean ~ Group`

`+ (1|ParticipantID)`. Substantial evidence was ascertained when Bayes factors exceeded

3 and when the 95% highest-density interval of the posterior was situated outside of a region

of practical equivalence of [-0.1, 0.1] (ROPE test, Makowski et al., 2019).

Results

Time-Frequency Analysis

Experiment 1

ΔP_{rel}. The cluster-based permutation test in the whole post-stimulus time-frequency range

(latency: 0 – 700 ms, frequency: 2 – 40 Hz) did not reveal any significant difference in ΔP_{rel}

between the cataract reversal groups and their respective control groups (Fig. 3 a – f); CC vs.

MCC, most negative cluster mass (± 95% confidence interval, CI): -369.014 ± 0.006, *p* = .118,

most positive cluster mass: 28.399, the parametric threshold for clustering was not exceeded;

DC vs. MDC, most negative cluster mass: -241.928 ± 0.009, *p* = .293, most positive cluster

mass: 37.595, the parametric threshold for clustering was not exceeded.

Running cluster-based permutation in the "alpha desynchronization" time-frequency window

(time: 0.3 – 0.7 s, frequency: 8 – 12 Hz) as in Bottari et al. (2016), we once more did not find

evidence for a difference in ΔP_{rel} in the CC vs. MCC analysis (most negative cluster mass =

-29.842 ± 0.009, *p* = .304, most positive cluster mass = 0.043, threshold for clustering not

exceeded), nor in the DC vs. MDC comparison (most negative cluster mass = -6.959 ± 0.010,

p = .609, for positive cluster formation no time-frequency point significant at individual level).

Inter-trial Phase Coherence (ITPC). In the CC group, significant evidence for a lower ITPC

compared to the matched typically sighted control group (MCC) emerged (Fig. 3 g – l); most

negative cluster mass (± 95% CI): -405.176 ± 0.003, *p* = .025, most positive cluster mass (±

95% CI): 93.893 ± 0.010, *p* = 0.422. In contrast, between the DC and the MDC group, we did

not observe a significant difference in ITPC; most negative cluster mass (± 95% CI): -32.622,

the parametric threshold for clustering was not exceeded, most positive cluster mass (± 95%

CI): 120.391 ± 0.008, *p* = .226.

Figure 3. Time-Frequency Analysis of Relative Power Change and Phase Coherence at Posterior Electrodes in Experiment 1.

a Relative power change (ΔP_{rel}, unitless) compared to baseline (-700 to -300 ms) in the matched control group for congenital cataract reversal individuals (MCC). **b** ΔP_{rel} in the group of congenital cataract reversal individuals (CC). **c** Relative power change difference between the MCC and the CC group, $\Delta P_{rel, MCC} - \Delta P_{rel, CC}$. **d – f**. Relative power changes in the matched control group for developmental cataract reversal individuals (MDC, $\Delta P_{rel, MDC}$), in the group of developmental cataract reversal individuals (DC, $\Delta P_{rel, DC}$), and their difference ($\Delta P_{rel, MDC} - \Delta P_{rel, DC}$). **g – i**. Inter-trial phase coherence (ITPC) respectively in the MCC group, the CC group, and their difference. **j – l**. ITPC respectively in the MDC and the DC group, and their difference. For cluster-based permutation tests spanning the time-frequency range of [0 – 0.7 s] × [2 – 40 Hz], the time-frequency extend of a cluster driving a significant result has been outlined. The vertical dotted lines at 0 s indicate stimulus onset. ΔP_{re} and ITPC shown in the plots are averaged over the 13 posterior electrodes (TP9/TP10, CP1/CP2, CP5/CP6, P7/P8, P3/P4, Pz, and O1/O2). Note that ΔP_{rel} and ITPC have different scales (respectively -0.8 – 0.8 and -0.05 – 0.3). For each variable, the difference between the groups is displayed on a scale that is half of that used for the variable.

Experiment 2

ΔP_{rel}. The cluster-based permutation test in the whole post-stimulus time-frequency range

(latency: 0 – 700 ms, frequency: 2 – 40 Hz) once again did not find any significant difference

in *ΔP_{rel}* between the CC and the MCC group (Fig. 4 a – f), most negative cluster mass (± 95%

confidence interval, CI): -39.6720, the parametric threshold for clustering was not exceeded;

most positive cluster mass = 155.132 ± 0.010, *p* = .443. In contrast to the CC vs. MCC group

comparison (and in contrast to experiment 1), we found a reduced *ΔP_{rel}* in the DC group

compared to the MDC group, most negative cluster mass = -863.053 ± 0.004, *p* = .040; most

positive cluster mass = 29.512, parametric threshold for clustering not exceeded.

Employing cluster-based permutation in the alpha desynchronization time-frequency window

(time: 0.3 – 0.7 s, frequency: 8 – 12 Hz) as in Bottari et al. (2016), as in experiment 1, no

evidence of a difference in *ΔP_{rel}* emerged between the CC and the MCC group (for negative

cluster formation, no time-frequency point significant at individual level; most positive cluster

mass = 40.989 ± 0.008, *p* = .197).

Inter-trial Phase Coherence (ITPC). In the CC group, significant evidence for a lower ITPC

compared to the matched sighted control group (MCC) once again was found in experiment 2

(Fig. 4 g – l); most negative cluster mass (± 95% CI): -233.112 ± 0.003, *p* = .031, most positive

cluster mass = 28.907, parametric threshold for clustering not exceeded. Similar as in

experiment 1, we did not observe a significant difference in ITPC between the DC and the

MDC groups; most negative cluster mass (± 95% CI) = -106.505 ± 0.009, *p* = .266; most

positive cluster mass = 28.672, parametric threshold for clustering not exceeded.

Figure 4. Time-Frequency Analysis of Relative Power Change and Phase Coherence at Posterior Electrodes in Experiment 2.

a Relative power change (ΔP_{rel}, unitless) compared to baseline (-700 to -300 ms) in the matched control group for congenital cataract reversal individuals (MCC). **b** ΔP_{rel} in the group of congenital cataract reversal individuals (CC). **c** Relative power change difference between the MCC and the CC group, $\Delta P_{\text{rel, MCC}} - \Delta P_{\text{rel, CC}}$. **d – f**. Relative power changes in the matched control group for developmental cataract reversal individuals (MDC, $\Delta P_{\text{rel, MDC}}$), in the group of developmental cataract reversal individuals (DC, $\Delta P_{\text{rel, DC}}$), and their difference ($\Delta P_{\text{rel, MDC}} - \Delta P_{\text{rel, DC}}$). **g – i**. Inter-trial phase coherence (ITPC) respectively in the MCC group, the CC group, and their difference. **j – l**. ITPC respectively in the MDC and the DC group, and their difference. For cluster-based permutation tests spanning the time-frequency range of [0 – 0.7 s] × [2 – 40 Hz], the time-frequency extend of a cluster driving a significant result has been outlined. The vertical dotted lines at 0 s indicate stimulus onset. ΔP_{re} and ITPC shown in the plots are averaged over the 13 posterior electrodes (TP9/TP10, CP1/CP2, CP5/CP6, P7/P8, P3/P4, Pz, and O1/O2). Note that ΔP_{rel} and ITPC have different scales (respectively -0.8 – 0.8 and -0.05 – 0.3). For each variable, the difference between the groups is displayed on a scale that is half of that used for the variable.

Classification Based on Time-Frequency Features

In addition to the cluster-based permutation tests for group differences in spectral dynamics, we investigated whether CC individuals could be classified based on biomarkers in the time-frequency domain. To prevent overfitting in the high-dimensional spectral feature space, the classifications were performed based on the loadings of the first three principal components at

posterior electrodes (39 features), as well as based on the average loadings across the

posterior electrodes (3 features). In total, 8 classifications were performed: 2 classification

types (CC vs. non-CC, and CC vs. DC) × 2 feature classes (ΔP_{rel} and ITPC) × 2 loading types

(loadings at all posterior electrodes and loadings averaged over posterior electrodes). All

classifiers were trained on data from experiment 1 and validated on data from experiment 2.

Validation results are all displayed in Table 5, and the corresponding ROC curves are depicted

in Fig. 6, panels a – h. Additionally, the SVM decision boundaries for classification based on

ITPC- and power-based component loadings averaged across posterior electrodes are

visualized in Fig. 6, panels i – j in the three-dimensional feature space.

Table 5.

Principal Component Based Classification.

CC vs. non-CC classification

Domain	*n*(Features)	AUC	[95% CI]	*p*-value	Sensitivity	Specificity
Power	39	.660	[.499; .821]	.084	.462	.814
Power	3	.528	[.362; .694]	.774	.538	.558
ITPC	39	.748	[.575; .921]	.006*	.692	.767
ITPC	3	.766	[.624; .907]	.003*	.462	.814

517 *CC vs. DC classification*

Domain	*n*(Features)	AUC	[95% CI]	*p*-value	Sensitivity	Specificity
Power	39	.718	[.521; .915]	.052	.769	.467
Power	3	.621	[.397; .844]	.235	.692	.400
ITPC	39	.718	[.509; .927]	.052	.692	.800
ITPC	3	.728	[.536; .921]	.041*	1.000	.200

CC vs. non-CC Classification

Classification of CC participants from non-CC participants (including DC and normally sighted

control individuals) in experiment 2 achieved statistically significant performance based on

ITPC component loadings at all posterior electrodes (AUC = .748, 95% CI = [.575 – .921], p =

.006), but not based on the component loadings derived from ΔP_{rel} (AUC = .660, 95% CI =

[.499 – .821], p = .084). Interestingly, a robust performance was observed as well when

classification was based on the ITPC component loadings averaged over all posterior

electrodes, i.e., based on 3 instead of 39 features (AUC = .766, 95% CI = [.624– .907], p =

.003). In contrast, ΔP_{rel}-based classification performance dropped close to chance level when

classifying employed average loadings (AUC = .528, 95% CI = [.362– .694], p = .774). The

visualization of the corresponding three-dimensional SVM decision boundaries demonstrates

that only the phase-based group differences generalize from the training (experiment 1) to the

validation dataset (experiment 2, see Fig. 6 i – j). The optimal thresholds derived from

experiment 1 translated to experiment 2 to a moderate degree (Power et al., 2013).

CC vs. DC Classification

Classification of the CC from the DC individuals in the validation experiment 2 was not

statistically significant using either the ITPC component loadings (AUC = .718, 95% CI = [.509

– .927], p = .052) or the ΔP_{rel} component loadings at all posterior electrodes (AUC = .718, 95%

CI = [.521 – .915], p = .052). However, when averaging the component loadings across the

posterior electrodes, ITPC information successfully classified the CC from the DC individuals

(AUC = .728, 95% CI = [.536 – .921], p = .041). In contrast, using average component loadings

for ΔP_{rel}, the classification performance remained close to chance level (AUC = .621, 95% CI

= [.397 – .844], p = .294).

Phase Information is the Main Driver of the P1 Wave Reduction After Sight Recovery Following Transient Congenital Blindness

In a novel application, we exchanged the magnitude and phase information between each

sight-recovery individual and their control participant. ERPs were derived from the trials of the

mutual magnitude-phase transplanted participants (MuTx) and were systematically compared

to those of the original participants (Fig. 6 – 7). We reasoned that if phase information were

the main driving factor behind the P1 wave reduction observed in the CC group, the MuTx

group comprising the magnitude information of the MCC individuals but the phase information

of the CC individuals (denoted |MCC|∠CC) would exhibit a P1 wave reduction. Conversely, the

MuTx group consisting of chimeric participants with the magnitude information from the CC

individuals but the phase information from the MCC individuals (denoted |CC|∠MCC) would be

expected to exhibit typical P1 waves. This was indeed the case: in experiment 1, the grand

average visual ERPs qualitatively resembled the group whose phase was used to derive the

visual ERP (Fig. 6a). This held true too for the P1 wave topography: in the |CC|∠MCC group's

Figure 5. Receiver Operating Characteristic (ROC) Curves for Principal Component Based Classification.

a-d ROC curves for the classification of CC vs. non-CC individuals based on component loadings for relative power change (ΔP_{rel}) and ITPC features at all posterior electrodes ('All Loadings') and loadings averaged over posterior electrodes ('Mean Loadings'). Black dots indicate the optimal classification threshold derived from experiment 1. **e-f** ROC curves for the CC vs. DC classification using the same features. **i** The SVM decision boundary for CC vs. non-CC classification based on ITPC component loadings averaged over posterior electrodes (i.e., 3 features) on the training and validation dataset. **j** The SVM decision boundary for CC vs. Non-CC classification based on component loadings averaged over posterior electrodes for ΔP_{rel} on the training and validation dataset.

P1 wave topography resembled that of the MCC group, and the |MCC|∠CC group's topography resembled that of the CC group (Fig. 6b). Finally, the Bayesian mixed model investigating the effect of oscillatory magnitude and phase indicated that the mean of the standardized P1 values over posterior electrodes (MPP1) depended on the phase, but not magnitude information content for the CC, MCC, |CC|∠MCC, and |MCC|∠CC constellation; there was a decisive main effect of phase, β_{phase} [95% credible interval] = 0.37 [0.23, 0.52], BF_{10} = 1478.71, positive ROPE test. In contrast, there was no substantial evidence for the role of magnitude influencing the MPP1 values in this constellation, β_{mag} = 0.01 [-0.14, 0.15], BF_{10} = 0.047, and

572   no substantial interaction between the factors magnitude and phase, $\beta_{mag \times phase}$ = -0.01 [-0.15,

573   0.14], BF$_{10}$ = 0.048.

Figure 6. P1 waves of the original groups vs. the mutual magnitude-phase transplant (MuTx) groups in Experiment 1.

a Grand average event-related potentials (ERP) at the posterior electrode P8 for the sight recovered bilateral congenital cataract group (CC), their matched control group (MCC), and the MuTx groups with the magnitude (|○|) and phase (∠○) information swapped, i.e. |CC|∠MCC and |MCC|∠CC. The color-coding is according to the group's magnitude information. Ribbons indicate the standard error of the mean. The gray bar indicates the P1 wave time range (120 – 170 ms). **b** Topographic plot of the grand average P1 wave for the CC, MCC, |CC|∠MCC, and |MCC|∠CC groups, averaged between 120 – 170 ms after stimulus presentation. **c** Grand average ERPs at the posterior electrode P8 for the developmental cataract reversal group (DC), their matched normally sighted control group (MDC), and the MuTx groups |DC|∠MDC and |MDC|∠DC. **d** Topographic plot of the grand average P1 wave for the DC, MDC, |DC|∠MDC, and |MDC|∠DC groups, averaged between 120 – 170 ms after stimulus presentation. **e** Violin plot of the means of the standardized P1 calculated over the posterior electrodes. The mean posterior P1 value (MPP1) had been validated as a retrospective biomarker of congenital visual deprivation (Sourav et al., 2020). The threshold for diagnosing a CC individual is shown as a horizontal dotted line, and the number of individuals above and below the diagnosis line is indicated near the top and bottom border. Red circles denote the means, and the error bars indicate 95% bootstrapped confidence intervals for the means. Asterisks indicate substantial evidence (or above). Individual data points were jittered horizontally for readability.

The same model equipped with custom contrasts provided very strong evidence that the MPP1

values of the CC group was lower than the MPP1 values of the MCC group, β = -0.74 [-1.13, -

0.34], BF$_{10}$ = 56.123, positive ROPE test. In addition, the |CC|∠MCC group had a higher MPP1

value than the CC group, β = 0.75 [0.34, 1.15], BF$_{10}$ = 71.263, positive ROPE test. Finally,

there was very strong evidence that |MCC|∠CC group's MPP1 values were lower compared to

the MPP1 values of the MCC group, β = -0.71 [-1.11, -0.31], BF$_{10}$ = 48.557, positive ROPE

test, indicating that difference in phase information likely caused the P1 wave attenuation in

the CC group.

These results were replicated in the validation dataset from experiment 2, both qualitatively

(Fig. 7 a – d) as well as quantitatively (Fig. 7e). Specifically, a Bayesian mixed model once

Figure 7. P1 waves of the original groups vs. the mutual magnitude-phase transplant (MuTx) groups in Experiment 2 (Validation Dataset)

a Grand average visual event-related potentials (VERP) at the posterior electrode P8 for the sight recovered bilateral congenital cataract group (CC), their matched control group (MCC), and the MuTx groups with the magnitude (|○|) and phase (∠○) information swapped, i.e. |CC|∠MCC and |MCC|∠CC. The color-coding is according to the group's magnitude information. Ribbons indicate the standard error of the mean. The gray bar indicates the P1 wave time range (120 – 170 ms). **b** Topographic plot of the grand average P1 wave for the CC, MCC, |CC|∠MCC, and |MCC|∠CC groups, averaged between 120 – 170 ms after stimulus presentation. **c** Grand average VERPs at the posterior electrode P8 for the typically sighted recovered bilateral developmental cataract group (DC), their matched control group (MDC), and the MuTx groups |DC|∠MDC and |MDC|∠DC. **d** Topographic plot of the grand average P1 wave for the DC, MDC, |DC|∠MDC, and |MDC|∠DC groups, averaged between 120 – 170 ms after stimulus presentation. **e** Violin plot of the mean of the standardized P1 topography, with the mean calculated over the posterior electrodes. The mean posterior P1 value (MPP1) was developed and validated as a biomarker based on the present experiments (Sourav et al., 2020). The threshold for diagnosing a CC individual (from experiment 1) is shown as a horizontal dotted line, and the number of individuals above and below the diagnosis line is indicated near the top and bottom border. Red circles denote the means, and the error bars indicate 95% bootstrapped confidence intervals for the means. Asterisks indicate substantial evidence (or above). Individual data points were jittered horizontally for readability.

again indicated that the MPP1 values for the CC, MCC, |CC|∠MCC, and |MCC|∠CC

constellation depended on the phase but not magnitude information; there was a decisive main

effect of phase, β_{phase} [95% credible interval] = 0.27 [0.14, 0.40], BF_{10} = 117.02, positive ROPE

test. In contrast, there was no substantial evidence for the role of magnitude influencing the

MPP1 values in this constellation, β_{mag} = 0.06 [-0.07, 0.19], BF_{10} = 0.075, and no substantial

interaction between the factors magnitude and phase, $\beta_{mag \times phase}$ = -0.01 [-0.14, 0.12], BF_{10} =

0.049.

The same model equipped with custom contrasts provided very strong evidence that the MPP1

values of the CC group were lower than those of the MCC group, β = -0.65 [-1.01, -0.28], BF_{10}

= 39.909, positive ROPE test. In addition, there was substantial evidence that the |CC|∠MCC

group had higher MPP1 values than the CC group, β = 0.55 [0.18, 0.91], BF_{10} = 9.893, positive

ROPE test. Finally, there was substantial evidence that the |MCC|∠CC group's MPP1 values

were lower compared to those of the MCC group, β = -0.52 [-0.88, -0.15], BF_{10} = 6.836, positive

ROPE test.

When we tested the DC, MDC, |DC|∠MDC, and |MDC|∠DC group constellation, we found no

evidence in any of the two experiments that the MPP1 values substantially differed between

the groups (all non-intercept βs had BF_{10} < 1, no positive ROPE tests). Unsurprisingly, there

was no substantial evidence that the MPP1 depended on phase or the magnitude information

in this constellation (all non-intercept BF_{10} < 1, no positive ROPE tests).

Source Localization of the P1 wave

In experiment 1, the Bayesian lognormal model indicated that the magnitude of mean posterior

sLORETA activation substantially differed between the CC and the MCC groups (Fig. 8), $\beta_{CC\text{-}}$

$_{MCC}$ [95% CrI] = -1.03 [-1.76, -0.30], BF_{10} = 6.19, positive ROPE test. In contrast, no substantial

difference in magnitude of mean posterior sLORETA activations was observed when

comparing the DC vs. the MDC group; $\beta_{DC\text{-}MDC}$ = -0.11 [-0.63, 0.41], BF_{10} = 0.111, negative

ROPE test (3.06 % of posterior distribution in the ROPE region).

In experiment 2, we did not find any substantial evidence that the magnitude of mean posterior

sLORETA activation substantially differed between the CC and the MCC groups (see Fig. 9),

$\beta_{CC\text{-}MCC}$ [95% CrI] = -0.58 [-1.19, -0.03], BF_{10} = 0.772, negative ROPE test (0.46% of posterior

distribution in the ROPE region). No substantial difference in magnitude of mean posterior

sLORETA activations was observed when comparing the DC vs. the MDC groups as well; $\beta_{DC\text{-}}$

$_{MDC}$ = -0.11 [-0.63, 0.41], BF_{10} = 0.111, negative ROPE test (3.06 % of posterior distribution in

the ROPE region).

Figure 8. Source Localization of the P1 Wave in Experiment 1.

a Magnitude (vector norm) of the grand average source localization solution for the P1 wave (mean over 120 – 170 ms post-stimulus) obtained with the sLORETA method (unitless; Pascual-Marqui, 2002; Tadel et al., 2011) respectively for the CC and the MCC groups (left and center columns), and the MCC – CC group difference of grand average norms (right column). **b** Magnitude of the grand average sLORETA solution for the P1 wave (mean over 120 – 170 ms post-stimulus) respectively for the DC and the MDC groups (left and center columns), and the MDC – DC group difference of grand average norms (right column). **c** Posterior region of interest (ROI) for testing group differences, which included the occipital, temporal and parietal regions excluding the postcentral sulcus and the limbic lobe. **d** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the CC and the MCC group (scaled to unit variance). **e** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the DC and the MDC group (scaled to unit variance). Asterisk (*) indicates substantial evidence for a group difference (Bayes factor > 3 and positive ROPE test, lognormal model).

The Bayesian models, being parametric, allowed evidence synthesis across the experiments, which on the one hand led to a more stable parameter estimation and on the other hand permitted reconciling partially different result patterns for the CC vs. MCC group difference across the experiments. Combining evidence across both experiments with Bayesian lognormal mixed models, we found substantial evidence for a group difference in the mean posterior sLORETA activation strength between the CC and the MCC group, β_{CC-MCC} [95% CrI] = -0.77 [-1.26, -0.27], BF_{10} = 9.01, positive ROPE test (Fig. 9). In contrast, once again no substantial difference in the norm of mean posterior sLORETA activations emerged when comparing the DC vs. the MDC group; β_{DC-MDC} = -0.10 [-0.46   0.27], BF_{10} = 0.085, negative ROPE test (3.97 % of posterior distribution in ROPE region).

Figure 9. Source Localization of the P1 Wave in Experiment 2 and Evidence Synthesis Across Experiments 1 and 2.

a Magnitude (vector norm) of the grand average source localization solution for the P1 wave (mean over 120 – 170 ms post-stimulus) obtained with the sLORETA method in experiment 2 (unitless; Pascual-Marqui, 2002; Tadel et al., 2011) respectively for the CC and the MCC groups (left and center columns), and the MCC – CC group difference of grand average norms (right column). **b** Magnitude of the grand average sLORETA solution for the P1 wave (mean over 120 – 170 ms post-stimulus) respectively for the DC and the MDC groups in experiment 2 (left and center columns), and the MDC – DC group difference of grand average norms (right column). **c** Posterior region of interest (ROI) for testing group differences, which included the occipital, temporal and parietal regions excluding the postcentral sulcus and the limbic lobe. **d** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the CC and the MCC group in experiment 2 (scaled to unit variance). **e** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the DC and the MDC group in experiment 2 (scaled to unit variance). **f** Grand average source localization solution across all participants and groups in experiment 1 and experiment 2. Black borders indicate the loci of the highest 5% sLORETA magnitude. Relevant cortical regions from the Desikan-Killiany atlas are depicted on the left side of the sLORETA solution; FuG: Fusiform gyrus, LNG: Lingual gyrus, LOC: Lateral occipital cortex. **g** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the CC and the MCC group across experiment 1 and experiment 2, scaled to unit variance. All unique participants are depicted, for participants who took part in both experiments, mean values are presented. **h** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the DC and the MDC group across experiment 1 and experiment 2, scaled to unit variance. All unique participants are depicted, for participants who took part in both experiments, mean values are presented. Asterisk (*) indicates substantial evidence for a group difference (Bayes factor > 3 and positive ROPE test, lognormal model)

The grand average sLORETA map across all participants and both experiments indicated posterior regions as the source of the P1 wave (Fig. 9). The maximum 5% regions of this activation were located in the right ventral extrastriate visual cortical regions, including the fusiform gyrus (FuG), lingual gyrus (LNG) and lateral occipital cortex (LOC) according to the Desikan-Killany atlas (Desikan et al., 2006).

Discussion

A transient phase of congenital visual deprivation seems to permanently alter visual cortical structure and function (Röder et al., 2021; Hölig et al., 2023; Sourav et al., 2024). Research on experience-dependent brain development has emphasized the investigation of cortical activation strength in (visual) brain areas in permanent congenital blindness (Wanet-Defalque et al., 1988; Ossandón et al., 2023), or after sight recovery (Bottari et al., 2016; Sourav et al., 2020; Pitchaimuthu et al., 2021; Ossandón et al., 2023). Yet, to the best of our knowledge, no study in this group has systematically examined the trial-to-trial temporal consistency, a critical component of the spatiotemporal organization of visual processing.

We investigated group differences in task-related oscillatory activation strength and timing consistency using three methodological approaches. First, in time-frequency representations of cortical activity during the processing of simple visual stimuli, the across-trial consistency of visual processing, indexed by the inter-trial phase coherence (ITPC), was significantly and replicably attenuated in the CC group compared to their matched normally sighted controls (MCC group), while no differences in relative power change from baseline between these groups was observed. Second, further supporting these findings, the CC individuals could only be classified from non-CC individuals (i.e., a combined group of sighted controls as well as DC individuals) using phase but not using oscillatory power information. Moreover, the successful classification of CC individuals from non-CC individuals was replicated after averaging the ITPC values across the posterior electrodes as well, indicating that the phase information which discriminated CC individuals was expressed reliably over the posterior electrodes. Finally, through a novel approach that we term *mutual magnitude-phase transplants* (MuTx), we swapped the oscillatory activation magnitude and phase information between each CC individual and their matched sighted controls. The MuTx analysis revealed that combining the phase information of the CC individuals with the magnitude information of the MCC individuals, but not vice versa, yielded the CC-typical attenuation of the extrastriate P1 wave in visual ERPs (Figs. 6 – 7). The convergent evidence from all three methodological approaches across the

two experiments indicates that task-related oscillatory timing information, but not oscillatory

strength is persistently affected by a period of transient congenital blindness.

Between the DC individuals and their matched control group (MDC), cluster-based permutation

tests did not reveal significant ITPC differences in any experiment, indicating that the decrease

in task-related oscillatory phase synchronization in the CC vs. MCC group comparison is linked

to visual deprivation during a sensitive period. This selective difference for CC individuals

excludes alternative explanations such as unspecific effects related to cataract surgery or

testing location (i.e., India vs. Germany). Additionally, phase, but not power information

averaged over the posterior electrodes allowed us to classify CC from DC individuals. Lastly,

MuTx analyses revealed no effect of exchanging the magnitude and phase information

between the DC and the MDC individuals, highlighting the specific temporal processing

impairment in the CC group.

Compared to striate cortical processing, extrastriate cortical functions have been reported to

exhibit higher vulnerability to a period of transient congenital visual deprivation (McKyton et

al., 2015; Sourav et al., 2018, 2020; Pitchaimuthu et al., 2021). Consistent with this line of

evidence, we have previously found that the P1 wave in visual ERPs, which has been

hypothesized to originate in extrastriate cortical areas (Di Russo et al., 2002; Miller et al.,

2015), was strongly reduced in CC individuals (Sourav et al., 2020). Moreover, this strong and

reliable reduction of the P1 wave in sight recovery individuals allowed a robust retrospective

diagnosis of a congenital vs. developmental etiology years after sight restoration (Sourav et

al., 2020). In the present study, source localization of the P1 wave indicated areas of maximum

activation in extrastriate cortical areas in the right hemisphere, including the fusiform gyrus,

lingual gyrus and lateral occipital cortex, results that are by and large consistent with previous

EEG/fMRI work (Di Russo et al., 2002). Corroborating the finding of a sensor-level reduction

of the standardized posterior P1 wave amplitude, evidence synthesis across the experiments

at source level revealed a substantial evidence for a reduced sLORETA magnitude in the CC

group.

Phase alignment of alpha (8 – 12 Hz) oscillatory activity has been argued to contribute to the

generation of the P1 wave (Klimesch et al., 2007), supported by the findings that alpha phase

alignment predicted the P1 wave latency (Gruber et al., 2005), and that alpha phase locking is

largest during the P1 wave latency (Klimesch et al., 2004; see Freunberger et al., 2008).

Consistent with this evidence, the principal component loading patterns of the ITPC features

for classifying CC individuals (Fig. 1) were predominantly located across the alpha and the

theta (4 – 8 Hz) range and covered the P1 wave latency. Additionally, significant ITPC

differences between the CC and the MCC group in the cluster-based permutation tests were

associated with time-frequency clusters that spanned the P1 latency and the alpha frequency

range. In contrast, the P1 wave amplitude has been argued to be less affected by oscillatory

power than by phase alignment (Gruber et al., 2005; Freunberger et al., 2008). In a related

visual deprivation model of amblyopia, a reduction of the P1 wave amplitude in the patients'

amblyopic eye compared to the fellow eye was associated with a broader trial-to-trial latency

(but not amplitude) distribution (Bankó et al., 2013). Consistent with these lines of evidence,

we did not find any significant difference in the post-stimulus relative power change in the CC

vs. MCC group comparison and observed no significant classification performance based on

relative power change as opposed to ITPC information. In fact, the MuTx analysis in the

present study provided compelling evidence of the driving role of phase information behind P1

reduction in the CC group: combining CC individuals' activations with the phase information of

typically sighted controls substantially recovered their P1 wave amplitude.

Along with relative power increases in the delta (1 – 4 Hz) and theta range, a post-stimulus

decrease of relative power is usually observed in the alpha oscillatory band, termed *alpha*

desynchronization (Pfurtscheller et al., 1994). Previously, using biological motion stimuli in a

group of largely non-overlapping CC individuals from the our lab (*n* = 12*)*, Bottari et al. (2016)

reported a reduction in alpha desynchronization compared to typically sighted control

participants, positing that alpha oscillation strength might not recover after sight recovery. In

contrast, we did not find any significant difference in relative power in the CC group compared

to the MCC group in any of the two experiments (see Figs. 3 – 4), neither in the whole post-

stimulus time-frequency region, nor in a region of interest identical to Bottari et al. (2016)

specifically employed to investigate alpha desynchronization differences. However, the

divergent results of the present study and Bottari et al. (2016) are in line with the hypothesis

that a reduced alpha desynchronization in CC individuals might not be general and task-

independent (Klimesch et al., 2011), but may be limited to specific visual stimulus classes, e.g.,

motion processing. Interestingly, Ossandón et al (2023) have recently observed that in resting

state (i.e., task-free) EEG recordings, alpha desynchronization due to eye opening was

indistinguishable between their CC and MCC groups, despite an overall lower alpha oscillatory

power level in the CC group. The results of the present study indicate that in contrast to phase

coherence, task-related modulation of oscillatory activity strength recovers to a considerable

extent after sight restoration in CC individuals.

Temporal stability, and its converse, temporal variability, is a critical marker of sensory

processing related to the alignment of dynamic neural states (Afraimovich et al., 2004; Jones

et al., 2007; Rabinovich et al., 2008). The converging results of the present study indicate that

a disrupted temporal orchestration of cortical activity (Yusuf et al., 2022), especially in higher

sensory areas, and not a general attenuation of cortical oscillation may be the most profound

consequence of a period of congenital sensory deprivation. In this context we note that

feedforward and feedback connection profiles mature asynchronously, with feedforward

connectivity leading (Burkhalter et al., 1993). Moreover, there is evidence that the elaboration

of feedback connectivity more extensively depends on experience (Röder and Kekunnaya,

2021). Early sensory experience, thus, might be crucial for temporally aligned cortical feedback

signaling, and in turn for the shaping of feedforward activity. The observation that alpha-band

activity, beyond its purported role in the generation of P1, has been proposed to be a major

mediator of cortical feedback (Fries, 2015), corroborate this account.

In conclusion, we provide to the best of our knowledge the first report of an impaired visual

cortical phase coherence in humans after a period of transient congenital (but not

developmental) pattern visual deprivation, indicating the presence of a sensitive period for the

development of temporally coherent visual cortical processing in humans. Our results suggest

that aberrant temporal tuning of extrastriate cortical activity might underlie many of the

persistent visual deficits observed after recovering from congenital blindness.

Acknowledgements

We thank D. Balasubramanian (L V Prasad Eye Institute, Hyderabad, India) for supporting this study. Seema Banerjee, Larissa Brockmann, Maria Guerreiro, Marlene Hense, Giulia Dormal, Siddhart Srivatsav Rajendran, Lisa Stockleben and Florian Süßer helped with data acquisition in India and/or Germany. The study was funded the German Research Foundation (DFG, Ro 2625/10-1) and by the European Research Council grant ERC-2009-AdG 249425-*CriticalBrainChanges* to B.R.

References

Afraimovich VS, Zhigulin VP, Rabinovich MI (2004) On the origin of reproducible sequential activity in neural circuits. Chaos: An Interdisciplinary Journal of Nonlinear Science 14:1123–1129.

Aydore S, Pantazis D, Leahy RM (2013) A Note on the Phase Locking Value and its Properties. Neuroimage 74:231–244.

Bankó ÉM, Körtvélyes J, Németh J, Weiss B, Vidnyánszky Z (2013) Amblyopic deficits in the timing and strength of visual cortical responses to faces. Cortex 49:1013–1024.

Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B (Methodological) 57:289–300.

Birch EE, Cheng C, Stager DR, Weakley DR, Stager DR (2009) The critical period for surgical treatment of dense congenital bilateral cataracts. Journal of American Association for Pediatric Ophthalmology and Strabismus 13:67–71.

Bottari D, Kekunnaya R, Hense M, Troje NF, Sourav S, Röder B (2018) Motion processing after sight restoration: No competition between visual recovery and auditory compensation. NeuroImage 167:284–296.

Bottari D, Troje NF, Ley P, Hense M, Kekunnaya R, Röder B (2016) Sight restoration after congenital blindness does not reinstate alpha oscillatory activity in humans. Scientific Reports 6:24683.

Burkhalter A, Bernardo KL, Charles V (1993) Development of local circuits in human visual cortex. J Neurosci 13:1916–1931.

Buzzell GA, Niu Y, Aviyente S, Bernat E (2022) A practical introduction to EEG Time-Frequency Principal Components Analysis (TF-PCA). Developmental Cognitive Neuroscience 55:101114.

De Heering A, Maurer D (2014) Face memory deficits in patients deprived of early visual input by bilateral congenital cataracts. Developmental Psychobiology 56:96–108.

Delorme A, Makeig S (2004) EEGLAB: An open source toolbox for analysis of single-trial
EEG dynamics including independent component analysis. Journal of Neuroscience
Methods 134:9–21.

Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale
AM, Maguire RP, Hyman BT, Albert MS, Killiany RJ (2006) An automated labeling
system for subdividing the human cerebral cortex on MRI scans into gyral based
regions of interest. NeuroImage 31:968–980.

Di Russo F, Martínez A, Sereno MI, Pitzalis S, Hillyard SA (2002) Cortical sources of the
early components of the visual evoked potential. Human Brain Mapping 15:95–111.

Freunberger R, Höller Y, Griesmayr B, Gruber W, Sauseng P, Klimesch W (2008) Functional
similarities between the P1 component and alpha oscillations. European Journal of
Neuroscience 27:2330–2340.

Fries P (2015) Rhythms for Cognition: Communication through Coherence. Neuron 88:220–
235.

Grami A (2016) Chapter 3 - Signals, Systems, and Spectral Analysis. In: Introduction to
Digital Communications (Grami A, ed), pp 41–150. Boston: Academic Press. Available
at: https://www.sciencedirect.com/science/article/pii/B978012407682200003X
[Accessed October 1, 2024].

Gruber WR, Klimesch W, Sauseng P, Doppelmayr M (2005) Alpha Phase Synchronization
Predicts P1 and N1 Latency and Amplitude Size. Cerebral Cortex 15:371–377.

Hölig C, Guerreiro MJS, Lingareddy S, Kekunnaya R, Röder B (2023) Sight restoration in
congenitally blind humans does not restore visual brain structure. Cerebral Cortex
33:2152–2161.

Huang Y, Parra LC, Haufe S (2016) The New York Head-A precise standardized volume
conductor model for EEG source localization and tES targeting. Neuroimage
140:150–162.

Hyvärinen J, Carlson S, Hyvärinen L (1981) Early visual deprivation alters modality of
neuronal responses in area 19 of monkey cortex. Neuroscience Letters 26:239–243.

Jones LM, Fontanini A, Sadacca BF, Miller P, Katz DB (2007) Natural stimuli evoke dynamic
sequences of states in sensory cortical ensembles. Proceedings of the National
Academy of Sciences 104:18772–18777.

Klimesch W (2012) Alpha-band oscillations, attention, and controlled access to stored
information. Trends in Cognitive Sciences 16:606–617.

Klimesch W, Fellinger R, Freunberger R (2011) Alpha Oscillations and Early Stages of Visual
Encoding. Front Psychol 2 Available at:
https://www.frontiersin.org/journals/psychology/articles/10.3389/fpsyg.2011.00118/full
[Accessed September 25, 2024].

Klimesch W, Hanslmayr S, Sauseng P, Gruber WR, Doppelmayr M (2007) P1 and traveling
alpha waves: evidence for evoked oscillations. J Neurophysiol 97:1311–1318.

Klimesch W, Schack B, Schabus M, Doppelmayr M, Gruber W, Sauseng P (2004) Phase-
locked alpha and theta oscillations generate the P1-N1 complex and are related to
memory performance. Cognitive Brain Research 19:302–316.

Lüdecke D (2018) Sjstats: statistical functions for regression models. Available at:
894   https://zenodo.org/record/1489175 [Accessed October 17, 2021].

895   Makowski D, Ben-Shachar MS, Chen SHA, Lüdecke D (2019) Indices of effect existence and
896   significance in the Bayesian framework. Frontiers in Psychology 10:2767.

897   Mason SJ, Graham NE (2002) Areas beneath the relative operating characteristics (ROC)
898   and relative operating levels (ROL) curves: Statistical significance and interpretation.
899   Quarterly Journal of the Royal Meteorological Society 128:2145–2166.

900   Maurer D, Lewis TL (2013) Sensitive Periods in Visual Development. In: The Oxford
901   Handbook of Developmental Psychology, Vol. 1: Body and Mind (Zelazo PD, ed), pp
902   0. Oxford University Press. Available at:
903   https://doi.org/10.1093/oxfordhb/9780199958450.013.0008 [Accessed August 28,
904   2023].

905   McKyton A, Ben-Zion I, Doron R, Zohary E (2015) The limits of shape recognition following
906   late emergence from blindness. Current Biology 25:2373–2378.

907   Miller CE, Shapiro KL, Luck SJ (2015) Electrophysiological measurement of the effect of
908   inter-stimulus competition on early cortical stages of human vision. NeuroImage
909   105:229–237.

910   Mitchell DE, Maurer D (2022) Critical Periods in Vision Revisited. Annual Review of Vision
911   Science 8:291–321.

912   Oostenveld R, Fries P, Maris E, Schoffelen J-M (2011) FieldTrip: Open source software for
913   advanced analysis of MEG, EEG, and invasive electrophysiological data.
914   Computational intelligence and neuroscience 2011:156869.

915   Ossandón JP, Stange L, Gudi-Mindermann H, Rimmele JM, Sourav S, Bottari D, Kekunnaya
916   R, Röder B (2023) The development of oscillatory and aperiodic resting state activity
917   is linked to a sensitive period in humans. Neuroimage 275:120171.

918   Pascual-Marqui RD (2002) Standardized low-resolution brain electromagnetic tomography
919   (sLORETA): technical details. Methods Find Exp Clin Pharmacol 24 Suppl D:5–12.

920   Peirce JW (2009) Generating stimuli for neuroscience using PsychoPy. Frontiers in
921   Neuroinformatics 2:1–8.

922   Pfurtscheller G, Neuper C, Mohl W (1994) Event-related desynchronization (ERD) during
923   visual processing. International Journal of Psychophysiology 16:147–153.

924   Piastra MC, Cruijsen J van der, Piai V, Jeukens FEM, Manoochehri M, Schouten AC, Selles
925   RW, Oostendorp T (2021) ASH: an Automatic pipeline to generate realistic and
926   individualized chronic Stroke volume conduction Head models. J Neural Eng
927   18:044001.

928   Pitchaimuthu K, Dormal G, Sourav S, Shareef I, Rajendran SS, Ossandón JP, Kekunnaya R,
929   Röder B (2021) Steady state evoked potentials indicate changes in nonlinear neural
930   mechanisms of vision in sight recovery individuals. Cortex 144:15–28.

931   Power M, Fell G, Wright M (2013) Principles for high-quality, high-value testing. BMJ
932   Evidence-Based Medicine 18:5–10.

933   Putzar L, Hötting K, Rösler F, Röder B (2007) The development of visual feature binding
934   processes after visual deprivation in early infancy. Vision Research.

R Core Team (2016) R: A language and environment for statistical computing. Available at:
https://www.r-project.org/.

Rabinovich M, Huerta R, Laurent G (2008) Transient Dynamics for Neural Processing.
Science 321:48–50.

Röder B, Kekunnaya R (2021) Visual experience dependent plasticity in humans. Current
Opinion in Neurobiology 67:155–162.

Röder B, Kekunnaya R, Guerreiro MJS (2021) Neural mechanisms of visual sensitive
periods in humans. Neuroscience and biobehavioral reviews 120:86–99.

Röder B, Ley P, Shenoy BH, Kekunnaya R, Bottari D (2013) Sensitive periods for the
functional specialization of the neural system for human face processing.
Proceedings of the National Academy of Sciences 110:16760–16765.

Roy S (1946) খিচুড়ি (Hodgepodge). In: আবোল তাবোল (Balderdash), 2nd ed. Kolkata: Signet Press.
Available at: https://archive.org/details/dli.scoerat.2010aboltabol/page/n7 [Accessed
January 8, 2025].

Schad DJ, Vasishth S, Hohenstein S, Kliegl R (2020) How to capitalize on a priori contrasts
in linear (mixed) models: A tutorial. Journal of Memory and Language 110:104038.

Schrader S, Westhoff A, Piastra MC, Miinalainen T, Pursiainen S, Vorwerk J, Brinck H,
Wolters CH, Engwer C (2021) DUNEuro—A software toolbox for forward modeling in
bioelectromagnetism. PLOS ONE 16:e0252431.

Sourav S, Bottari D, Kekunnaya R, Röder B (2018) Evidence of a retinotopic organization of
early visual cortex but impaired extrastriate processing in sight recovery individuals.
Journal of Vision 18:22.

Sourav S, Bottari D, Shareef I, Kekunnaya R, Röder B (2020) An electrophysiological
biomarker for the classification of cataract-reversal patients: A case-control study.
EClinicalMedicine 27:100559.

Sourav S, Kekunnaya R, Bottari D, Shareef I, Pitchaimuthu K, Röder B (2024) Sound
suppresses earliest visual cortical processing after sight recovery in congenitally blind
humans. Commun Biol 7:1–14.

Tadel F, Baillet S, Mosher JC, Pantazis D, Leahy RM (2011) Brainstorm: a user-friendly
application for MEG/EEG analysis. Comput Intell Neurosci 2011:879716.

Tallon-Baudry C, Bertrand O, Tallon-Baudry C, Bertrand O, Tallon-Baudry C, Bertrand O
(1999) Oscillatory gamma activity in humans and its role in object representation.
Trends in Cognitive Sciences 3:151–162.

Tytla ME, Lewis TL, Maurer D, Brent HP (1993) Stereopsis after congenital cataract.
Investigative Ophthalmology and Visual Science 34:1767–1773.

Wanet-Defalque M-C, Veraart C, De Volder A, Metz R, Michel C, Dooms G, Goffinet A (1988)
High metabolic activity in the visual cortex of early blind human subjects. Brain
Research 446:369–373.

Weber A (2018) The USC-SIPI image database: Version 5. Available at: http://sipi. usc.
edu/database/.

Weinhaus F (2011) Fourier Transforms -- IM v6 Examples. ImageMagick v6 Examples --
976   Fourier Transforms Available at: https://legacy.imagemagick.org/Usage/fourier/
977   [Accessed October 1, 2024].

978   Wiesel TN, Hubel DH (1965a) Extent of recovery from the effects of visual deprivation in
979   kittens. Journal of neurophysiology 28:1060–1072.

980   Wiesel TN, Hubel DH (1965b) Comparison of the effects of unilateral and bilateral eye
981   closure on cortical unit responses in kittens. Journal of neurophysiology.

982   World Medical Association (2013) World Medical Association Declaration of Helsinki: Ethical
983   Principles for Medical Research Involving Human Subjects. JAMA 310:2191–2194.

984   Yusuf PA, Lamuri A, Hubka P, Tillein J, Vinck M, Kral A (2022) Deficient Recurrent Cortical
985   Processing in Congenital Deafness. Front Syst Neurosci 16:806142.