# The Temporal Stability of Visual Cortical Processing in Humans Depends on Early Experience

Suddha Sourav [a, *], Max Emanuel Feucht [a], Ramesh Kekunnaya [b], and Brigitte Röder [a, b]

[a] Biological Psychology and Neuropsychology, University of Hamburg, 20146 Hamburg, Germany

[b] Jasti V Ramanamma Children's Eye Care Center, Child Sight Institute, L V Prasad Eye Institute, 500034 Hyderabad, India

**Author Information**

**Suddha Sourav**: https://orcid.org/0000-0002-4172-7711

Contributions (CRediT taxonomy): Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Visualization, Writing – original draft, Writing – review & editing

* Corresponding author. E-Mail: suddha.sourav@uni-hamburg.de

**Max Emanuel Feucht**: https://orcid.org/0009-0009-5929-9258

Contributions (CRediT taxonomy): Formal analysis, Methodology, Software, Validation, Visualization, Writing – original draft, Writing – review & editing

**Ramesh Kekunnaya:** https://orcid.org/0000-0001-5789-2300

Contributions (CRediT taxonomy): Investigation, Funding acquisition, Methodology, Project administration, Resources, Supervision, Writing – review & editing

**Brigitte Röder:** https://orcid.org/0000-0003-3088-8023

Contributions (CRediT taxonomy): Conceptualization, Investigation, Funding acquisition, Methodology, Project administration, Resources, Supervision, Writing – review & editing

The authors have no conflicts of interest to declare.

No generative artificial intelligence models were used to write or rephrase any part of this manuscript.

## 25 Abstract

Proper timing is essential for effective neural processing. Yet, how early postnatal vision

shapes the temporal stability of human visual cortical processing remains unknown. Here,

using electroencephalography, we examined cortical timing properties in individuals who were

born pattern vision blind due to congenital cataracts, but surgically recovered sight. While sight-

recovery individuals exhibited an attenuated cortical oscillatory phase coherence (i.e., higher

temporal variability) during visual processing, their oscillatory strength was unimpaired.

Moreover, phase coherence information, but not activation strength, allowed the classification

of sight-recovery from control individuals. Finally, exchanging phase information between

sight-recovery and control individuals indicated oscillatory timing impairments as the source of

group differences in higher-order visual cortical processing. Neural timing impairments were

specific for reversed congenital blindness, that is, were not observed in individuals with

reversed developmental (late-onset) cataracts. These results suggest that the development of

intricately temporally orchestrated visual cortical processing in humans requires early visual

experience.

## Significance Statement

Neural circuit functioning requires an efficient temporal orchestration. The present work investigated how the temporal stability of visual cortical dynamics depends on adequate experience. In rare individuals who were born with pattern vision blindness but later recovered sight, a marked increase in the temporal variability of oscillatory brain activity during visual processing was observed while oscillatory strength was surprisingly unimpaired. Therefore, the emergence of precisely timed visual processing in the human brain seems to crucially depend on early visual experience. We speculate that impaired timing of neural processing, cascading throughout the visual cortical hierarchy, might be a major source of the incomplete visual recovery in individuals with treated congenital blindness.

## Introduction

The typically developed brain exhibits a remarkable stability of sensory processing on a trial-

by-trial basis, which may appear surprising in light of considerable spontaneous activity and

diverse sources of biological noise in the brain (Rabinovich et al., 2008; Oby et al., 2025;

Trägenap et al., 2025). Although numerous studies have investigated changes to visual cortical

activation strength after sight loss (Wanet-Defalque et al., 1988; Rösler et al., 1993; Amedi et

al., 2003), and subsequently, after sight recovery (Mitchell and Maurer, 2022; Röder and

Kekunnaya, 2022), the neural mechanisms preventing complete behavioral recovery are yet

unknown. In fact, humans who had suffered from a period of congenital pattern visual blindness

due to the presence of total, bilateral congenital cataracts (CC), suffer from persistent visual

impairments following sight restoration, even after extended recovery periods. These deficits

include a permanently reduced visual acuity and stereoscopic ability (Tytla et al., 1993; Birch

et al., 2009), atypical mid- to high-level visual feature processing (Putzar et al., 2007; McKyton

et al., 2015), and altered face processing (Röder et al., 2013; De Heering and Maurer, 2014).

In contrast, a substantially higher recovery of visual functions has been observed if the onset

of visual loss was later (instead of congenital), e.g., through developmental cataracts (Röder

et al., 2013; Bottari et al., 2018; Sourav et al., 2020). These results indicate the presence of

*sensitive periods*, during which the effect of experience on the development of visual cortical

circuits and the associated functions is especially strong (Knudsen, 2004; Maurer and Lewis,

2013; Röder et al., 2021). However, to date, the temporal stability of visual cortical processing

as a sensitive period mechanism has not been studied in humans.

On the one hand, functional magnetic resonance imaging (fMRI) methods, indispensable for

deriving spatially well-resolved activation maps in sight-recovery individuals, cannot resolve

the fast, millisecond-level cortical temporal dynamics. On the other hand,

electro/magnetoencephalographic (EEG/MEG) research methods have emphasized event-

related (i.e., trial-averaged) indices of cortical activation (Makeig et al., 2004), discarding the

trial-by-trial temporal variation information necessary to investigate the stability of visual

cortical processing (Bottari et al., 2016; Ossandón et al., 2023; Sourav et al., 2024). For

example, an attenuated event-related potential (ERP) in CC individuals, that is, a scalp-

recorded marker of the summed electric field potentials in the brain, might result from a

generally reduced cortical activation across trials, or equivalently, from a lack of temporal

coherence across individual trials with unimpaired overall activation strength. At the level of

the neural generators, these two hypothetical extremes represent fundamentally different (and

to date not disentangled) effects of a period of transient visual deprivation, i.e., a generally

attenuated cortical activity vs. compromised temporal stability of visual cortical processing.

In two experiments, we here investigated the temporal stability of visual cortical processing

indexed by oscillatory phase coherence in sight recovery individuals with a history of transient

congenital blindness. In addition, we compared cortical activation strength in CC individuals to

normally sighted control individuals. First, time-frequency decomposition using wavelet

transforms was employed to calculate the activation magnitude, operationalized by the post-

stimulus power change of cortical oscillations compared to a pre-stimulus baseline, as well as

the inter-trial oscillatory phase coherence to assess the temporal stability of cortical visual

processing. These two markers were subsequently compared between individuals with

reversed congenital cataracts and normally sighted controls, and to ascertain the presence of

an early sensitive period, additionally between individuals with reversed developmental

(childhood) cataracts and normally sighted controls. Next, we used a classification approach

that attempted to identify CC individuals based on power changes relative to pre-stimulus

baseline, as well as based on inter-trial phase coherence information in the time-frequency

representations. Further, we derived what we term *"mutual magnitude-phase transplants",* a

novel method that we here introduce, which allowed us to exchange the per-trial activation

strength and timing information between a cataract reversal individual and their matched

normally sighted control participant. Previously, we have reported a robust and replicable

attenuation of the earliest electrophysiological marker of higher-order (extrastriate) visual

cortical processing, the P1 wave (~ 150 ms; Di Russo et al., 2002; Miller et al., 2015), in CC

individuals (Sourav et al., 2020). The method of mutual magnitude-phase transplants allowed

us to investigate the relative contribution of activation strength vs. timing information to the

extrastriate cortical changes observed after a period of transient congenital visual deprivation.

Finally, minimum-norm source estimation was performed to verify the early extrastriate cortical

sources of the P1 wave.

# Materials and Methods

## Participants

Datasets originating from two different experiments, termed experiment 1 and experiment 2,

were utilized. Both experiments included a group of congenital cataract reversal individuals

(CC) as well as a group of developmental cataract reversal individuals (DC), who had

undergone the same surgical procedures. In addition, normally sighted control participants with

a typical developmental history were recruited, and each normally sighted control participant

was matched to a cataract reversal individual for age, sex, and handedness.

Thirteen CC individuals took part in experiment 1, of whom the data of 1 participant were not

analyzed here due to a very short visual deprivation period (1 month). The remaining CC

participants, whose data entered further analysis ($n$ = 12, see Table 1 for participant details),

had a mean age of 18.67 years (range = 10 – 37 years), 2 were female (and 10 were male),

and 1 was left-handed.  In addition, 16 DC individuals took part in experiment 1, of whom the

data of 3 individuals were not analyzed respectively due to a diagnosis of seizure and

developmental delay, the participant's unwillingness to continue the experiment shortly after

the start and the inability of the participant to look straight ahead. The data of 13 DC individuals

(Table 2) entered further analysis (mean age = 16 years, range = 11 – 24 years, 5 females and

8 males, all right-handed)**.** The CC individuals had received surgical treatment after a mean of

41.33 months of blindness (range 4 – 213 months). The average age at treatment for DC

individuals was 117.38 months (range 24 – 208 months). The visual acuity (VA) of the CC

individuals (geometric mean, decimal = 0.23, range = 0.05 – 0.5) was significantly lower

compared to that of the DC individuals (geometric mean, decimal = 0.74, range = 0.44 – 1.00),

as revealed by a one-tailed $t$-test after converting the VAs to LogMAR units**,** $t(14.531)$ = 5.642,

*p* < .001. The mean age of the typically sighted control individuals matched to the CC

individuals (MCC group) was 18.08 years (range = 10 – 34 years, 2 female and 10 male, 1 left-

handed), and 16.85 years for the typically sighted control individuals matched to the DC

individuals (MDC group, range = 10 – 25 years, 5 female and 8 male, all right-handed).

In experiment 2, 15 participants with a history of bilateral, dense congenital cataracts took part.

The data of 2 of these participants were rejected because in one case, the participant was born

with incomplete cataract and thus did not suffer from complete pattern visual blindness since

birth, and in another case, because of a very short visual deprivation period (1 month). Thus,

the data of 13 CC individuals entered further analysis (Table 3, mean age = 17.69 years, range

6 – 39 years, 3 female and 10 male, 1 left-handed)**.** In addition, 16 DC individuals took part in

experiment 2. We did not analyze the data of 1 participant due to a history of seizures and

developmental delay. The data of 15 DC individuals entered further analysis (Table 4, mean

age = 14.47 years, range 9 – 24 years, 4 female and 11 male, 1 left-handed)**.**

The CC individuals in experiment 2 whose data were analyzed (*n* = 13) had been treated for

cataract after a mean of 45.31 months of blindness (range 4 – 213 mo.), and the DC individuals

whose data were analyzed had been treated for cataract with a mean age of 88.33 months

(range 23 – 173 mo.). Once again, the CC individuals exhibited a generally lower mean

geometric visual acuity (0.21, range = 0.051 – 0.5) than the DC individuals (0.618, range =

0.23 – 1.00; one-tailed t-test, *t*(21.532) = 5.297, p < .001). The MCC group in experiment 2 had

a mean age of 17 years (range = 7 – 38 years, 3 female and 10 male, 1 lefthanded), and the

MDC group had a mean age of 14.93 years (range = 11 – 26 years, 4 female and 11 male, 1

lefthanded).

Ten CC individuals, 8 DC individuals and 5 normally sighted control individuals took part in

both experiment 1 and experiment 2; 35.7% of cataract reversal individuals were therefore

unique to experiment 2 (i.e., 3 CC and 7 DC individuals). All cataract reversal individuals from

both experiments were recruited and tested at the L V Prasad Eye Institute, Hyderabad, India.

Control individuals were tested at the University of Hamburg, Hamburg, Germany. Adult

participants received a small monetary compensation for taking part in the study, whereas

minors received a small gift. Costs for travel and accommodation were reimbursed. The study

obtained approval from the local ethics commission of the faculty of Psychology and Human

Movement at the University of Hamburg and the institutional ethics review board of LVPEI, and

adhered to the ethical principles outlined in the Declaration of Helsinki (World Medical

Association, 2013) .

## Experimental Design

In both experiments, participants were presented monochromatic square wave grating stimuli

in front of a grey background, one at a time for 150 ms in one of the four visual field quadrants

with a mean interstimulus interval of 1.85 s (1.5 s – 2.2 s). The stimuli were presented with a

fixed eccentricity of 4 degrees from the center and subtended a visual angle of 2.5° as depicted

in Figure 2 in Sourav et al. (2020). The stimuli were presented at a polar angle of 25° in the

upper visual field and at a polar angle of −45° in the lower visual field. The gratings were either

horizontally (standard stimuli, $P$ = .80) or vertically oriented (target stimuli, $P$ = .20), with a

spatial frequency of 2 cycles/degree. Participants were instructed to attend to a central fixation

cross and to indicate when a rare target stimulus appeared in any of the locations by pressing

a foot pedal in experiment 1 and by clicking a computer mouse in experiment 2. Sight restored

cataract individuals were tested using a Dell IN2030 monitor at LVPEI, Hyderabad, and

typically sighted control individuals were tested on a Samsung P2370 monitor at the University

of Hamburg. Both monitors had a refresh rate of 60 Hz and a nominal luminance of 250 cdm⁻

$^2$. Participants sat at a distance of 45 cm to the monitor in experiment 1, and at a distance of

65 cm in experiment 2. The angular measurements described above were identical in both

experiments. Participants were presented additional stimuli in both experiments which are not

reported here.

For stimulus presentation we used the the PsychoPy framework (v1.83, Peirce, 2009).

**Table 1.** Participant characteristics for the sight recovery individuals with a history of dense bilateral congenital cataracts and subsequent surgery
(CC) in experiment 1. The table is based on Sourav et al. (2020).

| Participant ID | Group | Age (yrs.) | Blindness Duration (mo.) | Time Since Surgery (mo.) | Sex | Handedness | Visual Acuity (VA), decimal | VA, LogMAR | Pre-surgical VA (Better Eye) | Nystagmus | Strabismus | Family History |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| CC – 001 | CC | 26 | 5 | 308 | M | Right | 0.180 | 0.745 | NA | Yes | Esotropia | No |
| CC – 002 | CC | 37 | 24 | 428 | M | Right | 0.400 | 0.398 | NA | Yes | Esotropia | Yes |
| CC – 003 | CC | 11 | 7 | 124 | M | Right | 0.430 | 0.367 | Fixates and follows light | Yes | Exotropia | No |
| CC – 004 | CC | 13 | 4 | 158 | M | Right | 0.130 | 0.886 | Fixates and follows light | Yes | Exotropia | No |
| CC – 005 | CC | 33 | 72 | 330 | M | Right | 0.051 | 1.292 | NA | Yes | No | Yes |
| CC – 006 | CC | 23 | 4 | 279 | M | Left | 0.130 | 0.886 | NA | Yes | Esotropia | Yes |
| CC – 007 | CC | 13 | 15 | 150 | M | Right | 0.500 | 0.301 | Fixates and follows light | Yes | Exotropia | Yes |
| CC – 009 | CC | 10 | 11 | 116 | M | Right | 0.240 | 0.620 | Fixates and follows light | Yes | Esotropia | No |
| CC – 010 | CC | 18 | 25 | 198 | F | Right | 0.310 | 0.509 | Fixates and follows light | Yes | Esotropia | Yes |
| CC – 011 | CC | 19 | 213 | 25 | M | Right | 0.200 | 0.699 | Counting fingers at 0.5 m | Yes | Esotropia | Yes |
| CC – 012 | CC | 11 | 42 | 89 | F | Right | 0.450 | 0.347 | Fixates and follows light | Yes | Exotropia | Yes |
| CC – 013 | CC | 10 | 74 | 48 | M | Right | 0.250 | 0.602 | 0.017 | Yes | No | Yes |

187  **Table 2.** Participant characteristics for the sight recovery individuals with a history of bilateral developmental cataracts and subsequent surgery (DC)
188  in experiment 1. The table is based on Sourav et al. (2020).

| Participant ID | Group | Age (yrs.) | Age at Surgery (yrs.) | Time Since Surgery (mo.) | Sex | Handedness | Visual Acuity (VA), decimal | VA, LogMAR | Pre-surgical VA (Better Eye) | Nystagmus | Strabismus | Family History |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| DC – 001 | DC | 16 | 12 | 53 | M | Right | 1.000 | 0.000 | Counting fingers at 0.5 m | No | No | No |
| DC – 002 | DC | 19 | 14 | 55 | M | Right | 1.000 | 0.000 | 0.200 | No | No | No |
| DC – 003 | DC | 12 | 6 | 63 | F | Right | 1.000 | 0.000 | 0.500 | No | No | No |
| DC – 004 | DC | 11 | 8 | 39 | F | Right | 0.530 | 0.276 | 0.159 | No | No | No |
| DC – 005 | DC | 13 | 8 | 68 | M | Right | 0.710 | 0.149 | 0.100 | No | Exotropia | No |
| DC – 006 | DC | 12 | 8 | 51 | M | Right | 0.600 | 0.222 | 0.400 | No | No | No |
| DC – 007 | DC | 12 | 6 | 77 | M | Right | 1.000 | 0.000 | 0.250 | No | No | Yes |
| DC – 008 | DC | 18 | 12 | 74 | F | Right | 0.790 | 0.102 | Sees hand movement | No | No | No |
| DC – 009 | DC | 16 | 10 | 67 | M | Right | 0.740 | 0.131 | 0.400 | No | No | No |
| DC – 010 | DC | 24 | 2 | 264 | F | Right | 0.440 | 0.357 | Fixates and follows light | No | Exotropia | No |
| DC – 011 | DC | 16 | 7 | 110 | F | Right | 0.530 | 0.276 | Counting fingers @ 1 m | No | Exotropia | Yes |
| DC – 012 | DC | 20 | 12 | 97 | M | Right | 0.880 | 0.056 | 0.400 | No | No | No |
| DC – 013 | DC | 19 | 17 | 17 | M | Right | 0.730 | 0.137 | 0.200 | No | No | No |

190 **Table 3.** Participant characteristics for the sight recovery individuals with a history of dense bilateral congenital cataracts and subsequent surgery
191 (CC) in experiment 2 (Please note that age of the same participant might vary from Table 1 due to the experiments being run at different time points).
192 The table is based on Sourav et al. (2020).

| Participant ID | Group | Age (yrs.) | Blindness Duration (mo.) | Time Since Surgery (mo.) | Sex | Handedness | Visual Acuity (VA), decimal | VA, LogMAR | Pre-surgical VA (Better Eye) | Nystagmus | Strabismus | Family History |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| CC – 001 | CC | 26 | 5 | 308 | M | Right | 0.180 | 0.745 | NA | Yes | Esotropia | No |
| CC – 005 | CC | 33 | 72 | 330 | M | Right | 0.051 | 1.292 | NA | Yes | No | Yes |
| CC – 014 | CC | 6 | 5 | 70 | M | Right | 0.240 | 0.620 | Not fixating/following light | Yes | No | No |
| CC – 002 | CC | 39 | 24 | 444 | M | Right | 0.400 | 0.398 | NA | Yes | Esotropia | Yes |
| CC – 004 | CC | 14 | 4 | 175 | M | Right | 0.130 | 0.886 | Fixates and follows light | Yes | Exotropia | No |
| CC – 006 | CC | 24 | 4 | 294 | M | Left | 0.130 | 0.886 | NA | Yes | Esotropia | Yes |
| CC – 007 | CC | 15 | 15 | 166 | M | Right | 0.500 | 0.301 | Fixates and follows light | Yes | Exotropia | Yes |
| CC – 009 | CC | 11 | 11 | 129 | M | Right | 0.240 | 0.620 | Fixates and follows light | Yes | Esotropia | No |
| CC – 015 | CC | 8 | 48 | 48 | F | Right | 0.150 | 0.824 | Perception of light | Yes | No | Yes |
| CC – 013 | CC | 11 | 74 | 61 | M | Right | 0.250 | 0.602 | 0.017 | Yes | No | Yes |
| CC – 012 | CC | 11 | 42 | 90 | F | Right | 0.450 | 0.347 | Fixates and follows light | Yes | Exotropia | Yes |
| CC – 011 | CC | 21 | 213 | 40 | M | Right | 0.200 | 0.699 | Counting fingers at 0.5 m | Yes | Esotropia | Yes |
| CC – 016 | CC | 11 | 72 | 68 | F | Right | 0.260 | 0.585 | Fixates and follows light | Yes | No | No |

194 **Table 4.** Participant characteristics for the sight recovery individuals with a history of bilateral developmental cataracts and subsequent surgery (DC)
195 in experiment 2 (Please note that age of the same participant might vary from Table 2 due to the experiments being run at different time points). The
196 table is based on Sourav et al. (2020).

| Participant ID | Group | Age (yrs.) | Age at Surgery (yrs.) | Time Since Surgery (mo.) | Sex | Handedness | Visual Acuity (VA), decimal | VA, LogMAR | Pre-surgical VA (Better Eye) | Nystagmus | Strabismus | Family History |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| DC – 002 | DC | 19 | 14 | 55 | M | Right | 1.000 | 0.000 | 0.200 | No | No | No |
| DC – 011 | DC | 14 | 7 | 80 | F | Right | 0.530 | 0.276 | Finger counting at 1 m | No | Exotropia | Yes |
| DC – 003 | DC | 13 | 6 | 74 | F | Right | 1.000 | 0.000 | 0.500 | No | No | No |
| DC – 005 | DC | 14 | 8 | 80 | M | Right | 0.710 | 0.149 | 0.100 | No | Exotropia | No |
| DC – 007 | DC | 12 | 6 | 77 | M | Right | 1.000 | 0.000 | 0.250 | No | No | Yes |
| DC – 014 | DC | 12 | 7 | 66 | M | Right | 0.230 | 0.638 | 0.125 | No | No | No |
| DC – 015 | DC | 12 | 7 | 67 | M | Right | 0.460 | 0.337 | 0.100 | No | No | No |
| DC – 009 | DC | 16 | 10 | 67 | M | Right | 0.740 | 0.131 | 0.400 | No | No | No |
| DC – 010 | DC | 24 | 2 | 264 | F | Right | 0.440 | 0.357 | Fixates and follows light | No | Exotropia | No |
| DC – 008 | DC | 18 | 12 | 74 | F | Right | 0.790 | 0.102 | Perceives hand movement | No | No | No |
| DC – 016 | DC | 17 | 12 | 60 | M | Right | 0.310 | 0.509 | 0.160 | No | No | Yes |
| DC – 017 | DC | 9 | 2 | 86 | M | Right | 0.930 | 0.032 | 0.154 | No | No | No |
| DC – 018 | DC | 16 | 6 | 113 | M | Right | 0.520 | 0.284 | Finger counting at 1 m | No | No | No |
| DC – 019 | DC | 10 | 3 | 76 | M | Left | 0.730 | 0.137 | 0.065 | No | No | Yes |
| DC – 020 | DC | 11 | 1 | 115 | M | Right | 0.667 | 0.176 | 0.154 | No | No | No |

## EEG: Data Acquisition and Preprocessing

EEG data were acquired continuously during the whole experiment using 32 Ag/AgCl electrodes with a left earlobe online reference. Brain activity was recorded at the 10-10 locations FP1, FP2, F7, F3, Fz, F4, F8, FC5, FC1, FCz, FC2, FC6, T7, C3, Cz, C4, T8, TP9, CP5, CP1, CP2, CP6, TP10, P7, P3, Pz, P4, P8, O1, O2, F9, and F10, with an online high-pass filter with a cutoff frequency of 0.016 Hz, a low-pass filter with a cutoff frequency of 250 Hz, and a sampling rate of 1 kHz. To eliminate electrical line noise, the data were notch-filtered offline at 50 Hz and, if apparent in the frequency spectrum, at the corresponding harmonics. Subsequently, the voltages were average referenced and biological artifacts such as blinks, eye movements, muscle activity or heartbeats were removed from the signal with an independent component analysis (ICA), using EEGLAB for MATLAB (EEGLAB, version 11.5.4b, Delorme and Makeig, 2004; MATLAB version 2012b; MathWorks, Natick, MA). To prevent confounds due to eye movements or blinks during stimulus presentation, we excluded stimulus epochs that contained blinks or eye movements within the time window of −25 ms to 175 ms relative to the stimulus, as previously reported (Sourav et al., 2020, 2024). The data were then low pass filtered at 40 Hz and EEG epochs were generated from -1.7s before to 1.7 s after the stimulus onset. For increasing the statistical power, data from trials with a left hemifield stimulation were remapped; at each electrode, the data were swapped with the data of the electrode in its mirrored location with respect to the nasion-inion axis, ignoring the midline electrodes (see Sourav et al., 2020). Additionally, to prevent artifacts related to motor activation, epochs containing a motor response within ±500 ms of the stimulus onset were rejected.

Only EEG trials/ERPs generated by the frequent, standard stimuli were used for further analysis. Moreover, as extrastriate-generated P1 wave amplitudes elicited by upper visual field stimulation are larger than those elicited by lower visual field stimulations in the present paradigm, we only considered visual ERPs for upper visual field stimulation for the subsequent analyses, as in Sourav et al. (2020).

## Data Reuse

The dataset in the current study has been derived from Sourav et al. (2020). The only difference is the exclusion of one participant with a very short duration of visual deprivation prior to the commencement of data analysis (1 month in participant CC – 008; see supplemental materials for Sourav et al., 2020). Parts of the dataset have been included in two other publications (Sourav et al., 2018, 2024).

## Time-Frequency Analysis

Time-frequency representations of the EEG data were derived by wavelet transformation of individual trials at each electrode using the *FieldTrip* software in MATLAB (Oostenveld et al., 2011). Signals at each electrode were convolved with complex Morlet wavelets which have a Gaussian shape over time (with standard deviation $\sigma_t$) as well as over frequency (with standard deviation $\sigma_f$), centered around a time $t_0$ and a frequency $f_0$ (Tallon-Baudry et al., 1999). Non-constant wavelets were applied with a ratio $f_0/\sigma_f = 4$ for the lowest frequency and increasing linearly up to 8.5 for the highest, identical to Bottari et al. (2016, 2018). Thereafter, the relative power change compared to baseline ($\Delta P_{rel}$), and the inter-trial phase coherence (ITPC) across trials were computed at each frequency from 2 to 40 Hz with a step size of 1 Hz, and from 700 ms before to 700 ms after stimulus onset in steps of 20 ms. $\Delta P_{rel}$ was derived by normalizing the wavelet power with respect to a baseline interval from -700 ms to -300 ms, i.e., $\Delta P(t,f)_{rel} = [P(t,f) - P_{baseline}(f)]/P_{baseline}(f)$, see Bottari et al. (2016, 2018). The ITPC is a measure of oscillatory phase synchronization across trials at a given timepoint and frequency (Delorme & Makeig, 2004) and was derived by calculating the magnitude of the complex average of the normalized phase directions. The ITPC can take on values in the interval [0, 1], where ITPC = 1 indicates a perfect synchronization across trials and ITPC = 0 indicates complete desynchronization. Importantly, the ITPC is a biased but consistent statistical measure, e.g., for uniformly random, desynchronized phase values with a finite number of trials, the ITPC is expected to be positive (i.e., is overestimated), but approaches 0 as the number of trials is increased (Aydore et al., 2013). To remove this bias of the ITPC estimates for group comparisons, and to derive mutual magnitude-phase transplants as described in a following

253    section, we used an equal number of EEG epochs for each cataract reversal participant and

254    their control participant.

## Cluster Based Permutation Tests

The $\Delta P_{rel}$ and ITPC values were compared between each cataract reversal group and their

sighted controls (i.e., CC vs. MCC; DC vs. MDC) using two-tailed cluster-based permutation

tests. The tests were performed with the *FieldTrip* framework (Oostenveld et al., 2011) running

on MATLAB (R2022a, Natick, MA), over the post-stimulus time-frequency plane ($t$: 0 – 700 ms;

$f$: 2- 40 Hz) and over the 13 electrodes posterior to the midline, where the P1 wave is most

prominent. The cluster-based permutation tests were based on 10,000 randomizations and

used weighted cluster mass as the test metric.

After visual stimulus presentation, a task-related decrease of $\Delta P_{rel}$ is typically observed in the

alpha (8 – 12 Hz) frequency range ("alpha blocking", Klimesch, 2012). Based on a previous

report by Bottari et al. (2016) that task-related post-stimulus alpha blocking might differ in the

CC group, we additionally compared $\Delta P_{rel}$ in each sight recovery group to their control group

(i.e., CC vs. MCC; DC vs. MDC) with cluster-based permutation tests, with the same

parameters as above, in the "alpha blocking" time-frequency window (time: 0.3 – 0.7 s,

frequency: 8 – 12 Hz) .

## Classification Based on Time-Frequency Features

In addition to statistical tests for group differences in time-frequency representations, we tested

group discriminability of CC vs. Non-CC (i.e., DC and MCC) and CC vs. DC individuals in a

classification paradigm based on the time-frequency features, separately for $\Delta P_{rel}$ and ITPC.

Classification was performed based on oscillatory activity from 2 to 40 Hz in steps of 1 Hz (i.e.,

in 39 frequency bins) and from 100 ms before stimulus onset to 500 ms after stimulus onset,

with 20 ms step size (i.e., 31 temporal bins) at all 13 posterior electrodes, yielding 15717

classification features for both $\Delta P_{rel}$ and ITPC. To deal with the high dimensionality of the data,

we employed principal component analysis (PCA) prior to classification (Buzzell et al., 2022).

Using the principal component loadings, participants were then classified using linear support

vector machines (SVM). Data from experiment 1 were used for deriving the principal

components, for tuning the SVM hyperparameters, and for subsequent SVM training. In

contrast, data from experiment 2 were utilized to validate the previously trained SVMs. The

analysis was performed in R, version 4.0.3 (R Core Team, 2016).

Dimension Reduction of the Time-Frequency Features

The $\Delta P_{rel}$ and ITPC values at each frequency and timepoint at the 13 posterior electrodes were

used to derive principal components for $\Delta P_{rel}$ and ITPC separately. PCA was performed across

all participants of experiment 1 (including the normally sighted control individuals), without

centering or scaling the $\Delta P_{rel}$ and ITPC features, respectively. We chose the first three principal

components for classification which explained 84.84% of the variance of ITPC and 49.21% of

the variance of $\Delta P_{rel}$. Importantly, the PCA components do not necessarily reflect components

with the highest *intergroup* variance, rather they reflect the highest *interindividual* variance

regardless of the group membership. According to the suggestion by Buzzell et al. (2022),

varimax rotation was first applied to the selected components before computing the single

participants' loadings at each electrode, yielding 39 $\Delta P_{rel}$ and 39 ITPC features per participant

(13 electrodes × 3 principal component loadings). The varimax-rotated components are

displayed in Fig. 1.

To assess the robustness of potential group differences, classification was additionally

performed based on loadings of each of the first three components averaged over all posterior

electrodes, yielding 3 classification features for both $\Delta P_{rel}$ and ITPC. The principal components

were solely derived from data of experiment 1 and the same components were used for

classification of the data from experiment 2.

**Figure 1. Principal component loadings of relative power change and of ITPC.**

**a – c**. The first three principal components of the relative power change compared to baseline ($\Delta P_{rel}$), summed across all posterior electrodes for each time-frequency feature (unitless). **d – e**. The first three principal components of inter-trial phase coherence (ITPC), summed across all posterior electrodes for each time-frequency feature (unitless). Each pixel in each plot represents one time-frequency feature. Summation over all electrodes is performed to highlight overall positive and negative loading clusters for each component.

## Support Vector Machine Configuration

Using the features derived from the PCA-based dimensionality reduction, the regularizing hyperparameter $c$ of the SVM was tuned in a repeated cross-validation (CV) procedure, with a split of $k = 5$ (CC vs. non-CC classification) and $k = 3$ (CC vs. DC classification). The CV procedure was repeated 500 times, and the optimal $c$ that was chosen most often of all repetitions was used to fit a final SVM to all data from experiment 1. For each feature class (i.e., $\Delta P_{rel}$ and ITPC), each classification type (CC vs. non-CC, and CC vs. DC), an SVM was trained once using all 39 component loadings from experiment 1 and the most often chosen optimal value of $c$, summing to 4 training procedures (2 feature classes, × 2 classification types). This procedure was repeated for the component loadings averaged over the posterior electrodes, yielding 4 additional training procedures each only using 3 classification features. For each SVM, a receiver operating characteristic (ROC) analysis was conducted to extract the optimal classification threshold by maximizing the *Youden's J* parameter. To validate each trained classifier, the decision boundary derived from experiment 1 was used to classify the participants in experiment 2. ROC analyses were conducted for each classifier and the

corresponding areas under the ROC curves (AUCs) were extracted. Subsequently, the

significance of the AUCs was determined using the Mann-Whitney U-Statistic (Mason and

Graham, 2002), and the resulting *p*-values were adjusted using the Benjamini and Hochberg

procedure (Benjamini and Hochberg, 1995). The sensitivity and specificity of each

classification procedure was obtained through the optimal thresholds derived from experiment

1.

## Mutual Magnitude-Phase Transplants

The discrete Fourier transform decomposes a signal (e.g., scalp-recorded voltage at an

electrode) into a set ("spectrum") of complex sinusoidal oscillations, each possessing a

different frequency. A constituent oscillation has a magnitude, indicating the oscillation

strength, as well as a phase, indicating its relative position in time or space (Grami, 2016). The

magnitude and phase spectra of a signal generally carry distinct information about the signal.

For example, in an image, most of the global color information is held in the magnitude

spectrum of the color channels, whereas the phase spectrum contains most of the form

information (Weinhaus, 2011). It is well known that swapping the magnitude and phase

information between two images creates chimeric images that generally inherit the color

palette from the magnitude spectrum and the form information from the phase spectrum (Fig.

2a). To probe the relative contribution of signal strength vs. coherent timing to the P1 wave

reduction observed in CC individuals, we adapted this signal processing method for event-

related potentials at the level of individual trials. Here we briefly describe this method, which

we term *mutual magnitude-phase transplants* (MuTx) in the present context.

Event-related potentials (ERP) are derived by averaging across numerous trials (Fig. 2 b – c).

For each CC individual and their matched normally sighted control participant (MCC), we

derived the MuTx waves from a set balanced number of trials, as in all the other analyses

reported in this article. First, for both individuals, we applied a Hann window to each individual

trial and calculated the discrete Fourier transform with the FFT method in Matlab (R2022a,

Natick, MA). Then, the magnitude and the phase spectra of the FFT was calculated, e.g.,

351
$$x_i(t) \xrightarrow{FFT} X_i(\omega) := |X_i(\omega)|e^{j \cdot \angle X_i(\omega)}$$

352
$$y_i(t) \xrightarrow{FFT} Y_i(\omega) := |Y_i(\omega)|e^{j \cdot \angle Y_i(\omega)}$$

353      where $|X_i(\omega)|$ and $\angle X_i(\omega)$ are the magnitude and phase spectra of the $i^{th}$ trial of the CC

354      individual, $|Y_i(\omega)|$ and $\angle Y_i(\omega)$ are the magnitude and phase spectra of the $i^{th}$ trial of the MCC

355      individual, and $j$ is the imaginary unit, i.e. $j^2 = -1$.

356      After this step, the phase of the CC and the MCC individuals were exchanged, and an inverse

357      fast Fourier transform (IFFT) yielded two MuTx trials (Fig. 2 d – e):

358
$$|X_i(\omega)|e^{j \cdot \angle Y_i(\omega)} \xrightarrow{IFFT} |x_i| \angle y_i(t)$$

359
$$|Y_i(\omega)|e^{j \cdot \angle X_i(\omega)} \xrightarrow{IFFT} |y_i| \angle x_i(t)$$

The first of the two MuTx trials, $|x_i| \angle y_i(t)$, inherited the magnitude information from the CC

individual's $i^{th}$ trial, but the phase information from the MCC individual's $i^{th}$ trial. Conversely,

the second MuTx trial, $|y_i| \angle x_i(t)$, had the magnitude of the MCC individual's $i^{th}$ trial but the

phase of the CC individual's $i^{th}$ trial. Averaging across these trials provided two MuTx ERPs,

which we term $|CC| \angle MCC$ and $|MCC| \angle CC$. Here, $|CC| \angle MCC$ is the ERP derived from the trials

with the magnitude of CC individuals and the phase of MCC individuals. Conversely, $|MCC| \angle CC$

had the magnitude of MCC individuals and the phase of CC individuals. If a generally lower

activation magnitude across trials would be the main driver of the P1 reduction in CC

individuals, we would expect $|CC| \angle MCC$ to have lower P1 values than $|MCC| \angle CC$. In contrast,

if the P1 wave reduction would be driven mainly by a reduced phase alignment (i.e.,

coherence) in the CC individuals, then $|MCC| \angle CC$ would exhibit a reduced P1 wave compared

to the $|CC| \angle MCC$.

The role of magnitude and phase information in a constellation of sight recovery group and

their control participants (i.e., CC vs. MCC or DC vs. MCC) was investigated using the mean

of the standardized P1 wave activation (i.e., *z*-score) over the 13 posterior electrodes (120 –

170 ms post-stimulus at electrodes TP9/TP10, CP1/CP2, CP5/CP6, P7/P8, P3/P4, Pz, and

O1/O2) as the dependent variable (Sourav et al., 2020).

**Figure 2. Derivation of mutual magnitude-phase transplants (MuTx).**

**a** An image processing example motivating the MuTx approach. Two-dimensional fast Fourier transform (FFT) of signals, here public-domain images of a house (Weber, 2018) and an imaginary animal (Roy, 1946), decomposes each image into a set of sinusoidal spatial oscillations (here, in two spatial direction *u* and *v*). Each oscillation (i.e. each member of the set of sinusoidal oscillations derived by FFT) have a magnitude, indicating the strength of oscillation, as well as a phase, indicating its relative position. As an experiment, the magnitude and phases of the two images can be exchanged. After this interchange, inverse fast Fourier transforms (IFFT) yield the mutual magnitude-phase transplants (MuTx). It can be appreciated that for images, most of the shape information seems to be contained in the signal phase, whereas color information seems to rely more on the signal magnitude. **b − c** Derivation of magnitude and phase information via one-dimensional FFT of the individual trials of the EEG epochs respectively in a CC individual and their matched normally sighted control participant (MCC), with the mean across trials (unaltered event related potentials, ERPs) displayed. **d − e** After exchanging the magnitude and phase spectra, the inverse FFT yields two MuTx waves with the magnitude of the CC individual but the phase of the MCC individuals ($|CC|\angle MCC$) and vice versa ($|MCC|\angle CC$). It can be appreciated that phase information seems to drive most of the P1 magnitude difference (grey bar, 120 − 170 ms post-stimulus) between the CC and the MCC participant.

The *m*ean *p*osterior *P*1 activation, termed MPP1, was investigated in Bayesian linear models

of the form `MPP1 ~ magnitude × phase`. The factors *magnitude* and *phase* had two levels

each, which indicated the source group for the magnitude or phase information, namely *sight*

*recovery* and *control*. Separate models were run for the CC vs. MCC and the DC vs. MDC

comparisons.

In addition to this model, we used a reparametrized model with custom contrasts (Schad et al.,

2020) to test for three differences of interest. For the CC vs. MCC comparison, the first of these

differences tested whether the CC group exhibited a P1 wave reduction (contrast: MCC – CC).

The other two contrasts tested the effect of "transplanted" phase on the P1 wave amplitude

(contrasts: |CC|∠MCC – CC, and |MCC|∠CC – MCC). Analogous models were run for the DC

vs. MDC comparison. In all cases, weak priors were set with the auto_prior() function of the

*sjstat* library (Lüdecke, 2018).

## P1 Wave Source Analysis

Average-referenced ERP voltage topographies were converted to a 15,002-point,

unconstrained cortical source space using the sLORETA method for EEG inverse modeling

(Pascual-Marqui, 2002), implemented in the Brainstorm software package (Tadel et al., 2011).

For calculating the forward model used in the sLORETA modeling, we started with the *New*

*York Head* Model with six tissue types: scalp, skull, cerebrospinal fluid, gray matter, white

matter, and air cavities (Huang et al., 2016). The forward model was then calculated with the

DUNEuro partial differential equation solver package using realistic conductivity values for the

tissues and with the electrode locations in the present study (Piastra et al., 2021; Schrader et

al., 2021). An identity matrix was used as the noise covariance.

Similar to the P1 wave analysis at the sensor level, group comparisons (CC vs. MCC, DC vs.

MDC) of source activity were performed in a posterior cortical region of interest that included

the occipital, temporal and parietal regions, excluding the postcentral sulcus and the limbic

lobe. To this end, the mean sLORETA activation across the posterior region was extracted,

with automatic sign inversion implemented in the Brainstorm package for oppositely oriented

sLORETA solutions. The vector norm, i.e. magnitude of the mean sLORETA activation, served

as the dependent variable for a Bayesian lognormal model of the form norm_of_mean ~

Group; weakly informative priors were set for the intercept and groups, respectively $N(0, 10)$

and $N(0, 2.5)$. For evidence synthesis across experiments, we added a random intercept of

participants to take participation in both experiments into account, i.e., `norm_of_mean ~ Group`

`+ (1|ParticipantID)`. Substantial evidence was ascertained when Bayes factors exceeded

3 and when the 95% highest-density interval of the posterior was situated outside of a region

of practical equivalence of [-0.1, 0.1] (ROPE test, Makowski et al., 2019).

## Results

### Time-Frequency Analysis

### Experiment 1

*$\Delta P_{rel}$.* The cluster-based permutation test in the whole post-stimulus time-frequency range

(latency: 0 – 700 ms, frequency: 2 – 40 Hz) did not reveal any significant difference in $\Delta P_{rel}$

between the cataract reversal groups and their respective control groups (Fig. 3 a – f); CC vs.

MCC, most negative cluster mass (± 95% confidence interval, CI): -369.014 ± 0.006, *p* = .118,

most positive cluster mass: 28.399, the parametric threshold for clustering was not exceeded;

DC vs. MDC,  most negative cluster mass: -241.928 ± 0.009, *p* = .293, most positive cluster

mass: 37.595, the parametric threshold for clustering was not exceeded.

Running cluster-based permutation in the "alpha blocking" time-frequency window (time: 0.3 –

0.7 s, frequency: 8 – 12 Hz)  as in Bottari et al. (2016), we once more did not find  evidence

for a difference in $\Delta P_{rel}$ in the CC vs. MCC analysis (most negative cluster mass =

-29.842 ± 0.009, *p* = .304, most positive cluster mass = 0.043, threshold for clustering not

exceeded), nor in the DC vs. MDC comparison (most negative cluster mass = -6.959 ± 0.010,

*p* = .609, for positive cluster formation no time-frequency point significant at individual level).

*Inter-trial Phase Coherence (ITPC).* In the CC group, significant evidence for a lower ITPC

compared to the matched typically sighted control group (MCC) emerged (Fig. 3 g – l); most

negative cluster mass (± 95% CI): -405.176 ± 0.003, *p* = .025, most positive cluster mass (±

95% CI): 93.893 ± 0.010, *p* = 0.422. In contrast, between the DC and the MDC group, we did

not observe a significant difference in ITPC; most negative cluster mass (± 95% CI): -32.622,

the parametric threshold for clustering was not exceeded, most positive cluster mass (± 95%

CI): 120.391 ± 0.008, *p* = .226.

**Figure 3. Time-Frequency Analysis of Relative Power Change and Phase Coherence at Posterior Electrodes in Experiment 1**.

**a** Relative power change ($\Delta P_{rel}$, unitless) compared to baseline (-700 to -300 ms) in the matched control group for congenital cataract reversal individuals (MCC). **b** $\Delta P_{rel}$ in the group of congenital cataract reversal individuals (CC). **c** Relative power change difference between the MCC and the CC group, $\Delta P_{rel, MCC} - \Delta P_{rel, CC}$. **d – f**. Relative power changes in the matched control group for developmental cataract reversal individuals (MDC, $\Delta P_{rel, MDC}$), in the group of developmental cataract reversal individuals (DC, $\Delta P_{rel, DC}$), and their difference ($\Delta P_{rel, MDC} - \Delta P_{rel, DC}$). **g – i**. Inter-trial phase coherence (ITPC) respectively in the MCC group, the CC group, and their difference. **j – l**. ITPC respectively in the MDC and the DC group, and their difference. For cluster-based permutation tests spanning the time-frequency range of [0 – 0.7 s] × [2 – 40 Hz], the time-frequency extend of a cluster driving a significant result has been outlined. The vertical dotted lines at 0 s indicate stimulus onset. $\Delta P_{re}$ and ITPC shown in the plots are averaged over the 13 posterior electrodes (TP9/TP10, CP1/CP2, CP5/CP6, P7/P8, P3/P4, Pz, and O1/O2). Note that $\Delta P_{rel}$ and ITPC have different scales (respectively -0.8 – 0.8 and -0.05 – 0.3). For each variable, the difference between the groups is displayed on a scale that is half of that used for the variable.

Experiment 2

*$\Delta P_{rel}$.* The cluster-based permutation test in the whole post-stimulus time-frequency range

(latency: 0 – 700 ms, frequency: 2 – 40 Hz) once again did not find any significant difference

in $\Delta P_{rel}$ between the CC and the MCC group (Fig. 4 a – f), most negative cluster mass (± 95%

confidence interval, CI): -39.6720, the parametric threshold for clustering was not exceeded;

most positive cluster mass = 155.132 ± 0.010, *p* = .443. In contrast to the CC vs. MCC group

comparison (and in contrast to experiment 1), we found a reduced $\Delta P_{rel}$ in the DC group

compared to the MDC group, most negative cluster mass = -863.053 ± 0.004, *p* = .040; most

positive cluster mass = 29.512, parametric threshold for clustering not exceeded.

Employing cluster-based permutation in the "alpha blocking" time-frequency window (time: 0.3

– 0.7 s, frequency: 8 – 12 Hz)  as in Bottari et al. (2016), as in experiment 1, no evidence of a

difference in $\Delta P_{rel}$ emerged between the CC and the MCC group (for negative cluster

formation, no time-frequency point significant at individual level; most positive cluster mass =

40.989 ± 0.008, *p* = .197).

*Inter-trial Phase Coherence (ITPC).* In the CC group, significant evidence for a lower ITPC

compared to the matched sighted control group (MCC) once again was found in experiment 2

(Fig. 4 g – l); most negative cluster mass (± 95% CI): -233.112 ± 0.003, *p* = .031, most positive

cluster mass = 28.907, parametric threshold for clustering not exceeded. Similar as in

experiment 1, we did not observe a significant difference in ITPC between the DC and the

MDC groups; most negative cluster mass (± 95% CI) = -106.505 ± 0.009, *p* = .266; most

positive cluster mass = 28.672, parametric threshold for clustering not exceeded.

**Figure 4. Time-Frequency Analysis of Relative Power Change and Phase Coherence at Posterior Electrodes in Experiment 2**.

**a** Relative power change ($\Delta P_{rel}$, unitless) compared to baseline (-700 to -300 ms) in the matched control group for congenital cataract reversal individuals (MCC). **b** $\Delta P_{rel}$ in the group of congenital cataract reversal individuals (CC). **c** Relative power change difference between the MCC and the CC group, $\Delta P_{rel, MCC} - \Delta P_{rel, CC}$. **d – f**. Relative power changes in the matched control group for developmental cataract reversal individuals (MDC, $\Delta P_{rel, MDC}$), in the group of developmental cataract reversal individuals (DC, $\Delta P_{rel, DC}$), and their difference ($\Delta P_{rel, MDC} - \Delta P_{rel, DC}$). **g – i**. Inter-trial phase coherence (ITPC) respectively in the MCC group, the CC group, and their difference. **j – l**. ITPC respectively in the MDC and the DC group, and their difference. For cluster-based permutation tests spanning the time-frequency range of [0 – 0.7 s] × [2 – 40 Hz], the time-frequency extend of a cluster driving a significant result has been outlined. The vertical dotted lines at 0 s indicate stimulus onset. $\Delta P_{re}$ and ITPC shown in the plots are averaged over the 13 posterior electrodes (TP9/TP10, CP1/CP2, CP5/CP6, P7/P8, P3/P4, Pz, and O1/O2). Note that $\Delta P_{rel}$ and ITPC have different scales (respectively -0.8 – 0.8 and -0.05 – 0.3). For each variable, the difference between the groups is displayed on a scale that is half of that used for the variable.

## Classification Based on Time-Frequency Features

In addition to the cluster-based permutation tests for group differences in spectral dynamics, we investigated whether CC individuals could be classified based on biomarkers in the time-frequency domain. To prevent overfitting in the high-dimensional spectral feature space, the classifications were performed based on the loadings of the first three principal components at

posterior electrodes (39 features), as well as based on the average loadings across the

posterior electrodes (3 features). In total, 8 classifications were performed: 2 classification

types (CC vs. non-CC, and CC vs. DC) × 2 feature classes ($\Delta P_{rel}$ and ITPC) × 2 loading types

(loadings at all posterior electrodes and loadings averaged over posterior electrodes). All

classifiers were trained on data from experiment 1 and validated on data from experiment 2.

Validation results are all displayed in Table 5, and the corresponding ROC curves are depicted

in Fig. 6, panels a – h. Additionally, the SVM decision boundaries for classification based on

ITPC- and power-based component loadings averaged across posterior electrodes are

visualized in Fig. 6, panels i – j in the three-dimensional feature space.

**Table 5.**

Principal Component Based Classification.

*CC vs. non-CC classification*

| Domain | *n*(Features) | AUC | [95% CI] | *p*-value | Sensitivity | Specificity |
|--------|--------------|------|-----------|-----------|-------------|-------------|
| Power | 39 | .660 | [.499; .821] | .084 | .462 | .814 |
| Power | 3 | .528 | [.362; .694] | .774 | .538 | .558 |
| ITPC | 39 | .748 | [.575; .921] | .006* | .692 | .767 |
| ITPC | 3 | .766 | [.624; .907] | .003* | .462 | .814 |

*CC vs. DC classification*

| Domain | *n*(Features) | AUC | [95% CI] | *p*-value | Sensitivity | Specificity |
|--------|--------------|------|-----------|-----------|-------------|-------------|
| Power | 39 | .718 | [.521; .915] | .052 | .769 | .467 |
| Power | 3 | .621 | [.397; .844] | .235 | .692 | .400 |
| ITPC | 39 | .718 | [.509; .927] | .052 | .692 | .800 |
| ITPC | 3 | .728 | [.536; .921] | .041* | 1.000 | .200 |

## CC vs. non-CC Classification

Classification of CC participants from non-CC participants (including DC and normally sighted

control individuals) in experiment 2 achieved statistically significant performance based on

ITPC component loadings at all posterior electrodes (AUC = .748, 95% CI = [.575 – .921], p =

.006), but not based on the component loadings derived from $\Delta P_{rel}$ (AUC = .660, 95% CI =

[.499 – .821], p = .084). Interestingly, a robust performance was observed as well when

classification was based on the ITPC component loadings averaged over all posterior

electrodes, i.e., based on 3 instead of 39 features (AUC = .766, 95% CI = [.624– .907], p =

.003). In contrast, $\Delta P_{rel}$-based classification performance dropped close to chance level when

classifying employed average loadings (AUC = .528, 95% CI = [.362– .694], p = .774). The

visualization of the corresponding three-dimensional SVM decision boundaries demonstrates

that only the phase-based group differences generalize from the training (experiment 1) to the

validation dataset (experiment 2, see Fig. 6 i – j). The optimal thresholds derived from

experiment 1 translated to experiment 2 to a moderate degree (Power et al., 2013).

### CC vs. DC Classification

Classification of the CC from the DC individuals in the validation experiment 2 was not

statistically significant using either the ITPC component loadings (AUC = .718, 95% CI = [.509

– .927], $p$ = .052) or the $\Delta P_{rel}$ component loadings at all posterior electrodes (AUC = .718, 95%

CI = [.521 – .915], $p$ = .052). However, when averaging the component loadings across the

posterior electrodes, ITPC information successfully classified the CC from the DC individuals

(AUC = .728, 95% CI = [.536 – .921], $p$ = .041). In contrast, using average component loadings

for $\Delta P_{rel}$, the classification performance remained close to chance level (AUC = .621, 95% CI

= [.397 – .844], $p$ = .294).

### Phase Information is the Main Driver of the P1 Wave Reduction After Sight Recovery Following Transient Congenital Blindness

In a novel application, we exchanged the magnitude and phase information between each

sight-recovery individual and their control participant. ERPs were derived from the trials of the

mutual magnitude-phase transplanted participants (MuTx) and were systematically compared

to those of the original participants (Fig. 6 – 7). We reasoned that if phase information were

the main driving factor behind the P1 wave reduction observed in the CC group, the MuTx

group comprising the magnitude information of the MCC individuals but the phase information

of the CC individuals (denoted |MCC|∠CC) would exhibit a P1 wave reduction. Conversely, the

MuTx group consisting of chimeric participants with the magnitude information from the CC

individuals but the phase information from the MCC individuals (denoted |CC|∠MCC) would be

expected to exhibit typical P1 waves. This was indeed the case: in experiment 1, the grand

average visual ERPs qualitatively resembled the group whose phase was used to derive the

visual ERP (Fig. 6a). This held true too for the P1 wave topography: in the |CC|∠MCC group's

**Figure 5. Receiver Operating Characteristic (ROC) Curves for Principal Component Based Classification.**

**a-d** ROC curves for the classification of CC vs. non-CC individuals based on component loadings for relative power change ($\Delta P_{rel}$) and ITPC features at all posterior electrodes ('All Loadings') and loadings averaged over posterior electrodes ('Mean Loadings'). Black dots indicate the optimal classification threshold derived from experiment 1. **e-f** ROC curves for the CC vs. DC classification using the same features. **i** The SVM decision boundary for CC vs. non-CC classification based on ITPC component loadings averaged over posterior electrodes (i.e., 3 features) on the training and validation dataset. **j** The SVM decision boundary for CC vs. Non-CC classification based on component loadings averaged over posterior electrodes for $\Delta P_{rel}$ on the training and validation dataset.

P1 wave topography resembled that of the MCC group, and the |MCC|∠CC group's topography resembled that of the CC group (Fig. 6b). Finally, the Bayesian mixed model investigating the effect of oscillatory magnitude and phase indicated that the mean of the standardized P1 values over posterior electrodes (MPP1) depended on the phase, but not magnitude information content for the CC, MCC, |CC|∠MCC, and |MCC|∠CC constellation; there was a decisive main effect of phase, $\beta_{phase}$ [95% credible interval] = 0.37 [0.23, 0.52], $BF_{10}$ = 1478.71, positive ROPE test. In contrast, there was no substantial evidence for the role of magnitude influencing the MPP1 values in this constellation, $\beta_{mag}$ = 0.01 [-0.14, 0.15], $BF_{10}$ = 0.047, and

575    no substantial interaction between the factors magnitude and phase, $\beta_{mag \times phase}$ = -0.01 [-0.15,

576    0.14], $BF_{10}$ = 0.048.

The same model equipped with custom contrasts provided very strong evidence that the MPP1

values of the CC group was lower than the MPP1 values of the MCC group, $\beta$ = -0.74 [-1.13, -

0.34], $BF_{10}$ = 56.123, positive ROPE test. In addition, the |CC|∠MCC group had a higher MPP1

value than the CC group, $\beta$ = 0.75 [0.34, 1.15], $BF_{10}$ = 71.263, positive ROPE test. Finally,

there was very strong evidence that |MCC|∠CC group's MPP1 values were lower compared to

the MPP1 values of the MCC group, $\beta$ = -0.71 [-1.11, -0.31], $BF_{10}$ = 48.557, positive ROPE

test, indicating that difference in phase information likely caused the P1 wave attenuation in

the CC group.

These results were replicated in the validation dataset from experiment 2, both qualitatively

(Fig. 7 a – d) as well as quantitatively (Fig. 7e). Specifically, a Bayesian mixed model once

**Figure 7. P1 waves of the original groups vs. the mutual magnitude-phase transplant (MuTx) groups in Experiment 2 (Validation Dataset)**

**a** Grand average visual event-related potentials (VERP) at the posterior electrode P8 for the sight recovered bilateral congenital cataract group (CC), their matched control group (MCC), and the MuTx groups with the magnitude (|○|) and phase (∠○) information swapped, i.e. |CC|∠MCC and |MCC|∠CC. The color-coding is according to the group's magnitude information. Ribbons indicate the standard error of the mean. The gray bar indicates the P1 wave time range (120 – 170 ms). **b** Topographic plot of the grand average P1 wave for the CC, MCC, |CC|∠MCC, and |MCC|∠CC groups, averaged between 120 – 170 ms after stimulus presentation. **c** Grand average VERPs at the posterior electrode P8 for the typically sighted recovered bilateral developmental cataract group (DC), their matched control group (MDC), and the MuTx groups |DC|∠MDC and |MDC|∠DC. **d** Topographic plot of the grand average P1 wave for the DC, MDC, |DC|∠MDC, and |MDC|∠DC groups, averaged between 120 – 170 ms after stimulus presentation. **e** Violin plot of the mean of the standardized P1 topography, with the mean calculated over the posterior electrodes. The mean posterior P1 value (MPP1) was developed and validated as a biomarker based on the present experiments (Sourav et al., 2020). The threshold for diagnosing a CC individual (from experiment 1) is shown as a horizontal dotted line, and the number of individuals above and below the diagnosis line is indicated near the top and bottom border. Red circles denote the means, and the error bars indicate 95% bootstrapped confidence intervals for the means. Asterisks indicate substantial evidence (or above). Individual data points were jittered horizontally for readability.

again indicated that the MPP1 values for the CC, MCC, |CC|∠MCC, and |MCC|∠CC

constellation depended on the phase but not magnitude information; there was a decisive main

effect of phase, $\beta_{\text{phase}}$ [95% credible interval] = 0.27 [0.14, 0.40], $BF_{10}$ = 117.02, positive ROPE

test. In contrast, there was no substantial evidence for the role of magnitude influencing the

MPP1 values in this constellation, $\beta_{\text{mag}}$ = 0.06 [-0.07, 0.19], $BF_{10}$ = 0.075, and no substantial

interaction between the factors magnitude and phase, $\beta_{\text{mag} \times \text{phase}}$ = -0.01 [-0.14, 0.12], $BF_{10}$ =

0.049.

The same model equipped with custom contrasts provided very strong evidence that the MPP1

values of the CC group were lower than those of the MCC group, $\beta$ = -0.65 [-1.01, -0.28], $BF_{10}$

= 39.909, positive ROPE test. In addition, there was substantial evidence that the |CC|∠MCC

group had higher MPP1 values than the CC group, $\beta$ = 0.55 [0.18, 0.91], $BF_{10}$ = 9.893, positive

ROPE test. Finally, there was substantial evidence that the |MCC|∠CC group's MPP1 values

were lower compared to those of the MCC group, $\beta$ = -0.52 [-0.88, -0.15], $BF_{10}$ = 6.836, positive

ROPE test.

When we tested the DC, MDC, |DC|∠MDC, and |MDC|∠DC group constellation, we found no

evidence in any of the two experiments that the MPP1 values substantially differed between

the groups (all non-intercept $\beta$s had $BF_{10}$ < 1, no positive ROPE tests). Unsurprisingly, there

was no substantial evidence that the MPP1 depended on phase or the magnitude information

in this constellation (all non-intercept $BF_{10}$ < 1, no positive ROPE tests).

## Source Localization of the P1 wave

In experiment 1, the Bayesian lognormal model indicated that the magnitude of mean posterior

sLORETA activation substantially differed between the CC and the MCC groups (Fig. 8), $\beta_{CC\text{-}MCC}$

[95% CrI] = -1.03 [-1.76, -0.30], $BF_{10}$ = 6.19, positive ROPE test. In contrast, no substantial

difference in magnitude of mean posterior sLORETA activations was observed when

comparing the DC vs. the MDC group; $\beta_{DC\text{-}MDC}$ = -0.11 [-0.63, 0.41], $BF_{10}$ = 0.111, negative

ROPE test (3.06 % of posterior distribution in the ROPE region).

In experiment 2, we did not find any substantial evidence that the magnitude of mean posterior

sLORETA activation substantially differed between the CC and the MCC groups (see Fig. 9),

$\beta_{CC\text{-}MCC}$ [95% CrI] = -0.58 [-1.19, -0.03], $BF_{10}$ = 0.772, negative ROPE test (0.46% of posterior

distribution in the ROPE region). No substantial difference in magnitude of mean posterior

sLORETA activations was observed when comparing the DC vs. the MDC groups as well; $\beta_{DC\text{-}MDC}$

$_{MDC}$ = -0.11 [-0.63, 0.41], $BF_{10}$ = 0.111, negative ROPE test (3.06 % of posterior distribution in

the ROPE region).

**Figure 8. Source Localization of the P1 Wave in Experiment 1.**

**a** Magnitude (vector norm) of the grand average source localization solution for the P1 wave (mean over 120 – 170 ms post-stimulus) obtained with the sLORETA method (unitless; Pascual-Marqui, 2002; Tadel et al., 2011) respectively for the CC and the MCC groups (left and center columns), and the MCC – CC group difference of grand average norms (right column). **b** Magnitude of the grand average sLORETA solution for the P1 wave (mean over 120 – 170 ms post-stimulus) respectively for the DC and the MDC groups (left and center columns), and the MDC – DC group difference of grand average norms (right column). **c** Posterior region of interest (ROI) for testing group differences, which included the occipital, temporal and parietal regions excluding the postcentral sulcus and the limbic lobe. **d** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the CC and the MCC group (scaled to unit variance). **e** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the DC and the MDC group (scaled to unit variance). Asterisk (*) indicates substantial evidence for a group difference (Bayes factor > 3 and positive ROPE test, lognormal model).

The Bayesian models, being parametric, allowed evidence synthesis across the experiments, which on the one hand led to a more stable parameter estimation and on the other hand permitted reconciling partially different result patterns for the CC vs. MCC group difference across the experiments. Combining evidence across both experiments with Bayesian lognormal mixed models, we found substantial evidence for a group difference in the mean posterior sLORETA activation strength between the CC and the MCC group, $\beta_{CC\text{-}MCC}$ [95% CrI] = -0.77 [-1.26, -0.27], $BF_{10}$ = 9.01, positive ROPE test (Fig. 9). In contrast, once again no substantial difference in the norm of mean posterior sLORETA activations emerged when comparing the DC vs. the MDC group; $\beta_{DC\text{-}MDC}$ = -0.10 [-0.46    0.27], $BF_{10}$ = 0.085, negative ROPE test (3.97 % of posterior distribution in ROPE region).

**Figure 9. Source Localization of the P1 Wave in Experiment 2 and Evidence Synthesis Across Experiments 1 and 2.**

**a** Magnitude (vector norm) of the grand average source localization solution for the P1 wave (mean over 120 – 170 ms post-stimulus) obtained with the sLORETA method in experiment 2 (unitless; Pascual-Marqui, 2002; Tadel et al., 2011) respectively for the CC and the MCC groups (left and center columns), and the MCC – CC group difference of grand average norms (right column). **b** Magnitude of the grand average sLORETA solution for the P1 wave (mean over 120 – 170 ms post-stimulus) respectively for the DC and the MDC groups in experiment 2 (left and center columns), and the MDC – DC group difference of grand average norms (right column). **c** Posterior region of interest (ROI) for testing group differences, which included the occipital, temporal and parietal regions excluding the postcentral sulcus and the limbic lobe. **d** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the CC and the MCC group in experiment 2 (scaled to unit variance). **e** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the DC and the MDC group in experiment 2 (scaled to unit variance). **f** Grand average source localization solution across all participants and groups in experiment 1 and experiment 2. Black borders indicate the loci of the highest 5% sLORETA magnitude. Relevant cortical regions from the Desikan-Killiany atlas are depicted on the left side of the sLORETA solution; FuG: Fusiform gyrus, LNG: Lingual gyrus, LOC: Lateral occipital cortex. **g** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the CC and the MCC group across experiment 1 and experiment 2, scaled to unit variance. All unique participants are depicted, for participants who took part in both experiments, mean values are presented. **h** Violin plot of the magnitude of the mean sLORETA activation in the posterior ROI for the DC and the MDC group across experiment 1 and experiment 2, scaled to unit variance. All unique participants are depicted, for participants who took part in both experiments, mean values are presented. Asterisk (*) indicates substantial evidence for a group difference (Bayes factor > 3 and positive ROPE test, lognormal model)

The grand average sLORETA map across all participants and both experiments indicated posterior regions as the source of the P1 wave (Fig. 9). The maximum 5% regions of this activation were located in the right ventral extrastriate visual cortical regions, including the fusiform gyrus (FuG), lingual gyrus (LNG) and lateral occipital cortex (LOC) according to the Desikan-Killany atlas (Desikan et al., 2006).

## Discussion

Optimal functioning of neural circuits requires appropriate neural activation strength as well as precisely timed excitation and inhibition. Since sufficiently temporally resolved neural timing information is hard to access with brain imaging methods (e.g., fMRI), and as activation strength and timing information is intermingled in trial-averaged electrophysiological measures such as those derived from EEG/MEG, how neural circuit timing depends on early experience is still an unanswered question. A recent study involving ferrets has suggested that particularly temporal processing stability in the visual cortex may depend on visual input after birth (Trägenap et al., 2025). However, in non-human animal research the degree of recovery after visual deprivation and subsequent sight restoration is difficult to investigate. Thus, it is currently unknown whether a sensitive period for temporally tuned neural circuit processing exists; sensitive periods are defined by incomplete recovery after sensory restitution (Kundsen, 2004).

A transient phase of congenital visual deprivation in humans has been found to permanently alter visual cortical structure and function (Le Grand et al., 2003; Röder et al., 2021; Hölig et al., 2023; Pedersini et al., 2023; Sourav et al., 2024). Of specific relevance is the earliest electrophysiological maker of higher-order, extrastriate visual cortical activity, the P1 wave, which peaks at around 150 ms after the visual stimulation onset (Di Russo et al., 2002). The P1 wave was found to be markedly attenuated in sight-recovery individuals with a history of dense, bilateral congenital cataracts (Sourav et al., 2020). Yet, to the best of our knowledge, no study has systematically examined the trial-to-trial temporal consistency of cortical oscillations in individuals with reversed congenital blindness to disentangle to role of cortical activation strength vs. impaired temporal processing stability as a neural mechanism of their extrastriate visual processing impairment. A dissociation of both aspects of neural processing is crucial for the understanding of the precise mechanisms of experience-dependent brain development.

The present study investigated group differences in task-related oscillatory activation strength and timing stability in individuals with reversed congenital cataracts (CC), individuals with

reversed developmental, i.e., late-onset cataracts (DC), and matched normally sighted controls

(SC). Three methodological approaches were implemented: First, in time-frequency

representations of cortical activity during the processing of simple visual stimuli, the across-

trial consistency of visual processing, indexed by the inter-trial phase coherence, was

significantly attenuated in the CC group compared to their matched normally sighted controls

(MCC group). In contrast, no differences in relative power change from baseline between these

groups was observed. Second, further supporting these findings, the CC individuals could be

classified from non-CC individuals (i.e., a combined group of sighted controls as well as DC

individuals) only by using phase information but not by using oscillatory power information.

Third, through a novel approach that we term *mutual magnitude-phase transplants,* we

swapped the oscillatory activation magnitude and phase information between each CC

individual and their matched sighted controls. This analysis revealed that combining the phase

information of the CC individuals with the magnitude information of the MCC individuals, but

not vice versa, yielded the attenuation of the extrastriate P1 wave in visual event-related

potentials (ERPs) which is typical for the CC group. The convergent evidence from all three

methodological approaches was replicated in a second experiment. Thus, task-related

oscillatory timing information, but not oscillatory strength, appears to critically depend on early

experience and therefore seems to be a key mechanism of sensitive period plasticity.

Between the DC individuals and their matched control group (MDC), cluster-based permutation

tests did not reveal significant temporal stability differences in any experiment, indicating that

the decrease in task-related oscillatory phase synchronization in the CC vs. MCC group

comparison is linked to visual deprivation during an early sensitive period. This selective

impairment for CC individuals excludes alternative explanations such as unspecific effects

related to cataract surgery. Additionally, phase, but not power information averaged over the

posterior electrodes allowed us to classify CC from DC individuals. Lastly, *mutual magnitude-*

*phase transplant* analyses revealed no effect of exchanging the magnitude and phase

information between the DC and the MDC individuals, confirming that temporal processing

impairments were specific to the CC group.

Compared to striate cortical processing, extrastriate cortical functions have been reported to

exhibit higher vulnerability to a period of transient congenital visual deprivation (McKyton et

al., 2015; Sourav et al., 2018, 2020; Pitchaimuthu et al., 2021). Consistent with this line of

evidence, we previously found a robustly reduced P1 wave in CC individuals' visual ERPs

(Sourav et al., 2020), which has been hypothesized to originate in extrastriate cortical areas

(Di Russo et al., 2002; Miller et al., 2015). Moreover, this strong and reliable reduction of the

P1 wave in sight recovery individuals allowed a robust retrospective diagnosis of a congenital

vs. developmental etiology years after sight restoration (Sourav et al., 2020). In the present

study, source localization of the P1 wave indicated areas of maximum activation in extrastriate

cortical areas in the right (ipsilateral) hemisphere, including the fusiform gyrus, lingual gyrus

and lateral occipital cortex. These results are by and large consistent with previous EEG/fMRI

work (Di Russo et al., 2002). Moreover, augmenting the sensor-level findings, evidence

synthesis across the two experiments of the present study revealed substantial evidence for a

trial-averaged reduction of the P1 wave in the CC group at source level as well.

Phase alignment of alpha oscillatory activity (8 – 12 Hz) has been argued to contribute to the

generation of the P1 wave (Klimesch et al., 2007), supported by the findings that alpha phase

alignment predicts the P1 wave latency (Gruber et al., 2005), and that alpha phase locking is

largest during the P1 wave latency (Klimesch et al., 2004; see Freunberger et al., 2008).

Consistent with this evidence, the principal component loading patterns of the inter-trial phase

coherence features for classifying CC individuals were predominantly located across the alpha

(8 – 12 Hz) and the theta (4 – 8 Hz) range and covered the P1 wave latency. Additionally,

significant inter-trial phase coherence differences between the CC and the MCC group in the

cluster-based permutation tests were associated with time-frequency clusters that spanned

the P1 latency and the alpha frequency range. In contrast to phase alignment across trials, the

P1 wave amplitude has been argued to be less affected by oscillatory power (Gruber et al.,

2005; Freunberger et al., 2008). In a related visual deprivation model of amblyopia, a reduction

of the P1 wave amplitude in the patients' amblyopic eye compared to the fellow eye was

associated with a broader trial-to-trial latency (but not amplitude) distribution (Bankó et al.,

2013). Consistent with these lines of evidence, we did not find any significant difference in the

post-stimulus relative power change in the CC vs. MCC group comparison. We did not observe

a significant classification performance based on relative power change as opposed to inter-

trial phase coherence information either. In fact, the *mutual magnitude-phase transplants*

analysis in the present study provided compelling evidence of the driving role of phase

information behind P1 reduction in the CC group: combining CC individuals' activation

magnitudes with the phase information of typically sighted controls substantially recovered

their P1 wave amplitude.

In the present study, we did not observe any relative oscillatory power impairments in the CC

group compared to the sighted controls during the processing of simple visual stimuli (gratings)

in any of the two experiments. While these results are inconsistent with those of Bottari et al.

(2016) in a group of CC individuals ($n$ = 12) during biological motion processing, they are

corroborated by Ossandón et al (2023): The latter authors have recently reported a spared

alpha power modulation in resting state EEG due to eye opening (compared to an eyes closed

condition) in CC individuals, despite an overall lower alpha oscillatory power level compared

to sighted controls.

Temporal stability, and its converse, temporal variability, is a critical marker of sensory

processing related to the alignment of dynamic neural states (Afraimovich et al., 2004; Jones

et al., 2007; Rabinovich et al., 2008). The converging results of the two present experiments

indicate that a disrupted temporal orchestration of cortical activity (Yusuf et al., 2022),

especially in higher sensory areas, and not a general attenuation of cortical oscillation may be

the most profound consequence of a period of congenital sensory deprivation. In this context

we note that feedforward and feedback connection profiles develop asynchronously, with

feedforward connectivity maturing earlier (Burkhalter et al., 1993). Moreover, there is evidence

that the elaboration of feedback connectivity more extensively depends on experience (Ibrahim

et al., 2021; Röder and Kekunnaya, 2021). Early sensory experience, thus, might be crucial

for temporally aligned cortical feedback signaling and in turn for the shaping of feedforward

activity (Ibrahim et al., 2021). A recent study that examined the development of the ferret visual

cortex strongly supports this account: in ferret kits whose eyes were prematurely opened,

visual cortical responses recorded with calcium imaging were found to be similarly strong, yet

temporally less stable across trials compared to visually experienced kits (Trägenap et al.,

2025). Crucially, a week of lid suture in the ferret kits, starting shortly before the time of natural

eye opening, disrupted the typical development of the temporal stability of cortical processing

when tested shortly after lid suture removal. The present results, on the one hand, extend

these findings in ferrets to human cortical development. On the other hand, the temporal

resolution of the Trägenap et al. (2025) study was much lower than that of the EEG recordings

we employed: The specific calcium indicator utilized by the authors (GCaMP6s, see Trägenap

et al., 2025) had a time constant of over a hundred milliseconds, that is, a temporal resolution

two orders of magnitude less precise than EEG which provides millisecond-level temporal

resolution (Buzsáki et al., 2012; Jercog et al., 2016). Thus, the results of the present study

extend the findings of Trägenap et al. (2025): The disrupted temporal stability of cortical

processing, resulting from aberrant early visual experience seems to appear already within

150 ms after visual stimulation onset in markers of extrastriate and downstream visual cortical

processing. Crucially, we investigated sight-recovery individuals at least two years after sight

restitution surgery. Thus, we are able to report evidence for a sensitive period in humans, that

is, for a persistent loss of temporal processing stability in visual neural circuits despite sight

restitution and extensive post-surgical visual experience.

In conclusion, we provide to the best of our knowledge the first report that temporal processing

stability in visual neural circuits critically depends on early experience in humans. This finding

of an impaired visual cortical processing stability in humans after a period of transient

congenital (but not developmental) pattern visual deprivation provides strong evidence for the

existence of a sensitive period for the development of precisely temporally orchestrated visual

cortical processing in humans. Our results suggest that aberrant temporal tuning of the visual

cortex, especially in extrastriate and higher-order visual areas, might underlie the numerous

persistent visual deficits observed after reversing congenital blindness.

## Acknowledgements

We thank D. Balasubramanian (L V Prasad Eye Institute, Hyderabad, India) for supporting this study. We acknowledge Davide Bottari for helping with data acquisition and for the initial EEG preprocessing pipeline. Seema Banerjee, Larissa Brockmann, Maria Guerreiro, Marlene Hense, Giulia Dormal, Siddhart Srivatsav Rajendran, Lisa Stockleben and Florian Süßer additionally helped with data acquisition in India and/or Germany. The study was funded the German Research Foundation (DFG, Ro 2625/10-1) and by the European Research Council grant ERC-2009-AdG 249425-*CriticalBrainChanges* to B.R.

## References

Afraimovich VS, Zhigulin VP, Rabinovich MI (2004) On the origin of reproducible sequential activity in neural circuits. Chaos: An Interdisciplinary Journal of Nonlinear Science 14:1123–1129.

Amedi A, Raz N, Pianka P, Malach R, Zohary E (2003) Early "visual" cortex activation correlates with superior verbal memory performance in the blind. Nature Neuroscience.

Aydore S, Pantazis D, Leahy RM (2013) A Note on the Phase Locking Value and its Properties. Neuroimage 74:231–244.

Bankó ÉM, Körtvélyes J, Németh J, Weiss B, Vidnyánszky Z (2013) Amblyopic deficits in the timing and strength of visual cortical responses to faces. Cortex 49:1013–1024.

Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B (Methodological) 57:289–300.

Birch EE, Cheng C, Stager DR, Weakley DR, Stager DR (2009) The critical period for surgical treatment of dense congenital bilateral cataracts. Journal of American Association for Pediatric Ophthalmology and Strabismus 13:67–71.

Bottari D, Kekunnaya R, Hense M, Troje NF, Sourav S, Röder B (2018) Motion processing after sight restoration: No competition between visual recovery and auditory compensation. NeuroImage 167:284–296.

Bottari D, Troje NF, Ley P, Hense M, Kekunnaya R, Röder B (2016) Sight restoration after congenital blindness does not reinstate alpha oscillatory activity in humans. Scientific Reports 6:24683.

Burkhalter A, Bernardo KL, Charles V (1993) Development of local circuits in human visual cortex. J Neurosci 13:1916–1931.

Buzsáki G, Anastassiou CA, Koch C (2012) The origin of extracellular fields and currents — EEG, ECoG, LFP and spikes. Nat Rev Neurosci 13:407–420.

Buzzell GA, Niu Y, Aviyente S, Bernat E (2022) A practical introduction to EEG Time-
Frequency Principal Components Analysis (TF-PCA). Developmental Cognitive
Neuroscience 55:101114.

De Heering A, Maurer D (2014) Face memory deficits in patients deprived of early visual
input by bilateral congenital cataracts. Developmental Psychobiology 56:96–108.

Delorme A, Makeig S (2004) EEGLAB: An open source toolbox for analysis of single-trial
EEG dynamics including independent component analysis. Journal of Neuroscience
Methods 134:9–21.

Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale
AM, Maguire RP, Hyman BT, Albert MS, Killiany RJ (2006) An automated labeling
system for subdividing the human cerebral cortex on MRI scans into gyral based
regions of interest. NeuroImage 31:968–980.

Di Russo F, Martínez A, Sereno MI, Pitzalis S, Hillyard SA (2002) Cortical sources of the
early components of the visual evoked potential. Human Brain Mapping 15:95–111.

Freunberger R, Höller Y, Griesmayr B, Gruber W, Sauseng P, Klimesch W (2008) Functional
similarities between the P1 component and alpha oscillations. European Journal of
Neuroscience 27:2330–2340.

Grami A (2016) Chapter 3 - Signals, Systems, and Spectral Analysis. In: Introduction to
Digital Communications (Grami A, ed), pp 41–150. Boston: Academic Press. Available
at: https://www.sciencedirect.com/science/article/pii/B978012407682200003X
[Accessed October 1, 2024].

Gruber WR, Klimesch W, Sauseng P, Doppelmayr M (2005) Alpha Phase Synchronization
Predicts P1 and N1 Latency and Amplitude Size. Cerebral Cortex 15:371–377.

Hölig C, Guerreiro MJS, Lingareddy S, Kekunnaya R, Röder B (2023) Sight restoration in
congenitally blind humans does not restore visual brain structure. Cerebral Cortex
33:2152–2161.

Huang Y, Parra LC, Haufe S (2016) The New York Head-A precise standardized volume
conductor model for EEG source localization and tES targeting. Neuroimage
140:150–162.

Ibrahim LA, Huang S, Fernandez-Otero M, Sherer M, Qiu Y, Vemuri S, Xu Q, Machold R,
Pouchelon G, Rudy B, Fishell G (2021) Bottom-up inputs are required for
establishment of top-down connectivity onto cortical layer 1 neurogliaform cells.
Neuron 109:3473-3485.e5.

Jercog P, Rogerson T, Schnitzer MJ (2016) Large-Scale Fluorescence Calcium-Imaging
Methods for Studies of Long-Term Memory in Behaving Mammals. Cold Spring Harb
Perspect Biol 8:a021824.

Jones LM, Fontanini A, Sadacca BF, Miller P, Katz DB (2007) Natural stimuli evoke dynamic
sequences of states in sensory cortical ensembles. Proceedings of the National
Academy of Sciences 104:18772–18777.

Klimesch W (2012) Alpha-band oscillations, attention, and controlled access to stored
information. Trends in Cognitive Sciences 16:606–617.

Klimesch W, Hanslmayr S, Sauseng P, Gruber WR, Doppelmayr M (2007) P1 and traveling
alpha waves: evidence for evoked oscillations. J Neurophysiol 97:1311–1318.

Klimesch W, Schack B, Schabus M, Doppelmayr M, Gruber W, Sauseng P (2004) Phase-locked alpha and theta oscillations generate the P1-N1 complex and are related to memory performance. Cognitive Brain Research 19:302–316.

Knudsen EI (2004) Sensitive periods in the development of the brain and behavior. Journal of Cognitive Neuroscience 16:1412–1425.

Le Grand R, Mondloch CJ, Maurer D, Brent HP (2003) Expert face processing requires visual input to the right hemisphere during infancy. Nat Neurosci 6:1108–1112.

Lüdecke D (2018) Sjstats: statistical functions for regression models. Available at: https://zenodo.org/record/1489175 [Accessed October 17, 2021].

Makeig S, Debener S, Onton J, Delorme A (2004) Mining event-related brain dynamics. Trends in Cognitive Sciences 8:204–210.

Makowski D, Ben-Shachar MS, Chen SHA, Lüdecke D (2019) Indices of effect existence and significance in the Bayesian framework. Frontiers in Psychology 10:2767.

Mason SJ, Graham NE (2002) Areas beneath the relative operating characteristics (ROC) and relative operating levels (ROL) curves: Statistical significance and interpretation. Quarterly Journal of the Royal Meteorological Society 128:2145–2166.

Maurer D, Lewis TL (2013) Sensitive Periods in Visual Development. In: The Oxford Handbook of Developmental Psychology, Vol. 1: Body and Mind (Zelazo PD, ed), pp 0. Oxford University Press. Available at: https://doi.org/10.1093/oxfordhb/9780199958450.013.0008 [Accessed August 28, 2023].

McKyton A, Ben-Zion I, Doron R, Zohary E (2015) The limits of shape recognition following late emergence from blindness. Current Biology 25:2373–2378.

Miller CE, Shapiro KL, Luck SJ (2015) Electrophysiological measurement of the effect of inter-stimulus competition on early cortical stages of human vision. NeuroImage 105:229–237.

Mitchell DE, Maurer D (2022) Critical Periods in Vision Revisited. Annual Review of Vision Science 8:291–321.

Oby ER, Degenhart AD, Grigsby EM, Motiwala A, McClain NT, Marino PJ, Yu BM, Batista AP (2025) Dynamical constraints on neural population activity. Nat Neurosci 28:383–393.

Oostenveld R, Fries P, Maris E, Schoffelen J-M (2011) FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Computational intelligence and neuroscience 2011:156869.

Ossandón JP, Stange L, Gudi-Mindermann H, Rimmele JM, Sourav S, Bottari D, Kekunnaya R, Röder B (2023) The development of oscillatory and aperiodic resting state activity is linked to a sensitive period in humans. Neuroimage 275:120171.

Pascual-Marqui RD (2002) Standardized low-resolution brain electromagnetic tomography (sLORETA): technical details. Methods Find Exp Clin Pharmacol 24 Suppl D:5–12.

Pedersini CA, Miller NP, Gandhi TK, Gilad-Gutnick S, Mahajan V, Sinha P, Rokers B (2023) White matter plasticity following cataract surgery in congenitally blind patients. Proceedings of the National Academy of Sciences 120:e2207025120.

Peirce JW (2009) Generating stimuli for neuroscience using PsychoPy. Frontiers in
Neuroinformatics 2:1–8.

Piastra MC, Cruijsen J van der, Piai V, Jeukens FEM, Manoochehri M, Schouten AC, Selles
RW, Oostendorp T (2021) ASH: an Automatic pipeline to generate realistic and
individualized chronic Stroke volume conduction Head models. J Neural Eng
18:044001.

Pitchaimuthu K, Dormal G, Sourav S, Shareef I, Rajendran SS, Ossandón JP, Kekunnaya R,
Röder B (2021) Steady state evoked potentials indicate changes in nonlinear neural
mechanisms of vision in sight recovery individuals. Cortex 144:15–28.

Power M, Fell G, Wright M (2013) Principles for high-quality, high-value testing. BMJ
Evidence-Based Medicine 18:5–10.

Putzar L, Hötting K, Rösler F, Röder B (2007) The development of visual feature binding
processes after visual deprivation in early infancy. Vision Research.

R Core Team (2016) R: A language and environment for statistical computing. Available at:
https://www.r-project.org/.

Rabinovich M, Huerta R, Laurent G (2008) Transient Dynamics for Neural Processing.
Science 321:48–50.

Röder B, Kekunnaya R (2021) Visual experience dependent plasticity in humans. Current
Opinion in Neurobiology 67:155–162.

Röder B, Kekunnaya R (2022) Effects of Early Visual Deprivation. In: Oxford Research
Encyclopedia of Psychology Available at:
https://oxfordre.com/psychology/display/10.1093/acrefore/9780190236557.001.0001/
acrefore-9780190236557-e-839 [Accessed May 3, 2025].

Röder B, Kekunnaya R, Guerreiro MJS (2021) Neural mechanisms of visual sensitive
periods in humans. Neuroscience and biobehavioral reviews 120:86–99.

Röder B, Ley P, Shenoy BH, Kekunnaya R, Bottari D (2013) Sensitive periods for the
functional specialization of the neural system for human face processing.
Proceedings of the National Academy of Sciences 110:16760–16765.

Rösler F, Röder B, Heil M, Hennighausen E (1993) Topographic differences of slow event-
related brain potentials in blind and sighted adult human subjects during haptic
mental rotation. Cognitive Brain Research 1:145–159.

Roy S (1946) খিচুড়ি (Hodgepodge). In: আবোল তাবোল (Balderdash), 2nd ed. Kolkata: Signet Press.
Available at: https://archive.org/details/dli.scoerat.2010aboltabol/page/n7 [Accessed
January 8, 2025].

Schad DJ, Vasishth S, Hohenstein S, Kliegl R (2020) How to capitalize on a priori contrasts
in linear (mixed) models: A tutorial. Journal of Memory and Language 110:104038.

Schrader S, Westhoff A, Piastra MC, Miinalainen T, Pursiainen S, Vorwerk J, Brinck H,
Wolters CH, Engwer C (2021) DUNEuro—A software toolbox for forward modeling in
bioelectromagnetism. PLOS ONE 16:e0252431.

Sourav S, Bottari D, Kekunnaya R, Röder B (2018) Evidence of a retinotopic organization of
early visual cortex but impaired extrastriate processing in sight recovery individuals.
Journal of Vision 18:22.

Sourav S, Bottari D, Shareef I, Kekunnaya R, Röder B (2020) An electrophysiological
biomarker for the classification of cataract-reversal patients: A case-control study.
EClinicalMedicine 27:100559.

Sourav S, Kekunnaya R, Bottari D, Shareef I, Pitchaimuthu K, Röder B (2024) Sound
suppresses earliest visual cortical processing after sight recovery in congenitally blind
humans. Commun Biol 7:1–14.

Tadel F, Baillet S, Mosher JC, Pantazis D, Leahy RM (2011) Brainstorm: a user-friendly
application for MEG/EEG analysis. Comput Intell Neurosci 2011:879716.

Tallon-Baudry C, Bertrand O, Tallon-Baudry C, Bertrand O, Tallon-Baudry C, Bertrand O
(1999) Oscillatory gamma activity in humans and its role in object representation.
Trends in Cognitive Sciences 3:151–162.

Trägenap S, Whitney DE, Fitzpatrick D, Kaschube M (2025) The developmental emergence
of reliable cortical representations. Nat Neurosci 28:394–405.

Tytla ME, Lewis TL, Maurer D, Brent HP (1993) Stereopsis after congenital cataract.
Investigative Ophthalmology and Visual Science 34:1767–1773.

Wanet-Defalque M-C, Veraart C, De Volder A, Metz R, Michel C, Dooms G, Goffinet A (1988)
High metabolic activity in the visual cortex of early blind human subjects. Brain
Research 446:369–373.

Weber A (2018) The USC-SIPI image database: Version 5. Available at: http://sipi. usc.
edu/database/.

Weinhaus F (2011) Fourier Transforms -- IM v6 Examples. ImageMagick v6 Examples --
Fourier Transforms Available at: https://legacy.imagemagick.org/Usage/fourier/
[Accessed October 1, 2024].

World Medical Association (2013) World Medical Association Declaration of Helsinki: Ethical
Principles for Medical Research Involving Human Subjects. JAMA 310:2191–2194.

Yusuf PA, Lamuri A, Hubka P, Tillein J, Vinck M, Kral A (2022) Deficient Recurrent Cortical
Processing in Congenital Deafness. Front Syst Neurosci 16:806142.

1034