Environmental and temporal factors affecting record white-tailed deer antler characteristics in

Ontario, Canada

Brooklyn S. Cars, Environmental and Life Sciences Graduate Program, Trent University, 2140 East

Bank Drive, Peterborough, ON K9J 7B8, Canada

Joseph M. Northrup, Environmental and Life Sciences Graduate Program, Trent University, 2140 East

Bank Drive, Peterborough, ON K9J 7B8, Canada and Wildlife Research and Monitoring Section,

Ontario Ministry of Natural Resources, Peterborough, Ontario, Canada

Keith Beasley' The Foundation for the Recognition of Ontario Wildlife, Ontario, Canada

Paul Beasley, The Foundation for the Recognition of Ontario Wildlife, Ontario, Canada

Kevin Beasley, The Foundation for the Recognition of Ontario Wildlife, Ontario, Canada

Aaron B.A Shafer, Environmental and Life Sciences Graduate Program, Trent University, 2140 East

Bank Drive, Peterborough, ON K9J 7B8, Canada

Correspondence: Brooklyn S. Cars, Trent University, 2140 East Bank Drive, Peterborough, ON K9J

7B8, Canada. Email: brooklyncars@trentu.ca

ABSTRACT

White-tailed deer (*Odocoileus virginianus*) are an ecologically and economically important

species in North America. Their antlers, one of their most recognizable features, are used for dominance

displays, mate attraction, and defense, with size and morphology being key determinants of success.

Antler characteristics are influenced by a combination of genetic and environmental factors, including

habitat quality and resource availability. In this study, we explored how diverse environmental factors,

including climate and land cover composition, impact antler size, morphology, and the distribution of

record-scoring white-tailed deer across Ontario, Canada, using hunter-submitted data from long-term

antler scoring records. We used conditional autoregressive (CAR) models to examine these

relationships and found that warmer temperatures the year of harvest were positively associated with

larger antlers and more record deer in a given county, while winter precipitation the year of harvest was

negatively associated with these characteristics, likely due to reduced forage availability or increased

energy expenditure during more severe winters. Rangeland and forest land cover types were positively

associated with increased antler size and tine number. We observed no temporal changes in antler size

in Ontario, contrasting with broader trends observed in North America. These results show how local

environmental conditions and land cover composition influence antler traits and the distribution of

record white-tailed deer, highlighting the complexity of environmental influences on trait variation.

KEYWORDS white-tailed deer, *Odocoileus virginianus,* antler size, antler morphology, environmental

factors, Ontario, hunters, harvest, climate effects, land cover

The Cervidae (deer) family emerged during the late Miocene where they developed the

secondary sex characteristics that we associate with modern deer (Gilbert et al. 2006). One such

secondary sex characteristic, antlers, are bony extensions of the skull that generally shed every year, and

apart from caribou (*Rangifer tarandus*) are typically only found on males (Rue, 1989; Hewitt, 2011).

Antlers have multiple purposes including dominance displays and interactions, weapons against

predators and other male Cervidae, and indicators to females that males are of good genetic quality

(Clutton-Brock 1982, Goss 1983). In white-tailed deer (*Odocoileus virginianus*), males are the

dominant sex where social dominance is largely dictated by age and body mass, both of which are

closely tied to antler size (Townsend and Bailey 1981). Fully mature males tend to have larger antlers

due to the lower energy demands for body growth and thus their ability to allocate resources to antlers

(Goss, 1983). Likewise, in red deer, larger antlers are associated with increased reproductive success

(Kruuk et al. 2002). That being said, younger or smaller-antlered males can achieve mating success if

they are more dominant and in good overall health, particularly in populations with a female-skewed

sex ratio or a younger male age structure, or by avoiding direct competition with larger males and

engaging in opportunistic mating (Townsend and Bailey 1981, Turner et al. 2016, Partridge 2017).

Antler characteristics are influenced by both genetic and environmental factors. Jamieson et al.

(2020) found that in white-tailed deer >1.5 years old, the number of antler tines were approximately

33% heritable, while Michel et al. (2016) found that in captive white-tailed deer 2.5–5.5 years old antler

score and main beam length were 56% and 45% heritable, respectively. Accordingly, Anderson et al.

(2022) found a suite of genes, with some related to animal health, underlying large antlers. Cain et al.

(2019) showed that larger-antlered deer in the Midwest United States were associated with areas that

had an intermixed landscape with more fragmented patches of land cover, suggesting that habitat

composition and variety can shape antler growth through changes in resource availability. In the

Southern United States, Strickland and Demarais (2008) observed a positive association between mean

antler size and landcover types that favour early-stage herbaceous vegetation such as agriculture.

Intergenerational effects have also been seen to influence antler and other physical traits (Freeman et al.

2013), with, for example, maternal nutrition influencing offspring body mass and antler size in white-

tailed deer (Monteith et al. 2009). Similarly, Peterson et al. (2019) found that years with extreme

droughts, had lasting effects on antler development in white-tailed deer and that deer born during these

years had smaller antlers in later years, suggesting that environmental stress affecting maternal

investment and the early-life of deer can have persistent effects on antler growth. Collectively, these

studies highlight the need to consider multi-year and spatial environment covariates when modelling

trait variation.

In addition to understanding the genetic and environmental factors affecting antler development,

white-tailed deer management approaches can have an effect on antler size and distribution. Across

North America, hunter harvest is the primary means of managing white-tailed deer and management

strategies vary widely, ranging from highly regulated systems with antler point restrictions and

controlled harvests (Strickland and Demarais 2007) to more liberal frameworks with longer seasons and

fewer restrictions on age or sex class (Florida Fish And Wildlife Conservation Commission 2025). In

Ontario (Canada) hunters can purchase a white-tailed deer licence and an antlered deer (i.e. Buck) tag

valid for any open deer season, while antlerless deer (female/fawn buck) tags are awarded via random

draw for a specific Wildlife Management Unit (WMU) (Ontario Ministry of Natural Resources and

Forestry 2019). This system allows wildlife managers to regulate the population through antlerless tag

quotas as females are critical for population growth. Hunters that target male white-tailed deer are often

seeking trophy bucks with higher scoring antlers, which are evaluated based on characteristics such as

tine length and number, beam length and circumference, and antler spread (D'Angelo and Grund 2015,

Boone and Crockett Club 2025). As a result of high-scoring bucks often experiencing heavier harvests,

there has been a broad negative trend in the mean antler size of trophy white-tailed deer across North

America over the last century (Monteith et al. 2013). This decline is not only significant for hunters who

might experience reduced opportunities for harvesting large, trophy bucks, but the removal of larger

males could also disrupt mating structure, potentially resulting in smaller less dominant males having

more success (Langvatn and Loison 1999, Pigeon et al. 2016). This shift could in theory reduce antler

size, which would weaken sexual selection and could, in extreme cases, even lead to reduced

reproductive success and fitness. Evidence of this comes from Malo et al. (2005) who found that in red

deer, antler size and complexity were positively associated with sperm quality and quantity. Similarly,

Ditchkoff et al. (2001) found that in white-tailed deer, antler development was linked to elevated

testosterone during the breeding season, and pathogen resistance.

This study aims to determine which environmental factors influence antler size and morphology

in white-tailed deer across Ontario, Canada. Long-term antler scoring records provided us with a unique

opportunity to examine white-tailed deer antler characteristics and temporal patterns over time and

space in the province. We examined the influence of environmental factors on the spatial distribution,

tine number, antler symmetry, and overall antler score of record white-tailed deer across Ontario. We

hypothesized that antler size in white-tailed deer is positively associated with environmental factors that

promote forage availability and minimize energy expenditure. Specifically, we predicted larger antlers

in counties with higher proportions of agricultural, rangeland, and forest cover, which provide

accessible food and shelter. We also hypothesized that warmer temperatures during the growing season

would be associated with increased antler size, while higher winter precipitation would have a negative

effect due to increased energetic demands. Finally, we hypothesized that selective harvest of large-

antlered individuals over time has led to a temporal decline in antler size across Ontario.

STUDY AREA

White-tailed deer in Ontario, Canada, occupy a diverse landscape with varying anthropogenic

influences. Historically, white-tailed deer populations have been concentrated in southern Ontario,

where winters are milder and growing seasons are longer compared to more northern parts of the

province (Kennedy-Slaney et al. 2018). This area is dominated by agricultural and deciduous forests,

which provide high-quality forage such as grasses, forbs, and crops during spring and summer

(Ecological Stratification Working Group 1995, Government of Ontario 2021). However, white-tailed

deer exist throughout parts of more central and northern Ontario, including the boreal forest region, and

these deer experience harsher winters. To cope with these conditions, deer in these areas inhabit

coniferous and mixed woods forests, which offer shelter from excessive snowfall and winter browse

(Hewitt 2011). While browse is lower in nutritional quality, it is abundant during Ontario's colder

seasons, making up 74-91% of the white-tailed deer's diet when other vegetation is scarce (Mautz et al.

1976, Hewitt 2011). Where they overlap, wolves (*Canis lupus*) are the primary predators of white-tailed

deer in Ontario, while coyotes (*Canis latrans*) fill that role in their absence, though do not fully replace

wolves ecologically (Benson et al. 2017). In addition to climate variation, Ontario's expanding

infrastructure and growing population, which has tripled in the last 50 years and has led to an increase

in white-tailed deer density and a decline in the availability of their habitat and food per individual

(Armstrong et al. 1983, Government of Canada 2016, 2022). However, the proliferation of agricultural

landscapes might increase food availability in some regions further north (Hewitt 2011).

METHODS

Data collection

We obtained Ontario antler data from The Foundation for the Recognition of Ontario Wildlife

(FROW) for the years 1941 to 2022. FROW is a non-profit organization that manages the records of big

game animals in Ontario. Antler measurements for white-tailed deer were voluntarily submitted by

hunters to FROW from every county in Ontario, except for Temiskaming. The antler dataset included

date of kill (DOK), antler type (typical or atypical), tine numbers and multiple scores following

FROW's scoring system, which assigns scores based on additive measurements of the antler including

tine length, beam circumference, and spread. FROW has a minimum requirement that must be met in

order for a white-tailed deer to be entered into the record book (Foundation for the Recognition of

Ontario Wildlife 2015), so we refer to these deer as "record deer" hereafter. Because we were interested

in assessing the influence of spatial and temporal variables on records, we removed entries that did not

contain a county or DOK. Lack of landcover data dating back pre--1999 resulted in all record deer

entries prior to 1997 being removed in the spatial models. We then assessed the influence of a series of

environmental and temporal variables on tine number, gross score, net score, symmetry and the number

of record deer per county.

Predictor variables in our model included temperature (^{o}C), precipitation (mm), and number of

growing degree days for each county corresponding to the year a record deer was harvested (Table S1,

available in Supporting Information). These variables were obtained from Natural Resources Canada

(Hutchinson et al., 2009; Hopkinson et al., 2011; McKenney et al., 2011). We extracted temperature and

precipitation data from the months March to August as these are the months when antler growth occurs

(Hewitt 2011). For precipitation, we also extracted the months January to March to gauge winter

severity through snow fall. Precipitation data were averaged. For temperature, we followed Siegert *et al.*

(2013) and calculated the yearly temperature difference by subtracting each county's average

temperature over the time series from the temperature for each individual year. The temperature and

precipitation data for one year prior to the DOK and two years prior to the DOK were also extracted.

ArcGIS pro (version 3.1.3; Johnston and Environmental Systems Research Institute, 2004), was

used to extract landcover data. We used two land cover layers: the Ontario Land Cover Compilation

v.2.0 (Land Information Ontario, 2023a) which captured the years 1999 to 2011 and Ontario Land

Cover Version 1.0 (Ontario Land Cover Version 1.0 - Overview n.d.) which captured the years 2015 to

2020. Both land cover layers are raster datasets with a 15-metre spatial resolution and a minimum

mapping unit of 0.5 hectares. Depending on their DOK, the record deer were assigned to a landcover

dataset (Table S1). For the years not captured by the landcover databases (2012-2014), whichever layer

encompassed the year closest to the DOK was used. Reclassification of the land cover categories within

these databases was completed to ensure consistency between land cover types: reclassified categories

were; agriculture, infrastructure, deciduous forest, coniferous forest, mixed forest, other forest,

rangeland, water, wetlands, and other landcover types (Fig. 1, Table S2). The proportion of each

landcover type per county was calculated (Table S3), and "other landcover" was dropped in order to

interpret the relationship between the response variable and the proportion covariates (see Valle *et al.*,

2024).

The final three predictor variables used were average county elevation (meters; TessaDEM v1.2

2024), county size (km^2) and deer abundance (Table S1). Average county elevation and county size are

fixed variables, with the values used per county staying constant over the study period. A relative index

of white-tailed deer abundance per county was generated using data from provincial hunter surveys

(Government of Ontario 2025). Each year, the Ontario Ministry of Natural Resources collects

information through hunter surveys on hunting activities. As part of this survey, hunters are asked to

report the number of deer they saw during their hunt and the total number of days they hunted,

providing a metric of the deer seen per day. These values were averaged across all available years

(2008-2023) in each county to provide a single estimate per county.

Statistical analysis

The following predictor variables were standardized (centered to a mean of zero and scaled to a

standard deviation of one) prior to analysis: average county elevation, precipitation (winter and antler

growing season), estimated deer abundance, and growing degree days. Temperature predictors were not

standardized because they were already transformed to reflect deviations from long-term county-level

averages. We assessed pairwise correlations among the variables, removing those with a correlation

greater than 0.70 or less than -0.70 (i.e. Dormann et al. 2013). Landcover variables were excluded from

the correlation assessment among themselves since they are proportion data, therefore, changes in one

necessarily correspond to changes in others, making high correlations expected and not indicative of

problematic collinearity (Valle et al. 2024). However, we did assess correlations between landcover

variables and other covariates. We then fit a series of models to each response variable. All spatial

models were fitted using the S.CARmultilevel function from the CARBayes package in R (v.6.1.1Lee,

2013, R Core Team 2023). These models capture spatial autocorrelation by incorporating spatial

random effects modeled with a Leroux conditional autoregressive (CAR) prior, where neighboring

counties are defined by shared borders. Model fitting occurred within a Bayesian framework using

Markov chain Monte Carlo (MCMC) simulation with default priors applied to all model parameters. All

model specifications, including response variables and full lists of predictors, can be found in Table S4,

available in supporting information.

To assess the number of record deer harvested in each county we used a Poisson regression with

a log link and a county area offset to translate the analysis into one of rates of record deer per area. We

summed the total number of record deer harvested per county across all years, so DOK was not

considered when assigning temperature and precipitation in this model. Instead, average values for the

temperature and precipitation of each county over the specified time span were used. The averaging of

these environmental variables allowed us to observe relationships between the environment and the

spatial distribution of record deer harvests across Ontario counties over the entire study period.

To evaluate antler characteristics, we used DOK when assigning temperature, precipitation and

landcover values which will allow us to observe spatial-temporal relationships at the individual level.

Each model used one of three distributions and their standard link functions: Poisson regression model

with a log link to assess the number of antler tines, a binomial regression model with a logistic link

(logit) to assess antler symmetry (symmetrical = 0, asymmetrical = 1) and gaussian regression models

with identity links to assess antler score (Net and Gross). Antler score was also log-transformed to

normalize the distribution and ensure the assumptions of the model were met.

For CAR models, we evaluated parameter effects based on the probability of direction (pd)

which provided the probability each covariate was either positively or negatively related to antler

characteristics (Makowski et al. 2019). Parameters with a probability $\geq$ 97.5% were considered to have

a high probability of affecting the antler trait. Parameters with a probability of 95 – 97.5% were

considered to have a moderate probability of affecting the antler trait, and parameters with a probability

of 90 – 95% were considered to have a weak, but present, probability of affecting the antler trait. All

probabilities below 90% were considered to have no effect (Lohr et al. 2020, Dias et al. 2025). Spatial

autocorrelation (ρ) was interpreted such that values near 0 indicated spatial independence and values

near 1 indicated strong spatial dependence, so, ρ was considered high if > 0.7, moderate if 0.3–0.7, and

low if < 0.3. Spatial variance (τ^2) and residual variance (σ^2) were considered high if values approached

or exceeded 1 and low if around or below 0.01, based on the Inverse-Gamma (1, 0.01) priors, which

assume most variation is near zero unless the data strongly indicates otherwise (Lee 2013).

Finally, standard generalized linear models (GLMs) from the base r package (R Core Team

2023) were used to assess temporal trends in antler score and tine number, with year as the predictor

variable. Harvest data was sparse pre 1980 and so we used only entries from 1980 onward for these

GLMs. Additionally, minimum scoring requirements increased in 2006, so all records from 1980 to

2006 below this adjusted score were excluded (i.e. Atypical score 130 and Typical score 120;

Foundation for the Recognition of Ontario Wildlife, 2015).

No additional offsets or weights were applied to the models. Model assumptions were assessed

using standard analyses, including assessing normality and homoscedasticity of residuals for the

Gaussian models, and overdispersion for Poisson and Binomial models. It should also be noted that this

data is hunter-submitted and therefore is likely to be non-random and influenced by variation in hunting

effort or harvest efficiency, which are not explicitly captured in our models.

RESULTS

The final dataset after cleaning consisted of 2,413 entries between 1997 and 2022, and 3,072

entries between 1980 and 2022. Number of growing days was removed due to its high correlation with

temperature. Temperature was kept as it directly reflects seasonal climate variation and is easier to

interpret. Deer abundance was also removed due to its high correlation with multiple landcover

variables. Residuals from gaussian models were normal, and no overdispersion was detected in

binomial or Poisson models.

Average temperature had a strong probability of having a positive effect on the number of record

deer per county (Fig. 2; $\beta = 1.24$, pd $= 99.9\%$). Elevation showed a weak probability of having a

positive effect (Fig. 2; $\beta = 0.31$, pd $= 94.3\%$). This model had a strong τ^2 (0.98) and moderate ρ (0.46).

For tine number, the temperature 1 year prior to the year of harvest had a strong probability of having a

positive effect (Fig 2; $\beta = 0.02$, pd $= 99.1\%$). The temperature of the year of harvest and the % of

Rangeland per county both had weak probability of having positive effects (Fig 2 & 3; $\beta = 0.01$, pd $=$

91.1% and $\beta = 0.94$, pd $= 92.2\%$, respectively). This model had a low τ^2 (0.003) and moderate ρ (0.63).

For antler symmetry, precipitation during antler growth 2 years prior to the year of harvest had a strong

probability of having a negative effect (Fig 2; β = -0.12, pd = 98.6%), while the temperature of the year

of harvest had a moderate probability of having a positive effect (Fig 2; β = 0.10, pd = 95.6%). This

model had a low τ^2 (0.03) and moderate ρ (0.38).

For Gross Score, we found the temperature the year of harvest had a strong probability of having

a positive effect (Fig 2; β = 0.01, pd = 100%), while precipitation during the winter the year of harvest

had a strong probability of having a negative effect (Fig 2; β = -0.01, pd = 97.8%). Precipitation during

antler growth the year of harvest and precipitation during the winter 1 year prior to the year of harvest

both had moderate probabilities of having positive effects (Fig 2; β = 0.01, pd = 96.8% and β = 0.01,

96.5%, respectively), and temperature 2 years prior to harvest and the % of Deciduous Forest per county

had weak probabilities of positive effects (Fig 2 & 3; β = 0.003, pd = 91.4% and β = 0.22, 93.4%,

respectively). This model had τ^2 and v^2, both of which were relatively low (0.001 and 0.01 respectively),

and a moderate ρ (0.70).

For Net Score, temperature the year of harvest, temperature 2 years prior to the year of harvest

and the % of Rangeland per county had strong probabilities of having a positive effect (Fig 2 & 3; β =

0.01, pd = 100%, β = 0.01, pd = 99.2% and β = 0.58, 98.9%, respectively). The % of Deciduous Forest

per county had a moderate probability of having a positive effect (Fig 3, β = 0.27, pd = 96.0%). The

precipitation during antler growth the year of harvest and the % of Coniferous Forest per county both

had weak probabilities of positive effects (Fig 2 & 3; β = 0.004, pd = 92.8% and β = 0.28, pd = 94.8%,

respectively), while the precipitation during the winter the year of harvest had a probability of having a

weak but negative effect (Fig 2 & 3; β = -0.004, pd = 91.1%). τ^2 and v^2 were relatively low (0.002 and

0.01 respectively), and ρ was moderately high (β =0.74). Finally, there was no temporal trend in size

and tine number over time (Table S5, available in Supporting Information).

DISCUSSION

The recreational hunting of white-tailed deer brings in millions of dollars to the Canadian economy annually (Fish and Wildlife Ontario 2016). To the Indigenous peoples of North America, white-tailed deer are a food source whose antlers and hides provide material for clothing, tools, and shelter, while also playing an important role in spiritual ceremonies (Brown 2011). In addition, deer with large antlers are highly sought after as trophies by many recreational hunters. Despite their economic and cultural importance, the size of trophy antlers appears to be declining across North America (Monteith et al. 2013). This decline showcases the need to better understand the factors influencing antler growth on a regional level, with some studies suggesting that landscapes featuring a mixture of forest, grassland, and agriculture that create an abundance of edge habitats could increase the chances of a record deer being harvested (Cain et al. 2019). Accordingly, we investigated spatial drivers and temporal patterns of harvested record deer in Ontario, Canada. We observed that increases in temperature during the months that antler growth is occurring (March-August) positively influenced all antler characteristics (Fig 2), and that rangeland and forest are key landscape components, the amounts of which are correlated to overall greater antler size.

Interestingly, no change in antler size was detected over time in Ontario (Fig. 4), potentially due to the province's position at the northern edge of the white-tailed deer's range. Historically, harsher winters and limited forage could have constrained antler growth, but ongoing warming due to climate change is likely improving habitat suitability by reducing winter severity and extending growing seasons, which could be helping maintain antler size in Ontario. This data set is representative of adult antlered white-tailed deer within Ontario and although not a random sample, the data captures a large range of variation in antler size across the population (Fig S1, available in Supporting Information) and temporal, spatial and environmental influences should be reflected in this large sample of male white-

tailed deer (see also Strickland and Demarais, 2008; Monteith *et al.*, 2013; Cain *et al.*, 2019). It should

also be noted that, although some effect sizes presented are small, they nonetheless represent

measurable changes that will be considered in the interpretation of this study.

To account for spatial dependence in our data and better understand regional patterns, we used

CAR models, which revealed moderate to high spatial autocorrelation (ρ) in all models used suggesting

that nearby counties tend to have similar conditions (e.g., temperature). However, a moderate value

indicates that some unaccounted-for spatial covariates are still present but not explicitly modeled.

Possible missing covariates worth further exploration include land productivity, which affects forage

availability, landcover structure that shapes shelter and feeding opportunities, and the distribution of

minerals like phosphorus and calcium, which are involved in bone and antler development (Grasman

and Hellgren 1993, Cain et al. 2019), though the appropriate GIS layers currently do not exist for

Ontario. We also observed low τ^2 in models assessing antler characteristics indicating that, although the

predictors often have similar conditions in neighboring counties, antler size does not exhibit strong

spatial structure. However, a strong τ^2 when assessing the count of record deer per county does suggest

that counties with higher numbers of record deer are more likely to have neighboring counties with

similarly high counts, likely reflective of variation in regional hunting interest and behaviour. Notably,

hunter behaviour and success which will vary with the amount of roads, dwellings, and with

accessibility, while regional differences in the number of hunters and frequency of hunting might also

contribute to these spatial patterns (Kilgo et al. 1998, Bonnot et al. 2013).

Environmental drivers of record antlers

Temperature was the most consistent environmental driver of antler traits: higher temperature

was related to an increase in antler size metrics in all models (Fig 2). During the year of harvest a 1°C

increase in temperature was associated with a ~1% increase in tine number, a ~10% increase in

symmetry, and a ~1% increase in score. Warmer temperatures have been linked to larger antlers and

greater reproductive success in white-tailed deer and other Cervidae (Clements et al. 2010, Simard et al.

2010). Increased plant growth from warmer conditions means more forage availability for the deer and

therefore more energy is able to be allocated to antler growth (French et al. 1956, Hatfield and Prueger

2015). Here, the unique annual regrowth of antlers makes them strongly affected by yearly temperature

changes (Foley et al. 2012). When looking at tine number and score, the temperature one year prior to

harvest and 2 years prior to harvest respectively, also had positive effects on antler size and morphology

(Fig. 2). Antler growth depends on the accumulation of resources stored in the animal's body, warmer

environments, and in turn higher nutritional gain, in prior years can enhance fat reserves which are then

used in subsequent antler cycles when that energy is needed (Hewitt 2011).

Similar inference can be made from the influence of precipitation during antler growth the year

of harvest on antler size. Wetter conditions can lead to increased plant growth which could again affect

the annual regrowth of antlers (Foley et al. 2012, Scharwies and Dinneny 2019). A 1mm increase in

precipitation was associated with a ~1% increase in net score. Similarly, we saw negative effects from

precipitation during the winter the year of harvest on antler size (Fig 2). More snow in winter can make

it more difficult for the deer to move around and find forage, which causes them to congregate in deer

yards (Government of Ontario 2019). Deer yards are an area made up of coniferous trees that have a

large canopy for blocking excessive snowfall (Government of Ontario 2019). With deer being confined

to these yards deer density and the availability of forage for each individual animal decreases (Patterson

and Power 2002). This could then put individuals with poor body conditions at a disadvantage and

minimal fat reserves when antler regrowth commences (Lesage et al. 2001). Future work could

strengthen this finding by incorporating direct measurements of snow depth, snow cover duration, or

broader indices such as the Winter Severity Index (WSI) to capture more details on the energetic

challenges white-tailed deer face during the winter months.

Rangeland and Forest landcover had positive effects on antler characteristics, meaning

that as the percentages of these types of landcover in each county rose so too did antler score and

tine number (Fig 5). Rangeland and Forest together have been found to increase the probability

of harvesting a record deer and are ideal edge habitats (Cain et al. 2019). Forests will block

excessive snowfall and contain large amounts of northern deer's primary food source, browse

(Hewitt 2011). Further, rangeland allows for greater visibility of large antlered bucks from

further away, potentially increasing hunter success or improving the effectiveness of dominance

displays and mate attraction, while also providing food such as grasses (Gilbert et al. 2006,

Hewitt 2011, Houde et al. 2020). Collectively, these combined edge habitats are preferred by

white-tailed deer as it allows them to have a larger variety of food sources and provides both

more space for movement (rangeland) while still having adequate cover (forest) (Halls 1984,

Alverson et al. 1988).

Number of harvested deer and antler symmetry

We observed that increased average temperatures were associated with more record white-tailed

deer being harvested in a given Ontario county with a 1°C increase in temperature during the year of

harvest being associated with an increase in the number of record-sized deer by 1.24 deer in a given

county. Similar to antler size, warmer temperatures allow for more forage availability. With more food

available, more energy can be allocated to individual deer in a county, presumably allowing more

individuals to reach record deer status. The number of record deer harvested was not significantly

affected by any landcover covariates (Fig 3); so, while landcover type positively impacts antler size, the

overall effect is minor such that it does not increase the net number of large animals harvested in a given

year. Curiously, record deer numbers were positively affected by elevation (Fig. 2). It is tempting to

suggest this being reflective of mature deer finding refuge from human activity via fewer roadways at

these elevations. Less trafficked zones allow deer to move more freely and access critical resources with

minimal human interference (Hamilton et al. 2024) and behaviourally deer learn to avoid humans

(Kilgo et al. 1998, Marantz et al. 2016). However, the elevational range is small in the province (0 m –

700 m) and obfuscated when averaged across the county as done here (66 m – 415 m), and thus finer

scale analysis is warranted to better understand this effect.

The symmetry of antlers appears to be impacted by precipitation, with higher precipitation levels

2 years prior to antler growth resulting in less symmetrical antlers. Under uniform environmental

conditions, white-tailed deer exhibit a genetic predisposition for antlers to grow symmetrically

(Ditchkoff et al. 2001). White-tailed deer can develop antler asymmetry when exposed to stressors such

as environmental conditions, disease, or injury that affect energy allocation before and during antler

growth (Marburger et al. 1972, Goss 1983, Wild et al. 2022). The mechanism by which precipitation 2

years prior impacts asymmetry is still unclear. However, maternal effects on antler size are documented

in cervids (Freeman et al. 2013), so early-in-life conditions can impact adult traits, with the impact to

antlers seemingly long-lasting (Freeman et al. 2013). Asymmetry has also been seen to increase as trait

size increases (De Coster et al. 2013), which we observed in our data set (Fig S2, available in

Supporting Information); however, we see no strong relationship between precipitation and antler size.

It is thus still conceivable that high environmental stressors might have long-term impacts on symmetry,

even for antlers that are shed annually.

Temporal patterns of record deer

Over the last 40 years, there has been no significant change in antler size (or score) and

the tine number of harvested record white-tailed deer in Ontario (Table S5). This contrasts the

findings of Monteith et al. (2013) who reported a negative trend across North America in antler

size in white-tailed deer from 1950 to 2008, based on trophy records of mature males submitted

to the Boone and Crockett Club's *Records of North American Big Game* (Boone and Crockett

Club 2025). These negative trends were hypothesized to be a combination of harvest-induced

reduction in age structure, sociological effects and density dependence (Monteith et al. 2013).

Our finding suggests that antler trends in white-tailed deer are region specific and vary

depending on the environment and pressures present.

In Ontario, the number of active hunters has stayed relatively consistent since 2008 while the

number of records submitted to FROW has declined (Fig 6). This could be indicative of a possible shift

in hunting priorities (Monteith et al. 2013), but multiple tough winters in 2007/08 and 2010/11, have

coincided with apparent population declines (Ontario Outdoors Magazine Game Forecast 2009, 2011)

which would explain a decrease in entries. Still, the number of white-tailed deer in Ontario is relatively

low especially compared to other regions of North America where population numbers are substantially

higher. For example, Michigan with similar environmental conditions to Ontario and an area of ~

250,493 km^2 has a population of ~2 million white-tailed deer, compared to Ontario with an area of over

1 million km^2 and a white-tailed deer population of ~400 thousand (Mason 2016, Conner 2021). This

lower density is likely due to Ontario being the northern edge of the white-tailed deer's range, where

limited suitable habitat likely has more influence on antlers than harvest pressure or hunting regulations.

While Ontario's hunting regulations of being allotted one antlered tag and allocating antlerless tags by

lottery could result in lower overall harvest rates compared to other regions, it is not clear if these

restrictions make any substantial difference in population dynamics or antler development. Instead,

environmental factors associated with climate and habitat likely have a greater influence on population

density and antler growth in the province.

In much of North America, warmer climates lead to longer growing seasons and an abundance

of high-quality food, which allows white-tailed deer to reach their maximum physical size during antler

growth (Mysterud et al. 2005, Sontheimer et al. 2024). In contrast, Ontario's colder winters limit access

to nutritious food, which could be preventing deer from achieving their full antler growth potential; this

limitation could mean there is still room for growth. Our models indicate that warming temperatures in

Ontario are positively influencing antler development, as milder winters and longer growing seasons

improve food availability. Combined these conditions could be helping bolster antler size despite

pressures that are causing declines elsewhere (e.g.Monteith et al. 2013). With ongoing climate change,

we would predict a positive trend in overall antler size if environmental conditions became more

favourable for deer and harvest pressures remain like those during our time series.

MANAGEMENT IMPLICATIONS

Our findings highlight how variation in habitat conditions across Ontario counties are related to

differences in antler size and morphology as well as record white-tailed deer distributions.

Understanding these patterns and relationships can help identify conditions more likely to produce

larger antlered deer, and by communicating these findings, hunters can better manage their expectations

and recognize aspects that could cause certain regions to have higher large-antlered white-tailed deer

concentrations, leading to greater satisfaction (D'Angelo and Grund 2015). An increase in hunter

satisfaction should, in turn, generate revenue that supports the funding of conservation efforts, wildlife

management programs, and local economies. With no significant temporal trend in antler size, the

current harvest practices in Ontario are likely effective as it relates to maintaining antler size, but we

note increased harvest pressure on males in other regions has led to a reduction in age structure, which

could impact antler size in the future(Langvatn and Loison 1999, Mysterud 2011). Density dependence

also can disproportionately affect large males, as they often enter the winter months with low fat

reserves from energy expenditure during the mating season (Lesage et al. 2001, Keyser et al. 2005),

which could have implications for survival and future antler growth. This reinforces the need to

continue to monitor harvest practices to ensure that population dynamics and habitat limitations do not

negatively impact the long-term sustainability of Ontario's white-tailed deer population. It is important

to note that these findings are based on hunter-reported data, which could be subject to sampling biases,

and are most directly applicable within the Ontario context. Nonetheless, they provide valuable insight

into how environmental variation is shaping antler characteristics and white-tailed deer distributions,

contributing to a deeper understanding of population dynamics and habitat influences.

ACKNOWLEDGMENTS

This work was supported by NSERC Discovery Grant (ABAS; 410 SDC); Canadian Foundation

for Innovation (ABAS): John R. Evans Leaders Fund (ABAS) and the Ontario Early Researcher Award

(ABAS). Trent University is located on the traditional territory of the Michi Saagiig Anishnaabeg.

ETHICS STATEMENT

No approval of research ethics committees was required to accomplish the goals of this study.

DATA AVAILABILITY

The antler data that support the findings of this study are available from The Foundation for the

Recognition of Ontario Wildlife, Ontario, Canada and were acquired under research agreements for this

study. All other model covariates are provided.

CONFLICT OF INTEREST

Authors declare no competing interests.

REFERENCES

Alverson, W. S., D. M. Waller, and S. L. Solheim. 1988. Forests Too Deer: Edge Effects in
     Northern Wisconsin. Conservation Biology 2:348–358.

Anderson, S. J., S. D. Côté, J. H. Richard, and A. B. A. Shafer. 2022. Genomic
     architecture of phenotypic extremes in a wild cervid. BMC Genomics 23:NA-NA.

Armstrong, E., D. Euler, and G. Racey. 1983. White-Tailed Deer Habitat and Cottage
     Development in Central Ontario. The Journal of Wildlife Management 47:605–612.

Benson, J. F., K. M. Loveless, L. Y. Rutledge, and B. R. Patterson. 2017. Ungulate predation and
     ecological roles of wolves and coyotes in eastern North America. Ecological
     Applications: A Publication of the Ecological Society of America 27:718–733.

Bonnot, N., N. Morellet, H. Verheyden, B. Cargnelutti, B. Lourtet, F. Klein, and A. J. M.
     Hewison. 2013. Habitat use under predation risk: hunting, roads and human dwellings
     influence the spatial behaviour of roe deer. European Journal of Wildlife Research
     59:185–193.

Boone and Crockett Club. 2025. Scoring Your Trophy: typical whitetail deer. Boone and
     Crockett Club. <https://www.boone-crockett.org/scorechart/add/whitetail_deer>.
     Accessed 3 Mar 2025.

Brown, A. 2011. Traditional relationship of Cherokees and Whitetail Deer. The Cherokee One
     Feather. <https://theonefeather.com/2011/11/28/traditional-relationship-of-cherokees-
     and-whitetail-deer/>. Accessed 8 Jan 2025.

Cain, R. L., N. P. Snow, J. C. Crawford, D. M. Williams, and W. F. Porter. 2019. Spatial
     distribution and landscape associations of large-antlered deer. The Journal of Wildlife
     Management 83:1762–1772.

Clements, M. N., T. H. Clutton-Brock, S. D. Albon, J. M. Pemberton, and L. E. B. Kruuk. 2010.

    Getting the timing right: antler growth phenology and sexual selection in a wild red deer

    population. Oecologia 164:357–369.

Clutton-Brock, T. H. 1982. The Functions of Antlers. Behaviour 79:108–125.

Conner, M. 2021. What is the deer population in Ontario? - ruggedthuglife.com. Canada Online

    Guide. <https://ruggedthuglife.com/ontario/what-is-the-deer-population-in-ontario/>.

    Accessed 12 Jan 2025.

D'Angelo, G. J., and M. D. Grund. 2015. Evaluating Competing Preferences of Hunters and

    Landowners for Management of Deer Populations. https://doi.org/10.26077/N707-1D90.

De Coster, G., S. Van Dongen, P. Malaki, M. Muchane, A. Alcántara-Exposito, H. Matheve, and

    L. Lens. 2013. Fluctuating Asymmetry and Environmental Stress: Understanding the

    Role of Trait History. PLoS ONE 8:e57966.

Dias, T., N. P. Lemoine, S. W. Yanco, M. Zimova, R. A. Bay, and B. C. Weeks. 2025. Long-

    term increases in wing length occur independently of changes in climate and climate-

    driven shifts in body size. Proceedings of the Royal Society B: Biological Sciences

    292:20242556.

Ditchkoff, S. S., R. L. Lochmiller, R. E. Masters, S. R. Hoofer, and R. A. Van Den Bussche.

    2001. Major-histocompatibility-complex-associated variation in secondary sexual traits

    of white-tailed deer (Odocoileus virginianus): evidence for good-genes advertisement.

    Evolution; International Journal of Organic Evolution 55:616–625.

Dormann, C. F., J. Elith, S. Bacher, C. Buchmann, G. Carl, G. Carré, J. R. G. Marquéz, B.

    Gruber, B. Lafourcade, P. J. Leitão, T. Münkemüller, C. McClean, P. E. Osborne, B.

    Reineking, B. Schröder, A. K. Skidmore, D. Zurell, and S. Lautenbach. 2013.

Collinearity: a review of methods to deal with it and a simulation study evaluating their

performance. Ecography 36:27–46.

Ecological Stratification Working Group. 1995. National Ecological Framework for Canada.

Agriculture and Agri-Food Canada: Ottawa, ON.

Fish and Wildlife Ontario. 2016. Fish and Wildlife Special Purpose Account Annual Report

2016-17.

Florida Fish And Wildlife Conservation Commission. 2025. Species Season Dates and Bag

Limits. <https://myfwc.com/hunting/season-dates/>. Accessed 21 Jul 2025.

Foley, A. M., R. W. DeYoung, S. D. Lukefahr, J. S. Lewis, D. G. Hewitt, M. W. Hellickson, D.

A. Draeger, C. A. DeYoung, and H. Beck. 2012. Repeatability of antler characteristics in

mature white-tailed deer in South Texas: consequences of environmental effects. Journal

of Mammalogy 93:1149–1157.

Foundation for the Recognition of Ontario Wildlife. 2015. Big Game Records of Ontario.

Seventh edition. Beasley Brothers Outdoors Inc.

Freeman, E. D., R. T. Larsen, K. Clegg, and B. R. McMillan. 2013. Long-Lasting Effects of

Maternal Condition in Free-Ranging Cervids. PLoS ONE 8:e58373.

French, C. E., L. C. McEwen, N. D. Magruder, R. H. Ingram, and R. W. Swift. 1956. Nutrient

Requirements for Growth and Antler Development in the White-Tailed Deer. The Journal

of Wildlife Management 20:221–232.

Gilbert, C., A. Ropiquet, and A. Hassanin. 2006. Mitochondrial and nuclear phylogenies of

Cervidae (Mammalia, Ruminantia): Systematics, morphology, and biogeography.

Molecular Phylogenetics and Evolution 40:101–117.

Goss, R. J. 1983. Deer antlers: regeneration, function, and evolution. Academic Press, New

York.

Government of Canada. 2016. Table 24.1 Population, by province and territory, selected years,

1861 to 2011. <https://www150.statcan.gc.ca/n1/pub/11-402-

x/2012000/chap/pop/tbl/tbl01-eng.htm>. Accessed 26 Nov 2024.

Government of Canada, S. C. 2022. Profile table, Census Profile, 2021 Census of Population -

Ontario [Province]. <https://www12.statcan.gc.ca/census-recensement/2021/dp-

pd/prof/index.cfm?Lang=E>. Accessed 19 Sep 2023.

Government of Ontario. 2019. 4 Identifying habitats of seasonal concentrations of animals | 1

Significant wildlife habitat technical guide | ontario.ca.

<http://www.ontario.ca/document/significant-wildlife-habitat-technical-guide/4-

identifying-habitats-seasonal>. Accessed 4 Feb 2025.

Government of Ontario. 2021. Forest regions. <http://www.ontario.ca/page/forest-regions>.

Accessed 11 Oct 2023.

Government of Ontario. 2025. Hunter reporting. <http://www.ontario.ca/page/hunter-reporting>.

Accessed 17 Jun 2025.

Grasman, B. T., and E. C. Hellgren. 1993. Phosophorus Nutrition in White-Tailed Deer: Nutrient

Balance, Physiological Responses, and Antler Growth. Ecology 74:2279–2296.

Halls, L. K., editor. 1984. White-Tailed Deer: Ecology and Management. Stackpole Books,

Harrisburg, Pa.

Hamilton, K. M., T. Bommarito, and J. S. Lewis. 2024. Human activities influence mule deer use

of overpasses across multiple scales. The Journal of Wildlife Management 88:e22560.

Hatfield, J. L., and J. H. Prueger. 2015. Temperature extremes: Effect on plant growth and

development. Weather and Climate Extremes 10:4–10.

Hewitt, D. 2011. Biology and management of white-tailed deer. CRC Press, Boca Raton, FL.

Houde, N., J.-P. Tremblay, N. Thiffault, and S. D. Côté. 2020. Manipulating forage and risk

avoidance to increase white-tailed deer vulnerability to hunters. Wildlife Biology 2020.

Jamieson, A., S. J. Anderson, J. Fuller, S. D. Côté, J. M. Northrup, and A. B. A. Shafer. 2020.

Heritability Estimates of Antler and Body Traits in White-Tailed Deer (Odocoileus

virginianus) From Genomic-Relatedness Matrices. Journal of Heredity 111:429–435.

Kennedy-Slaney, L., J. Bowman, A. A. Walpole, and B. A. Pond. 2018. Northward bound: the

distribution of white-tailed deer in Ontario under a changing climate. Wildlife Research

45:220–228.

Keyser, P. D., D. C. Guynn, and H. S. Hill. 2005. Population Density: Physical Condition

Relationships in White-Tailed Deer. The Journal of Wildlife Management 69:356–365.

Kilgo, J., R. Labisky, and D. Fritzen. 1998. Influences of Hunting on the Behavior of

White□Tailed Deer: Implications for Conservation of the Florida Panther. Conservation

Biology 12:1359–1364.

Kruuk, E. B., J. Slate, J. M. Pemberton, S. Brotherstone, F. Guinness, and T. Clutton-Brock.

2002. Antler size in red deer: heritability and selection but no evolution. Evolution;

International Journal of Organic Evolution 56:1683–1695.

Langvatn, R., and A. Loison. 1999. Consequences of harvesting on age structure, sex ratio and

population dynamics of red deer Cervus elaphus in central Norway. Wildlife Biology

5:213–223.

Lee, D. 2013. CARBayes: An R Package for Bayesian Spatial Modeling with Conditional

Autoregressive Priors. Journal of Statistical Software 55:1–24.

Lesage, L., M. Crête, J. Huot, and J.-P. Ouellet. 2001. Evidence for a Trade-off between Growth

and Body Reserves in Northern White-Tailed Deer. Oecologia 126:30–41.

Lohr, A. K., J. A. Martin, G. T. Wann, B. S. Cohen, B. A. Collier, and M. J. Chamberlain. 2020.

Behavioral strategies during incubation influence nest and female survival of Wild

Turkeys. Ecology and Evolution 10:11752–11765.

Makowski, D., M. S. Ben-Shachar, and D. Lüdecke. 2019. bayestestR: Describing Effects and

their Uncertainty, Existence and Significance within the Bayesian Framework. Journal of

Open Source Software 4:1541.

Marantz, S. A., J. A. Long, S. L. Webb, K. L. Gee, A. R. Little, and S. Demarais. 2016. Impacts

of human hunting on spatial behavior of white-tailed deer (Odocoileus virginianus).

Canadian Journal of Zoology 94:853–862.

Marburger, R. G., R. M. Robinson, J. W. Thomas, M. J. Andregg, and K. A. Clark. 1972. Antler

malformation produced by leg injury in white-tailed deer. Journal of Wildlife Diseases

8:311–314.

Mason, R. 2016. Michigan Deer Management Plan.

Mautz, W. W., T. W. Walski, and W. E. Urban. 1976. Digestibility of Fresh Frozen versus

Pelleted Browse by Snowshoe Hares. The Journal of Wildlife Management 40:496–499.

Michel, E. S., S. Demarais, B. K. Strickland, T. Smith, and C. M. Dacus. 2016. Antler

characteristics are highly heritable but influenced by maternal factors. 80:1420–1426.

Monteith, K. L., L. E. Schmitz, J. A. Jenks, J. A. Delger, and R. T. Bowyer. 2009. Growth of
Male White-Tailed Deer: Consequences of Maternal Effects. Journal of Mammalogy
90:651–660.

Monteith, K., R. Long, V. Bleich, J. Heffelfinger, P. Krausman, and R. Bowyer. 2013. Effects of
harvest, culture, and climate on trends in size of horn-like structures in trophy ungulates.
Wildlife Monographs 183:1–26.

Mysterud, A. 2011. Selective harvesting of large mammals: how often does it result in
directional selection? Journal of Applied Ecology 48:827–834.

Mysterud, A., E. Meisingset, R. Langvatn, N. G. Yoccoz, and N. Chr. Stenseth. 2005. Climate-
dependent allocation of resources to secondary sexual traits in red deer. Oikos 111:245–
252.

Ontario Land Cover Version 1.0 - Overview. n.d.
<https://www.arcgis.com/home/item.html?id=667367a759214a089917adccdbae7cb2>.
Accessed 1 May 2024.

Ontario Ministry of Natural Resources and Forestry. 2019. White-tailed deer population
objective setting and harvest management guidelines. <http://www.ontario.ca/page/white-
tailed-deer-population-objective-setting-and-harvest-management-guidelines>. Accessed
30 Dec 2024.

Ontario Outdoors Magazine Game Forecast. 2009. Hunting Forecast. 2009.

Ontario Outdoors Magazine Game Forecast. 2011. Hunting Forecast. 2011.

Partridge, C. 2017. Sneak Copulation as an Alternative Mating Strategy. Pages 1–3 *in* T. K.
Shackelford and V. A. Weekes-Shackelford, editors. Encyclopedia of Evolutionary
Psychological Science. Springer International Publishing, Cham.

Patterson, B. R., and V. A. Power. 2002. Contributions of Forage Competition, Harvest, and

Climate Fluctuation to Changes in Population Growth of Northern White-Tailed Deer.

Oecologia 130:62–71.

Peterson, B. C., C. W. Schoenebeck, and N. J. Fryda. 2019. Effects of Extreme Environmental

Conditions on White-tailed Deer Antlers. Wildlife Society Bulletin (2011-) 43:708–716.

Pigeon, G., M. Festa-Bianchet, D. W. Coltman, and F. Pelletier. 2016. Intense selective hunting

leads to artificial evolution in horn size. Evolutionary Applications 9:521–530.

R Core Team. 2023. R: The R Project for Statistical Computing. <https://www.r-project.org/>.

Accessed 19 Jan 2023.

Scharwies, J. D., and J. R. Dinneny. 2019. Water transport, perception, and response in plants.

Journal of Plant Research 132:311–324.

Siegert, S., J. Broecker, and H. Kantz. 2013. Skill of data based predictions versus dynamical

models -- case study on extreme temperature anomalies.

Simard, M. A., T. Coulson, A. Gingras, and S. D. Côté. 2010. Influence of Density and Climate

on Population Dynamics of a Large Herbivore Under Harsh Environmental Conditions.

The Journal of Wildlife Management 74:1671–1685.

Sontheimer, W. P., D. Pfeffer, and F. W. Weckerly. 2024. Ontogenetic and static scaling of

antler mass in White-tailed Deer (Odocoileus virginianus). Journal of Mammalogy

105:134–142.

Strickland, B. K., and S. Demarais. 2007. Using Antler Restrictions to Manage for Older-aged

Bucks.

Strickland, B. K., and S. Demarais. 2008. Influence of Landscape Composition and Structure on

Antler Size of White-Tailed Deer. The Journal of Wildlife Management 72:1101–1108.

TessaDEM v1.2. 2024. Near-global 30-meter Digital Elevation Model combining AW3D30,

MERIT DEM, Forest Height, World Settlement Footprint, and Global Surface Water

datasets. Available under Open Database License (ODbL) v1.0. <https://tessadem.com/>.

Accessed 16 Jun 2025.

Townsend, T. W., and E. D. Bailey. 1981. Effects of Age, Sex and Weight on Social Rank in

Penned White-tailed Deer. The American Midland Naturalist 106:92–101.

Turner, M. M., C. S. Deperno, W. Booth, E. L. Vargo, M. C. Conner, and R. A. Lancia. 2016.

The mating system of white-tailed deer under Quality Deer Management: Mating System

of White-Tailed Deer Under QDM. The Journal of Wildlife Management 80:935–940.

Valle, D., J. Mintz, and I. V. Brack. 2024. Estimation and interpretation problems and solutions

when using proportion covariates in linear regression models. Ecology 105:e4256.

Wild, M. A., G. A. Sargeant, K. Garrison, and D. Conradson. 2022. Association of antler

asymmetry with hoof disease in elk. The Journal of Wildlife Management 86:e22245.

FIGURES

Figure 1. Approximate Ontario land cover distribution (Left) and elevation (Right) divided by county.

Figure 2. Probability of direction (pd) of elevation, temperature, and precipitation on the response variables. Variables in pink have a high probability affecting the antler trait (97.5%). Variables in Orange have a moderate probability (95 – 97.5%). And Variables in green have a weak, but present probability (90 – 95%).

Figure 3. Probability of direction (pd) of landcover proportion on the response variables. Variables in pink have a high probability affecting the antler trait (97.5%). Variables in Orange have a moderate probability (95 – 97.5%). And Variables in green have a weak, but present probability (90 – 95%).

Figure 4. Change in antler score and tine number over time. Cutoffs are representative of Boone and Crockett data, depicting the higher cutoff size for record antlers in other parts of North America.

Figure 5. Percent of significant landcover per predicted antler score and tine number

Figure 6. Number of records submitted to the Foundation for the Recognition of Ontario Wildlife (FROW) from 1997 to 2020 (blue) and Number of active hunters from 2008 to 2022 (pink)

APPENDIX A. SUPPLEMENTAL