Environmental and temporal factors affecting record white-tailed deer antler characteristics in

Ontario, Canada

Brooklyn S. Cars, Environmental and Life Sciences Graduate Program, Trent University, 2140 East

Bank Drive, Peterborough, ON K9J 7B8, Canada

Joseph M. Northrup, Environmental and Life Sciences Graduate Program, Trent University, 2140 East

Bank Drive, Peterborough, ON K9J 7B8, Canada and Wildlife Research and Monitoring Section,

Ontario Ministry of Natural Resources, Peterborough, Ontario, Canada

Keith Beasley, The Foundation for the Recognition of Ontario Wildlife, Ontario, Canada

Paul Beasley, The Foundation for the Recognition of Ontario Wildlife, Ontario, Canada

Kevin Beasley, The Foundation for the Recognition of Ontario Wildlife, Ontario, Canada

Aaron B.A Shafer, Environmental and Life Sciences Graduate Program, Trent University, 2140 East

Bank Drive, Peterborough, ON K9J 7B8, Canada

Correspondence: Brooklyn S. Cars, Trent University, 2140 East Bank Drive, Peterborough, ON K9J

7B8, Canada. Email: brooklyncars@trentu.ca

ABSTRACT

White-tailed deer (*Odocoileus virginianus*) are an ecologically and economically important species in North America. Their antlers, one of their most recognizable features, are used for dominance displays, mate attraction, and defense, with size and shape being key determinants of success. Antler characteristics are influenced by a combination of genetics, age and environmental factors, notably habitat quality and resource availability. In this study, we explored how diverse environmental factors, including climate and land cover composition, impact antler size, tine configuration, and the distribution of record-scoring white-tailed deer across Ontario, Canada, using hunter-submitted data from long-term antler scoring records. We used conditional autoregressive (CAR) models to examine these relationships and found that warmer temperatures the year of harvest were positively associated with larger antlers and more record deer in a given county, while winter precipitation the year of harvest was negatively associated with these characteristics, likely due to reduced forage availability or increased energy expenditure during more severe winters. Rangeland and forest land cover types were positively associated with increased antler size and tine number. We observed no temporal changes in antler size in Ontario, contrasting with broader trends observed in North America. These results show how local environmental conditions and land cover composition influence antler traits and the distribution of record white-tailed deer, highlighting the complexity of environmental influences on trait variation.

KEYWORDS white-tailed deer, *Odocoileus virginianus,* antler size, tine configuration, environmental factors, Ontario, hunters, harvest, climate effects, land cover

The Cervidae (deer) family emerged during the late Miocene where they developed the

secondary sex characteristics that we associate with modern deer (Gilbert et al. 2006). One such

secondary sex characteristic, antlers, are bony extensions of the skull that generally shed every year, and

apart from caribou (*Rangifer tarandus*) are typically only found on males (Rue 1989, Hewitt 2011) .

Antlers have multiple purposes including dominance displays and interactions, weapons against

predators and other male Cervidae, and indicators used in mate selection (Clutton-Brock 1982, Goss

1983, Morina et al. 2018). In white-tailed deer (*Odocoileus virginianus*), males are the dominant sex

where social dominance is largely dictated by age and body mass, both of which are closely tied to

antler size (Townsend and Bailey 1981). Fully mature males tend to have larger antlers due to the lower

energy demands for body growth and thus their ability to allocate more resources (Goss, 1983).

Likewise, in red deer (*Cervus elaphus*), larger antlers are associated with increased reproductive success

(Kruuk et al. 2002). But while larger-antlered males may have an advantage when it comes to breeding

opportunities, mating success is not limited to them (DeYoung et al. 2006). Younger or smaller-antlered

white-tailed deer can achieve mating success if they are more dominant and in good overall health, in

populations with a female-skewed sex ratio or a younger male age structure, or by avoiding direct

competition with larger males and engaging in opportunistic mating (Townsend and Bailey 1981,

DeYoung et al. 2006, Turner et al. 2016).

Although antler traits have a heritable component (e.g. Michel *et al.*, 2016; Jamieson *et al.*,

2020), environmental conditions and habitat composition affect their expression, with the quality and

quantity of forage among landcover types playing a large role. Agricultural and rangeland areas provide

nutrient rich, easily accessible forage, including plants such as soybeans, alfalfa, and clover (Harper

2019), while forests offer cover from predators and winter conditions but the quality of forage varies

with tree species and understory composition (Voigt et al. 1997). Consistent with these patterns,

Strickland and Demarais (2008) observed a positive association in the southern United States between

mean antler size and landcover types that favour forage-producing vegetation such as agriculture.

Further, Cain et al. (2019) showed that larger-antlered deer in the Midwest United States were

associated with an intermixed landscape with more fragmented patches of land cover, suggesting that

habitat composition and variety can shape antler growth through changes in resource availability. In

addition to landcover, seasonal environmental conditions such as heavy winter snowfall can reduce

forage availability and increase energetic costs for white-tailed deer populations. For example,

Garroway and Broders (2005) found that deer in areas with deeper snow exhibited lower body

condition, suggesting that winter severity directly affects white-tailed deer energy reserves.

Intergenerational effects can also influence antler and other physical traits (Freeman et al. 2013),

with, for example, maternal nutrition influencing offspring body mass and antler size in white-tailed

deer (Monteith et al. 2009). Strickland et al. (2020) showed that summer droughts during gestation and

early lactation caused maternal nutritional stress that limited resource allocation to developing offspring

and resulted in reduced antler mass in young males. Similarly, Peterson et al. (2019) found that male

white-tailed deer born during years with extreme droughts had smaller antler metrics compared with

those born in non-extreme years. Collectively, these studies highlight the need to consider multi-year

and spatial environmental covariates when modelling trait variation.

Across North America, hunter harvest is the primary means of managing white-tailed deer and

management strategies vary widely, ranging from highly regulated systems with antler□point

restrictions and controlled harvests (Strickland and Demarais 2007) to more liberal frameworks with

longer seasons and fewer restrictions on age or sex class (Florida Fish And Wildlife Conservation

Commission 2025). In Ontario, Canada, the hunting system leans toward a moderately liberal

framework; hunters can purchase a white-tailed deer licence and an antlered deer (i.e. buck) tag valid

for any open deer season, with the option to purchase additional tags depending on the management

unit. Ontario regulations define an antlered deer as one with at least one antler measuring 7.5 cm or

more in length, which limits harvest of younger or smaller-antlered bucks helping preserve future

breeding males. Antlerless deer (female/fawn buck) tags are awarded via random draw for a specific

Wildlife Management Unit (WMU) which allow wildlife managers to regulate population growth by

controlling female harvest (Government of Ontario 2019*a*). Both residents and non-residents can hunt in

Ontario, with harvest levels fluctuating among WMUs based on deer density, land access, and hunter

participation. Approximately 50,000–75,000 deer are harvested annually across the province

(Government of Ontario 2025).

With the allocation of antlered deer tags being more liberal than antlerless tags, hunters have

greater opportunity to harvest male white-tailed deer. Regardless of hunting motivation (e.g. food,

trophy), hunters generally prefer larger bucks, including trophy bucks with larger antlers (D'Angelo and

Grund 2015, Boone and Crockett Club 2025). With larger, higher-scoring bucks often experiencing

heavier harvests, there has been a broadscale negative trend in the mean antler size of trophy white-

tailed deer across North America over the last century, largely attributed to reductions in male age

structure resulting from harvest pressure (Monteith et al. 2013). But while selective harvest can

influence age structure, the effects on population-level antler size and mating structure are complex.

Males of all ages and antler sizes typically breed successfully, and management strategies often allow

younger males to reach prime antler size before harvest (Webb et al. 2012, Turner et al. 2016). Thus, the

potential evolutionary impacts of removing larger males on mating structure and antler size are likely

limited (Webb et al. 2012).

This study aims to determine which environmental factors influence antler size and tine

configuration in white-tailed deer across Ontario, Canada. Long-term antler scoring records provided us

with a unique opportunity to examine white-tailed deer antler characteristics and temporal patterns over

time and space in the province. We examined the influence of environmental factors on the spatial

distribution, tine number, antler symmetry, and overall antler score of record white-tailed deer across

Ontario. We hypothesized that antler size in white-tailed deer is positively associated with

environmental factors that promote forage availability and minimize energy expenditure during harsh

winter conditions, such as adequate cover and reduced snow depth. Specifically, we predicted larger

antlers in counties with higher proportions of agricultural, rangeland, and forest cover, which provide

accessible food and shelter. We also hypothesized that warmer temperatures during the growing season

would be associated with increased antler size, while higher winter precipitation would have a negative

effect due to increased energetic demands. Finally, we hypothesized that selective harvest of large-

antlered individuals over time has led to a temporal decline in antler size across Ontario through

reductions in age structure.

STUDY AREA

White-tailed deer in Ontario, Canada, occupy a diverse landscape with varying anthropogenic

influences. Historically, white-tailed deer populations have been concentrated in southern Ontario,

where winters are milder and growing seasons are longer compared to more northern parts of the

province (Kennedy-Slaney et al. 2018). This area is dominated by agricultural and deciduous forests,

which provide high-quality forage such as forbs, legumes, and other crops during spring and summer

(Ecological Stratification Working Group 1995, Government of Ontario 2021). However, white-tailed

deer exist throughout parts of central and northern Ontario, including the boreal forest region, where

they experience harsher winters. Their summer habitats consist of early successional forests with

abundant food sources which shifts to coniferous and mixed-wood forests in winter that offer shelter

from excessive snowfall and winter browse (Voigt et al. 1997, Hewitt 2011). Though browse is lower in

nutritional quality, it is readily used by white-tailed deer during Ontario's colder seasons, making up 74-

91% of their diet when other vegetation is scarce (Mautz *et al.*, 1976; Gray and Servello, 1995; Hewitt,

2011). Where they overlap, wolves (*Canis lupus*) are the primary predators of white-tailed deer in

Ontario, while coyotes (*Canis latrans*) fill that role in their absence, though do not fully replace wolves

ecologically (Benson et al. 2017). In addition to climate variation, Ontario's expanding infrastructure

and growing population has tripled in the last 50 years (Government of Canada 2016, 2022),

consequently reducing the available habitat and forage available for deer. As a result, deer are

increasingly concentrated in the habitats that remain suitable, including forest patches, crop fields, and

early successional areas (Voigt et al. 1997, Chen et al. 2015). However, the proliferation of agricultural

landscapes might increase food availability in some regions further north (Hewitt 2011).

METHODS

Data collection

We obtained Ontario antler data from The Foundation for the Recognition of Ontario Wildlife

(FROW) for the years 1941 to 2020. FROW is a non-profit organization that manages the records of big

game animals in Ontario. Antler measurements for white-tailed deer were voluntarily submitted by

hunters to FROW from every county in Ontario, except for Temiskaming. The antler dataset included

date of kill (DOK), antler type (typical or atypical), tine numbers and multiple scores following the

Boone and Crockett scoring system, which assigns scores based on additive measurements of the antler

including tine length, beam circumference, and spread. FROW has a minimum requirement that must be

met in order for a white-tailed deer to be entered into the record book (atypical net score 130 and typical

net score 120; Foundation for the Recognition of Ontario Wildlife 2015), so we refer to these deer as

"record deer" hereafter. Minimum scoring requirements increased in 2006, so all records prior to 2006

below this adjusted score were excluded. Because we were interested in assessing the influence of

spatial and temporal variables on records, we removed entries that did not contain a county or DOK.

Lack of landcover data dating back pre--1999 resulted in all record deer entries prior to 1997 being

removed in the spatial models. We then assessed the influence of a series of environmental and

temporal variables on tine number, gross score, net score, symmetry and the number of record deer per

county.

Predictor variables in our model included temperature ($^{\circ}$C), precipitation (mm), and number of

growing degree days (GDD, a measure of accumulated warmth above a threshold needed for plant

growth) for each county corresponding to the year a record deer was harvested (Table S1, available in

Supporting Information). These variables were obtained from Natural Resources Canada (Hutchinson et

al. 2009, Hopkinson et al. 2011, McKenney et al. 2011). We extracted temperature and precipitation

data from the months March to August as these are the months when antler growth occurs (Hewitt

2011). For precipitation, we also extracted the months January to March to gauge winter severity

through snow fall. Precipitation data were averaged. For temperature, we followed Siegert *et al.* (2013)

and calculated the yearly temperature difference by subtracting each county's average temperature over

the time series from the temperature for each individual year. The temperature and precipitation data for

one year prior to the DOK and two years prior to the DOK were also extracted.

ArcGIS pro (version 3.1.3;(Johnston and Environmental Systems Research Institute 2004), was

used to extract landcover data. We used two land cover layers: the Ontario Land Cover Compilation

v.2.0 (Land Information Ontario 2023) which captured the years 1999 to 2011 and Ontario Land Cover

Version 1.0 (Ontario Land Cover Version 1.0 - Overview n.d.) which captured the years 2015 to 2020.

Both land cover layers are raster datasets with a 15-metre spatial resolution and a minimum mapping

unit of 0.5 hectares. Depending on their DOK, the record deer were assigned to a landcover dataset

(Table S1). For the years not captured by the landcover databases (2012-2014), whichever layer

encompassed the year closest to the DOK was used. Reclassification of the land cover categories within

these databases was completed to ensure consistency between land cover types: reclassified categories

were; agriculture, infrastructure, deciduous forest, coniferous forest, mixed forest, other forest,

rangeland, water, wetlands, and other landcover types (Fig. 1, Table S2). The proportion of each

landcover type per county was calculated (Table 1), and "other landcover" was dropped in order to

interpret the relationship between the response variable and the proportion covariates (see Valle *et al.*,

2024).

The final three predictor variables used were average county elevation (meters; TessaDEM v1.2

2024), county size (km^2) and deer abundance (Table S1). Average county elevation and county size are

fixed variables, with the values used per county staying constant over the study period. A relative index

of white-tailed deer abundance per county was generated using data from provincial hunter surveys

(Government of Ontario 2025). Each year, the Ontario Ministry of Natural Resources collects

information online or by phone through mandatory hunter surveys on hunting activities. As part of this

survey, hunters are asked to report the number of deer they saw during their hunt and the total number

of days they hunted, providing a metric of the deer seen per day (Government of Ontario 2025). These

values were averaged across all available years (2008-2023) in each county to provide a single estimate

per county.

Statistical analysis

The following predictor variables were standardized (centered to a mean of zero and scaled to a

standard deviation of one) prior to analysis: average county elevation, precipitation (winter and antler

growing season - March to August), estimated deer abundance, and GDD. Temperature predictors were

not standardized because they were already transformed to reflect deviations from long-term county-

level averages. We assessed pairwise correlations among the variables, removing those with a

correlation greater than 0.70 or less than -0.70 (i.e. Dormann et al. 2013). Landcover variables were

excluded from the correlation assessment among themselves since they are proportion data, therefore,

changes in one necessarily correspond to changes in others, making high correlations expected and not

indicative of problematic collinearity (Valle et al. 2024). However, we did assess correlations between

landcover variables and other covariates. We then fit a series of models to each response variable. All

spatial models were fitted using the S.CARmultilevel function from the CARBayes package in R (

v.6.1.1Lee, 2013, R Core Team 2023). These models capture spatial autocorrelation by incorporating

spatial random effects modeled with a Leroux conditional autoregressive (CAR) prior, where

neighboring counties are defined by shared borders. Model fitting occurred within a Bayesian

framework using Markov chain Monte Carlo (MCMC) simulation with default priors applied to all

model parameters. All model specifications, including response variables and full lists of predictors, can

be found in Table S3, available in supporting information.

To assess the number of record deer harvested in each county we used a Poisson regression with

a log link and a county area offset to translate the analysis into one of rates of record deer per area. We

summed the total number of record deer harvested per county across all years, so DOK was not

considered when assigning temperature and precipitation in this model. Instead, average values for the

temperature and precipitation of each county over the specified time span were used. The averaging of

these environmental variables allowed us to observe relationships between the environment and the

spatial distribution of record deer harvests across Ontario counties over the entire study period.

To evaluate antler characteristics, we used DOK when assigning temperature, precipitation and

landcover values which allowed us to observe spatial-temporal relationships at the individual level.

Each model used one of three distributions and their standard link functions: Poisson regression model

with a log link to assess the number of antler tines, a binomial regression model with a logistic link

(logit) to assess antler symmetry (symmetrical = 0, asymmetrical = 1), with asymmetry defined as

unequal tine counts between antlers rather than the broader Boone and Crockett "non-typical"

classification, and gaussian regression models with identity links to assess antler score (Net and Gross).

Antler score was also log-transformed to normalize the distribution and ensure the assumptions of the

model were met.

For CAR models, we evaluated parameter effects based on the probability of direction (pd)

which provided the probability each covariate was either positively or negatively related to antler

characteristics (Makowski et al. 2019). Parameters with a probability $\geq 97.5\%$ were considered to have

a high probability of affecting the antler trait. Parameters with a probability of 95 – 97.5% were

considered to have a moderate probability of affecting the antler trait, and parameters with a probability

of 90 – 95% were considered to have a weak, but present, probability of affecting the antler trait. All

probabilities below 90% were considered to have no effect (Lohr et al. 2020, Dias et al. 2025). Spatial

autocorrelation (ρ) was interpreted such that values near 0 indicated spatial independence and values

near 1 indicated strong spatial dependence, so, ρ was considered high if > 0.7, moderate if 0.3–0.7, and

low if < 0.3. Spatial variance (τ^2) and residual variance (σ^2) were considered high if values approached

or exceeded 1 and low if around or below 0.01, based on the Inverse-Gamma (1, 0.01) priors, which

assume most variation is near zero unless the data strongly indicates otherwise (Lee 2013).

Finally, standard generalized linear models (GLMs) from the base r package (R Core Team

2023) were used to assess temporal trends in antler score and tine number, with year as the predictor

variable. Harvest data was sparse pre 1980 and so we used only entries from 1980 onward for these

GLMs. No additional offsets or weights were applied to the models. Model assumptions were assessed

using standard analyses, including assessing normality and homoscedasticity of residuals for the

Gaussian models, and overdispersion for Poisson and Binomial models. It should also be noted that this

data was hunter-submitted and therefore likely to be non-random and influenced by variation in hunting

effort or harvest efficiency, which was not explicitly captured in our models.

RESULTS

The final dataset after cleaning consisted of 2,413 entries between 1997 and 2020, and 3,072

entries between 1980 and 2020. Number of growing days was removed due to its high correlation with

temperature (Fig S1, available in Supporting Information). Temperature was kept as it directly reflects

seasonal climate variation and is easier to interpret. Deer abundance was also removed due to its high

correlation with agricultural landcover (Fig S1, available in Supporting Information). Residuals from

gaussian models were normal, and no overdispersion was detected in binomial or Poisson models.

Average temperature had a strong probability of having a positive effect on the number of record

deer per county (Fig. 2; $\beta = 1.24$, pd $= 99.9\%$), with years 1 °C warmer than the county average

associated with 3.5 times more record deer, corresponding to a 1.24 increase on the log scale. Elevation

showed a weak probability of having a positive effect (Fig. 2; $\beta = 0.31$, pd $= 94.3\%$), with a 0.35%

increase in the expected number of record deer per 1 m increase in elevation. This model had a strong τ^2

(0.98) and moderate ρ (0.46). For tine number, the temperature one year prior to the year of harvest had

a strong probability of having a positive effect (Fig 2; $\beta = 0.02$, pd $= 99.1\%$), with a 1 °C increase in the

average temperature during the year prior to harvest associated with 2% more tines. The temperature of

the year of harvest and the % of Rangeland per county both had weak probability of having positive

effects (Fig 2 & 3; $\beta = 0.01$, pd $= 91.1\%$ and $\beta = 0.94$, pd $= 92.2\%$), suggesting that years 1 °C warmer

than the county average and counties with a higher percentage of rangeland were associated with 1%

and 156% (2.6-fold) more tines, respectively. This model had a low τ^2 (0.003) and moderate ρ (0.63).

For antler symmetry, precipitation during antler growth two years prior to the year of harvest had a

strong probability of having a negative effect (Fig 2; $\beta = -0.12$, pd $= 98.6\%$), corresponding to a 0.85%

decrease in the odds of asymmetrically antlered deer per 1 mm increase in precipitation, while the

temperature of the year of harvest had a moderate probability of having a positive effect (Fig 2; β =

0.10, pd = 95.6%) , corresponding to an 11% increase in the odds of asymmetrically antlered deer per 1

°C warmer year. This model had a low τ^2 (0.03) and moderate ρ (0.38).

For Gross Score, we found the temperature the year of harvest had a strong probability of having

a positive effect (Fig 2; β = 0.01, pd = 100%), with years 1 °C warmer than the county average

associated with a 1% increase in Gross Score. Winter precipitation during the year of harvest had a

strong probability of having a negative effect (Fig 2; β = -0.01, pd = 97.8%), with a 1 mm increase in

winter precipitation associated with a 0.05% decrease in Gross Score. Precipitation during antler growth

the year of harvest and precipitation during the winter one year prior to the year of harvest both had

moderate probabilities of having positive effects (Fig 2; β = 0.01, pd = 96.8% and β = 0.01, 96.5%,

respectively), with a 1 mm increase in precipitation during antler growth the year of harvest being

associated with a 0.07% increase in Gross Score and a 0.05% increase in Gross Score for winter

precipitation one year prior to harvest. Temperature two years prior to harvest and the % of Deciduous

Forest per county then had weak probabilities of positive effects (Fig 2 & 3; β = 0.003, pd = 91.4% and

β = 0.22, 93.4%), with 0.3% increase per 1 °C above the county average and 25% increase per 1%

more Deciduous Forest, respectively. This model had τ^2 and v^2, both of which were relatively low

(0.001 and 0.01 respectively), and a moderate ρ (0.70).

For Net Score, temperature the year of harvest and temperature two years prior to the year of

harvest had strong probabilities of having a positive effect (Fig 2, β = 0.01, pd = 100%, β = 0.01, pd =

99.2% respectively), with each suggesting a 1% increase in Net Score per 1 °C increase above the

county average. The % of rangeland per county also had strong probabilities of having a positive effect

(β = 0.58, pd = 98.9%), with a 79% increase in Net Score per 1% more rangeland. The % of Deciduous

Forest per county then had a moderate probability of having a positive effect (Fig 3, β = 0.27, pd =

96.0%), with a 31% increase in Net Score per 1% more deciduous forest. The precipitation during antler

growth the year of harvest and the % of Coniferous Forest per county both had weak probabilities of

positive effects (Fig 2 & 3; β = 0.004, pd = 92.8% and β = 0.28, pd = 94.8%, respectively), with a 1 mm

increase in precipitation during antler growth the year of harvest being associated with a 0.03% increase

in Net Score and a 1% increase in coniferous forest being associated with a 32% increase in Net Score.

The precipitation during the winter the year of harvest then had a probability of having a weak but

negative effect (Fig 2; β = -0.004, pd = 91.1%), with a 1 mm increase in precipitation during the winter

the year of harvest being associated with 0.02% decrease in Net Score. τ^2 and v^2 were relatively low

(0.002 and 0.01 respectively), and ρ was moderately high (β =0.74). Finally, there was no temporal

trend in size and tine number over time (Table S4, available in Supporting Information).

DISCUSSION

We investigated spatial drivers and temporal patterns of harvested record deer in Ontario,

Canada. We observed that increases in temperature during the months that antler growth is occurring

(March-August) positively influenced all antler characteristics (Fig 2), and that rangeland and forest are

key landscape components, the amounts of which are correlated to overall greater antler size. These

patterns are consistent with our hypothesis that conditions providing abundant forage and adequate

cover will promote antler growth. Higher winter precipitation had negative effects, consistent with our

prediction that conditions increasing energetic demands, such as deep snow and limited forage, would

hinder antler growth. Interestingly, no change in antler size was detected over time in Ontario (Fig. 4),

which does not support the idea of regular harvest of large-antlered individuals leading to a temporal

decline through reductions in age structure. This finding is likely due to the province's position at the

northern edge of the white-tailed deer's range and a relatively stable population age structure.

Historically, harsher winters and limited forage could have constrained antler growth, but ongoing

warming due to climate change is likely improving habitat suitability by reducing winter severity and

extending growing seasons, which could be helping maintain antler size in Ontario (Lesage et al. 2001,

Dawe and Boutin 2016).

This data set is representative of adult antlered white-tailed deer within Ontario and although not

a random sample, the data captures a large range of variation in antler size across the population (Fig

S2, available in Supporting Information) and temporal, spatial and environmental influences should be

reflected in this large sample of male white-tailed deer (see also Strickland and Demarais, 2008;

Monteith *et al.*, 2013; Cain *et al.*, 2019). However, because participation in record programs is

voluntary and biased toward larger antlered deer, those with smaller antlers are likely underrepresented

(LaSharr et al. 2019). As a result, the dataset might not fully represent population wide variation in

antler size, but relative patterns and relationships with environmental variables that are still informative

and can help to understand the factors influencing antler development across Ontario (LaSharr et al.

2019). It should also be noted that, although some effect sizes presented are small, they nonetheless

represent measurable changes that will be considered in the interpretation of this study.

To account for spatial dependence in our data and better understand regional patterns, we used

CAR models, which revealed moderate to high spatial autocorrelation (ρ) in all models, revealing that

nearby counties tend to have similar conditions (e.g., temperature). However, these values also indicate

that some unaccounted-for spatial covariates are still present but not explicitly modeled. Possible

missing covariates worth further exploration include land productivity, which affects forage availability,

landcover structure that shapes shelter and feeding opportunities, and the distribution of minerals like

phosphorus and calcium, which is influenced by vegetation type and tissue age and affects bone and

antler development (Grasman and Hellgren 1993, Cain et al. 2019, Turner et al. 2025), though the

appropriate GIS layers currently do not exist for Ontario. We also observed low τ^2 in models assessing

antler characteristics indicating that, although the predictors often have similar conditions in

neighboring counties, antler size does not exhibit strong spatial structure. However, a strong τ^2 when

assessing the count of record deer per county does suggest that counties with higher numbers of record

deer are more likely to have neighboring counties with similarly high counts, likely reflective of similar

forage resources and variation in regional hunting interest and behaviour. Notably, hunter behaviour and

success which will vary with the amount of roads, dwellings, and with accessibility, while regional

differences in the number of hunters and frequency of hunting might also contribute to these spatial

patterns (Kilgo et al. 1998, Bonnot et al. 2013).

Environmental drivers of record antlers

Temperature was the most consistent environmental driver of antler traits: higher temperature

was related to an increase in antler size metrics in all models (Fig 2). Warmer temperatures have been

linked to larger antlers and greater reproductive success in white-tailed deer and other Cervidae

(Clements et al. 2010, Simard et al. 2010). Increased plant growth from warmer conditions means more

forage availability for the deer, which will supply the nutrients and energy needed for antler growth

(French et al. 1956, Hatfield and Prueger 2015). Here, the unique annual regrowth of antlers makes

them strongly affected by yearly environmental changes (Foley et al. 2012). When looking at tine

number and score, the temperature one year prior to harvest and two years prior to harvest respectively,

also had positive effects on antler size and shape (Fig. 2). Antler growth depends on the accumulation of

resources stored in the animal's body (i.e. mineral and fat reserves) (French et al. 1956, Lesage et al.

2001). Warmer conditions that promote forage growth in prior years can enhance fat reserves, which are

then used in subsequent antler cycles when these nutrients is needed (Hewitt 2011).

Similar inference can be made from the influence of precipitation during antler growth the year

of harvest on antler size. Wetter conditions can lead to increased plant growth which could again affect

the annual regrowth of antlers (Foley et al. 2012, Scharwies and Dinneny 2019). Conversely, we saw

negative effects from precipitation during the winter the year of harvest on antler size (Fig 2). More

snow in winter can make it more difficult for the deer to move around and find forage, which causes

them to congregate in deer yards (Government of Ontario 2019*b*). Deer yards are an area made up of

coniferous trees that have a large canopy for blocking excessive snowfall (Government of Ontario

2019*b*). With deer being confined to these yards deer density and the availability of forage for each

individual animal decreases (Patterson and Power 2002). This could then put individuals with poor body

conditions at a disadvantage and minimal fat reserves when antler regrowth commences (Lesage et al.

2001). Future work could strengthen this finding by incorporating direct measurements of snow depth,

snow cover duration, or broader indices such as the Winter Severity Index (WSI) to capture more details

on the energetic challenges white-tailed deer face during the winter months.

Rangeland and forest landcover had positive effects on antler characteristics, meaning

that as the percentages of these types of landcover in each county rose so too did antler score and

tine number (Fig 5). Areas combining forest with regions that provide abundant forage have been

associated with higher harvests of record deer, likely due to greater amounts of high-contrast

edges, which create habitat interfaces that offer the resources necessary for male deer to produce

large antlers (Cain et al. 2019). In our case, forests provide cover, which block excessive

snowfall, and contain the northern deer's primary winter food source, browse (Hewitt 2011), and

rangeland provides a more variable and nutritious food sources such as forbs, while also

allowing for greater visibility of large antlered bucks from further away, potentially increasing

hunter success or improving the effectiveness of dominance displays and mate attraction (Gilbert

et al. 2006, Hewitt 2011, Houde et al. 2020). Collectively, these habitat interfaces are preferred

by white-tailed deer as it allows them to have a larger variety of food sources throughout the year

and provides both more space for movement among resources (rangeland) while still having

adequate cover from predators and severe winters (forest) (Voigt et al. 1997, Bartel et al. 2025).

Number of harvested deer and antler symmetry

We observed that increased average temperatures were associated with more record white-tailed

deer being harvested in a given Ontario county with years 1 °C warmer than the county average

associated with 3.5 times more record-scoring deer in a given county. Similar to our assessment of this

trend seen in our antler size and configuration models, warmer temperatures result in more forage at the

northern edge of the white-tailed deer's range and thus more energy is able to be allocated to individual

deer in a county, presumably allowing more individuals to reach record deer status. The number of

record deer harvested was not significantly affected by any landcover covariates (Fig 3); so, while

landcover type positively impacts antler size, the overall effect is minor such that it does not increase

the net number of large animals harvested in a given year. Curiously, record deer numbers were

positively affected by elevation (Fig. 2). It is tempting to suggest this being reflective of mature deer

finding refuge from human activity via fewer roadways at these elevations. Less trafficked zones allow

deer to move more freely and access critical resources with minimal human interference (Hamilton et al.

2024) and behaviourally deer learn to avoid humans (Kilgo et al. 1998, Marantz et al. 2016). However,

the elevational range is small in the province (0 m – 700 m) and obfuscated when averaged across the

county as done here (66 m – 415 m), and thus finer scale analysis is warranted to better understand this

effect.

The symmetry of antlers appears to be impacted by precipitation, with higher precipitation levels

two years prior to antler growth resulting in more symmetrically antlered deer. Since white-tailed deer

can develop antler asymmetry when exposed to stressors such as environmental conditions, disease, or

injury that affect energy allocation before and during antler growth (Marburger et al. 1972, Goss 1983,

Wild et al. 2022), the mechanism by which precipitation two years prior impacts symmetry is still

unclear. However, there is the potential for lagged environmental effects on antler morphology which

has been observed in other studies, with Strickland et al. (2020) demonstrating that conditions

experienced by male white-tailed deer two years prior were associated with variation in antler mass.

Likewise, Patterson and Power (2002) suggested that antler beam size was affected by lagged climatic

effects through influences on habit and population density. Symmetry has also been seen to decrease as

trait size increases (De Coster et al. 2013), which we observed in our data set (Fig S3, available in

Supporting Information); however, we see no strong relationship between precipitation and antler size.

It is thus still conceivable that environmental conditions might have long-term impacts on symmetry,

even for antlers that are shed annually.

Temporal patterns of record deer

Over the last 40 years, there has been no significant change in antler size (or score) and

the tine number of harvested record white-tailed deer in Ontario (Table S4). This contrasts the

findings of Monteith et al. (2013) who reported a negative trend across North America in antler

size in white-tailed deer from 1950 to 2008, based on trophy records of mature males submitted

to the Boone and Crockett Club's *Records of North American Big Game* (Boone and Crockett

Club 2025). These negative trends were hypothesized to be a combination of harvest-induced

reduction in age structure, sociological effects and density dependence (Monteith et al. 2013).

Our finding suggests that antler trends in white-tailed deer are region specific and vary

depending on the environment and pressures present.

In Ontario, the number of active hunters has stayed relatively consistent since 2008 while the

number of records submitted to FROW has declined (Fig 6). This could be indicative of a possible shift

in hunting priorities (Monteith et al. 2013), but multiple severe winters in 2007/08 and 2010/11, have

coincided with apparent population declines (Ontario Outdoors Magazine Game Forecast 2009, 2011)

which would explain a decrease in entries. Still, the number of white-tailed deer in Ontario is relatively

low especially compared to other regions of North America where population numbers are substantially

higher. For example, Michigan with similar environmental conditions to Ontario and an area of ~

250,493 km^2 has a population of ~2 million white-tailed deer, compared to Ontario with an area of over

1 million km^2 and a white-tailed deer population of ~400 thousand (Mason 2016, Conner 2021). This

lower density is likely due to Ontario being the northern edge of the white-tailed deer's range, where

limited suitable habitat likely has more influence on antlers than harvest pressure or hunting regulations.

While Ontario's hunting regulations could result in lower overall harvest rates compared to other

regions, it is not clear if these restrictions make any substantial difference in population dynamics or

antler development. Instead, environmental factors associated with climate and habitat likely have a

greater influence on population density and antler growth in the province.

In some parts of North America, warmer climates lead to longer growing seasons allowing

white-tailed deer to have more time to access forage and accumulate resources, supporting antler

development allowing them to reach their maximum physical size during antler growth (Mysterud et al.

2005, Sontheimer et al. 2024). In contrast, Ontario's relatively shorter growing seasons and colder

winters limit access to nutritious food and deplete body reserves, which could reduce the amount

resources deer can allocate to antler growth in the subsequent spring; this limitation could mean there is

still room for growth. Our models indicate that warmer temperatures in Ontario are positively

influencing antler development, likely because milder winters and longer growing seasons improve food

availability. Combined these conditions could be helping bolster antler size despite pressures that are

causing declines elsewhere (e.g. Monteith et al. 2013).

MANAGEMENT IMPLICATIONS

Our findings highlight how variation in habitat conditions across Ontario counties are related to differences in antler size and tine configuration as well as record white-tailed deer distributions. This emphasizes the importance of interpreting phenotypic trends relative to specific areas rather than generalizing across entire populations. Because environmental factors such as land cover composition and forage availability influence antler growth, where larger antlered bucks are desired, management practices could focus on maintaining or enhancing the quality of white-tailed deer habitats in Ontario, particularly in areas where forage is limited. Efforts such as sustaining forested habitat interfaces and managing deer densities to balance browse pressure could help improve nutritional conditions that support antler development.

With no significant temporal trend in antler size, the current harvest practices in Ontario are likely effective as it relates to maintaining antler size, but continued monitoring could be carried out to ensure that population dynamics remain stable. The long-term collection of harvest and antler data will be critical for tracking population health and evaluating the effects of management practices. Finally, by communicating these findings, hunters can better manage their expectations and recognize aspects of climate that could cause certain regions and years to have higher large-antlered white-tailed deer concentrations.

ACKNOWLEDGMENTS

This work was supported by NSERC Discovery Grant (ABAS; 410 SDC); Canadian Foundation for Innovation (ABAS): John R. Evans Leaders Fund (ABAS) and the Ontario Early Researcher Award (ABAS). Trent University is located on the traditional territory of the Michi Saagiig Anishnaabeg. We thank 4 anonymous reviewers for their helpful comments.

ETHICS STATEMENT

No approval of research ethics committees was required to accomplish the goals of this study.

DATA AVAILABILITY

The antler data that support the findings of this study are available from The Foundation for the

Recognition of Ontario Wildlife, Ontario, Canada and were acquired under research agreements for this

study. All other model covariates are provided.

CONFLICT OF INTEREST

Authors declare no competing interests.

REFERENCES

Bartel, S. L., L. Hakkila, and J. L. Orrock. 2025. Deer Vigilance and Movement Behavior Are
        Affected by Edge Density and Connectivity. Ethology 131:22–30.

Benson, J. F., K. M. Loveless, L. Y. Rutledge, and B. R. Patterson. 2017. Ungulate predation and
        ecological roles of wolves and coyotes in eastern North America. Ecological
        Applications: A Publication of the Ecological Society of America 27:718–733.

Bonnot, N., N. Morellet, H. Verheyden, B. Cargnelutti, B. Lourtet, F. Klein, and A. J. M.
        Hewison. 2013. Habitat use under predation risk: hunting, roads and human dwellings
        influence the spatial behaviour of roe deer. European Journal of Wildlife Research
        59:185–193.

Boone and Crockett Club. 2025. Scoring Your Trophy: typical whitetail deer. Boone and
        Crockett Club. <https://www.boone-crockett.org/scorechart/add/whitetail_deer>.
        Accessed 3 Mar 2025.

Cain, R. L., N. P. Snow, J. C. Crawford, D. M. Williams, and W. F. Porter. 2019. Spatial
        distribution and landscape associations of large-antlered deer. The Journal of Wildlife
        Management 83:1762–1772.

Chen, D., H. Wong, P. Belanger, K. Moore, M. Peterson, and J. Cunningham. 2015. Analyzing
        the Correlation between Deer Habitat and the Component of the Risk for Lyme Disease
        in Eastern Ontario, Canada: A GIS-Based Approach. ISPRS International Journal of Geo-
        Information 4:105–123.

Clements, M. N., T. H. Clutton-Brock, S. D. Albon, J. M. Pemberton, and L. E. B. Kruuk. 2010.
        Getting the timing right: antler growth phenology and sexual selection in a wild red deer
        population. Oecologia 164:357–369.

Clutton-Brock, T. H. 1982. The Functions of Antlers. Behaviour 79:108–125.

Conner, M. 2021. What is the deer population in Ontario? - ruggedthuglife.com. Canada Online
        Guide. <https://ruggedthuglife.com/ontario/what-is-the-deer-population-in-ontario/>.
        Accessed 12 Jan 2025.

D'Angelo, G. J., and M. D. Grund. 2015. Evaluating Competing Preferences of Hunters and
        Landowners for Management of Deer Populations. https://doi.org/10.26077/N707-1D90.

Dawe, K. L., and S. Boutin. 2016. Climate change is the primary driver of white-tailed deer
        (Odocoileus virginianus) range expansion at the northern extent of its range; land use is
        secondary. Ecology and Evolution 6:6435–6451.

De Coster, G., S. Van Dongen, P. Malaki, M. Muchane, A. Alcántara-Exposito, H. Matheve, and
        L. Lens. 2013. Fluctuating Asymmetry and Environmental Stress: Understanding the
        Role of Trait History. PLoS ONE 8:e57966.

DeYoung, R. W., S. Demarais, R. L. Honeycutt, K. L. Gee, and R. A. Gonzales. 2006. Social
        Dominance and Male Breeding Success in Captive White-Tailed Deer. Wildlife Society
        Bulletin 34:131–136.

Dias, T., N. P. Lemoine, S. W. Yanco, M. Zimova, R. A. Bay, and B. C. Weeks. 2025. Long-
        term increases in wing length occur independently of changes in climate and climate-
        driven shifts in body size. Proceedings of the Royal Society B: Biological Sciences
        292:20242556.

Dormann, C. F., J. Elith, S. Bacher, C. Buchmann, G. Carl, G. Carré, J. R. G. Marquéz, B.
        Gruber, B. Lafourcade, P. J. Leitão, T. Münkemüller, C. McClean, P. E. Osborne, B.
        Reineking, B. Schröder, A. K. Skidmore, D. Zurell, and S. Lautenbach. 2013.

Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. Ecography 36:27–46.

Ecological Stratification Working Group. 1995. National Ecological Framework for Canada. Agriculture and Agri-Food Canada: Ottawa, ON.

Florida Fish And Wildlife Conservation Commission. 2025. Species Season Dates and Bag Limits. <https://myfwc.com/hunting/season-dates/>. Accessed 21 Jul 2025.

Foley, A. M., R. W. DeYoung, S. D. Lukefahr, J. S. Lewis, D. G. Hewitt, M. W. Hellickson, D. A. Draeger, C. A. DeYoung, and H. Beck. 2012. Repeatability of antler characteristics in mature white-tailed deer in South Texas: consequences of environmental effects. Journal of Mammalogy 93:1149–1157.

Foundation for the Recognition of Ontario Wildlife. 2015. Big Game Records of Ontario. Seventh edition. Beasley Brothers Outdoors Inc.

Freeman, E. D., R. T. Larsen, K. Clegg, and B. R. McMillan. 2013. Long-Lasting Effects of Maternal Condition in Free-Ranging Cervids. PLoS ONE 8:e58373.

French, C. E., L. C. McEwen, N. D. Magruder, R. H. Ingram, and R. W. Swift. 1956. Nutrient Requirements for Growth and Antler Development in the White-Tailed Deer. The Journal of Wildlife Management 20:221–232.

Garroway, C. J., and H. G. Broders. 2005. The quantitative effects of population density and winter weather on the body condition of white-tailed deer (Odocoileus virginianus) in Nova Scotia, Canada. Canadian Journal of Zoology 83:1246–1256.

Gilbert, C., A. Ropiquet, and A. Hassanin. 2006. Mitochondrial and nuclear phylogenies of Cervidae (Mammalia, Ruminantia): Systematics, morphology, and biogeography. Molecular Phylogenetics and Evolution 40:101–117.

Goss, R. J. 1983. Deer antlers: regeneration, function, and evolution. Academic Press, New York.

Government of Canada. 2016. Table 24.1 Population, by province and territory, selected years, 1861 to 2011. <https://www150.statcan.gc.ca/n1/pub/11-402-x/2012000/chap/pop/tbl/tbl01-eng.htm>. Accessed 26 Nov 2024.

Government of Canada, S. C. 2022. Profile table, Census Profile, 2021 Census of Population - Ontario [Province]. <https://www12.statcan.gc.ca/census-recensement/2021/dp-pd/prof/index.cfm?Lang=E>. Accessed 19 Sep 2023.

Government of Ontario. 2019*a*. Ontario Hunting Regulations Summary. <http://www.ontario.ca/document/ontario-hunting-regulations-summary>. Accessed 26 Oct 2025.

Government of Ontario. 2019*b*. 4 Identifying habitats of seasonal concentrations of animals | 1 Significant wildlife habitat technical guide | ontario.ca. <http://www.ontario.ca/document/significant-wildlife-habitat-technical-guide/4-identifying-habitats-seasonal>. Accessed 4 Feb 2025.

Government of Ontario. 2021. Forest regions. <http://www.ontario.ca/page/forest-regions>. Accessed 11 Oct 2023.

Government of Ontario. 2025. Hunter reporting. <http://www.ontario.ca/page/hunter-reporting>. Accessed 17 Jun 2025.

Grasman, B. T., and E. C. Hellgren. 1993. Phosophorus Nutrition in White-Tailed Deer: Nutrient Balance, Physiological Responses, and Antler Growth. Ecology 74:2279–2296.

Gray, P. B., and F. A. Servello. 1995. Energy Intake Relationships for White-Tailed Deer on Winter Browse Diets. The Journal of Wildlife Management 59:147–152.

Hamilton, K. M., T. Bommarito, and J. S. Lewis. 2024. Human activities influence mule deer use of overpasses across multiple scales. The Journal of Wildlife Management 88:e22560.

Harper, C. A. 2019. LANDOWNERS' GUIDE TO WILDLIFE FOOD PLOTS. The University of Tennessee Institute of Agriculture.

Hatfield, J. L., and J. H. Prueger. 2015. Temperature extremes: Effect on plant growth and development. Weather and Climate Extremes 10:4–10.

Hewitt, D. 2011. Biology and management of white-tailed deer. CRC Press, Boca Raton, FL.

Hopkinson, R. F., D. W. McKenney, E. J. Milewska, M. F. Hutchinson, P. Papadopol, and L. A. Vincent. 2011. Impact of Aligning Climatological Day on Gridding Daily Maximum-Minimum Temperature and Precipitation over Canada. Journal of Applied Meteorology & Climatology 50:1654–1665.

Houde, N., J.-P. Tremblay, N. Thiffault, and S. D. Côté. 2020. Manipulating forage and risk avoidance to increase white-tailed deer vulnerability to hunters. Wildlife Biology 2020.

Hutchinson, M. F., D. W. McKenney, K. Lawrence, J. H. Pedlar, R. F. Hopkinson, E. Milewska, and P. Papadopol. 2009. Development and Testing of Canada-Wide Interpolated Spatial Models of Daily Minimum–Maximum Temperature and Precipitation for 1961–2003. Journal of Applied Meteorology & Climatology 48:725–741.

Jamieson, A., S. J. Anderson, J. Fuller, S. D. Côté, J. M. Northrup, and A. B. A. Shafer. 2020. Heritability Estimates of Antler and Body Traits in White-Tailed Deer (Odocoileus virginianus) From Genomic-Relatedness Matrices. Journal of Heredity 111:429–435.

Johnston, K. and Environmental Systems Research Institute. 2004. Using ArcGIS geostatistical analyst. ESRI, Redlands, CA.

Kennedy-Slaney, L., J. Bowman, A. A. Walpole, and B. A. Pond. 2018. Northward bound: the distribution of white-tailed deer in Ontario under a changing climate. Wildlife Research 45:220–228.

Kilgo, J., R. Labisky, and D. Fritzen. 1998. Influences of Hunting on the Behavior of White☐Tailed Deer: Implications for Conservation of the Florida Panther. Conservation Biology 12:1359–1364.

Kruuk, E. B., J. Slate, J. M. Pemberton, S. Brotherstone, F. Guinness, and T. Clutton-Brock. 2002. Antler size in red deer: heritability and selection but no evolution. Evolution; International Journal of Organic Evolution 56:1683–1695.

Land Information Ontario. 2023. Ontario Land Cover Compilation v.2.0. <https://www.arcgis.com/home/item.html?id=7aa998fdf100434da27a41f1c637382c>. Accessed 22 Aug 2023.

Langvatn, R., and A. Loison. 1999. Consequences of harvesting on age structure, sex ratio and population dynamics of red deer Cervus elaphus in central Norway. Wildlife Biology 5:213–223.

LaSharr, T. N., R. A. Long, J. R. Heffelfinger, V. C. Bleich, P. R. Krausman, R. T. Bowyer, J. M. Shannon, E. M. Rominger, C. P. Lehman, M. Cox, and K. L. Monteith. 2019. Biological relevance of antler, horn, and pronghorn size in records programs. Journal of Mammalogy 100:1447–1458.

Lee, D. 2013. CARBayes: An R Package for Bayesian Spatial Modeling with Conditional Autoregressive Priors. Journal of Statistical Software 55:1–24.

Lesage, L., M. Crête, J. Huot, and J.-P. Ouellet. 2001. Evidence for a Trade-off between Growth and Body Reserves in Northern White-Tailed Deer. Oecologia 126:30–41.

Lohr, A. K., J. A. Martin, G. T. Wann, B. S. Cohen, B. A. Collier, and M. J. Chamberlain. 2020.
Behavioral strategies during incubation influence nest and female survival of Wild
Turkeys. Ecology and Evolution 10:11752–11765.
Makowski, D., M. S. Ben-Shachar, and D. Lüdecke. 2019. v. Journal of Open Source Software
4:1541.
Marantz, S. A., J. A. Long, S. L. Webb, K. L. Gee, A. R. Little, and S. Demarais. 2016. Impacts
of human hunting on spatial behavior of white-tailed deer (Odocoileus virginianus).
Canadian Journal of Zoology 94:853–862.
Marburger, R. G., R. M. Robinson, J. W. Thomas, M. J. Andregg, and K. A. Clark. 1972. Antler
malformation produced by leg injury in white-tailed deer. Journal of Wildlife Diseases
8:311–314.
Mason, R. 2016. Michigan Deer Management Plan.
Mautz, W. W., T. W. Walski, and W. E. Urban. 1976. Digestibility of Fresh Frozen versus
Pelleted Browse by Snowshoe Hares. The Journal of Wildlife Management 40:496–499.
McKenney, D. W., M. F. Hutchinson, P. Papadopol, K. Lawrence, J. Pedlar, K. Campbell, E.
Milewska, R. F. Hopkinson, D. Price, and T. Owen. 2011. Customized Spatial Climate
Models for North America. Bulletin of the American Meteorological Society 92:1611–
1622.
Michel, E. S., S. Demarais, B. K. Strickland, T. Smith, and C. M. Dacus. 2016. Antler
characteristics are highly heritable but influenced by maternal factors. 80:1420–1426.
Monteith, K. L., L. E. Schmitz, J. A. Jenks, J. A. Delger, and R. T. Bowyer. 2009. Growth of
Male White-Tailed Deer: Consequences of Maternal Effects. Journal of Mammalogy
90:651–660.
Monteith, K., R. Long, V. Bleich, J. Heffelfinger, P. Krausman, and R. Bowyer. 2013. Effects of
harvest, culture, and climate on trends in size of horn-like structures in trophy ungulates.
Wildlife Monographs 183:1–26.
Morina, D. L., S. Demarais, G. D. Chesser Jr., J. W. Lowe, and B. K. Strickland. 2018. Antler
manipulation procedures for use in social and behavioral studies of deer. Wildlife
Biology 2018:wlb.00452.
Mysterud, A. 2011. Selective harvesting of large mammals: how often does it result in
directional selection? Journal of Applied Ecology 48:827–834.
Mysterud, A., E. Meisingset, R. Langvatn, N. G. Yoccoz, and N. Chr. Stenseth. 2005. Climate-
dependent allocation of resources to secondary sexual traits in red deer. Oikos 111:245–
252.
Ontario Land Cover Version 1.0 - Overview. n.d.
<https://www.arcgis.com/home/item.html?id=667367a759214a089917adccdbae7cb2>.
Accessed 1 May 2024.
Ontario Outdoors Magazine Game Forecast. 2009. Hunting Forecast. 2009.
Ontario Outdoors Magazine Game Forecast. 2011. Hunting Forecast. 2011.
Patterson, B. R., and V. A. Power. 2002. Contributions of Forage Competition, Harvest, and
Climate Fluctuation to Changes in Population Growth of Northern White-Tailed Deer.
Oecologia 130:62–71.
Peterson, B. C., C. W. Schoenebeck, and N. J. Fryda. 2019. Effects of Extreme Environmental
Conditions on White-tailed Deer Antlers. Wildlife Society Bulletin (2011-) 43:708–716.
R Core Team. 2023. R: The R Project for Statistical Computing. <https://www.r-project.org/>.
Accessed 19 Jan 2023.

Rue, L. L. 1989. The deer of North America. Guilford, CT□: Lyons Press.
Scharwies, J. D., and J. R. Dinneny. 2019. Water transport, perception, and response in plants.
Journal of Plant Research 132:311–324.
Siegert, S., J. Broecker, and H. Kantz. 2013. Skill of data based predictions versus dynamical
models -- case study on extreme temperature anomalies.
Simard, M. A., T. Coulson, A. Gingras, and S. D. Côté. 2010. Influence of Density and Climate
on Population Dynamics of a Large Herbivore Under Harsh Environmental Conditions.
The Journal of Wildlife Management 74:1671–1685.
Sontheimer, W. P., D. Pfeffer, and F. W. Weckerly. 2024. Ontogenetic and static scaling of
antler mass in White-tailed Deer (Odocoileus virginianus). Journal of Mammalogy
105:134–142.
Strickland, B. K., and S. Demarais. 2007. Using Antler Restrictions to Manage for Older-aged
Bucks.
Strickland, B. K., and S. Demarais. 2008. Influence of Landscape Composition and Structure on
Antler Size of White-Tailed Deer. The Journal of Wildlife Management 72:1101–1108.
Strickland, B. K., P. G. Dixon, P. D. Jones, S. Demarais, N. O. Owen, D. A. Cox, K. Landry-
Guyton, W. M. Baldwin, and W. T. McKinley. 2020. Cohort antler size signals
environmental stress in a moderate climate. International Journal of Biometeorology
64:611–621.
TessaDEM v1.2. 2024. Near-global 30-meter Digital Elevation Model combining AW3D30,
MERIT DEM, Forest Height, World Settlement Footprint, and Global Surface Water
datasets. Available under Open Database License (ODbL) v1.0. <https://tessadem.com/>.
Accessed 16 Jun 2025.
Townsend, T. W., and E. D. Bailey. 1981. Effects of Age, Sex and Weight on Social Rank in
Penned White-tailed Deer. The American Midland Naturalist 106:92–101.
Turner, M. A., C. A. Harper, M. A. Lashley, B. K. Strickland, and M. Q. Wilber. 2025. Plant
nutrient concentrations inform white-tailed deer diet limitations. Journal of
Environmental Management 375:124186.
Turner, M. M., C. S. Deperno, W. Booth, E. L. Vargo, M. C. Conner, and R. A. Lancia. 2016.
The mating system of white-tailed deer under Quality Deer Management: Mating System
of White-Tailed Deer Under QDM. The Journal of Wildlife Management 80:935–940.
Valle, D., J. Mintz, and I. V. Brack. 2024. Estimation and interpretation problems and solutions
when using proportion covariates in linear regression models. Ecology 105:e4256.
Voigt, D. R., J. D. Broadfoot, and J. A. Baker. 1997. Forest Management Guidelines for the
Provision of White-tailed Deer Habitat.
Webb, S. L., S. Demarais, B. K. Strickland, R. W. DeYoung, B. P. Kinghorn, and K. L. Gee.
2012. Effects of selective harvest on antler size in white-tailed deer: A modeling
approach. The Journal of Wildlife Management 76:48–56.
Wild, M. A., G. A. Sargeant, K. Garrison, and D. Conradson. 2022. Association of antler
asymmetry with hoof disease in elk. The Journal of Wildlife Management 86:e22245.
714

TABLES

Table 1. Percentage of land cover types in Ontario, 1999 – 2011 vs 2015 – 2020.

FIGURES

Figure 1. Approximate Ontario land cover distribution (Left) and elevation (Right) divided by county.

Figure 2. Probability of direction (pd) of elevation, temperature, and precipitation on the response variables. Variables in pink have a high probability affecting the antler trait (97.5%). Variables in Orange have a moderate probability (95 – 97.5%). And Variables in green have a weak, but present probability (90 – 95%).

Figure 3. Probability of direction (pd) of landcover proportion on the response variables. Variables in pink have a high probability affecting the antler trait (97.5%). Variables in Orange have a moderate probability (95 – 97.5%). And Variables in green have a weak, but present probability (90 – 95%).

Figure 4. Change in antler score and tine number over time. Cutoffs are representative of Boone and Crockett data, depicting the higher cutoff size for record antlers in other parts of North America.

Figure 5. Percent of significant landcover per predicted antler score and tine number.

Figure 6. Number of records submitted to the Foundation for the Recognition of Ontario Wildlife (FROW) from 1997 to 2020 (blue) and Number of active hunters from 2008 to 2023 (pink).

APPENDIX A. SUPPLEMENTAL

Landcover Type
- Water
- Coniferous Forest
- Deciduous Forest
- Mixed Forest
- Other Forest
- Wetlands
- Rangeland
- Infrastructure
- Agriculture
- Other

Elevation (meters)
- 699
- 0

0   175   350   700 Kilometers