Complete genome sequences of *Fusobacterium watanabei* sp. isolates

Martha A. Zepeda-Rivera[1], Falk Ponath[2], Kaitlyn N. Lewis[1], Rutika P. Gavate[1], Floyd E. Dewhirst[3,4], Junko Tomida[5], Yoshiaki Kawamura[5], Kaori Tanaka[6], Susan Bullman[2], Christopher D. Johnston[1#]

[1]Genomic Medicine, MD Anderson Cancer Center, Houston, Texas, USA

[2]Immunology, James P. Allison Institute, MD Anderson Cancer Center, Houston, Texas, USA

[3]ADA Forsyth Institute, Cambridge, MA, USA

[4]Department of Oral Medicine, Infection and Immunity, Harvard School of Dental Medicine, Boston, Massachusetts, USA

[5]Department of Microbiology, School of Pharmacy, Aichi Gakuin University, Nagoya, Aichi, Japan

[6]Division of Anaerobe Research, Life Science Research Center, Gifu University, Yanagido, Gifu, Japan

[#]**Address Correspondence to**:

Christopher D. Johnston, PhD, cdjohnston@mdanderson.org

Keywords: Fusobacterium, nomenclature, Illumina sequencing

Abstract

We report the complete genome sequences of eight *Fusobacterium watanabei* clinical isolates,

ranging from 1.95 to 2.09 Mbp. Analysis against the Genome Taxonomy Database (GTDB)

indicates that *Fusobacterium watanabei* genomes are part of the "*Fusobacterium nucleatum_J*"

group, which also encompasses the previously published FNU strain and *Fna* C1 isolates.

Announcement

Fusobacterium watanabei was initially described as a novel species closely related to the

Fusobacterium nucleatum sensu lato subspecies: *F. nucleatum* subsp. *animalis, F. nucleatum*

subsp. *nucleatum, F. nucleatum* subsp. *polymorphum, F. nucleatum* subsp. *vincentii*[1]. Species

designation was based on whole genome comparisons of eight *Fusobacterium watanabei* clinical

isolates against *Fusobacterium nucleatum* sensu lato subspecies members. Analysis indicated

average nucleotide identity (ANI) and digital DNA-DNA hybridization (dDDH) metrics consistent

with species-level assignment[1]. Here, we report and release the genome sequences for these eight

clinical isolates of *Fusobacterium watanabei* (Table 1). Genome assembly indicates a genome size

range from 1.95 to 2.09 Mbp (average of 2.01 Mbp) with GC content ranging from 26.82% to

26.99% (average of 26.92%).

Although the PAGU 1796 (type strain) 16S rRNA gene and partial sequences of three

marker genes were available at National Center for Biotechnology Information (NCBI), the draft

genome was only available in the Short Read Archive. Here, analysis of these eight *Fusobacterium*

watanabei assembled genomes, including PAGU 1796 (type strain), against the Genome

Taxonomy Database using GTDB-tk[2] assigned them all to the currently designated

"*Fusobacterium nucleatum_J*" group, a member of the *Fusobacterium nucleatum* sensu lato group

of taxa. "*Fusobacterium nucleatum_J*" also encapsulates the strain 13-08-02, strain

"*Fusobacterium nucleatum* subsp. *unique* (FNU)" previously described in Ma et al.[3], and members

of the group we recently described as "*Fusobacterium nucleatum* subsp. *animalis* clade 1 (*Fna*

C1)"[4]. ANI values between "*Fusobacterium watanabei*", "*FNU*", and "*Fna* C1" genomes range

from 98.1228% to 99.9995%, supporting the observation that all three of these labels represent the

same *Fusobacterium* lineage (**Figure 1**). The lack of an annotated genome for *Fusobacterium*

watanabei in NCBI limited the ability of investigators to recognize and apply the name

"*Fusobacterium watanabei*" to describe FNU[3] and *Fna* C1[4] genomes. Nevertheless, despite the

use of different names, observations regarding this lineage are in agreement across publications[1,3,4].

The taxonomy of *Fusobacterium* species has recently undergone significant revision[5].

Based on the proposal by Kook et al.[6] to elevate *Fusobacterium nucleatum* sensu lato subspecies

to the species level assignments, the four historic subspecies have been revised to individual

species: *Fusobacterium animalis, Fusobacterium nucleatum, Fusobacterium polymorphum* and

Fusobacterium vincentii. Within this context of changing nomenclature, it is fortuitous that

Fusobacterium watanabei is a recognized species name under the List of Prokaryotic names with

Standing in Nomenclature (LPSN)[7] and can now be used for genomes in databases designated as

"*FNU*", "*Fna* C1" and "*F. nucleatum_J*". Here, we have publicly released genomes for strains

PAGU 1795, PAGU 1796, PAGU 1797, PAGU 1798, PAGU 1799, PAGU 1800, PAGU 1801, and

PAGU 1802. This increases the number of representative genomes to 34 for the *Fusobacterium*

watanabei ("*FNU*"/"*Fna* C1"/"*F. nucleatum_J*") group, which remains the closest phylogenetic

lineage to the colorectal cancer-associated *Fusobacterium animalis*[4]. Finally, it is noteworthy that

the PAGU 1796 *Fusobacterium watanabei* type strain, previously deposited to DSMZ under

CCUG 74246, now has an available assembled genome deposited under PRJEB87340.

Figure

 Figure 1: Average nucleotide identity between "*Fusobacterium watanabei*", "*FNU*", "*Fna*
 C1" genomes
 Clustered dendrogram depicts the FastANI values between *Fusobacterium watanabei* genomes,
 including the previously published "*FNU*"[3] (‡) , and "*Fna* C1"[4] (*) strains. Color of each
 individual box represents the fastANI value, ranging from 98.0% (white) to 100% (dark green).

Table

Genome Name	Size (bp)	GC (%)	No. Contigs	No. Reads	N50
PAGU 1795	2,015,742	26.89	48	1,310,315	177,963
PAGU 1796	1,951,877	26.86	44	1,292,750	144,056
PAGU 1797	2,005,769	26.98	34	1,235,122	278,257
PAGU 1798	2,012,989	26.85	44	1,422,737	317,217
PAGU 1799	2,087,850	26.82	53	1,421,568	217,125
PAGU 1800	2,009,578	26.90	92	1,342,333	120,039
PAGU 1801	2,004,759	26.99	41	1,461,651	278,257
PAGU 1802	2,005,282	26.98	42	1,595,261	159,188

Table 1: *Fusobacterium watanabei* genomes
Table shows each *Fusobacterium watanabei* isolate name, with the size and the percentage of GC content of each resulting assembly.

Data Availability

Previously deposited draft genome data is available under DRA009376 and DRA009930. Previously deposited partial gene sequences for PAGU 1796 are available under accession numbers NR_179326.1, LC514068.1, LC514067.1, LC514066.1, and LC514065.1. Whole-genome sequences for PAGU 1795, PAGU 1796, PAGU 1797, PAGU 1798, PAGU 1799, PAGU 1800, PAGU 1801, and PAGU 1802 are available in ENA under the study accession number PRJEB87340.

Acknowledgements

Research reported in this publication was supported by the National Institute of Dental and Craniofacial Research of the National Institutes of Health under award number [R01 DE027850] (to C.D.J.), and National Cancer Institute [R01CA288827] (to C.D.J and S.B], as well as start-up funds provided by UT MD Anderson Cancer Center (to S.B and C.D.J).

Methods

Bacterial culturing

Strains PAGU 1795, PAGU 1796, PAGU 1797, PAGU 1798, PAGU 1799, PAGU 1800, PAGU 1801, and PAGU 1802 were grown on anaerobic blood agar plates (Remel[TM], CDC formulation, R01036) at 37 °C degree under anaerobic conditions (Baker Concept 1000; N_2:H_2:CO_2 90:5:5).White-greyish colonies appeared between 24h to 48h which were used for the downstream isolation of genomic DNA.

Genomic DNA isolation

Genomic DNA was extracted using a modified protocol with the Monarch® Spin gDNA Extraction Kit (New England Biolabs, Ipswich, MA, USA). Briefly, the resuspended bacterial pellet and lysis buffer were combined and transferred to an MP Lysing Matrix B tube (MP Biomedicals, Santa Ana, CA, USA), followed by mechanical disruption using the MP Biomedicals Fastprep-24 Bead Beater (MP Biomedicals, Santa Ana, CA, USA) at 4.0 m/s for 20 seconds. Subsequent steps followed the standard procedure outlined in the Monarch® Spin gDNA Extraction Kit protocol.

Illumina DNA Sequencing

Illumina sequencing libraries were prepared using the tagmentation-based and PCR-based Illumina DNA Prep kit (Illumina, San Diego, CA, USA) and custom IDT 10bp unique dual indices (UDI) (Integrated DNA Technologies, Inc., Coralville, Iowa, USA) with a target insert size of 280bp. No additional DNA fragmentation or size selection steps were performed. Illumina sequencing was performed on an Illumina Novaseq X Plus sequencer in one or more multiplexed shared-flow-cell runs, producing 2X151bp paired-end reads. Demultiplexing, quality control and adapter trimming was performed with bcl-convert (v.4.2.4). Short read assembly was performed with Unicycler[8].

Phylogenetic classification in Genome Database Taxonomy (GTDB)

Phylogenetic classifications were assessed using GTDB-tk [2] as listed in Table 2

(https://github.com/Ecogenomics/GTDBTk).

Average Nucleotide Identity

Pairwise average nucleotide identity values were calculated using FastANI [9] via the DOE Systems

Biology Knowledgebase (KBase) [10]. Clustered dendogram generated with pheatmap R package,

RStudio version 2024.12.0+467.

126
References

1   Tomida, J. *et al. Fusobacterium watanabei* sp. nov. As additional species within the genus
Fusobacerium, isolated from human clinical specimens. *Anaerobe* **69**, 102323 (2021).
https://doi.org/10.1016/j.anaerobe.2021.102323
2   Chaumeil, P. A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to
classify genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925-1927
(2019). https://doi.org/10.1093/bioinformatics/btz848
3   Ma, X. *et al.* Pangenomic Study of *Fusobacterium nucleatum* Reveals the Distribution of
Pathogenic Genes and Functional Clusters at the Subspecies and Strain Levels. *Microbiol*
Spectr **11**, e0518422 (2023). https://doi.org/10.1128/spectrum.05184-22
4   Zepeda-Rivera, M. *et al.* A distinct *Fusobacterium nucleatum* clade dominates the
colorectal cancer niche. *Nature* **628**, 424-432 (2024). https://doi.org/10.1038/s41586-024-
07182-w
5   Krieger, M., Guo, M. & Merritt, J. Reexamining the role of *Fusobacterium nucleatum*
subspecies in clinical and experimental studies. *Gut Microbes* **16**, 2415490 (2024).
https://doi.org/10.1080/19490976.2024.2415490
6   Kook, J. K. *et al.* Genome-Based Reclassification of *Fusobacterium nucleatum* Subspecies
at the Species Level. *Curr Microbiol* **74**, 1137-1147 (2017).
https://doi.org/10.1007/s00284-017-1296-9
7   Oren, A. & Garrity, G. M. Valid publication of new names and new combinations
effectively published outside the IJSEM. *Int J Syst Evol Microbiol* **71** (2021).
https://doi.org/10.1099/ijsem.0.004846
8   Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial
genome assemblies from short and long sequencing reads. *PLoS Comput Biol* **13**, e1005595
(2017). https://doi.org/10.1371/journal.pcbi.1005595
9   Jain, C., Rodriguez, R. L., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat*
Commun **9**, 5114 (2018). https://doi.org/10.1038/s41467-018-07641-9

10   Arkin, A. P. *et al.* KBase: The United States Department of Energy Systems Biology
158   Knowledgebase. *Nat Biotechnol* **36**, 566-569 (2018). https://doi.org/10.1038/nbt.4163
159