Complete genome sequences of *Fusobacterium watanabei* sp. isolates

Martha A. Zepeda-Rivera[1], Falk Ponath[2], Kaitlyn N. Lewis[1], Rutika P. Gavate[1], Floyd E. Dewhirst[3,4], Junko Tomida[5], Yoshiaki Kawamura[5], Kaori Tanaka[6], Susan Bullman[2], Christopher D. Johnston[1#]

[1]Genomic Medicine, MD Anderson Cancer Center, Houston, Texas, USA

[2]Immunology, James P. Allison Institute, MD Anderson Cancer Center, Houston, Texas, USA

[3]ADA Forsyth Institute, Cambridge, MA, USA

[4]Department of Oral Medicine, Infection and Immunity, Harvard School of Dental Medicine, Boston, Massachusetts, USA

[5]Department of Microbiology, School of Pharmacy, Aichi Gakuin University, Nagoya, Aichi, Japan

[6]Division of Anaerobe Research, Life Science Research Center, Gifu University, Yanagido, Gifu, Japan

[#]**Address Correspondence to**:

Christopher D. Johnston, PhD, cdjohnston@mdanderson.org

Keywords: Fusobacterium, nomenclature, Illumina sequencing

Abstract

We report the complete genome sequences of eight *Fusobacterium watanabei* clinical isolates, ranging from 1.95 to 2.09 Mbp. Analysis against the Genome Taxonomy Database (GTDB) indicates that *Fusobacterium watanabei* genomes are part of the "*Fusobacterium nucleatum_J*" group, which also encompasses the previously published FNU strain and *Fna* C1 isolates.

Announcement

Fusobacterium watanabei was initially described as a novel species closely related to the *Fusobacterium nucleatum* sensu lato subspecies: *F. nucleatum* subsp. *animalis, F. nucleatum* subsp. *nucleatum, F. nucleatum* subsp. *polymorphum, F. nucleatum* subsp. *vincentii*[1]. Species designation was based on whole genome comparisons of eight *Fusobacterium watanabei* clinical isolates against *Fusobacterium nucleatum* sensu lato subspecies members. Analysis indicated average nucleotide identity (ANI) and digital DNA-DNA hybridization (dDDH) metrics consistent with species-level assignment[1]. Here, we report and release the genome sequences for these eight clinical isolates of *Fusobacterium watanabei* (**Table 1**). Genome assembly indicates a genome size range from 1.95 to 2.09 Mbp (average of 2.01 Mbp) with GC content ranging from 26.82% to 26.99% (average of 26.92%).

Although the 16S rRNA gene and partial sequences of four marker genes for the PAGU 1796 (type strain) were available at National Center for Biotechnology Information (NCBI), the draft genome was only available in the Short Read Archive (DRA009376 and DRA009930). Here, analysis of assembled genomes from the eight *Fusobacterium watanabei* strains originally described by Tomida et. al[1], including PAGU 1796 (type strain), against the Genome Taxonomy Database using GTDB-tk[2] assigned them all to the currently designated "*Fusobacterium*

nucleatum_J" group, a member of the *Fusobacterium nucleatum* sensu lato group of taxa.

"*Fusobacterium nucleatum_J*" also encapsulates the strain 13-08-02, strain "*Fusobacterium*

nucleatum subsp. *unique* (FNU)" previously described in Ma et al.[3], and members of the group we

recently described as "*Fusobacterium nucleatum* subsp. *animalis* clade 1 (*Fna* C1)"[4]. ANI values

between "*Fusobacterium watanabei*", "*FNU*", and "*Fna* C1" genomes range from 98.1228% to

99.9995% (**Figure 1**). This supports a previous observation based on single marker gene analysis

that isolates of "*Fusobacterium watanabei*" and "*Fna* C1" may represent the same *Fusobacterium*

lineage (Dr. Øyvind Kommedal, personal communication, October 2024). Until now, the lack of

an annotated genome for *Fusobacterium watanabei* in publicly available databases limited the

ability of investigators, NCBI, and GTDB to recognize and apply the name "*Fusobacterium*

watanabei" to describe FNU[3] and *Fna* C1[4] genomes. Nevertheless, observations regarding this

lineage are in agreement across publications[1,3,4].

The taxonomy of *Fusobacterium* species has recently undergone significant revision[5].

Based on the proposal by Kook et al.[6] to elevate *Fusobacterium nucleatum* sensu lato subspecies

to the species level assignments, the four historic subspecies have been revised to individual

species: *Fusobacterium animalis, Fusobacterium nucleatum, Fusobacterium polymorphum* and

Fusobacterium vincentii. Within this context of changing nomenclature, it is fortuitous that

Fusobacterium watanabei is already a recognized species name under the List of Prokaryotic

names with Standing in Nomenclature (LPSN)[7] and can now be used for genomes in databases

designated as "*FNU*", "*Fna* C1" and "*F. nucleatum_J*". Here, we have publicly released genomes

for strains PAGU 1795, PAGU 1796, PAGU 1797, PAGU 1798, PAGU 1799, PAGU 1800, PAGU

1801, and PAGU 1802. This increases the number of representative genomes to 34 for the

Fusobacterium watanabei ("*FNU*"/"*Fna* C1"/"*F. nucleatum_J*") group, currently the closest

known phylogenetic clade to the colorectal cancer-associated *Fusobacterium animalis*[4]. Finally, it

is noteworthy that the PAGU 1796 *Fusobacterium watanabei* type strain was previously deposited

to DSMZ under CCUG 74246, with sequencing read data subsequently deposited under

DRA009376 and DRA009930. More recently, a genome for the type strain has been made publicly

available (PRJEB85514, GCA_965119615.1). These sequences and the PAGU 1796 genome

assembly released at ENA under GCA_965217535.1 are technically duplicate sequencing efforts

of the same isolate and not to be considered as distinct strains in future analyses.

 Figure

79 **Figure 1: Average nucleotide identity between "*Fusobacterium watanabei*", "*FNU*", "*Fna***
80 **C1" genomes**
81 Clustered dendrogram depicts the FastANI values between *Fusobacterium watanabei* genomes,
82 including the previously published "*FNU*"[3] (‡) , and "*Fna* C1"[4] (*) strains. Color of each
83 individual box represents the fastANI value, ranging from 98.0% (white) to 100% (dark green).
84

Table

Genome Name	Size (bp)	GC (%)	No. Contigs	No. Reads	N50
PAGU 1795	2,015,742	26.89	48	1,310,315	177,963
PAGU 1796	1,951,877	26.86	44	1,292,750	144,056
PAGU 1797	2,005,769	26.98	34	1,235,122	278,257
PAGU 1798	2,012,989	26.85	44	1,422,737	317,217
PAGU 1799	2,087,850	26.82	53	1,421,568	217,125
PAGU 1800	2,009,578	26.90	92	1,342,333	120,039
PAGU 1801	2,004,759	26.99	41	1,461,651	278,257
PAGU 1802	2,005,282	26.98	42	1,595,261	159,188

Table 1: *Fusobacterium watanabei* genomes
Table shows each *Fusobacterium watanabei* isolate name, with the size and the percentage of GC content of each resulting assembly.

Data Availability

Previously deposited sequencing read data is available under DRA009376 and DRA009930.

Previously deposited partial gene sequences for PAGU 1796 are available under accession numbers NR_179326.1, LC514068.1, LC514067.1, LC514066.1, and LC514065.1. A more recent genome sequence for the PAGU 1796 type strain is available under PRJEB85514 and GCA_965119615.1. Whole-genome assemblies for PAGU 1795, PAGU 1796, PAGU 1797, PAGU 1798, PAGU 1799, PAGU 1800, PAGU 1801, and PAGU 1802 are available in ENA under the study number PRJEB87340 with the following sample accessions: GCA_965217525 (PAGU 1795), GCA_965217535 (PAGU 1796), GCA_965217465 (PAGU 1797), GCA_965217495 (PAGU 1798), GCA_965217515 (PAGU 1799), GCA_965217485 (PAGU 1800), GCA_965217505 (PAGU 1801), and GCA_965217475 (PAGU 1802).

Acknowledgements

Research reported in this publication was supported by the National Institute of Dental and Craniofacial Research of the National Institutes of Health under award number [R01 DE027850]

106   (to C.D.J.), and National Cancer Institute [R01CA288827] (to C.D.J and S.B], as well as start-up

107   funds provided by UT MD Anderson Cancer Center (to S.B and C.D.J).

Methods

Bacterial culturing

Strains PAGU 1795, PAGU 1796, PAGU 1797, PAGU 1798, PAGU 1799, PAGU 1800, PAGU 1801, and PAGU 1802 were grown on anaerobic blood agar plates (Remel^{TM}, CDC formulation, R01036) at 37 °C degree under anaerobic conditions (Baker Concept 1000; $N_2:H_2:CO_2$ 90:5:5).White-greyish colonies appeared between 24h to 48h which were used for the downstream isolation of genomic DNA.

Genomic DNA isolation

Genomic DNA was extracted using a modified protocol with the Monarch® Spin gDNA Extraction Kit (New England Biolabs, Ipswich, MA, USA). Briefly, the resuspended bacterial pellet and lysis buffer were combined and transferred to an MP Lysing Matrix B tube (MP Biomedicals, Santa Ana, CA, USA), followed by mechanical disruption using the MP Biomedicals Fastprep-24 Bead Beater (MP Biomedicals, Santa Ana, CA, USA) at 4.0 m/s for 20 seconds. Subsequent steps followed the standard procedure outlined in the Monarch® Spin gDNA Extraction Kit protocol.

Illumina DNA Sequencing

Illumina sequencing libraries were prepared using the tagmentation-based and PCR-based Illumina DNA Prep kit (Illumina, San Diego, CA, USA) and custom IDT 10bp unique dual indices (UDI) (Integrated DNA Technologies, Inc., Coralville, Iowa, USA) with a target insert size of 280bp. No additional DNA fragmentation or size selection steps were performed. Illumina sequencing was performed on an Illumina Novaseq X Plus sequencer in one or more multiplexed shared-flow-cell runs, producing 2X151bp paired-end reads. Demultiplexing, quality control and adapter trimming was performed with bcl-convert (v.4.2.4). Short read assembly was performed with Unicycler[8].

Phylogenetic classification in Genome Database Taxonomy (GTDB)

Phylogenetic classifications were assessed using GTDB-tk [2] as listed in Table 2

(https://github.com/Ecogenomics/GTDBTk).

Average Nucleotide Identity

Pairwise average nucleotide identity values were calculated using FastANI [9] via the DOE Systems

Biology Knowledgebase (KBase) [10]. Clustered dendogram generated with pheatmap R package,

RStudio version 2024.12.0+467.

139
140   **References**
141
References

1   Tomida, J. *et al.* Fusobacterium watanabei sp. nov. As additional species within the
genus Fusobacerium, isolated from human clinical specimens. *Anaerobe* **69**, 102323
(2021). https://doi.org/10.1016/j.anaerobe.2021.102323
2   Chaumeil, P. A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to
classify genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925-
1927 (2019). https://doi.org/10.1093/bioinformatics/btz848
3   Ma, X. *et al.* Pangenomic Study of *Fusobacterium nucleatum* Reveals the Distribution
of Pathogenic Genes and Functional Clusters at the Subspecies and Strain Levels.
Microbiol Spectr **11**, e0518422 (2023). https://doi.org/10.1128/spectrum.05184-22
4   Zepeda-Rivera, M. *et al.* A distinct *Fusobacterium nucleatum* clade dominates the
colorectal   cancer   niche.   *Nature*   **628**,   424-432   (2024).
https://doi.org/10.1038/s41586-024-07182-w
5   Krieger, M., Guo, M. & Merritt, J. Reexamining the role of *Fusobacterium nucleatum*
subspecies in clinical and experimental studies. *Gut Microbes* **16**, 2415490 (2024).
https://doi.org/10.1080/19490976.2024.2415490
6   Kook, J. K. *et al.* Genome-Based Reclassification of *Fusobacterium nucleatum*
Subspecies   at   the   Species   Level.   *Curr   Microbiol* **74**,   1137-1147   (2017).
https://doi.org/10.1007/s00284-017-1296-9
7   Oren, A. & Garrity, G. M. Valid publication of new names and new combinations
effectively   published   outside   the   IJSEM. *Int J Syst Evol Microbiol* **71** (2021).
https://doi.org/10.1099/ijsem.0.004846
8   Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial
genome assemblies from short and long sequencing reads. *PLoS Comput Biol* **13**,
e1005595 (2017). https://doi.org/10.1371/journal.pcbi.1005595
9   Jain, C., Rodriguez, R. L., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species

boundaries. *Nat Commun* **9**, 5114 (2018). https://doi.org/10.1038/s41467-018-
07641-9
10 Arkin, A. P. *et al*. KBase: The United States Department of Energy Systems Biology
Knowledgebase. *Nat Biotechnol* **36**, 566-569 (2018).
https://doi.org/10.1038/nbt.4163
174