Title:

A novel allele of *Mitogen Activated Protein Kinase 6* is linked to disease resistance and
constitutive immunity in rice.

Authors:

Gokulan C.G.[1,6,†], Mohammed Jamaloddin[1,†], Deepti Rao[1], Namami Gaur[1,2], Amita Rani Supriyo[1],
Nisha Sao[1], Deepak Niranjan[1,7], Shrish Tiwari[1], Gouri S. Laha[3], Kalyani M. Barbadikar[3], Raman
Meenakshi Sundaram[3], Sheshu Madhav Maganti[3,4,*], Hitendra K. Patel[1,5,*], Ramesh V. Sonti[1,7,*]

Affiliations:

[1]CSIR-Centre for Cellular and Molecular Biology, Telangana, India.

[3]ICAR-Indian Institute of Rice Research, Telangana, India.

[5]Academy of Scientific and Industrial Research, Uttar Pradesh, India.

[†]These authors should be considered joint first author.

[*]Addresses for correspondence: Sheshu Madhav Maganti - sheshu24@gmail.com; Hitendra K.
Patel - hkpatel@ccmb.res.in; Ramesh V. Sonti – rameshvsonti@gmail.com

Present addresses:

[2]The James Hutton Institute, University of Dundee, Scotland.

[4]ICAR-Central Tobacco Research Institute, Andhra Pradesh, India.

[6]Centre for DNA Fingerprinting and Diagnostics, Telangana, India.

[7]International Centre for Genetic Engineering and Biotechnology, New Delhi, India.

Word count: 6614

SUMMARY

Bacterial Blight (BB) is the most serious bacterial disease of rice. Usage of resistant cultivars has been a successful strategy in controlling BB. However, constant selection pressure causes the evolution of hypervirulent pathogenic strains that break down resistance, thus necessitating additional sources of resistance. An induced-mutant line derived from the elite rice cultivar Samba Mahsuri was identified to exhibit broad-spectrum resistance to BB. Using next-generation gene mapping approaches, we identified a genomic interval in chromosome 6 to be linked to BB resistance. Analysis of the SNPs in the locus and subsequent linkage analysis indicated that a missense SNP located in the second exon of the *Mitogen-Activated Protein Kinase 6* (*OsMAPK6*) could be the causal mutation. A detailed examination of *OsMAPK6* gene sequences in over 4000 rice genomes revealed that the mutant allele identified in this study is novel to rice germplasm. The candidate SNP causes the substitution of an invariant Serine residue with a Proline residue (S84P), thus likely affecting the protein function. Global transcriptome, metabolome, biochemical, and molecular analyses suggested that BB42 exhibit constitutive immunity. On the mechanistic front, the mutation seems to result in diminished brassinosteroid signaling in BB42. Taken together, we report a novel allele of a highly conserved *MAPK* gene and provide evidence for its possible association with constitutive immunity and disease resistance in rice.

KEYWORDS: Bacterial blight, Disease resistance, Gene mapping, MAP kinase, Plant immunity, Rice, *Xanthomonas*.

INTRODUCTION

Plants possess a potent innate immunity which every cell that faces a threat can elicit. The two-tiered innate immunity of plants consists of Pattern-Triggered Immunity (PTI) and Effector-Triggered Immunity (ETI) (Jones and Dangl, 2006). PTI is elicited by the recognition of the Pathogen-Associated Molecular Patterns (PAMPs) or Damage-Associated Molecular Patterns (DAMPs). PAMPs include conserved microbial-derived molecules like flagellin, peptidoglycan, LPS, and chitin; DAMPs include damage-associated molecules like extracellular ATP, extracellular DNA, and oligogalacturonides (Chen and Ronald, 2011; Pontiggia *et al.*, 2020). The patterns are perceived by the membrane-bound receptor kinases, that are collectively called as Pattern Recognition Receptors (PRRs), that relay signals to the signaling proteins like the MITOGEN-ACTIVATED PROTEIN KINASE (MAPK), that typically transduce the signals to transcription factors. PTI induces an extensive transcriptional and translational reprogramming that results in the production of defense-related proteins, phytohormones, and metabolites, thus limiting pathogen entry and proliferation (D Couto, 2016; Yuan *et al.*, 2021; Zipfel and Felix, 2005). On the other hand, ETI is a stronger defense response that is activated by the recognition of pathogen-secreted proteins (also known as effectors) by intracellular receptors mainly belonging to the Nucleotide-Binding and Leucine-rich Repeat containing Receptors (NB-LRRs/NLRs) superfamily. ETI is manifested by the Hypersensitive Response (HR) which results in a localized cell-death, thus restricting the spread of pathogens (Jones *et al.*, 2016). On the other hand, uncontrolled regulation of various arms of the plant immune system results in constitutive immunity which in turn results in various traits like stunted growth, spontaneous lesions, and disease resistance among others (Freh *et al.*, 2022).

Bacterial Blight (BB) is the most serious bacterial disease of rice that can result in up to 70% yield loss depending on the infection severity (Goto, 1992; K. Reddy, 1979; Ou, 1973, 1985). Rice cultivation in India is affected by BB with ~8.5% loss in annual yield (Savary *et al.*, 2019). BB is caused by the Gram-negative bacterium *Xanthomonas oryzae* pv. *oryzae* (*Xoo*). *Xoo* invades rice leaves through hydathodes and colonizes the extracellular spaces in xylem parenchyma tissue (Hilaire *et al.*, 2007). Under high humidity conditions, the bacteria ooze out of the openings such as hydathodes or wounds and form beads or strands of exudates that potentially act as inoculum for further infections (Mew, 1993). Being a prolific colonizer of the vascular tissue, *Xoo* spreads systemically along the veins and results initially in chlorotic yellow and later in necrotic lesions

on the infected leaves, ultimately leading to drying of the leaves (Joshi *et al.*, 2020; Niño-Liu *et*
al., 2006)**.**

Previous studies have associated several genes and their alleles with BB resistance, of
which a few alleles are widely used in breeding programme to develop BB resistant rice varieties
(Fiyaz *et al.*, 2022; Jamaloddin *et al.*, 2021; Jiang *et al.*, 2020). Such resistant varieties have been
deployed in the field with ample success (Dasari *et al.*, 2022; Huang *et al.*, 1997; Jamaloddin *et*
al., 2020; Oliva *et al.*, 2019; Pradhan *et al.*, 2015; Sundaram *et al.*, 2008). It was observed that a
combination of two to three *R*-genes was effective against majority of the *Xoo* pathotypes.
However, a constant selection pressure on pathogens combined with their rapid generation time is
expected to result in the evolution of adapted pathotypes that breakdown existing sources of
disease resistance. Studies have identified multiple strains of *Xoo* that breakdown individual
resistance genes (Kaur *et al.*, 2020; Mishra *et al.*, 2013; Quibod *et al.*, 2020; Yugander *et al.*, 2017,
2022). Adding another layer to this serious issue is the migration of hypervirulent strains from one
country to another (Carpenter *et al.*, 2020). Despite the existence of several *R*-genes, the practical
deployment of the genes needs to be rigorously tested on a case-to-case basis to ensure the
compatibility of the genes to different agroclimatic zones. For instance, although *Xa7* was
identified to be a broad-spectrum R-gene, a study in India found that 94% of the 71 strains tested
were virulent on a *Xa7*-positive rice line (Yugander *et al.*, 2022). Therefore, to counter resistance
gene breakdown and to curb disease outbreaks, it is essential to identify new sources of broad-
spectrum resistance that are suitable for given agroclimatic conditions and the cultivar background.

Mutation breeding is an efficient method to obtain novel sources of resistance in relatively
less time, while preserving the genetic architecture of the desired cultivar. Further, the availability
of advanced gene mapping approaches enables forward genetics using populations derived from
the induced mutant lines (Abe *et al.*, 2012; Takagi *et al.*, 2013). Earlier, we reported the generation
and screening of a large-scale ethyl methanesuphonate-induced mutant lines of the rice mega-
variety Samba Mahsuri for various traits of economic importance (Potupureddi *et al.*, 2021). Here,
we report the characterization of one of the mutant lines (henceforth BB42) that exhibited a broad-
spectrum resistance to various hypervirulent isolates of *Xoo*. Through a genetic mapping study,
we observed that resistance in BB42 is controlled by a single recessive locus. We further identified
a SNP located in the second exon of *OsMAPK6* gene that causes Serine to Proline substitution at
the 84[th] position and cosegregates with the resistance trait. Notably, the mutant allele identified

here represents a novel genetic variation in the rice germplasm. Transcriptome profiling in
combination with multiple biochemical and molecular assays revealed a constitutively active basal
defense signaling in BB42 when compared to SM. Particularly, several kinase-encoding genes
were significantly upregulated, and several members of the major transcription factor families
were differentially expressed. Overall, our study reports a novel allele of *OsMAPK6* that is linked
to bacterial blight resistance, and constitutive immunity and presents a potential genetic source for
breeding BB resistance in rice.

RESULTS

BB42 exhibits broad-spectrum resistance to highly virulent *Xoo* isolates

Earlier, we identified BB42 from an EMS-induced mutant library of SM as a line exhibiting
enhanced resistance to the bacterial blight disease of rice (Potupureddi *et al.*, 2021). Here, we
characterized BB42 further through a combination of phenotyping, genomic, molecular,
transcriptomics and metabolomics approaches. Morphologically, BB42 exhibited reduced height
compared to SM (Fig. 1A, B). Further, to understand the spectrum of resistance, we challenged
BB42 with several hypervirulent strains of *Xoo* that were shown to breakdown resistance provided
by major resistance genes including *xa5*, *xa13* and *Xa21* (Mishra *et al.*, 2013). The results revealed
that BB42 exhibits enhanced resistance to all the tested *Xoo* strains (Fig. 1C, D). Notably,
challenging SM and BB42 with *Xanthomonas oryzae* pv. *oryzicola* (*Xoc*) – a pathovar of *X. oryzae*
that causes bacterial leaf streak (BLS) in rice - showed no difference in the susceptibility.
Intriguingly, the susceptibility level of SM to *Xoc* was significantly lesser than that of Nipponbare
- another BLS susceptible rice variety (Figure S1A). Taken together, these results suggest that
BB42, a semi-dwarf mutant of SM, is resistant to *Xoo* in what appears to be a non-race-specific
manner.

A missense mutation in *OsMAPK6* is linked to bacterial blight resistance

We next aimed to identify the genetic basis of bacterial blight resistance in BB42. To this
end, we generated two F$_2$ populations, each with 244 plants, and screened them with a virulent
Xoo strain, IXO1221. Phenotype segregation in both the populations followed the Mendelian ratio
for a monogenic recessive trait (Table S4; Figure S2A, B). Progenies from both the populations
were used for gene mapping through the QTL-sequencing pipeline (Takagi *et al.*, 2013). The
analysis revealed a genomic interval in the short arm of chromosome 6, in both the populations, to

be linked to resistance (Fig. 2A, B). We identified five Single Nucleotide Polymorphisms (SNPs) within the interval, of which, one SNP was annotated as a 'missense and splice region' variant and was predicted to be deleterious (Table 1). This SNP (T → C at SM-Chr6:2664062) is located in the second exon of the rice *Mitogen Activated Protein Kinase 6* (*OsMAPK6*) gene and results in the substitution of a Serine by Proline (Fig. 2C). Analysis of the RNA reads mapped to OsMAPK6 indicated no anomalies in the splicing of the gene between SM and BB42 (Figure S3). Multiple sequence alignment of one-hundred MAPK6 homologues across different plant species indicated a 100% conservation of the affected Serine residue (Fig. 2D). Notably, when searched in the RiceVarMap V2.0 database (Zhao *et al.*, 2021) which contains sequences of about 4500 rice accessions, we found no accession to carry any variation at this position. To further ascertain the co-segregation of the observed SNP with BB resistance, we developed PCR-based markers that differentiate the SM and the BB42 alleles (Figure S2C). We tested randomly selected F_2 progenies that were resistant or susceptible to *Xoo* IXO1221 and found that the BB42 allele of *OsMAPK6* co-segregated with the resistant progenies, whereas the susceptible progenies were homozygous for SM allele or were heterozygous (Fig. 2E). Overall, these results suggest that the variation in the *OsMAPK6* gene likely confers BB resistance. The identified allele is novel to rice germplasm and the affected Serine residue is evidently invariable in MAPK6 homologues that were assessed, indicating a likely role of this novel variant in BB resistance.

Xoo **induces an overlapping transcriptome change in SM and BB42**

Given the central role of MAPK6 in orchestrating cellular signalling during defense and development, we reasoned that BB42 and SM are likely to have different transcriptional profiles which might explain the observed resistance to *Xoo*. Therefore, we performed a global transcriptome profiling of leaves from SM and BB42 that were sampled immediately post infection (Mock) or at 24 hours post infection with *Xoo* IXO1221 (*Xoo*). The samples were analysed to identify the treatment-based transcriptome changes in both the genotypes (i.e., *Xoo* vs Mock in SM and BB42) as well as genotype-specific transcriptome differences in both the treatments (i.e., BB42 vs SM in Mock and *Xoo*). Four combinations of samples were analysed in total and the results indicated that *Xoo* treatment caused a major transcriptome reprogramming in both the genotypes (1974 DEGs in SM and 1477 DEGs in BB42) (Data S1). On the other hand, the genotype-specific differences in the transcriptome were relatively smaller (810 DEGs in Mock

treatment and 493 DEGs in *Xoo* treatment) (Fig. 3A; Figure S4; Table S5). Further comparative

analyses revealed that about 60% of the upregulated genes in SM and BB42 upon *Xoo* infection

were common (Fig. 3B). With regards to the downregulated genes, about 87% of the genes

downregulated in BB42 were also downregulated in SM upon *Xoo* infection (Fig. 3B), while

several additional genes were uniquely downregulated in SM. This is suggestive of a largely

similar *Xoo*-mediated transcriptional landscape in SM and BB42 at gene level, which is also

reflected at pathway levels (Figure S5).

We then performed a *k*-means clustering analysis of the top 2000 genes exhibiting highly

variable expression across different samples and treatments. We observed 1628 genes showing

similar expression trend in both SM and BB42, whereas 372 genes showed an increased expression

in BB42 irrespective of the treatment, when compared to SM (Fig. 3C; Cluster B). Intrigued by

this observation, we did a cluster-wise GO term enrichment analysis and noted that cluster B was

enriched with genes that are involved in defense response and cell signalling. It is mention-worthy

that the higher expression of defense genes is observed even in control BB42 plants (Fig. 3C),

which points towards the likelihood of a constitutively active defense in BB42. Therefore, we

probed deeper to understand the nature of constitutive defense in BB42 using DEGs from the

mock-treated samples.

BB42 exhibits constitutive immunity

We considered the DEGs between mock-treated BB42 and SM for further analysis to

understand the basal genotype-specific differences. A GO term enrichment analysis revealed

higher expression levels of genes involved in defense response, receptor signalling, and protein

phosphorylation (Fig. 4A). Conversely, certain genes involved in wounding response, response to

jasmonic acid, and regulation of transcription were expressed at lower levels in BB42 in

comparison to SM (Fig. 4A). We first looked into the genes present in the highly expressed

categories and noted the presence of various classes of defense related genes (n=41) encoding

receptor kinases, receptor-like cytoplasmic kinases, a catalase, pathogenesis-related proteins and

a JA biosynthetic protein (Table S6). Very importantly, amongst the above set of genes were

fourteen genes encoding Nucleotide-Binding adaptor shared by APAF-1, R proteins, and CED-4

(NB-ARC) domain containing proteins and Nucleotide-Binding Leucine-rich Repeat containing

Receptor proteins (NB-LRR or NLRs), which is one of the characteristic traits observed in

autoimmune mutants in plants (Freh *et al.*, 2022) (Figure S6). While this suggests an enhanced

defense signalling in BB42, we were puzzled to observe downregulation of JA responsive genes.

However, a detailed inspection indicated that such genes encoded the JASMONATE-ZIM

DOMAIN (JAZ or TIFY) proteins, which act as repressors of JA signalling (Table S7). This further

suggests the possibility of an active and/or enhanced JA signalling in BB42. To validate this

observation and also to check if this scenario persists through different developmental stages, we

performed quantitative PCR using defense marker genes (PR genes) in SM and BB42 seedlings.

In accordance with the previous observation, we found significant upregulation of defense marker

genes in BB42 seedlings (Figure S76A). Moreover, we found other hallmarks of defense activation

like H_2O_2 accumulation and MAPK activation to be enhanced in BB42 even in unelicited

conditions (Figure S7B, C). Further, as we observed an upregulation of various types of kinases,

we used Phospho-Serine and Phospho-Threonine/Tyrosine antibodies to assess the qualitative

differences in the phosphorylated proteins in SM and BB42. There was an observable difference

in the signal patterns in all forms of phosphorylated proteins between SM and BB42 (Figure S7D).

By and large, the transcriptomics data in combination with other assays and the dwarf stature of

BB42 are indicative of constitutive immunity and a differential phosphorylation-mediated

signalling between the two lines.

We performed an untargeted metabolite profiling of SM and BB42 leaves at the tillering

stage and observed a general difference in the metabolite profiles of SM and BB42 (Figure S8A).

Among the detected metabolites, 92 metabolites were differentially accumulated (DAMs), of

which 40 were relatively abundant in BB42 and 52 were relatively abundant in SM (Figure S8B;

Data S2). Enrichment analysis of the DAMs indicated linoleic acid metabolism, vitamin B6

metabolism, and nicotinate metabolism to be upregulated and starch and sucrose metabolism, and

glutathione metabolism to be downregulated in BB42 (Figure S9). On that note, levels of some of

the compounds associated with the above pathways showed clear difference between SM and

BB42. Notably, compounds that are associated with defense like linoleic acid and coniferaldehyde

were abundant in BB42, while compounds that were associated with disease susceptibility like

sucrose, *trans*-Zeatin, and amino acids were abundant in SM (Fig. 4B).

Brassinosteroid signalling is affected in BB42

Earlier, Liu *et al.*, (2015) identified an allele of *OsMAPK6* that resulted in dwarf stature and small grain size (*dsg1*) and showed that the observed traits were due to imbalance in brassinosteroid (BR) homeostasis. Interestingly, analysis of various morphological traits showed slight similarity between *dsg1* and BB42. For instance, BB42 grains were slightly yet significantly shorter than those of SM, but there were no difference in grain width (Fig. 5A, B, C). We then tested the BR sensitivity of SM and BB42 plants by measuring coleoptile elongation following treatment with 24-epibrassinolide (epiBL). The elongation of BB42 coleoptile in response to epiBL was not as significant as that of SM, suggesting a reduced sensitivity to BR in BB42 (Fig. 5D). Intriguingly, brassinolide was detected in the metabolome profiling and its level was slightly, albeit not significantly, higher in BB42 than in SM (Fig. 5E). It is possible that the reduced sensitivity to BR in BB42 might have negatively impacted the feedback inhibition of BR biosynthesis, thus resulting in higher basal BR levels. Additionally, expression of several genes involved in BR signalling and response was diminished in BB42 as compared to SM (Table S8). On the whole, it seems like BB42 is impaired in BR sensitivity and a role of the mutated OsMAPK6 in this impairment could be speculated.

DISCUSSION

In this study, we report an EMS-induced allele of *Mitogen-Activated Protein Kinase 6* and show its involvement in resistance to bacterial blight, which is a serious disease of rice. The variant of OsMAPK6 identified here is new to rice germplasm wherein an otherwise invariant Serine at the 84[th] position is replaced by Proline. Not just in plants, the affected Serine was observed to be invariant in MAPK6 homologues of different kingdoms (Figure S10A, B). Further, the serine residue is conserved in ten of the seventeen rice MAPKs (Figure S11A, B). These results signify the importance of the Serine residue for the function of MAPK6 homologues and perhaps in majority of the rice MAPKs as well. The causal SNP is a T to C transition, a non-canonical type of EMS-induced variation. EMS is believed to predominantly cause G/C to A/T type transitions, although exceptions are reported (Abe *et al.*, 2012; Greene *et al.*, 2003; Jhingan *et al.*, 2023; Loveless, 1969; Till *et al.*, 2007).

Effect of the variation on MAPK6 function

MAPK6 homologues are highly conserved across eukaryotes and play vital roles in disease and development (Meng and Zhang, 2013; Sun and Zhang, 2022). Previous studies have associated OsMAPK6 with various growth processes (Liu *et al.*, 2015; Xu *et al.*, 2018; Yi *et al.*, 2016) as well as defense response and resistance to diseases like blast, blight and bacterial leaf streak (Chen *et al.*, 2021; Chujo *et al.*, 2014; Kim *et al.*, 2012; Kishi-Kaboshi *et al.*, 2010; Lieberherr *et al.*, 2005; Xu *et al.*, 2018). It is interesting to note that natural variations in *OsMAPK6* differentially affect the protein activity and resistance to *Xoc* (Ma *et al.*, 2021). In line with this, the observed difference between SM and Nipponbare in their response to *Xoc* (Figure S1A) might be explained by the natural variation in the N-terminus of MAPK6 between SM and Nipponbare (Figure S1B). Further, certain loss-of-function alleles of *OsMAPK6* were reported to cause embryo lethality (Ma *et al.*, 2021; Yi *et al.*, 2016). It was also observed that a heterozygously inherited loss-of-function mutation in OsMAPK6 could confer moderate resistance to *Xoo* and *Xoc* (Ma *et al.*, 2021). Hence, the effect of the S84P mutation on OsMAPK6 function becomes a primary question. The fact that the activation of OsMAPK6 is not affected in BB42 (Figure S6C) suggests that the mutation has not abolished the protein activation and function, rather the protein's function could have been altered. Intriguingly, one of resistant progenies showed a heterozygous genotype in our analysis. Noteworthy, the SNP-index of this variation is not 1.0 in either of the population. This might indicate an effect of the environment, that could have resulted in one or a few heterozygous individuals showing resistance trait getting included in our sequencing and downstream analyses. Nevertheless, none of the susceptible individuals tested were homozygous for the mutant allele and all of the lines that were homozygous for the mutant allele were resistant. On balance, it is possible that the presence of the mutant allele and allele dosage in combination with the environment might contribute to BB resistance.

MAPK6 integrity could be crucial to prevent constitutive immunity in rice

The phenotypes observed in BB42 including dwarf stature and constitutive immunity, seem to partly resemble a defective MAPK cascade reported in the model plant *Arabidopsis thaliana*. In Arabidopsis, three major MAPKs that are involved in plant immunity and pathogen response are AtMPK3, AtMPK4, and AtMPK6 (Sun and Zhang, 2022). Several studies have elucidated the signalling cascade involving AtMPK4 and showed that the MEKK1-MKK1/2/6-MPK4 cascade is

involved in the negative regulation of innate immunity (Gao *et al.*, 2008; Lian *et al.*, 2018;

Rodriguez *et al.*, 2010). The mutants *mekk1, mkk1/2,* and *mpk4* exhibit stunted growth, severe

dwarfism, and enhanced basal defense responses. Subsequent research showed that the MPK4

cascade is guarded by NLR proteins SUMM2 and RPS6, which get activated when the cascade is

disrupted (Lian *et al.*, 2018; Zhang *et al.*, 2012, 2017). In this context, it is possible that S84P

mutation in OsMAPK6 may modify its interactome, which in turn might activate the immune

components guarding it.

As activation of MAPKs is critical for proper elicitation of plant immunity, the observation

of BB42 exhibiting constitutive defense traits which includes MAPK6 activation raises the

following question: Is OsMAPK6 activation in unelicited BB42 a cause or consequence of the

active defense? A remarkable phenomenon observed in BB42 was the basal upregulation of several

NLR-encoding genes, which appears to phenocopy the Arabidopsis plants that express a

constitutively active form of MPK3 (CA-MPK3). Studies have revealed that expression of CA-

MPK3 causes enhanced expression of NLR encoding genes in a salicylic acid-dependent manner

(Genot *et al.*, 2017; Lang *et al.*, 2017, 2022). Expression of CA-MPK3 also resulted in resistance

to the bacterial pathogen *Pseudomonas syringae* (Lang *et al.*, 2017). Another study showed that

long-lasting activation of MPK3/MPK6 is necessary for ETI-mediated reduction in photosynthetic

efficiency in Arabidopsis (Su *et al.*, 2018). In line with this, we observed a reduction in Chlorophyll

A in BB42 as compared to SM (Fig. 4B). Further, prolonged activation of MPK3/MPK6 was

observed during ETI and such activation was critical for the regulation of SA-responsive genes

(Tsuda *et al.*, 2013). This study suggested that the activation duration of MAPK is a determinant

of the robustness of immunity. Of note, the consequence of CA-MAPK expression is dependent

on the MAPK in question. For instance, overexpression of CA-MPK4 dampened immune

responses and abolished pathogen resistance in Arabidopsis (Berriri *et al.*, 2012). Considering the

above-discussed points, it is possible that the activation status of the OsMAPK6-S84P variant and

NLR expression could be correlated. Nonetheless, the directionality of these two observations is

not certain yet. Also, unlike SA in Arabidopsis, our data suggests a possible involvement of JA in

the execution of MAPK6-mediated defense responses in rice. Future investigations are needed to

draw a firm conclusion.

It appears that BB42 is a unique mutant which possesses most hallmarks of autoimmunity

except the spontaneous lesions that occur due to localised cell death. In rice, almost all the reported

autoimmune mutants display spontaneous lesions on the leaves, thereby earning the name lesion mimic mutants (LMMs); some LMMs show reduced photosynthetic capacity as well (Freh *et al.*, 2022; Xiaobo *et al.*, 2020). Interestingly, loss of function mutations in components of MAPK signalling including OsMKK6 and OsMAPK4 have been reported to cause spontaneous lesions and autoimmunity in rice (Jiang *et al.*, 2023; Wang *et al.*, 2021). Although the molecular mechanisms of autoimmunity in rice is still unclear, based on the data presented here, we propose that a module involving MAPK6, NLRs, and JA signalling proteins might govern autoimmunity.

Potential link between MAPK6, BR, and disease resistance

The relationship between OsMAPK6 function and brassinosteroid homeostasis has been established earlier (Liu *et al.*, 2015) and has also been observed in our study (Fig. 5). However, whether the perturbed BR homeostasis has any direct effect on disease resistance remains to be explored. Involvement of BRs in plant-pathogen interaction as well as growth and development has been demonstrated earlier. BRs can act both synergistically and antagonistically with the immune components to regulate growth and defense (Albrecht *et al.*, 2012; Belkhadir *et al.*, 2012; Heese *et al.*, 2007; Liao *et al.*, 2020). In rice, BRs were shown to suppress rice immunity against a root pathogen (De Vleesschauwer *et al.*, 2012). Another study showed exogenous treatment of rice leaves with brassinolide enhanced resistance to *Xoo* (Nakashita *et al.*, 2003). The *Xanthomonas* effector Xoo2875 was shown to suppress immunity by interacting with OsBAK1, a component of PTI and BR signalling (Yamaguchi *et al.*, 2013). In Arabidopsis, BRs were shown to antagonize JA-mediated plant defense (Liao *et al.*, 2020). Strikingly, several components of BR signalling were downregulated whereas JA signalling was positively regulated in BB42 (Tables S6, S7, S8; Fig. 4A). Whether and how OsMAPK6-S84P affect BR homeostasis and its relation with disease resistance warrants further investigations.

Our study leads to several questions such as: Is the novel MAPK6 variant constitutively active? Does the variation affect the interactome of OsMAPK6? Are rice MAPK cascades guarded by NLRs? Is rice NLR expression controlled by OsMAPK6-mediated signalling? How could OsMAPK6 govern BR homeostasis, and what is the role of BR signalling in rice-*Xoo* interaction? Do rice pathogens directly target the components of MAPK cascade to promote virulence? BB42 can also serve as a novel source of BB resistance as initial observations suggest that there is no drastic effect on yield. However, further studies are needed to determine if BB42 has more subtle

reductions in yield as compared to the parental line. More research directed towards the above
questions would provide deeper insights into rice MAPK signalling, broaden our understanding of
plant immunity in crop species and also indicate whether BB42 can be a source of resistance that
can be used for rice improvement.

EXPERIMENTAL PROCEDURES

Plant materials and bacterial strains

Plant materials used for this study include the wildtype/parental cultivar Samba Mahsuri
(SM; BPT5204) and the EMS-induced mutant line named BB42. Nipponbare rice cultivar was
used as a susceptibility check for bacterial leaf streak disease scoring. *Xanthomonas oryzae* pv.
oryzae (*Xoo*) and *X. oryzae* pv. *oryzicola* (*Xoc*) isolates were cultured from the lab stocks and the
stocks obtained from Indian Institute of Rice Research, Hyderabad. The bacterial strains that were
used are listed in Table S1. The seeds were germinated on Petri plates at 28°C in darkness for 72
hours and under 12-hour photoperiod from days 4 to 7. Later the seedlings were potted onto soil
before transplanting to the field after 14-21 days of potting and grown in greenhouse conditions
under natural lighting.

Plant inoculation

Plants in the maximum tillering stage (50-60 days) were inoculated with *Xoo* isolates using
the clip inoculation method (Kauffman *et al.*, 1973). The *Xoo* isolates were grown in Peptone-
Sucrose (PS) broth (1% (w/v) peptone and 1% (w/v) sucrose) for 24 hours and the cells were
pelleted. The cells were further washed in sterile Milli-Q water one to two times and adjusted to
an optical density at 600nm (OD_{600}) of 1.0. Sterile scissors were used to dip into the bacterial
suspension and clip the fully expanded leaves ~3-4 cm from the leaf tip. The disease severity was
scored by measuring the lesion length from the site of inoculation after 14 days of inoculation. The
scoring was performed following the Rice Standard Evaluation System (IRRI, 2014). Inoculations
were done in green house with temperature maintained under 30°C for transcriptome analysis
studies and multiple-isolate screening. The parental lines and F_2 progenies were screened in the
rice field facility of CSIR-CCMB (17.41° N, 78.55° E) during Kharif 2020 with the *Xoo* strain
IXO1221. *Xoc* inoculation was performed in greenhouse conditions on 2-3 weeks old seedlings
through syringe infiltration of *Xoc* suspension at OD_{600} = 1.0. Lesion lengths were measured at 5
days post inoculation.

Generation of mapping populations

Two mapping populations were generated in this study. The populations were obtained by using SM as recipient and BB42 as donor (SM x BB42) and vice versa (BB42 x SM). The F_1 plants, so generated, were tested for hybridity by Sanger sequencing of known SNPs between SM and BB42, which were identified using whole genome sequencing data. The confirmed F_1 plants were let to self-fertilize to obtain two independent segregating populations. F_2 populations with 244 progenies per population were screened for BB resistance as described above.

Bulk sequencing and analysis

F_2 plants from both the populations showing the extreme phenotypes i.e., resistance and susceptibility to BB infection were selected to constitute the resistant and susceptible bulks, respectively. An equal amount of leaves (15mg per leaf) were pooled from F_2 plants showing identical phenotype for DNA isolation. DNA from the bulked leaves as well as the parental leaves were isolated using the CTAB method and quantified using Qubit^{TM} High-Sensitivity dsDNA Assay in a Qubit^{TM} 4 fluorometer (Thermo Fisher Scientific, USA). The quality of the DNA was assessed on 0.8% agarose gel through electrophoresis. An amount of 1µg DNA from all the samples was used to prepare sequencing libraries with TruSeq Nano DNA HT Sample Preparation Kit (Illumina, USA) and qualified libraries were sequenced on Illumina NovaSeq 6000 platform to obtain 2x150 bp reads at approximately 40X coverage. The library preparation and sequencing were performed at the NGS facility of CSIR-CCMB. The sequenced reads were quality-checked using *FastQC* and were provided as input for QTLseq pipeline (Sugihara *et al.*, 2020) to identify the causal loci. SM reference genome (assembled in-house) was used for the gene mapping analysis (Rao *et al.*, *in preparation*). *AUGUSTUS* in the *Galaxy* platform was used for annotating the interval of interest (Afgan *et al.*, 2018; Stanke *et al.*, 2004). *SnpEff* was used for annotating the variants (Cingolani *et al.*, 2012).

Development of SNP markers and linkage analysis

Two derived Cleaved Amplification Polymorphic Site (dCAPS) markers were developed to distinguish the *OsMAPK6* alleles of SM and BB42. One marker was developed by introducing a SacI restriction site in the SM allele while the other marker was developed by introducing a SmaI restriction site in the BB42 allele. The primer sequences were designed based on the SM reference sequence using dCAPS Finder 2.0 (Neff *et al.*, 2002). All the reactions were performed using 2X

KAPA Taq ReadyMix (Roche, Switzerland) with 500-1000 pmoles/µl primers and 20-40ng of the

template. The reaction conditions are as follows: 95℃ for 5 minutes, 35 cycles of 95℃ for 30

seconds, 58℃ for 30 seconds and 72℃ for 30 seconds, and 72℃ for 5 minutes. The PCR samples

were then added with 0.5-1µl of corresponding restriction enzymes and incubated at 37℃ for 6

hours to overnight before resolving them on a 2.5% agarose gel for 1 hour at 100V. Gel images

were captured on a gel documentation system (AlphaInnotech, USA). Further, another marker was

developed based on the tetra-primer amplification refractory mutation system (tetra-ARMS) PCR

(Ye *et al.*, 2001). A typical tetra-ARMS-PCR reaction contains four primers among which each of

the two inner primers is specific to the two alleles of interest. The tetra-ARMS reaction to

distinguish the *OsMAPK6* alleles was standardized, and the reaction composition is as follows:

500 nM of each primer, PCR buffer – 10mM tris-HCl (pH8.3), 20mM ammonium sulphate, 2mM

magnesium chloride, and 0.25mM of each dNTP, 0.2µl of 5U/µl Taq DNA polymerase (G-

biosciences), and 40ng of the template. The reaction conditions are as follows: 95℃ for 5 minutes,

35 cycles of 95℃ for 30 seconds, 58℃ for 20 seconds and 72℃ for 30 seconds, and 72℃ for 5

minutes. The samples were resolved on a 2.5% agarose gel for one hour at 100V. To test the

linkage between the OsMAPK6 allele and BB resistance, we tested the genotype of seven resistant

progenies and seven susceptible progenies from the F_2 generation using the dCAPS marker that

was developed as mentioned above.

Protein sequence analysis

Homologous sequences of OsMAPK6 were retrieved from NCBI through a BLASTp

search. Sequences of all the rice MAPKs were obtained from RGAP database (Hamilton *et al.*,

2025). Multiple sequence alignment (MSA) was performed using Multalin (Corpet, 1988). ESPript

3 was used to visualize MSA results (Robert and Gouet, 2014). WebLogo was used to generate

sequence alignment logos (https://weblogo.berkeley.edu/logo.cgi) (Crooks *et al.*, 2004). One-click

mode of Phylogeny.fr server was used for constructing the maximum likelihood trees of MAPK6

homologues and the rice MAPKs (Dereeper *et al.*, 2008). Phylogenetic trees were annotated using

iTOL (http://itol.embl.de/) (Letunic and Bork, 2021).

RNA sequencing and analysis

SM and BB42 plants were inoculated with the *Xoo* isolate IXO1221 during the maximum

tillering stage. One-centimeter-long leaf samples from the site of clipping were collected at the

time of infection as control (0 hour/Mock) and 24-hour post infection (Xoo) from both SM and

BB42. The samples were collected from three independent plants serving as biological replicates.

The samples were flash-frozen immediately after collection and stored at -80°C until further

processing. RNA was isolated from the samples using RNeasy Plant RNA Isolation Kit (Qiagen,

Germany) with on-column DNA digestion using RNase-Free DNase Set (Qiagen, Germany).

Purified RNA was quantified using Qubit^{TM} RNA HS Assay Kit (Thermo Fisher Scientific, USA).

RNA integrity was assessed using agarose gel electrophoresis and BioAnalyzer 2100 (Agilent

Technologies, USA). 1 µg of intact total RNA samples were used for library preparation using

Illumina TruSeq Stranded Total RNA with Ribo-Zero Plant (Illumina, USA). Sequencing was

performed on Illumina's NovaSeq6000 platform with cycling conditions to obtain 2x100bp reads

on an S2 flow cell.

The raw data were obtained, and the quality was assessed using *FastQC*. Adapters and

low-quality reads were trimmed using *Trim Galore! RNA-STAR* was used to align the trimmed

reads to the rice genome (MSU Rice Genome Annotation Project - version 7). Reads that mapped

to the genes were quantified using *featureCounts*. Differential gene expression was obtained using

the R library *DEseq2*. Genes with a false discovery rate ≤ 0.05 and $\log_2$ fold change ≤ -1 or ≥ 1

were considered differentially expressed genes (DEGs). For gene ontology, and pathway analyses

ShinyGO was used. iDEP.96 was used for *k*-means clustering and further analysis (Ge *et al.*, 2018).

Venny was used for generating Venn diagrams (Oliveros, J.C., 2007). Protein domain enrichment

among the DEGs was performed in STRING v12.0 (Szklarczyk *et al.*, 2023). RNA sequence

statistics are provided in Table S3.

Quantitative PCR for defense marker genes

Complementary DNA (cDNA) was synthesized from 1-5 microgram RNA using EcoDry

cDNA Synthesis Kit (Takara) containing Oligo(dT). The reaction condition is as follows: 42°C for

60 minutes; 70°C for 10 minutes. The synthesized cDNA samples were diluted to 15-25ng/µl for

quantitative PCR (qPCR). qPCR was performed using PowerSYBR kit (Invitrogen) in triplicates

per primer-sample combination. The reaction composition is as follows: 5µl of 2X master mix,

0.5µl each of forward primer (5µM) and reverse primer (5µM), 1µl template, 3µl nuclease-free

water. The reactions were performed in 384-well plates at the following conditions: 50°C for 2

minutes, 95°C for 10minutes, 35 cycles of 95°C for 10 seconds and 60°C for 30seconds and melt

curve analysis between 60℃ and 95℃. The reactions were set in Viaa7 (Applied Biosystems) or

CFX384 (BioRad) instruments. The primers for qPCR reactions were designed using QuantPrime

(Arvidsson *et al.*, 2008). The list of primers and corresponding sequences are provided in Table

S2.

Immunoblotting

For MAPK activation assay, one-week old SM and BB42 roots were treated with mock or 1μM

flg22 for 15 minutes. For assay in seedlings, 1-week laboratory-grown seedlings were harvested

and processed further. Protein samples were isolated from the above samples, resolved on a 12%

polyacrylamide SDS gel, transferred onto PVDF membrane and probed using the anti-Phospho-

erk1/2 antibody (#9101; 1:2000 dilution; Cell Signaling Technologies). The blots were stained

with Ponceau-S-stained for loading control.

For qualitative assessment of total phosphorylated proteins, SM and BB42 leaf samples from

plants in maximum tillering stage (~60 days) were processed as mentioned above and probed using

anti-Phosphoserine (ab9332; 1:1000; Abcam) and anti-Phosphothreonine/tyrosine (#9381; 1:3000;

Cell Signaling Technologies) antibodies. Ponceau-S-stained blot is used as loading control blots.

The immuno signals were visualized using HRP-conjugated anti-rabbit secondary antibody

(ab98440; 1:20000; Abcam) and ECL Western high-sensitivity substrate (Thermo Fisher

Scientific, USA).

BR sensitivity assays

All the assays pertinent to brassinosteroid treatment were performed as described earlier (Liu *et*

al., 2015). Briefly, for coleoptile elongation assay, seeds of SM and BB42 were soaked overnight

in fresh water, washed and plated on Petri plates containing solvent control or 5μM 24-

epibrassinolide (Real-Gene Labs, UP, India) and grown further. The hypocotyl length was

measured one-week post germination in dark conditions. Grain parameters like length and width

were measured using a graduated ruler.

Metabolite profiling

Metabolite profiling and analysis was performed on field-grown SM and BB42 leaves during

maximum tillering stage essentially by following the steps as mentioned elsewhere (Gokulan *et*

al., 2024). Sample preparation and processing were outsourced to Novelgene Technologies Pvt.

Ltd., Hyderabad, India.

Statistical analyses

The relevant statistical tests and considerations are provided in appropriate figure legends. Box

plots, bar plots, and multivariate dot plots were plotted on R4.2.2 using *ggplot* package and the

statistical parameters were integrated to the plots using the *ggpubr* package.

Author contributions

RVS, MSM, and HKP: conceptualization; RVS, MSM, HKP, and GCG: methodology; GCG:

formal analysis; GCG, MJ, DR, NG, ARS, NS, and DN: investigation; RVS, HKP, ST, MSM,

RMS, KMB, and GSL: resources; GCG: data curation; GCG: writing - original draft; GCG, RVS,

HKP, MSM, KMB, and GSL: writing - review & editing; GCG: visualization; RVS, HKP, and

MSM: supervision; RVS, HKP, and MSM: funding acquisition.

Acknowledgements

We thank the CCMB NGS Facility for nucleic acid sequencing services. We thank the skilled

workers at ICAR-IIRR and CSIR-CCMB for their support in field activities and maintenance. We

thank Sebastien Santini (CNRS/AMU IGS UMR7256) and the PACA Bioinfo platform for the

availability and management of the phylogeny.fr website. GCG, MJ, DR, NG, ARS, NS, and DN

thank the Council of Scientific and Industrial Research for fellowship. GCG thanks the Department

of Biotechnology for the MK Bhan - Young Researcher Fellowship.

Supplementary information

Figure S1: SM and BB42 are equally susceptible to *Xoc.*

Figure S2: F_2 phenotyping and marker development.

Figure S3: Mutation in OsMPAK6 does not affect gene splicing.

Figure S4: PCA plot of gene expression.

Figure S5: Pathway enrichment analysis among DEGs.

Figure S6: Protein domain enrichment among DEGs.

Figure S7: BB42 shows signs of autoimmunity.

Figure S8: Metabolite analysis of SM and BB42.

Figure S9: Pathway enrichment analysis of differentially accumulated metabolites.

Figure S10: Phylogeny of MAPK6 homologues across different forms of life.

Figure S11: Phylogeny of seventeen OsMAPK proteins.

Table S1: List of strains.

Table S2: List of primers.

Table S3: RNA-seq read statistics.

Table S4: Phenotype segregation analysis.

Table S5: DEG statistics.

Table S6: Differential expression of defense related genes.

Table S7: Differential expression of JAZ-encoding genes.

Table S8: Differential expression of BR-related genes.

Data S1: List of DEGs in across different comparisons.

Data S2: List of differentially accumulated metabolites.

Conflict of interest

No conflict of interest declared.

Data availability

544 **Funding Statement**

545 This work was supported by Council of Scientific and Industrial Research's grant [MLP0121-

546 Phase I and Phase II] and the Scientific and Engineering Research Board's JC Bose fellowship to

547 RVS [SB/S2/JCB-12/2014].

REFERENCES

Abe, A., Kosugi, S., Yoshida, Kentaro, Natsume, S., Takagi, H., Kanzaki, H., et al. (2012) *Genome sequencing reveals agronomically important loci in rice using MutMap. Nature Biotechnology*, **30**, 174–178.

Afgan, E., Baker, D., Batut, B., Van Den Beek, M., Bouvier, D., Ech, M., et al. (2018) *The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic Acids Research*, **46**, W537–W544.

Albrecht, C., Boutrot, F., Segonzac, C., Schwessinger, B., Gimenez-Ibanez, S., Chinchilla, D., et al. (2012) *Brassinosteroids inhibit pathogen-associated molecular pattern-triggered immune signaling independent of the receptor kinase BAK1. Proc Natl Acad Sci U S A*, **109**, 303–308.

Arvidsson, S., Kwasniewski, M., Riaño-Pachón, D.M., and Mueller-Roeber, B. (2008) *QuantPrime - A flexible tool for reliable high-throughput primer design for quantitative PCR. BMC Bioinformatics*, **9**, 1–15.

Belkhadir, Y., Jaillais, Y., Epple, P., Balsemão-Pires, E., Dangl, J.L., and Chory, J. (2012) *Brassinosteroids modulate the efficiency of plant immune responses to microbe-associated molecular patterns. Proc Natl Acad Sci U S A*, **109**, 297–302.

Berriri, S., Garcia, A.V., Frei dit Frey, N., Rozhon, W., Pateyron, S., Leonhardt, N., et al. (2012) *Constitutively active mitogen-activated protein kinase versions reveal functions of Arabidopsis MPK4 in pathogen defense signaling. Plant Cell*, **24**, 4281–4293.

Carpenter, S.C.D., Mishra, P., Ghoshal, C., Dash, P.K., Wang, L., Midha, S., et al. (2020) *An xa5 Resistance Gene-Breaking Indian Strain of the Rice Bacterial Blight Pathogen Xanthomonas oryzae pv. oryzae Is Nearly Identical to a Thai Strain. Frontiers in Microbiology*, **11**, 2328.

Chen, J., Wang, L., Yang, Z., Liu, H., Chu, C., Zhang, Z., et al. (2021) *The rice Raf-like MAPKKK OsILA1 confers broad-spectrum resistance to bacterial blight by suppressing the OsMAPKK4–OsMAPK6 cascade. Journal of Integrative Plant Biology*, **63**, 1815–1842.

Chen, X. and Ronald, P.C. (2011) *Innate immunity in rice. Trends in Plant Science*, **16**, 451–459.

Chujo, T., Miyamoto, K., Ogawa, S., Masuda, Y., Shimizu, T., Kishi-Kaboshi, M., et al. (2014) *Overexpression of phosphomimic mutated OsWRKY53 leads to enhanced blast resistance in rice. PLoS One*, **9**, e98737.

Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., et al. (2012) *A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly*, **6**, 80–92.

Corpet, F. (1988) *Multiple sequence alignment with hierarchical clustering. Nucleic acids research*, **16**, 10881–10890.

Crooks, G.E., Hon, G., Chandonia, J.-M., and Brenner, S.E. (2004) *WebLogo: a sequence logo generator. Genome Res*, **14**, 1188–1190.

D Couto, C.Z. (2016) *Regulation of pattern recognition receptor signalling in plants. Nat. Rev. Immunol.*, **16**, 537–552.

Dasari, A., Vemulapalli, P., Gonuguntla, R., Thota, D.K., Elumalai, P., Muppavarapu, K., et al. (2022) *Improvement of bacterial blight resistance of the popular variety, Nellore Mahsuri (NLR34449) through marker-assisted breeding. J Genet*, **101**, 7.

De Vleesschauwer, D., Van Buyten, E., Satoh, K., Balidion, J., Mauleon, R., Choi, I.-R., et al. (2012) *Brassinosteroids Antagonize Gibberellin- and Salicylate-Mediated Root Immunity in Rice. Plant Physiology*, **158**, 1833–1846.

Dereeper, A., Guignon, V., Blanc, G., Audic, S., Buffet, S., Chevenet, F., et al. (2008)
Phylogeny.fr: robust phylogenetic analysis for the non-specialist. Nucleic Acids Res, **36**,
W465-469.

Fiyaz, R.A., Shivani, D., Chaithanya, K., Mounika, K., Chiranjeevi, M., Laha, G.S., et al. (2022)
Genetic Improvement of Rice for Bacterial Blight Resistance: Present Status and Future
Prospects. Rice Science, **29**, 118–132.

Freh, M., Gao, J., Petersen, M., and Panstruga, R. (2022) *Plant autoimmunity—fresh insights into*
an old phenomenon. Plant Physiology, **188**, 1419–1434.

Gao, M., Liu, J., Bi, D., Zhang, Z., Cheng, F., Chen, S., and Zhang, Y. (2008) *MEKK1,*
MKK1/MKK2 and MPK4 function together in a mitogen-activated protein kinase cascade
to regulate innate immunity in plants. Cell Res, **18**, 1190–1198.

Ge, S.X., Son, E.W., and Yao, R. (2018) *iDEP: an integrated web application for differential*
expression and pathway analysis of RNA-Seq data. BMC Bioinformatics, **19**, 534.

Genot, B., Lang, J., Berriri, S., Garmier, M., Gilard, F., Pateyron, S., et al. (2017) *Constitutively*
Active Arabidopsis MAP Kinase 3 Triggers Defense Responses Involving Salicylic Acid
and SUMM2 Resistance Protein. Plant Physiology, **174**, 1238–1249.

Gokulan, C.G., Bangale, U., Balija, V., Ballichatla, S., Potupureddi, G., Rao, D., et al. (2024)
Multiomics-assisted characterization of rice-Yellow Stem Borer interaction provides
genomic and mechanistic insights into stem borer resistance in rice. Theor Appl Genet,
137, 122.

Goto, M. (1992) *Fundamentals of bacterial plant pathology*, 1st edn. Academic Press.

Greene, E.A., Codomo, C.A., Taylor, N.E., Henikoff, J.G., Till, B.J., Reynolds, S.H., et al. (2003)
Spectrum of Chemically Induced Mutations From a Large-Scale Reverse-Genetic Screen
in Arabidopsis. Genetics, **164**, 731–740.

Hamilton, J.P., Li, C., and Buell, C.R. (2025) *The rice genome annotation project: an updated*
database for mining the rice genome. Nucleic Acids Research, **53**, D1614–D1622.

Heese, A., Hann, D.R., Gimenez-Ibanez, S., Jones, A.M.E., He, K., Li, J., et al. (2007) *The*
receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants. Proc.
Natl. Acad. Sci. U.S.A., **104**, 12217–12222.

Hilaire, E., Young, S.A., Willard, L.H., McGee, J.D., Sweat, T., Chittoor, J.M., et al. (2007)
Vascular Defense Responses in Rice: Peroxidase Accumulation in Xylem Parenchyma
Cells and Xylem Wall Thickening. https://doi.org/10.1094/MPMI.2001.14.12.1411, **14**,
1411–1419.

Huang, N., Angeles, E.R., Domingo, J., Magpantay, G., Singh, S., Zhang, G., et al. (1997)
Pyramiding of bacterial blight resistance genes in rice: marker-assisted selection using
RFLP and PCR. Theoretical and Applied Genetics 1997 95:3, **95**, 313–320.

IRRI (2014) *Standard evaluation system for rice. International Rice Research Institute*. IRRI.

Jamaloddin, M., Mahender, A., Gokulan, C.G., Balachiranjeevi, C., Maliha, A., Patel, H.K., and
Ali, J. (2021) *Molecular Approaches for Disease Resistance in Rice*. In: *Rice Improvement*
, pp. 315–378. Cham: Springer International Publishing.

Jamaloddin, Md., Durga Rani, Ch.V., Swathi, G., Anuradha, Ch., Vanisri, S., Rajan, C.P.D., et al.
(2020) *Marker Assisted Gene Pyramiding (MAGP) for bacterial blight and blast resistance*
into mega rice variety "Tellahamsa." PLOS ONE, **15**, e0234088.

Jhingan, S., Kumar, A., Harloff, H.-J., Dreyer, F., Abbadi, A., Beckmann, K., et al. (2023) *Direct*
access to millions of mutations by whole genome sequencing of an oilseed rape mutant
population. The Plant Journal, **113**, 866–880.

Jiang, N., Yan, J., Liang, Y., Shi, Y., He, Z., Wu, Y., et al. (2020) *Resistance Genes and their Interactions with Bacterial Blight/Leaf Streak Pathogens (Xanthomonas oryzae) in Rice (Oryza sativa L.)—an Updated Review*. Rice, **13**, 1–12.

Jiang, R., Zhou, S., Da, X., Yan, P., Wang, K., Xu, J., and Mo, X. (2023) *OsMKK6 Regulates Disease Resistance in Rice. International Journal of Molecular Sciences*, **24**, 12678.

Jones, J.D.G. and Dangl, J.L. (2006) *The plant immune system. Nature 2006 444:7117*, **444**, 323–329.

Jones, J.D.G., Vance, R.E., and Dangl, J.L. (2016) *Intracellular innate immune surveillance devices in plants and animals. Science*, **354**.

Joshi, J.B., Arul, L., Ramalingam, J., and Uthandi, S. (2020) *Review Advances in the Xoo-rice pathosystem interaction and its exploitation in disease management. Journal of Biosciences*, **45**.

K. Reddy, A.P. (1979) *Relationship of Bacterial Leaf Blight Severity to Grain Yield of Rice. Phytopathology*, **69**, 967.

Kauffman, H.E., Reddy, A.P.K., Hsieh, S.P.Y., and Merca, S.D. (1973) *An improved technique for evaluating resistance of rice varieties to Xanthomonas oryzae. Plant Dis. Rep*, **57**, 737–741.

Kaur, A., Bansal, K., and Patil, P.B. (2020) *Extensive Genomic Rearrangements along with Distinct Mobilome and TALome are Associated with Extreme Pathotypes of a Rice Pathogen. Genome Biology and Evolution*, **12**, 3951–3956.

Kim, S.-H., Oikawa, T., Kyozuka, J., Wong, H.L., Umemura, K., Kishi-Kaboshi, M., et al. (2012) *The bHLH Rac Immunity1 (RAI1) Is Activated by OsRac1 via OsMAPK3 and OsMAPK6 in Rice Immunity. Plant Cell Physiol*, **53**, 740–754.

Kishi-Kaboshi, M., Okada, K., Kurimoto, L., Murakami, S., Umezawa, T., Shibuya, N., et al. (2010) *A rice fungal MAMP-responsive MAPK cascade regulates metabolic flow to antimicrobial metabolite synthesis. Plant J*, **63**, 599–612.

Lang, J., Genot, B., Bigeard, J., and Colcombet, J. (2022) *MPK3 and MPK6 control salicylic acid signaling by up-regulating NLR receptors during pattern- and effector-triggered immunity. Journal of Experimental Botany*, **73**, 2190–2205.

Lang, J., Genot, B., Hirt, H., and Colcombet, J. (2017) *Constitutive activity of the Arabidopsis MAP Kinase 3 confers resistance to Pseudomonas syringae and drives robust immune responses. Plant Signaling & Behavior*, **12**, e1356533.

Letunic, I. and Bork, P. (2021) *Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. Nucleic Acids Research*, **49**, W293–W296.

Lian, K., Gao, F., Sun, T., van Wersch, R., Ao, K., Kong, Q., et al. (2018) *MKK6 Functions in Two Parallel MAP Kinase Cascades in Immune Signaling. Plant Physiol*, **178**, 1284–1295.

Liao, K., Peng, Y.-J., Yuan, L.-B., Dai, Y.-S., Chen, Q.-F., Yu, L.-J., et al. (2020) *Brassinosteroids Antagonize Jasmonate-Activated Plant Defense Responses through BRI1-EMS-SUPPRESSOR1 (BES1)1. Plant Physiol*, **182**, 1066–1082.

Lieberherr, D., Thao, N.P., Nakashima, A., Umemura, K., Kawasaki, T., and Shimamoto, K. (2005) *A sphingolipid elicitor-inducible mitogen-activated protein kinase is regulated by the small GTPase OsRac1 and heterotrimeric G-protein in rice 1[w]. Plant Physiol*, **138**, 1644–1652.

Liu, S., Hua, L., Dong, S., Chen, H., Zhu, X., Jiang, J., et al. (2015) *OsMAPK6, a mitogen-activated protein kinase, influences rice grain size and biomass production. Plant Journal*, **84**, 672–681.

Loveless, A. (1969) *Possible relevance of O-6 alkylation of deoxyguanosine to the mutagenicity*
and carcinogenicity of nitrosamines and nitrosamides. Nature, **223**, 206–207.

Ma, Z., Qin, G., Zhang, Y., Liu, C., Wei, M., Cen, Z., et al. (2021) *Bacterial leaf streak 1 encoding*
a mitogen-activated protein kinase confers resistance to bacterial leaf streak in rice. The
Plant Journal, **107**, 1084–1101.

Meng, X. and Zhang, S. (2013) *MAPK Cascades in Plant Disease Resistance Signaling. Annu.*
Rev. Phytopathol., **51**, 245–266.

Mew, T.W. (1993) *Focus on Bacterial Blight of Rice. Plant Disease*, **77**, 5.

Mishra, D., Vishnupriya, M.R., Anil, M.G., Konda, K., Raj, Y., and Sonti, R.V. (2013) *Pathotype*
and Genetic Diversity amongst Indian Isolates of Xanthomonas oryzae pv. oryzae. PLOS
ONE, **8**, e81996.

Nakashita, H., Yasuda, M., Nitta, T., Asami, T., Fujioka, S., Arai, Y., et al. (2003) *Brassinosteroid*
functions in a broad range of disease resistance in tobacco and rice. The Plant Journal,
33, 887–898.

Neff, M.M., Turk, E., and Kalishman, M. (2002) *Web-based primer design for single nucleotide*
polymorphism analysis. Trends in genetics : TIG, **18**, 613–615.

Niño-Liu, D.O., Ronald, P.C., and Bogdanove, A.J. (2006) *Xanthomonas oryzae pathovars: model*
pathogens of a model crop. Molecular Plant Pathology, **7**, 303–324.

Oliva, R., Ji, C., Atienza-Grande, G., Huguet-Tapia, J.C., Perez-Quintero, A., Li, T., et al. (2019)
Broad-spectrum resistance to bacterial blight in rice using genome editing. Nature
Biotechnology, **37**, 1344–1350.

Oliveros, J.C. (2007) *Venny. An interactive tool for comparing lists with Venn's diagrams.*

Ou, S.H. (1973) *A handbook of rice diseases in the tropics.* University of Illinois at Urbana-
Champaign: International Rice Research Institute.

Ou, S.H. (1985) *Rice diseases*, 2nd edn. Commonwealth Mycological Institute.

Pontiggia, D., Benedetti, M., Costantini, S., De Lorenzo, G., and Cervone, F. (2020) *Dampening*
the DAMPs: How Plants Maintain the Homeostasis of Cell Wall Molecular Patterns and
Avoid Hyper-Immunity. Frontiers in Plant Science, **0**, 2010.

Potupureddi, G., Balija, V., Ballichatla, S., Gokulan, C.G., Awalellu, K., Lekkala, S., et al. (2021)
Mutation resource of Samba Mahsuri revealed the presence of high extent of variations
among key traits for rice improvement. PLOS ONE, **16**, e0258816.

Pradhan, S.K., Nayak, D.K., Mohanty, S., Behera, L., Barik, S.R., Pandit, E., et al. (2015)
Pyramiding of three bacterial blight resistance genes for broad-spectrum resistance in
deepwater rice variety, Jalmagna. Rice, **8**, 19.

Quibod, I.L., Atieza-Grande, G., Oreiro, E.G., Palmos, D., Nguyen, M.H., Coronejo, S.T., et al.
(2020) *The Green Revolution shaped the population structure of the rice pathogen*
Xanthomonas oryzae pv. oryzae. ISME J, **14**, 492–505.

Robert, X. and Gouet, P. (2014) *Deciphering key features in protein structures with the new*
ENDscript server. Nucleic Acids Research, **42**.

Rodriguez, M.C.S., Petersen, M., and Mundy, J. (2010) *Mitogen-activated protein kinase*
signaling in plants. Annu Rev Plant Biol, **61**, 621–649.

Savary, S., Willocquet, L., Pethybridge, S.J., Esker, P., McRoberts, N., and Nelson, A. (2019) *The*
global burden of pathogens and pests on major food crops. Nature Ecology and Evolution,
3, 430–439.

Stanke, M., Steinkamp, R., Waack, S., and Morgenstern, B. (2004) *AUGUSTUS: a web server for*
gene finding in eukaryotes. Nucleic Acids Research, **32**, W309–W312.

Su, J., Yang, L., Zhu, Q., Wu, H., He, Y., Liu, Y., et al. (2018) *Active photosynthetic inhibition*
mediated by MPK3/MPK6 is critical to effector-triggered immunity. PLOS Biology, **16**,
e2004122.
Sugihara, Y., Young, L., Yaegashi, H., Natsume, S., Shea, D.J., Takagi, H., et al. (2020) *High-*
performance pipeline for MutMap and QTL-seq. bioRxiv.
Sun, T. and Zhang, Y. (2022) *MAP kinase cascades in plant development and immune signaling.*
EMBO reports, **23**, e53817.
Sundaram, R.M., Vishnupriya, M.R., Biradar, S.K., Laha, G.S., Reddy, G.A., Rani, N.S., et al.
(2008) *Marker assisted introgression of bacterial blight resistance in Samba Mahsuri, an*
elite indica rice variety. Euphytica, **160**, 411–422.
Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., et al. (2023) *The*
STRING database in 2023: protein-protein association networks and functional
enrichment analyses for any sequenced genome of interest. Nucleic Acids Res, **51**, D638–
D646.
Takagi, H., Abe, A., Yoshida, K., Kosugi, S., Natsume, S., Mitsuoka, C., et al. (2013) *QTL-seq:*
Rapid mapping of quantitative trait loci in rice by whole genome resequencing of DNA
from two bulked populations. Plant Journal, **74**, 174–183.
Till, B.J., Cooper, J., Tai, T.H., Colowit, P., Greene, E.A., Henikoff, S., and Comai, L. (2007)
Discovery of chemically induced mutations in rice by TILLING. BMC Plant Biology, **7**, 1–
12.
Tsuda, K., Mine, A., Bethke, G., Igarashi, D., Botanga, C.J., Tsuda, Y., et al. (2013) *Dual*
Regulation of Gene Expression Mediated by Extended MAPK Activation and Salicylic Acid
Contributes to Robust Innate Immunity in Arabidopsis thaliana. PLOS Genetics, **9**,
e1004015.
Wang, D., Wang, H., Liu, Q., Tu, R., Zhou, X., Zhang, Y., et al. (2021) *Reduction of OsMPK6*
activity by a R89K mutation induces cell death and bacterial blight resistance in rice. Plant
Cell Rep, **40**, 835–850.
Xiaobo, Z., Mu, Z., Mawsheng, C., Xuewei, C., and Jing, W. (2020) *Deciphering Rice Lesion*
Mimic Mutants to Understand Molecular Network Governing Plant Immunity and Growth.
Rice Science, **27**, 278–288.
Xu, R., Duan, P., Yu, H., Zhou, Z., Zhang, B., Wang, R., et al. (2018) *Control of Grain Size and*
Weight by the OsMKKK10-OsMKK4-OsMAPK6 Signaling Pathway in Rice. Molecular
Plant, **11**, 860–873.
Yamaguchi, K., Nakamura, Y., Ishikawa, K., Yoshimura, Y., Tsuge, S., and Kawasaki, T. (2013)
Suppression of rice immunity by Xanthomonas oryzae type III effector Xoo2875. Biosci
Biotechnol Biochem, **77**, 796–801.
Ye, S., Dhillon, S., Ke, X., Collins, A.R., and Day, I.N.M. (2001) *An efficient procedure for*
genotyping single nucleotide polymorphisms. Nucleic acids research, **29**, 88–88.
Yi, J., Lee, Y.-S., Lee, D.-Y., Cho, M.-H., Jeon, J.-S., and An, G. (2016) *OsMPK6 plays a critical*
role in cell differentiation during early embryogenesis in Oryza sativa. J Exp Bot, **67**,
2425–2437.
Yuan, M., Jiang, Z., Bi, G., Nomura, K., Liu, M., Wang, Y., et al. (2021) *Pattern-recognition*
receptors are required for NLR-mediated plant immunity. Nature, **592**, 105–109.
Yugander, A., Ershad, M., Muthuraman, P.P., Prakasam, V., Ladhalakshmi, D., Sheshu Madhav,
M., et al. (2022) *Understanding the variability of rice bacterial blight pathogen,*

Xanthomonas oryzae pv. oryzae in Andhra Pradesh, India. Journal of Basic Microbiology,
62, 185–196.
Yugander, A., Sundaram, R.M., Ladhalakshmi, D., Hajira, S.K., Prakasam, V., Prasad, M.S., et al.
(2017) *Virulence profiling of Xanthomonas oryzae pv. oryzae isolates, causing bacterial*
blight of rice in India. European Journal of Plant Pathology, **149**, 171–191.
Zhang, Z., Liu, Y., Huang, H., Gao, M., Wu, D., Kong, Q., and Zhang, Y. (2017) *The NLR protein*
SUMM2 senses the disruption of an immune signaling MAP kinase cascade via CRCK3.
EMBO Rep, **18**, 292–302.
Zhang, Z., Wu, Y., Gao, M., Zhang, J., Kong, Q., Liu, Y., et al. (2012) *Disruption of PAMP-*
Induced MAP Kinase Cascade by a Pseudomonas syringae Effector Activates Plant
Immunity Mediated by the NB-LRR Protein SUMM2. Cell Host & Microbe, **11**, 253–263.
Zhao, H., Li, J., Yang, L., Qin, G., Xia, C., Xu, X., et al. (2021) *An inferred functional impact map*
of genetic variants in rice. Molecular Plant, **14**, 1584–1599.
Zipfel, C. and Felix, G. (2005) *Plants and animals: A different taste for microbes? Current*
Opinion in Plant Biology, **8**, 353–360.
793
794

795   **Table 1:** List of high-frequency SNPs identified within the mapped genomic interval, their SNP

796   indices, and effect.

Chr	Pos[a]	SNP index (SM x BB42)	SNP index (BB42 x SM)	Effect
Chr6	2243785	0.76	0.72	Intron variant
Chr6	2358077	0.81	1.00	Intron variant
Chr6	2379923	0.84	0.94	Downstream gene variant
Chr6	**2664062**	**0.94**	**0.90**	**Missense variant & splice region variant;** SIFT - deleterious (0)
Chr6	3057969	0.88	0.81	Downstream gene variant

[a] Coordinates in SM reference genome

Figure 1: **BB42 is a dwarf and bacterial blight resistant derivative of SM.** (A) and (B)
Comparison of the morphological features of SM and BB42 showing a significantly reduced height
of the latter. The statistical significance was calculated using unpaired two-tailed Student's *t* test.
Number of samples: SM=15, BB42=13. (C) Representative leaves of SM and BB42 showing the
disease symptoms at 14 days post inoculation (dpi). (D) Lesion length readings of *Xoo* infected
SM and BB42 plants at 14 days post inoculation. Fully expanded leaves were clip inoculated with
Xoo suspension at OD_{600} 1.0 (n>5). Statistical significance was calculated using one-way Anova
and comparisons between groups were performed using Tukey's HSD (Honest Significant
Difference) test with a 95% family-wise confidence level. $P<0.05$ was considered statistically
significant. In (B) and (D) boxes represent the 25th to 75th percentile while the line within the box
indicates the median. The whiskers extend to maximum and minimum data points that are within
1.5 times the interquartile range. Red dots indicate outliers.

Figure 2: A novel mutation in *OsMAPK6* is linked to BB resistance. Allele frequency plots of
chromosome 6 indicating the mapped regions in two F_2 population: (A) SM x BB42 and (B) BB42
x SM. The top plot represents the regression analysis of SNP-index of resistant bulk, middle plot
represents that of susceptible bulk, and the bottom plot indicates the delta SNP-index plot. The red
line is the moving average line plotted by averaging the SNP or delta SNP indices in 2Mb windows
and a 100kb step size. Each dot represents an SNP. The transparent orange bar indicates the
mapped genomic interval. (C) Schematic gene diagram of *OsMAPK6* indicating the affected base
(red arrow) and the amino acid substitution caused. (D) Multiple sequence alignment logo of 100
plant homologues of MAPK6 spanning the Serine residue of interest (red arrowhead). (E)
Genotyping of selected F_2 progenies using MAPK6 allele-specific PCR markers showing co-
segregation of the mutant allele with resistance.

Figure 3: SM and BB42 share a substantial *Xoo*-mediated transcriptional landscape. (A)
Number of Differentially Expressed Genes (DEGs) identified in various comparisons of the
transcriptome data. (B) Venn diagram showing common and unique differential gene expression
patterns in SM and BB42 samples challenged with *Xoo*. Up and Down mean upregulation or
downregulation of genes in *Xoo* treated condition as compared to mock treated condition in the
mentioned genotypes. (C) k-means clustering of gene expression profiles of the top 2000 genes.
Lines with different colours indicate different clusters based on the gene expression values.
Number of genes within each cluster is provided in the legend. Enriched terms adjacent to each
cluster are the GO terms that are enriched within corresponding clusters.

Figure 4: BB42 exhibits constitutive immunity. (A) Gene ontology term enrichment of the genes
that are upregulated (top panel) or downregulated (bottom panel) in mock samples of BB42 as
compared to the corresponding SM samples. (B) Box plots showing the abundance of certain
primary and secondary metabolites that are differentially accumulated between SM and BB42. The
raw peak intensities were normalised and plotted on the y-axis. The numbers above the horizontal
lines indicate the *P*-values as calculated using unpaired Student's *t*-test considering equal variance
between samples. *P*<0.05 is considered statistically significant. Boxes represent the 25th to 75th
percentile while the line within the box indicates the median. The whiskers extend to maximum
and minimum data points.

Figure 5: Brassinosteroid signalling is impaired in BB42. (A) Comparison of grain size of SM
and BB42. Scale bar = 1 cm. (B) and (C) bar graphs depicting the grain length and grain width,
respectively, of SM and BB42. N=30. Bars are drawn to average and the error bars indicate
standard deviation. The dots represent the data points. (D) Box plots showing the length of the
hypocotyl upon treatment of the seeds with mock solution or solution containing 5µM 24-
epibrassinolide. Boxes represent the 25th to 75th percentile while the line within the box indicates
the median. The whiskers extend to maximum and minimum data points that are within 1.5 times
the interquartile range. Solid black dots outside the whiskers, if any, indicate outliers. N>10. The
numbers above the horizontal line indicates *P*-values as calculated using unpaired Student's *t*-test
considering equal sample variance. *P*<0.05 is considered statistically significant. (E) Box plot
showing the abundance of brassinolide in SM and BB42 leaves. Boxes represent the 25th to 75th
percentile while the line within the box indicates the median. The whiskers extend to maximum
and minimum data points.

Figures

Figure 1 to 5.

A novel allele of *Mitogen Activated Protein Kinase 6* is linked to disease resistance and constitutive immunity in rice.

Gokulan C.G.[1,6,†], Mohammed Jamaloddin[1,†], Deepti Rao[1], Namami Gaur[1,2], Amita Rani Supriyo[1], Nisha Sao[1], Deepak Niranjan[1,7], Shrish Tiwari[1], Gouri S. Laha[3], Kalyani M. Barbadikar[3], Raman Meenakshi Sundaram[3], Sheshu Madhav Maganti[3,4,*], Hitendra K. Patel[1,5,*], Ramesh V. Sonti[1,7,*]

Figure 1: **BB42 is a dwarf and bacterial blight resistant derivative of Samba Mahsuri (SM).** (A) and (B) Comparison of the morphological features of SM and BB42 showing a significantly reduced height of the latter. The statistical significance was calculated using unpaired two-tailed Student's *t* test. Number of samples: SM=15, BB42=13. (C) Representative leaves of SM and BB42 showing the disease symptoms at 14 days post inoculation (dpi). (D) Lesion length readings of *Xoo* infected SM and BB42 plants at 14 days post inoculation. Fully expanded leaves were clip inoculated with *Xoo* suspension at OD_{600} 1.0 (n>5). Statistical significance was calculated using one-way Anova and comparisons between groups were performed using Tukey's HSD (Honest Significant Difference) test with a 95% family-wise confidence level. *P*<0.05 was considered statistically significant. In (B) and (D) boxes represent the 25th to 75th percentile while the line within the box indicates the median. The whiskers extend to maximum and minimum data points that are within 1.5 times the interquartile range. Red dots indicate outliers.

Figure 2: A novel mutation in *OsMAPK6* is linked to BB resistance. Allele frequency plots of chromosome 6 indicating the mapped regions in two F₂ population: (A) SM x BB42 and (B) BB42 x SM. The top plot represents the regression analysis of SNP-index of resistant bulk, middle plot represents that of susceptible bulk, and the bottom plot indicates the delta SNP-index plot. The red line is the moving average line plotted by averaging the SNP or delta SNP indices in 2Mb windows and a 100kb step size. Each dot represents an SNP. The transparent orange bar indicates the mapped genomic interval. (C) Schematic gene diagram of *OsMAPK6* indicating the affected base (red arrow) and the amino acid substitution caused. (D) Multiple sequence alignment logo of 100 plant homologues of MAPK6 spanning the Serine residue of interest (red arrowhead). (E)

Genotyping of selected F_2 progenies using MAPK6 allele-specific PCR markers showing co-segregation of the mutant allele with resistance.

Figure 3: SM and BB42 share a substantial *Xoo*-mediated transcriptional landscape. (A) Number of Differentially Expressed Genes (DEGs) identified in various comparisons of the transcriptome data. (B) Venn diagram showing common and unique differential gene expression patterns in SM and BB42 samples challenged with *Xoo*. Up and Down mean upregulation or downregulation of genes in *Xoo* treated condition as compared to mock treated condition in the mentioned genotypes. (C) k-means clustering of gene expression profiles of the top 2000 genes. Lines with different colours indicate different clusters based on the gene expression values. Number of genes within each cluster is provided in the legend. Enriched terms adjacent to each cluster are the GO terms that are enriched within corresponding clusters.

Figure 4: BB42 exhibits constitutive immunity. (A) Gene ontology term enrichment of the genes that are upregulated (top panel) or downregulated (bottom panel) in mock samples of BB42 as compared to the corresponding SM samples. (B) Box plots showing the abundance of certain primary and secondary metabolites that are differentially accumulated between SM and BB42. The raw peak intensities were normalised and plotted on the y-axis. The numbers above the horizontal lines indicate the *P*-values as calculated using unpaired Student's *t*-test considering equal variance between samples. *P*<0.05 is considered statistically significant. Boxes represent the 25th to 75th

percentile while the line within the box indicates the median. The whiskers extend to maximum and minimum data points.

Figure 5: Brassinosteroid signalling is impaired in BB42. (A) Comparison of grain size of SM and BB42. Scale bar = 1 cm. (B) and (C) bar graphs depicting the grain length and grain width, respectively, of SM and BB42. N=30. Bars are drawn to average and the error bars indicate standard deviation. The dots represent the data points. (D) Box plots showing the length of the hypocotyl upon treatment of the seeds with mock solution or solution containing 5μM 24-epibrassinolide. Boxes represent the 25th to 75th percentile while the line within the box indicates the median. The whiskers extend to maximum and minimum data points that are within 1.5 times the interquartile range. Solid black dots outside the whiskers, if any, indicate outliers. N>10. The numbers above the horizontal line indicates *P*-values as calculated using unpaired Student's *t*-test

considering equal sample variance. $P<0.05$ is considered statistically significant. (E) Box plot showing the abundance of brassinolide in SM and BB42 leaves. Boxes represent the 25th to 75th percentile while the line within the box indicates the median. The whiskers extend to maximum and minimum data points.