Growth charts of infant visual neurodevelopment generalize across global contexts

Emma T. Margolis[1,2]*#, Chris Camp[3]*, Ana C. Sobrino[1,2], Guilherme V. Polanczyk[4], Daniel Fatori[4,5], Khula South Africa Team, Germina Team, LABS Team, GABA Team, 1kD EEG Working Group, 1kD Machine Learning Working Group, Laura Cornelissen[6], Charles B. Berde[6], Takao K. Hensch[7], Charles A. Nelson[8,9], Elizabeth Shephard[4,5,10], Kirsten A. Donald[11,12], Dustin Scheinost[13-17]^, Laurel J. Gabard-Durnam[1,2]^#

* indicates equal contributions; ^ indicates equal contributions; # indicates corresponding authors

Author Affiliations
1. Department of Psychology, Northeastern University, Boston, MA, USA
2. Center for Cognitive and Brain Health, Northeastern University, Boston, MA, USA
3. Interdepartmental Neuroscience Program, Yale School of Medicine, Yale University, New Haven, CT, USA
4. Division of Child & Adolescent Psychiatry, Department & Institute of Psychiatry, Faculdade de Medicina FMUSP, Universidade de São Paulo, São Paulo, Brazil
5. Laboratory of Psychopathology and Psychiatric Therapeutics LIM-23, Institute of Psychiatry, Clinical Hospital HCFMUSP, School of Medicine, University of São Paulo, São Paulo, Brazil
6. Department of Anesthesiology, Critical Care and Pain Medicine, Boston Children's Hospital, Harvard Medical School, Boston, MA, USA
7. F.M. Kirby Neurobiology Center, Department of Neurology, Boston Children's Hospital, Harvard Medical School, Boston, MA, USA.
8. Laboratories of Cognitive Neurosciences, Department of Pediatrics, Boston Children's Hospital, Harvard Medical School, Boston, MA, USA
9. Harvard Graduate School of Education, Harvard University, Cambridge, MA, USA
10. Department of Developmental Learning and Personality Psychology, Institute of Psychology, University of São Paulo, São Paulo, Brazil
11. Division of Developmental Paediatrics, Department of Paediatrics and Child Health, Red Cross War Memorial Children's Hospital, University of Cape Town, Cape Town, South Africa
12. The Neuroscience Institute, University of Cape Town, Cape Town, South Africa
13. Department of Radiology & Biomedical Imaging, Yale School of Medicine, New Haven, CT, USA
14. Department of Biomedical Engineering, Yale University, New Haven, CT, USA
15. Department of Statistics & Data Science, Yale University, New Haven, CT, USA
16. Child Study Center, Yale School of Medicine, New Haven, CT, USA
17. Wu Tsai Institute, Yale University, New Haven, CT, USA

40 **Corresponding Authors:** Emma Margolis, margolis.emm@northeastern.edu; Laurel Gabard-
41 Durnam, l.gabard-durnam@northeastern.edu
42

Abstract (142/50 words)
Normative brain growth charts in early life hold great promise for furthering basic and clinical
science. We leverage the rapid, substantial development of visual cortex function that is indexed
by visual-evoked potentials (VEP) in electroencephalography to create longitudinal normative
growth curves of task-related brain function with 1374 observations contributed by 802 infants
(57 to 579 days old) from South Africa, Brazil, and the United States. Site-specific models were
cross-validated and showed excellent fits to other sites' samples, demonstrating functional
growth curves generalize across contexts robustly. Deviations from the normative growth models
associated with early environmental and behavioral measures such as prenatal exposures and
postnatal cognition. These findings demonstrate the utility of using functional growth charts to
understand and potentially act on individual neurodevelopmental trajectories. VEP brain function
growth charts represent a new direction for EEG research to support healthy brain development
globally.
56

Acknowledgements: The authors thank the caregivers, families, and infants who made this work
possible. The Khula South Africa Team tag includes the following authors: Lauren Davel,
Bokang Methola, Donna Herr, Marlie Miles, Michal Zieff, Thandeka Mazubane and Zayaan
Goolam Nabi. The Germina Team tag includes the following authors: Alline Cristina de
Campos, André Fujita, Carla R. Taddei, Maria Rita Passos-Bueno, Patricia Beltrão-Braga. The
LABS Team tag includes the following authors: Trevor Bissert and Berit Hartjen. The GABA
Team tag includes the following authors: Siobhan Coffman, Isabelle Kim, Alice Tao, Ellen
Underwood, and Maria Maloney. The 1kD EEG Working Group tag includes the following
authors: Nicolò Pini, William Fifer, Sarah A. McCormick, Priyanka Ghosh, Cara Bosco, and
Michal R. Zieff. The 1kD Machine Learning Working Group tag includes the following authors:
Peter Wijeratne, Kevin S. Bonham, Daniel C. Alexander, and Guilherme Zainotti Miguel Fahur
Bottino. This research is supported by the Wellcome LEAP 1kD Program (KAD, GVP, CAN,
ES, DS, LJGD), Boston Children's Hospital (Trailblazer Research Award to LC and Sara Page
Mayo Endowment for Pediatric Pain Research, Treatment, and Education to CBB), and the
World Premier International Research Center Initiative–International Research Center for
Neurointelligence (TKH). The views expressed in this article are those of the authors only.
73

74 **Keywords:** normative modeling; visual-evoked potential; developmental trajectories; cross-
75 cultural; task-based EEG; multi-site
76

77   **Introduction**

Normative changes quantified through physical growth charts have transformed science,

public health, and pediatric care[1–5]. The brain undergoes tremendous functional and structural

change across the lifespan[6–12]. Normative brain growth charts also hold great promise for

furthering basic and clinical science[13–17]. Growth charts may reveal milestones in brain

development and provide individual-level neural indicators facilitating risk identification,

prognosis, stratification, or intervention evaluation[13,14,18,19]. Several candidate growth charts for

normative brain structure[13,18,20] and function at rest or in sleep are emerging for the lifespan[21–23].

Growth charts characterizing awake brain function during task paradigms can offer unique

information about how brains function and change. This approach may also improve chart utility

as task-related activity, compared to resting activity, has shown greater sensitivity and specificity

in associating individual differences in brain function with cognition, symptoms, and

behavior[24,25]. However, normative growth charts of task-related brain function have yet to be

constructed.

There are multiple challenges in identifying which charts of task-related brain function

are feasible and useful[13,17]. These include identifying standardized, scalable, and geoculturally-

appropriate paradigms to elicit specific neural activity across contexts and development[26–33].

Statistical strategies must also facilitate external validation of growth curves to test their

robustness across contexts and integrate information from multiple features generated during the

task[17]. Finally, while initial brain growth charts have taken a lifespan approach, growth curves

have poorer performance at their tails, meaning brain changes in infancy may be less well-

characterized[34]. However, elevated early neuroplasticity enables rapid, substantial brain

development[35–38]. As mandated for physical growth charts, targeted brain charts with dense

longitudinal sampling in early life can best capture these changes[4,39].

Here, we address these challenges to construct and evaluate brain function growth charts

in awake infants. We focus on characterizing the rapid, early development of visual cortical

function that can be precisely and efficiently measured through a standardized phase-reversal

visual-evoked potential (VEP) paradigm across contexts and ages[29–32,40–43]. Charting visual

neurodevelopment provides critical information about foundational sensory development. It also

provides early functional indicators of risk, scaffolding for subsequent higher-order cognition,

and opportunities for translational research across species[37,43–46]. We chart longitudinal data

(1374 observations) over the first two years of life from four cohorts across three continents that

represent distinct cultural, linguistic, geographic, and socioeconomic contexts (Cape Town,

South Africa, São Paulo, Brazil, and Boston, USA) with *a priori* harmonization of paradigm,

data acquisition, and processing parameters[47–49]. We leverage electroencephalography (EEG) as

an ideal method to capture standardized features of awake brain function in early life[26,28,29] and

generalized additive models of location, scale, and space (GAMLSS) with external cross-

validation to create and generalize growth curves[4,17,50].We demonstrate that brain function

growth charts in infancy are feasible, stable, and generalizable. We also show that they capture

individual deviations that are both sensitive to early developmental risk factors and relate to
emerging behavior.
118
Results
We constructed developmental growth curves of visual brain function from 57 to 579
days of age with participants (n=802) from four longitudinal cohorts (i.e., Khula South Africa,
Germina, LABS, GABA) across three continents (i.e., Africa, South America, and North
America) (Fig. 1). Cohorts' characteristics are summarized in Table 1, and demographic
information for each cohort is summarized in Table S1. Paradigm, acquisition, and processing
were harmonized *a priori* across all sites, and processing parameters are available in Tables S2
and S3. Visual brain function was indexed by 128-channel electroencephalography (EEG)-
derived pattern-reversal visual-evoked potentials (VEP) using a standard black-white
checkerboard paradigm on an external monitor while infants sat on their caregiver's lap[29,51].
VEPs were extracted and quantified from an occipital ROI (Fig. 2B) over the primary visual
cortex generator[52–57] with respect to six standard morphological features—peak amplitudes and
latencies to those peak amplitudes (i.e., latencies) for the N1, P1, and N2 components (Fig.
2A)[58]. These morphological features reflect distinct functional and structural processes that
characterize primary visual cortical development. Specifically, N1 and N2 components reflect
parvocellular visual pathway development (sensitive to color and spatial detail), while the P1
component is generated by the magnocellular pathway (sensitive to motion)[59–64]. Amplitude
features reflect cortical neurotransmission, especially the balance of excitatory/inhibitory
postsynaptic potentials[58,65–67], while latencies largely reflect the structural integrity and
myelination of the visual pathways[51,68,69]. We modeled the six VEP features independently using
generalized additive models of location, scale, and space (GAMLSS), a World Health
Organization-recommended framework to model non-linear growth trajectories[4].
141

142
Figure 1: *Study Overview.* We created longitudinal growth curves using data from 802 infants (*n*=1374 EEG
observations) between 57 and 579 days old across three continents. 128-channel EEG data were *a priori* harmonized
for paradigm, acquisition, and processing parameters across sites. We independently trained GAMLSS curves for
each site and VEP feature—amplitudes and latencies for the N1, P1, and N2 components. The GAMLSS curves
were externally cross-validated across sites. Individual centiles were extracted and associated with a known

148 developmental risk factor (prenatal alcohol exposure) and subsequent cognitive scores on a global public health
149 measure (Global Scales of Early Development, Long Form).
150

Cohort	Site	Age Range (Days)	N infants	N infants Providing >1 Useable Timepoint	N observations
Khula	Cape Town, South Africa	[58, 516]	303	214 (70.63%)	622
Germina	São Paulo, Brazil	[94, 312]	390	110 (28.21%)	500
LABS	Boston, USA	[85, 579]	56	38 (67.86%)	157
GABA	Boston, USA	[57, 366]	53	33 (62.26%)	95

151 **Table 1:** *Cohort Characteristics.*
152

Generalizing Growth Curves of Visual Neurodevelopment

We trained GAMLSS curves for each VEP feature to model normative development of
visual cortex function across sites. The fitting and testing procedures are outlined in Fig. S1.
Models were fit with p-spline smoothing terms and fixed effects for each feature. Amplitudes
were shifted by their minimum to have the range $[1, \infty)$. We prioritized developing models
generalizable across all three sites. Distributions for each VEP feature model were chosen by
minimizing the Bayesian information criterion (BIC) in the held-out site of a cross-validation
fold. Smoothing penalties were tuned by optimizing the generalized Akaike information criterion
penalty k to minimize the total validation BIC. Using the established hyperparameters, we fit
models on all pairs of site datasets. Models were assessed by examining the residuals' worm
plots and Q statistics and comparing the predicted and actual VEP feature values with Pearson r
correlation and root mean square error (RMSE). 95% confidence intervals were generated for
correlations with 1000-fold nonparametric bootstraps.
All VEP feature curves exhibited high model fits, with an average RMSE of 1.13 μV for
amplitudes and 13.24 ms for latencies. Similarly, the average Pearson correlation across site
datasets between the predicted and observed z-scores ranged between $r=.68$ (N2 Latency, 95%
CI: .64–.72) to $r=1$ (N1 Amplitude, 95% CI: .99–1). Additional model statistics are in Table S5.
We did not observe significant differences in correlations across site pairs with an omnibus
ANOVA ($F(2)=.15$, $p=.86$), suggesting that the models fit site pairs equally well. Amplitudes
had higher correlations between models and data than latencies (two-sided t test: $t(14.36)=2.95$,
$p<.05$), likely due to their simpler distributions. Including sex as a covariate did not change
model fit for any of the VEP features. Therefore, it was not included in any of the models.

Models Generalize Across Sites

To assess generalizability across sites, we then evaluated model fit in the held-out dataset
for each site pair. All feature curves exhibited high model fits on the held-out datasets, with an
average RMSE of 2.16 μV for amplitudes and 15.94 ms for latencies (Table S5). The average
Pearson correlation across datasets between the predicted and observed z-scores ranged between
$r=.71$ (N2 Latency, 95% CI: .66–.76) to $r=.99$ (N1 Amplitude, 95% CI: .99–1). As with the
within-dataset fits, we did not observe significant differences in correlations across held-out
datasets with an omnibus ANOVA ($F(2)=.21$, $p=.81$).
183

Figure 2: *Characterizing the VEP Over Infancy.* (A) Demonstrative VEP waveform with labeled components (i.e., N1, P1, N2) and features (i.e., amplitude, latency). (B) 128-channel electrode configuration with highlighted occipital region of interest (i.e., E70, E71, E75, E76, E83). (C) Averaged VEP waveforms across infants (n=15 each) for the mean age of our dataset (in days) and ± 1 SD (i.e., 227.66, 353.23, 102.09 days). (D) Growth curves for VEP features trained on all three sites (*n* = 1371 observations). Nonlinear changes are primarily observed in the P1 amplitude and the P1 and N2 latencies.

Characterizing VEP Development

Next, we characterized the shape of the VEP growth curves to identify developmental trends in visual function across individuals. We trained a model for each feature on all available data to provide the best-powered representation of the growth curves (Table S6-7). The six VEP features exhibited different growth shapes, representing distinct developmental patterns in their trajectories (Fig. 2D). The best underlying distributions as determined by cross-validation were as follows: sinh-arcsinh (SHASH) for N1 amplitude; Johnson's S_U for P1 amplitude, N2 amplitude, and P1 latency; *t*-family (TF) for N1 latency, and generalized inverse Gaussian (GIG) for N2 Latency. We plotted the fitted terms to examine the behavior of each parameter for the

models and used numerical derivation to determine slope minima and maxima (Table S8, Fig.
S14-23). We observed several trends across the features. The central tendency (μ, the 'location'
parameter) of the amplitude peaks diminished with age, with the negative component amplitudes
(N1, N2) increasing and P1 decreasing. Negative component amplitudes (N1, N2) demonstrated
steeper slopes of change than the P1 component during infancy. Latency central tendency
parameters also decreased with age, with N2 showing the steepest slopes over infancy. Variance
(σ, the 'scale' parameter) decreased with age in all features except P1 latency as the components
matured and converged. The remaining two parameters, ν and τ, were more variable across
features as they represent different moments depending on the distribution across individuals.
We also examined differences in VEP development to identify milestones in early visual
function (Fig. 2C, S24-25). We estimated the 50th percentile of each feature across the age range
and calculated the first and second derivatives (Fig. S26-29). As is expected, we observed
decreasing latencies over development, indicating increased processing speed and myelination as
infants get older[58]. P1 and N2 latencies had nonlinear shapes, with P1 latency maturing faster.
We observed minimal change in N1 latencies over infancy. The magnitudes of amplitudes
decreased with age, with negative amplitudes (i.e., N1 and N2) becoming less negative and the
positive amplitude (i.e., P1) becoming less positive. P1 amplitude exhibited a nonlinear shape
with the greatest changes in the first months of infancy. In contrast, the rates of change in N1 and
N2 amplitudes (decreasing) were relatively constant over infancy, suggesting protracted change.
220
An Individual's Curve is Stable Across Development
We leveraged the longitudinal study design to characterize the stability of each
individual's growth centiles. For a normative growth curve, most individuals should stay near or
within their growth centile across timepoints. Growth curve stability is frequently quantified by
binning centile lines at the 5th, 10th, 25th, 50th, 75th, 90th, and 95th percentiles, since volatility
is attenuated at the fringes of the growth curves. Centiles for all VEP features were generally
stable, with 75% of participants shifting no more than one centile line across the measured period
(Fig. 3). The mean centile change per timepoint across all features was .75 lines (SD=0.62), or
14.45 centiles (SD=10.68). We observed no significant differences in stability across VEP
features ($F(5, 1272)$=.87, p=.50), suggesting that features were consistent in their stability. We
explored centile change as a function of age by fitting a locally estimated scatterplot smoothing
(LOESS) curve to the relationship between age and centile change per day (Fig. S29). The point
of greatest change was at the earliest age for all features. There was a consistent decrease in
volatility across the first year of development.
235

236
237 **Figure 3:** *Stability of Individual VEP Centile Scores by Feature.* Violin plot representing the number of centile lines
238 (located at the 5th, 10th, 25th, 50th, 75th, 90th, and 95th percentiles) crossed by each individual per observation.
239 75% of participants shifted less than one centile line, indicating that most individuals remained near the same centile
240 across infancy. No significant differences in stability across feature were observed.
241

Assessing Utility of VEP Growth Charts: Associations with Cognitive Development and
Early Exposures
Individual VEP Scores are Associated with Cognitive Development
We investigated whether individual differences in VEP development were associated
with a globally scalable measure of neurodevelopment and emerging cognition designed for
public health contexts, the Global Scales for Early Development (Long Form; GSED-LF),
developed and validated by the WHO[70–72]. We tested whether deviation from the VEP GAMLSS
curves related to the GSED-LF age-adjusted D (DAZ) scores measured at 14 months (mean
age=429±30.66 days) in the South Africa site. As the VEP waveform includes six distinct
features, each with a unique growth trajectory, we integrated the information contained within
each feature into a composite deviation score per infant. We extracted z-scores for every
observation and feature for each individual. These z-scores measure an individual's deviation
from the normative curve for that feature. We then summed all z-scores to derive an individual's
total deviation score. Using the models trained on the full dataset across all three sites, the VEP
deviation scores were positively associated with GSED scores ($r(89)=.27$, $p=.0089$; Fig. 4A).
This association remained when correcting for potential covariates of age at GSED collection

and sex (r(89)=.22, p=.036), as well as percent VEP trials excluded during EEG quality control
at each VEP observation (r(89)=.27, p=.012). To test the generalizability of this association, we
applied the GAMLSS model trained in the Brazil and US sites to the held-out South Africa site
to derive total deviation scores. Deviation scores were nearly identical to those derived from the
model trained on all three sites (all r's>.93, p<.001). Accordingly, the deviation scores were still
associated with GSED scores (r(89)=.28, p=.0072). These results further demonstrate the ability
of the VEP growth curves to generalize across populations.
We conducted several *post hoc* analyses to better understand the association between
growth curve deviations and neurocognitive development. We separated the total deviation
scores by VEP latency and amplitude. Deviations from the latency feature curves were
significantly associated with GSED scores (r(89)=.30; p=.0036), while amplitude deviations
were not (r(89)=.090; p=.40), suggesting that the latency deviations were driving the association
with GSED scores (Fig. 4B). We also separated the total deviation scores by age based on the
target VEP observation timepoints. Deviations at a single age group were not associated with
cognitive outcomes measured by the GSED (r's<.12, p's>.12), highlighting the improved power
by integrating longitudinal data (Fig. 4C). This remained true when only using the latency
measurements (r's<.094, p's>.23). Associations between all individual feature measurements
and GSED scores are shown in Fig. S31-33.
Prenatal Alcohol Exposure is Associated with Deviations from Normative VEP Development
Finally, to test the sensitivity of VEP normative growth curve centiles to known
environmental influences, we have replicated results from a prior study with the South African
site that used conventional VEP analyses and a subset of the current sample. Specifically, we
have previously shown that prenatal alcohol exposure during the first trimester was associated
with longer P1 latencies in the first 6 months of life that neurally resolve by the end of the first
year[51]. Here, we extracted P1 latency centile scores in the Khula South Africa dataset to replicate
this finding. Corrected P1 latency centile scores for infants under 6 months positively associated
with drinks per week during the first trimester (r(131)=.43, p<.001). Similarly to the original
analysis, this effect was not observed at later ages (r(130)=-.022, p=.80). Results were identical
when using the models with the South Africa site data held out of the training process
(r(130)=.43, p<.001). This replication demonstrates the ability of centile curves to detect
developmentally-sensitive effects of environmental influences, including those relevant to public
health efforts, even in held-out site data.
290

Figure 4: *Associations of VEP Deviation scores and Cognitive Outcomes.* Scatter plots representing associations between deviation composite scores and Global Scales for Early Development Long Form (GSED-LF) age-adjusted D (DAZ) scores measured at 14 months. A) Deviation scores across all features and measurements; B) Deviation composites separated by amplitude *vs.* latency; C) Deviations composites separated by age.

Discussion

We have created and validated brain function growth charts in awake infants over the first two years of life. In contrast to other extant brain charts[13,18,21], we have used task activity and *a priori* harmonized, longitudinal data, demonstrating the feasibility of this approach to understand brain development at a global scale. In keeping with established practice for physical growth charts that single out limited, early periods of rapid growth[1-5], we focused on the narrow but critical age range of the first two postnatal years to rigorously map the substantial brain changes underlying visual function during this period[58,73,74]. We targeted the visual-evoked potential (VEP) index of sensory brain function, which is neurobiologically richly-characterized, clinically relevant, and widely collected[29-32,40-43,51,51-57]. The growth charts generalized well across sites with distinct sociocultural and economic profiles from three continents. Deviations from the normative growth models associated with both environmental influences (prenatal alcohol exposure) and postnatal cognitive scores on a global WHO-validated public health measure, demonstrating growth charts' utility to understand individual neurodevelopment and potential to

integrate with early identification and intervention efforts. Overall, our brain function growth
charts represent a new direction for EEG research and have broad impacts in the following ways.

Generalized reference growth charts for early brain function

We prioritized scalability and generalizability across global contexts with diverse
populations in this investigation so that our growth charts may serve as reference models for the
broader community of neurodevelopmental researchers. Most of our participants live in global
majority contexts (e.g., global South), which are underrepresented in neuroscience research[75–79],
including functional brain chart efforts[21,22]. Critically, the three sites have sufficient age range
overlap and sample size to distinguish between site and age effects. Overall, this diversity
provides a thorough test of the generalizability of our models. Other researchers can leverage
these models to estimate centiles for each VEP feature from smaller cohorts, special populations,
or after new data collection efforts without creating new growth charts. These centiles scores can
be combined using our composite deviation score approach to increase power in downstream
analyses. To facilitate these efforts, we integrated these growth charts directly into HAPPE—an
open-source, user-friendly software for processing EEG data used in the present study[80,81]. We
also make VEP stimuli, paradigm scripts, and processing parameter files available so researchers
can replicate the VEP procedure and processing precisely and seamlessly progress within
HAPPE from raw EEG data to growth-chart-derived centile scores for new data. The cost-
effectiveness, tolerance, and scalability of EEG relative to other early life neuroimaging
approaches further make this a promising approach to expand early neurodevelopmental research
efforts.

Insights into Early Visual Functional Neurodevelopment

The six VEP features showed divergent developmental patterns that invite future
mechanistic work. The temporal pattern of changes differed between amplitude and latency
features, with peak amplitudes exhibiting gentle slopes over infancy, while latencies show
distinct nonlinear decreases. This pattern is consistent with separable underlying developmental
mechanisms. Specifically, amplitude changes reflect postsynaptic dynamics in local circuits,
including the balance of excitatory and inhibitory neurotransmission (E/I balance), which
changes over infancy[58,65,66]. Synaptogenesis rates also increase in primary visual cortex during
this infant window and impact local circuit dynamics[82,83]. VEP amplitude changes may provide
functional indices of the interplay between rapid synaptogenesis and changing E/I balance within
the primary visual cortex. In contrast, latency reductions reflect myelination of the geniculate-
occipital pathways[68,69,84,85], and MRI-derived myelin-sensitive measures have been related to
infant VEP latency in a subsample of the study participants previously[51]. Here we show these
latency changes largely resolve within the first year.

With respect to the different VEP components, the slopes of the N1 component peak
amplitude and latency are largely flat within the featured age range, although the N1 latency
exhibits a very gentle negative slope before postnatal day 200. This pattern suggests a very early
and brief window of postnatal development for the N1 component within the parvocellular
pathway[58]. In contrast, the P1 and N2 features showed significant developmental changes over

the infant period, consistent with prior studies of visual function across imaging modalities[86].
Notably, the shallower slope of P1 amplitude change and earlier stabilization of the P1 latency
relative to the N2 amplitude and latency patterns is consistent with earlier functional and
structural maturation of the magnocellular relative to the parvocellular visual pathway in infancy
as others have posited[60,61]. Further research is required to precisely link the underlying
neurobiological mechanisms to observed developmental trajectories in these VEP growth charts.
All features stabilized by 2 years, supporting our targeted focus on this developmental window
(though protracted changes at smaller scales occur through childhood).
Additionally, we found empirical support for the role of early, basic visual
neurodevelopment in scaffolding higher-level cognition. Though prior research has linked infant
visual function behaviorally (e.g., visual attention) with cognition[87–89], studies testing early visual
brain function directly through the VEP have been inconclusive with respect to whether these
early functional properties relate to cognition[41,90,91]. We found that deviations from normative
VEP development were associated with a globally-validated public health index of cognition (the
Global Scales of Early Development Long Form). This finding in the context of prior studies
suggests that it is the trajectory of VEP development rather than discrete features or timepoints
that relate to emerging cognition. Alterations in timing (i.e., precocious or delayed maturation) of
the neurobiological processes underlying VEP development can have long-term consequences
for behavior[45,92–96]. For example, alterations in the timing of sensory neurodevelopment are
implicated in disorders such as Rett Syndrome[45,92,94,95,97], Down Syndrome[92,95], and Autism
Spectrum Disorders[45,92,95]. Our results using normative modeling highlight the power of this
approach to uncover early brain-behavior relations at the individual level during the infant period
when early intervention support is most efficacious.
Methodological Advances
Methodologically, we make several contributions to mapping normative growth charts in
neuroimaging data. First, we borrowed modeling approaches from machine learning, excluding
subsets of data (e.g., a site) during initial fitting to improve generalizability. Training and
evaluating models in disparate populations prevents overfitting[98]. Our unique dataset with
overlapping age ranges and *a priori* harmonization but distinct cultural, linguistic, geographic,
and socioeconomic contexts facilitates this testing. Second, our longitudinal data allowed us to
estimate centile stability by tracking changes within individuals following the existing literature.
Our curves demonstrated better individual centile stability than other developmental growth
curves, like weight[99]. In contrast, previous work with mainly cross-section neuroimaging data
reported the linearly interpolated interquartile range, which discards individual level data by
averaging[13]. Finally, we combined centiles from multiple timepoints and features to improve
effect sizes with developmental outcomes. Simple multivariate approaches improve effect sizes
beyond individual univariate features[100,101]. Notably, no single feature or timepoint was
associated with cognitive outcomes on its own. Future work can assess how additional composite
score calculations may further increase power across brain growth charts.
Limitations

Unlike other brain growth charts, but in keeping with established physical growth charts
like weight and height, we intentionally used a narrow age range (57–579 days) within the first
two postnatal years. The visual cortex most rapidly changes structurally and functionally during
this period[58,102,103]. Our results reinforce this premise, as all VEP features significantly slowed
rates of change by the end of the second year of life. Nevertheless, changes in the VEP
waveform also occur before 57 days and are not captured in our models. Our sample size
(n=1374 observations) might appear limited compared to the quantity suggested by previous
simulations[34]. However, we use a narrow age range (about one-fiftieth to one-third in size), task
activity, and longitudinal data, which would all modify simulation estimates downwards. For
example, growth charts using longitudinal data require much less data than those with cross-
sectional data[15,17]. Together, these design choices increase the generalizability of our curves when
using smaller samples. Though we left sites out when building our growth curves, our models
were not fully externally validated. Lastly, while we demonstrate these growth charts' utility
through associations with prenatal exposures and postnatal outcomes, additional research is
needed to assess widerspread utility, sensitivity in early life, and associations with longer-term
neurocognitive outcomes. Integration of these growth charts into HAPPE facilitates such wider
future testing.
408
Conclusion
We demonstrate the feasibility and utility of task-derived brain function growth charts
using the EEG-derived visual-evoked potential in awake infants from cohorts spanning three
continents. The growth charts exhibit high levels of fit, stability within individual trajectories,
and generalize well across different global cultural, linguistic, and socioeconomic contexts.
These normative models are sensitive to environmental factors and emerging cognitive
development and increase power to discover new results. Overall, we demonstrate the potential
of the brain function growth charts in awake infants as an impactful tool to study
neurodevelopment at scale.
418

Methods

Participants

All study procedures were approved by the relevant university Health Research Ethics

Committees (Protocol Nos.: Khula - HREC REF 666/2021, Germina - CAAE

49671221.2.0000.0068, LABS - P00039731, GABA - P00028129). Informed consent was

collected from caregivers on behalf of their infants. Cohort characteristics are summarized in

Table 1, and demographic information by cohort in Table S1.

Cohort Descriptions

Khula Cohort from South Africa. Infants from the Khula South Africa cohort[49] were

recruited from the Gugulethu Midwife Obstetrics Unit, an antenatal clinic in an informal

settlement, Gugulethu, in the Western Cape province of South Africa, as part of the Wellcome

LEAP 1kD project. Data collection occurred at the University of Cape Town's Neuroscience

Institute. 97.46% of caregivers from this site spoke Xhosa as their primary language. 82.30% of

caregivers included in this analysis were recruited prenatally, with the remaining 17.70% being

recruited shortly after birth. Women were eligible to be enrolled in the study if they were (i)

pregnant, (ii) in their third trimester of pregnancy (28-36 weeks) or up to three months

postpartum, and (iii) over the age of 18 years at the time of recruitment. Inclusion criteria for the

study included (a) singleton pregnancy, (b) no psychotropic drug endorsement during pregnancy,

(c) no infant congenital malformation or abnormalities (e.g., spina bifida, Down's syndrome),

and (d) no significant delivery complications (e.g., uterine rupture, birth asphyxia). Infants from

this cohort contributed Visual-Evoked Potential EEG, cognitive outcome (i.e., GSED-LF), and

prenatal alcohol exposure data to this paper.

Germina Cohort from Brazil. Infants from the Germina cohort[48] were recruited from

São Paulo, Brazil, as part of the Wellcome LEAP 1kD project. Data collection occurred at the

University of São Paulo Medical School. 100% of caregivers from this site spoke Brazilian

Portuguese, and assessments were administered in Brazilian Portuguese. Infants included in this

analysis were recruited postnatally, with the average age at recruitment being 2.4 months.

Inclusion criteria were (a) maternal age of 20-45 years, (b) infant age of 3 months 0 days to 3

months and 29 days, (c) gestational age of >37 weeks, (d) infant birth weight >2000 grams, (e)

no substance endorsement during pregnancy, (f) no history of severe maternal mental disorders

(e.g., psychosis, bipolar disorder), (g) no delivery complications requiring medical intervention

(e.g., perinatal asphyxia, shoulder dystocia, excessive bleeding), (h) no infant genetic syndrome

or auditory/visual impairment diagnoses, and (i) availability for in-person lab visits. Infants from

this cohort contributed visual-evoked potential EEG data to this paper.

LABS Cohort from USA. Infants from the LABS cohort were recruited from the greater

Boston area in the United States of America as part of the Wellcome LEAP 1kD project. Data

collection occurred at Boston Children's Hospital. Based on inclusion criteria, English was the

primary language for all caregivers from this site. Infants included in this analysis were recruited

postnatally. The average age at recruitment varies by wave, with waves beginning at 3 months

(+/- 7 days), 6 months (+/- 14 days), 12 months (+/- 14 days), 18 months (+/- 28 days), and 24

months (+/- 42 days). Inclusion criteria were (a) English was the primary language spoken at
home (greater than 75% of the language spoken), (b) parental age was 18 years or older, (c)
gestational age of 36-40 weeks, (d) infant birth weight of 2500-4500 grams, (e) no prenatal,
postnatal, or birth-related complications, (f) no diagnosis of genetic, metabolic, syndromic, or
neurological disorders, (g) no hearing or vision problems (unless corrected). Infants from this
cohort contributed visual-evoked potential EEG data to this paper.

GABA Cohort from USA. Infants from the GABA cohort[47] were recruited from the
greater Boston area in the United States of America as part of a larger study investigating the
effects of early general anesthesia on brain development, and data collection occurred at Boston
Children's Hospital. 92.06% of caregivers from this site reported speaking English with their
child. All infants included in this analysis were recruited postnatally. Inclusion criteria included
(a) parents/legal guardians ≥18 years old without developmental disabilities, without being wards
of the state and willing to complete all study visits, (b) infant age ≤280 postnatal days, (c)
gestational age of ≥32 weeks, (d) lack of confirmed or suspected clinical seizures, (e) lack of
confirmed or suspected auditory or visual impairment, (f) no diagnosis of a genetic disorder
affecting development, (g) no ongoing medical condition that would interfere with the conduct
and assessment of the study, and (h) fluent English speakers, in which English is spoken to the
infant ≥75% awake hours. Infants recruited for the comparison group without general anesthetic
exposure or hospital admittance and who opted to have their data used in future studies were
included in this analysis. Infants from this cohort contributed visual-evoked potential EEG data
to this paper.
480
Measures
Electroencephalography (EEG)
EEG Acquisition. At all sites, EEG data were acquired from infants seated in their
caregiver's lap in a dimly lit, quiet room. A 128-channel high-density HydroCel Geodesic Sensor
Net (EGI, Eugene, OR), amplified with a NetAmps 400 high-input amplifier (GABA cohort used
the 300 series) and recorded via an Electrical Geodesics, Inc. (EGI, Eugene, OR) system was
used. A 1000 Hz sampling rate was used at the South Africa and U.S.A. sites. A 500 Hz
sampling rate was used at the Brazil site. EEG data were online referenced to the vertex (channel
Cz) through the EGI Netstation software. Impedances were kept below 100KΩ following the
impedance capabilities of the high-impedance amplifiers. For the Khula cohort, Geodesic Sensor
Nets with modified tall pedestals designed to improve the inclusion of infants with
thick/curly/tall hair were used as needed across participants. Shea moisture leave-in castor oil
conditioner with insulating ingredients was applied to hair across the scalp before net placement
to improve impedances and participant comfort[104]. At all sites, the Visual-Evoked Potential
(VEP) task was presented using Eprime software (Psychology Software Tools, Pittsburgh, PA)
on a desktop computer with an external monitor ranging from 19.5 to 21.5 inches on the diagonal
facing the infant. The monitor was approximately 60-65 cm away from the infant. A standard
phase-reversal VEP was induced with a black and white checkerboard (1cm x 1 cm squares

within the board) stimulus that alternated presentation (black squares became white, white
squares became black) every 500 milliseconds for a total of 100 trials (up to 200 trials in GABA
cohort). Participant looking was monitored by video and by an assistant throughout data
collection. The task was rerun if the participant looked away during the VEP task or was overly
fussy. For the GABA cohort, stimulus presentation proceeded if a Tobii Eye Tracker system
registered the infant's gaze on the monitor or the experimenter could see the infant looking at the
screen. For the Germina cohort, video recording of the experiment was manually reviewed.
Periods when the infant looked away or was overly fussy were removed from the data.
EEG Data Pre-Processing. EEG data were exported from native Netstation .mff format
to .raw format and then pre-processed using HAPPE v3.3 software (HAPPE v3.2 used for pre-
processing in GABA, but updates between versions do not affect ERP measures), an automated
open-source EEG processing software validated for infant data[80]. A subset of the 128 channels
was selected for pre-processing, which excluded the rim electrodes as these are typically artifact-
laden (channels excluded from pre-processing are included in Table S2). The HAPPE Pre-
Processing pipeline was run with user-selected specifications outlined in Table S3. Parameters
were largely consistent across sites except for line noise frequency, which differed by geographic
region.
Pre-processed EEG data were considered usable and moved forward to VEP extraction if
HAPPE pre-processing ran successfully, at least 20 trials were retained following bad trial
rejection, and at least two good channels were kept within the visual ROI.
VEP Extraction. VEP waveforms were extracted and quantified using the HAPPE v3.3
Generate ERPs script from the HAPPE+ER pipeline[81]. Electrodes in the occipital region were
selected as a region of interest (i.e., E70, E71, E75, E76, E83). The VEP waveform has three
main components to be quantified: a negative N1 peak, a positive P1 peak, and a negative N2
peak. The features of interest for these components were peak latency and amplitude.
Developmentally appropriate windows were selected for each age group with an overall N1
window range of 40 to 120 milliseconds, an overall P1 window range of 75 to 175 milliseconds,
and an overall N2 window range of 100 to 325 milliseconds. Parameters (which were consistent
across sites) used in extracting the ERPs with HAPPE+ER are summarized in Table S3.
All VEPs were visually inspected to ensure the automatically extracted values were
correct and adjusted if observable peaks occurred outside the automated window bounds. Data
quality for included files at each site is summarized in Table S4.
Cognitive Outcomes
Global Scales for Early Development, Long Form (GSED-LF). To investigate how
deviation from the GAMLSS curves relates to emerging behavior, we collected the Global Scales
for Early Development, Long Form (GSED-LF) from infants in the Khula cohort around age 428
days (*SD*=36.61). The GSED Long Form (GSED-LF) is a directly administered tool developed
by the World Health Organization to collect population-level behavioral development at scale
globally across diverse cultural and linguistic contexts[70–72]. The GSED-LF provides an overall

"d-score" that represents overall cognitive, communication, and motor functioning holistically.
The standardized daz-score was used as our cognitive outcome.

Prenatal Alcohol Exposure

As previously described[51], mothers in the Khula South Africa cohort self-reported alcohol
endorsement during pregnancy using the Alcohol Exposure Questionnaire (AEQ)[105] at the first
research visit. We extracted the number of standardized alcohol units consumed per week in each
trimester from the AEQ (number of drinking episodes per week in each trimester multiplied by
number of drinks consumed per episode). Mothers also self-reported non-alcohol substance
endorsement (i.e., tobacco, cannabis, cocaine, amphetamines, inhalants, sedatives/sleeping pills,
hallucinogens, opioids, or other) in the three months before enrollment using the World Health
Organization Alcohol, Smoking, and Substance Involvement Screening Test [106]. Infants with
prenatal alcohol *and* non-alcohol substance exposures were excluded from the analysis to isolate
the effects of alcohol[51,107,108]. 46 infants provided usable data at 112 days ($SD = 26.86$ days)
whose mothers reported use of alcohol but no other comorbid substances.

Analyses

Generalized Additive Models for Location, Scale, and Shape

Growth curves were modeled using generalized additive models for location, scale, and
shape (GAMLSS), a framework developed to model complex data distributions[50]. GAMLSS is
an extension of generalized additive models that allows independent functions to fit up to four
parameters of the response distribution. As a result, myriad distribution families can be captured
by modeling the mean, standard deviation, skewness, and kurtosis. GAMLSS is the World
Health Organization's recommended method for developing growth curves due to its flexibility
in modeling the relationship between age and higher-order moments[4].

GAMLSS models were fit to model the relationship between age and individual VEP
features. We used p-splines as the smoothing function for each parameter. Distributions were
chosen for each feature by minimizing the total validation Bayesian Information Criterion (BIC)
when training the model in two datasets and validating in the third held-out dataset. We
compared the standard forms of the following 3-4 parameter distributions: generalized *t* (GT),
power exponential (PE), power exponential 2 (PE2), Sinh-Arcsinh (SHASH), *t*-family (TF and
TF2), Box-Cox Cole and Green (BCCGo), Box-Cox power exponential (BCPEo), Box-Cox *t*
(BCTo), gamma family (GAF), generalized beta type 2 (GB2), generalized gamma (GG),
Johnson's S_U (JSUo), generalized inverse gamma (GIG), skew normal types 1 and 2 (SN1 and
SN2), skew exponential power type 1 (SEP1), and skew *t* (ST1). Degrees of freedom were
optimized using maximum likelihood during the distribution comparison. With the distributions
chosen for each feature, we selected the optimal generalized Akaike criterion penalty *k* using the
same process. All models were fitted using up to 30 iterations of the Rigby-Stasinopoulos (RS)[109]
algorithim and up to 20 iterations of the Cole-Green (CG)[110] algorithm. To evaluate biological
sex as a covariate, we refit models with all features with sex included as an additional smoothing

term for the μ term (i.e., the location or mean parameter) and compared the total validation BIC
across all dataset combinations.
Model fit was quantified within-dataset by assessing the correspondence of the centiles
predicted by the model and the actual quantiles of each data point. Specifically, we first predict
the z-scores from the training data values and then convert them to their corresponding centile
value. Using these centile scores, we predict the feature value given each age value from the
training set and compare these to the actual values from the data. We report the Pearson
correlation and root mean squared error (RMSE). Models were also assessed for fit through
worm plots and Q-statistics. To understand the change in each parameter as a function of age, we
created term plots representing *mu* (location), *sigma* (scale), *nu* (shape, skewness, or degrees of
freedom, depending on the distribution), and *tau* (kurtosis) of each model (if applicable) over age
(Fig. S14-19). Additional model details are reported in Table S8. We used numerical derivation
to identify peaks and troughs by predicting each term over the range of ages and by predicting
the 50th percentile for each feature.
Centile Stability
To assess the stability of centiles within individuals, we binned centile scores into "lines"
as is typically done to measure movement on growth curves. This is done because movement at
the extreme centiles is not equivalent to movement near the center. We binned centiles at 5%,
10%, 25%, 50%, 75%, 90%, and 95%. Stability was measured in several ways based on how
many centile lines an individual crossed. The primary metric was distance per timepoint,
calculated by taking the absolute value of centile change between timepoints and dividing it by
the number of timepoints. We also measured the variance, standard deviation, and range.
Associations with Cognitive Outcomes
We associated deviation from the GAMLSS curves associated with the GSED-LF z-
scores measured at 14 months in the South Africa site. The z-scores from each feature were
summed to generate composite deviation scores for each subject. This analysis was restricted to
the South Africa site, as only these participants had available the GSED-LF outcome measures.
Participants were only included if they had viable data for all three sessions (*n*=90). Associations
between deviation scores and cognitive outcomes were quantified with Pearson's *r* correlation.
We used partial correlations to assess the impact of potential covariates, including sex, age at
VEP measurement, and percent frames removed during EEG preprocessing. We also fit a linear
regression model predicting cognitive outcomes from VEP features to determine the relative
influence of each feature. We used a 1000-fold bootstrap to determine the included features and
coefficients of the optimal model with confidence intervals. While the R^2 of the resultant model
was high, the results were unstable due to the limited sample size (Fig. S34).
Associations with Prenatal Alcohol Exposure
Finally, we show that deviation scores can replicate results from conventional analyses.
We associated increased prenatal alcohol exposure during the first trimester with longer P1
latencies in the first 6 months of life. Because P1 is preceded by N1, P1 latency is commonly
corrected by subtracting N1 latency. We followed this approach with the z-scores to create

corrected P1 latency z-scores. Associations between these scores and alcohol consumption were

assessed by Pearson's *r* correlation.

Data Availability

The EEG and behavioral data are not publicly available due to ethical concerns. Access to the

data is subject to a data sharing agreement as described in the original data papers [47–49]. The

GAMLSS curves are publicly available through the HAPPE software and can be found at

https://github.com/PINE-Lab/HAPPE.

Code Availability

HAPPE can be found at https://github.com/PINE-Lab/HAPPE. The GAMLSS R package can be

found at https://cran.r-project.org/web/packages/gamlss/index.html. Other associated code can be

found at https://github.com/chrisclaycamp/VEP-GAMLSS.

References

1. de Onis, M., Wijnhoven, T. M. A., & Onyango, A. W. (2004). Worldwide practices in child growth monitoring. *The Journal of Pediatrics*, *144*(4), 461–465. https://doi.org/10.1016/j.jpeds.2003.12.034

2. Kuczmarski, R. J., Ogden, C. L., Guo, S. S., Grummer-Strawn, L. M., Flegal, K. M., Mei, Z., Wei, R., Curtin, L. R., Roche, A. F., & Johnson, C. L. (2002). *2000 CDC growth charts for the United States: Methods and development* (11(246); Vital Health Statistics). National Center for Health Statistics.

3. Thompson, L. A., & Moreno, M. A. (2018). Growth and Growth Charts in Children. *JAMA Pediatrics*, *172*(6), 604. https://doi.org/10.1001/jamapediatrics.2018.0601

4. WHO Multicentre Growth Reference Study Group, & de Onis, M. (2006). WHO Child Growth Standards based on length/height, weight and age. *Acta Paediatrica*, *95*(S450), 76–85. https://doi.org/10.1111/j.1651-2227.2006.tb02378.x

5. World Health Organization. (1978). *A Growth Chart for International Use in Maternal and Child Health Care: Guidelines for Primary Health Care Personnel*. https://iris.who.int/bitstream/handle/10665/37294/9241541296.pdf?sequence=1

6. Dustman, R. E., Emmerson, R. Y., & Shearer, D. E. (1996). Life Span Changes in Electrophysiological Measures of Inhibition. *Brain and Cognition*, *30*(1), 109–126. https://doi.org/10.1006/brcg.1996.0007

7. Giedd, J. N., Raznahan, A., Alexander-Bloch, A., Schmitt, E., Gogtay, N., & Rapoport, J. L. (2015). Child Psychiatry Branch of the National Institute of Mental Health Longitudinal Structural Magnetic Resonance Imaging Study of Human Brain Development. *Neuropsychopharmacology*, *40*(1), 43–49. https://doi.org/10.1038/npp.2014.236

8. Hedman, A. M., van Haren, N. E. M., Schnack, H. G., Kahn, R. S., & Hulshoff Pol, H. E. (2012). Human brain changes across the life span: A review of 56 longitudinal magnetic resonance imaging studies. *Human Brain Mapping*, *33*(8), 1987–2002. https://doi.org/10.1002/hbm.21334

9. Huang, H., Shu, N., Mishra, V., Jeon, T., Chalak, L., Wang, Z. J., Rollins, N., Gong, G., Cheng, H., Peng, Y., Dong, Q., & He, Y. (2015). Development of Human Brain Structural Networks Through Infancy and Childhood. *Cerebral Cortex*, *25*(5), 1389–1404. https://doi.org/10.1093/cercor/bht335

10. Mueller, V., Brehmer, Y., von Oertzen, T., Li, S.-C., & Lindenberger, U. (2008). Electrophysiological correlates of selective attention: A lifespan comparison. *BMC Neuroscience*, *9*(1), 18. https://doi.org/10.1186/1471-2202-9-18

11. Murphy, K. M., & Monteiro, L. (2024). Anatomical and molecular development of the human primary visual cortex. *Frontiers in Cellular Neuroscience*, *18*. https://doi.org/10.3389/fncel.2024.1427515

12. Wierenga, L. M., van den Heuvel, M. P., van Dijk, S., Rijks, Y., de Reus, M. A., & Durston, S. (2016). The development of brain network architecture. *Human Brain Mapping*, *37*(2), 717–729. https://doi.org/10.1002/hbm.23062

13. Bethlehem, R. a. I., Seidlitz, J., White, S. R., Vogel, J. W., Anderson, K. M., Adamson, C., Adler, S., Alexopoulos, G. S., Anagnostou, E., Areces-Gonzalez, A., Astle, D. E., Auyeung, B., Ayub, M., Bae, J., Ball, G., Baron-Cohen, S., Beare, R., Bedford, S. A., Benegal, V., … Alexander-Bloch, A. F. (2022). Brain charts for the human lifespan. *Nature*, *604*(7906), 525–533. https://doi.org/10.1038/s41586-022-04554-y

14. Chen, L.-Z., Holmes, A. J., Zuo, X.-N., & Dong, Q. (2021). Neuroimaging brain growth charts: A road to mental health. *Psychoradiology*, *1*(4), 272–286. https://doi.org/10.1093/psyrad/kkab022

15. Di Biase, M. A., Tian, Y. E., Bethlehem, R. A. I., Seidlitz, J., Alexander-Bloch, Aaron. F., Yeo, B. T. T., & Zalesky, A. (2023). Mapping human brain charts cross-sectionally and longitudinally. *Proceedings of the National Academy of Sciences*, *120*(20), e2216798120. https://doi.org/10.1073/pnas.2216798120

16. Rutherford, S., Kia, S. M., Wolfers, T., Fraza, C., Zabihi, M., Dinga, R., Berthet, P., Worker, A., Verdi, S., Ruhe, H. G., Beckmann, C. F., & Marquand, A. F. (2022). The normative modeling framework for computational psychiatry. *Nature Protocols*, *17*(7), 1711–1734. https://doi.org/10.1038/s41596-022-00696-5

17. Zhou, Z.-X., Chen, L.-Z., Milham, M. P., Zuo, X.-N., & Lifespan Brain Chart Consortium (LBCC). (2023). Six cornerstones for translational brain charts. *Science Bulletin*, *68*(8), 795–799. https://doi.org/10.1016/j.scib.2023.03.047

18. Alex, A. M., Aguate, F., Botteron, K., Buss, C., Chong, Y.-S., Dager, S. R., Donald, K. A., Entringer, S., Fair, D. A., Fortier, M. V., Gaab, N., Gilmore, J. H., Girault, J. B., Graham, A. M., Groenewold, N. A., Hazlett, H., Lin, W., Meaney, M. J., Piven, J., … Knickmeyer, R. C. (2024). A global multicohort study to map subcortical brain development and cognition in infancy and early childhood. *Nature Neuroscience*, *27*(1), 176–186. https://doi.org/10.1038/s41593-023-01501-6

19. World Health Organization. (2022). *Optimizing brain health across the life course: WHO Position Paper*. https://iris.who.int/bitstream/handle/10665/361251/9789240054561-eng.pdf?sequence=1

20. Rutherford, S., Fraza, C., Dinga, R., Kia, S. M., Wolfers, T., Zabihi, M., Berthet, P., Worker, A., Verdi, S., Andrews, D., Han, L. K., Bayer, J. M., Dazzan, P., McGuire, P., Mocking, R. T., Schene, A., Sripada, C., Tso, I. F., Duval, E. R., … Marquand, A. F. (2022). Charting brain growth and aging at high spatial precision. *eLife*, *11*, e72904. https://doi.org/10.7554/eLife.72904

21. Iyer, K. K., Roberts, J. A., Waak, M., Vogrin, S. J., Kevat, A., Chawla, J., Haataja, L. M., Lauronen, L., Vanhatalo, S., & Stevenson, N. J. (2024). A growth chart of brain function from infancy to adolescence based on EEG. *eBioMedicine*, *102*. https://doi.org/10.1016/j.ebiom.2024.105061

22. Rutherford, S., Barkema, P., Tso, I. F., Sripada, C., Beckmann, C. F., Ruhe, H. G., & Marquand, A. F. (2023). Evidence for embracing normative modeling. *eLife*, *12*, e85082. https://doi.org/10.7554/eLife.85082

23. Sun, L., Zhao, T., Liang, X., Xia, M., Li, Q., Liao, X., Gong, G., Wang, Q., Pang, C., Yu, Q., Bi, Y., Chen, P., Chen, R., Chen, Y., Chen, T., Cheng, J., Cheng, Y., Cui, Z., Dai, Z., … He, Y. (2025). Human lifespan changes in the brain's functional connectome. *Nature Neuroscience*, *28*(4), 891–901. https://doi.org/10.1038/s41593-025-01907-4

24. Greene, A. S., Gao, S., Scheinost, D., & Constable, R. T. (2018). Task-induced brain state manipulation improves prediction of individual traits. *Nature Communications*, *9*(1), 2807. https://doi.org/10.1038/s41467-018-04920-3

25. Rosenberg, M. D., & Finn, E. S. (2022). How to establish robust brain–behavior relationships without thousands of individuals. *Nature Neuroscience*, *25*(7), 835–837. https://doi.org/10.1038/s41593-022-01110-9

26. Barry-Anwar, R., Riggins, T., & Scott, L. S. (2020). Electrophysiology in developmental populations: Key methods and findings. In K. Cohen Kadosh (Ed.), *The Oxford Handbook of Developmental Cognitive Neuroscience*. Oxford University Press. https://doi.org/10.1093/oxfordhb/9780198827474.013.3

27. Finn, E. S. (2021). Is it time to put rest to rest? *Trends in Cognitive Sciences*, *25*(12), 1021–1032. https://doi.org/10.1016/j.tics.2021.09.005

28. Lopez, K. L., Monachino, A. D., Vincent, K. M., Peck, F. C., & Gabard-Durnam, L. J. (2023). Stability, change, and reliable individual differences in electroencephalography measures: A lifespan perspective on progress and opportunities. *NeuroImage*, *275*, 120116. https://doi.org/10.1016/j.neuroimage.2023.120116

29. Norton, E. S., MacNeill, L. A., Harriott, E. M., Allen, N., Krogh-Jespersen, S., Smyser, C. D., Rogers, C. E., Smyser, T. A., Luby, J., & Wakschlag, L. (2021). EEG/ERP as a pragmatic method to expand the reach of infant-toddler neuroimaging in HBCD: Promises and challenges. *Developmental Cognitive Neuroscience*, *51*, 100988. https://doi.org/10.1016/j.dcn.2021.100988

30. Sokol, S. (1978). Measurement of infant visual acuity from pattern reversal evoked potentials. *Vision Research*, *18*(1), 33–39. https://doi.org/10.1016/0042-6989(78)90074-3

31. Sokol, S. (1980). Pattern Visual Evoked Potentials: Their Use in Pediatric Ophthalmology. *International Ophthalmology Clinics*, *20*(1), V.

32. Taylor, M. J., & McCulloch, D. L. (1992). Visual Evoked Potentials in Infants and Children. *Journal of Clinical Neurophysiology*, *9*(3), 357.

33. Xu, W., Monachino, A. D., McCormick, S. A., Margolis, E. T., Sobrino, A., Bosco, C., Franke, C. J., Davel, L., Zieff, M. R., Donald, K. A., Gabard-Durnam, L. J., & Morales, S. (2024). Advancing the Reporting of Pediatric EEG Data: Tools for Estimating Reliability, Effect Size, and Data Quality Metrics. *Developmental Cognitive Neuroscience*, 101458. https://doi.org/10.1016/j.dcn.2024.101458

34. Bozek, J., Griffanti, L., Lau, S., & Jenkinson, M. (2023). Normative models for neuroimaging markers: Impact of model selection, sample size and evaluation criteria. *NeuroImage*, *268*, 119864. https://doi.org/10.1016/j.neuroimage.2023.119864

35. Luby, J. L., Baram, T. Z., Rogers, C. E., & Barch, D. M. (2020). Neurodevelopmental Optimization after Early-Life Adversity: Cross-Species Studies to Elucidate Sensitive Periods and Brain Mechanisms to Inform Early Intervention. *Trends in Neurosciences*, *43*(10), 744–751. https://doi.org/10.1016/j.tins.2020.08.001

36. McLaughlin, K. A., & Gabard-Durnam, L. (2022). Experience-driven plasticity and the emergence of psychopathology: A mechanistic framework integrating development and the environment into the Research Domain Criteria (RDoC) model. *Journal of Psychopathology and Clinical Science*, *131*(6), 575–587. https://doi.org/10.1037/abn0000598

37. Morishita, H., & Hensch, T. K. (2008). Critical period revisited: Impact on vision. *Current Opinion in Neurobiology*, *18*(1), 101–107. https://doi.org/10.1016/j.conb.2008.05.009

38. Zwaigenbaum, L., Bauman, M. L., Choueiri, R., Kasari, C., Carter, A., Granpeesheh, D., Mailloux, Z., Smith Roley, S., Wagner, S., Fein, D., Pierce, K., Buie, T., Davis, P. A., Newschaffer, C., Robins, D., Wetherby, A., Stone, W. L., Yirmiya, N., Estes, A., … Natowicz, M. R. (2015). Early Intervention for Children With Autism Spectrum Disorder Under 3 Years of Age: Recommendations for Practice and Research. *Pediatrics*, *136*(Supplement_1), S60–S81. https://doi.org/10.1542/peds.2014-3667E

39. WHO Multicentre Growth Reference Study Group. (2006). *WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age*. World Health Organization. https://www.who.int/publications/i/item/924154693X

40. Hoyt, C. S. (1984). The Clinical Usefulness of the Visual Evoked Response. *Journal of Pediatric Ophthalmology & Strabismus*, *21*(6), 231–234. https://doi.org/10.3928/0191-3913-19841101-08

41. Jensen, S. K. G., Kumar, S., Xie, W., Tofail, F., Haque, R., Petri, W. A., & Nelson, C. A. (2019). Neural correlates of early adversity among Bangladeshi infants. *Scientific Reports*, *9*(1), 3507. https://doi.org/10.1038/s41598-019-39242-x

42. McCulloch, D. L. (2007). Visual evoked potentials in infants. In *Infant EEG and Event-Related Potentials*. Psychology Press.

43. Varcin, K. J., & Nelson, C. A. I. (2016). A developmental neuroscience approach to the search for biomarkers in autism spectrum disorder. *Current Opinion in Neurology*, *29*(2), 123. https://doi.org/10.1097/WCO.0000000000000298

44. Hensch, T. K., & Fagiolini, M. (2005). Excitatory–inhibitory balance and critical period plasticity in developing visual cortex. In *Progress in Brain Research* (Vol. 147, pp. 115–124). Elsevier. https://doi.org/10.1016/S0079-6123(04)47009-5

45. LeBlanc, J. J., & Fagiolini, M. (2011). Autism: A "Critical Period" Disorder? *Neural Plasticity*, *2011*(1), 921680. https://doi.org/10.1155/2011/921680

46. Saby, J. N., Peters, S. U., Roberts, T. P. L., Nelson, C. A., & Marsh, E. D. (2020). Evoked Potentials and EEG Analysis in Rett Syndrome and Related Developmental

Encephalopathies: Towards a Biomarker for Translational Research. *Frontiers in Integrative Neuroscience*, *14*. https://doi.org/10.3389/fnint.2020.00030

47. Berde, C., Coffman, S., Cornelissen, L., Gabard-Durnam, L. J., Nelson, C. A., & Hensch, T. K. (2024). *General Anesthesia and Brain Activity (the GABA Study) [Study Protocol]*. OSF. https://doi.org/10.17605/OSF.IO/XS6MJ

48. Fatori, D., Shephard, E., Benette, D., Naspolini, N. F., Guzman, G. C., Wang, J. Y. T., Tótolo, P., Mafra, A. L., Isaias, C., Santos, D. P. dos, Russo, F. B., Kobayashi, G., Argeu, A., Teixeira, M., Mattiello-Sverzut, A. C., Fernandes, M. T. B., Petian-Alonso, D. C., Brentani, H., Scliar, M., … Polanczyk, G. V. (2025). Identifying biomarkers and trajectories of executive functions and language development in the first 3 years of life: Design, methods, and findings of the Germina cohort study. *Development and Psychopathology*, 1–11. https://doi.org/10.1017/S0954579425000069

49. Zieff, M. R., Miles, M., Mbale, E., Eastman, E., Ginnell, L., Williams, S. C. R., Jones, D. K., Alexander, D. C., Wijeratne, P. A., Gabard-Durnam, L. J., Klepac-Ceraj, V., Bonham, K. S., Pini, N., Sania, A., Lucchini, M., Deoni, S., Fifer, W. P., Gladstone, M., Amso, D., & Donald, K. A. (2024). Characterizing developing executive functions in the first 1000 days in South Africa and Malawi: The Khula Study [Version 1; Peer review: 2 approved with reservations]. *Wellcome Open Research*, *9*, 157. https://doi.org/10.12688/wellcomeopenres.19638.1

50. Rigby, R. A., & Stasinopoulos, D. M. (2005). Generalized Additive Models for Location, Scale and Shape. *Journal of the Royal Statistical Society Series C: Applied Statistics*, *54*(3), 507–554. https://doi.org/10.1111/j.1467-9876.2005.00510.x

51. Margolis, E. T., Davel, L., Bourke, N. J., Bosco, C., Zieff, M. R., Monachino, A. D., Mazubane, T., Williams, S. R., Miles, M., Jacobs, C. A., Williams, S., Bradford, L., Knipe, C., Madi, Z., Methola, B., Mhlakwaphalwa, T., Mlandu, N., Nkubungu, K., Goolam Nabi, Z., … Gabard-Durnam, L. J. (2024). Longitudinal effects of prenatal alcohol exposure on visual neurodevelopment over infancy. *Developmental Psychology*, *60*(9), 1673–1698. https://doi.org/10.1037/dev0001727

52. Di Russo, F., Martínez, A., Sereno, M. I., Pitzalis, S., & Hillyard, S. A. (2002). Cortical sources of the early components of the visual evoked potential. *Human Brain Mapping*, *15*(2), 95–111. https://doi.org/10.1002/hbm.10010

53. Hashimoto, T., Kashii, S., Kikuchi, M., Honda, Y., Nagamine, T., & Shibasaki, H. (1999). Temporal profile of visual evoked responses to pattern-reversal stimulation analyzed with a whole-head magnetometer. *Experimental Brain Research*, *125*(3), 375–382. https://doi.org/10.1007/s002210050693

54. Marcar, V. L., & Wolf, M. (2021). An investigation into the relationship between stimulus property, neural response and its manifestation in the visual evoked potential involving retinal resolution. *European Journal of Neuroscience*, *53*(8), 2612–2628. https://doi.org/10.1111/ejn.15112

55. Mohr, K. S., Geuzebroek, A. C., & Kelly, S. P. (2024). Visual cortical area contributions to the transient, multifocal and steady-state VEP: A forward model-informed analysis. *Imaging Neuroscience*, *2*, 1–26. https://doi.org/10.1162/imag_a_00152

56. Shigeto, H., Tobimatsu, S., Yamamoto, T., Kobayashi, T., & Kato, M. (1998). Visual evoked cortical magnetic responses to checkerboard pattern reversal stimulation: A study on the neural generators of N75, P100 and N145. *Journal of the Neurological Sciences*, *156*(2), 186–194. https://doi.org/10.1016/S0022-510X(98)00026-4

57. Tobimatsu, S. (2002). Chapter 39 Neurophysiologic tools to explore visual cognition. In R. C. Reisin, M. R. Nuwer, M. Hallett, & C. Medina (Eds.), *Supplements to Clinical Neurophysiology* (Vol. 54, pp. 261–265). Elsevier. https://doi.org/10.1016/S1567-424X(09)70459-3

58. Lippé, S., Roy, M.-S., Perchet, C., & Lassonde, M. (2007). Electrophysiological Markers of Visuocortical Development. *Cerebral Cortex*, *17*(1), 100–107. https://doi.org/10.1093/cercor/bhj130

59. Ellemberg, D., Hammarrenger, B., Lepore, F., Roy, M. S., & Guillemot, J. P. (2001). Contrast dependency of VEPs as a function of spatial frequency: The parvocellular and magnocellular contributions to human VEPs. *Spatial Vision*, *15*(1), 99–111. https://doi.org/10.1163/15685680152692042

60. Hammarrenger, B., Leporé, F., Lippé, S., Labrosse, M., Guillemot, J.-P., & Roy, M.-S. (2003). Magnocellular and parvocellular developmental course in infants during the first year of life. *Documenta Ophthalmologica*, *107*(3), 225–233. https://doi.org/10.1023/B:DOOP.0000005331.66114.05

61. Hammarrenger, B., Roy, M.-S., Ellemberg, D., Labrosse, M., Orquin, J., Lippe, S., & Lepore, F. (2007). Developmental delay and magnocellular visual pathway function in very-low-birthweight preterm infants. *Developmental Medicine & Child Neurology*, *49*(1), 28–33. https://doi.org/10.1017/S0012162207000084.x

62. Livingstone, M., & Hubel, D. (1988). Segregation of Form, Color, Movement, and Depth: Anatomy, Physiology, and Perception. *Science*, *240*(4853), 740–749. https://doi.org/10.1126/science.3283936

63. McCulloch, D. L., Orbach, H., & Skarf, B. (1999). Maturation of the pattern-reversal VEP in human infants: A theoretical framework. *Vision Research*, *39*(22), 3673–3680. https://doi.org/10.1016/S0042-6989(99)00091-7

64. Wadhwa, S., & Bijlani, V. (1988). Cytodifferentiation and developing neuronal circuitry in the human lateral geniculate nucleus. *International Journal of Developmental Neuroscience*, *6*(1), 59–75. https://doi.org/10.1016/0736-5748(88)90028-7

65. Morrone, M. C., & Burr, D. C. (1986). Evidence for the existence and development of visual inhibition in humans. *Nature*, *321*(6067), 235–237. https://doi.org/10.1038/321235a0

66. Muthukumaraswamy, S. D., Myers, J. F. M., Wilson, S. J., Nutt, D. J., Hamandi, K., Lingford-Hughes, A., & Singh, K. D. (2013). Elevating Endogenous GABA Levels with GAT-1 Blockade Modulates Evoked but Not Induced Responses in Human Visual

Cortex. *Neuropsychopharmacology*, *38*(6), 1105–1112.
https://doi.org/10.1038/npp.2013.9

67. Šuštar Habjan, M., Bach, M., van Genderen, M., Li, S., Mizota, A., Nilsson, J., Thompson, D., & Robson, A. (in press). ISCEV standard for clinical visual evoked potentials (2025 update). *Documenta Ophthalmologica*.
https://iscev.wildapricot.org/resources/Documents/StandardsEtc/VEP%20standard%202020 25.pdf

68. Dubois, J., Dehaene-Lambertz, G., Soarès, C., Cointepas, Y., Bihan, D. L., & Hertz-Pannier, L. (2008). Microstructural Correlates of Infant Functional Development: Example of the Visual Pathways. *Journal of Neuroscience*, *28*(8), 1943–1948.
https://doi.org/10.1523/JNEUROSCI.5145-07.2008

69. Heidari, M., Radcliff, A. B., McLellan, G. J., Ver Hoeve, J. N., Chan, K., Kiland, J. A., Keuler, N. S., August, B. K., Sebo, D., Field, A. S., & Duncan, I. D. (2019). Evoked potentials as a biomarker of remyelination. *Proceedings of the National Academy of Sciences*, *116*(52), 27074–27083. https://doi.org/10.1073/pnas.1906358116

70. McCray, G., McCoy, D., Kariger, P., Janus, M., Black, M. M., Chang, S. M., Tofail, F., Eekhout, I., Waldman, M., Buuren, S. van, Khanam, R., Sazawal, S., Nizar, A., Schönbeck, Y., Zongo, A., Brentani, A., Zhang, Y., Dua, T., Cavallera, V., … Gladstone, M. (2023). The creation of the Global Scales for Early Development (GSED) for children aged 0–3 years: Combining subject matter expert judgements with big data. *BMJ Global Health*, *8*(1), e009827. https://doi.org/10.1136/bmjgh-2022-009827

71. World Health Organization. (2023). *Global Scales for Early Development v1.0: Long Form (Directly Administered) Item Guide*.
https://apps.who.int/iris/bitstream/handle/10665/366269/WHO-MSD-GSEDpackage-v1.0-2023.6-eng.pdf

72. World Health Organization. (2023). *Global Scales for Early Development v1.0: Adaptation and Translation Guide*. https://apps.who.int/iris/bitstream/handle/10665/366278/WHO-MSD-GSEDpackage-v1.0-2023.9-eng.pdf

73. Eimas, P. D., & Quinn, P. C. (1994). Studies on the Formation of Perceptually Based Basic-Level Categories in Young Infants. *Child Development*, *65*(3), 903–917.
https://doi.org/10.1111/j.1467-8624.1994.tb00792.x

74. Slater, A. (Ed.). (1998). *Perceptual Development: Visual, Auditory, and Speech Perception in Infancy*. Psychology Press.

75. Chiao, J. Y., & Cheon, B. K. (2010). The weirdest brains in the world. *The Behavioral and Brain Sciences*, *33*(2–3), 88–90. https://doi.org/10.1017/S0140525X10000282

76. Duncan, N. W., & Rae, C. L. (2024). Geographical and economic influences on neuroimaging modality choice. *Royal Society Open Science*, *11*(5), 231496.
https://doi.org/10.1098/rsos.231496

77. Khan, T., Abimbola, S., Kyobutungi, C., & Pai, M. (2022). How we classify countries and people—And why it matters. *BMJ Global Health*, *7*(6), e009704. https://doi.org/10.1136/bmjgh-2022-009704

78. Margolis, E. T., Nelson, P. M., Fiske, A., Champaud, J. L. Y., Olson, H. A., Gomez, M. J. C., Dineen, Á. T., Bulgarelli, C., Troller-Renfree, S. V., Donald, K. A., Spann, M. N., Howell, B., Scheinost, D., & Korom, M. (2025). Modality-level obstacles and initiatives to improve representation in fetal, infant, and toddler neuroimaging research samples. *Developmental Cognitive Neuroscience*, *72*, 101505. https://doi.org/10.1016/j.dcn.2024.101505

79. Morales, S., Oh, L., Cox, K., Rodriguez-Sanchez, R., Nadaya, G., Buzzell, G., & Troller-Renfree, S. (2024). *Diversity and Representation in Developmental EEG: Participant Recruitment and Reporting*. OSF. https://doi.org/10.31234/osf.io/t2he5

80. Gabard-Durnam, L. J., Mendez Leal, A. S., Wilkinson, C. L., & Levin, A. R. (2018). The Harvard Automated Processing Pipeline for Electroencephalography (HAPPE): Standardized Processing Software for Developmental and High-Artifact Data. *Frontiers in Neuroscience*, *12*. https://doi.org/10.3389/fnins.2018.00097

81. Monachino, A. D., Lopez, K. L., Pierce, L. J., & Gabard-Durnam, L. J. (2022). The HAPPE plus Event-Related (HAPPE+ER) software: A standardized preprocessing pipeline for event-related potential analyses. *Developmental Cognitive Neuroscience*, *57*, 101140. https://doi.org/10.1016/j.dcn.2022.101140

82. Huttenlocher, P. R., de Courten, C., Garey, L. J., & Van der Loos, H. (1982). Synaptogenesis in human visual cortex—Evidence for synapse elimination during normal development. *Neuroscience Letters*, *33*(3), 247–252. https://doi.org/10.1016/0304-3940(82)90379-2

83. Siu, C. R., & Murphy, K. M. (2018). The development of human visual cortex and clinical implications. *Eye and Brain*, *10*, 25–36. https://doi.org/10.2147/EB.S130893

84. Caffarra, S., Joo, S. J., Bloom, D., Kruper, J., Rokem, A., & Yeatman, J. D. (2021). Development of the visual white matter pathways mediates development of electrophysiological responses in visual cortex. *Human Brain Mapping*, *42*(17), 5785–5797. https://doi.org/10.1002/hbm.25654

85. Cordano, C., Sin, J. H., Timmons, G., Yiu, H. H., Stebbins, K., Guglielmetti, C., Cruz-Herranz, A., Xin, W., Lorrain, D., Chan, J. R., & Green, A. J. (2022). Validating visual evoked potentials as a preclinical, quantitative biomarker for remyelination efficacy. *Brain*, *145*(11), 3943–3952. https://doi.org/10.1093/brain/awac207

86. Fornari, E., Knyazeva, M. G., Meuli, R., & Maeder, P. (2007). Myelination shapes functional activity in the developing brain. *NeuroImage*, *38*(3), 511–518. https://doi.org/10.1016/j.neuroimage.2007.07.010

87. Hendry, A., Johnson, M. H., & Holmboe, K. (2019). Early Development of Visual Attention: Change, Stability, and Longitudinal Associations. *Annual Review of Developmental Psychology*, *1*(Volume 1, 2019), 251–275. https://doi.org/10.1146/annurev-devpsych-121318-085114

88. Rose, S. A., Feldman, J. F., & Wallace, I. F. (2025). *Infant Information Processing in Relation to Six-Year Cognitive Outcomes*.

89. Rosen, M. L., Amso, D., & McLaughlin, K. A. (2019). The role of the visual association cortex in scaffolding prefrontal cortex development: A novel mechanism linking socioeconomic status and executive function. *Developmental Cognitive Neuroscience*, *39*, 100699. https://doi.org/10.1016/j.dcn.2019.100699

90. Hansen, R. L., Struthers, J. M., & Gospe Jr., S. M. (1993). Visual Evoked Potentials and Visual Processing in Stimulant Drug-Exposed Infants. *Developmental Medicine & Child Neurology*, *35*(9), 798–805. https://doi.org/10.1111/j.1469-8749.1993.tb11731.x

91. Otten, K., Edgar, J. C., Green, H. L., Mol, K., McNamee, M., Kuschner, E. S., Kim, M., Liu, S., Huang, H., Nordt, M., Konrad, K., & Chen, Y. (2024). The maturation of infant and toddler visual cortex neural activity and associations with fine motor performance. *Developmental Cognitive Neuroscience*, *71*, 101501. https://doi.org/10.1016/j.dcn.2024.101501

92. Baroncelli, L., Braschi, C., Spolidoro, M., Begenisic, T., Maffei, L., & Sale, A. (2011). Brain Plasticity and Disease: A Matter of Inhibition. *Neural Plasticity*, *2011*(1), 286073. https://doi.org/10.1155/2011/286073

93. Hensch, T. K. (2005). Critical period plasticity in local cortical circuits. *Nature Reviews Neuroscience*, *6*(11), Article 11. https://doi.org/10.1038/nrn1787

94. Ismail, F. Y., Fatemi, A., & Johnston, M. V. (2017). Cerebral plasticity: Windows of opportunity in the developing brain. *European Journal of Paediatric Neurology*, *21*(1), 23–48. https://doi.org/10.1016/j.ejpn.2016.07.007

95. Sale, A., Berardi, N., Spolidoro, M., Baroncelli, L., & Maffei, L. (2010). GABAergic inhibition in visual cortical plasticity. *Frontiers in Cellular Neuroscience*, *4*. https://doi.org/10.3389/fncel.2010.00010

96. Takesian, A. E., & Hensch, T. K. (2013). Chapter 1—Balancing Plasticity/Stability Across Brain Development. In M. M. Merzenich, M. Nahum, & T. M. Van Vleet (Eds.), *Progress in Brain Research* (Vol. 207, pp. 3–34). Elsevier. https://doi.org/10.1016/B978-0-444-63327-9.00001-1

97. LeBlanc, J. J., DeGregorio, G., Centofante, E., Vogel-Farley, V. K., Barnes, K., Kaufmann, W. E., Fagiolini, M., & Nelson, C. A. (2015). Visual evoked potentials detect cortical processing deficits in Rett syndrome. *Annals of Neurology*, *78*(5), 775–786. https://doi.org/10.1002/ana.24513

98. Tejavibulya, L., Rolison, M., Gao, S., Liang, Q., Peterson, H., Dadashkarimi, J., Farruggia, M. C., Hahn, C. A., Noble, S., Lichenstein, S. D., Pollatou, A., Dufford, A. J., & Scheinost, D. (2022). Predicting the future of neuroimaging predictive models in mental health. *Molecular Psychiatry*, *27*(8), 3129. https://doi.org/10.1038/s41380-022-01635-2

99. Bennett, W. E., Jr, Hendrix, K. S., Thompson, R. T., Carroll, A. E., & Downs, S. M. (2014). The Natural History of Weight Percentile Changes in the First Year of Life. *JAMA Pediatrics*, *168*(7), 681–682. https://doi.org/10.1001/jamapediatrics.2014.345

100. Marek, S., Tervo-Clemmens, B., Calabro, F. J., Montez, D. F., Kay, B. P., Hatoum, A. S., Donohue, M. R., Foran, W., Miller, R. L., Hendrickson, T. J., Malone, S. M., Kandala, S., Feczko, E., Miranda-Dominguez, O., Graham, A. M., Earl, E. A., Perrone, A. J., Cordova, M., Doyle, O., … Dosenbach, N. U. F. (2022). Reproducible brain-wide association studies require thousands of individuals. *Nature*, *603*(7902), Article 7902. https://doi.org/10.1038/s41586-022-04492-9

101. Shen, X., Finn, E. S., Scheinost, D., Rosenberg, M. D., Chun, M. M., Papademetris, X., & Constable, R. T. (2017). Using connectome-based predictive modeling to predict individual behavior from brain connectivity. *Nature Protocols*, *12*(3), 506–518. https://doi.org/10.1038/nprot.2016.178

102. Dean, D. C., O'Muircheartaigh, J., Dirks, H., Waskiewicz, N., Lehman, K., Walker, L., Han, M., & Deoni, S. C. L. (2014). Modeling healthy male white matter and myelin development: 3 through 60months of age. *NeuroImage*, *84*, 742–752. https://doi.org/10.1016/j.neuroimage.2013.09.058

103. Lippe, S., Kovacevic, N., & McIntosh, R. (2009). Differential maturation of brain signal complexity in the human auditory and visual system. *Frontiers in Human Neuroscience*, *3*. https://doi.org/10.3389/neuro.09.048.2009

104. Mlandu, N., McCormick, S. A., Davel, L., Zieff, M. R., Bradford, L., Herr, D., Jacobs, C. A., Khumalo, A., Knipe, C., Madi, Z., Mazubane, T., Methola, B., Mhlakwaphalwa, T., Miles, M., Nabi, Z. G., Negota, R., Nkubungu, K., Pan, T., Samuels, R., … Gabard-Durnam, L. J. (2024). Evaluating a novel high-density EEG sensor net structure for improving inclusivity in infants with curly or tightly coiled hair. *Developmental Cognitive Neuroscience*, *67*, 101396. https://doi.org/10.1016/j.dcn.2024.101396

105. Kwiatkowski, M. A., Donald, K. A., Stein, D. J., Ipser, J., Thomas, K. G. F., & Roos, A. (2018). Cognitive outcomes in prenatal methamphetamine exposed children aged six to seven years. *Comprehensive Psychiatry*, *80*, 24–33. https://doi.org/10.1016/j.comppsych.2017.08.003

106. WHO ASSIST Working Group. (2002). The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): Development, reliability and feasibility. *Addiction*, *97*(9), 1183–1194. https://doi.org/10.1046/j.1360-0443.2002.00185.x

107. Coles, C. D., & Li, Z. (2011). Functional Neuroimaging in the Examination of Effects of Prenatal Alcohol Exposure. *Neuropsychology Review*, *21*(2), 119–132. https://doi.org/10.1007/s11065-011-9165-y

108. Donald, K. A., Eastman, E., Howells, F. M., Adnams, C., Riley, E. P., Woods, R. P., Narr, K. L., & Stein, D. J. (2015). Neuroimaging effects of prenatal alcohol exposure on the developing human brain: A magnetic resonance imaging review. *Acta Neuropsychiatrica*, *27*(5), 251–269. https://doi.org/10.1017/neu.2015.12

109. Rigby, R. A., & Stasinopoulos, D. M. (1996). A semi-parametric additive model for variance heterogeneity. *Statistics and Computing*, *6*(1), 57–65. https://doi.org/10.1007/BF00161574

110. Cole, T. J., & Green, P. J. (1992). Smoothing reference centile curves: The lms method and penalized likelihood. *Statistics in Medicine*, *11*(10), 1305–1319. https://doi.org/10.1002/sim.4780111005