The evolution of gene expression in seasonal environments

Shuichi N. Kudo[1], Yuka Ikezaki[2], Junko Kusumi[3], Hideki Hirakawa[4,5], Sachiko Isobe[4,6], Akiko Satake[2]*

1. Graduate School of Systems Life Sciences, Kyushu University, Fukuoka 819-0395, Japan

2. Department of Biology, Faculty of Science, Kyushu University, Fukuoka 819-0395, Japan

3. Department of Environmental Changes, Faculty of Social and Cultural Studies, Kyushu University, Fukuoka 819-0395, Japan

4. Kazusa DNA Research Institute, 2-6-7 Kazusa-kamatari, Kisarazu, Chiba 292-0818, Japan

5. Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University, Fukuoka 819-0395, Japan

6. Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, 113-8657, Japan

*Corresponding author: Akiko Satake

Email: satake.akiko.269@m.kyushu-u.ac.jp

Author Contributions: S.N.K. and A.S. designed research; S.N.K., Y.I., J.K., H.H., S.I., and A.S. performed research; S.N.K., Y.I., J.K., H.H., S.I., and A.S. analyzed data; S.N.K. and A.S. wrote the paper with inputs from all the authors.

Competing Interest Statement: The authors declare no conflict of interest.

Keywords: Phenology, Gene expression evolution, Seasonal constraints

Abstract

The biological activities of organisms are closely linked to seasonality. Phenology, the temporal orchestration of biological activities, is governed by gene expression, yet the evolutionary dynamics shaping seasonal gene expression remain unclear. To investigate the evolutionary patterns of gene expression in seasonal environments, we compared genome-wide expression dynamics (molecular phenology) in four dominant evergreen Fagaceae species in Asia (*Quercus glauca*, *Q. acuta*, *Lithocarpus edulis*, and *L. glaber*), using leaf and bud tissues over two seasonal cycles. We assembled high-quality reference genomes for *Q. glauca* and *L. edulis*, identifying 11749 single-copy orthologous genes. Seasonal

transcriptomic profiling of these orthologous genes revealed highly conserved gene expression across species in winter when temperatures fall below ~10°C. Rhythmic gene expression with significant periodic oscillations was more prevalent in buds (51.9%) than in leaves (40.6%), with most rhythmic genes (78.4–92.0%) exhibiting annual periodicity, while a smaller fraction (1.2–11.9%) followed half-annual cycles. The seasonal peaks of rhythmic genes were synchronized across species in winter but diverged during the growing season, aligning with phenological events such as leaf flushing and flowering. Genes with conserved seasonal peaks exhibited slower evolutionary rates, suggesting weak selective pressures maintaining seasonal gene expression. However, the evolutionary rates were similarly slow regardless of whether genes were expressed in winter or the growing season. These findings suggest that winter gene expression is more conserved than that in the growing season, imposing a seasonal constraint on gene expression evolution. This constraint may limit temporal niche partitioning and slow species divergence rates in seasonal environments.

Introduction
Fundamental biological processes, such as growth, mortality, and reproduction, rarely remain constant over time; instead, they exhibit pronounced variation in response to seasonal environmental changes (1). This temporal organization of a series of biological activities, known as phenology, plays a critical role in the evolution of life histories in response to seasonal fluctuating environments (2–5). Phenology is regulated by the coordinated expression of genes that enable organisms to exhibit essential functions at the appropriate times (6–8). However, gene expression dynamics underpinning seasonal responses in organisms remain complex and not fully understood.

To explore the evolution of gene expression, transcriptome analyses are widely employed to compare transcript profiles across organs, sexes, species, and developmental stages within a given environment. Such studies have been conducted in mammals (9–11), *Drosophila* (12, 13), and Tanzanian cichlids (14), providing key insights into the evolutionary dynamics of gene expression and its central role in shaping phenotypic diversity. These studies have revealed that gene expression profiles tend to be more conserved across species within the same tissue, whereas the evolutionary rate of gene expression varies among organs. In mammals, for instance, nervous tissue exhibits a slower rate of gene expression change compared to the testis (9–11), suggesting the presence of tissue-specific functional constraints that influence the evolution of gene expression.

In addition to tissue-dependent constraints, seasonal environmental changes may play a crucial role in shaping the evolution of gene expression, particularly by influencing the temporal coordination of gene expression under natural conditions. Gene expression responses to certain seasons may be more resistant to change, whereas in other seasons, divergence in gene expression timing across species may occur, contributing to temporal niche differentiation. Despite its potential importance, the influence of seasonal environmental

fluctuations on the gene expression evolution remains poorly understood, largely
because most transcriptomic analyses have been conducted under tightly
controlled laboratory conditions.
To investigate the evolution of gene expression in seasonal environments,
we performed a molecular phenology analysis on Fagaceae tree species, which
are widely distributed across the Northern Hemisphere, with its center of diversity
in subtropical Southeast Asia (15–18). Molecular phenology refers to
transcriptome analysis conducted under natural fluctuating conditions (6, 19–21)
and provides a valuable framework for studying gene–environment interactions
and their role in orchestrating the temporal biological functions essential for
adaptation to environmental fluctuations (7). In this study, we selected four
evergreen *Fagaceae* species—two *Quercus* species (*Q. glauca* and *Q. acuta*) and
two *Lithocarpus* species (*L. edulis* and *L. glaber*)—that co-occur in the same
locality in Japan. *Q. glauca*, *L. edulis* and *L. glaber* inhabit nearly identical
environment, while *Q. acuta* is restricted to the high-altitude regions with slightly
colder environment compared to the other three species. Leaf flushing and
flowering phenology vary across species, with *Q. glauca* and *Q. acuta* flowering in
early spring, almost simultaneously with leaf flushing, *L. edulis* flowering in late
spring, and *L. glaber* flowering in autumn, approximately one to two months after
leaf flushing. This setting enabled us to compare seasonal gene expression
profiles across species with diverse phenology, both within and between genera,
under nearly identical environmental fluctuations. By assessing and comparing
seasonal transcriptomic dynamics across *Fagaceae* species, our study provides
new insights into the evolution of gene expression and adaptation to seasonal
natural environments.
105
Results
Genome assemblies of *Q. glauca* and *L. edulis*. To quantify transcript
abundance of non-model Fagaceae species, we first constructed reference
genomes for *Q. glauca*, as a representative of genus *Quercus*, and *L. edulis,* as a
representative of genus *Lithocarpus* (Fig. 1*A*). To obtain high-quality chromosome-
level reference genomes, we conducted a hybrid genome assembly for the HiFi
reads (total length of subreads: 273.6 Gb, 165× coverage) and the high-throughput
chromosome conformation capture (Hi-C) reads (total length: 48.8 Gb, 54.8×
coverage). The resultant scaffolds were 867.7 and 835.7 Mb and consisted of 797
and 750 scaffolds including 12 chromosome-level scaffolds with contig N50 of 71.7
Mb and 68.8 Mb for *Q. glauca* and *L. edulis*, respectively (Table S1). Approximately
94.8% and 97.2% of the final assemblies were assigned to 12
pseudochromosomes of *Q. glauca* and *L. edulis*, respectively (Figure 1*B*). The
number of chromosome-scale scaffolds were identical to the known numbers in
other *Quercus* (22) and *Lithocarpus* species (23). The genomes were estimated to
contain 39758 and 34059 protein-coding genes (33947 and 30246 genes were
annotated within 12 chromosomes), covering 98.3% and 95.0% of complete
Benchmarking Universal Single Copy Orthologs (BUSCO) genes
(eudicots_odb10) for *Q. glauca* and *L. edulis*, respectively (Table S1). The genome

sizes of *Q. glauca* and *L. edulis* were estimated 891.2 Mb and 868.7 Mb by
KmerGenie (24) and GenomeScope2 (25), respectively.
Overall, the abundance and distribution of protein-coding genes, GC
content, and repeat sequences (Fig. 1*C*, *D;* Table S1) and syntenic structure (Fig.
1*B*) were similar between *Q. glauca* and *L. edulis*. Using OrthoFinder2 (26), we
clustered genes from the two newly assembled genomes into orthologous groups.
We identified 13427 single-copy orthologs, enabling pairwise comparisons
between species (Fig. 1*E*). Single-copy genes exhibited significantly higher
average expression levels than multi-copy genes (Steel-Dwass test, $p < 0.05$, Fig.
1*F*, *G*). These single-copy orthologous genes were used for further analyses.
135
Seasonal gene expression dynamics. To characterize the evolution of seasonal
gene expressions, we generated a total of 483 transcriptome profiles every four
weeks over two years using leaf and bud tissues from *Q. glauca*, *Q. acuta*, *L.
edulis*, and *L. glaber* during 2021–2023. Reads from the two *Quercus* species were
mapped to the newly constructed *Q. glauca* reference genome, achieving a mean
mapping rate of 87.0 ± 5.6%. Similarly, reads from the two *Lithocarpus* species
were mapped to the newly constructed *L. edulis* reference genome, with a mean
mapping rate of 90.0 ± 3.0%. The number of reads uniquely mapped on the
reference genome was 19.9 million reads per sample, on average. Among the
single copy orthologous genes, we excluded genes with low expression levels
(mean of Reads Per Kilobase (RPK) across samples for each species < 1) and
normalized the transcript count data using the GeTMM method (27). This process
resulted in a high-quality expression dataset for 11749 genes (Fig. S1*A*). For
further analyses, we used the mean expression levels across three biological
replicates (three individuals per species), except for *L. edulis*, where only one
individual was available.
To gain an initial overview of gene expression dynamics in a seasonal
environment, we assessed the relative magnitude of seasonal variation in
expression levels for each gene by calculating the standard deviation divided by
the mean for each gene (σ_i/μ_i) and compared it to the overall variation across all
genes in the genome (σ_g/μ_g) (Fig. S1*B*). A prior study using 3025 yeast RNA-seq
experiments reported that the dynamic range of individual genes across
experiments was considerably smaller than the overall genomic variation under
laboratory conditions (28). Consistently, our results demonstrate that despite
substantial seasonal fluctuations in environmental conditions (Fig. S2), individual
gene expression remains relatively conserved. On average, the magnitude of
genome-wide variation was 15.8 times greater than the seasonal variation
observed for individual genes (Fig. S1*B*).
To assess the similarity of gene expression profiles across tissues, species,
and seasons, we performed a hierarchical clustering of seasonal transcriptome
profiles from leaf and bud tissues across four species and identified five distinct
clusters (cluster L1, L2, B1–B3) according to the Elbow method (Fig. 2*A*; Fig. S3*A*;
Datasets S1). Transcriptional profiles first grouped into leaves (cluster L1 and L2)
and buds (cluster B1–B3), irrespective of seasons and species (Fig. 2*A*), except

that newly flushed leaf samples were grouped closely with buds (Fig. 2*A*; Fig. S4).
Within the leaf cluster, transcriptome profiles were first clustered by genus and
then by species in *Lithocarpus* (cluster L1; Fig. 2*A*). In contrast, samples from
Quercus species formed a mixture of two species (cluster L2; Fig. 2*A*). Notably,
the winter samples of *Q. acuta*, collected during periods of low mean temperatures,
were distinctly separated from those collected in other seasons (cluster L2; Fig.
2*A*). Within the bud cluster, transcriptome profiles were predominantly grouped by
season—winter (cluster B3; November to February, with the exception of a March
sample from *L. edulis*; Datasets S1) versus the other growing seasons (cluster B1
and B2; March to November; Datasets S1)—and subsequently organized by genus
and species. This suggests that the winter season exerts a stronger influence on
gene expression in buds than phylogenetic relationships.
Within the bud cluster, transcriptional profiles in the winter cluster were
observed under significantly lower temperatures, approximately below 10°C,
compared to other seasons (Fig. 2*B*; Fig. S5*A*). In contrast, no such temperature-
related distinction was evident within the leaf cluster (Fig. 2*B*). Additionally,
photoperiods were markedly shorter in the winter cluster in buds (Fig. S5*B*). These
findings suggest that the buds of the four species exhibit a conserved response to
environmental factors characteristic of the winter season. To further investigate the
relationships among all samples, we conducted a principal component analysis
(PCA). The first principal component (PC1) clearly separated samples by tissue
type, while the second (PC2) and the third principal component (PC3) gradually
differentiated winter samples from those collected during other seasons (Fig. S6).
To investigate seasonal expression dynamics of each gene, we conducted
hierarchical clustering on the transcriptional dynamics of 11749 genes and
identified five distinct gene sets characterizing different transcriptional modes
(gene set I–V; Fig. 2*C, D*; Fig. S3*B*; Datasets S2). Gene set I exhibited
predominant expression in buds, with seasonal peaks during summer, whereas
gene set II was primarily expressed in leaves, highlighting clear tissue-specificity
(Fig. 2*C, D*). Gene sets III and V demonstrated genus-specific expression patterns,
with higher expression levels observed in *Lithocarpus* and *Quercus*, respectively
(Fig. 2*C, D*). In contrast, gene set IV was expressed during winter across both leaf
and bud tissues (Fig. 2*C, D*).
203
Detection of rhythmic genes. To identify genes revealing seasonal expression
dynamics and determine the period and seasonal peaks in gene expression, we
applied the RAIN algorithm (29). This approach enabled us to detect genes with
rhythmic expression (hereafter rhythmic genes) and analyze their periods and
phases (peak months). Our analysis revealed that 44.6–58.8% of genes in bud
tissues and 33.9–45.9% in leaf tissues exhibited significant rhythmic expression
across species (Fig. S7*A*; Datasets S3). Notably, bud tissues demonstrated a
higher proportion of rhythmic genes, with 51.9% across four species on average,
compared to 40.6% in leaf tissues (Fig. S7*A*).
Among the rhythmic genes, the majority (78.4–92.0%) exhibited annual
periodicity, with a period of 12 ± 1 month observed across species and tissues

(Fig. 3*A-B*, Datasets S3). Additionally, hundreds of genes (1.2–11.9%) displayed
half-annual periodicity (6 ± 1 month) with these genes being most frequently
observed in the bud samples of *L. edulis* (Fig. 3*A*, *B*). The fraction of genes
exhibiting other periodicities was relatively small (5.7–14.1%; Fig. 3*A-B*). Based on
the observed period distributions, we classified genes with periods within the range
of 8–16 months as exhibiting annual periodicity (96.0%) and those with periods
shorter than 8 months as having half-annual periodicity (4.0%), accounting for
detection noise (Fig. S7*A*). Rhythmic genes with longer periods (>17 months) were
also identified, although their periodicity may be less reliable due to the limited
number of time points. Many genes with annual periodicity were shared across all
species, including 1280 genes in leaves and 1894 genes in buds (Fig. S7*B*). In
contrast, most genes exhibiting half-annual periodicity were species-specific, with
no genes shared among all the species (Fig. S7*C*).
228
Peak month distribution of rhythmic genes and intra-genus and inter-genera
comparison. To determine the seasonal timing of gene expression peaks, we
focused on genes with annual or half-annual periodicity and visualized the density
and direction of these peaks throughout the year using a circular histogram (Fig.
3*C*, *D*; Datasets S3). On average, 34.9% and 41.3% of annual rhythmic genes
peaked in winter (December–February) in leaf and bud samples, respectively.
Additionally, another peak corresponding to or occurring slightly after leaf flushing
and flowering events was observed, with its timing varying by species: April in *Q.*
glauca (15.6%), July in *Q. acuta* (15.3%), June in *L. edulis* (16.0%), September in
L. glaber (18.1%), for leaf tissues (Fig. 3*C*). Similar peaks corresponding to leaf
flushing or flowering were also detected in bud samples, occurring approximately
one to two months earlier than in leaf samples (Fig. 3*D*). For genes with half-annual
periodicity, the majority (63.2 ± 4.8%) exhibited expression peaks both in spring
(March–May) and autumn (September–November) in both tissues (Fig. S7*D*;
Dataset S3), with the exception of *L. glaber*, an autumn flowering species.
Interestingly, *WRKY22*, a transcription factor regulating ethylene biosynthesis and
senescence (30), exhibited half-annual periodicity in *L. glaber* (Fig. S7*E*). In
addition, *SNF1-RELATED PROTEIN KINASE REGULATORY SUBUNIT GAMMA*
1 (*KINγ1*), a subunit of SNF1-related kinase (*SnRK*) involved in stress response
such as starvation (31), showed two distinct peaks in *Q. glauca* and *L. edulis*,
occurring in both summer (July) and winter (January) (Fig. S7*E*). This result
suggests that both summer and winter pose stressful conditions for these species.
To compare the peak month of gene expression with annual periodicity
across species, we quantified the number of genes within each peak combination
for both intra-genus (Fig. 4*A*) and inter-genera species pairs in buds (Fig. 4*B*) and
leaves (Fig. S8*A*, *B*). During winter, a large number of genes exhibited highly
synchronized expression peaks across species, with high pairwise Pearson's
correlation coefficients observed (Fig. 4*A*, *B*). In contrast, the peak timing of gene
expression from spring to autumn varied across species, aligning with differences
in leaf flushing and flowering times (Fig. 4*A*, *B*). Some genes exhibited pronounced
divergence, showing negative Pearson's correlation coefficients (< -0.3) between

species. Notably, the number of such genes was larger in inter-genera
comparisons (11–77 genes; Fig. 4*B*) than in intra-genus comparisons (4–12
genes; Fig. 4*A*). A similar pattern was observed in leaf samples (Fig. S8*A, B*).
When comparing seasonal expression peaks between tissues, the majority of
genes were synchronously activated in winter in both tissues (Fig. S8*C*). During
spring to summer, slight variations in seasonal expression peaks were observed
among tissues (Fig. S8*C*).
We quantified the degree of divergence in seasonal gene expression
patterns across species by calculating the peak month differences between two
species, where one species exhibits an expression peak in the given month. We
defined the molecular phenology divergence index (*D*) for each month by
quantifying the proportion of genes with peak month differences greater than 2
months between two species, relative to the total number of genes with expression
peaks in that month. From spring to autumn, the mean of *D* over all species pairs
in buds ranged from 0.47 to 0.66, but it declined sharply during winter, reaching a
minimum of 0.12 in January in buds (Nemenyi test, $p < 0.05$; Fig. 4*C*). A similar
trend was observed in leaf samples, with another trough in summer (Fig. S8*D*).
The values of *D* were not consistently smaller in inter-genera pairs compared to
intra-genus pairs. These findings suggest that there is a seasonal constraint in
gene expression in which the degree of molecular phenology divergence is not
constant but varies significantly across seasons, with the least divergence
occurring in winter.
282
Phylogenetic constraints in the evolution of seasonal gene expression. We
next investigated whether phylogenetic relationship influences the evolution of
seasonal gene expression by calculating pairwise Pearson's correlations of
seasonal expression for each gene across all species pairs (Datasets S4). The
median of Pearson's correlation coefficients across all genes were significantly
higher between species within the same genus compared to those between
species from different genera in both leaves (Steel-Dwass test, $p < 0.05$; Fig. S9*A*)
and buds (Steel-Dwass test, $p < 0.05$; Fig. 5*A*). The intra-genus comparison in
genus *Lithocarpus* exhibited the highest correlation in their seasonal gene
expression, despite their notable differences in flowering phenology, with *L. edulis*
flowering in spring and *L. glaber* in autumn. In contrast, the intra-genus comparison
in genus *Quercus* resulted in a relatively low correlation in their gene expression
regardless of relatively similar phenology in leaf flushing and flowering. This
difference may be attributed to environmental factors, as *Q. acuta* is the only
species found in the high-altitude study site with colder climates compared to the
other three species.
Rhythmic genes exhibited more conserved expression patterns than
arrhythmic genes in both tissues (Fig. 5*A*; Fig. S9*A*). In particular, genes with
seasonal expression peaks in winter showed high correlation across species in
both tissues (Fig. 5*B*; Fig. S9*B*). The proportion of rhythmic genes with annual
periodicity declined as the mean correlation coefficient across species decreased,
with a more rapid decline observed for genes peaking in winter compared to those

with peak expression during the growing seasons (Fig. 5*B*; Fig. S9*B*). Additionally,
the mean absolute difference in seasonal gene expression dynamics was
significantly smaller between species within the same genus than between species
from different genera in both tissues (Fig. S10*A*, *B*). The relationship between the
mean absolute difference and the proportion of rhythmic genes was less clear
compared to that of the mean correlation coefficient (Fig. S10*C*, *D*). These findings
suggest that, in addition to seasonal constraints, the overall dynamics of seasonal
gene expression are also shaped by phylogenetic constraints.
313
Functional annotation of genes with conserved or divergent expression
across species. To functionally annotate genes with conserved or divergent
expression across species, we extracted genes with the top and bottom 5% based
on the distribution of mean correlation (Fig. 5*B*; Fig. S9*B*) and conducted Gene
Ontology (GO) enrichment analysis. GO analysis revealed that the top 5% of genes
in leaves were associated with photosynthesis, while those in buds were linked to
cell fate determination (Fig. S11*A*; Datasets S5). The gene with the highest
correlation in leaves was *STRESS RESPONSE SUPPRESSOR1* (*STRS1*; mean
of pairwise correlation = 0.91; Fig. 5*C*), a DEAD-box RNA helicase previously
reported to enhance tolerance to abiotic stresses such as cold (32). In buds, the
most highly correlated gene was the histone deacetylase *HDA19* (mean of
pairwise correlation = 0.93; Fig. 5*C*), which plays a key role in environmental stress
responses through histone modification and chromatin remodeling (33). These
findings suggest that genes associated with photosynthesis during the growing
season and stress responses during winter exhibit highly conserved seasonal
expression patterns across species.
The bottom 5% of genes were predominantly associated with DNA
replication and cell division in leaves (Fig. S11*B*), aligning with the divergent leaf
phenology observed (Fig. 3*B*). In contrast, GO terms related to immune response
were weakly enriched in buds (Fig. S11*B*). Notably, we identified MADS-box genes
linked to floral transition among the bottom 5% of genes: *BROTHER OF FT AND*
TFL1 (*BFT*) in leaves (mean of pairwise correlation = -0.094) and *FRUITFULL*
(*FUL*) in buds (mean of pairwise correlation = -0.069). *BFT*, a floral repressor (34),
was expressed in autumn in the spring-flowering species *L. edulis*, and in spring
in the autumn-flowering species *L. glaber* leaves (Fig. 5*D*). Meanwhile,
FRUITFULL (*FUL*), a positive regulator of floral transition (35), was expressed in
spring in *L. edulis* and in autumn in *L. glaber* (Fig. 5*D*), mirroring the timing of
flowering. In *Q. glauca* and *Q. acuta*, *FUL* was expressed in winter (Fig. 5*D*), nearly
out of phase with their expressions in *Lithocarpus* species. These results suggest
that genes involved in reproductive processes exhibit species-specific divergence
in expression patterns in response to seasonal environmental changes.
345
Relating seasonal gene expression divergence to sequence divergence. If
divergence of seasonal timing of gene expression is related to selective constraint,
we would expect a correlation between sequence divergence patterns and the
divergence in seasonal expression peaks. To explore the relationship between the

divergence of seasonal gene expression and protein sequence evolution, we
calculated synonymous (dS) and non-synonymous substitution rates (dN) and their
ratio (dN/dS) for each gene using coding sequences from newly assembled
genomes of *Q. glauca* and *L. edulis*. We first confirmed that the highly expressed
genes evolve more slowly by identifying a significant negative correlation between
evolutionary rate (dN/dS) and mean of the gene expression (r = -0.21, p = 1.8 ×
10^{-118} for dN/dS; Fig. S12*A*; Datasets S6). This finding aligns with previous studies
in yeasts, mammals, and plants (36–39). By targeting the genes that exhibited
annual periodicity across both species and tissues, we examined the relationship
between these evolutionary measures and peak month differences in seasonal
gene expression between species (Δ*φ*) in leaves and bud tissues using linear
regression. Assessment of the significance of the regression coefficient using a *t*-
test revealed that genes with smaller peak month differences between species
exhibited lower values of dN/dS in buds (r = 0.072, p =2.6 × 10^{-2}) and marginally
lower values of dN/dS in leaves (r = 0.062, p = 5.7 × 10^{-2}). The lower values were
also found in dN in both leaves (r = 0.086, p = 8.1 × 10^{-3}) and buds (r = 0.086, p =
7.8 × 10^{-3}), whereas no significant trend was observed for dS in both tissues (Fig.
5*E*; Fig. S12*B*,*C*; Datasets S6). This finding suggests that genes with conserved
seasonal expression patterns tend to evolve at a slower rate. We further
investigated whether the association between the divergence in seasonal
expression peaks and sequence divergence patterns varies between genes
expressed in winter and in the growing season. We classified genes that exhibited
annual periodicity across both species and tissues into four groups based on their
expression peak timing (winter or growing seasons) and tissue specificity (leaves
or buds) and compared the extent of protein sequence divergence across these
groups. We found no significant differences in dN/dS (Kruskal Wallis test, p = 2.8
× 10^{-1}; Fig. S12*D*; Datasets S6) between genes expressed in winter in both tissues
and those in growing seasons. The same result was observed even when focusing
only on genes with conserved peaks (peak month difference < 2), with no
significant differences in dN/dS between these gene sets (Kruskal Wallis test, p =
3.5 × 10^{-1}; Fig. S12*E*; Datasets S6). These results suggest that, regardless of the
season, genes with conserved expression peaks tend to evolve at slower rates.
382
Discussion
A central goal in biology is to unravel the molecular basis of phenotypic evolution,
with gene expression serving as a fundamental driver of this process. In this study,
we investigated the impact of seasonal environmental fluctuations on the evolution
of gene expression and its relationship with phenological traits in four dominant
Fagaceae tree species in forest ecosystems. We found that gene expression
profiles were more resistant to change during winter compared to the growing
season, indicating the presence of a seasonal constraint on the evolution of gene
expression. This constraint becomes particularly pronounced in buds during winter
when temperatures fall below approximately 10°C (Fig. 2*B*). This temperature
threshold aligns with previous findings, where winter-specific transcript profiles
were reported at temperatures below 8.7–11.3°C in cherry trees (40), 11.3°C in

ever-green *Fagaceae* (20), and 10.5°C in perennial *Arabidopsis* (41). These results suggest the presence of a conserved mechanism for gene expression regulation under low-temperature environments across different species. The greater conservation of gene expression patterns in buds compared to leaves across species may be attributed to their functional role protecting essential meristematic tissues, including the shoot apex and leaf and flower primordia. Additionally, buds likely play a crucial role in maintaining common physiological functions necessary for the resumption of growth in spring. This functional consistency may impose evolutionary constraints on gene expression, contributing to its conservation across species.

It is intriguing to compare the seasonal constraints identified in this study with developmental constraints in gene expression evolution, where evolutionary conservation and variation across developmental stages during the life cycle have been observed, such as in the early conservation model and the developmental hourglass model (Fig. S13*A*)(42–44). Previous studies have demonstrated that certain developmental stages exhibit remarkably similar gene expression patterns (45–47). Our findings highlight that such highly similar gene expression patterns are observed during the winter season (Fig. S13*B*), suggesting that a common physiological state is essential for enduring harsh winter conditions. This seasonal constraint on gene expression evolution may have the potential to limit temporal niche partitioning at the phenotypic level and slow species divergence rates in seasonal environments (48, 49). Although completely disentangling the effects of developmental stages and seasonal changes on gene expression is challenging when using molecular phenology data collected from natural environments, the observation that species with different phenology—i.e., different developmental timing in seasonal environments—exhibit convergent gene expression patterns during winter underscores the profound influence of winter environmental conditions on the evolution of gene expression. Future research is needed to elucidate the mechanisms driving the establishment of such conserved gene expression patterns, including the dynamic reorganization of chromatin architecture in response to cold (50–52).

We found that the evolution of coding-sequences was weakly correlated with the divergence of seasonal peaks in gene expression across species. This suggests that weak functional constraints exist for specific proteins that are expressed in certain seasons, such as photosynthesis in summer and stress response to winter. However, among genes with conserved seasonal peaks, the evolutionary rate of coding-sequences was similarly slow regardless of whether they were expressed in winter or during the growing season. Therefore, to understand the evolutionary mechanism underlying the seasonal constraint on gene expression that results in similar expression patterns during winter, it is essential to consider evolution of non-coding sequences that modulate gene expression levels in response to seasonal environmental changes. Regulatory variation has been recognized as a major driver of evolutionary innovation (53, 54). Recent deep learning studies utilizing genome and transcriptome data from yeast and humans have highlighted the significance of co-evolution between coding and

non-coding sequences in shaping gene expression levels (28, 55). Understanding
how seasonal environments drive the co-evolution of coding and regulatory
sequences, as well as the divergence of molecular phenology and its phenological
outputs, will require expanding analyses to a broader range of species. Such
efforts will facilitate the development of predictive models for evolutionary
processes influenced by seasonal dynamics. Comparison of molecular phenology
across wide climatic zones, including both tropical and temperate zones, will also
be effective to identify key genes for the phenological response to on-going climate
change.
449
Materials and Methods
Sample collection for genome sequencing
452 To construct genomes of *Q. glauca* and *L. edulis*, we collected approximately 5g
453 of fresh leaves from one individual in spring, respectively. We shipped these
454 samples to Dovetail Genomics, LLC (Scotts Valley, CA, USA) where Dovetail Staff
455 performed DNA extraction, library preparation, sequencing, and assembly steps.
456
Genome sequencing and assembly
DNA extraction was performed using the QIAamp DNA Mini Kit (Qiagen,
Germantown, MD, USA) using the manufacturer's recommended protocol. Mean
fragment length of the extracted DNA was 60 kbp for *Q. glauca* and 65 kbp for *L.
edulis*. DNA samples were quantified using Qubit 2.0 Fluorometer (Life
Technologies, Carlsbad, CA, USA). The library preparation, sequencing and
scaffolding were carried out by Dovetail Genomics (California, USA) according to
their standard genome assembly pipeline (https://dovetailgenomics.com/). The
PacBio SMRTbell library (~20kb) for PacBio Sequel was constructed using
SMRTbell Express Template Prep Kit 2.0 (PacBio, Menlo Park, CA, USA) using
the manufacturer's recommended protocol. The library was bound to polymerase
using the Sequel II Binding Kit 2.0 (PacBio) and loaded onto PacBio Sequel II.
Sequencing was performed on PacBio Sequel II 8M SMRT cells. PacBio CCS
reads were used as an input to Hifiasm1 v0.15.4-r347 with default parameters. For
Q. glauca, we also generated sequence data using PacBio Revio. BLASTn results
of the contigs obtained by Hifiasm assembly against the NCBI's NT database were
used as input for blobtools2 v1.1.1 and contigs identified as possible contamination
were removed from the assembly. Finally, purge_dups3 v1.2.5 was used to
remove haplotigs and contig overlaps.
476
Assembly scaffolding with HiRise
A proximity ligation library was generated by the Dovetail's Omni-C technique,
followed by sequencing on an Illumina HiSeqX platform. Chromatin was fixed in
place in the nucleus with formaldehyde before extraction
(https://dovetailgenomics.com/wp-content/uploads/2021/09/Omni-C-Tech-
Note.pdf). Fixed chromatin was digested with DNase I, fragmented chromatin ends
were repaired and biotinylated to adapters followed by proximity ligation.
Crosslinks were then reversed, the DNA purified, and the biotin subsequently

removed. The DNA library was prepared and sequenced to produce 2 × 150 bp
paired-end reads (around 30× coverage). The Omni-C technology uses a
sequence-independent endonuclease which provides even, unbiased genome
coverage. The input *de novo* assembly and Omni-C library reads were used as
input data for HiRise, a software pipeline designed specifically for using proximity
ligation data to scaffold genome assemblies (56). The Omni-C library sequences
were aligned to the contigs assembled by Hifiasm using bwa (57). The separations
of Omni-C reads from paired-end reads mapped within the contigs were analyzed
by HiRise to produce a likelihood model for genomic distance between paired-end
reads, and the model was used to identify and break putative misjoins, to score
prospective joins, and make joins above a threshold.
496
Genome size estimation
The genome size of *Q. glauca* was estimated by GenomeScope2 v2.0 (25) using
k-mer frequency plot obtained by kmc v3.2.4 (58) with *k*-mer size 61 against the
HiFi reads. The paired-end reads of four individuals of *L. edulis* were merged and
trimmed by fastp v0.23.2 (59). The genome size of *L. edulis* was estimated by
KmerGenie (24) using *k*-mer size 55 against the trimmed paired-end reads.
503
Sample collection for RNA sequencing
To monitor the genome-wide gene expression profiles, we collected leaf, bud, and
flower samples from four Fagaceae species, *Q. glauca*, *Q. acuta*, *L. edulis* and *L.*
glaber, every four weeks spanning from May 2019 to June 2023 across Fukuoka
and Saga prefectures in Kyushu, Japan. We targeted three individuals, and each
sample was collected from three branches per individual, except for *L. edulis*.
Given the consistent expression patterns observed among individuals in a previous
study (20), we focused on only one individual for *L. edulis*. Study sites for *Q. glauca*
is the biodiversity reserve on the Ito campus of Kyushu University (33º35′ N,
130º12′ E, 20 to 57 m a.s.l). *Q. acuta* was monitored in Sefuri mountains (33º26′
N, 130º22′ E, 970 to 974 m a.s.l). *L. edulis* and *L. glaber* were studied at the
Imajuku Field Activity Center (33º33′ N, 130º16′ E, 84 to 111 m a.s.l).
Environmental data from Ito Campus and the Imajuku Field Activity Center were
obtained from the Japan Meteorological Agency, while data from Sefuri Mountain
were collected using a logger installed at the sampling site.
Samples for RNA-seq were taken from the sun-exposed crown
(approximately 2–5 m from the ground) using long pruning shears from 11:30 to
12:30 h. For each sample, 0.2–0.4 g of tissue was preserved in a 2 ml microtube
containing 1.5 ml of RNA-stabilizing reagent (RNAlater; Thermo Fisher Scientific,
Waltham, MA, USA) immediately after harvesting. The samples were transferred
to the laboratory within 2 hr after sampling, stored at 4°C overnight and then stored
at −80°C until RNA extraction. During transport to the laboratory, the samples were
kept in a cooler box with ice to maintain a low temperature.
527
RNA extraction and RNA-seq analysis

RNA was extracted independently from bud and leaf samples of each tree. Total
RNA of the leaf sample was extracted in accordance with the method described in
a previous study (60). Total RNA of the bud sample was extracted using the
PureLink™ Plant RNA Reagent (Thermo Fisher Scientific, Waltham, MA, USA)
and subsequently processed with DNase using the TURBO DNA-free kit™
(Thermo Fisher Scientific) according to their protocols and purified using the
RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) according to its RNA cleanup
protocol. RNA integrity was examined using the Agilent RNA 6000 Nano kit on a
2100 Bioanalyzer (Agilent Technologies), and the RNA yield was determined using
a NanoDrop ND-2000 spectrophotometer (Thermo Fisher Scientific). Five to six
micrograms of RNA extracted from each sample were sent to Hangzhou Veritas
Genetics Medical Institute Co., Ltd., where a cDNA library was prepared with a
NEBNext Ultra II RNA Library Prep Kit for Illumina, and 150 base paired-end
transcriptome sequencing of each sample was conducted using an Illumina
NovaSeq6000 sequencer (Illumina).
The quantification of gene expression followed a three-step process. First,
quality filtering was performed using the fastp v0.23.2 with default options: phred
quality ≥15, the maximum number of N base in one read = 5 (59). Second, RNA-
seq reads were mapped to the reference genome using STAR v2.7.10b (61). Third,
RNA-seq reads were quantified using RSEM v1.3.1 (62). We excluded the genes
with low expression level using the threshold, the value of mean Reads Per
Kilobase (RPK) < 1 for each species. Then, we further normalized the expression
level using the value of Gene length corrected Trimmed Mean of M-values
(GeTMM) (27) using an R package "edgeR" v3.42.4. For the downstream analysis
we calculated the mean expression levels of GeTMM of each gene across multiple
individuals and transformed them to $\log_2$(GeTMM+1).
555
Genome annotation
The RNA-seq reads of *Q. glauca* (8 samples obtained by a previous study (20)
collected from May to December 2017; GEO's accession number: GSE211384)
and *L. edulis* (20 samples covering leaf, bud and flower tissues collected in
January, April, July, and November 2021) were respectively mapped against the
scaffolds of *Q. glauca* and *L. edulis* obtained by HiRise with HISAT2 v2.1.0 (63).
The bam files generated by the mappings and amino acid sequences of the 9
species obtained from NCBI (*Castanea mollissima*: GCA_000763605.2 (NCBI's
accession number), *Q. lobata*: GCF_001633185.1, *Q. robur*: GCF_932294415.1,
Q. suber: GCF_002906115.1, *Carpinus fangiana*: GCA_006937295.1, *Carya
illinoinensis*: GCF_018687715.1, *Juglans macrocarpa × J. regia*:
GCF_004785595.1, *Juglans regia*: GCF_001411555.2, *Morella rubra*:
GCA_003952965.2) were used for ab initio gene prediction by Braker2 v2.1.6 (64),
and the genes with highest-score were selected to exclude splicing variants. In
parallel, the gene models were respectively constructed in *Q. glauca* and *L. edulis*
by GeMoMa v1.7.1 (65) with RNA-seq reads and amino acid sequences used for
the gene prediction in Braker2, and the genes with highest-score were selected to
exclude splicing variants. The duplicated genes between Braker2 and GeMoMa

were excluded by GFFcompare v0.11.2 (66). The remaining genes were searched
against UniProtKB (67) by DIAMOND v2.0.5 (68) in 'more-sensitive' option with E-
value ≤10^{-20} and identity ≥25%. Next, gene expression analysis was performed by
Salmon v1.4.0 (69), and the genes with TPM values >0.1 were considered as
expressed. The orthologous genes were searched against EggNOG (70) using
eggNOG-mapper v2.1.8 (71) with E-value ≤10^{-10}. The genes were searched
against protein sequences of *Castanea mollissima* (Accession:
GCA_000763605.2), *Quercus lobata* (GCF_001633185.2), and *Arabidopsis*
thaliana (Araport11 (72)) by BLASTp v2.13.0 (73) with E-value ≤10^{-20}. According
to these similarity searches, the genes whose product names contain keywords
related to transposable elements (TEs) were categorized into TE. The genes
having similarity hits and being expressed by Salmon were classified into high-
confidence (HC) genes, while remaining genes were categorized into low-
confidence (LC) genes. The genes categorized into HC were used for further
analyses.
589
Genome synteny analysis
To investigate the syntenic relationship between *Q. glauca* and *L. edulis*, we first
performed an all-vs-all BLASTp search of the protein sequences, using a cut-off
E-value of 10^{-5}. Collinearity blocks within and between the genomes were then
identified using MCScanX (74) based on the results of the BLASTp searches.
Finally, the collinearity blocks were visualized using SynVisio
(https://synvisio.github.io).
597
Illumina short-read sequencing and library preparation
To draw molecular phylogeny, we also generated short-read sequencing of *Q.*
glauca, *Q. acuta*, *L. edulis*, and *L. glaber.* DNA extraction was performed using a
modified version of the method described previously (75). The DNA samples were
sent to Macrogen Inc. (Republic of Korea) for sequencing on the Illumina HiseqX
platform (Illumina, San Diego, CA, USA). DNA was sheared to around 500 bp
fragments in size using dsDNA fragmentase (New England BioLabs, Ipswich, MA,
USA). Library preparation was performed using the NEBNext Ultra II DNA Library
Prep Kit (New England BioLabs) according to the manufacturer's protocol, and the
libraries were individually indexed with the NEBNext Multiplex Oligos for Illumina
(New England BioLabs) by PCR. The quality and quantity of each amplified library
were analyzed using the Bioanalyzer 2100 (Agilent Technologies, Santa Clara,
CA, USA), the High Sensitivity DNA kit (Agilent Technologies), and the NEBNext
Library Quant Kit for Illumina (New England BioLabs).
612
Phylogenetic analysis
We constructed a phylogenetic tree using an alignment-free method (76) based on
short-read data from *Q. acuta* and *L. glaber* and genome assembly data from *Q.*
glauca and *L. edulis*. Additionally, genomic data for *Fagus sylvatica* was obtained
from beechgenome.net (77) and included as an outgroup. Due to the large volume
of short-read data for *Q. acuta* and *L. glaber* (18 Gb each), we randomly

subsampled half of the reads. Bootstrap values were estimated through 100
resampling of the k-mer table.
621

Identification of orthologs

The prediction of orthogroups was based on a BLASTp all-against-all comparison
of the protein sequences (E-value < 10^{-5}) of 5 Fagaceae species: *L. edulis*, *Q.*
glauca, *Q. robur*, *Fagus sylvatica* and *Juglans regia*. Clustering was conducted
with OrthoFinder2, using version 2.2.6 for BLAST data generation and version
2.5.5 for orthogroup determination (26). The genomic data of *Q. robur* and *J. regia*
were downloaded from NCBI (dhQueRobu3.1 and Walnut 2.0), while *F. sylvatica*
data were sourced from beechgenome.net (77). Since the *F. sylvatica* dataset
contained numerous predicted transposable elements (TEs), we re-annotated
genes using three databases—EggNOG5 (EggNOG-mapper), Pfam35.0 (HMMER
v3.3.2), and UniRef90_2023_0628 (DIAMOND v2.1.8.162)—and excluded genes
identified as TEs in at least one database. In total, we identified 26692 orthogroups
and focused on single-copy orthogroups after filtering out genes mapped outside
the 12 chromosomes. This resulted in a final dataset of 11749 genes for further
analysis (Dataset S2).
637

Hierarchical clustering and principal component analysis (PCA)

To assess the similarity of the genome-wide transcriptional profiles across
orthologous genes and 483 samples collected in different seasons, we performed
hierarchical clustering using the monthly time series data of 11749 orthologous
genes from October 2019 to April 2023. For each orthologous gene, there were 26
or 27 time points, with three individuals each for *Q. glauca*, *Q. acuta*, *L. glaber* and
one individual for *L. edulis*. We calculated the mean expression levels of each
orthologous gene across three individuals in each species and subsequently
normalized the values by adjusting the mean to zero and the standard deviation to
one. We performed hierarchical clustering using the Ward method and the
Euclidian distance using the pheatmap function in R (v 1.0.12). To determine the
number of the clusters, we calculated Within-Cluster-Sum of Squared Error (WSS)
and determined the number of clusters as an elbow point of it (Fig. S3). To assess
the seasonal expression dynamics of 11749 orthologous genes, we also
performed PCA of the gene expression profiles from all samples using the prcomp
function in R (v 4.3.1).
654

Identification of genes with rhythmic expression

To identify genes with rhythmic expression and determine the period and peak
month of gene expression, we applied the RAIN algorithm (29) using the rain
function in R (v 1.34.0). We set the following settings: deltat = 1, period = 12,
period.delta = 11, peak.border = (0,1), nr.series = 1, method= "independent",
na.rm= T , adjp.method = "ABH". Based on the results of the RAIN analysis, we
identified genes with significant periodicity using a Benjamini-Hochberg (BH)
adjusted *q*-value threshold of < 0.01. Since the period and peak month in the RAIN
output are expressed in terms of the number of time points, we converted them

into months using the sampling span (four weeks) and sampling dates.
Subsequently, we categorized the rhythmic genes into three categories based on
the period length T (in months): "half-annual" ($0 \leqq T < 8$), "annual" ($8 \leqq T \leqq 16$),
and "long" ($T > 16$).
668
Calculation of Pearson's correlation coefficient across species
To quantify the similarity of seasonal gene expression pattern across species, we
calculated pairwise Pearson's correlations of seasonal expression for each gene
across all the species pairs. For species within the same genus, the monitoring
period was identical, allowing direct comparison. On the other hand, the monitoring
periods differed between *Quercus* and *Lithocarpus* species: *Q. glauca* and *Q.*
acuta were monitored from May 2021 to April 2023, while *L. edulis* and *L.*
glaber were monitored from October 2019 to October 2021 (Fig. S2). To ensure
comparable time series data for inter-genera comparisons, we extracted the time
series from *Q. glauca* and *Q. acuta* spanning May 19, 2021, to November 2, 2022,
and from *L. edulis* and *L. glaber* spanning May 13, 2020, to October 20, 2021, for
correlation analysis. The mean absolute differences for each gene across all the
species pairs were also calculated in a similar manner.
682
Molecular phenology divergence index (*D*)
To quantify the seasonal difference of peak month divergence across species, we
defined the molecular phenology divergence index (*D*) for each month *t* and
species pair A-B as the proportion of genes that peak in month *t* in the reference
species A but exhibit a peak month difference ($\Delta\varphi$) greater than two months in
species B using following formula:
689

690    $$D := \frac{\text{\# Genes with peak month } t \text{ in species A and } t \pm \Delta\phi (\geq 2) \text{ in species B}}{\text{\# Genes with peak month } t \text{ in species A}}.$$

691
Gene Ontology (GO) enrichment analysis
First, we assigned GO terms for each gene in *Lithocarpus* and *Quercus* genomes
based on the annotation data of *Arabidopsis thaliana* orthologs downloaded from
PLAZA Database (http://bioinformatics.psb.ugent.be/plaza/). Then, GO
enrichment analysis was conducted using the GOstats (78) function in R (v 2.66.0)
with the following settings: ontology = BP, pvalueCutoff = 0.05, conditional =
TRUE, and testDirection = over. The universe gene set was defined as genes
associated with GO terms common to both of the genomes. Genes without
annotated GO terms were removed from the universe set, and those with differing
GO term annotations between the two genomes were also excluded. The BH
adjusted *q*-values were calculated from the *p*-values generated by GOstats to
account for multiple testing.
704
Phenology measurement: leaf unfolding and flowering
To compare seasonal changes in genome-wide gene expression profiles with
phenology, we conducted observations of leaf unfolding and flowering phenology

on the three terminal branches of each of the three individuals used for gene
expression monitoring. Leaf unfolding was defined as the stage where both the
entire leaf blade and the leaf stalk were visible (Fig. S4). Flowering was defined as
the stage where either male or female flowers were visible (Fig. S4). We
subsequently plotted the proportion of branches exhibiting leaf unfolding and
flowering. The data are provided in Dataset S2.
714

Estimation of protein-coding sequence divergence
Pairwise distances for non-synonymous (dN) and synonymous (dS) substitutions,
as well as the dN/dS ratio, were estimated for each orthologous gene pair between
L. edulis and *Q. glauca* using codeml (PAML 4.0)(79). The analysis was performed
with the following settings: seqtype = 1, CodonFreq = 2, Runmode = -2, and the
transition/transversion ratio estimated from the data. To ensure reliability, genes
exhibiting signs of saturated divergence or underestimated dS values were
excluded, as codeml produces robust results only within a moderate range of
sequence divergence. Specifically, we removed 43 orthologs with dN/dS $\geqq$ 99 and
483 genes with dS > 10, dS < 0.001, or dN < 0.0003. The thresholds of dN and dS
were determined by examining the relationship between dN and dS and excluding
outliers that deviated from the main distribution. After filtering, the final dataset
comprised 11223 orthologous genes.
Linear regression analyses were performed to examine the relationship
between dN, dS, dN/dS and peak month differences ($\Delta\varphi$) in gene expression
between *Q. glauca* and *L. edulis*, as well as mean expression levels. These
analyses were conducted using the lm function in R software (v 4.3.1). Specifically,
we modeled the evolutionary rates (dN, dS, and dN/dS) as the response variable
and the peak month difference ($\Delta\varphi$) or the mean levels in gene expression as the
predictor variable. To account for the skewed distributions of evolutionary rates,
logarithm transformations with base 2 were applied.
736

Acknowledgments
We thank Atsuko Miyawaki-Kuwakado for her valuable technical advice on RNA-
seq analysis. We also appreciate the technical support provided by Kayoko Ohta.
Additionally, we thank Koharu Yamaguchi, Yuta Aoyagi, and Ryosuke Imai for their
help with sample collection.

References
1. SD. Fretwell, Populations in a seasonal environment. *Monogr. Popul. Biol.* 5, 1–217
(1972).

2. J. Forrest, A. J. Miller-Rushing, Toward a synthetic understanding of the role of
phenology in ecology and evolution. *Philos. Trans. R. Soc.* 365, 3101–3112 (2010).

3. L. P. C. Morellato, M. G. G. Camargo, E. Gressler, "A review of plant phenology in
South and Central America." in Phenology: An Integrative Environmental Science,
M. D. Schwartz Eds. (Springer, 2013), 91-113.

4.   M. D. Schwartz (Eds.), Phenology: An Integrative Environmental Science, 2nd Ed.
(Springer 2013).

5.   S. Piao, *et al.*, Plant phenology and global climate change: Current progresses and
challenges. *Glob. Chang. Biol.* 25, 1922–1940 (2019).

6.   H. Kudoh, Molecular phenology in plants: *in natura* systems biology for the
comprehensive understanding of seasonal responses under natural environments.
New Phytol. 210, 399–412 (2016).

7.   A. Satake, A. Nagahama, E. Sasaki, A cross-scale approach to unravel the
molecular basis of plant phenology in temperate and tropical climates. *New Phytol.*
233, 2340–2353 (2022).

8.   A. Satake, *et al.*, Plant molecular phenology and climate feedbacks mediated by
BVOCs. *Annu. Rev. Plant Biol.* 75, 605–627 (2024).

9.   D. Brawand, *et al.*, The evolution of gene expression levels in mammalian organs.
Nature 478, 343–348 (2011).

10.   S. Naqvi, *et al.*, Conservation, acquisition, and functional impact of sex-biased gene
expression in mammals. *Science* 365 (2019).

11.   M. Cardoso-Moreira, *et al.*, Gene expression across mammalian organ
development. *Nature* 571, 505–509 (2019).

12.   S. V. Nuzhdin, M. L. Wayne, K. L. Harmon, L. M. McIntyre, Common pattern of
evolution of gene expression level and protein sequence in *Drosophila*. *Mol. Biol.*
Evol. 21, 1308–1317 (2004).

13.   B. Lemos, B. R. Bettencourt, C. D. Meiklejohn, D. L. Hartl, Evolution of proteins and
gene expression levels are coupled in *Drosophila* and are independently associated
with mRNA abundance, protein length, and number of protein-protein interactions.
Mol. Biol. Evol. 22, 1345–1354 (2005).

14.   A. El Taher, *et al.*, Gene expression dynamics during rapid organismal
diversification in African cichlid fishes. *Nat. Ecol. Evol.* 5, 243–250 (2021).

15.   A. Kremer, *et al.*, Genomics of Fagaceae. *Tree Genet. Genomes* 8, 583–610 (2012).

16.   B. F. Zhou, *et al.*, Phylogenomic analyses highlight innovation and introgression in
the continental radiations of Fagaceae across the Northern Hemisphere. *Nat.*
Commun. 13 (2022).

17.   P. S. Manos, Z. Zhou, C. H. Cannon, Systematics of Fagaceae: phylogenetic tests
of reproductive trait evolution. *Int. J. Plant Sci.* 162, 1361–1379 (2001).

18.   A. L. Hipp, *et al.*, Genomic landscape of the global oak phylogeny. *New Phytol.* 226,
1198–1212 (2020).

19.   H. Komoto, *et al.*, The transcriptional changes underlying the flowering phenology
787   shift of Arabidopsis halleri in response to climate warming. *Plant Cell Environ.* 47,
788   174–186 (2024).

789   20.   A. Satake, K. Ohta, N. Takeda-Kamiya, K. Toyooka, J. Kusumi, Seasonal gene
790   expression signatures of delayed fertilization in Fagaceae. *Mol. Ecol.* 32, 4801–
791   4813 (2023).

792   21.   A. J. Nagano, *et al.*, Annual transcriptome dynamics in natural environments
793   reveals plant seasonal adaptation. *Nat. Plants* 5, 74–83 (2019).

794   22.   V. Zoldos, *et al.*, Molecular-cytogenetic studies of ribosomal genes and
795   heterochromatin reveal conserved genome organization among 11 Quercus
796   species. *Theor. Appl. Genet.* 99, 969–977 (1999).

797   23.   H. Liu, *et al.*, Chromosome-scale genome assembly of sweet tea (Lithocarpus
798   polystachyus Rehder). *Sci Data* 10 (2023).

799   24.   R. Chikhi, P. Medvedev, Informed and automated k-mer size selection for genome
800   assembly. *Bioinformatics* 30, 31–37 (2014).

801   25.   T. R. Ranallo-Benavidez, K. S. Jaron, M. C. Schatz, GenomeScope 2.0 and
802   Smudgeplot for reference-free profiling of polyploid genomes. *Nat. Commun.* 11
803   (2020).

804   26.   D. M. Emms, S. Kelly, OrthoFinder: Phylogenetic orthology inference for
805   comparative genomics. *Genome Biol.* 20 (2019).

806   27.   M. Smid, *et al.*, Gene length corrected trimmed mean of M-values (GeTMM)
807   processing of RNA-seq data performs similarly in intersample analyses while
808   improving intrasample comparisons. *BMC Bioinformatics* 19 (2018).

809   28.   J. Zrimec, F. Buric, M. Kokina, V. Garcia, A. Zelezniak, Learning the regulatory code
810   of gene expression. *Front. Mol. Biosci.* 8 (2021).

811   29.   P. F. Thaben, P. O. Westermark, Detecting rhythms in time series with rain. *J. Biol.*
812   *Rhythms* 29, 391–400 (2014).

813   30.   G. Q. Zhu, L. Qu, H. W. Xue, Casein kinase 1 AELs promote senescence by
814   enhancing ethylene biosynthesis through phosphorylating WRKY22 transcription
815   factor. *New Phytol.* 244, 116–130 (2024).

816   31.   S. Emanuelle, M. S. Doblin, D. I. Stapleton, A. Bacic, P. R. Gooley, Molecular
817   insights into the enigmatic metabolic regulator, SnRK1. *Trends Plant Sci.* 21, 341–
818   353 (2016).

819   32.   P. Kant, S. Kant, M. Gordon, R. Shaked, S. Barak, Stress Response Suppressor1
820   and Stress Response Suppressor2, two Dead-box RNA helicases that attenuate
821   *Arabidopsis* responses to multiple abiotic stresses. *Plant Physiol.* 145, 814–830
822   (2007).

823   33.   L. T. Chen, K. Wu, Role of histone deacetylases HDA6 and HDA19 in ABA and
824   abiotic stress response. *Plant Signal. Behav.* 5, 1318–1320 (2010).

825   34.   S. J. Yoo, *et al.*, BROTHER of FT and TFL1 (BFT) has TFL1-like activity and
826   functions redundantly with TFL1 in inflorescence meristem development in
827   *Arabidopsis*. *Plant J.* 63, 241–253 (2010).

828   35.   C. Smaczniak, *et al.*, Characterization of MADS-domain transcription factor
829   complexes in *Arabidopsis* flower development. *Proc. Natl. Acad. Sci. U. S. A.* 109,
830   1560–1565 (2012).

831   36.   D. A. Drummond, J. D. Bloom, C. Adami, C. O. Wilke, F. H. Arnold, Why highly
832   expressed proteins evolve slowly. *Proc. Natl. Acad. Sci. U. S. A.* 102, 14338–14343
833   (2005).

834   37.   B. Y. Liao, J. Zhang, Low rates of expression profile divergence in highly expressed
835   genes and tissue-specific genes during mammalian evolution. *Mol. Biol. Evol.* 23,
836   1119–1128 (2006).

837   38.   B. S. Gaut, The patterns and causes of variation in plant nucleotide substitution
838   rates. *Annu. Rev. Ecol. Evol. Syst.* 42, 245-266 (2011).

839   39.   L. Yang, B. S. Gaut, Factors that contribute to variation in evolutionary rate among
840   *Arabidopsis* genes. *Mol. Biol. Evol.* 28, 2359–2369 (2011).

841   40.   A. Miyawaki-Kuwakado, Q. Han, K. Kitamura, A. Satake, Impacts of climate change
842   on the transcriptional dynamics and timing of bud dormancy release in Yoshino-
843   cherry tree. *Plants People Planet* 6, 1505–1521 (2024).

844   41.   S. Aikawa, M. J. Kobayashi, A. Satake, K. K. Shimizu, H. Kudoh, Robust control of
845   the seasonal expression of the *Arabidopsis* FLC gene in a fluctuating environment.
846   *Proc. Natl. Acad. Sci. U. S. A.* 107, 11632–11637 (2010).

847   42.   N. Irie, S. Kuratani, Comparative transcriptome analysis reveals vertebrate
848   phylotypic period during organogenesis. *Nat. Commun.* 2 (2011).

849   43.   M. Quint, *et al.*, A transcriptomic hourglass in plant embryogenesis. *Nature* 490,
850   98–101 (2012).

851   44.   J. S. Lotharukpong, *et al.*, A transcriptomic hourglass in brown algae. *Nature* 635,
852   129–135 (2024).

853   45.   A. T. Kalinka, *et al.*, Gene expression divergence recapitulates the developmental
854   hourglass model. *Nature* 468, 811–816 (2010).

855   46.   H. G. Drost, *et al.*, Post-embryonic hourglass patterns mark ontogenetic transitions
856   in plant development. *Mol. Biol. Evol.* 33, 1158–1163 (2016).

857   47.   M. Levin, *et al.*, The mid-developmental transition and the evolution of animal body
858   plans. *Nature* 531, 637–641 (2016).

48. T. Kubo, Y. Iwasa, Phenological pattern of tree regeneration in a model for forest
species diversity. *Theor. Popul. Biol.* 49, 90-117 (1996).

49. E. E. Cleland, E. M. Wolkovich, Effects of phenology on plant community assembly
and structure. *Annu. Rev Ecol. Evol. Syst.* 55, 471-492 (2025).

50. H. Nishio, A. J. Nagano, T. Ito, Y. Suzuki, H. Kudoh, Seasonal plasticity and diel
stability of H3K27me3 in natural fluctuating environments. *Nat. Plants* 6, 1091–1097
(2020).

51. H. Fischl, *et al.*, Cold-induced chromatin compaction and nuclear retention of clock
mRNAs resets the circadian rhythm. *EMBO J.* 39 (2020).

52. X. Zhang, *et al.*, High-resolution Hi-C maps highlight multiscale chromatin
architecture reorganization during cold stress in *Brachypodium distachyon*. *BMC
Plant Biol.* 23 (2023).

53. J. R. True, E. S. Haag, Developmental system drift and flexibility in evolutionary
trajectories. *Evol. Dev.* 3, 109–119 (2001).

54. B. Prud'homme, N. Gompel, S. B. Carroll, Emerging principles of regulatory
evolution. *Proc. Natl. Acad. Sci. U. S. A.* 104, 8605–8612 (2007).

55. J. Zrimec, *et al.*, Deep learning suggests that gene expression is encoded in all
parts of a co-evolving interacting gene regulatory structure. *Nat. Commun.* 11
(2020).

56. N. H. Putnam, *et al.*, Chromosome-scale shotgun assembly using an in vitro method
for long-range linkage. *Genome Res.* 26, 342–350 (2016).

57. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* 25, 1754–1760 (2009).

58. M. Kokot, M. Dlugosz, S. Deorowicz, KMC 3: counting and manipulating k-mer
statistics. *Bioinformatics* 33, 2759–2761 (2017).

59. S. Chen, Y. Zhou, Y. Chen, J. Gu, "Fastp: An ultra-fast all-in-one FASTQ
preprocessor" in *Bioinformatics*, (Oxford University Press, 2018), pp. i884–i890.

60. Y. Miyazaki, *et al.*, Nitrogen as a key regulator of flowering in *Fagus crenata*:
understanding the physiological mechanism of masting by gene expression
analysis. *Ecol. Lett.* 17, 1299–1309 (2014).

61. A. Dobin, *et al.*, STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* 29, 15–
21 (2013).

62. B. Li, C. N. Dewey, RSEM: Accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* 12 (2011).

63. D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* 37,
907–915 (2019).

64.   T. Brůna, K. J. Hoff, A. Lomsadze, M. Stanke, M. Borodovsky, BRAKER2:
Automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS
supported by a protein database. *NAR Genom. Bioinform.* 3, 1–11 (2021).

65.   J. Keilwagen, F. Hartung, J. Grau, "GeMoMa: Homology-based gene prediction
utilizing intron position conservation and RNA-seq data" in *Methods in Molecular
Biology*, (Humana Press Inc., 2019), pp. 161–177.

66.   M. Pertea, G. Pertea, GFF Utilities: GffRead and GffCompare. *F1000Res.* 9 (2020).

67.   A. Bateman, *et al.*, UniProt: the Universal Protein Knowledgebase in 2025. *Nucleic
Acids Res.* 53, D609–D617 (2025).

68.   B. Buchfink, C. Xie, D. H. Huson, Fast and sensitive protein alignment using
DIAMOND. *Nat. Methods* 12, 59–60 (2014).

69.   R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, C. Kingsford, Salmon provides fast
and bias-aware quantification of transcript expression. *Nat. Methods* 14, 417–419
(2017).

70.   A. Hernández-Plaza, *et al.*, eggNOG 6.0: enabling comparative genomics across
12 535 organisms. *Nucleic Acids Res.* 51, D389–D394 (2023).

71.   C. P. Cantalapiedra, A. Herṇandez-Plaza, I. Letunic, P. Bork, J. Huerta-Cepas,
eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. *Mol. Biol. Evol.* 38, 5825–5829 (2021).

72.   V. Krishnakumar, *et al.*, Araport: The Arabidopsis information portal. *Nucleic Acids
Res.* 43, D1003–D1009 (2015).

73.   C. Camacho, *et al.*, BLAST+: Architecture and applications. *BMC Bioinformatics* 10
(2009).

74.   Y. Wang, *et al.*, MCScanX: A toolkit for detection and evolutionary analysis of gene
synteny and collinearity. *Nucleic Acids Res.* 40 (2012).

75.   A. Satake, *et al.*, Somatic mutation rates scale with time not growth rate in long-
lived tropical trees. *Elife* 12 (2024).

76.   H. Fan, A. R. Ives, Y. Surget-Groba, C. H. Cannon, An assembly and alignment-
free method of phylogeny reconstruction from next-generation sequencing data.
BMC Genomics 16 (2015).

77.   B. Mishra, *et al.*, A chromosome-Level genome assembly of the European beech
(*Fagus sylvatica*) reveals anomalies for organelle DNA integration, repeat Content
and distribution of SNPs. *Front. Genet.* 12 (2022).

78.   S. Falcon, R. Gentleman, Using GOstats to test gene lists for GO term association.
Bioinformatics 23, 257–258 (2007).

79.   Z. Yang, PAML 4: Phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* 24,
1586–1591 (2007).

933
934

 Figures and Tables

 Figure 1. Synteny and distribution of genomic features of *Q. glauca* and *L. edulis*
 genomes. (*A*) A phylogenetic tree of five Fagaceae species from three genera,
 Quercus, *Lithocarpus*, and *Fagus*. Bootstrap values for all branches are 100. (*B*)
 Synteny relationship between the *Q. glauca* and *L. edulis* genomes. The collinear
 blocks within the genomes were displayed by lines. Black lines show reverse
 alignment and gray lines represent regular. Numbers correspond to the 12
 chromosomes. (*C–D*) Circos plots of the *Q. glauca* (*C*) and *L. edulis* genomes (*D*).
 Lanes depict circular representation of chromosome length (Mb) (a), gene density
 (b), GC density (c), and repeat (d). Lines in the inner circle represent links between
 synteny-selected paralogs. (*E*) Gene copy-number variations between the *Q.*
 glauca and *L. edulis* genomes. The bar plot illustrates the number of genes in gene
 families categorized into 1:1 orthologs, N:N (many-to-many), 1:N (one-to-many),
 N:1 (many-to-one), and species-specific unique genes. (*F–G*) Mean gene
 expression levels in each copy number variation category for *Q. glauca* (*F*) and *L.*

Figure 2. Seasonal gene expression dynamics. (*A*) Hierarchical clustering of 213
gene expression profiles encompassing two tissues (leaf and bud) and four
species (*Q. glauca*, *Q. acuta*, *L. edulis*, and *L. glaber*), collected over two years.
(*B*) Distribution of daily mean temperature on each sampling date across clusters,
from L1 (top) to B3 (bottom). The Mann-Whitney U test was used to compare daily
mean temperature between cluster L1 and L2 (*p* = 0.136). The Steel-Dwass test
was applied for multiple comparisons across clusters B1–B3 (*p* < 0.05). Different
letters denote statistically significant differences (*p* < 0.05). (*C–D*) Hierarchical
clustering of 11749 genes based on the similarity of gene expression dynamics.
Daily mean temperature, flowering and leaf flushing phenology (proportion) are
shown in the upper three panels of the heatmaps.

Figure 3. Distribution of period and peak month in rhythmic gene expression. (*A*–*B*) Distribution of period length (month) among the genes with significant rhythmicity calculated by RAIN for leaf (*A*) and bud (*B*) samples of each species. (*C*–*D*) Distribution of seasonal peaks among the genes with annual periodicity. The angle of each bar represents the peak month, while the bar height indicates the number of genes. Purple and green squares around each rose plot denote the observed phenology of flowering and leaf flushing (proportion), respectively.

Figure 4. Comparison of peak months in seasonal gene expression across species in buds. (*A*) Intra-genus and (*B*) inter-genera comparisons of peak months in seasonal gene expression. The size and color of the circles represent the number of genes and the mean of Pearson's correlation, respectively. Genes with negative correlations lower than -0.3 are highlighted in blue, while the month of leaf flushing is highlighted in green. (*C*) Plot of molecular phenology divergence index *D* across months. Triangles and circles indicate the values of *D* calculated from intra-genus and inter-genera comparison, respectively. The line and shaded envelope indicate the mean and standard deviation (s.d.). Different letters denote statistically significant differences (Nemenyi test, *p* < 0.05).

Figure 5. Phylogenetic constraints in the evolution of seasonal gene expression
and the relationship between seasonal gene expression divergence and sequence
evolution. (*A*) Distribution of pairwise Pearson's correlation coefficients for gene
expression across species (Le: *L. edulis*, Lg: *L. glabe*r, Qg: *Q. glauca*, Qa: *Q.*
acuta; *n* = 11749). Colors indicate rhythmicity categories. Black bars show median
correlation values, and different letters indicate significant differences (Nemenyi
test, *p* < 0.05). (*B*) Heatmap of pairwise correlation coefficients for each gene, with
mean gene counts in four seasonal expression categories: (1) annual rhythmic
genes peaking in the growing season (March–November), (2) annual rhythmic
genes peaking in winter (December–February), (3) half-annual rhythmic genes,
and (4) arrhythmic genes. Genes are ordered by mean correlation across species
pairs. (*C–D*) Conserved (*C*) and diverged (*D*) gene expression patterns, based on
the top and bottom 5% of pairwise correlations. Solid lines and error bars indicate
mean ± SD. Numbers in parentheses show peak month differences between *Q.*
glauca and *L. edulis*. Peak month difference for *BFT* was not calculated due to
arrhythmic expression in *Q. glauca*. (*E*) Relationship between evolutionary rate
(dN/dS) and peak month difference (Δφ) for genes with annual periodicity in *Q.*
glauca and *L. edulis* (*n* = 951). Black lines and transparent bands represent
regression lines and 95% confidence intervals.

1007
1008

29