Evolution of gene expression in seasonal environments

Shuichi N. Kudo[1], Yuka Ikezaki[2], Junko Kusumi[3], Hideki Hirakawa[4,5], Sachiko Isobe[4,6], Akiko Satake[2]*

1. Graduate School of Systems Life Sciences, Kyushu University, Fukuoka 819-0395, Japan

2. Department of Biology, Faculty of Science, Kyushu University, Fukuoka 819-0395, Japan

3. Department of Environmental Changes, Faculty of Social and Cultural Studies, Kyushu University, Fukuoka 819-0395, Japan.

4. Kazusa DNA Research Institute, 2-6-7 Kazusa-kamatari, Kisarazu, Chiba 292-0818, Japan

5. Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University, Fukuoka 819-0395, Japan

6. Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, 113-8657, Japan

*Corresponding author: Akiko Satake

Email: satake.akiko.269@m.kyushu-u.ac.jp

Abstract

The biological activities of organisms are closely linked to seasonality. Phenology, the temporal orchestration of biological activities, is governed by gene expression, yet the evolutionary dynamics underlying seasonal gene expression remain unclear. To investigate these dynamics, we compared genome-wide expression dynamics (molecular phenology) in four dominant evergreen Fagaceae species in Asia (*Quercus glauca*, *Q. acuta*, *Lithocarpus edulis*, and *L. glaber*), using leaf and bud tissues over two seasonal cycles. We assembled high-quality reference genomes, identifying 11749 single-copy orthologous genes. Seasonal transcriptomic profiling of these orthologous genes revealed highly conserved gene expression across species in winter when temperatures fall below ~10°C. Rhythmic gene expression with significant periodic oscillations was more prevalent in buds (51.9%) than in leaves (40.6%), with most rhythmic genes (78.4–92.0%) exhibiting annual periodicity, while a smaller fraction (1.2–11.9%) followed half-annual cycles. The seasonal peaks of rhythmic genes were highly synchronized across species in winter but diverged during the growing season, reflecting species-specific timing of leaf flushing and flowering. These findings suggest that the four species share a common molecular calendar in winter, which constrains the evolution of gene expression under seasonal environments.

Impact Statement

A comparative analysis of genome-wide seasonal gene expression dynamics across four forest tree species revealed that a shared "molecular calendar" emerges in winter, constraining gene expression evolution and potentially limiting temporal niche partitioning and species divergence in seasonal environments.

Introduction

Fundamental biological processes, such as growth, mortality, and reproduction, rarely remain constant over time; instead, they exhibit pronounced variation in response to seasonal environmental changes (Fretwell, 1972). This temporal organization of a series of biological activities, known as phenology, plays a critical role in the evolution of life histories in response to seasonal fluctuating environments (Forrest and Miller-Rushing, 2010; Morellato et al., 2013; Piao et al., 2019; Schwartz, 2013). Phenology is regulated by the coordinated expression of genes that enable organisms to exhibit essential functions at the appropriate times (Kudoh, 2016; Satake et al., 2024a, 2022). However, gene expression dynamics underpinning seasonal responses in organisms remain not fully understood.

To explore the evolution of gene expression, transcriptome analyses are widely employed to compare transcript profiles across organs, sexes, species, and developmental stages within a given environment. Such studies have been conducted in mammals (Brawand et al., 2011; Cardoso-Moreira et al., 2019; Naqvi et al., 2019), *Drosophila* (Lemos et al., 2005; Nuzhdin et al., 2004), and Tanzanian cichlids (El Taher et al., 2021), providing key insights into the evolutionary dynamics of gene expression and its central role in shaping phenotypic diversity. These studies have revealed that gene expression profiles tend to be more

conserved across species within the same tissue, whereas the evolutionary rate of
gene expression varies among organs. In mammals, for instance, nervous tissue
exhibits a slower rate of gene expression change compared to the testis (Brawand
et al., 2011; Cardoso-Moreira et al., 2019; Naqvi et al., 2019), suggesting the
presence of tissue-specific functional constraints that influence the evolution of
gene expression.
In addition to tissue-dependent constraints, seasonal environmental
changes may play a crucial role in shaping the evolution of gene expression,
particularly by influencing the temporal coordination of gene expression under
natural conditions. Gene expression responses to certain seasons may be more
resistant to change, whereas in other seasons, divergence in gene expression
timing across species may occur, contributing to temporal niche differentiation.
Despite its potential importance, the influence of seasonal environmental
fluctuations on the gene expression evolution remains poorly understood, largely
because most transcriptomic analyses have been conducted under tightly
controlled laboratory conditions.
To investigate the evolution of gene expression in seasonal environments,
we performed a molecular phenology analysis and phenological observation of leaf
flushing and flowering in Fagaceae tree species, which are widely distributed
across the Northern Hemisphere, with its center of diversity in subtropical
Southeast Asia (Hipp et al., 2020; Kremer et al., 2012; Manos et al., 2001; Zhou et
al., 2022). Molecular phenology refers to transcriptome analysis conducted under
natural fluctuating conditions (Komoto et al., 2024; Kudoh, 2016; Nagano et al.,
2019; Satake et al., 2023) and provides a valuable framework for studying gene–
environment interactions and their role in orchestrating the temporal biological
functions essential for adaptation to environmental fluctuations (Satake et al.,
2022). In this study, we selected four evergreen Fagaceae species—two *Quercus*
species (*Q. glauca* and *Q. acuta*) and two *Lithocarpus* species (*L. edulis* and *L.
glaber*)—that co-occur in the same locality in Japan. *Q. glauca*, *L. edulis* and *L.
glaber* inhabit nearly identical environment, while *Q. acuta* is restricted to the high-
altitude regions with slightly colder environment compared to the other three
species. Leaf flushing and flowering phenology vary across species, with *Q. glauca*
and *Q. acuta* flowering in early spring, almost simultaneously with leaf flushing, *L.
edulis* flowering in late spring, and *L. glaber* flowering in autumn, approximately
one to two months after leaf flushing. This setting enabled us to compare seasonal
gene expression profiles across species with diverse phenology, both within and
between genera, under nearly identical environmental fluctuations. By assessing
and comparing seasonal transcriptomic dynamics across Fagaceae species, our
study provides new insights into the evolution of gene expression and adaptation
to seasonal natural environments.
105
Results
Genome assemblies of *Q. glauca* and *L. edulis*. To quantify transcript
abundance of non-model Fagaceae species, we first constructed reference
genomes for *Q. glauca*, as a representative of genus *Quercus*, and *L. edulis,* as a

representative of genus *Lithocarpus* (Figure 1A). To obtain high-quality
chromosome-level reference genomes, we conducted a hybrid genome assembly
for the HiFi reads and the high-throughput chromosome conformation capture (Hi-
C) reads (Supplementary file 1).The resultant scaffolds were 867.7 and 835.7 Mb
and consisted of 797 and 750 scaffolds including 12 chromosome-level scaffolds
with contig N50 of 71.7 Mb and 68.8 Mb for *Q. glauca* and *L. edulis*, respectively
(Supplementary file 1). Approximately 94.8% and 97.2% of the final assemblies
were assigned to 12 pseudochromosomes of *Q. glauca* and *L. edulis*, respectively
(Figure 1B). The number of chromosome-scale scaffolds were identical to the
known numbers in other *Quercus* (Zoldos et al., 1999) and *Lithocarpus* species
(Liu et al., 2023). The genomes were estimated to contain 39758 and 34059
protein-coding genes (33947 and 30246 genes were annotated within 12
chromosomes), covering 98.3% and 95.0% of complete Benchmarking Universal
Single Copy Orthologs (BUSCO) genes (eudicots_odb10) for *Q. glauca* and *L.
edulis*, respectively. The genome sizes of *Q. glauca* and *L. edulis* were estimated
to be 891.2 Mb and 868.7 Mb, respectively, using KmerGenie (Chikhi and
Medvedev, 2014) and GenomeScope2 (Ranallo-Benavidez et al., 2020).
Overall, the abundance and distribution of protein-coding genes, GC
content, and repeat sequences (Figure 1C, D*;* Supplementary file 1) and syntenic
structure (Figure 1B) were similar between *Q. glauca* and *L. edulis*. This pattern of
synteny conservation appears to be broadly shared across Fagaceae, except for
the genus *Fagus*, which shows a distinct genome structure (Ikezaki et al., 2025).
The homology between different chromosomes within the genome suggests that
small-scale segmental duplications likely occurred, as previously proposed
(Chen et al., 2014). Using OrthoFinder2 (Emms and Kelly, 2019), we clustered
genes from the two newly assembled genomes into orthologous groups. We
identified 13427 single-copy orthologs, enabling pairwise comparisons between
species (Figure 1E).
138
Seasonal gene expression dynamics. To characterize the evolution of seasonal
gene expressions, we generated a total of 483 transcriptome profiles every four
weeks over two years using leaf and bud tissues from *Q. glauca*, *Q. acuta*, *L.
edulis*, and *L. glaber* during 2021–2023. Reads from the two *Quercus* species were
mapped to the newly constructed *Q. glauca* reference genome, achieving a mean
mapping rate of 92.6 ± 2.2% and 88.6 ± 2.3% for *Q. glauca* and *Q. acuta*,
respectively. Similarly, reads from the two *Lithocarpus* species were mapped to
the newly constructed *L. edulis* reference genome, with a mean mapping rate of
89.3 ± 2.7% and 84.3 ± 5.4% for *L. edulis* and *L. glaber*, respectively. The number
of reads uniquely mapped on the reference genome was 19.9 million reads per
sample, on average. We used single-copy orthologous genes for further analyses
because these genes exhibited significantly higher average expression levels than
multi-copy genes (Steel-Dwass test, $p < 0.05$; Figure 1F, G). Among the single
copy orthologous genes, we excluded genes with low expression levels (mean of
Reads Per Kilobase (RPK) across samples for each species < 1) and normalized
the transcript count data using the GeTMM method (Smid et al., 2018). This

process resulted in a high-quality expression dataset for 11749 genes (Figure 2–
figure supplement 1A). For further analyses, we used the mean expression levels
across three biological replicates (three individuals per species), except for *L.*
edulis, where only one individual was available. We confirmed that expression
patterns of individual genes were highly consistent across all the three biological
replicates (Figure 2–figure supplement 2) in *Q. glauca* (Pearson's correlation r =
0.947), *Q. acuta* (r = 0.948) and *L. glaber* (r = 0.948).
To gain an initial overview of gene expression dynamics in a seasonal
environment, we assessed the relative magnitude of seasonal variation in
expression levels for each gene by calculating the standard deviation divided by
the mean for each gene (σ_i/μ_i) and compared it to the overall variation across all
genes in the genome (σ_g/μ_g) (Figure 2–figure supplement 1B). A prior study using
3025 yeast RNA-seq experiments reported that the dynamic range of individual
genes across experiments was considerably smaller than the overall genomic
variation under laboratory conditions (Zrimec et al., 2021). Consistently, our results
demonstrate that despite substantial seasonal fluctuations in environmental
conditions (Figure 2–figure supplement 3), individual gene expression remains
relatively conserved. On average, the magnitude of genome-wide variation was
15.8 times greater than the seasonal variation observed for individual genes
(Figure 2–figure supplement 1B).
To assess the similarity of gene expression profiles across tissues, species,
and seasons, we performed a hierarchical clustering of seasonal transcriptome
profiles from leaf and bud tissues across four species and identified five distinct
clusters (cluster L1, L2, B1–B3) according to the Elbow method (Figure 2A; Figure
2–figure supplement 4A; Supplementary file 2). Transcriptional profiles first
grouped into leaves (cluster L1 and L2) and buds (cluster B1–B3), irrespective of
seasons and species (Figure 2A), except that newly flushed leaf samples were
grouped closely with buds (Figure 2A; Figure 2–figure supplement 5). Within the
leaf cluster, transcriptome profiles were first clustered by genus and then by
species in *Lithocarpus* (cluster L1; Figure 2A). In contrast, samples from *Quercus*
species formed a mixture of two species (cluster L2; Figure 2A). Notably, the winter
samples of *Q. acuta*, collected during periods of low mean temperatures, were
distinctly separated from those collected in other seasons (cluster L2; Figure 2A).
Within the bud cluster, transcriptome profiles were predominantly grouped by
season—winter (cluster B3; November to February, with the exception of a March
sample from *L. edulis*; Supplementary file 2) versus the other growing seasons
(cluster B1 and B2; March to November; Supplementary file 2)—and subsequently
organized by genus and species. This suggests that the winter season exerts a
stronger influence on gene expression in buds than phylogenetic relationships. We
confirmed the robustness of our findings to differences in sampling periods
between *Quercus* and *Lithocarpus* species by reanalyzing expression data limited
to common sampling periods (Figure 2–figure supplement 6). Even when using
only the data from this shared period, the clustering patterns were consistently
reproduced. To assess whether the higher-elevation site of *Q. acuta* introduced
confounding environmental effects, we reanalyzed the data after excluding this

species. Hierarchical clustering still revealed that winter bud samples formed a
distinct cluster regardless of species identity (Figure 2–figure supplement 7),
consistent with our original findings.
To assess the biological relevance of the clusters identified, we tested
whether the temperatures and photoperiod at the time of sample collection differed
significantly among the clusters. Within the bud cluster, transcriptional profiles in
the winter cluster (cluster B3) were associated with significantly lower
temperatures—approximately below 10 °C—compared to those in other seasons
(Figure 2B; Figure 2–figure supplement 8A). In contrast, no such temperature-
related distinction was evident within the leaf cluster (Figure 2B). Additionally,
photoperiods were markedly shorter in the winter cluster in buds (Figure 2–figure
supplement 8B). These findings suggest that the buds of the four species exhibit
a conserved transcriptional response to environmental cues, namely cold
temperatures and short photoperiods, characteristic of the winter season. To
further investigate the relationships among all samples, we conducted a principal
component analysis (PCA). The first principal component (PC1) clearly separated
samples by tissue type, while the second (PC2) and the third principal component
(PC3) gradually differentiated winter samples from those collected during other
seasons (Figure 2–figure supplement 9).
To investigate seasonal expression dynamics of each gene, we conducted
hierarchical clustering on the transcriptional dynamics of 11749 genes and
identified five distinct gene sets characterizing different transcriptional modes
(gene set I–V; Figure 2C, D; Figure 2–figure supplement 4B; Supplementary file
3). Gene set I exhibited predominant expression in buds, with seasonal peaks
during summer, whereas gene set II was primarily expressed in leaves,
highlighting clear tissue-specificity (Figure 2C, D). Gene sets III and V
demonstrated genus-specific expression patterns, with higher expression levels
observed in *Lithocarpus* and *Quercus*, respectively (Figure 2C, D). In contrast,
gene set IV was expressed during winter across both leaf and bud tissues (Figure
2C, D).
230
Detection of rhythmic genes. To identify genes revealing seasonal expression
dynamics and determine the period and seasonal peaks in gene expression, we
applied the RAIN algorithm (Thaben and Westermark, 2014). This approach
enabled us to detect genes with rhythmic expression (hereafter rhythmic genes)
and analyze their periods and phases (peak months). Our analysis revealed that
44.6–58.8% of genes in bud tissues and 33.9–45.9% in leaf tissues exhibited
significant rhythmic expression across species (Figure 3–figure supplement 1A;
Supplementary file 4). Notably, bud tissues demonstrated a higher proportion of
rhythmic genes, with 51.9% across four species on average, compared to 40.6%
in leaf tissues (Figure 3–figure supplement 1A).
Among the rhythmic genes, the majority (78.4–92.0%) exhibited annual
periodicity, with a period of 12 ± 1 month observed across species and tissues
(Figure 3A, B, Supplementary file 4). Additionally, hundreds of genes (1.2–11.9%)
displayed half-annual periodicity (6 ± 1 month) with these genes being most

frequently observed in the bud samples of *L. edulis* (Figure 3A, B). The fraction of
genes exhibiting other periodicities was relatively small (5.7–14.1%; Figure 3A, B).
Based on the observed period distributions, we classified genes with periods within
the range of 8–16 months as exhibiting annual periodicity (96.0%) and those with
periods shorter than 8 months as having half-annual periodicity (4.0%), accounting
for detection noise (Figure 3–figure supplement 1A). Rhythmic genes with longer
periods (>17 months) were also identified, although their periodicity may be less
reliable due to the limited number of time points. Many genes with annual
periodicity were shared across all species, including 1280 genes in leaves and
1894 genes in buds (Figure 3–figure supplement 1B). In contrast, most genes
exhibiting half-annual periodicity were species-specific, with no genes shared
among all the species Figure 3–figure supplement 1C).
257
Peak month distribution of rhythmic genes and intra-genus and inter-genera
comparison. To determine the seasonal timing of gene expression peaks, we
focused on genes with annual or half-annual periodicity and visualized the density
and direction of these peaks throughout the year using a circular histogram (Figure
3C, D; Supplementary file 4). On average, 34.9% and 41.3% of annual rhythmic
genes peaked in winter (December–February) in leaf and bud samples,
respectively. Additionally, another peak corresponding to or occurring slightly after
leaf flushing and flowering events was observed, with its timing varying by species:
April in *Q. glauca* (15.6%), July in *Q. acuta* (15.3%), June in *L. edulis* (16.0%),
September in *L. glaber* (18.1%), for leaf tissues (Figure 3C). Similar peaks
corresponding to leaf flushing or flowering were also detected in bud samples,
occurring approximately one to two months earlier than in leaf samples (Figure
3D). For genes with half-annual periodicity, the majority (63.2 ± 4.8%) exhibited
expression peaks both in spring (March–May) and autumn (September–
November) in both tissues (Figure 3–figure supplement 1D; Supplementary file 4),
with the exception of *L. glaber*, an autumn flowering species. Interestingly,
WRKY22, a transcription factor regulating ethylene biosynthesis and senescence
(Zhu et al., 2024), exhibited half-annual periodicity in *L. glaber* (Figure 3–figure
supplement 1*E*). In addition, *SNF1-RELATED PROTEIN KINASE REGULATORY*
SUBUNIT GAMMA 1 (*KINγ1*), a subunit of SNF1-related kinase (*SnRK*) involved
in stress response such as starvation (Emanuelle et al., 2016), showed two distinct
peaks in *Q. glauca* and *L. edulis*, occurring in both summer (July) and winter
(January) (Figure 3–figure supplement 1E). This result suggests that both summer
and winter pose stressful conditions for these species.
To compare the peak month of gene expression with annual periodicity
across species, we quantified the number of genes within each peak combination
for both intra-genus (Figure 4A) and inter-genera species pairs in buds (Figure 4B)
and leaves (Figure 4–figure supplement 1A, B). During winter, a large number of
genes exhibited highly synchronized expression peaks across species, with high
pairwise Pearson's correlation coefficients observed (Figure 4A, B). In contrast,
the peak timing of gene expression from spring to autumn varied across species,
aligning with differences in leaf flushing and flowering times (Figure 4A, B). Some

genes exhibited pronounced divergence, showing negative Pearson's correlation
coefficients (< -0.3) between species. Notably, the number of such genes was
larger in inter-genera comparisons (11–77 genes; Figure 4B) than in intra-genus
comparisons (4–12 genes; Figure 4A). A similar pattern was observed in leaf
samples (Figure 4–figure supplement 1A, B). When comparing seasonal
expression peaks between tissues, the majority of genes were synchronously
activated in winter in both tissues (Figure 4–figure supplement 1C). During spring
to summer, slight variations in seasonal expression peaks were observed among
tissues (Figure 4–figure supplement 1C).
We quantified the degree of divergence in seasonal gene expression
patterns across species by calculating the peak month differences between two
species, where one species exhibits an expression peak in the given month. We
defined the molecular phenology divergence index (D) for each month by
quantifying the proportion of genes with peak month differences greater than 2
months between two species, relative to the total number of genes with expression
peaks in that month. From spring to autumn, the mean of D over all species pairs
in buds ranged from 0.47 to 0.66, but it declined sharply during winter, reaching a
minimum of 0.12 in January in buds (Nemenyi test, $p < 0.05$; Figure 4C). A similar
trend was observed in leaf samples, with another trough in summer (Figure 4–
figure supplement 1D). This seasonal trend remained consistent even after
excluding $Q. acuta$, which was sampled from a slightly cooler environment than
the other three species, from the analysis (Figure 4–figure supplement 2),
indicating that the observed patterns are not driven by environmental differences
associated with elevation. The values of D were not consistently smaller in inter-
genera pairs compared to intra-genus pairs. These findings suggest that there is a
seasonal constraint in gene expression in which the degree of molecular
phenology divergence is not constant but varies significantly across seasons, with
the least divergence occurring in winter.
318
Phylogenetic constraints in the evolution of seasonal gene expression. We
next investigated whether phylogenetic relationship influences the evolution of
seasonal gene expression by calculating pairwise Pearson's correlations of
seasonal expression for each gene across all species pairs (Supplementary file 5).
The median of Pearson's correlation coefficients across all genes were
significantly higher between species within the same genus compared to those
between species from different genera in both leaves (Steel-Dwass test, $p < 0.05$;
Figure 5–figure supplement 1A; Figure 5–figure supplement 2) and buds (Steel-
Dwass test, $p < 0.05$; Figure 5A; Figure 5–figure supplement 2). The intra-genus
comparison in genus $Lithocarpus$ exhibited the highest correlation in their seasonal
gene expression, despite their notable differences in flowering phenology, with $L.$
$edulis$ flowering in spring and $L. glaber$ in autumn. In contrast, the intra-genus
comparison in genus $Quercus$ resulted in a relatively low correlation in their gene
expression regardless of relatively similar phenology in leaf flushing and flowering.
This difference may be attributed to environmental factors, as $Q. acuta$ is the only

species found in the high-altitude study site with colder climates compared to the other three species.

Rhythmic genes exhibited more conserved expression patterns than arrhythmic genes in both tissues (Figure 5*A*; Figure 5–figure supplement 1A). In particular, genes with seasonal expression peaks in winter showed high correlation across species in both tissues (Figure 5B; Figure 5–figure supplement 1B). The proportion of rhythmic genes with annual periodicity declined as the mean correlation coefficient across species decreased, with a more rapid decline observed for genes peaking in winter compared to those with peak expression during the growing seasons (Figure 5B; Figure 5–figure supplement 1B). Additionally, the mean absolute difference in seasonal gene expression dynamics was significantly smaller between species within the same genus than between species from different genera in both tissues (Figure 5–figure supplement 3A, B). The relationship between the mean absolute difference and the proportion of rhythmic genes was less clear compared to that of the mean correlation coefficient (Figure 5–figure supplement 3C, D). These findings suggest that, in addition to seasonal constraints, the overall dynamics of seasonal gene expression are also shaped by phylogenetic constraints.

Functional annotation of genes with conserved or divergent expression across species. To functionally annotate genes with conserved or divergent expression across species, we extracted genes with the top and bottom 5% based on the distribution of mean correlation (Figure 5B; Figure 5–figure supplement 1B) and conducted Gene Ontology (GO) enrichment analysis. GO analysis revealed that the top 5% of genes in leaves were associated with photosynthesis, while those in buds were linked to cell fate determination (Figure 5–figure supplement 4A; Supplementary file 6). The gene with the highest correlation in leaves was *STRESS RESPONSE SUPPRESSOR1* (*STRS1*; mean of pairwise correlation = 0.91; Figure 5C), a DEAD-box RNA helicase previously reported to enhance tolerance to abiotic stresses such as cold (Kant et al., 2007). In buds, the most highly correlated gene was the histone deacetylase *HDA19* (mean of pairwise correlation = 0.93; Figure 5C), which plays a key role in environmental stress responses through histone modification and chromatin remodeling (Chen and Wu, 2010). These findings suggest that genes associated with photosynthesis during the growing season and stress responses during winter exhibit highly conserved seasonal expression patterns across species.

The bottom 5% of genes were predominantly associated with DNA replication and cell division in leaves (Figure 5–figure supplement 4B), aligning with the divergent leaf phenology observed (Figure 3B). In contrast, GO terms related to immune response were weakly enriched in buds (Figure 5–figure supplement 4B). Notably, we identified MADS-box genes linked to floral transition among the bottom 5% of genes: *BROTHER OF FT AND TFL1* (*BFT*) in leaves (mean of pairwise correlation = -0.094) and *FRUITFULL* (*FUL*) in buds (mean of pairwise correlation = -0.069). *BFT*, a floral repressor (Yoo et al., 2010), was expressed in autumn in the spring-flowering species *L. edulis* and in spring in the

autumn-flowering species *L. glaber* leaves (Figure 5D). Meanwhile, *FUL*, a positive
regulator of floral transition (Smaczniak et al., 2012), was expressed in spring in *L.*
edulis and in autumn in *L. glaber* (Figure 5D), mirroring the timing of flowering. In
Q. glauca and *Q. acuta*, *FUL* was expressed in winter (Figure 5D), nearly out of
phase with their expressions in *Lithocarpus* species. These results suggest that
genes involved in reproductive processes exhibit species-specific divergence in
expression patterns in response to seasonal environmental changes.
386
Relating seasonal gene expression divergence to sequence divergence. If
divergence of seasonal timing of gene expression is related to selective constraint,
we would expect a correlation between sequence divergence patterns and the
divergence in seasonal expression peaks. To explore the relationship between the
divergence of seasonal gene expression and protein sequence evolution, we
calculated synonymous (dS) and non-synonymous substitution rates (dN) and their
ratio (dN/dS) for each gene using coding sequences from newly assembled
genomes of *Q. glauca* and *L. edulis*. We first confirmed that the highly expressed
genes evolve more slowly by identifying a significant negative correlation between
evolutionary rate (dN/dS) and mean of the gene expression (r = -0.21, p = 1.8 ×
10^{-118} for dN/dS; Figure 5–figure supplement 5A; Supplementary file 7). This
finding aligns with previous studies in yeasts, mammals, and plants (Drummond et
al., 2005; Gaut et al., 2011; Liao and Zhang, 2006; Yang and Gaut, 2011). By
targeting the genes that exhibited annual periodicity across both species and
tissues, we examined the relationship between these evolutionary measures and
peak month differences in seasonal gene expression between species ($\Delta\varphi$) in
leaves and bud tissues using linear regression. Assessment of the significance of
the regression coefficient using a *t*-test revealed that genes with smaller peak
month differences between species exhibited lower values of dN/dS in buds (p
=2.6 × 10^{-2}) and marginally lower values of dN/dS in leaves (p = 5.7 × 10^{-2}). The
lower values were also found in dN in both leaves (p = 8.1 × 10^{-3}) and buds (p =
7.8 × 10^{-3}), whereas no significant trend was observed for dS in both tissues
(Figure 5E; Figure 5–figure supplement 5B, C; Supplementary file 7). We further
investigated whether the association between the divergence in seasonal
expression peaks and sequence divergence patterns varies between genes
expressed in winter and in the growing season. We classified genes that exhibited
annual periodicity across both species and tissues into four groups based on their
expression peak timing (winter or growing seasons) and tissue specificity (leaves
or buds) and compared the extent of protein sequence divergence across these
groups. We found no significant differences in dN/dS (Kruskal Wallis test, p = 2.8
× 10^{-1}; Figure 5–figure supplement 5*D*; Supplementary file 7) between genes
expressed in winter in both tissues and those in growing seasons. The same result
was observed even when focusing only on genes with conserved peaks (peak
month difference < 2), with no significant differences in dN/dS between these gene
sets (Kruskal Wallis test, p = 3.5 × 10^{-1}; Figure 5–figure supplement 5E;

Supplementary file 7). These results suggest that coding-sequence variation is
unlikely to be the primary driver of seasonal gene expression patterns.
424

Discussion

A central goal in biology is to unravel the molecular basis of phenotypic evolution,
with gene expression serving as a fundamental driver of this process. In this study,
we investigated the impact of seasonal environmental fluctuations on the evolution
of gene expression and its relationship with phenological traits in four dominant
Fagaceae tree species in forest ecosystems. We found that gene expression
profiles were more resistant to change during winter compared to the growing
season, indicating the presence of a seasonal constraint on the evolution of gene
expression. This constraint becomes particularly pronounced in buds during winter
when temperatures fall below approximately 10°C (Figure 2B). This temperature
threshold aligns with previous findings, where winter-specific transcript profiles
were reported at temperatures below 8.7–11.3°C in cherry trees (Miyawaki‑
Kuwakado et al., 2024), 11.3°C in ever-green *Fagaceae* (Satake et al., 2023), and
10.5°C in perennial *Arabidopsis* (Aikawa et al., 2010). These results suggest the
presence of a conserved mechanism for gene expression regulation under low-
temperature environments across different species. The greater conservation of
gene expression patterns in buds compared to leaves across species may be
attributed to their functional role protecting essential meristematic tissues,
including the shoot apex and leaf and flower primordia. Additionally, buds likely
play a crucial role in maintaining common physiological functions necessary for the
resumption of growth in spring. This functional consistency may impose
evolutionary constraints on gene expression, contributing to its conservation
across species.
It is intriguing to compare the seasonal constraints identified in this study
with developmental constraints in gene expression evolution, where evolutionary
conservation and variation across developmental stages during the life cycle have
been observed, such as in the early conservation model and the developmental
hourglass model (Figure 5–figure supplement 6A)(Irie and Kuratani, 2011;
Lotharukpong et al., 2024; Quint et al., 2012). Previous studies have demonstrated
that certain developmental stages exhibit remarkably similar gene expression
patterns (Drost et al., 2016; Kalinka et al., 2010; Levin et al., 2016). Our findings
highlight that such highly similar gene expression patterns are observed during the
winter season (Figure 5–figure supplement 6B), suggesting that a common
physiological state is essential for enduring harsh winter conditions. This seasonal
constraint on gene expression evolution may have the potential to limit temporal
niche partitioning at the phenotypic level and slow species divergence rates in
seasonal environments (Cleland and Wolkovich, 2024; Kubo and Iwasa, 1996).
Although completely disentangling the effects of developmental stages and
seasonal changes on gene expression is challenging when using molecular
phenology data collected from natural environments, the observation that species
with different phenology—i.e., different developmental timing in seasonal
environments—exhibit convergent gene expression patterns during winter

underscores the profound influence of winter environmental conditions on the
evolution of gene expression. Future research is needed to elucidate the
mechanisms driving the establishment of such conserved gene expression
patterns, including the dynamic reorganization of chromatin architecture in
response to cold (Fischl et al., 2020; Nishio et al., 2020; Zhang et al., 2023).
We found that the evolution of coding-sequences was only weakly
correlated with the divergence of seasonal peaks in gene expression across
species. To understand the evolutionary mechanism underlying the seasonal
constraint on gene expression that results in similar expression patterns during
winter, an important future direction is to explore the evolution of non-coding
sequences that modulate gene expression levels in response to seasonal
environmental changes. Regulatory variation has been recognized as a major
driver of evolutionary innovation (Prud'homme et al., 2007; True and Haag, 2001).
Recent deep learning studies utilizing genome and transcriptome data from yeast
and humans have highlighted the significance of co-evolution between coding and
non-coding sequences in shaping gene expression levels (Zrimec et al., 2021,
2020). Understanding how seasonal environments drive the co-evolution of coding
and regulatory sequences, as well as the divergence of molecular phenology and
its phenological outputs, will require expanding analyses to a broader range of
species. Such efforts will facilitate the development of predictive models for
evolutionary processes influenced by seasonal dynamics. Comparison of
molecular phenology across wide climatic zones, including both tropical and
temperate zones, will also be effective to identify key genes for the phenological
response to on-going climate change.
491
492 **Materials and Methods**

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Addition al informat ion**
biological sample (*Quercus. glauca*)	Leaf	This paper		Freshly isolated from *Q. glauca*
biological sample (*Quercus. glauca*)	Bud	This paper		Freshly isolated from *Q. glauca*

biological sample (*Quercus. acuta*)	Leaf	This paper		Freshly isolated from *Q. acuta*
biological sample (*Quercus. acuta*)	Bud	This paper		Freshly isolated from *Q. acuta*
biological sample (*Lithocar pus. edulis*)	Leaf	This paper		Freshly isolated from L. *edulis*
biological sample (*Lithocar pus. edulis*)	Bud	This paper		Freshly isolated from *L. edulis*
biological sample (*Lithocar pus. glaber*)	Leaf	This paper		Freshly isolated from L. *glaber*
biological sample (*Lithocar pus. glaber*)	Bud	This paper		Freshly isolated from *L. glaber*
commerci al assay or kit	QIAamp DNA Mini Kit	Qiagen	Cat. #: 51304	
commerci al assay or kit	SMRTbell Express Template Prep Kit 2.0	PacBio	Cat. #: 100-938-900	
commerci al assay or kit	Sequel II Binding Kit 2.0	PacBio	Cat. #: 101-789-500	

commercial assay or kit	PureLink™ Plant RNA Reagent	Thermo Fisher Scientific	Cat. #: 12322012	
commercial assay or kit	TURBO DNA-free kit™	Thermo Fisher Scientific	Cat. #: AM1907	
commercial assay or kit	NEBNext Ultra II RNA Library Prep Kit for Illumina	New England BioLabs	Cat. #: E7770	
commercial assay or kit	NEBNext Ultra II DNA Library Prep Kit	New England BioLabs	Cat. #: E7645	
commercial assay or kit	NEBNext Multiplex Oligos for Illumina	New England BioLabs	Cat. #: E7335, E7500, E7710	
commercial assay or kit	RNeasy Plant Mini Kit	Qiagen	Cat. #: 74904	
commercial assay or kit	Agilent RNA 6000 Nano kit	Agilent Technologies	Cat. #: 5067-1511	
commercial assay or kit	High Sensitivity DNA kit	Agilent Technologies	Cat. #: 5067-4626	
commercial assay or kit	NEBNext Library Quant Kit for Illumina	New England BioLabs	Cat. #: E7630	
chemical compound, drug	RNA-stabilizing reagent (RNAlater)	Thermo Fisher Scientific	Cat. #: AM7021	
software, algorithm	Hifiasm1	Hifiasm	Hifiasm1 v0.15.4-r347	

software, algorithm	blobtools2	BlobToolKit	RRID:SCR_023351	
software, algorithm	purge_dups3	purge_dups	RRID:SCR_021173	
software, algorithm	HiRise	Putnam et al., 2016	RRID:SCR_023037	
software, algorithm	fastp	Chen et al., 2018	RRID:SCR_016962	
software, algorithm	STAR	Dobin et al., 2013	RRID:SCR_004463	
software, algorithm	edgeR	edgeR	RRID:SCR_012802	
software, algorithm	Braker2	Brůna et al., 2021	RRID:SCR_018964	
software, algorithm	GeMoMa	Keilwagen et al., 2019	RRID:SCR_017646	
software, algorithm	GFFcompare	Pertea and Pertea, 2020	DOI: 10.12688/f1000research.23297.1	
software, algorithm	DIAMOND	Buchfink et al., 2014	RRID:SCR_016071	
software, algorithm	Salmon	Patro et al., 2017	RRID:SCR_017036	
software, algorithm	eggNOG-mapper	Cantalapiedra et al., 2021	RRID:SCR_021165	

software, algorithm	BLASTp	Camacho et al., 2009	RRID:SCR_001010	
software, algorithm	MCScanX	Wang et al., 2012	RRID:SCR_022067	
software, algorithm	GenomeScope2	Ranallo-Benavidez et al., 2020	DOI:10.1038/s41467-020-14998-3	
software, algorithm	KmerGenie	Chikhi and Medvedev, 2014	DOI:10.1093/bioinformatics/btt310	
software, algorithm	OrthoFinder2	Emms and Kelly, 2019	RRID:SCR_017118	
software, algorithm	R	R project	RRID:SCR_001905	
software, algorithm	PAML	Yang, 2007	RRID:SCR_014932	

Sample collection for genome sequencing

495 To construct genomes of *Q. glauca* and *L. edulis*, we collected approximately 5g
496 of fresh leaves from one individual in spring, respectively. We shipped these
497 samples to Dovetail Genomics, LLC (Scotts Valley, CA, USA) where Dovetail Staff
498 performed DNA extraction, library preparation, sequencing, and assembly steps.

Genome sequencing and assembly

DNA extraction was performed using the QIAamp DNA Mini Kit (Qiagen,
Germantown, MD, USA) using the manufacturer's recommended protocol. Mean
fragment length of the extracted DNA was 60 kbp for *Q. glauca* and 65 kbp for *L.*
edulis. DNA samples were quantified using Qubit 2.0 Fluorometer (Life
Technologies, Carlsbad, CA, USA). The library preparation, sequencing and
scaffolding were carried out by Dovetail Genomics (California, USA) according to
their standard genome assembly pipeline (https://dovetailgenomics.com/). The
PacBio SMRTbell library (~20kb) for PacBio Sequel was constructed using
SMRTbell Express Template Prep Kit 2.0 (PacBio, Menlo Park, CA, USA) using
the manufacturer's recommended protocol. The library was bound to polymerase

using the Sequel II Binding Kit 2.0 (PacBio) and loaded onto PacBio Sequel II.
Sequencing was performed on PacBio Sequel II 8M SMRT cells. PacBio CCS
reads were used as an input to Hifiasm1 v0.15.4-r347 with default parameters. For
Q. glauca, we also generated sequence data using PacBio Revio. BLASTn results
of the contigs obtained by Hifiasm assembly against the NCBI's NT database were
used as input for blobtools2 v1.1.1 and contigs identified as possible contamination
were removed from the assembly. Finally, purge_dups3 v1.2.5 was used to
remove haplotigs and contig overlaps.
519
Assembly scaffolding with HiRise
A proximity ligation library was generated by the Dovetail's Omni-C technique,
followed by sequencing on an Illumina HiSeqX platform. Chromatin was fixed in
place in the nucleus with formaldehyde before extraction
(https://dovetailgenomics.com/wp-content/uploads/2021/09/Omni-C-Tech-
Note.pdf). Fixed chromatin was digested with DNase I, and the fragmented
chromatin ends were repaired and biotinylated to adapters followed by proximity
ligation. Crosslinks were then reversed, the DNA purified, and the biotin
subsequently removed. The DNA library was prepared and sequenced to produce
2 × 150 bp paired-end reads (around 30× coverage). The Omni-C technology uses
a sequence-independent endonuclease which provides even, unbiased genome
coverage. The input *de novo* assembly and Omni-C library reads were used as
input data for HiRise, a software pipeline designed specifically for using proximity
ligation data to scaffold genome assemblies (Putnam et al., 2016). The Omni-C
library sequences were aligned to the contigs assembled by Hifiasm using bwa (Li
and Durbin, 2009). The separations of Omni-C reads from paired-end mapped
within the contigs were analyzed by HiRise to produce a likelihood model for
genomic distance between paired-end reads, and the model was used to identify
and break putative misjoins, to score prospective joins, and make joins above a
threshold.
540
Sample collection for RNA sequencing
To monitor the genome-wide gene expression profiles, we collected leaf, bud, and
flower samples from four Fagaceae species, *Q. glauca*, *Q. acuta*, *L. edulis* and *L.*
glaber, every four weeks for two years (October 2019 to October 2021 for the
Lithocarpus species and April 2021 to April 2023 for the *Quercus species*) across
Fukuoka and Saga prefectures in Kyushu, Japan. We targeted three individuals,
and each sample was collected from three branches per individual, except for *L.*
edulis. Given the consistent expression patterns observed among individuals in a
previous study (Satake et al., 2023), we focused on only one individual for *L. edulis*.
Study sites for *Q. glauca* is the biodiversity reserve on the Ito campus of Kyushu
University (33º35′ N, 130º12′ E, 20 to 57 m a.s.l). *L. edulis* and *L. glaber* were
studied at the Imajuku Field Activity Center (33º33′ N, 130º16′ E, 84 to 111 m a.s.l).
The Imajuku and Ito Campus study sites, located only 7.3 km apart, experience
nearly identical temperature conditions. *Q. acuta* was monitored in Sefuri
mountains (33º26′ N, 130º22′ E, 970 to 974 m a.s.l), a higher-altitude, cooler site.

Environmental data from Ito Campus and the Imajuku Field Activity Center were
obtained from the Japan Meteorological Agency, while data from Sefuri Mountain
were collected using a logger installed at the sampling site. Additionally, to assess
the potential impact of different sampling periods between the *Lithocarpus* and
Quercus species on the gene expression analysis, we conducted an additional
year of sampling for *L. edulis* and *L. glaber*, collecting leaf and bud samples from
November 2021 to November 2022, and used the samples collected within the
same period for all the four species (May 2021 to November 2022).
Samples for RNA-seq were taken from the sun-exposed crown
(approximately 2–5 m from the ground) using long pruning shears from 11:30 to
12:30 h. For each sample, 0.2–0.4 g of tissue was preserved in a 2 ml microtube
containing 1.5 ml of RNA-stabilizing reagent (RNAlater; Thermo Fisher Scientific,
Waltham, MA, USA) immediately after harvesting. The samples were transferred
to the laboratory within 2 hr after sampling, stored at 4°C overnight and then stored
at −80°C until RNA extraction. During transport to the laboratory, the samples were
kept in a cooler box with ice to maintain a low temperature.
572
RNA extraction and RNA-seq analysis
RNA was extracted independently from bud and leaf samples of each tree. Total
RNA of the leaf sample was extracted in accordance with the method described in
a previous study (Miyazaki et al., 2014). Total RNA of the bud sample was
extracted using the PureLink™ Plant RNA Reagent (Thermo Fisher Scientific,
Waltham, MA, USA) and subsequently processed with DNase using the TURBO
DNA-free kit™ (Thermo Fisher Scientific) according to their protocols and purified
using the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) according to its RNA
cleanup protocol. RNA integrity was examined using the Agilent RNA 6000 Nano
kit on a 2100 Bioanalyzer (Agilent Technologies), and the RNA yield was
determined using a NanoDrop ND-2000 spectrophotometer (Thermo Fisher
Scientific). Five to six micrograms of RNA extracted from each sample were sent
to Hangzhou Veritas Genetics Medical Institute Co., Ltd., where a cDNA library
was prepared with a NEBNext Ultra II RNA Library Prep Kit for Illumina, and 150
base paired-end transcriptome sequencing of each sample was conducted using
an Illumina NovaSeq6000 sequencer (Illumina).
The quantification of gene expression followed a three-step process. First,
quality filtering was performed using the fastp v0.23.2 with default options: phred
quality ≥15, the maximum number of N base in one read = 5 (Chen et al., 2018).
Second, RNA-seq reads were mapped to the reference genome using STAR
v2.7.10b (Dobin et al., 2013). Third, RNA-seq reads were quantified using RSEM
v1.3.1 (Li and Dewey, 2011). We excluded the genes with low expression level
using the threshold, the value of mean Reads Per Kilobase (RPK) < 1 for each
species. Then, we further normalized the expression level using the value of Gene
length corrected Trimmed Mean of M-values (GeTMM) (Smid et al., 2018) using
an R package "edgeR" v3.42.4. For the downstream analysis we calculated the
mean expression levels of GeTMM of each gene across multiple individuals and
transformed them to $\log_2(\text{GeTMM}+1)$. To confirm that the effect of the difference

in sampling period is negligible, we added the RNA-Seq data of *L. edulis* and *L.*
glaber collected from December 2021 to December 2022. We extracted the same
genes in the existing data that curated based on the mean RPK values across two-
year sampling periods. Then we further normalized the expression level using
GeTMM.
606
Genome annotation
The RNA-seq reads of *Q. glauca* (8 samples obtained by a previous study (20)
collected from May to December 2017; GEO's accession number: GSE211384)
and *L. edulis* (20 samples covering leaf, bud and flower tissues collected in
January, April, July, and November 2021) were respectively mapped against the
scaffolds of *Q. glauca* and *L. edulis* obtained by HiRise with HISAT2 v2.1.0 (Kim
et al., 2019). The bam files generated by the mappings and amino acid sequences
of the 9 species obtained from NCBI (*Castanea mollissima*: GCA_000763605.2
(NCBI's accession number), *Q. lobata*: GCF_001633185.1, *Q. robur*:
GCF_932294415.1, *Q. suber*: GCF_002906115.1, *Carpinus fangiana*:
GCA_006937295.1, *Carya illinoinensis*: GCF_018687715.1, *Juglans macrocarpa*
× *J. regia*: GCF_004785595.1, *Juglans regia*: GCF_001411555.2, *Morella rubra*:
GCA_003952965.2) were used for ab initio gene prediction by Braker2 v2.1.6
(Brůna et al., 2021), and the genes with highest-score were selected to exclude
splicing variants. In parallel, the gene models were respectively constructed in *Q.*
glauca and *L. edulis* by GeMoMa v1.7.1 (Keilwagen et al., 2019) with RNA-seq
reads and amino acid sequences used for the gene prediction in Braker2, and the
genes with highest-score were selected to exclude splicing variants. The
duplicated genes between Braker2 and GeMoMa were excluded by GFFcompare
v0.11.2 (Pertea and Pertea, 2020). The remaining genes were searched against
UniProtKB (Bateman et al., 2025) by DIAMOND v2.0.5 (Buchfink et al., 2014) in
'more-sensitive' option with E-value ≤10^{-20} and identity ≥25%. Next, gene
expression analysis was performed by Salmon v1.4.0 (Patro et al., 2017), and the
genes with TPM values >0.1 were considered as expressed. The orthologous
genes were searched against EggNOG (Hernández-Plaza et al., 2023) using
eggNOG-mapper v2.1.8 (Cantalapiedra et al., 2021) with E-value ≤10^{-10}. The
genes were searched against protein sequences of *Castanea mollissima*
(Accession: GCA_000763605.2), *Quercus lobata* (GCF_001633185.2), and
Arabidopsis thaliana (Araport11 (Krishnakumar et al., 2015)) by BLASTp v2.13.0
(Camacho et al., 2009) with E-value ≤10^{-20}. According to these similarity searches,
the genes whose product names contain keywords related to transposable
elements (TEs) were categorized into TE. The genes having similarity hits and
being expressed by Salmon were classified into high-confidence (HC) genes, while
remaining genes were categorized into low-confidence (LC) genes. The genes
categorized into HC were used for further analyses.
642
Genome synteny analysis
To investigate the syntenic relationship between *Q. glauca* and *L. edulis*, we first
performed an all-vs-all BLASTp search of the protein sequences, using a cut-off

E-value of 10^{-5}. Collinearity blocks within and between the genomes were then
identified using MCScanX (Wang et al., 2012) based on the results of the BLASTp
searches. Finally, the collinearity blocks were visualized using SynVisio
(https://synvisio.github.io).
650
Illumina short-read sequencing and library preparation
For genome size estimation and phylogenetic reconstruction, we also generated
short-read sequencing of *Q. glauca*, *Q. acuta*, *L. edulis*, and *L. glaber.* DNA
extraction was performed using a modified version of the method described
previously (Satake et al., 2024b). The DNA samples were sent to Macrogen Inc.
(Republic of Korea) for sequencing on the Illumina HiseqX platform (Illumina, San
Diego, CA, USA). DNA was sheared to around 500 bp fragments in size using
dsDNA fragmentase (New England BioLabs, Ipswich, MA, USA). Library
preparation was performed using the NEBNext Ultra II DNA Library Prep Kit (New
England BioLabs) according to the manufacturer's protocol, and the libraries were
individually indexed with the NEBNext Multiplex Oligos for Illumina (New England
BioLabs) by PCR. The quality and quantity of each amplified library were analyzed
using the Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA, USA), the
High Sensitivity DNA kit (Agilent Technologies), and the NEBNext Library Quant
Kit for Illumina (New England BioLabs).
666
Genome size estimation
The genome size of *Q. glauca* was estimated by GenomeScope2 v2.0 (Ranallo-
Benavidez et al., 2020) using *k*-mer frequency plot obtained by kmc v3.2.4 (Kokot
et al., 2017) with *k*-mer size 61 against the HiFi reads. We also used pair-end short
reads for genome size estimation of *Q. glauca* and confirmed that resulted
estimates were not different. For *L. edulis*, paired-end short reads from four
individuals were merged and quality-trimmed using fastp v0.23.2 (Chen et al.,
2018). The genome size was then estimated using KmerGenie (Chikhi and
Medvedev, 2014) with a k-mer size of 55 applied to the trimmed reads.
676
Phylogenetic reconstruction
We constructed a phylogenetic tree using an alignment-free method (Fan et al.,
2015) based on short-read data from *Q. acuta* and *L. glaber* and genome assembly
data from *Q. glauca* and *L. edulis*. Additionally, genomic data for *Fagus sylvatica*
was obtained from beechgenome.net (Mishra et al., 2022) and included as an
outgroup. Due to the large volume of short-read data for *Q. acuta* and *L. glaber*
(18 Gb each), we randomly subsampled half of the reads. Bootstrap values were
estimated through 100 resampling of the k-mer table.
685
Identification of orthologs
The prediction of orthogroups was based on a BLASTp all-against-all comparison
of the protein sequences (E-value < 10^{-5}) of 5 Fagaceae species: *L. edulis*, *Q.*
glauca, *Q. robur*, *Fagus sylvatica* and *Juglans regia*. Clustering was conducted
with OrthoFinder2, using version 2.2.6 for BLAST data generation and version

2.5.5 for orthogroup determination (Emms and Kelly, 2019). The genomic data of
Q. robur and *J. regia* were downloaded from NCBI (dhQueRobu3.1 and Walnut
2.0), while *F. sylvatica* data were sourced from beechgenome.net (Mishra et al.,
2022). Since the *F. sylvatica* dataset contained numerous predicted transposable
elements (TEs), we re-annotated genes using three databases—EggNOG5
(EggNOG-mapper), Pfam35.0 (HMMER v3.3.2), and UniRef90_2023_0628
(DIAMOND v2.1.8.162)—and excluded genes identified as TEs in at least one
database. In total, we identified 26692 orthogroups and focused on single-copy
orthogroups after filtering out genes mapped outside the 12 chromosomes. This
resulted in a final dataset of 11749 genes for further analysis (Data S2).
701
Hierarchical clustering and principal component analysis (PCA)
To assess the similarity of the genome-wide transcriptional profiles across
orthologous genes and 483 samples collected in different seasons, we performed
hierarchical clustering using the monthly time series data of 11749 orthologous
genes from October 2019 to April 2023. For each orthologous gene, there were 26
or 27 time points, with three individuals each for *Q. glauca*, *Q. acuta*, *L. glaber* and
one individual for *L. edulis*. We calculated the mean expression levels of each
orthologous gene across three individuals in each species and subsequently
normalized the values by adjusting the mean to zero and the standard deviation to
one. We performed hierarchical clustering using the Ward method and the
Euclidian distance using the pheatmap function in R (v 1.0.12). To determine the
number of the clusters, we calculated Within-Cluster-Sum of Squared Error (WSS)
and determined the number of clusters as an elbow point of it (Figure 2–figure
supplement 4). To assess the seasonal expression dynamics of 11749 orthologous
genes, we also performed PCA of the gene expression profiles from all samples
using the prcomp function in R (v 4.3.1).
718
Identification of genes with rhythmic expression
To identify genes with rhythmic expression and determine the period and peak
month of gene expression, we applied the RAIN algorithm (Thaben and
Westermark, 2014) using the rain function in R (v 1.34.0). We set the following
settings: deltat = 1, period = 12, period.delta = 11, peak.border = (0,1), nr.series =
1, method= "independent", na.rm= T , adjp.method = "ABH". Based on the results
of the RAIN analysis, we identified genes with significant periodicity using a
Benjamini-Hochberg (BH) adjusted *q*-value threshold of < 0.01. Since the period
length and peak month in the RAIN output are expressed in terms of the number
of time points, we converted them into months using the sampling span (four
weeks) and sampling dates. Subsequently, we categorized the rhythmic genes into
three categories based on the period length *T* (in months): "half-annual" ($0 \leqq T <$
8), "annual" ($8 \leqq T \leqq 16$), and "long" ($T > 16$).
732
Calculation of Pearson's correlation coefficient across species
To quantify the similarity of seasonal gene expression pattern across species, we
calculated pairwise Pearson's correlations of seasonal expression for each gene

across all the species pairs. For species within the same genus, the monitoring
period was identical, allowing direct comparison. On the other hand, the monitoring
periods differed between *Quercus* and *Lithocarpus* species: *Q. glauca* and *Q.*
acuta were monitored from May 2021 to April 2023, while *L. edulis* and *L.*
glaber were monitored from October 2019 to October 2021 (Figure 2–figure
supplement 3). To ensure comparable time series data for inter-genera
comparisons, we extracted the time series from *Q. glauca* and *Q. acuta* spanning
May 19, 2021, to November 2, 2022, and from *L. edulis* and *L. glaber* spanning
May 13, 2020, to October 20, 2021, for correlation analysis. The mean absolute
differences for each gene across all the species pairs were also calculated in a
similar manner.
747
Molecular phenology divergence index (*D*)
To quantify the seasonal difference of peak month divergence across species, we
defined the molecular phenology divergence index (*D*) for each month *t* and
species pair A-B as the proportion of genes that peak in month *t* in the reference
species A but exhibit a peak month difference ($\Delta\varphi$) greater than two months in
species B using following formula:
754
755 $$D := \frac{\text{\# Genes with peak month } t \text{ in species A and } t \pm \Delta\phi (\geq 2) \text{ in species B}}{\text{\# Genes with peak month } t \text{ in species A}}.$$
756
Gene Ontology (GO) enrichment analysis
First, we assigned GO terms for each gene in *Lithocarpus* and *Quercus* genomes
based on the annotation data of *Arabidopsis thaliana* orthologs downloaded from
PLAZA Database (http://bioinformatics.psb.ugent.be/plaza/). Then, GO
enrichment analysis was conducted using the GOstats (Falcon and Gentleman,
2007) function in R (v 2.66.0) with the following settings: ontology = BP,
pvalueCutoff = 0.05, conditional = TRUE, and testDirection = over. The universe
gene set was defined as genes associated with GO terms common to both of the
genomes. Genes without annotated GO terms were removed from the universe
set, and those with differing GO term annotations between the two genomes were
also excluded. The BH adjusted *q*-values were calculated from the *p*-values
generated by GOstats to account for multiple testing.
769
Phenology measurement: leaf unfolding and flowering
To compare seasonal changes in genome-wide gene expression profiles with
phenology, we conducted observations of leaf unfolding and flowering phenology
on the three terminal branches of each of the three individuals used for gene
expression monitoring. Leaf unfolding was defined as the stage where both the
entire leaf blade and the leaf stalk were visible (Figure 2–figure supplement 5).
Flowering was defined as the stage where either male or female flowers were
visible (Figure 2–figure supplement 5). We subsequently plotted the proportion of
branches exhibiting leaf unfolding and flowering. The data are provided in
Supplementary file 3.

Estimation of protein-coding sequence divergence

Pairwise distances for non-synonymous (dN) and synonymous (dS) substitutions, as well as the dN/dS ratio, were estimated for each orthologous gene pair between *L. edulis* and *Q. glauca* using codeml (PAML 4.0)(Yang, 2007). The analysis was performed with the following settings: seqtype = 1, CodonFreq = 2, Runmode = -2, and the transition/transversion ratio estimated from the data. To ensure reliability, genes exhibiting signs of saturated divergence or underestimated dS values were excluded, as codeml produces robust results only within a moderate range of sequence divergence. Specifically, we removed 43 orthologs with dN/dS $\geqq$ 99 and 483 genes with dS > 10, dS < 0.001, or dN < 0.0003. The thresholds of dN and dS were determined by examining the relationship between dN and dS and excluding outliers that deviated from the main distribution. After filtering, the final dataset comprised 11223 orthologous genes.

Linear regression analyses were performed to examine the relationship between dN, dS, dN/dS and peak month differences ($\Delta\varphi$) in gene expression between *Q. glauca* and *L. edulis*, as well as mean expression levels. These analyses were conducted using the lm function in R software (v 4.3.1). Specifically, we modeled the evolutionary rates (dN, dS, and dN/dS) as the response variable and the peak month difference ($\Delta\varphi$) or the mean levels in gene expression as the predictor variable. To account for the skewed distributions of evolutionary rates, logarithm transformations with base 2 were applied.

Data deposition and code availability

All sequencing data generated in this study have been deposited in the DNA Data Bank of Japan (DDBJ) and are available under the BioProject accession number PRJDB20498. RNA-seq reads used for expression analyses are available under the DDBJ Sequence Read Archive (DRA) run accession numbers DRR659185-DRR659374, DRR664882-DRR665873, and DRR724385-DRR724496, while DNA sequences used for genome assembly, genome size estimation and phylogenetic reconstruction are available under DRR659899-DRR659929. The assembled genomes are also available from DDBJ: the *Q. glauca* genome under accession numbers BAAHOF010000001-BAAHOF010000797, and the *L. edulis* genome under BAAHOE010000001-BAAHOE010000750. All codes used in the analysis are available at https://github.com/ku-biomath/Evolution_of_gene_expression_in_seasonal_envrionment.

Acknowledgments

We thank Atsuko Miyawaki-Kuwakado for her valuable technical advice on RNA-seq analysis. We also appreciate the technical support provided by Kayoko Ohta. Additionally, we thank Koharu Yamaguchi, Yuta Aoyagi, and Ryosuke Imai for their help with sample collection.

Additional information

Funding

825 This work was supported by JSPS KAKENHI Grant Numbers JP23H04965 and
826 JP23H04966.
827

828 **Author Contributions**
829 S.N.K. and A.S. designed research; S.N.K., Y.I., J.K., H.H., S.I., and A.S.
830 performed research; S.N.K., Y.I., J.K., H.H., S.I., and A.S. analyzed data; S.N.K.
831 and A.S. wrote the paper with inputs from all the authors.
832
833 **Competing Interest Statement**
834 The authors declare no conflict of interest.
835

References
Aikawa S, Kobayashi MJ, Satake A, Shimizu KK, Kudoh H. 2010. Robust control
of the seasonal expression of the Arabidopsis FLC gene in a fluctuating
environment. *PNAS* **107**:11632–11637. doi:10.1073/pnas.0914293107

Bateman A, Martin M-J, Orchard S, Magrane M, Adesina A, Ahmad S, Bowler-
Barnett EH, Bye-A-Jee H, Carpentier D, Denny P, Fan J, Garmiri P, Gonzales
LJ da C, Hussein A, Ignatchenko A, Insana G, Ishtiaq R, Joshi V, Jyothi D,
Kandasaamy S, Lock A, Luciani A, Luo J, Lussi Y, Marin JSM, Raposo P, Rice
DL, Santos R, Speretta E, Stephenson J, Totoo P, Tyagi N, Urakova N,
Vasudev P, Warner K, Wijerathne S, Yu CW-H, Zaru R, Bridge AJ, Aimo L,
Argoud-Puy G, Auchincloss AH, Axelsen KB, Bansal P, Baratin D, Batista
Neto TM, Blatter M-C, Bolleman JT, Boutet E, Breuza L, Gil BC, Casals-Casas
C, Echioukh KC, Coudert E, Cuche B, de Castro E, Estreicher A, Famiglietti
ML, Feuermann M, Gasteiger E, Gaudet P, Gehant S, Gerritsen V, Gos A,
Gruaz N, Hulo C, Hyka-Nouspikel N, Jungo F, Kerhornou A, Mercier P Le,
Lieberherr D, Masson P, Morgat A, Paesano S, Pedruzzi I, Pilbout S, Pourcel
L, Poux S, Pozzato M, Pruess M, Redaschi N, Rivoire C, Sigrist CJA,
Sonesson K, Sundaram S, Sveshnikova A, Wu CH, Arighi CN, Chen C, Chen
Y, Huang H, Laiho K, Lehvaslaiho M, McGarvey P, Natale DA, Ross K,
Vinayaka CR, Wang Y, Zhang J. 2025. UniProt: the Universal Protein
Knowledgebase in 2025. *Nucleic Acids Research* **53**:D609–D617.
doi:10.1093/nar/gkae1010

Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, Weier M,
Liechti A, Aximu-Petri A, Kircher M, Albert FW, Zeller U, Khaitovich P,
Grützner F, Bergmann S, Nielsen R, Pääbo S, Kaessmann H. 2011. The
evolution of gene expression levels in mammalian organs. *Nature* **478**:343–
348. doi:10.1038/nature10532

Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. 2021. BRAKER2:
Automatic eukaryotic genome annotation with GeneMark-EP+ and
AUGUSTUS supported by a protein database. *NAR Genomics and*
Bioinformatics **3**:1–11. doi:10.1093/nargab/lqaa108

Buchfink B, Xie C, Huson DH. 2014. Fast and sensitive protein alignment using
DIAMOND. *Nature Methods*. doi:10.1038/nmeth.3176

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden
TL. 2009. BLAST+: Architecture and applications. *BMC Bioinformatics* **10**.
doi:10.1186/1471-2105-10-421

Cantalapiedra CP, Herṇandez-Plaza A, Letunic I, Bork P, Huerta-Cepas J. 2021.
eggNOG-mapper v2: functional annotation, orthology assignments, and
domain prediction at the metagenomic scale. *Molecular Biology and Evolution*
38:5825–5829. doi:10.1093/molbev/msab293

Cardoso-Moreira M, Halbert J, Valloton D, Velten B, Chen C, Shao Y, Liechti A,
Ascenção K, Rummel C, Ovchinnikova S, Mazin P V., Xenarios I, Harshman
K, Mort M, Cooper DN, Sandi C, Soares MJ, Ferreira PG, Afonso S, Carneiro
M, Turner JMA, VandeBerg JL, Fallahshahroudi A, Jensen P, Behr R, Lisgo
S, Lindsay S, Khaitovich P, Huber W, Baker J, Anders S, Zhang YE,
Kaessmann H. 2019. Gene expression across mammalian organ
development. *Nature* **571**:505–509. doi:10.1038/s41586-019-1338-5

Chen LT, Wu K. 2010. Role of histone deacetylases HDA6 and HDA19 in ABA and
abiotic stress response. *Plant Signaling & Behavior* **5**:1318–1320.
doi:10.4161/psb.5.10.13168

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics* **34**:i884–i890. doi:10.1093/bioinformatics/bty560

Chen SC, Cannon CH, Kua CS, Liu JJ, Galbraith DW. 2014. Genome size variation
in the Fagaceae and its implications for trees. *Tree Genetics and Genomes*
10:977–988. doi:10.1007/s11295-014-0736-y

Chikhi R, Medvedev P. 2014. Informed and automated k-mer size selection for
genome assembly. *Bioinformatics* **30**:31–37.
doi:10.1093/bioinformatics/btt310

Cleland EE, Wolkovich EM. 2024. Effects of phenology on plant community
assembly and structure. *Annual Review of Ecology, Evolution, and*
Systematics **55**:471–492. doi:10.1146/annurev-ecolsys-102722-011653

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson
M, Gingeras TR. 2013. STAR: Ultrafast universal RNA-seq aligner.
Bioinformatics **29**:15–21. doi:10.1093/bioinformatics/bts635

Drost HG, Bellstädt J, Ó'Maoiléidigh DS, Silva AT, Gabel A, Weinholdt C, Ryan
PT, Dekkers BJW, Bentsink L, Hilhorst HWM, Ligterink W, Wellmer F, Grosse
I, Quint M. 2016. Post-embryonic hourglass patterns mark ontogenetic
transitions in plant development. *Mol Biol Evol* **33**:1158–1163.
doi:10.1093/molbev/msw039

Drummond DA, Bloom JD, Adami C, Wilke CO, Arnold FH. 2005. Why highly expressed proteins evolve slowly. *PNAS* **102**:14338–14343. doi:10.1073/pnas.0504070102

El Taher A, Böhne A, Boileau N, Ronco F, Indermaur A, Widmer L, Salzburger W. 2021. Gene expression dynamics during rapid organismal diversification in African cichlid fishes. *Nature Ecology and Evolution* **5**:243–250. doi:10.1038/s41559-020-01354-3

Emanuelle S, Doblin MS, Stapleton DI, Bacic A, Gooley PR. 2016. Molecular insights into the enigmatic metabolic regulator, SnRK1. *Trends in Plant Science* **21**:341–353. doi:10.1016/j.tplants.2015.11.001

Emms DM, Kelly S. 2019. OrthoFinder: Phylogenetic orthology inference for comparative genomics. *Genome Biology* **20**. doi:10.1186/s13059-019-1832-y

Falcon S, Gentleman R. 2007. Using GOstats to test gene lists for GO term association. *Bioinformatics* **23**:257–258. doi:10.1093/bioinformatics/btl567

Fan H, Ives AR, Surget-Groba Y, Cannon CH. 2015. An assembly and alignment-free method of phylogeny reconstruction from next-generation sequencing data. *BMC Genomics* **16**. doi:10.1186/s12864-015-1647-5

Fischl H, McManus D, Oldenkamp R, Schermelleh L, Mellor J, Jagannath A, Furger A. 2020. Cold-induced chromatin compaction and nuclear retention of clock mRNAs resets the circadian rhythm. *EMBO Journal* **39**. doi:10.15252/embj.2020105604

Forrest J, Miller-Rushing AJ. 2010. Toward a synthetic understanding of the role of phenology in ecology and evolution. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**:3101–3112. doi:10.1098/rstb.2010.0145

Fretwell SD. 1972. Populations in a seasonal environment. *Monographs in Population Biology* **5**:1–217.

Gaut B, Yang L, Takuno S, Eguiarte LE. 2011. The patterns and causes of variation in plant nucleotide substitution rates. *Annual Review of Ecology, Evolution, and Systematics* **42**:245–266. doi:10.1146/annurev-ecolsys-102710-145119

Hernández-Plaza A, Szklarczyk D, Botas J, Cantalapiedra CP, Giner-Lamia J, Mende DR, Kirsch R, Rattei T, Letunic I, Jensen LJ, Bork P, von Mering C, Huerta-Cepas J. 2023. eggNOG 6.0: enabling comparative genomics across 12 535 organisms. *Nucleic Acids Research* **51**:D389–D394. doi:10.1093/nar/gkac1022

Hipp AL, Manos PS, Hahn M, Avishai M, Bodénès C, Cavender-Bares J, Crowl AA, Deng M, Denk T, Fitz-Gibbon S, Gailing O, González-Elizondo MS, González-

Rodríguez A, Grimm GW, Jiang XL, Kremer A, Lesur I, McVay JD, Plomion C, Rodríguez-Correa H, Schulze ED, Simeone MC, Sork VL, Valencia-Avalos S. 2020. Genomic landscape of the global oak phylogeny. *New Phytologist* **226**:1198–1212. doi:10.1111/nph.16162

Ikezaki Y, Kudo SN, Nakata T, Koita S, Munakata R, Yazaki K, Torimaru T, Tomaru N, Isobe S, Hirakawa H, Kusumi J, Satake A. 2025. Molecular evolution of terpene synthase underlying the diversification of isoprene emission in Fagaceae. doi:10.1101/2025.07.31.667835

Irie N, Kuratani S. 2011. Comparative transcriptome analysis reveals vertebrate phylotypic period during organogenesis. *Nature Communications* **2**. doi:10.1038/ncomms1248

Kalinka AT, Varga KM, Gerrard DT, Preibisch S, Corcoran DL, Jarrells J, Ohler U, Bergman CM, Tomancak P. 2010. Gene expression divergence recapitulates the developmental hourglass model. *Nature* **468**:811–816. doi:10.1038/nature09634

Kant P, Kant S, Gordon M, Shaked R, Barak S. 2007. Stress Response Suppressor1 and Stress Response Suppressor2, two Dead-box RNA helicases that attenuate Arabidopsis responses to multiple abiotic stresses. *Plant Physiology* **145**:814–830. doi:10.1104/pp.107.099895

Keilwagen J, Hartung F, Grau J. 2019. GeMoMa: Homology-based gene prediction utilizing intron position conservation and RNA-seq data *Methods in Molecular Biology*. Humana Press Inc. pp. 161–177. doi:10.1007/978-1-4939-9173-0_9

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* **37**:907–915. doi:10.1038/s41587-019-0201-4

Kokot M, Dlugosz M, Deorowicz S. 2017. KMC 3: counting and manipulating k-mer statistics. *Bioinformatics* **33**:2759–2761. doi:10.1093/bioinformatics/btx304

Komoto H, Nagahama A, Miyawaki-Kuwakado A, Hata Y, Kyozuka J, Kajita Y, Toyama H, Satake A. 2024. The transcriptional changes underlying the flowering phenology shift of *Arabidopsis halleri* in response to climate warming. *Plant Cell & Environment* **47**:174–186. doi:10.1111/pce.14716

Kremer A, Abbott AG, Carlson JE, Manos PS, Plomion C, Sisco P, Staton ME, Ueno S, Vendramin GG. 2012. Genomics of Fagaceae. *Tree Genetics & Genomes* **8**:583–610. doi:10.1007/s11295-012-0498-3

Krishnakumar V, Hanlon MR, Contrino S, Ferlanti ES, Karamycheva S, Kim M, Rosen BD, Cheng CY, Moreira W, Mock SA, Stubbs J, Sullivan JM, Krampis K, Miller JR, Micklem G, Vaughn M, Town CD. 2015. Araport: The Arabidopsis

Information Portal. *Nucleic Acids Research* **43**:D1003–D1009.
doi:10.1093/nar/gku1200

Kubo T, Iwasa Y. 1996. Phenological pattern of tree regeneration in a model for
forest species diversity. *Theoretical Population Biology* **49**:90–117.
doi:10.1006/tpbi.1996.0004

Kudoh H. 2016. Molecular phenology in plants: *in natura* systems biology for the
comprehensive understanding of seasonal responses under natural
environments. *New Phytologist* **210**:399–412. doi:10.1111/nph.13733

Lemos B, Bettencourt BR, Meiklejohn CD, Hartl DL. 2005. Evolution of proteins
and gene expression levels are coupled in *Drosophila* and are independently
associated with mRNA abundance, protein length, and number of protein-
protein interactions. *Molecular Biology and Evolution* **22**:1345–1354.
doi:10.1093/molbev/msi122

Levin M, Anavy L, Cole AG, Winter E, Mostov N, Khair S, Senderovich N, Kovalev
E, Silver DH, Feder M, Fernandez-Valverde SL, Nakanishi N, Simmons D,
Simakov O, Larsson T, Liu SY, Jerafi-Vider A, Yaniv K, Ryan JF, Martindale
MQ, Rink JC, Arendt D, Degnan SM, Degnan BM, Hashimshony T, Yanai I.
2016. The mid-developmental transition and the evolution of animal body
plans. *Nature* **531**:637–641. doi:10.1038/nature16994

Li B, Dewey CN. 2011. RSEM: Accurate transcript quantification from RNA-Seq
data with or without a reference genome. *BMC Bioinformatics* **12**.
doi:10.1186/1471-2105-12-323

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-
Wheeler transform. *Bioinformatics* **25**:1754–1760.
doi:10.1093/bioinformatics/btp324

Liao BY, Zhang J. 2006. Low rates of expression profile divergence in highly
expressed genes and tissue-specific genes during mammalian evolution.
Molecular Biology and Evolution **23**:1119–1128. doi:10.1093/molbev/msj119

Liu H, Zhang R, Zhou BF, Shen Z, Chen XY, Gao J, Wang B. 2023. Chromosome-
scale genome assembly of sweet tea (Lithocarpus polystachyus Rehder).
Scientific Data **10**. doi:10.1038/s41597-023-02791-y

Lotharukpong JS, Zheng M, Luthringer R, Liesner D, Drost H-G, Coelho SM. 2024.
A transcriptomic hourglass in brown algae. *Nature* **635**:129–135.
doi:10.1038/s41586-024-08059-8

Manos PS, Zhou Z, Cannon CH. 2001. Systematics of Fagaceae: phylogenetic
tests of reproductive trait evolution. *International Journal of Plant Sciences*
162:1361–1379. doi:10.1086/322949

Mishra B, Ulaszewski B, Meger J, Aury JM, Bodénès C, Lesur-Kupin I, Pfenninger
M, Da Silva C, Gupta DK, Guichoux E, Heer K, Lalanne C, Labadie K,
Opgenoorth L, Ploch S, Le Provost G, Salse J, Scotti I, Wötzel S, Plomion C,
Burczyk J, Thines M. 2022. A chromosome-level genome assembly of the
European Beech (Fagus sylvatica) reveals anomalies for organelle DNA
integration, repeat content and distribution of SNPs. *Frontiers in Genetics* **12**.
doi:10.3389/fgene.2021.691058

Miyawaki‐Kuwakado A, Han Q, Kitamura K, Satake A. 2024. Impacts of climate
change on the transcriptional dynamics and timing of bud dormancy release
in Yoshino‐cherry tree. *PLANTS, PEOPLE, PLANET* **6**:1505–1521.
doi:10.1002/ppp3.10548

Miyazaki Y, Maruyama Y, Chiba Y, Kobayashi MJ, Joseph B, Shimizu KK, Mochida
K, Hiura T, Kon H, Satake A. 2014. Nitrogen as a key regulator of flowering in
Fagus crenata: Understanding the physiological mechanism of masting by
gene expression analysis. *Ecology Letters* **17**:1299–1309.
doi:10.1111/ele.12338

Morellato LPC, Camargo MGG, Gressler E. 2013. "A review of plant phenology in
South and Central America." in Phenology: An Integrative Environmental
Science. Springer.

Nagano AJ, Kawagoe T, Sugisaka J, Honjo MN, Iwayama K, Kudoh H. 2019.
Annual transcriptome dynamics in natural environments reveals plant
seasonal adaptation. *Nature Plants* **5**:74–83. doi:10.1038/s41477-018-0338-
z

Naqvi S, Godfrey AK, Hughes JF, Goodheart ML, Mitchell RN, Page DC. 2019.
Conservation, acquisition, and functional impact of sex-biased gene
expression in mammals. *Science (1979)* **365**. doi:10.1126/science.aaw7317

Nishio H, Nagano AJ, Ito T, Suzuki Y, Kudoh H. 2020. Seasonal plasticity and diel
stability of H3K27me3 in natural fluctuating environments. *Nature Plants*
6:1091–1097. doi:10.1038/s41477-020-00757-1

Nuzhdin S V., Wayne ML, Harmon KL, McIntyre LM. 2004. Common pattern of
evolution of gene expression level and protein sequence in Drosophila.
Molecular Biology and Evolution **21**:1308–1317. doi:10.1093/molbev/msh128

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast
and bias-aware quantification of transcript expression. *Nature Methods*
14:417–419. doi:10.1038/nmeth.4197

Pertea M, Pertea G. 2020. GFF Utilities: GffRead and GffCompare. *F1000Res* **9**.
doi:10.12688/f1000research.23297.1

Piao S, Liu Q, Chen A, Janssens IA, Fu Y, Dai J, Liu L, Lian X, Shen M, Zhu X.
2019. Plant phenology and global climate change: Current progresses and
challenges. *Global Change Biology* **25**:1922–1940. doi:10.1111/gcb.14619

Prud'homme B, Gompel N, Carroll SB. 2007. Emerging principles of regulatory
evolution. *PNAS* **104**:8605–8612. doi:10.1073/pnas.0700488104

Putnam NH, O'Connell BL, Stites JC, Rice BJ, Blanchette M, Calef R, Troll CJ,
Fields A, Hartley PD, Sugnet CW, Haussler D, Rokhsar DS, Green RE. 2016.
Chromosome-scale shotgun assembly using an in vitro method for long-range
linkage. *Genome Research* **26**:342–350. doi:10.1101/gr.193474.115

Quint M, Drost HG, Gabel A, Ullrich KK, Bönn M, Grosse I. 2012. A transcriptomic
hourglass in plant embryogenesis. *Nature* **490**:98–101.
doi:10.1038/nature11394

Ranallo-Benavidez TR, Jaron KS, Schatz MC. 2020. GenomeScope 2.0 and
Smudgeplot for reference-free profiling of polyploid genomes. *Nature*
Communications **11**. doi:10.1038/s41467-020-14998-3

Satake A, Hagiwara T, Nagano AJ, Yamaguchi N, Sekimoto K, Shiojiri K, Sudo K.
2024a. Plant molecular phenology and climate feedbacks mediated by
BVOCs. *Annual Review of Plant Biology* **75**:605–627. doi:10.1146/annurev-
arplant-060223-032108

Satake A, Imai R, Fujino T, Tomimoto S, Ohta K, Na'iem M, Indrioko S, Widiyatno
W, Purnomo S, Morales AM, Nizhynska V, Tani N, Suyama Y, Sasaki E,
Kasahara M. 2024b. Somatic mutation rates scale with time not growth rate in
long-lived tropical trees. *Elife* **12**. doi:10.7554/eLife.88456

Satake A, Nagahama A, Sasaki E. 2022. A cross‐scale approach to unravel the
molecular basis of plant phenology in temperate and tropical climates. *New*
Phytologist **233**:2340–2353. doi:10.1111/nph.17897

Satake A, Ohta K, Takeda-Kamiya N, Toyooka K, Kusumi J. 2023. Seasonal gene
expression signatures of delayed fertilization in Fagaceae. *Molecular Ecology*
32:4801–4813. doi:10.1111/mec.17079

Schwartz MD. 2013. Phenology: An Integrative Environmental Science, 2nd ed.
Springer Dordrecht.

Smaczniak C, Immink RGH, Muiño JM, Blanvillain R, Busscher M, Busscher-
Lange J, Dinh QD, Liu S, Westphal AH, Boeren S, Parcy F, Xu L, Carles CC,
Angenent GC, Kaufmann K. 2012. Characterization of MADS-domain
transcription factor complexes in *Arabidopsis* flower development. *PNAS*
109:1560–1565. doi:10.1073/pnas.1112871109

Smid M, Coebergh van den Braak RRJ, van de Werken HJG, van Riet J, van Galen
A, de Weerd V, van der Vlugt-Daane M, Bril SI, Lalmahomed ZS, Kloosterman
WP, Wilting SM, Foekens JA, IJzermans JNM, Martens JWM, Sieuwerts AM.
2018. Gene length corrected trimmed mean of M-values (GeTMM) processing
of RNA-seq data performs similarly in intersample analyses while improving
intrasample comparisons. *BMC Bioinformatics* **19**. doi:10.1186/s12859-018-
2246-7

Thaben PF, Westermark PO. 2014. Detecting rhythms in time series with rain.
Journal of Biological Rhythms **29**:391–400. doi:10.1177/0748730414553029

True JR, Haag ES. 2001. Developmental system drift and flexibility in evolutionary
trajectories. *Evolution & Development* **3**:109–119. doi:10.1046/j.1525-
142x.2001.003002109.x

Wang Y, Tang H, Debarry JD, Tan X, Li J, Wang X, Lee TH, Jin H, Marler B, Guo
H, Kissinger JC, Paterson AH. 2012. MCScanX: A toolkit for detection and
evolutionary analysis of gene synteny and collinearity. *Nucleic Acids
Research* **40**. doi:10.1093/nar/gkr1293

Yang L, Gaut BS. 2011. Factors that contribute to variation in evolutionary rate
among arabidopsis genes. *Molecular Biology and Evolution* **28**:2359–2369.
doi:10.1093/molbev/msr058

Yang Z. 2007. PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular
Biology and Evolution* **24**:1586–1591. doi:10.1093/molbev/msm088

Yoo SJ, Chung KS, Jung SH, Yoo SY, Lee JS, Ahn JH. 2010. BROTHER of FT
and TFL1 (BFT) has TFL1-like activity and functions redundantly with TFL1 in
inflorescence meristem development in *Arabidopsis*. *Plant Journal* **63**:241–
253. doi:10.1111/j.1365-313X.2010.04234.x

Zhang X, Yu G, Dai Y, Zhang H, Wang K, Han J. 2023. High-resolution Hi-C maps
highlight multiscale chromatin architecture reorganization during cold stress in
Brachypodium distachyon. *BMC Plant Biology* **23**. doi:10.1186/s12870-023-
04269-w

Zhou BF, Yuan S, Crowl AA, Liang YY, Shi Y, Chen XY, An QQ, Kang M, Manos
PS, Wang B. 2022. Phylogenomic analyses highlight innovation and
introgression in the continental radiations of Fagaceae across the Northern
Hemisphere. *Nature Communications* **13**. doi:10.1038/s41467-022-28917-1

Zhu GQ, Qu L, Xue HW. 2024. Casein kinase 1 AELs promote senescence by
enhancing ethylene biosynthesis through phosphorylating WRKY22
transcription factor. *New Phytologist* **244**:116–130. doi:10.1111/nph.19785

Zoldos V, Papes D, Cerbah M, Panaud O, Besendorfer V, Siljak-Yakovlev S. 1999.
Molecular-cytogenetic studies of ribosomal genes and heterochromatin reveal

conserved genome organization among 11 Quercus species. *Theoretical and*
Applied Genetics **99**:969–977. doi:10.1007/s001220051404

Zrimec J, Börlin CS, Buric F, Muhammad AS, Chen R, Siewers V, Verendel V,
Nielsen J, Töpel M, Zelezniak A. 2020. Deep learning suggests that gene
expression is encoded in all parts of a co-evolving interacting gene regulatory
structure. *Nature Communications* **11**. doi:10.1038/s41467-020-19921-4

Zrimec J, Buric F, Kokina M, Garcia V, Zelezniak A. 2021. Learning the Regulatory
Code of Gene Expression. *Frontiers in Molecular Biosciences* **8**.
doi:10.3389/fmolb.2021.673363

1136
1137

Figures
Figure 1. Synteny and distribution of genomic features of *Q. glauca* and *L. edulis* genomes. (A) A phylogenetic tree of five Fagaceae species from three genera, *Quercus*, *Lithocarpus*, and *Fagus*. Bootstrap values for all branches are 100. (B) Synteny relationship between the *Q. glauca* and *L. edulis* genomes. The collinear blocks within the genomes were displayed by lines. Black lines show reverse alignment and gray lines represent regular. Numbers correspond to the 12 chromosomes. (C–D) Circos plots of the *Q. glauca* (C) and *L. edulis* genomes (D). Lanes depict circular representation of chromosome length (Mbp) (a), gene density (b), GC density (c), and repeat (d). Lines in the inner circle represent links between synteny-selected paralogs. (E) Orthogroup classification based on the orthogroup size difference between the *Q. glauca* and *L. edulis* genomes. The bar plot illustrates the number of genes in gene families categorized into 1:1 orthologs, N:N (many-to-many), 1:N (one-to-many), N:1 (many-to-one), and species-specific unique genes. (F–G) Mean gene expression levels in each copy number variation

category for *Q. glauca* (F) and *L. edulis* (G). Different letters denote statistically significant differences (*p* < 0.05, Steel-Dwass test).

Figure 2. Seasonal gene expression dynamics. (A) Hierarchical clustering of 213 time points encompassing two tissues (leaf and bud) and four species (*Q. glauca*, *Q. acuta*, *L. edulis*, and *L. glaber*), collected over two-year. Newly flushed leaf samples are indicated as asterisks. (B) Distribution of daily mean temperature on each sampling date across clusters, from L1 (top) to B3 (bottom). The Mann-Whitney U test was used to compare daily mean temperature between cluster L1 and L2 (*p* = 0.136). The Steel-Dwass test was applied for multiple comparisons across clusters B1–B3 (*p* < 0.05). Different letters denote statistically significant differences (*p* < 0.05). (C–D) Hierarchical clustering of 11749 genes based on the similarity of gene expression dynamics. Daily mean temperature, flowering and

leaf flushing phenology (proportion) are shown in the upper three panels of the heatmaps.

Figure 3. Distribution of period and peak month in rhythmic gene expression. (A–B) Distribution of period length (month) among the genes with significant rhythmicity calculated by RAIN for leaf (A) and bud (B) samples of each species. (C–D) Distribution of seasonal peaks among the genes with annual periodicity. The angle of each bar represents the peak month, while the bar height indicates the number of genes. Blue arrows indicate the month with the highest number of genes peaking during winter (December–February), while red arrows indicate the month with the highest number of genes peaking during the growing season (March–November). Purple and green squares around each rose plot denote the observed phenology of flowering and leaf flushing (proportion), respectively.

Figure 4. Comparison of peak months in seasonal gene expression across
species in buds. (A) Intra-genus and (B) inter-genera comparisons of peak months
in seasonal gene expression. The size and color of the circles represent the
number of genes and the mean of Pearson's correlation, respectively. Genes with
negative correlations lower than -0.3 are highlighted in blue, while the month of
leaf flushing is highlighted in green. (C) Plot of molecular phenology divergence
index D across months. Triangles and circles indicate the values of D calculated
from intra-genus and inter-genera comparison, respectively. The line and shaded
envelope indicate the mean and standard deviation (s.d.). Different letters denote
statistically significant differences (Nemenyi test, $p < 0.05$).

1191
1192

Figure 5. Phylogenetic constraints in the evolution of seasonal gene expression
and the relationship between seasonal gene expression divergence and sequence
evolution. (A) Distribution of pairwise Pearson's correlation coefficients for gene
expression across species (Le: *L. edulis*, Lg: *L. glabe*r, Qg: *Q. glauca*, Qa: *Q.*
acuta; n = 11749). Colors indicate rhythmicity categories. Black bars show median
correlation values, and different letters indicate significant differences (Nemenyi

test, *p* < 0.05). (B) Heatmap of pairwise correlation coefficients for each gene, with mean gene counts in four seasonal expression categories: (1) annual rhythmic genes peaking in the growing season (March–November), (2) annual rhythmic genes peaking in winter (December–February), (3) half-annual rhythmic genes, and (4) arrhythmic genes. Genes are ordered by mean correlation across species pairs. (C–D) Conserved (C) and diverged (D) gene expression patterns, based on the top and bottom 5% of pairwise correlations. Solid lines and error bars indicate mean ± SD. Numbers in parentheses show peak month differences between *Q. glauca* and *L. edulis*. Peak month difference for *BFT* was not calculated due to arrhythmic expression in *Q. glauca*. (E) Relationship between evolutionary rate (dN/dS) and peak month difference (Δφ) for genes with annual periodicity in *Q. glauca* and *L. edulis* (*n* = 951). Black lines and transparent bands represent regression lines and 95% confidence intervals. The regression coefficient (β) and corresponding *p*-value are shown in each panel.

1213
1214
1215

 Figure supplements

Figure 2–figure supplement 1. Dynamic range of gene expression in seasonal
environments. (A) Dynamic range of seasonal gene expression in leaves and buds
for *Q. glauca*, *Q. acuta*, *L. edulis*, and *L. glaber*. Each dot represents an expression
level (log$_2$(GeTMM+1)) on each sampling date, while black lines indicate the mean
expression level of each gene over the entire sampling period. The bars on the left
side of the panels distinguish between rhythmic (blue) and arhythmic (gray) genes.
(B) Density plots showing the distribution of the ratio of standard deviation divided
by the mean for each gene (σ_i / μ_i) and the overall variation across all genes in
the genome (σ_g/μ_g) in leaves and buds for *Q. glauca*, *Q. acuta*, *L. edulis*, and *L.*
glaber.

1227
1228

1229 **Figure 2–figure supplement 2.** Comparison of gene expression across
1230 individuals. Scatter plots showing the expression levels of each gene measured
1231 from different individuals for *L. glaber* (A), *Q. glauca* (B), and *Q. acuta* (C). The
1232 Pearson's correlation coefficient for each pair of individuals is shown in the
1233 corresponding panel.

1234
1235

Figure 2–figure supplement 3. Seasonal fluctuations in environmental factors at
the three study sites. Daily mean temperature and photoperiods at the three study
sites (Ito campus, Kyushu university; Imajuku Field Activity Center; Mt. Sefuri in
Kyushu, Japan) during the monitoring periods. The solid lines indicate the daily
mean temperature, and the shaded area indicates the photoperiod. The study sites
are located within a relatively small region, with distances between them ranging
from approximately 10 to 20 km. The elevations vary, with Ito Campus (20–57 m
a.s.l.) and Imajuku Field Activity Center (84–111 m a.s.l.) at lower altitudes, while
Mt. Sefuri is at a higher elevation (970–974 m a.s.l.). Environmental data from Ito
Campus and the Imajuku Field Activity Center were obtained from the Japan
Meteorological Agency, while data from Sefuri Mountain were collected using a
logger installed at the sampling site since July 15, 2021.

1248
1249

Figure 2–figure supplement 4. Determining the optimal number of clusters in
hierarchical clustering. (A) Relationship between the number of clusters k and total
Within-Cluster-Sum of Squared Error (WSS) on the clustering of 213 samples. We
adopted the optimal $k = 5$ because it was the elbow point of the curve. (B)
Relationship between the number of clusters k and total WSS on the clustering of
11749 genes. The elbow point was $k = 4$ or 5 and we selected the number of cluster
$k = 5$. If $k = 4$ is selected, gene set IV and V (Figure 2D, E in main text) are merged.

1257
1258

1259 **Figure 2–figure supplement 5.** Phenological observations of leaf flushing,
1260 flowering (male and female flower) and winter buds of the four target species.

1261
1262

Figure 2–figure supplement 6. Seasonal gene expression dynamics observed
within the same sampling period (from May 2021 to October 2022). (A) Hierarchical
clustering of 160 time points encompassing two tissues (leaf and bud) and four
species (*Q. glauca*, *Q. acuta*, *L. edulis*, and *L. glaber*). Newly flushed leaf samples
are indicated as asterisks. (C–D) Hierarchical clustering of 11749 genes based on
the similarity of gene expression dynamics. Daily mean temperature is shown in
the top of the heatmaps.

1270
1271

Figure 2–figure supplement 7. Seasonal gene expression dynamics under the identical environmental condition. (A) Hierarchical clustering of 161 time points from two tissues (leaf and bud) across three species (*Q. glauca*, *L. edulis*, and *L. glaber*) excluding *Q. acuta* sampled at higher elevation. Newly flushed leaf samples are indicated as asterisks. (C–D) Hierarchical clustering of 11749 genes based on the similarity of their gene expression dynamics. Daily mean temperature is shown in in the top of the heatmaps.

Figure 2–figure supplement 8. Distribution of environmental factors across
clusters. Distribution of mean temperature for the 7 days prior to the sampling (A)
and photoperiod for each sampling date (B) across clusters identified through
hierarchical clustering in Figure 2A. The Mann-Whitney U test was used to
compare climatic factors between leaf clusters (L1 and L2), while the Steel-Dwass
test was applied for multiple comparisons among bud clusters (B1, B2, and B3).
Newly flushed leaves were included within cluster B1. Different letters denote
statistically significant differences ($p < 0.05$).

1289
1290

Figure 2–figure supplement 9. 3D plot of PC1–PC3 resulting from the PCA of
11749 orthologous genes in four Fagaceae species. Leaf samples are represented
in a color gradient from green to blue, while bud samples range from orange to
yellow. The color gradients indicate the mean temperature on the respective
sampling dates for bud and leaf samples. The percentages in parentheses on each
axis denote the variance explained by each principal component. The numbers
within the plot indicate sampling months.

1298
1299

Figure 3–figure supplement 1. Number of rhythmic genes across species and distribution of peak months across half-annual genes. (A) Number of genes classified into annual (period range: 8-16 months), half-annual (< 8 month), and arrhythmic periodicity. (B–C) Number of common rhythmic genes shared among species within each species combination, categorized by annual (B) and half-annual (C) periodicity. Black dots beneath each bar indicate the species included in each combination. (D) Distribution of seasonal peaks across genes with half-annual periodicity. (E) Seasonal expression patterns of *WRKY22* in leaves and *KINγ1* in buds. For each gene, the species exhibiting a half-annual periodicity is highlighted with a thick, dark line, while the expression patterns in the other species are shown with thin, light lines.

Figure 4–figure supplement 1. Comparison of peak months in seasonal gene
expression across species in leaves. (A) Intra-genus and (B) inter-genera
comparison of peak months in seasonal gene expression. (C) Comparison of peak
months between leaves and buds. The size and color of the circles represent the
number of genes and the mean of Pearson's correlation, respectively. Genes with
negative correlations lower than -0.3 are highlighted in blue, while the month of
leaf flushing is highlighted in green. (D) Plot of molecular phenology divergence
index D across months. Triangles and circles indicate the D values calculated from
intra- and inter-genera comparison, respectively. The line and shaded envelope
indicate the mean and standard deviation (s.d.). Different letters denote statistically
significant differences (Nemenyi test, $p < 0.05$).

1324

Figure 4–figure supplement 2. Seasonal difference of molecular phenology
divergence index *D* calculated for species under the identical environmental
condition. Plot of molecular phenology divergence index *D* across months in buds
(A) and leaves (B) calculated from the gene expression in *Q. glauca*, *L. edulis*, and
L. glaber. Triangles and circles indicate the values of *D* calculated from intra-genus
and inter-genera comparison, respectively. The line and shaded envelope indicate
the mean and standard deviation (s.d.). Different letters denote statistically
significant differences (Nemenyi test, $p < 0.05$).

A Bud

B Leaf

1333
1334

 Figure 5–figure supplement 1. Phylogenetic constraints in the evolution of
seasonal gene expression in leaves. (A) Distribution of pairwise Pearson's
correlation coefficients for gene expression across species (Le: *L. edulis*, Lg: *L.*
glaber, Qg: *Q. glauca*, Qa: *Q. acuta*; *n* = 11749). Colors indicate three categories
based on the presence or absence of rhythmicity in gene expression. The median
values of Pearson's correlation across all the genes were depicted as black bars.
Different letters denote significantly different values (Nemenyi test, *p* < 0.05). (B)
Heatmap of pairwise Pearson's correlation coefficients for each gene, along with
the mean count of genes in four categories of seasonal gene expression: (1)
rhythmic genes with annual periodicity peaking in the growing season (March–
November), (2) rhythmic genes with annual periodicity peaking in winter
(December–February), (3) rhythmic genes with half-annual periodicity, and (4)
arrhythmic genes. The mean count was calculated for each block of 100 genes.
Genes were ordered based on the mean Pearson's correlation of their expression
across all species pairs.

1350
1351

Figure 5–figure supplement 2. Distribution of pairwise Pearson's correlation
coefficients for gene expression across species under the identical environmental
condition. Density plot of pairwise Pearson's correlation coefficients in buds (A)
and leaves (B), calculated from the gene expression in *Q. glauca*, *L. edulis*, and *L.*
glaber inhabiting nearly identical environments. Colors indicate rhythmicity
categories. Black bars show median correlation values, and different letters
indicate significant differences (Nemenyi test, *p* < 0.05).

1359
1360

Figure 5–figure supplement 3. Pairwise mean absolute difference of gene
expression dynamics. (A–B) Distribution of mean absolute difference of gene
expression dynamics (Log$_2$(GeTMM+1)) in leaf (A) and bud samples (B) across
species (Le: *L. edulis*, Lg: *L. glaber*, Qg: *Q. glauca*, Qa: *Q. acuta*; n = 11749).
Colors indicate three categories based on the presence or absence of rhythmicity
in gene expression. The median values across all the genes were depicted as
black bars. Different letters denote significantly different values (Nemenyi test, *p* <
0.05). (C–D) Heatmap of the mean absolute difference for each gene, along with
the mean count of genes in four categories of seasonal gene expression: (1)
rhythmic genes with annual periodicity peaking in the growing season (March–
November), (2) rhythmic genes with annual periodicity peaking in winter
(December–February), (3) rhythmic genes with half-annual periodicity, and (4)
arrhythmic genes. The mean count was calculated for each block of 100 genes.
Genes were ordered based on the mean absolute difference of their expression
across all the species pairs.

1376

Figure 5–figure supplement 4. Gene Ontology (GO) enrichment analysis. (A)
Top 5 GO terms (ranked by the raw *p*-values) for the genes with conserved
expression patterns across species (a mean correlation in gene expression
ranking in the top 5%) in leaves and buds. (B) Top 5 GO terms for the genes with
diverged expression patterns across species (a mean correlation in gene
expression ranking in the bottom 5%) in leaves and buds. Genes used for the
analysis are listed in Supplementary file 6.

1384
1385

 Figure 5–figure supplement 5. Relationship between sequence divergence and
 gene expression divergence. (A) Relationships between evolutionary measures
 (dN/dS, dN, dS), and mean expression levels of genes except for the outliers (*n* =
 11223). (B–C) Relationships between the evolutionary measures (dN, dS) and
 peak differences (Δφ) between *Q. glauca* and *L. edulis* in leaves (B) and buds (C)
 using the same gene set as in Figure 5E (n = 951). The regression slope and
 corresponding p-value are shown in each panel. Black lines and transparent bands
 represent regression lines and 95% confidence intervals. (D–E) Comparison of the
 evolutionary measures (dN/dS, dN, dS) among genes with conserved annual
 periodicity across four categories (*n* = 951). Genes were classified into four
 categories: those expressed in winter in both tissues, those expressed in winter
 only in buds, those expressed in winter only in leaves, and others. (E) shows a
 subset of (D), including only genes with conserved seasonal peaks (peak month
 difference < 2).

Figure 5–figure supplement 6. Developmental constraints and seasonal
constraints in the evolution of gene expression. (A)The hourglass model of
developmental constraints, illustrating similar gene expression during the mid-
embryonic stage (40–45). (B) Seasonal constraints identified in this study. Similar
gene expression patterns were observed during the winter season in both leaves
and buds, as well as during the summer in leaves. The vertical axis represents
time (developmental stages or seasons), while the horizontal axis represents the
divergence of gene expression patterns.

1409
1410

1411 **Supplementary files**
1412 **Supplementary file 1**
1413 Statistics for genome assemblies of two new sequenced species
1414
1415 **Supplementary file 2**
1416 Summary of clustering 213 gene expression profiles
1417
1418 **Supplementary file 3**
1419 Hierarchical clustering of 11749 gene expression dynamics
1420
1421 **Supplementary file 4**
1422 Identification of genes with rhythmic expression
1423
1424 **Supplementary file 5**
1425 Comparison of correlation and absolute difference among the species pairs
1426
1427 **Supplementary file 6**
1428 List of genes with conserved and diverged gene expression across species.
1429
1430 **Supplementary file 7**
1431 Statistical test of evolutionary rates
1432