Experimental approach to assessing the impact of the intensity of near-natural heatwaves on three marine phanerogam species in Mediterranean lagoon environment

Constance Bourdier[1*], Rutger De Wit[2], Laura Soissons[3], Elisa De Ronne[1], Vincent Ouisse[1]

[1] MARBEC, Univ. Montpellier, CNRS, Ifremer, IRD – Sète, France

[2] MARBEC, Univ. Montpellier, CNRS, Ifremer, IRD – Montpellier, France

[3] DECOD, Institut Agro, Inrae, Ifremer – Rennes, France

*Corresponding author
Correspondence: constance.bourdier@ifremer.fr

ABSTRACT

The increased intensity, frequency and duration of extreme climatic events (ECEs) in the context of climate change have been recognised as additional pressures for seagrass habitat in lagoon ecosystems. Among these ECE, this study focuses on the impact of marine heatwaves (MHWs) on seagrass species in Mediterranean coastal lagoons, *i.e.,* the seagrasses *Cymodocea nodosa, Nanozostera noltei,* and *Ruppia cirrhosa*. We experimentally assessed with a near-natural mesocosm, the effects of a 20-day heatwave at three intensities (+2, +4, +6°C above control near-natural conditions) in spring and summer on these seagrass species and quantified changes in growth performance, respiration, and gross primary production. Responses to these MHW varied by species and season. *C. nodosa* responded differently to MHWs in spring and summer, its net foliar growth rate being stimulated and maintained, respectively. *N. noltei* responded positively to moderate MHWs (+2, +4°C) in spring but showed vulnerability to the high-intensity MHWs (+6°C) in summer. *R. cirrhosa* remained resilient in spring and benefited from MHWs in summer, particularly at +4°C intensity, to stimulate growth. Hence, all three species were able to tolerate a single 20-day MHW with the species being increasingly better adapted in the following order *N. noltei, R. cirrhosa*, and *C. nodosa*. Increase of the initiation and growth of new leaves was a common phenological response observed in all three species, albeit at a lesser extent for *R. cirrhosa*. Our near-natural experimental approach reveals species- and season-dependent responses to MHWs, highlighting the complexity of seagrass resilience in lagoon ecosystems. These findings provide valuable insights to inform conservation strategies and improve the management of coastal habitats under future climate scenarios.

Keywords: Extreme events, Seagrass, coastal lagoon, mesocosm, heat stress, Growth performance, *Cymodocea nodosa, Ruppia cirrhosa, Nanozostera noltei.*

Introduction

The semi-enclosed nature of the Mediterranean Sea causes it to warm two to three times faster
than the global ocean (Diffenbaugh et al., 2007), making it one of the most affected bodies of
saltwater in the world (Jordà et al., 2012; Templado, 2014), and a well-known hot-spot region for
climate change (Giorgi, 2006). A rapid increase in the mean sea surface temperature (SST) of the
Mediterranean sea is expected in the near future (Giorgi & Lionello, 2008). By the end of the 21^e
century, surface temperatures in the Mediterranean Sea are predicted to rise by 0.8°C to 3.8°C
(IPCC, 2023). Besides these gradual changes, climate change is expected to increase the
intensity, the frequency and the duration of sudden and relatively short-term events known as
Extreme Climatic Events (ECEs) (Hobday et al., 2018; Jentsch & Beierkuhnlein, 2008). The
sudden nature of ECEs leaves ecosystems with little time to adapt, making it crucial to consider
them when understanding how ecosystems respond to climate change (Jentsch et al., 2007).
Among ECEs, this study focusses on MHWs, characterized by sea temperatures exceeding
seasonal thresholds for at least five consecutive days (Hobday et al., 2016), which can occur in all
seasons. While not representing a new phenomenon, MHWs have increased in frequency,
intensity, and duration over the past century, primarily due to anthropogenic climate change (IPCC,
2021). The Mediterranean Sea is expected to be highly sensitive to MHWs (Cramer et al., 2018;
Darmaraki et al., 2019; Diffenbaugh et al., 2007; Lejeusne et al., 2010). Notably, the growing
frequency and intensity of MHWs in Mediterranean sea have already been linked to mass mortality
events (Darmaraki et al., 2019; Garrabou et al., 2009). Mediterranean coastal zone ecosystems
are likely to be particularly sensitive to MHWs, as these events may potentially interact negatively
with anthropogenic pressures such as chemical contamination, eutrophication and shoreline
artificialization (de Wit, 2011; Lloret et al., 2008).

The Mediterranean coast is also characterized by numerous lagoons, transitional systems that
link terrestrial and marine environments. These highly productive systems play a crucial role in
regulating biogeochemical cycles and supporting biodiversity. This is related to their complex,
mosaic-like landscapes, which provide refuge for numerous species by offering a diversity of
habitats (de Wit, 2011; Iotti et al., 2023; Pérez-Ruzafa, Campillo, et al., 2019). Mediterranean
coastal lagoons are subject to frequent environmental disturbances and fluctuations due to their
shallowness and low thermal inertia (Nixon et al., 2004; Trombetta et al., 2019). Seasonal
temperature variations are also more pronounced than in the adjacent sea (de Wit, 2011). These
dynamics define them as naturally stressed environments, compelling their resident populations to
develop greater tolerance and adaptability compared to their counterparts in open sea
environments (Pérez-Ruzafa, Pérez-Ruzafa, et al., 2019).

The coastal lagoons comprise various linked habitats (Ouisse et al., 2023) that help species
and communities thrive (Iotti et al., 2023). Within these lagoon environments, seagrass meadows
play a key ecological role through their important biogeochemical and ecological functions. They
contribute to climate regulation by storing carbon in sediments (Fourqurean et al., 2012) and
support biodiversity (Colvin and Sne*LGR*ove, 2025), serving as nursery areas for many organisms
and commercial fish species (Tournois et al., 2017). Four species of marine phanerogams are
found in Mediterranean coastal lagoons. *Zostera marina* Linnaeus, 1753 is widely distributed in
the northern hemisphere, i.e., in the temperate North Atlantic and North Pacific bioregions up to
boreal altitudes. Its Southern limit is in the Mediterranean Sea (Green & Short, 2003; Short et al.,
2007). *Nanozostera noltei* (Hornemann) (Tomlinson & Posluzny, 2001) (synonym *Zostera noltei*)
ranges from the coasts of Norway in the temperate North Atlantic to Mauritania and occurs widely
in the Mediterranean Sea (Green & Short, 2003). *Cymodocea nodosa* (Ucria) Ascherson, 1870 is
mainly distributed along the Mediterranean coast but also occurs along the North Atlantic coast of
Africa reaching Senegal, and is, therefore, considered as a southern subtropical-Mediterranean
species (Green & Short, 2003; Mascaró et al., 2009). *Ruppia cirrhosa* (Petagna) Grande, 1918
mainly found in coastal waters and salt ponds in the Mediterranean, but also in estuaries and

brackish lagoons in temperate zones of the northern hemisphere (Mannino et al., 2015). Each species has a different geographic distribution, and it can thus be expected that they will show different responses both towards steadily increasing temperatures and MHW. For instance, seawater temperature rise has been reported to enhance mechanisms such as respiration, growth and flowering in seagrasses (Duarte, 2002).

Global warming and MHWs can have widespread impacts on seagrass communities and the ecosystem functions and services they provide (Orth et al., 2006; Strydom et al., 2020). Globally, significant seagrass meadow mortalities and subsequent biodiversity losses have been associated with MHWs occurrences (Lefcheck et al., 2017; Shields et al., 2019; Smale et al., 2019; Strydom et al., 2020). The loss of seagrass habitat will have consequences on the associated functions and services they provide to coastal zones (Duarte, 2002).

Zostera marina is vulnerable with respect to increasing temperatures, both for the populations in its southern range of distribution but also in cooler climates. Shields et al., (2019) showed high temperature driven die-offs of *Z. marina* in Chesapeake Bay (USA) and its replacement by more tolerant *Ruppia* species. Numerous studies have highlighted the negative impact of MHW on *Z. marina*, including reduced growth (Kim et al., 2020) and a decreased photosynthetic performance (Winters et al., 2011). *Cymodocea nodosa,* which is considered as a thermo-tolerant species resilient to high temperatures is expected by some authors to thrive and potentially benefit from global warming in the coming decades (Savva et al., 2018). Modelling studies show that in the Mediterranean Sea under a scenario of rising seawater temperatures, *C. nodosa* could replace more than half of the *Posidonia oceanica* meadows (Papaki et al., 2020). Nevertheless, not all studies converge on this point of view. Hence, it was reported that MHW could have a negative impact on *C. nodosa,* several days after perturbation, i.e., the study of Deguette et al., (2022) shows a decline in the area/DW ratio due to a reduction of the photosynthetic capacity that could reduce growth. On the other hand, it was reported that *C. nodosa* is able to increase photosynthesis and adjust its photo- physiological processes to cope with a MHW (Costa et al., 2021; Marin-Guirao et al., 2016). Cardoso et al., (2008) reported that a MHW during the summer of 2003, had a negative impact by reducing growth production and biomass of *N. noltei* and also on the resilience of a restored population of this species, which ceased to recover after the disturbance. *N. noltei* is negatively impacted by warming conditions, leading to important reduction in shoot density (Repolho et al., 2017). In contrast, another study shows that shoot density was not affected by the simulated MHW (Simão, 2021). In temperate regions, *R. cirrhosa* exhibits resilience to warming (Rasmusson et al., 2021), and the authors suggest that it could outcompete heat-sensitive species. Temperature rise significantly affects *Ruppia* species' germination, growth, and photosynthesis. Studies indicate that *R. cirrhosa* thrives between 20-30°C and maintains viability even at temperatures as low as 10°C, demonstrating greater temperature flexibility compared to other abovementioned seagrass species (Tsioli et al., 2019). The species reaches its thermal limit at temperatures between 32-34°C, beyond which it cannot survive (Tsioli et al., 2019). However, there are no studies on the effects of MHW on *R. cirrhosa* growth.

Experimental laboratory studies have proven essential for understanding the effects of isolated parameters. However, outdoor near-natural mesocosm studies that adopt a more realistic approach by mimicking seasonality, diel cycles, and stochastically fluctuating temperature regimes in Mediterranean lagoons remain absent. The scale of the study is also important to consider, as different responses can be detected depending on the population studied (Marin-Guirao et al., 2018; Nguyen et al., 2021). It is therefore essential to experimentally assess the impact of MHWs in Mediterranean coastal lagoons on seagrass populations, which are naturally adapted to these variable environments. These shallow ecosystems experience significant temperature variations, ranging from 4°C to over 30°C annually, with pronounced diel fluctuations that can reach 16–29°C in spring and 19–40°C in summer. Such strong fluctuations in shallow coastal lagoons are related to heat exchanges driven by the sun and the wind that vary strongly during a diel cycle. So far,

however, most experimental studies (Deguette et al., 2022; Marin-Guirao et al., 2018; Rasmusson
et al., 2021; Repolho et al., 2017; Tsioli et al., 2019), often indoor experiments, have used fixed
temperature settings, which do not reflect the natural conditions. In these cases, the temperatures
of both the controls and the warming treatments remained constant during the day. Consequently,
direct comparisons with populations thriving in their natural environments are tricky. Similarly,
comparisons with findings from the scarce literature remain challenging, as the reported responses
often varied depending on the study location or focused on different variables than those examined
in this study.

This study aims at examining for the first time how three seagrass species (*C. nodosa, N. noltei*,
and *R. cirrhosa*) from a Mediterranean coastal lagoon respond to marine heatwaves of varying
intensities during spring and summer. Because the negative effects of MHWs on *Z. marina* have
been well-documented in the literature as abovementioned, this species is not included in the
present study. We specifically focus on key traits including growth, productivity, and respiration.
Based on two outdoor *ex-situ* near-natural mesocosm experiments that simulated typical
Mediterranean lagoon temperature fluctuations, we evaluated the species' responses to 20-day
marine heatwaves at three intensity levels (+2°C, +4°C, and +6°C above control conditions),
intensity and duration was determined based on preliminary studies conducted by our team (data
from Petton et al., 2024) and aligned with IPCC projections. Our outdoor setup closely mimics a
natural system, reproducing the temperature fluctuations of the shallow coastal lagoon while
preserving natural incident light conditions. Hence, our study allowed us to replicate these
fluctuations for baseline conditions and study the impact of +2 - +6 °C increases above these
fluctuating baselines for the MHWs of different intensities. Applying fixed temperature upon a
naturally fluctuating background as a single factor, while maintaining the other conditions as closely
similar to *in situ* conditions, provides a clearer understanding of the impact of MHWs. We predict
that heatwave intensity will affect growth rates and metabolism differently across species,
depending on their individual thermal tolerances and the seasonal timing of heat exposure.
Furthermore, we expect that the responses observed in this study will differ from those reported in
other ecosystems, as seagrasses in Mediterranean coastal lagoons experience a particularly high
level of environmental variability, which may influence their resilience to thermal stress.

Materials and methods

Study area and biological material collection

The Thau lagoon is a restricted coastal lagoon with two inlets on average 5 metres deep,
(Fiandrino et al., 2017). The averaged water temperature varies during the year between 8 ± 1.8°C
(winter) and 26 ± 1.8°C (summer), while salinity varies between 35 ± 2.0 in early spring and 41 ±
1.5 in summer (REPHY network, 2021: January 2016 - May 2021).

The taxonomy of the Mediterranean lagoon seagrass species is in agreement with the World
Register of Marine Species (WoRMS - https://www.marinespecies.org), which integrated the recent
taxonomic revision in the Zosteraceae family (Tomlinson & Posluzny, 2001; Sullivan & Short,
2023). *Nanozostera noltei* was sampled in the southern part of the Thau lagoon (3.57330 /
43.35990) at approximately 2.5-meter depth. *Cymodocea nodosa* was sampled in the northern
part of the lagoon (3.65503 / 43.40541) at 1.3-meter depth. *Ruppia cirrhosa* was sampled in the
eastern part of the lagoon (3.63793 / 43.39355) at 1.3 meters depth. Since *Zostera marina* reaches
its southern distribution limit in Mediterranean lagoon (Green & Short, 2003; Rueda et al., 2008)
and numerous study have demonstrate its vulnerability to MHWs (see Introduction), it was not
included in the present study.

For each species, a total of 144 shoots with their roots and rhizomes were meticulously
collected by scuba diving on 18 April 2023 and on 18 July 2023 for the spring and summer
experiments, respectively. The collection process involved hand-picking the shoots underwater

from different locations within the seagrass meadow to ensure a representative sampling of the
area. The shoots were then transported to the laboratory using buckets filled with the lagoon water.
The collected shoots were replanted in experimental units (see below) in reconstituted sediments
of the same granulometry. Therefore, sediments were collected from each sampling site and upon
return to the laboratory dried (60°C for 24 hours) and subsequently sieved (2 mm mesh) to remove
fauna and plant residues.

Experimentation

We conducted an outdoor near-natural mesocosm using four fiberglass tanks (200L) filled with
Thau lagoon's water. The experimental setup consisted of four tanks reflecting four treatments:
• A control tank regulated to follow the natural temperature variations of the Thau lagoon.
• A tank with temperature elevated by +2°C above the control.
• A tank with temperature elevated by +4°C above the control.
• A tank with temperature elevated by +6°C above the control
Before heatwave exposure, all specimens were maintained under the same regulated
experimental conditions, reflecting the natural conditions of the Thau lagoon during 15 days.
Heatwaves of varying intensities were applied over a period of 20 days. After heatwave exposure
period, all specimens were maintained under the same conditions as the Control treatment for an
additional 15 days to simulate the natural conditions of the lagoon (Figure 1). We conducted this
experimental design across two seasons: spring and summer 2023.

Figure 1 : Timelines for the experimental conditions and timing of sampling in the
mesocosms showing the main events with respect to natural conditions, i.e.,
acclimation period before heatwave exposure (blue), during heatwave exposure
(orange), and return period after heatwave exposure (green). Arrows with their
associated pictograms indicate the timing of samplings for morphometry
measurements on seagrass shoots and oxygen fluxes measurements.

To regulate the temperature, we used a heater and a cooler in each tank. A sensor was
deployed in the Thau lagoon to measure the real-time water temperature (readings taken every 10
seconds). This data was automatically transmitted to a program, enabling real-time adjustments of
the water temperature in the control tank to the same temperature as measured in the lagoon. In
parallel, in the tanks used for the other treatments, the temperatures were adjusted following the
timelines shown in Figure 1. Three pumps per tank recirculated the water to ensure a
homogeneous water bath.

In each experimental unit, six shoots, along with their rhizomes and roots, were selected. The
rhizome segments were standardized according to species-specific protocols: for *N. noltei*,
rhizomes were adjusted to three segments; for *R. cirrhosa*, rhizomes were adjusted to two

segments in spring experiments and three segments in summer experiments; and for *C. nodosa*,
only horizontal rhizomes were retained. The standardized shoots were then placed in individual 3L
cylindrical glass experimental units (10 cm in diameter, 30 cm in height) that had been pre-filled
with 550 cm³ of sediment (approximately 7 cm in depth). This resulted in a total of 36 shoots per
treatment (control, +2°C, +4°C, and +6°C) for each species. Subsequently, each set of six
experimental units per species was arranged in experimental tanks, leading to a total of 18
experimental units per tank.

Water temperature was monitored continuously (a reading every 5 minutes) with autonomous
sensors (HOBO, UA-002-64, ± 0,2°C) placed in each experimental tank. The experimental system
was set up outdoors to take advantage of the natural variation in light, with the aim of reproducing
in-situ conditions as closely as possible. The available light intensity was continuously measured
in close proximity to the experimental system using an LI-190 planar sensor probe connected to a
data acquisition unit (Campbell, CR1000).

The water level and salinity in the experimental tanks were monitored three times a day, and
adjustments were made by refilling with fresh water to offset evaporation. Additionally, the water in
each treatment was refreshed weekly to maintain adequate nutrient levels. The samples for
nutrients concentrations of nutrients (NH_4^+, NO_3^-+NO_2^- and PO_4^{3-}) was estimated from spot
samples taken twice a week.

Analysis

Morphological, physiological and metabolic responses of three seagrass species were
measured throughout the experiment (Figure 1). Nutrients sample was analysed in the laboratory
using the method described by David et al. (2019).

At the onset of the experiment, we conducted initial morphological measurements for each
species. Leaf area was measured (length × width) and the number of leaves per shoot was counted
for all species. Leaf measurements were performed on individual shoots, involving the temporary
removal of each specimen from the sediment. Subsequently, measurements were taken for each
specimen, including the number of shoots, leaves, length and width of leaves, length and width of
rhizomes, ensuring a comprehensive assessment of morphological characteristics. After the
measurements, each specimen was carefully placed back into the experimental units during each
measurement session.

These measurements, which reflect the creation of new plant matter, were first used to calculate
individual leaf growth rates. For each leaf, the growth rate was determined using the following
formula:

(1) Individual leaf growth rate = $iLGR\ (mm^2 d^{-1}) = LAtf - LAti\ /\ tf - ti$

where LA is the leaf area (length * width) at final time (tf) and initial time (ti). For newly formed
leaves, corrections were applied. Negative values, which represented cut leaves, were adjusted to
zero. To obtain the growth rate ($mm^{-2}.d^{-1}$) at the shoot level, these individual leaf growth rates were
then summed for all leaves within each shoot:

(2) shoot leaf growth rate = $LGR\ (mm^2 d^{-1}) = \sum (iLGR)$

New leaf gain rate (mm-2.d-1) was quantified by tracking individual leaf appearance between
consecutive measurements (ti and tf). For each shoot, new leaf gain rate was calculated as:

(3) New leaf gain rate = $NewLeafGR\ (mm^2 d^{-1}) = \sum (LA[new\ leaves]\ /\ tf - ti)$

where LA is the recorded area of new leaf. Shoots without leaves creation were assigned a
zero rate.

Old leaf gain rate (mm^{-2}.d^{-1}) was quantified by tracking the retention of two leaves between
consecutive measurement times (ti and tf). For each shoot, old leaf gain rate was calculated as:

(4)   Old leaf gain rate = OldLeafGR (mm²d^{-1}) = $\sum$ (LA[*common leaves*] / tf – ti)

where LA is the recorded area of common leaf.

The Leaf Growth Rate (*LGR*) was defined as the sum of *NewLGR* and *OldLeafGR*, hence

(5)   Leaf growth rate = *LGR (mm²d^{-1})* = *NewLeafGR + OldLeafGR*

In addition, we calculated a *ratio* $\frac{NewLeafGR}{OldLeafGR}$ *(hereafter referred to as: New-Old leaf* ratio) to
characterise the proportions of growth allocated to the formation of new and old leaves,
respectively.

Because the leaf growth rate does not take into account the loss of leaf area due to the falling
leaves, the leaf loss rate (mm^{-2}.d^{-1}) were quantified by tracking individual leaf disappearance
between consecutive measurements (ti and tf). For each shoot, leaf loss rate was calculated as:

(6)   Leaf loss rate = *LLR* (mm²d^{-1}) = $\sum$ (LA[*lost leaves*] / tf – ti)

where LA is the last recorded leaf area before loss. Shoots without leaf loss were assigned a
zero rate.

In each experimental unit, at the end of heatwave exposure, oxygen fluxes were measured
using optical sensor (WTW FDO 925) through short-term incubations of 1 to 1.5 hours per session
(Figure 1). This process involved calculating the difference between initial and final oxygen
concentrations in both dark and light conditions to determine community respiration (CR: negative
or zero) and Net Community Production (NCP: negative, positive or zero), respectively. Gross
Community Productivity (GCP, positive or zero) was calculated by adjusting NCP for CR and
normalizing it to the total leaf area (*tot LA*) of seagrass shoots in each experimental unit at the time
of measurement.

(7)   Gross Community Productivity = GCP (O² mg.mm^{-2} h^{-1}) = (NCP – CR / tot LA)

Statistical analysis

All statistical analyses were conducted using Rstudio (R version 4.4.1) with significance set at
p=0.05. To assess the impact of different experimental conditions on temperature, daily averages
of the measurements were first calculated. Due to non-normality, daily temperature averages were
assessed across experimental periods using the Kruskal-Wallis test, with subsequent pairwise
comparisons conducted using the Dunn test and Benjamini-Hochberg correction. Similar Kruskal-
Wallis tests were applied to salinity, light, and nutrient concentration data. Leaf Growth rate (*LGR*)
and Leaf Loss rate (*LLR*) were analysed using linear mixed-effect models, with treatments and
experimental periods as fixed effects and experimental units (replicate) in random effects. Post-
hoc pairwise comparisons were using emmeans package, with p-value adjusted for multiple
comparisons using the Tuckey method. Graphic diagnostics were used to verify normality and
homogeneity of variances. Oxygen fluxes were analysed using Kruskal-Wallis test. Occasionally,
when during the acclimation period, treatments presented a *LGR* significantly different from the
control, we did not further report the pairwise comparisons between these treatments and the
control during heatwave exposure (noted as "*nc*" in Table 2-a), as their response might have

depended on prior conditions rather than solely on the experimental treatment. Nevertheless, within these treatments it was still meaningful to compare *LGR* during the heatwave and return periods with the acclimation period (Table 2-b).

Results

Experimental conditions

Table 1: Environmental conditions in the incubation system, including photoperiod, daily light integrate (DLI), salinity, and concentrations of nutrients (NH4, NOx, and PO4) before, during, and after heatwave exposure, for both spring and summer seasons. Photoperiod and DLI data were calculated in proximity to the experimental system, while the other parameters were measured on the experimental tanks water. The values represent the means ± standard deviation of all conditions (control, +2, +4, and +6) since no statistical differences were observed.

		Before	Heatwave	After
		15 days	*20 days*	*15 days*
Spring	Photoperiod (h)	13.40 - 13.95	14.02 - 14.57	14.53 - 14.92
	DLI (mol photons.m^{-2}.d^{-1})	23.5 ± 16.4	32.7 ± 11.8	43.4 ± 6.1
	Salinity	38.5 ± 0.7	39.3 ± 0.9	38.8 ± 0.7
	NH4 (µmoL.L^{-1})	0.63 ± 0.95	0.50 ± 0.85	0.63 ± 1.03
	NOx (µmoL.L^{-1})	0.40 ± 0.34	0.38 ± 0.29	0.52 ± 0.57
	PO4 (µmoL.L^{-1})	0.075 ± 0.05	0.109 ± 0.08	0.163 ± 0.12
Summer	Photoperiod (h)	14.77- 14.30	14.48 - 13.47	13.43 - 12.83
	DLI (mol photons.m^{-2}.d^{-1})	39.8 ± 12.25	32.6 ± 9.73	27.9 ± 11.16
	Salinity (psu)	40.67 ± 1.76	39.70 ± 1.45	40.53 ± 1.63
	NH4 (µmoL.L^{-1})	0.43 ± 0.64	0.59 ± 1.09	0.55 ± 0.84
	NOx (µmoL.L^{-1})	0.49 ± 0.36	0.57 ± 0.66	0.45 ± 0.44
	PO4 (µmoL.L^{-1})	0.06 ± 0.05	0.07 ± 0.05	0.07 ± 0.06

Light, salinity and inorganic nutrient concentrations in the outdoor experimental set-up during experiments are show in Table 1 for the spring and summer experiments, respectively. The length of the photoperiod increased by 11.3 % (1.52 hours) from the start to the end of the spring experiment, accompanied by a rise in the average daily light integral (DLI) by 84.7 % (Table 1; DLI = 23.5 ± 16.4 to 43.4 ± 6.1 mol photons.m^{-2}.d^{-1}). In contrast, during the summer experiment, the length of the photoperiod decreased by 13.1 % (1.93 hours), leading to a corresponding decline in average DLI of 29.9 % (39.8 ± 12.25 to 27.9 ± 11.16 mol photons.m^{-2}.d^{-1}). Salinity was homogeneous among treatments and experimental periods with an overall mean of 38.9 ± 0.9 psu

337 in spring and 40.2 ± 1.6 psu in summer. Nutrient concentrations remained stable during both
338 experiments (Table 1).

339   Before heatwave exposure (acclimation periods), water temperatures for both spring and

Figure 2 : Temperature variations in spring (left) and in summer (right) before heatwave exposure (in blue), during heatwave exposure (in orange), and after heatwave exposure (in green) for the 'control', '+2', '+4', and '+6' treatments. Kruskal-Wallis tests indicate the significant differences between treatments. Data are represented as box plots, with the mean shown in red. The horizontal line within each box represents the median, and the edges of the boxes correspond to the 25th and 75th percentiles. The ends of the whiskers indicate the 5th and 95th percentiles, and the dots represent outliers.

summer experiments were uniform across treatments (Figure 2). Following seasonal patterns,
water temperatures were consequently higher in summer than in spring, i.e., 27.80 ± 3.80 °C, and
20.63 ± 3.03 °C, respectively. During the heatwave exposure periods, both in spring and summer,
the temperature in the control treatments (+ 0 °C) did not vary significantly with respect to the
acclimation period. In contrast, there were significant temperature differences between the control
and the three heatwave treatments. Post hoc comparisons indicated that water temperature
increased significantly for the +2°C, +4°C, and +6°C treatments compared to the control, with mean
differences of 1.69 ± 0.55°C (p = 0.02), 4.07 ± 1.16°C (p < 0.001), 5.78 ± 1.02°C (p < 0.001),
respectively in spring and 1.51 ± 1.40°C (p = 0.06), 3.02 ± 1.65°C (p < 0.001), 4.39 ± 1.66°C (p <
0.001) in summer. During the final experimental period (the return phase), each treatment returned
to follow the natural temperature fluctuations in Thau lagoon (Figure 2). Unexpectedly, the return
phase in spring coincided with an uncommon natural heatwave that was comparable to a heatwave
treatment of approximately +4°C (25.12 ± 2.46 °C). In contrast for the summer experiment, the
temperatures were lower than during the acclimation and heatwave exposure periods (24.93 ±
3.67 °C), as expected for the season.

Seagrass responses to experimental conditions

While all the statements based on pairwise comparisons of experimental results for Leaf
Growth rate (*LGR*) and Leaf Loss rate (*LLR*) reported in this section are supported by statistical

tests (p < 0.05), we did neither report their results in the text nor in the corresponding figures
(Figures. 2- 4, panels a-d) to maintain their readability, but rather refer to Table 2 (Post-hoc pairwise
comparisons with the p-values adjusted for multiple comparisons using the Tuckey method, see
methods).

Table 2: Pairwise comparisons of the effects of heatwave treatment and experimental period on the *LGR* and *LLR* of *Cymodocea nodosa* (CN), *Nanozostera noltei* (ZN), and *Ruppia cirrhosa* (RC). "nc" indicates that statistics were not calculated when there was a significant difference (p > 0.05) during the acclimation phase ("before").*Ruppia cirrhosa* (RC).

(a) Pairwise comparison – treatments effect

| | | | Spring | | | | | | | | Summer | | | | | | | |
| | | | before | | | heatwave | | | after | | | before | | | heatwave | | | after | | |
			df	t.ratio	p-value	df	t.ratio	p-value	df	t.ratio	p-value	df	t.ratio	p-value	df	t.ratio	p-value	df	t.ratio	p-value
LGR	CN	"control"-"+2°C"	379	-0,44	0,97	379	-2,85	**0,024**	379	2,40	0,08	371	1,37	0,52	371	1,52	0,42	372	0,46	0,97
		"control"-"+4°C"	379	-0,61	0,93	379	-4,53	**<0,001**	379	1,01	0,74	371	0,38	0,98	371	0,06	1,00	371	0,40	0,98
		"control"-"+6°C"	379	-3,66	**0,002**	nc	nc	nc	nc	nc	nc	371	1,39	0,51	371	-0,55	0,95	372	0,98	0,76
	ZN	"control"-"+2°C"	366	-0,63	0,92	366	-1,73	0,31	366	2,09	0,16	351	1,83	0,26	351	-0,02	1,00	351	0,77	0,87
		"control"-"+4°C"	366	0,79	0,86	366	-1,58	0,39	366	0,94	0,78	351	4,05	**<0,001**	nc	nc	nc	nc	nc	nc
		"control"-"+6°C"	366	0,88	0,81	366	0,96	0,77	366	2,72	**0,03**	351	0,91	0,80	351	1,99	0,20	352	2,53	**0,05**
	RC	"control"-"+2°C"	297	-1,41	0,49	298	-0,80	0,85	298	-0,03	1,00	343	-0,46	0,97	343	-2,20	0,13	343	-1,51	0,43
		"control"-"+4°C"	297	-0,50	0,96	298	-1,03	0,73	298	1,30	0,57	343	4,70	**<0,001**	nc	nc	nc	nc	nc	nc
		"control"-"+6°C"	297	-1,97	0,20	297	-2,13	0,14	298	-0,83	0,84	343	3,63	**<0,001**	nc	nc	nc	nc	nc	nc
LLR	CN	"control"-"+2°C"	403	0,36	0,98	403	0,62	0,92	403	5,61	**<0,001**	403	2,47	0,07	403	-0,32	0,99	403	-0,51	0,96
		"control"-"+4°C"	403	-0,36	0,98	403	0,80	0,85	403	6,65	**<0,001**	403	-0,24	1,00	403	0,22	1,00	403	1,21	0,62
		"control"-"+6°C"	403	-0,13	1,00	403	1,98	0,20	403	7,67	**0,00**	403	0,39	0,98	403	2,68	**0,04**	403	0,69	0,90
	ZN	"control"-"+2°C"	398	-0,18	1,00	398	0,94	0,78	398	0,75	0,88	397	0,31	0,99	397	-0,85	0,83	397	0,71	0,89
		"control"-"+4°C"	398	-0,61	0,93	398	1,30	0,56	398	-0,85	0,83	397	-2,31	0,10	397	-1,12	0,68	398	0,21	1,00
		"control"-"+6°C"	398	-1,91	0,23	398	1,13	0,67	398	-1,99	0,19	397	-0,13	1,00	397	-1,26	0,59	397	2,02	0,18
	RC	"control"-"+2°C"	369	1,14	0,66	369	0,70	0,90	370	0,45	0,97	390	0,30	0,99	390	0,23	1,00	390	-1,54	0,42
		"control"-"+4°C"	369	0,39	0,98	369	-0,58	0,94	371	-0,43	0,97	390	-1,19	0,63	390	0,94	0,79	391	0,49	0,96
		"control"-"+6°C"	369	0,01	1,00	369	1,46	0,46	369	1,07	0,71	390	-0,15	1,00	390	0,43	0,97	391	0,45	0,97

(b) Pairwise comparison : experimental period effect

| | | | Spring | | | | | | | | | | | | Summer | | | | | | | | | | | |
| | | | "control" | | | "+2°C" | | | "+4°C" | | | "+6°C" | | | "control" | | | "+2°C" | | | "+4°C" | | | "+6°C" | | |
			df	t.ratio	p-value	df	t.ratio	p-value	df	t.ratio	p-value	df	t.ratio	p-value	df	t.ratio	p-value	df	t.ratio	p-value	df	t.ratio	p-value	df	t.ratio	p-value
LGR	CN	before-heatwave	379	-0,61	0,82	379	-3,09	**0,006**	379	-4,63	**<0,001**	379	-5,47	**<0,001**	371	2,98	**0,009**	371	3,19	**0,004**	371	2,66	**0,02**	371	1,05	0,54
		heatwave-after	379	-4,19	**<0,001**	379	1,04	0,55	379	1,27	0,42	379	4,58	**<0,001**	371	4,28	**<0,001**	371	6,24	**<0,001**	371	4,68	**<0,001**	371	6,84	**<0,001**
		before-after	379	-4,87	**<0,001**	379	-1,95	0,13	379	-3,23	**0,004**	379	-0,66	0,79	371	7,24	**<0,001**	371	3,19	**0,004**	371	7,37	**0**	371	5,73	**<0,001**
	ZN	before-heatwave	366	-1,13	0,49	366	-2,29	**0,05**	366	-3,51	**<0,001**	366	-0,97	0,60	351	2,89	**0,01**	351	1,09	0,52	351	0,43	0,90	351	4,07	**<0,001**
		heatwave-after	366	1,83	0,16	366	5,57	**<0,001**	366	4,26	**<0,001**	366	3,54	**<0,001**	352	1,16	0,48	351	2,00	0,11	351	1,40	0,34	352	2,05	0,10
		before-after	366	0,77	0,72	366	3,52	**<0,001**	366	0,99	0,58	366	2,76	**0,02**	352	3,85	**<0,001**	352	3,07	**0,007**	351	1,86	0,15	351	5,89	**<0,001**
	RC	before-heatwave	297	-0,64	0,80	297	-0,14	0,99	298	-1,19	0,46	297	-1,07	0,54	343	3,06	**0,007**	343	1,24	0,43	343	-3,12	**0,006**	344	-1,26	0,42
		heatwave-after	298	-0,65	0,79	298	0,07	1,00	297	1,61	0,24	298	0,43	0,90	343	1,38	0,35	343	1,98	0,12	344	2,38	**0,047**	344	2,78	0,02
		before-after	297	-1,25	0,42	298	-0,05	1,00	297	0,69	0,77	298	-0,49	0,88	343	4,27	**<0,001**	343	3,24	**0,004**	344	-0,33	0,94	344	1,74	0,19
LLR	CN	before-heatwave	403	0,63	0,80	403	0,90	0,64	403	1,79	0,17	403	2,74	**0,02**	403	3,21	**0,004**	403	0,42	0,91	403	3,67	**<0,001**	403	5,50	**<0,001**
		heatwave-after	403	3,75	**<0,001**	403	8,87	**0**	403	9,76	**0**	403	9,65	**0**	403	-1,12	0,50	403	-1,33	0,38	403	-0,10	0,99	403	-3,04	0,007
		before-after	403	4,37	**<0,001**	403	9,75	**0**	403	11,52	**0**	403	12,33	**0**	403	2,01	0,11	403	-0,92	0,63	403	3,48	**<0,001**	403	2,34	0,05
	ZN	before-heatwave	398	-1,92	0,14	398	-0,80	0,71	398	-0,0042	1,00	398	1,12	0,50	397	-2,58	0,03	397	-3,74	**<0,001**	397	-1,38	0,35	397	-3,71	**<0,001**
		heatwave-after	398	3,45	**0,002**	398	3,28	**0,003**	398	1,29	0,40	398	0,26	0,96	397	-3,22	**0,004**	398	-1,66	0,22	397	-1,89	0,14	397	0,06	1,00
		before-after	398	1,58	0,26	398	2,50	**0,03**	398	1,29	0,40	398	1,33	0,38	397	-5,72	**<0,001**	398	-5,25	**<0,001**	397	-3,20	**0,004**	397	-3,56	**<0,001**
	RC	before-heatwave	369	1,35	0,37	369	0,92	0,63	369	0,37	0,93	369	2,81	**0,01**	390	1,19	0,46	390	1,26	0,42	390	-0,93	0,62	390	0,62	0,81
		heatwave-after	370	4,22	**<0,001**	369	3,87	**<0,001**	371	3,67	**<0,001**	370	3,65	**<0,001**	390	-0,11	0,99	390	1,69	0,21	390	0,26	0,96	391	-0,17	0,98
		before-after	370	5,52	**<0,001**	369	4,72	**<0,001**	371	3,99	**<0,001**	370	6,09	**<0,001**	390	1,04	0,55	390	2,92	**0,01**	390	-0,60	0,82	391	0,41	0,91

Acclimation periods before heatwave exposure

366   Prior to heatwave exposure (acclimation period), leaf growth rate (*LGR*) with 4 exceptions (see Table 2) and all leaf loss rates (*LLR*) were homogeneous across treatments for all species in both spring and summer. The *New-Old leaf* ratio was close to one for the three species in both spring and summer, which showed an equilibrium between growth of new and old leaves. On average, the rate of gain of new leaves was higher in summer for *C. nodosa* and *R. cirrhosa* but lower for *N. noltei*. Meanwhile, the old leaf gain rate was higher in summer for *C. nodosa* and *N. noltei* and remained similar for *R. cirrhosa* (Figures 3b–e, 4b–e, and 5b–e).

Figure 3: Effects of experimental marine heatwaves on the growth and physiological responses of *Cymodocea nodosa*. (a, d) Leaf Growth Rate (*LGR*) and Leaf Loss Rates (*LLR* - mm² d⁻¹) measured in spring and summer across four experimental treatments (Control, +2°C, +4°C, +6°C). Statistical significance is indicated for

experimental effects (E), period effects (P), and their interaction (E:P) with their p-
values derived from the ANOVA of a linear mixed-effects model (lmerTest), post-hoc
pairwise comparisons are listed in Table 2. (b, e) *New-Old leaf* ratio across
experimental periods (Before, Heatwave, After). Data points are colored by period,
and different symbols indicate temperature treatments. The dashed line represents
a 1:1 relationship. Larger points represent the means of the respective data clouds.
(c, f) Gross oxygen production and respiration rates (mg O_2 mm^{-2} h^{-1}) under
different temperature treatments. Kruskal-Wallis tests indicate no significant
differences between treatments. Boxplots display the distribution of the data, with
the mean shown in red. The horizontal line within each box represents the median,
while the box edges correspond to the 25th and 75th percentiles. Whiskers extend
to the 5th and 95th percentiles, and individual points represent extreme values

Cymodocea nodosa

Cymodocea nodosa response during the spring experiment

No mortality was observed for *C. nodosa* throughout the entire spring experiment. During
heatwave exposure, *LGR* in Controls remained constant compared to the acclimation period (14.11
± 6.19 mm².d-1), whereas it significantly increased in the heatwave treatments (+2°C = 19.85 ±
7.78 mm².d^{-1}; +4°C = 23.23 ± 8.31 mm².d^{-1}; +6°C = 30.76 ± 8.80 mm².d^{-1}). After heatwave
exposure, *LGR* was homogeneous across treatments. Hence, upon the transition from heatwave
exposure to the return period, *LGR* increased in controls (22.71 ± 8.90 mm².d^{-1}), remained steady
for the +2°C and +4°C treatments (17.65 ± 10.48 mm².d^{-1} and 20.58 ± 10.23 mm².d^{-1}, respectively)
and decreased for the +6°C treatment (21.26 ± 12.56 mm².d^{-1}). The variability of these *LGR* was
highest in the +6°C treatment (Figure 3a, Table 2). *LLR* of *C. nodosa*, in spring (Figure 3a),
increased during the experiment, the *LLR* was homogeneous between treatments during heatwave
exposure (Figure 3a). After heatwave exposure, *LLR* was significantly higher for those individuals
who had experienced a heatwave (Figure 3a). The relationship between the *NewLeafGR* and the
OldLeafGR remained stable for the control, but showed an increasing *New-Old leaf* ratio for
treatments during heatwave exposure (Figure 3b). This increase was more pronounced in the +4°C
and +6°C treatments, where the ratio exceeded 3. After heatwave exposure, the *New-Old leaf* ratio
increased for controls (ratio = 2.6) but decreased for treatment (still higher than 1, Figure 3b).
During heatwave exposure both gross oxygen production and respiration rates were similar across
all treatments (Figure 3c).

Cymodocea nodosa response during the summer experiment

No mortality was observed before heatwave exposure, whatever the treatment. During
heatwave exposure, one shoot died in the controls, three shoots at +2°C, none at +4°C and one
at +6°C. After heatwave exposure, mortality remained low with one shoot in the controls, two at
+2°C, five at +4°C and three at +6°C. During heatwave exposure, no significant differences were
observed in *LGR* across treatments. The *LGR* of the Control treatment and the +2°C and the +4°C
treatments significantly decreased compared to the period before heatwave exposure (19.52 ±
7.54 mm².d^{-1}, 15.89 ± 8.72 mm².d^{-1} and 19.38 ± 9.43 mm².d^{-1}, respectively), while the *LGR*
remained similar for the +6°C treatment (20.84 ± 8.91 mm².d^{-1}). After heatwave exposure, *LGR*
was homogeneous across treatments. Compared to heatwave exposure, *LGR* drastically
decreased for all treatments (Figure 3d, Table 2b). During heatwave exposure, *LLR* significantly
increased for the Control, +4°C, and +6°C treatments, while the *LLR* remained steady for the +2°C
treatment. *LLR* during heatwave exposure was highest for the +6°C treatment (Figure 3d). After
heatwave exposure, *LLR* was similar across treatments and comparable to the period during
heatwave exposure, except for the +6°C treatment where it significantly decreases (-24.02 ± 22.38
mm².d^{-1}; Figure 3d). During heatwave exposure, the relationship between *NewLeafGR* and
OldLeafLGR of all treatments deviated from equilibrium by increasing the *New-Old leaf* ratio. The

mean *OldLeafLGR* decreased sharply and approaches zero for the +4°C and +6°C treatments, which directly impacts the ratio, making it particularly high for these two treatments (ratio = 13.7 ± 18.00; 56.56 ± 87.46, respectively, Figure 3e). The mean *NewLeafGR* remained similar to the Control and +2°C treatments. After heatwave exposure, in all treatments the *New-Old leaf* ratio returned to close to 1, however with lower *NewLeafGR* and *OldLeafLGR* than observed before heatwave exposure (Figure 3e). During heatwave exposure, both gross oxygen production and respiration rates were similar across all treatments (Figure 3f).

Figure 4: Effects of experimental marine heatwaves on the growth and physiological responses of *Nanozostera noltei*. (a, d) Leaf Growth Rate (*LGR*) and Leaf Loss Rates (*LLR* - mm² d⁻¹) measured in spring and summer across four experimental treatments (Control, +2°C, +4°C, +6°C). Statistical significance is indicated for experimental effects (E), period effects (P), and their interaction (E:P) with their p-values derived from the ANOVA of a linear mixed-effects model (lmerTest), post-hoc

pairwise comparisons are listed in Table 2. (b, e). *New-Old leaf* ratio (growth ratio of
new to old leaves) across experimental periods (Before, Heatwave, After). Data
points are colored by period, and different symbols indicate temperature treatments.
The dashed line represents a 1:1 relationship. Larger points represent the means of
the respective data clouds. (c, f) Gross oxygen production and respiration rates (mg
O_2 mm^{-2} h^{-1}) under different temperature treatments. Kruskal-Wallis tests indicate
no significant differences between treatments. Boxplots display the distribution of
the data, with the mean shown in red. The horizontal line within each box represents
the median, while the box edges correspond to the 25th and 75th percentiles.
Whiskers extend to the 5th and 95th percentiles, and individual points represent
extreme values.
451

Nanozostera noltei

Nanozostera noltei response during the spring experiment

Overall shoot mortality was very low for the 36 shoots used for each treatment. In the control
treatments, no shoot was lost. At +2°C one shoot died during heatwave exposure and another
after. At +4°C, one shoot died during this period, and two afterwards. At +6°C, three shoots died
during exposure, and two more afterwards. During heatwave exposure, *LGR* of the Control
treatment remained stable compared to the acclimation period, while the *LGR* of the other
treatments was not statistically different from the Control (Figure 4a - Table 2a). However, pairwise
comparisons within each treatment comparing the acclimation and subsequent heatwave
exposure periods showed an increase in the +4°C treatment between the before and heatwave
exposure periods (Figure 4a – Table 2b). After heatwave exposure, the *LGR* values remained
similar between treatments, with only the +6°C treatment showing significantly lower rates.
Specifically, while the Control treatment maintained stable *LGR* values, all experimental treatments
showed decreased rates (Figure 4a). The *LLR* remained homogeneous with the Control across all
treatments throughout the entire experiment (Figure 4a – Table 2a). However, pairwise
comparisons within each experiment indicate a tendency for decrease in the Control and +2°C
treatment between heatwave exposure and after (Figure 4a – Table 2b). During heatwave
exposure, *NewLeafGR* increased across all treatments, with the most substantial rise observed at
+2°C and +4°C, where it more than doubled compared to before heatwave exposure (Figure 4b).
OldLeafGR generally decreased, though this reduction was less pronounced in the +6°C treatment
(Figure 4b). The *New-Old leaf* ratio rise significantly in all treatments, especially at +4°C, indicating
a shift toward new leaf growth over old leaf maintenance. After heatwave exposure, *NewLeafGR*
returned to levels similar to or below exposure period, while *OldLeafGR* recovered slightly across
all treatments. Consequently, the ratio dropped back close to pre-heatwave values, reflecting a
return toward equilibrium. During heatwave exposure, both gross oxygen production and
respiration rates were similar across all treatments (Figure 4c).

Nanozostera noltei response during the summer experiment

One shoot died before heatwave exposure in the controls, two shoots at +2°C, three at +4°C,
and one shoot at +6°C. During heatwave exposure, six shoots died in the controls group, five at
+2°C, four at +4°C, and four plants at +6°C. After heatwave exposure, two plants died in the
controls, one at +2°C, one at +4°C, and six at +6°C. During heatwave exposure, no significant
differences were observed in *LGR* across treatments (Figure 4d – Table 2a). The *LGR* of the
Control treatment and the +6°C treatment significantly decreased compared to the period before
heatwave exposure (10.55 ± 6.75 mm².d^{-1} and 7.05 ± 4.07 mm².d^{-1}, respectively), while the *LGR*
remained similar for the +2°C and the +4°C treatment (10.64 ± 6.38 mm².d^{-1} and 7.97 ± 4.69 mm².d^{-1}
- Figure 4d – Table 2b). After heatwave exposure, the *LGR* was homogeneous across treatments,
with a tendency to be lower in the +6°C treatment (Figure 4d). During heatwave exposure, *LLR*
significantly decreased for the Control, +2°C, and +6°C treatments, while the *LLR* remained steady
for the +4°C treatment. The *LLR* during heatwave exposure were significantly highest for the +6°C

treatment (Figure 4d). After heatwave exposure, *LLR* was still decreasing for the Control treatment (-8.17 ± 11.28 mm².d^{-1}), and remained homogeneous for the treatments (+2°C = -13.31 ± 19.47 mm².d^{-1}; +4°C = -9.73 ± 15.27 mm².d^{-1}; +6°C = -22.65 ± 28.39 mm².d^{-1}). During heatwave exposure, *NewLeafGR* increased across all treatments, with the highest rise in the Control and +2°C treatments (Figure 4e). *OldLeafGR* decreased sharply in the control and +2°C treatments. The *New-Old leaf* ratio increased in the Control and +2°C treatments, indicating a shift toward new leaf growth, whereas it remained much lower for +4°C and +6°C (Figure 4e). After heatwave exposure, the *New-Old leaf* ratio returned close to 1 with lower *NewLeafGR* and *OldLeafLGR* values than before heatwave. During heatwave exposure, both gross oxygen production and respiration rates were similar across all treatments (Figure 4f).

Figure 5: Effects of experimental marine heatwaves on the growth and physiological responses of *Ruppia cirrhosa*. (a, b) Leaf Growth Rate (*LGR*) and Leaf Loss Rates (*LLR* - mm² d⁻¹) measured in spring and summer across four experimental treatments (Control, +2°C, +4°C, +6°C). Statistical significance is indicated for experimental effects (E), period effects (P), and their interaction (E:P) with their p-values derived from the ANOVA of a linear mixed-effects model (lmerTest). Post-hoc pairwise comparisons are listed in Table 2. (b, e) *New-Old leaf* ratio (growth ratio of

new to old leaves) across experimental periods (Before, Heatwave, After). Data points are colored by period, and different symbols indicate temperature treatments. The dashed line represents a 1:1 relationship. Larger points represent the means of the respective data clouds. (c, f) Gross oxygen production and respiration rates (mg O_2 mm^{-2} h^{-1}) under different temperature treatments. Kruskal-Wallis tests indicate the significant differences between treatments. Boxplots display the distribution of the data, with the mean shown in red. The horizontal line within each box represents the median, while the box edges correspond to the 25th and 75th percentiles. Whiskers extend to the 5th and 95th percentiles, and individual points represent extreme values.

Ruppia cirrhosa

Ruppia cirrhosa response during the spring experiment

   Ruppia cirrhosa showed substantially more mortality than the other two species. Ten shoots died in the controls in total (5 during heatwave, 5 after). At +2°C, 13 shoots died (1 before, 6 during heatwave, 6 after). At +4°C, 16 shoots died (4 before, 10 during heatwave, 2 after). At +6°C, 14 shoots died (11 during heatwave, 3 after). The *LGR* stayed homogeneous during and after heatwave exposure for all treatments (Figure 5a – Table 2a). During heatwave exposure, *LLR* remained homogeneous with the Control across all treatments. However, pairwise comparisons within each treatment indicated an increase in *LLR* in the +6°C treatment compared to before heatwave exposure (+6°C = -3.07 ± 3.42 mm².d^{-1}; Table 2b). After heatwave exposure, *LLR* was homogeneous across all treatment, but showed an overall increase compared to the heatwave period (Figure 5a – Table 2b). During heatwave exposure, *NewLeafGR* increased across all treatments, with the most pronounced rise at +4°C, where it nearly doubles. *OldLeafGR* decreases in all conditions, with the sharpest decline in +4°C treatments (Figure 5b). *The New-Old leaf* ratio increased across treatments, especially in +4°C, indicating a shift toward growth of new leaves over old leaves. After heatwave exposure, *NewLeafGR* remains high or continued increasing, *OldLeafGR* showed a slight increase in some treatments, and the *New-Old leaf* ratio remained elevated, particularly in +4°C (Figure 5b). During heatwave exposure. both gross oxygen production and respiration rates were similar across all treatments (Figure 5c).

Ruppia cirrhosa response during the summer experiment

   In the controls, 9 shoots died in total (5 during heatwave, 4 after). At +2°C, 7 shoots died (1 during heatwave, 6 after). At +4°C, 12 shoots died (4 before, 6 during heatwave, 2 after). At +6°C, 11 shoots died (3 before, 4 during heatwave, 4 after). During heatwave exposure, Control *LGR* decreased compared to before exposure (2.64± 1.40 mm².d^{-1}). *LGR* was homogeneous across treatments, but increased compared to before exposure for the +4°C treatment (3.45 ± 1.90 mm².d^{-1}). After heatwave exposure, *LGR* was homogeneous across all treatments, but decreased for +4°C and +6°C treatment compared to during heatwave exposure (Figure 5d – Table 2a). During heatwave exposure, *NewLeafGR* increased, particularly in the +4°C treatment, where it more than tripled compared to before heatwave exposure. In contrast, *OldLeafGR* generally decreased across all treatments. After heatwave exposure, the *New-Old leaf* ratio remained above 1, with *NewLeafGR* staying stable in the +2°C treatment and decreasing in the others. *OldLeafGR* remained stable compared to its levels during heatwave exposure (Figure 5e). During heatwave exposure, both gross oxygen production and respiration rates were similar across all treatments except for the +2°C treatment, where gross oxygen production was significantly lower than the Control (Figure 5f).

Discussion

Summary of Species Responses

The results of these two experiments highlight different responses depending on the intensity of the simulated heatwave, the season, and the studied species. *C. nodosa* behaved differently in spring and summer. In spring, this species showed a clear stimulation of *LGR* during all MHW, which coincided with a shift to new leaves growth (increase of the *New-Old leaf* ratio); albeit these positive effects were off-set during the return phase by increased *LLR*. Nevertheless, the species appeared very well adapted to both moderate (i.e., +2°C) and severe (+ 6°C) MHW in spring, by increasing the leaves' turnover while maintaining high net growth rates. During summer, this species was also well adapted to the MHW, also showing a shift towards new leaves formation, but without an increase of *LGR*. The return phase was again characterized by an increase of leave losses *LLR*, with a concomitant reduction of *LGR*. This was particularly the result of lower *NewLeafGR* than during the previous MHW, which may be related to the decreasing light conditions towards the end of the summer season (see Table 1). *N. noltei* showed more subtle differences between these two seasons. In spring, the +4 °C treatments resulted in a slight stimulation of *LGR* during the MHW, with no effects at +6 °C, while in contrast in summer the +2 °C and +4 °C treatments showed no effect and there was a reduction of *LGR* at + 6 °C. The increase of the *New-Old leaf* ratio during the MHW could compensate for the increased *LLR* during MHW in summer. Hence, *N. noltei* appears very well adapted to the moderate heatwaves both in spring and summer; in spring it can even tolerate an extreme (+ 6°C) MHW while in summer it is more vulnerable to such an extreme (+ 6°C) MHW. *R. cirrhosa* was the only species that showed significant mortality during these experiments and occasionally 34 % of the shoots died in one of the experimental units. On average 48 % of the mortality occurred during a MHW treatment period. Hence MHW could have negative impacts on this species inducing increased mortality, although it is more likely that the experimental conditions were not favourable for certain shoots of this species. In contrast, the remaining shoots of *R. cirrhosa* appeared very well adapted to MHW, as both during spring and summer there was hardly any effect on *LGR*. It's worth noting that *R. cirrhosa* generally exhibited lower growth rates than the other species, which may be partially attributed to its narrower leaves resulting in less surface area per leaf. Similar to the other species, *R. cirrhosa* showed an increased the *New-Old leaf* ratio during MHW, indicating a shift towards initiation of growth of new leaves.

Influence of thermal tolerance, seasonal cycles and link with biogeographical distributions

Our study indicates that the three species differed with respect to their sensitivity towards MHW in the following order of increasing vulnerability, i.e. *C. nodosa, R. cirrhosa, N. noltei*. Such differences could be related to lethal thermal thresholds, seasonal patterns and is linked to their biogeographic distribution along latitudinal gradients. The lethal thermal thresholds vary considerably between these three species: up to 40°C for *C. nodosa* according to Bennett et al., (2022), up to 38°C for *R. cirrhosa* according to Tsioli et al., (2019), as well as for *N. noltei* (Massa et al., 2009). This thermal hierarchy largely explains their contrasting responses to the MHW simulated in our study, where *C. nodosa* maintained its growth despite occasional peaks of up to 44°C in summer. MHW do not affect species uniformly across the seasons, but interact with their intrinsic life cycles. Hence, seasonal variations in the response of seagrass to different intensities of simulated marine heatwaves are linked to the natural cycle of this species within the Mediterranean lagoon. In spring, which represents the active growth period for *N. noltei*, growth was stimulated by moderate marine heatwaves (+2°C and +4°C), while extreme marine heatwaves (+6°C) had no effect. This response is in line with the observations of Lee et al., (2007), who demonstrated a positive correlation between temperature and growth in spring, but a negative one in summer. Conversely, *C. nodosa* maintains its growth during spring and summer. *R. cirrhosa* follows a different seasonal pattern, with peak development in late spring and summer, followed by a marked reduction in autumn and virtual disappearance in winter (Mannino & Graziano, 2016).

This difference could explain why *R. cirrhosa*'s growth was hardly impacted during MHWs in
summer, simply corresponding to its natural development peak. While occurring ubiquitously in
coastal environments, the precise biogeographic distribution is not known (Green & Short, 2003).
Both *C. nodosa* and *N. noltei* are Mediterranean species that also occur in the North Atlantic along
the coasts of Western Europe and West Africa. However, the distribution of *N. noltei* extends further
north than that of *C. nodosa*, while the southern distribution limit of *C. nodosa* is located further
south than that of *N. noltei*. Also at the regional scale, *C. nodosa* is widely distributed in Corsican
lagoons along the Tyrrhenian Sea (Garrido et al., 2013) and uncommon in the lagoons along the
colder Gulf of Lion (Le Fur et al., 2018). Hence, the biogeographic distribution of both species is
coherent with their different susceptibilities towards MHW.

Ecological plasticity of the three species

The three species studied show varying degrees of ecological plasticity, reflecting their
respective evolutionary histories and ecological niches. *C. nodosa* stands out for its remarkable
plasticity, which enables it not only to tolerate, but sometimes to benefit from high temperatures.
This characteristic, combined with its pronounced annual cycle (Agostini et al., 2003; Cancemi et
al., 2002), gives it an adaptive advantage in highly variable Mediterranean lagoon environments.
As pointed out by Egea et al., (2018), *C. nodosa* may even benefit from future climate scenarios,
although this increase in production may be accompanied by a depletion of carbon reserves,
making it potentially vulnerable to additional stresses. *R. cirrhosa* reveals a different environmental
specialization, favoring hyperhaline environments (Verhoeven, 1975) rather than the
Mediterranean Sea itself. This adaptation to extreme conditions may explain its remarkable
tolerance to MHW, as our study detected no significant change in its growth or oxygen flux under
the experimental conditions. This resilience corroborates the observations of Rasmusson et al.,
(2021) and suggests that *R. cirrhosa* could maintain its viability in the face of global warming.

It should, however, be highlighted that the abovementioned study by Rasmusson et al., (2021)
focused on *R. cirrhosa* in coastal environments along the Baltic Sea in Sweden. This raises the
question, whether ecotypes of these species do exist that are genetically adapted to their thermal
environments. For instance, one study compared individuals from the Northern Hemisphere
(considered cold-adapted) to those from the Southern Hemisphere (considered warm-adapted)
and found that the heatwave stimulated the growth of individuals from cold environments while
inhibiting the growth of those from warm environments (Nguyen et al., 2021). However, another
study compared two Spanish populations of *C. nodosa* from the Mediterranean Sea, separated by
700 km. Due to contrasting summer temperature regimes, one population is considered warm-
adapted and the other cold-adapted. The authors found that the cold-environment population was
more sensitive to warming, whereas the warm-environment population, for example, increased its
growth during the heatwave (Marin-Guirao et al., 2018). Significant interpopulation variability in
thermal responses has also been observed for southern populations of *N. noltei* (Massa et al.,
2011), which showed a moderate response to heat, while northern populations underwent more
marked changes, mainly linked to down-regulation of genes during heat shock.

Phenological responses to MHW

The systematic increase in the *New-Old leaf* ratio observed in all three species, albeit less
pronounced in *R. cirrhosa*, suggests their common phenological response to MHWs. This shift
likely reflects an acceleration of natural leaf turnover in response to thermal stress, particularly in
C. nodosa, a fast-growing opportunistic species where MHW exposure intensifies tissue renewal.
The production of new leaves may also involve structural adaptations to heat stress, such as
changes in leaf anatomy. For instance, studies on other seagrass species have shown that high
temperatures can increase mesophyll tissue thickness (Purnama et al., 2018), a response that, in
terrestrial plants, is associated with enhanced plasma membrane permeability—an adaptation that
improves gas exchange and heat dissipation while limiting exposure to thermal stress. Additionally,

modifications in leaf architecture, particularly the production of smaller leaves, may further optimize gas exchange and heat dissipation, reducing the impact of heat stress. The weaker response observed in *R. cirrhosa* could be attributed to its intrinsic morphology, as its naturally thin leaves require fewer structural adjustments. In *Posidonia oceanica*, young leaves under heat stress halt their growth to redirect energy toward protective mechanisms, notably through the activation of genes such as HSP90, AOX, and BI, which help prevent cellular damage and programmed cell death, thus preserving essential tissues for future survival (Ruocco et al., 2019). This response, combined with an increased production of new leaves, is likely linked to the expression of heat shock proteins (HSP), which enhance thermal tolerance and protect critical cellular functions. Moreover, the activation of photoprotective mechanisms involving heat-responsive genes helps maintain photosynthetic and respiratory activity (Beca-Carretero et al., 2018, 2018; Marin-Guirao et al., 2016). These morpho-physiological adjustments, combined with the species' ecological strategies—*Ruppia* as a colonizing species, *Nanozostera* as a colonizing/opportunistic species, and *Cymodocea* as an opportunistic species—suggest that species with short life cycles and high regeneration potential exhibit greater resilience to heat stress episodes.

Post-heatwave responses

Our study also included the post-heatwave period (15 days), as MHW effects often emerge after initial stress. While MHWs can reduce biomass by limiting leaf growth (Deguette et al., 2022), temperature fluctuations in our near-natural system led to a +4°C heatwave in late spring, preventing us from assessing full recovery. Cumulative heat events can negatively affect growth even if individual MHWs do not (Saha et al., 2020), raising concerns as MHWs become more frequent. Initial observations suggest *C. nodosa* can exploit moderate MHWs for growth, maintaining resilience unless extreme heat persists. In contrast, *N. noltei* showed limited capacity to sustain growth under repeated stress, especially in summer, while *R. cirrhosa* remained stable, demonstrating strong thermal resilience.

These findings highlight species-specific adaptation strategies. *C. nodosa* benefits from moderate warming but struggles under prolonged heat, while *N. noltei*'s increased sensitivity suggests a lower resilience threshold, potentially impacting its competitiveness in future climates. *R. cirrhosa*'s ability to maintain stable growth despite repeated stress reinforces its potential for persistence in fluctuating environments. As MHW frequency increases, a key question arises: how will seagrasses respond to multiple consecutive events in short periods, and what are the long-term consequences for ecosystem stability?

Conclusions and perspectives

Our study investigates for the first time the effect of MHW intensity in spring and summer on three key structuring species in a Mediterranean lagoon, it is very interesting that in our incubation systems, all three species were able to cope with the experimental MHW of 20 days in spring and summer, and only for *N. noltei* a clearly negative impact on growth was observed for the highest intensity (+ 6 °C) in summer. This study supports the importance of considering the provenance of the shoots used in experiments and confirms that organisms inhabiting these lagoons have developed robust survival strategies to withstand more extreme and variable conditions than marine species in more stable ecosystems. With intensification of ECEs, MHWs will become more frequent, our study investigates the effect of one MHWs with different intensities. However, the after period show us the importance of designing studies on the effect of cumulative MHWs on the resilience of seagrass. This is even more relevant for Mediterranean lagoon that are naturally stressed environment.

Acknowledgements

The authors sincerely thank Nicolas Cimiterra for his support in plant collection, as well as Jean-Yves Coail and Patrick Berriet for their invaluable technical assistance with the creation of the mesocosm system. They also extend their gratitude to Elodie Foucault and Gregory Messiaen for nutrient analysis with special thanks to Gregory Messiaen for his significant contribution to instrumentation. Finally, they thank Elise Lacoste for her help with statistical analyses.

Author contributions

C.B, RdW, L.S and VO. conceived the study and designed the experimental approach; C.B and E.D/R performed the laboratory analyses; C.B, RdW, L.S, V.O analysed the data and performed the statistical analyses; C.B, RdW writing-original draft preparation; RdW, L.S, V.O writing-review and editing. VO funding acquisition. All authors reviewed the manuscript.

Funding

This study is part of C.B.'s doctoral research and was funded by the Agence de l'Eau Rhône-Méditerranée-Corse and Ifremer.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

Data and script are available online on bioRxiv : https://doi.org/10.1101/2025.03.26.645255 ; (Bourdier et al., 2025).

REFERENCES

Agostini, S., Pergent, G., & Marchand, B. (2003). Growth and primary production of Cymodocea nodosa in a coastal lagoon. *Aquatic Botany*, *76*(3), 185-193. https://doi.org/10.1016/S0304-3770(03)00049-4

Beca-Carretero, P., Guiheneuf, F., Marin-Guirao, L., Bernardeau-Esteller, J., Garcia-Munoz, R., Stengel, D. B., & Ruiz, J. M. (2018). Effects of an experimental heat wave on fatty acid composition in two Mediterranean seagrass species. In *Marine Pollution Bulletin* (Vol. 134, Numéro SI, p. 27-37). PERGAMON-ELSEVIER SCIENCE LTD. https://doi.org/10.1016/j.marpolbul.2017.12.057

Bennett, S., Vaquer-Sunyer, R., Jorda, G., Forteza, M., Roca, G., & Marba, N. (2022). Thermal Performance of Seaweeds and Seagrasses Across a Regional Climate Gradient. In *Frontiers in Marine Science* (Vol. 9). FRONTIERS MEDIA SA. https://doi.org/10.3389/fmars.2022.733315

Bourdier, C., De Wit, R., Soissons, L., De Ronne, E., & Ouisse, V. (2025). *Experimental approach to assessing the impact of the intensity of near-natural heatwaves on three marine phanerogam species in Mediterranean lagoon environment*. Ecology. https://doi.org/10.1101/2025.03.26.645255

Cancemi, G., Buia, M. C., & Mazzella, L. (2002). Structure and growth dynamics of *Cymodocea nodosa* meadows. *Scientia Marina*, *66*(4), 365-373. https://doi.org/10.3989/scimar.2002.66n4365

Cardoso, P. G., Raffaelli, D., & Pardal, M. A. (2008). The impact of extreme weather events on the seagrass Zostera noltii and related Hydrobia ulvae population. In *Marine Pollution Bulletin* (Vol.

56, Numéro 3, p. 483-492). PERGAMON-ELSEVIER SCIENCE LTD. https://doi.org/10.1016/j.marpolbul.2007.11.006

Colvin, T., & Snelgrove, P. (2025). Role of seagrass physical structure in macrofaunal biodiversity-ecosystem functioning relationships. *Marine Ecology Progress Series*, *754*, 35-49. https://doi.org/10.3354/meps14774

Costa, M. M., Silva, J., Barrote, I., & Santos, R. (2021). Heatwave Effects on the Photosynthesis and Antioxidant Activity of the Seagrass Cymodocea nodosa under Contrasting Light Regimes. *Oceans*, *2*(3), 448-460. https://doi.org/10.3390/oceans2030025

Cramer, W., Guiot, J., Fader, M., Garrabou, J., Gattuso, J.-P., Iglesias, A., Lange, M. A., Lionello, P., Llasat, M. C., Paz, S., Peñuelas, J., Snoussi, M., Toreti, A., Tsimplis, M. N., & Xoplaki, E. (2018). Climate change and interconnected risks to sustainable development in the Mediterranean. *Nature Climate Change*, *8*(11), 972-980. https://doi.org/10.1038/s41558-018-0299-2

Darmaraki, S., Somot, S., Sevault, F., Nabat, P., Cabos Narvaez, W. D., Cavicchia, L., Djurdjevic, V., Li, L., Sannino, G., & Sein, D. V. (2019). Future evolution of Marine Heatwaves in the Mediterranean Sea. *Climate Dynamics*, *53*(3-4), 1371-1392. https://doi.org/10.1007/s00382-019-04661-z

David, M., Bailly-Comte, V., Munaron, D., Fiandrino, A., & Stieglitz, T. C. (2019). Groundwater discharge to coastal streams – A significant pathway for nitrogen inputs to a hypertrophic Mediterranean coastal lagoon. *Science of The Total Environment*, *677*, 142-155. https://doi.org/10.1016/j.scitotenv.2019.04.233

Deguette, A., Barrote, I., & Silva, J. (2022). Physiological and morphological effects of a marine heatwave on the seagrass Cymodocea nodosa. *Scientific Reports*, *12*(1). https://doi.org/10.1038/s41598-022-12102-x

de Wit, R. (2011). Biodiversity of Coastal Lagoon Ecosystems and Their Vulnerability to Global Change. In O. Grillo (Éd.), *Ecosystems Biodiversity*. InTech. https://doi.org/10.5772/24995

Diffenbaugh, N. S., Pal, J. S., Giorgi, F., & Gao, X. (2007). Heat stress intensification in the Mediterranean climate change hotspot. *Geophysical Research Letters*, *34*(11), 2007GL030000. https://doi.org/10.1029/2007GL030000

Duarte, C. M. (2002). The future of seagrass meadows. *Environmental Conservation*, *29*(2), 192-206. https://doi.org/10.1017/S0376892902000127

Egea, L. G., Jiménez-Ramos, R., Vergara, J. J., Hernández, I., & Brun, F. G. (2018). Interactive effect of temperature, acidification and ammonium enrichment on the seagrass Cymodocea nodosa. *Marine Pollution Bulletin*, *134*, 14-26. https://doi.org/10.1016/j.marpolbul.2018.02.029

Fiandrino, A., Ouisse, V., Dumas, F., Lagarde, F., Pete, R., Malet, N., Le Noc, S., & De Wit, R. (2017). Spatial patterns in coastal lagoons related to the hydrodynamics of seawater intrusion. *Marine Pollution Bulletin*, *119*(1), 132-144. https://doi.org/10.1016/j.marpolbul.2017.03.006

Fourqurean, J. W., Duarte, C. M., Kennedy, H., Marba, N., Holmer, M., Angel Mateo, M., Apostolaki, E. T., Kendrick, G. A., Krause-Jensen, D., McGlathery, K. J., & Serrano, O. (2012). Seagrass ecosystems as a globally significant carbon stock. In *Nature Geoscience* (Vol. 5, Numéro 7, p. 505-509). NATURE PORTFOLIO. https://doi.org/10.1038/ngeo1477

Garrabou, J., Coma, R., Bensoussan, N., Bally, M., Chevaldonné, P., Cigliano, M., Diaz, D., Harmelin, J. G., Gambi, M. C., Kersting, D. K., Ledoux, J. B., Lejeusne, C., Linares, C., Marschal, C., Pérez, T., Ribes, M., Romano, J. C., Serrano, E., Teixido, N., … Cerrano, C. (2009). Mass mortality in Northwestern Mediterranean rocky benthic communities : Effects of the 2003 heat wave. *Global Change Biology*, *15*(5), 1090-1103. https://doi.org/10.1111/j.1365-2486.2008.01823.x

Garrido, M., Lafabrie, C., Torre, F., Fernandez, C., & Pasqualini, V. (2013). Resilience and stability of Cymodocea nodosa seagrass meadows over the last four decades in a Mediterranean lagoon. *Estuarine, Coastal and Shelf Science*, *130*, 89-98. https://doi.org/10.1016/j.ecss.2013.05.035

Giorgi, F. (2006). Climate change hot-spots. *Geophysical Research Letters*, *33*(8), 2006GL025734. https://doi.org/10.1029/2006GL025734

Giorgi, F., & Lionello, P. (2008). Climate change projections for the Mediterranean region. *Global*
and Planetary Change, *63*(2-3), 90-104. https://doi.org/10.1016/j.gloplacha.2007.09.005
Green, E. P., & Short, F. T. (2003). World Atlas of Seagrasses. *(Eds.)*, 324.
Hobday, A. J., Alexander, L. V., Perkins, S. E., Smale, D. A., Straub, S. C., Oliver, E. C. J.,
Benthuysen, J. A., Burrows, M. T., Donat, M. G., Feng, M., Holbrook, N. J., Moore, P. J.,
Scannell, H. A., Sen Gupta, A., & Wernberg, T. (2016). A hierarchical approach to defining
marine heatwaves. *Progress in Oceanography*, *141*, 227-238.
https://doi.org/10.1016/j.pocean.2015.12.014
Hobday, A. J., Oliver, E. C. J., Sen Gupta, A., Benthuysen, J. A., Burrows, M. T., Donat, M. G.,
Holbrook, N. J., Moore, P. J., Thomsen, M. S., Wernberg, T., & Smale, D. A. (2018). Categorizing
and naming MARINE HEATWAVES. *Oceanography*, *31*(2, SI), 162-173.
https://doi.org/10.5670/oceanog.2018.205
Iotti, M., Darnaude, A. M., Bouriat, A., & Ouisse, V. (2023). Spatio-temporal Variation of Shallow
Microhabitats and Associated Juvenile Fish Assemblages in a Mediterranean Lagoon.
Estuaries and Coasts, *46*(1), 198-226. https://doi.org/10.1007/s12237-022-01102-9
IPCC. (2021). *Climate Change 2021 – The Physical Science Basis : Working Group I Contribution*
to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change (1ʳᵉ éd.).
Cambridge University Press. https://doi.org/10.1017/9781009157896
IPCC. (2023). *Climate Change 2022 – Impacts, Adaptation and Vulnerability : Working Group II*
Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate
Change (1ʳᵉ éd.). Cambridge University Press. https://doi.org/10.1017/9781009325844
Jentsch, A., & Beierkuhnlein, C. (2008). Research frontiers in climate change : Effects of extreme
meteorological events on ecosystems. *Comptes Rendus. Géoscience*, *340*(9-10), 621-628.
https://doi.org/10.1016/j.crte.2008.07.002
Jentsch, A., Kreyling, J., & Beierkuhnlein, C. (2007). A new generation of climate-change
experiments : Events, not trends. *Frontiers in Ecology and the Environment*, *5*(7), 365-374.
https://doi.org/10.1890/1540-9295(2007)5[365:ANGOCE]2.0.CO;2
Jordà, G., Marbà, N., & Duarte, C. M. (2012). Mediterranean seagrass vulnerable to regional
climate warming. *Nature Climate Change*, *2*(11), 821-824. https://doi.org/10.1038/nclimate1533
Kim, M., Qin, L.-Z., Kim, S. H., Song, H.-J., Kim, Y. K., & Lee, K.-S. (2020). Influence of Water
Temperature Anomalies on the Growth of Zostera marina Plants Held Under High and Low
Irradiance Levels. *Estuaries and Coasts*, *43*(3), 463-476. https://doi.org/10.1007/s12237-019-
00578-2
Le Fur, I., De Wit, R., Plus, M., Oheix, J., Simier, M., & Ouisse, V. (2018). Submerged benthic
macrophytes in Mediterranean lagoons : Distribution patterns in relation to water chemistry and
depth. *Hydrobiologia*, *808*(1), 175-200. https://doi.org/10.1007/s10750-017-3421-y
Lee, K.-S., Park, S. R., & Kim, Y. K. (2007). Effects of irradiance, temperature, and nutrients on
growth dynamics of seagrasses : A review. *Journal of Experimental Marine Biology and*
Ecology, *350*(1-2), 144-175. https://doi.org/10.1016/j.jembe.2007.06.016
Lefcheck, J. S., Wilcox, D. J., Murphy, R. R., Marion, S. R., & Orth, R. J. (2017). Multiple stressors
threaten the imperiled coastal foundation species eelgrass (*Zostera marina*) in Chesapeake
Bay, USA. *Global Change Biology*, *23*(9), 3474-3483. https://doi.org/10.1111/gcb.13623
Lejeusne, C., Chevaldonné, P., Pergent-Martini, C., Boudouresque, C. F., & Pérez, T. (2010).
Climate change effects on a miniature ocean : The highly diverse, highly impacted
Mediterranean Sea. *Trends in Ecology & Evolution*, *25*(4), 250-260.
https://doi.org/10.1016/j.tree.2009.10.009
Lloret, J., Marin, A., & Marin-Guirao, L. (2008). Is coastal lagoon eutrophication likely to be
aggravated by global climate change? *Estuarine Coastal and Shelf Science*, *78*(2), 403-412.
https://doi.org/10.1016/j.ecss.2008.01.003
Mannino, A. M., & Graziano, M. (2016). Differences in the growth cycle of *Ruppia cirrhosa*
(Petagna) Grande in a Mediterranean shallow system. *Plant Biosystems - An International*
Journal Dealing with All Aspects of Plant Biology, *150*(1), 54-61.
https://doi.org/10.1080/11263504.2014.906511

Mannino, A. M., Menéndez, M., Obrador, B., Sfriso, A., & Triest, L. (2015). The genus Ruppia L.
(Ruppiaceae) in the Mediterranean region : An overview. *Aquatic Botany*, *124*, 1-9.
https://doi.org/10.1016/j.aquabot.2015.02.005
Marin-Guirao, L., Bernardeau-Esteller, J., Garcia-Munoz, R., Ramos, A., Ontoria, Y., Romero, J.,
Perez, M., Ruiz, J. M., & Procaccini, G. (2018). Carbon economy of Mediterranean seagrasses
in response to thermal stress. In *Marine Pollution Bulletin* (Vol. 135, p. 617-629). PERGAMON-
ELSEVIER SCIENCE LTD. https://doi.org/10.1016/j.marpolbul.2018.07.050
Marin-Guirao, L., Ruiz, J. M., Dattolo, E., Garcia-Munoz, R., & Procaccini, G. (2016). Physiological
and molecular evidence of differential short-term heat tolerance in Mediterranean seagrasses.
In *Scientific Reports* (Vol. 6). NATURE PORTFOLIO. https://doi.org/10.1038/srep28615
Mascaró, O., Oliva, S., Pérez, M., & Romero, J. (2009). Spatial variability in ecological attributes
of the seagrass *Cymodocea nodosa*. *Botm*, *52*(5), 429-438.
https://doi.org/10.1515/BOT.2009.055
Massa, S. I., Arnaud-Haond, S., Pearson, G. A., & Serrao, E. A. (2009). Temperature tolerance
and survival of intertidal populations of the seagrass Zostera noltii (Hornemann) in Southern
Europe (Ria Formosa, Portugal). In *Hydrobiologia* (Vol. 619, p. 195-201). SPRINGER.
https://doi.org/10.1007/s10750-008-9609-4
Massa, S. I., Pearson, G. A., Aires, T., Kube, M., Olsen, J. L., Reinhardt, R., Serrao, E. A., &
Arnaud-Haond, S. (2011). Expressed sequence tags from heat-shocked seagrass Zostera noltii
(Hornemann) from its southern distribution range. In *Marine Genomics* (Vol. 4, Numéro 3, p.
181-188). ELSEVIER. https://doi.org/10.1016/j.margen.2011.04.003
Nguyen, H. M., Bulleri, F., Marin-Guirao, L., Pernice, M., & Procaccini, G. (2021). Photo-physiology
and morphology reveal divergent warming responses in northern and southern hemisphere
seagrasses. In *Marine Biology* (Vol. 168, Numéro 8). SPRINGER HEIDELBERG.
https://doi.org/10.1007/s00227-021-03940-w
Nixon, S. W., Granger, S., Buckley, B. A., Lamont, M., & Rowell, B. (2004). A one hundred and
seventeen year coastal water temperature record from Woods Hole, Massachusetts. *Estuaries*,
27(3), 397-404. https://doi.org/10.1007/BF02803532
Orth, R. J., Carruthers, T. J. B., Dennison, W. C., Duarte, C. M., Fourqurean, J. W., Heck, K. L.,
Hughes, A. R., Kendrick, G. A., Kenworthy, W. J., Olyarnik, S., Short, F. T., Waycott, M., &
Williams, S. L. (2006). A Global Crisis for Seagrass Ecosystems. *BioScience*, *56*(12), 987.
https://doi.org/10.1641/0006-3568(2006)56[987:AGCFSE]2.0.CO;2
Ouisse, V., Miramont, A., Bajjouk, T., Cimiterra, N., Fontes, H., Giraud, A., & Vaz, S. (2023).
Description méthodologique et analyse de la distribution des habitats en milieu lagunaire
méditerrannéen. Projet CHAMILA – Partie 1. 64p.
Papaki, D., Kokkos, N., & Sylaios, G. (2020). A Fuzzy Inference System for Seagrass Distribution
Modeling in the Mediterranean Sea : A First Approach. In *Water* (Vol. 12, Numéro 10). MDPI.
https://doi.org/10.3390/w12102949
Pérez-Ruzafa, A., Campillo, S., Fernández-Palacios, J. M., García-Lacunza, A., García-Oliva, M.,
Ibañez, H., Navarro-Martínez, P. C., Pérez-Marcos, M., Pérez-Ruzafa, I. M., Quispe-Becerra,
J. I., Sala-Mirete, A., Sánchez, O., & Marcos, C. (2019). Long-Term Dynamic in Nutrients,
Chlorophyll a, and Water Quality Parameters in a Coastal Lagoon During a Process of
Eutrophication for Decades, a Sudden Break and a Relatively Rapid Recovery. *Frontiers in*
Marine Science, *6*, 26. https://doi.org/10.3389/fmars.2019.00026
Pérez-Ruzafa, A., Pérez-Ruzafa, I. M., Newton, A., & Marcos, C. (2019). Coastal Lagoons :
Environmental Variability, Ecosystem Complexity, and Goods and Services Uniformity. In
Coasts and Estuaries (p. 253-276). Elsevier. https://doi.org/10.1016/B978-0-12-814003-
1.00015-0
Petton, S., Pouvreau, S., Fleury, E., & Lagarde, F. (2024). *ECOSCOPA network : High frequency*
environmental database (SEANOE, Éd.). https://doi.org/10.17882/86131
Purnama, P. R., Purnama, E. R., Manuhara, Y. S. W., Hariyanto, S., & Purnobasuki, H. (2018).
Effect of high temperature stress on changes in morphology, anatomy and chlorophyll content
in tropical seagrass Thalassia hemprichii. *11*(6).

Rasmusson, L. M., Nualla-ong, A., Wutiruk, T., Bjork, M., Gullstrom, M., & Buapet, P. (2021). Sensitivity of Photosynthesis to Warming in Two Similar Species of the Aquatic Angiosperm Ruppia from Tropical and Temperate Habitats. *Sustainability*, *13*(16). https://doi.org/10.3390/su13169433

REPHY network. (2021). *REPHY - French Observation and Monitoring program for Phytoplankton and Hydrology in coastal waters (2023)* (Dataset No. 10.17882/47248). SEANOE. https://doi.org/10.17882/47248

Repolho, T., Duarte, B., Dionisio, G., Paula, J. R., Lopes, A. R., Rosa, I. C., Grilo, T. F., Cacador, I., Calado, R., & Rosa, R. (2017). Seagrass ecophysiological performance under ocean warming and acidification. In *Scientific Reports* (Vol. 7). NATURE PUBLISHING GROUP. https://doi.org/10.1038/srep41443

Rueda, J. L., Salas, C., & Marina, P. (2008). Seasonal variation in a deep subtidal *Zostera marina* L. bed in southern Spain (western Mediterranean Sea). *Botm*, *51*(2), 92-102. https://doi.org/10.1515/BOT.2008.016

Ruocco, M., De Luca, P., Marín-Guirao, L., & Procaccini, G. (2019). Differential Leaf Age-Dependent Thermal Plasticity in the Keystone Seagrass Posidonia oceanica. *Frontiers in Plant Science*, *10*, 1556. https://doi.org/10.3389/fpls.2019.01556

Saha, M., Barboza, F. R., Somerfield, P. J., Al-Janabi, B., Beck, M., Brakel, J., Ito, M., Pansch, C., Nascimento-Schulze, J. C., Jakobsson Thor, S., Weinberger, F., & Sawall, Y. (2020). Response of foundation macrophytes to near-natural simulated marine heatwaves. *Global Change Biology*, *26*(2), 417-430. https://doi.org/10.1111/gcb.14801

Savva, I., Bennett, S., Roca, G., Jorda, G., & Marba, N. (2018). Thermal tolerance of Mediterranean marine macrophytes : Vulnerability to global warming. In *Ecology and Evolution* (Vol. 8, Numéro 23, p. 12032-12043). WILEY. https://doi.org/10.1002/ece3.4663

Shields, E. C. C., Parrish, D., & Moore, K. A. A. (2019). Short-Term Temperature Stress Results in Seagrass Community Shift in a Temperate Estuary. *Estuaries and Coasts*, *42*(3), 755-764. https://doi.org/10.1007/s12237-019-00517-1

Short, F., Carruthers, T., Dennison, W., & Waycott, M. (2007). Global seagrass distribution and diversity : A bioregional model. *Journal of Experimental Marine Biology and Ecology*, *350*(1-2), 3-20. https://doi.org/10.1016/j.jembe.2007.06.012

Simão, S. A. (2021). *Resilience of dwarf seagrass meadows to simulated marine heat waves* [MS thesis].

Smale, D. A., Wernberg, T., Oliver, E. C. J., Thomsen, M., Harvey, B. P., Straub, S. C., Burrows, M. T., Alexander, V, L., Benthuysen, J. A., Donat, M. G., Feng, M., Hobday, A. J., Holbrook, N. J., Perkins-Kirkpatrick, S. E., Scannell, H. A., Sen Gupta, A., Payne, B. L., & Moore, P. J. (2019). Marine heatwaves threaten global biodiversity and the provision of ecosystem services. *NATURE CLIMATE CHANGE*, *9*(4), 306+. https://doi.org/10.1038/s41558-019-0412-1

Strydom, S., Murray, K., Wilson, S., Huntley, B., Rule, M., Heithaus, M., Bessey, C., Kendrick, G. A., Burkholder, D., Fraser, M. W., & Zdunic, K. (2020). Too hot to handle : Unprecedented seagrass death driven by marine heatwave in a World Heritage Area. *Global Change Biology*, *26*(6), 3525-3538. https://doi.org/10.1111/gcb.15065

Sullivan, B. K., & Short, F. T. (2023). Taxonomic revisions in Zosteraceae (Zostera, Nanozostera, Heterozostera and Phyllospadix). *Aquatic Botany*, *187*, 103636. https://doi.org/10.1016/j.aquabot.2023.103636

Templado, J. (2014). Future Trends of Mediterranean Biodiversity. In S. Goffredo & Z. Dubinsky (Éds.), *The Mediterranean Sea* (p. 479-498). Springer Netherlands. https://doi.org/10.1007/978-94-007-6704-1_28

Tomlinson, P. B., & Posluzny, U. (2001). Generic limits in the seagrass family *Zosteraceae. TAXON*, *50*(2), 429-437. https://doi.org/10.2307/1223890

Tournois, J., Darnaude, A. M., Ferraton, F., Aliaume, C., Mercier, L., & McKenzie, D. J. (2017). Lagoon nurseries make a major contribution to adult populations of a highly prized coastal fish. *Limnology and Oceanography*, *62*(3), 1219-1233. https://doi.org/10.1002/lno.10496

Trombetta, T., Vidussi, F., Mas, S., Parin, D., Simier, M., & Mostajir, B. (2019). Water temperature
drives phytoplankton blooms in coastal waters. *Plos One*, *14*(4).
https://doi.org/10.1371/journal.pone.0214933

Tsioli, S., Orfanidis, S., Papathanasiou, V., Katsaros, C., & Exadactylos, A. (2019). Effects of
salinity and temperature on the performance of Cymodocea nodosa and Ruppia cirrhosa : A
medium-term laboratory study. In *Botanica Marina* (Vol. 62, Numéro 2, p. 97-108). WALTER DE
GRUYTER GMBH. https://doi.org/10.1515/bot-2017-0125

Verhoeven, J. T. A. (1975). Ruppia-communities in the camargue, France. Distribution and
structure in relation to salinity and salinity fluctuations. *Aquatic Botany*, *1*, 217-241.
https://doi.org/10.1016/0304-3770(75)90024-8

Winters, G., Nelle, P., Fricke, B., Rauch, G., & Reusch, T. (2011). Effects of a simulated heat wave
on photophysiology and gene expression of high- and low-latitude populations of Zostera
marina. *Marine Ecology Progress Series*, *435*, 83-95. https://doi.org/10.3354/meps09213

Cymodocea nodosa

Ruppia cirrhosa