AlphaFold3 and RoseTTAFold All-Atom structures enable radiosensitizers discovery by targeting multiple DNA damage repair proteins

Junjie Tong[1,2], Xixi Wang[1,3], Miaoshan Lu[1], Tong Wang[1], Jingying Chen[1], Jingjing Chen[1], Yaohan Li[1], Cong Xie[1,2], Yanhui Fu[2], Changbin Yu[1,2, *].

1 Central Hospital Affiliated to Shandong First Medical University, Jinan 250000 Shandong Province, China.

2 Key Laboratory of Tropical Medicinal Plant Chemistry of Ministry of Education, College of Chemistry and Chemical Engineering, Hainan Normal University, Haikou 571158, Hainan Province, China

3 Xi'an Jiaotong-Liverpool University, Suzhou 215000, Jiangsu Province, China

∗ Corresponding author

Email: yu_lab@sdfmu.edu.cn

Abstract

Radioresistance remains a primary obstacle in tumor radiotherapy, yet no radiosensitizers have been approved for clinical use due to toxicity concerns. To identify effective and safe radiosensitizers, a natural products (NP) database comprising 79,263 compounds were docked against a hybrid target library of four DNA damage repair (DDR)-related proteins, including both experimental and AI-predicted structures generated by AlphaFold3 and RoseTTAFold All-Atom models. It demonstrated that hit compounds identified by AI-modeled structures accounted for 80% of all hits in ataxia telangiectasia mutated (ATM), 88.89% in ATM- and Rad3-related (ATR), 92.31% in DNA-dependent protein kinase catalytic subunit (DNA-PKcs), and 70% in Poly (ADP-ribose) polymerase 1 (PARP1), expanding compounds diversity recognized by experimental structures. Seven promising multi-target radiosensitizer candidates were identified, among which two compounds were validated as effective radiosensitizers in vitro. Detailed proteomic analyses revealed that both compounds activate TP53-associated apoptosis and senescence as cellular endpoints by modulating the synergistic interplay between DDR and spindle checkpoints. The integration of AI-predicted and experimental structures highlighted the structural complementarity advantages of this approach for small-scale compound libraries, where limited computational resources are required. This study provided new radiosensitizers and strategies to overcome tumor radioresistance in future research.

Key words: AF3, RFAA, Natural products, DNA damage repair, Radiosensitizers

Introduction

Radiotherapy (RT) is a versatile and essential modality in the management of non-small cell lung cancer (NSCLC). However, consecutive exposures to ionizing radiation (IR) can lead to the incidence of radioresistance, resulting in the failure of RT and poor prognosis[1]. DNA damage is the primary mechanism by which IR exerts its cytotoxic effects on tumor cells. Accordingly, DNA damage repair (DDR) pathways are the main factors contributing to the development of radioresistance[2]. DDR kinases, such as ataxia telangiectasia mutated (ATM), ATM- and Rad3-related (ATR), DNA-dependent protein kinase catalytic subunit (DNA-PKcs), and the nuclear enzyme of Poly (ADP-ribose) polymerase 1 (PARP1), are principal regulators among these pathways[3, 4]. Therefore, inhibitors of ATM, ATR, DNA-PKcs, and PARP-1 have become promising therapeutic agents for overcoming tumor radioresistance[1]. Besides, since the genomic instability in cancer cells, synthetic lethality has emerged as a promising therapeutic strategy[5]. Several studies demonstrated that synthetic lethal interactions are widely prevalent among these four DDR targets[6-9]. Collectively, inhibitors targeting these four regulators not only act as potent radiosensitizers in RT but also induce cancer cell death through synthetic lethal interactions. However, no radiosensitizer has yet been approved for clinical use due to its toxicities[10]. Thus, there is an urgent need to develop effective and safe novel radiosensitizers.

Natural products (NP) serve as a major source of lead compounds in drug discovery, with rich structural diversity, high bioactivity, and considerable safety. To date, several NP-sourced radiosensitizers are being under investigation in clinical trials, including papaverine, curcumin, genistein, paclitaxel, and resveratrol[11]. However, discovering bioactive compounds from NPs is still challenging, since the identification of their biological targets involves randomness. Structure-based virtual screening (SBVS) has emerged as a powerful tool to address this challenge. It aims to enrich ligands formed favorable interactions with target proteins by screening compounds libraries using docking tools, such as AutoDock Vina[12], QVina2[13], and DOCK3.8[14].

The success of SBVS consists of two key components, the structure of targets and the library of compounds. Traditionally, experimental and homology modeled structures are widely used in SBVS[15]. However, for many targets, experimental structures are either unavailable or have low resolution, while homology models depend heavily on sequence similarity to templates. Recent advancements in artificial intelligence (AI) have revolutionized protein structure prediction. Following the emergence of AlphaFold2 (AF2)[16]and RoseTTAFold (RF)[17] models, numerous retrospective studies were conducted to assess the performance of AI-predicted structures in SBVS[18, 19]. These studies concluded that unprocessed AI-predicted structures were not ideal for predicting target-ligand conformations. However, more recently, a study reported that unrefined AF2 structures retrospectively failed to reproduce ligands screened by experimental structures, but prospectively identified novel high-affinity compounds that differ from experimental ones[20]. These findings supported the notion that bias arises from the use of single-source protein structure, since co-crystallographic ligands may influence the protein conformations, especially in binding pockets, thereby restricting the diversity of hit compounds in SBVS. In 2024, the developers of AF2 and RF models, recipients of the Nobel Prize in Chemistry, introduced their latest models, AlphaFold3 (AF3)[21] and RoseTTAFold All-Atom (RFAA)[22]. These models can simulate assemblies of proteins and other biomolecular systems, showing great promise for drug discovery applications. Nevertheless, it remains to be seen whether structures generated by these latest models will deliver enhanced performance in SBVS applications.

Meanwhile, a high-quality library of compounds is critical for SBVS. Recent studies identified numerous potential inhibitors and agonists with high binding affinities for various targets from the ultra-large ZINC library[23-25], containing over 1.3 billion purchasable compounds. The ultra-large library, with diverse chemotypes, can improve docking results by identifying more hit compounds with significant affinities[26, 27]. However, the expansion of libraries typically incurs exponentially greater computational and temporal costs. For instance, screening 1.2 billion compounds may require over 380,000 core-hours on a computational cluster exceeding 2,000 CPU cores[26], which is often impractical for most laboratories. Consequently, many research groups or pharmaceutical companies preferred small and specialized databases, such as libraries of NP (most contain fewer than 100,000

compounds)[28], marketed drugs (around three to four thousands)[29], or even in-house compound collections (normally less than 10,000 compounds). The limited size of these libraries restrict their utility in SBVS[27]. AI-predicted structures could discover bioactive compounds with novel structural features compared to those identified through experimental structures[20]. Building on this, we hypothesized that integrating AI-predicted structures with experimental ones for SBVS could expand the list of hit compounds for a given protein. This hybrid screening strategy could be especially valuable when working with small-size libraries.

We assembled a hybrid target library containing experimental, AF3-predicted, and RFAA-predicted structures of four key DNA damage repair regulators: ATM, ATR, DNA-PKcs, and PARP1. We evaluated the performance of these structures by retrospectively docking them against known ligands and prospectively docking the NP subset of ZINC20 database[29]. For the top-ranked ligands, we applied several filters to check their druglikeness, toxicity, and novelty. Ligands identified by multiple proteins were considered as potential multi-target compounds. Furthermore, we validated them computationally and selected two compounds to test their radiosensitization mechanisms by cell assays and proteomics analysis.

Result

Retrospective docking for experimental and AI-modeled structures

The experimentally determined structures of ATM (PDB ID: 7NI4, 3.00 Å), ATR (PDB ID: 5YZ0, 4.70 Å), DNA-PKcs (PDB ID: 7OTY, 2.96 Å), and PARP1 (PDB ID: 7AAC, 1.59 Å) were retrieved from the Protein Data Bank (PDB). Structures with superior resolutions and co-bound inhibitors were preferred. Only one experimental structure of ATR has been published to date, and it lacks a co-bound inhibitor. We focused on their primary binding domains: the kinase domains (KD) for ATM, ATR, and DNA-PKcs, and the catalytic domain (CAT) for PARP1. We first evaluated the structural differences between AI-predicted and experimental structures in terms of global structures (Fig.1) and local binding pockets (Figs.1), i.e., residues within 5Å distance from the

co-bound ligand. Three AI models (ATM-AF3, PARP1-AF3, and PARP1-RFAA) demonstrated high structural similarity to their experimental counterparts, with global structures and most local residues had root-mean-square deviation (RMSD)<2Å. In contrast, ATR-AF3, ATR-RFAA, and ATM-RFAA had global RMSD exceeding 4Å and more residues with RMSD over 2Å. The variations of ATR structures were likely generated from the poor resolution of its PDB structure, while the ATM PDB structure was excluded from the RFAA training data. Additionally, the absence of DNA-PKcs in the training data of both models likely contributed to the differences observed for this protein.

Fig. 1: The global differences between experimental and AI-predicted structures for ATM, ATR, DNA-PKcs, and PARP1. Structures were either downloaded from PDB or generated from AF3 and RFAA models. It demonstrated the global RMSD between experimental and AI-modeled structures.

We retrospectively docked ATM, ATR, DNA-PKcs, and PARP1 against two

types of known ligands: (1) the co-bound inhibitors from experimental assemblies; (2) the known inhibitors retrieved from ChEMBL database[30]. We first redocked M4076[31] back into the ATM experimental structure (RMSD=0.12Å) and observed that the interactions were highly consistent with the original assembly (Figs.2A). We then docked M4076 to ATM_AF3 and ATM_RFAA models, with RMSD<1Å and relatively lower affinities (-8.4 $kcal/mol$ for AF3 and -7.1 $kcal/mol$ for RFAA) than the redocked assembly (-9.5 $kcal/mol$) (Figs.2B). For the ATR protein, VX-970[32] was docked to the experimental structure with RMSD of 1.96Å; however, it was well predicted in both the AF3 (RMSD: 1.54Å) and RFAA (RMSD: 1.52Å) models (Figs.3); For the rest two proteins, M3814[33] and veliparib[34] were redocked to DNA-PKcs (Figs.4) and PARP1 (Figs.5) structures, respectively, which showed accurate prediction of ligands conformations with RMSD less than 2Å in all assemblies.

Fig. 2: Enrichment analysis of actives against decoys for ATM, ATR, DNA-PKcs, and PARP1. (A) Four structural sources of ATM (Experimental, AF3, AF2, RFAA) were used for retrospective docking screening. The AUC values were used to define the general ability of structures to enrich actives from decoys, while the $logAUC$ values described enrichment at early stages. Here, we identified the actives were recognized at top 10% (Gray area) of docked results. The $logAUC$ represented the difference between the model's semilogarthmic AUC and the random semilogarthmic AUC (dashed lines). (B) The AUC and $logAUC$ values among different ATR structures. (C) The AUC and $logAUC$ values among different DNA-PKcs structures. (D) The AUC and $logAUC$ values among different PARP1 structures.

We prepared known active ligands and their property-matched decoys for each target. We docked the active-decoy databases against four structural sources (AF3, RFAA, AF2, and experimental) for each protein (Fig.2). The area under curve (AUC) was calculated to evaluate the overall performance in

distinguishing actives from decoys, while the $logAUC$ was explored to check their enrichments at the early stages, i.e., 10% top-ranked molecules. For ATM protein (Fig.2A), the experimental ($logAUC$=0.48), RFAA ($logAUC$=0.44), AF2 ($logAUC$=0.49) models showed comparable enrichment, which was higher than the AF3 model ($logAUC$=0.31). In contrast, for the ATR protein (Fig.2B), the experimental structure performed the worst with a $logAUC$=0.21, while the AF3 ($logAUC$=0.55) and AF2 ($logAUC$=0.55) models successfully ranked the active compounds. For DNA-PKcs (Fig. 2C), the AF3 model ($logAUC$ = 0.03) exhibited significantly lower enrichment than other models. We checked the binding pocket of the DNA-PKcs_AF3 model and observed a smaller pocket than the experimental structure (Figs.1). Furthermore, we calculated the detail parameters of their binding pockets showed that AF3 had the smallest pocket volume (768Å^3) and the lowest depth (18.73Å), but the highest surface-to-volume ratio (SVR) of 1.34 (Figs.6). These results suggested that the AF3 structure may require different compounds to achieve effective binding. For PARP1 (Fig.2D), three structures (AF3, experimental, and AF2) had close $logAUC$ values ranged 0.377-0.395, while the RFAA returned a relative lower $logAUC$ of 0.272.

Prospective docking against the Natural Products database

We prospectively docked 12 hybrid protein structures from ATM, ATR, DNA-PKcs and PARP1, against the ZINC20 NP subset, with 79,263 secondary metabolites and its derivatives. We first checked the overlap rate between AI models and experimental structure as the size of top-ranked molecules gradually increased (step size=500) (Figs.7). Among the top-ranked 1000 compounds (around 1% of the NP database), ATM_AF3 (48.3%), PARP1_AF3 (40.8%), and PARP1_RFAA (37.3%) models exhibited high intersection ratios with their experimental models. As mentioned earlier, these three AI-predicted structures exhibited the lowest RMSD values (<2Å) (Fig.1), indicating high similarity to the experimental structures. In contrast, the ATR_AF3 model had the least overlap rate of 18.9%. Other models had comparable ratios, ranging from 22.6% to 28.5%.

We used the affinity score of the 1000th-ranked compound as the cutoff, selecting approximately 1,000 compounds for subsequent evaluation and screening (Table1). Top-ranked molecules were clustered using Morgan fingerprints, with Tanimoto coefficient (Tc) = 0.5, resulting in clusters numbers

ranged from 154 (ATM_RFAA model) to 277 (ATR_AF3 model). Molecules with the highest affinity for each cluster were kept as representative compounds. These representatives were further filtered using stringent druggability and toxicity predictions as described in the Methods section. By focusing on AI-predicted structures, we identified eight hit compounds for ATM (Figs.8), eight for ATR (Figs.9), twelve for DNA-PKcs (Figs.10), and seven for PARP1 (Figs.11). After merging compounds recognized by their corresponding experimental structures (Table 1), the list of hit compounds expanded: to ten compounds for ATM, to nine compounds for ATR, to thirteen compounds for DNA-PKcs, to ten compounds for PARP1. This result reflected a low overlap of hit compounds between AI-predicted and experimental structures. To identify compounds with novel chemotypes, we removed molecules with a Tc value > 0.35 compared to known ligands in the ChEMBL database. We then identified seven potential multi-target compounds: ZINC000001123725 ('3725), ZINC000003797417 ('7417), ZINC000000602850 ('2850), ZINC000000952701 ('2701), ZINC000004085325 ('5325), ZINC000000621966 ('1966), ZINC000004260672 ('0672), by calculating the intersections of four proteins' candidate lists (Figs.12). These multi-target compounds formed 24 assemblies with different protein structures.

Table 1. Changes in the number of compounds in target structures with different filtering criteria. Top-ranked compounds were clustered using their morgan fingerprints, with Tanimoto coefficient (Tc) = 0.5. The compound with the highest docking score was selected as representative for the following druggability, toxicity, and novelty screening. These filtered compounds were regarded as hit compounds. Hit compounds identified by different sourced structures were merged as total hit compounds. Ultimately, seven promising multi-target compounds identified by simultaneously in at least two targets were selected.

Target Structures		Top-ranked compounds	Represent compounds	Filtered Hit compounds	Total Hit compounds	Multi-target compounds
ATM	Experimental	1176	250	3		
	AF3	1090	270	7	10	
	RFAA	1008	154	1		
ATR	Experimental	1075	211	3		
	AF3	1162	277	2	9	
	RFAA	1109	264	6		

DNA-PKcs	Experimental	1250	196	2	7
	AF3	1066	237	8	13
	RFAA	1142	184	4	
PARP 1	Experimental	1058	217	7	
	AF3	1143	267	4	10
	RFAA	1249	275	5	

Molecular dynamics and MMPBSA calculations

We conducted 100 nanoseconds(ns) molecular dynamics simulations (MDS) and the binding free energy (BFE) calculations for all 24 protein-ligand assemblies. We initially performed 100ns MDS on Apo forms of proteins to set the baselines (Figs.13), and experimental assemblies were considered as positive controls. Subsequently, root-mean-square deviation (RMSD) analyses were conducted: RMSD of the protein backbone was calculated to assess conformational fluctuations, and RMSD of ligands relative to the protein backbone was to evaluate simulation convergence. Additionally, the root-mean-square fluctuation (RMSF) was used to evaluate dynamic characteristics of flexible regions in the proteins.

MDS on ligands screened by ATM protein

We conducted 100ns MDS on five ligands ('2850, '3725, '7417, '2701, and '5325) filtered by ATM structures (Figs.14). M4076 and most compounds had backbone stability comparable to the Apo forms of corresponding ATM structures; however, '3725 exhibited a higher backbone RMSD when bound to ATM_AF3. Regarding the residues RMSF, significant flexibility was observed in the C-terminal, N-terminal, and PRD regions of KD depending on ligand binding. During the simulations, M4076 exhibited the most stable binding states, with RMSD below 0.4nm, while other ligands varied between 0.4 and 0.7nm.

MDS on ligands screened by DNA-PKcs protein

The DNA-PKcs models identified five ligands ('1966, '0672, '3725, '5325, and '7417) (Figs.15), of which only '1966 was bound to the experimental and AF3

models in parallel. Compared with the Apo form, compounds '1966 and M3814 can stabilize the experimental structure during the last 40ns simulation, maintaining a more compact or ordered state. For the RMSF values, M3814 demonstrated a more flexible region around the P-loop (residues 3729-3735) than the Apo form. The movement of P-loop was recognized as a key conformational change when DNA-PKcs bounded with inhibitors[35]. Meanwhile, '1966 displayed high flexibility around the P-loop region and the comparable ligand stability similar to those observed with M3814.

MDS on ligands screened by PARP1 protein

There were five ligands ('0672, '1966, '2701, '3725, and '7417) screened by three PARP1 models (Figs.16). These ligands kept protein backbones at a stable state throughout the simulations. Concerning the protein residues, the Apo and Holo forms of the PARP1 demonstrated significant flexibility throughout the entire CAT domain. When analyzing the RMSD of ligands relative to the protein backbone, veliparib, serving as the reference inhibitor, demonstrated the most stable relative displacement, with an RMSD below 0.1nm. Other ligands showed greater variability, with RMSD values ranging from 0.2 to 0.5nm.

MDS on ligands screened by ATR protein

The ATR protein identified only two compounds ('2850 and '3725) via its AF3 model (Figs.17). The '3725 maintained a backbone RMSD similar to its Apo state, whereas compound '2850 increased the instability of the protein structure. Regarding RMSF, the flexibility of specific regions was not significantly affected by whether the protein was bound to a ligand or not. Finally, the RMSD of both ligands approached a similar equilibrium state at the end of simulations.

MMPBSA calculations

We used the Molecular mechanics/Poisson–Boltzmann surface area (MM/PBSA) model to calculate the BFE for all complexes and three experimental assemblies, i.e., ATM-M4076 (-28.35 $\pm$ 6.7 $kcal/mol$), DNA-PKcs-M3814 (-30.46 $\pm$ 4.5 $kcal/mol$), and PARP1-Veliparib (-46.88 $\pm$ 5.0 $kcal/mol$). Four compounds ('3725, '7417, '5325, and '1966) formed six assemblies, exhibiting improved BFE compared to those experimentally co-bound inhibitors, labeled as references (Table 2). We proceeded to select two compounds ('1966 and '7417) for further experimental

investigation based on their availability for follow-up studies. The '1966 showed better BFE than DNA-PKcs-M3814, '1966-DNA-PKcs_PDB ($-46.02\pm9.4kcal/mol$) and '1966-DNA-PKcs_AF3 ($-43.03\pm13.2kcal/mol$. For the '7417, it formed assembly with DNA-PKcs_RFAA, with a BFE of $-31.35\pm6.6kcal/mol$.

Table 2. The binding free energy and molecular docking affinity of protein-ligand assemblies. Three reference assemblies were downloaded from PDB database and set as the threshold. Seven potential multi-target compounds formed 24 assemblies with different protein structures.

NO.	Ligand ID	Protein Name	Structure Source	Binding Free Energy ($kcal/mol$)	Molecular Docking ($kcal/mol$)
ref	M4076	**ATM**	**PDB**	-28.35 ± 6.7	**-9.5**
ref	M3814	**DNA-PKcs**	**PDB**	-30.46 ± 4.5	**-10.4**
ref	Veliparib	**PARP1**	**PDB**	-46.88 ± 5.0	**-9.1**
1	'2850	ATM	PDB	-10.52 ± 8.9	-10.2
2	'2850	ATR	AF3	-24.26 ± 5.6	-10.9
3	**'3725**	**ATM**	**PDB**	-33.96 ± 4.4	**-10.3**
4	'3725	ATM	AF3	-19.30 ± 5.4	-10.6
5	'3725	ATR	AF3	-27.67 ± 5.9	-11
6	'3725	DNA-PKcs	AF3	-21.37 ± 4.8	-10.5
7	'3725	PARP1	PDB	-25.81 ± 8.1	-13.6
8	'3725	PARP1	AF3	-38.70 ± 6.8	-13.2
9	'3725	PARP1	RFAA	-37.97 ± 6.2	-12.7
10	'7417	ATM	PDB	-15.45 ± 5.0	-9.0
11	**'7417**	**DNA-PKcs**	**RFAA**	-31.35 ± 6.6	**-10**
12	'7417	PARP1	PDB	-36.85 ± 7.6	-11.6
13	'2701	ATM	AF3	-21.80 ± 5.4	-10.8
14	'2701	PARP1	PDB	-21.36 ± 8.8	-12.6

15	'2701	PARP1	AF3	-20.30±8.0	-11.8
16	'2701	PARP1	RFAA	-22.64±7.2	-11.6
17	'5325	**ATM**	**RFAA**	**-34.16±12.1**	**-8.7**
18	'5325	**DNA-PKcs**	**AF3**	**-48.31±7.6**	**-9.2**
19	'1966	**DNA-PKcs**	**PDB**	**-46.02±9.4**	**-9.2**
20	'1966	**DNA-PKcs**	**AF3**	**-43.03±13.2**	**-9.2**
21	'1966	PARP1	PDB	-18.02±7.7	-13.2
22	'1966	PARP1	RFAA	-12.46±8.2	-12.4
23	'0672	DNA-PKcs	AF3	-14.75±4.0	-9.4
24	'0672	PARP1	PDB	-16.42±5.5	-12.4

Assessing radiosensitivity properties of '1966 and '7417 compounds

We selected '1966 and '7417 as potential multi-target inhibitors and radiosensitizers. The '1966 is a tyrosine derivative featuring a protecting group. It is primarily used as a synthetic intermediate for the synthesis of bioactive peptidomimetics[36]. While the '7417 compound is a derivative developed and designed from the NP asperlicin[37], demonstrating as a non-peptide gastrin antagonist targeting brain cholecystokinin receptor (CCK-B)[38]. To date, no studies have reported on the relationship between these two compounds and their potential roles as radiosensitizers.

Fig. 3: Cell assays of compounds 1966 and 7417. (A) A549 cell viability at compounds concentrations of $6.25\,\mu M$, $12.5\,\mu M$, $25\,\mu M$, $50\,\mu M$, and $100\,\mu M$. (B) Colony formation under different conditions. (C) Cell viability under varying radiation doses, i.e., $2Gy$, $4Gy$, $6Gy$, and $8Gy$, for compound-treated and untreated A549 cells. (D) Flow cytometry was used to detect cells apoptosis under different treatment conditions. The treatment conditions were divided into four groups: normal A549 cells (Control), compound-treated cells, radiation-treated cells (IR), and the combination of compound and radiation-treated cells.

We initially assessed the inhibitory effects of '1966 and compound '7417 on A549 cells, revealing that both compounds exhibited dose-dependent suppression of cell viability, with IC50 values exceeding $50\,\mu M$ (Fig.3A). We next investigated their potential as radiosensitizer. To maintain cell viability while preserving the inhibitory effects of the compounds, we used 25 μM for '1966 and 12.5 μM for '7417 in subsequent experiments. Upon the additional

radiation exposures, we checked the cell viability of compound-treated tumor cells versus untreated cells. The combination of compounds treatment and radiation demonstrated significantly enhanced tumor cell inhibition at doses of $2Gy$, $4Gy$, $6Gy$, and $8Gy$ (Fig.3C). Colony formation assays further confirmed that both compounds, when combined with $4Gy$ radiation, exhibited superior clonogenic inhibition compared to other groups (Fig.3B). Additionally, flow cytometry assay demonstrated a significant increase in both early and late apoptotic cells in the compound-plus-radiation treatment group (Fig.3D).

Proteomics of '1966

To investigate the mechanistic responses of tumor cells to '1966, we employed both Data-dependent Acquisition (DDA) and Data-Independent Acquisition (DIA) proteomics across four groups: (1) Untreated tumor cells (Label as "Control"), (2) Cells under radiation (Label as "IR"), (3) Cells treated with compound '1966(Label as "ZINC1966"), and (4) Cells treated with the compound and radiation (Label as "ZINC1966IR"). We performed Principal Component Analysis (PCA), which effectively distinguished these groups at the proteome level (Fig.4A). We identified 6,645 proteins in the Control group, 6,307 proteins in the ZINC1966 group, 6,380 proteins in the IR group, and 6,261 proteins in the ZINC1966IR group (Fig.4B). The Venn diagram revealed that 5,695 proteins were shared among all four groups (Fig.4C). To explore the differences of compound '1966 under non-irradiated and irradiated conditions, we extracted the differentially expressed proteins (DEPs), with thresholds of $log2FC$>2, $log2FC$<-2 and p<0.05, between：

- the ZINC1966 and Control groups, identifying 205 upregulated proteins and 224 downregulated proteins (Fig.4D).

- the ZINC1966IR and IR groups, identifying 163 upregulated proteins and 310 downregulated proteins (Fig.4E).

Fig. 4: The proteomics analysis of compound 1966. (A) The principal component analysis of four groups (n=5 per group): Control, ZINC1966, IR, and ZINC1966_IR. (B) The number of detected proteins in four groups. (C) The Veen diagram showed overlaps between these four groups. (D) The differentially expressed proteins (DEPs) between ZINC1966 and Control. DEPs were selected based on $log2FC$>2, $log2FC$<-2 and p<0.05. (E) The DEPs between ZINC1966IR and IR. (F) Enrichment analysis of DEPs between ZINC1966 and Control using the Gene Ontology (GO), KEGG, and Reactome databases, with p<0.05. (G) The protein-protein interactions (PPI) analysis of DEPs between ZINC1966 and Control. These hub genes were identified by the CytoNCA plugin of Cytoscape. (H) Enrichment analysis of hub genes of ZINC1966 vs Control, with p<0.05. (I) Enrichment analysis of DEPs between ZINC1966IR and IR using the GO, KEGG, and Reactome databases, with p<0.05. (G) The PPI analysis of DEPs between ZINC1966IR and IR. (H) Enrichment analysis of hub genes of ZINC1966IR vs IR, with p<0.05.

Next, we performed the enrichment analysis of all DEPs using the Gene Ontology (GO), KEGG, and Reactome databases (Fig.4F). By comparing ZINC1966 against Control, we found that GO-Biological process (BP) was significantly enriched in pathways such as cell division, cell cycle, and double-strand break repair. GO-Cellular component (CC) analysis revealed that the proteins were primarily localized in the nucleoplasm, nucleus, and cytosol. Meanwhile, GO-Molecular function (MF) analysis indicated associations with RNA binding, phosphoprotein binding, and enzyme binding. Further pathway enrichment using KEGG and Reactome identified cell cycle checkpoints, particularly, spindle checkpoints. Subsequently, we conducted Protein-Protein Interaction (PPI) analysis to identify hub genes (Fig.4G), such as CHEK1, EP300, RB1, and MAD2L1. Pathway enrichment of these hub

DEPs confirmed significant associations with RNA polymerase II regulation, cell cycle-related pathways, DNA damage response and repair, cellular senescence, and TP53 signaling (Fig.4H).

In the absence of irradiation, the level of DNA damage in cells is limited. To evaluate mechanisms behind the compound under conditions of amplified DNA damage, we analyzed the enrichment results of DEPs between the ZINC1966IR and IR groups (Fig.4I). For the overall DEPs, GO-BP enrichment revealed more associations with DNA damage response, DNA repair (particularly double-strand break repair), cell cycle regulation, cellular senescence, and apoptosis. GO-CC and GO-MF analyzes indicated that these proteins were primarily localized in the nucleus and sites of cellular damage, with functions closely related to DNA binding. KEGG and Reactome Pathway enrichment analysis identified signaling pathways associated with cellular senescence and programmed cell death as markers of cellular fate, as well as cell cycle checkpoint-related pathways that were downstream of the DNA damage repair. Following PPI analysis (Fig.4J-K), we found that the hub genes remained enriched in pathways related to DNA damage response and repair, cell cycle regulation, TP53-induced apoptosis, and DNA damage-induced cellular senescence.

Based on the PPI analysis of pathways identified by global DEPs and hub genes, we hypothesized that, as a potential radiosensitizer, '1966 interfered with DNA repair pathway, leading to cell cycle arrest. This ultimately led to TP53-induced apoptosis and cellular senescence as the endpoints of cellular fate. Thus, we annotated the DNA damage response, DDR, Cell cycle checkpoints, Cellular apoptosis, and Cellular senescence pathways by GO_BP, KEGG, Reactome enrichment results of DEPs (Fig.5). Under unirradiated conditions (Fig.5 left panel), we observed the enrichment of DNA damage response-related proteins, such as H2AX ($log2FC$=-7.74), a core protein whose downregulation was previously shown to be associated with increased radiaosensitivity[39]. In DNA repair, by targeting DNA-PKcs and PARP1, part of double-strand and single-strand repair pathways were suppressed, alternative pathways were selectively activated, such as ERCC1($log2FC$=3.53), which participated in nucleotide excision repair (NER), and MLH1 ($log2FC$=2.34), which was involved in mismatch repair (MMR). The cell cycle checkpoints were primarily enriched in spindle checkpoints. With persistent damage or irreparable DNA damage, the TP53-induced apoptosis

and cellular senescence may be triggered. Upon irradiated conditions (Fig.5 right panel), we observed significantly enhanced involvement of the DNA damage response mechanisms, with the activation of alternative DNA repair pathways such as the ATM ($log2FC$ =2.25)-mediated pathway, ERCC1 ($log2FC$ =4.64)-associated NER, and FANCI ($log2FC$ =2.38)-mediated interstrand crosslink (ICL) repair. Similarly, a greater number of cell cycle checkpoints, TP53, apoptosis-, and senescence-related proteins were enriched.

Fig. 5: The diagram of the mechanism of compound '1966 under irradiated and non-irradiated conditions. Interactions of these pathways were identified by EnrichmentMap plugin in Cytoscape. Five pathways were focused on DNA damage response, DNA damage repair, Cell cycle checkpoints, Cell apoptosis and cellular senescence. DEPs were categorized into these pathways and annotated based on $log2FC$ values. On the left panel, DEPs between ZINC1966 and Control were annotated. On the right panel, DEPs between ZINC1966IR and IR were annotated.

In summary, our analysis revealed that DDR and cell cycle regulation represented core pathways in both irradiated and unirradiated conditions following treatment with '1966, while cell apoptosis and senescence acted as downstream endpoints of these cellular responses.

Proteomics of '7417

Compound '7417 shared similar targets with '1966, including DNA-PKcs and PARP1, and the additional target ATM protein. PCA demonstrated significant differences among four groups: Control, IR, ZINC7417, and ZINC7417IR (Fig.6A). We identified 6,418 proteins in the Control group, 6,362 proteins in the IR group, 6,166 proteins in the ZINC7417 group, and 6,109 proteins in the

ZINC7417IR group, with 5,571 proteins shared among all groups (Fig.6B-C). In the DEPs analysis, ZINC7417 exhibited 177 upregulated and 248 downregulated proteins compared to the Control group (Fig.6D). The DEPs between ZINC7417IR and IR groups, we identified 177 upregulated and 213 downregulated proteins (Fig.6E).

Fig. 6: The proteomics analysis of compound 7417. (A) The principal component analysis of four groups (n=5 per group): Control, ZINC7417, IR, and ZINC7417_IR. (B) The number of detected proteins in four groups. (C) The Veen diagram showed overlaps between these four groups. (D) The DEPs between ZINC7417 and Control. DEPs were selected based on $log2FC$>2, $log2FC$<-2 and p<0.05. (E) The DEPs between ZINC7417IR and IR. (F) Enrichment analysis of DEPs between ZINC7417 and Control using the GO, KEGG, and Reactome databases, with p < 0.05. (G) PPI analysis of DEPs between ZINC7417 and Control. (H) Enrichment analysis of hub genes of ZINC7417 vs Control, with p < 0.05. (I) Enrichment analysis of DEPs between ZINC7417IR and IR using the GO, KEGG, and Reactome databases, with p < 0.05. (G) The PPI analysis of DEPs between ZINC7417IR and IR. (H) Enrichment analysis of hub genes of ZINC7417IR vs IR, with p < 0.05.

In the GO enrichment analysis of DEPs between ZINC7417 and Control (Fig.6F), the BP category included cell cycle pathways and checkpoints, as well as DNA damage response and repair processes. The CC analysis revealed that the DEPs were primarily localized in the cytosol, nucleoplasm, and nucleus. Meanwhile, the MF analysis indicated that the DEPs were predominantly associated with various kinase activities, among which DNA-PKcs and ATM belong to the Ser/Thr protein kinase family. Pathway

enrichment analysis using KEGG and Reactome identified significant enrichment in cell cycle-related pathways, cellular senescence, autophagy, TP53 pathway, and colorectal cancer-related pathways. PPI analysis identified hub genes related to DDR (Fig.6G), such as ATM, CHEK1, and MTOR. Further pathway enrichment of these hub genes focused on DNA damage response and checkpoint regulation, double-strand break repair, cell cycle checkpoints, cellular senescence, apoptosis, and autophagy (Fig.6H). Additionally, the gastric cancer pathway was identified, since '7417, a known CCK-B antagonist, had been associated with gastric cancer and colorectal cancer[37, 38].

Upon additional radiation, we conducted enrichment analysis on the DEPs between ZINC7417IR and IR (Fig.6I). In the GO-BP category, the DEPs were enriched in various DNA damage response and repair-related pathways, as well as chromatin remodeling. GO-CC and GO-MF analysis revealed that the DEPs were primarily localized in the nucleus, nucleoplasm, and cytosol, with functions mainly related to RNA binding, DNA binding, and ATP binding activity. Pathway enrichment using KEGG and Reactome demonstrated that DEPs were involved in cellular senescence, cell cycle-related pathways, DNA repair, and Fanconi anemia pathway. Subsequently, we identified hub genes and performed enrichment analysis on them (Fig.6J-K). This revealed enrichment in cell cycle, TP53 activity, cellular senescence, DNA damage response and additional DNA repair pathways, such as NER, ICL repair, mainly Fanconi anemia pathway. Of note, we observed that core enriched pathways were associated with gastric cancer under unirradiated conditions; however, under radiation conditions, we did not observe the enrichment of gastric cancer-related pathways.

Since '7417 shared similar targets to '1966, we hypothesized that they exhibit similar mechanisms of action, specifically by regulating DNA repair and cell cycle checkpoints through the DNA damage response, ultimately leading to cell apoptosis and senescence. We next illustrated the mechanism of action for '7417(Fig.7). Without radiation (Fig.7 left panel), we observed a significant downregulation of ATM ($log2FC$=-2.01) protein, the direct target of '7417, accompanied by reduced expression of other proteins associated with DSBs, such as XRCC4 ($log2FC$=-3.43). Notably, XRCC4 functions downstream of DNA-PKcs[4]. Consequently, DEPs were significantly enriched in the TP53 pathway and apoptosis and senescence-related pathways. Following radiation exposure (Fig.7 right panel), we did not observe activation of the gastric

cancer pathway; however, there were further increases in proteins associated with DNA damage response and repair. Notably, specific repair pathways were activated, such as the Fanconi anemia pathway, including ERCC4 ($log2FC$ =2.47), FANCD2 ($log2FC$ =3.79), and FAAP100 ($log2FC$ =4.53). Additionally, we identified downregulation of numerous proteins involved in negative regulation of apoptotic processes, such as ERCC5 ($log2FC$=-2.63), CBX4 ($log2FC$=-2.04), BABAM2 ($log2FC$=-2.18), MALT1 ($log2FC$=-3.63), and PLK1 ($log2FC$=-3.04).

Fig. 7: The diagram of the mechanism of compound '7417 under irradiated and non-irradiated conditions. Interactions of these pathways were identified by EnrichmentMap plugin in Cytoscape. Pathways of Gastrin cancer, DNA damage response, DNA damage repair, Cell cycle checkpoints, Cell apoptosis, and cellular senescence were focused. DEPs were categorized into these pathways and annotated based on $log2FC$ values. On the left panel, DEPs between ZINC7417 and Control were annotated in six pathways. On the right panel, DEPs between ZINC7417IR and IR were annotated in five pathways, without the Gastrin cancer.

In summary, the above proteomic analysis demonstrated the potential of both compounds '1966 and '7417 as tumor radiosensitizers. Although they are distinct compounds, they exhibit similarities in their mechanisms of action at the proteomic level based on their shared targets.

Discussion

AI models like AF2, AF3, and RFAA now enable structure prediction for SBVS. While ultra-large libraries (millions to billions of compounds) can improve hit rates, smaller databases (e.g., market drugs, NP, or in-house isolated

molecules, typically under 100,000 compounds) often limit SBVS efficiency. Recent work shows AF2-predicted structures identify novel hits not found by experimental structures[20]. We therefore proposed that combining experimental and AI-predicted structures could enhance the diversity of hit compounds. To validate this approach, we selected the NP subset from ZINC20, with 79,263 molecules, and performed SBVS against four DDR-associated proteins: ATM, ATR, DNA-PKcs, and PARP1. These targets are crucial for tumor radiosensitizers discovery, and their synergistic lethal mechanisms represent a promising strategy in cancer therapy[6-9, 40]. We first retrospectively docked co-bound inhibitors and active-decoys libraries against the integrated targets library. Unlike previous studies[18-20], experimental structures were not superior to AI models in AUC and $logAUC$ values in all cases (Fig.2). Notably, ATR structures modeled by AI conducted superior performance in enriching active inhibitors from decoys (Fig.2B), implying AI modeled structures may be closer to the authentic ATR protein conformations.

In the prospective study, compounds identified by AI-predicted structures represented a large subset of hit compounds for each target (Table 1), with eight of total ten compounds for ATM (AI hit ratio=80%), eight of total nine compounds for ATR (AI hit ratio=88.89%), twelve of total thirteen compounds for DNA-PKcs (AI hit ratio=92.31%), and seven of total ten compounds for PARP1 (AI hit ratio=70%). Moreover, the number of compounds identified simultaneously by AI-predicted and experimental structures was kept small: one for ATM (Figs.8), two for ATR (Figs.9), one for DNA-PKcs (Figs.10), and four for PARP1 (Figs.11). This suggested that relying exclusively on experimental structures may restrict the discovery of hit compounds. It is recognized that co-bound inhibitors can induce conformational changes in protein binding pockets[20]. Since protein structures were normally treated as rigid entities in SBVS, the structural diversity of compounds screened by SBVS would be constrained within a defined range. Therefore, compounds screened using a single protein structure source exhibited specific structural biases in SBVS. While more AI-sourced structures could complement the limitations of small-scale compound libraries. We then selected two compounds, '1966 and '7417, based on their identification across multiple targets and lower BFE compared to co-bound inhibitors. Both compounds exhibited dose-dependent cytotoxic effects in cellular assays, likely due to synergistic lethal effects from multi-target engagement. Furthermore, both compounds demonstrated

potential as radiosensitizers, as cell viability decreased further with additional radiation exposure (Fig.3C). These findings provide preliminary evidence supporting the radiosensitizing potential of '1966 and '7417.

Furthermore, we aimed to validate the underlying mechanisms of action for '1966 and '7417, which could provide critical insights for future development of compounds. Proteomics analysis identified hundreds of DEPs in each drug-treated condition. PPI network analysis revealed hub genes. Pathway enrichment analysis of DEPs and hub genes showed both compounds were strongly linked to DNA damage response, DDR pathways, cell cycle regulation (notably at checkpoints), along with apoptosis and senescence, key outcomes of failed DNA repair. These results suggested that '1966 and '7417 shared similar mechanisms due to overlapping targets. Furthermore, these pathways were consistently enriched regardless of radiation exposure. However, radiation noticeably increased the number of DEPs within pathways, since more severe DNA damage was induced, rendering cells more dependent on DNA repair-related pathways. Notably, mechanistic analysis revealed '7417, a CCK-B antagonist, was involved in gastric cancer-associated pathways without radiation, but under radiation conditions, those pathways were not enriched. Instead, upstream and downstream DDR pathways became significantly enriched, supporting its potential synergistic effect with radiation.

Overall, we assessed the SBVS capabilities of AI-predicted and experimental structures in retrospective and prospective studies. Ultimately, we identified radiosensitizers and multi-target compounds from a small-scale NP database and validated their radiosensitizing effects at both cellular and proteomic levels. Nevertheless, several questions remain to be addressed, such as whether this strategy remains effective in local laboratory databases containing fewer than 10,000 compounds. While the proteomic analysis of compounds '1966 and '7417 demonstrated their impact on DNA repair pathways, in vivo efficacy and general adaptability across different tumor types have not been fully elucidated. To address these issues, future testing on laboratory-isolated NP databases will be essential. Additionally, more detailed evaluations of tumor radiosensitizers with promising prospects across different tumor types and in vivo efficacy are required.

Method

Preparation and assessment of four targets structures

The experimental structures of ATM (PDB ID: 7NI4), ATR (PDB ID: 5YZ0), DNA-PKcs (PDB ID: 7OTY), and PARP1 (PDB ID: 7AAC) were retrieved from the Protein Data Bank (PDB). They formed assemblies with known inhibitors, ATM-M4076, DNA-PKcs-M3814, PARP1-veliparib, except for the ATR which has no bounded inhibitors. We separated the protein structures from ligands and retained the kinase domains (KD) for ATM, ATR, DNA-PKcs, and the catalytic domain (CAT) for PARP1 via PyMol (v2.5.5). The amino acids sequences of key domains of these four proteins were acquired from the UniProt database (ATM ID: Q13315, ATR ID: Q13535, DNA-PKcs ID: P78527, PARP1 ID: P09874). The AF3 structures were generated from the online web tool, AlphaFold Server (https://alphafoldserver.com/). While the RFAA model (https://github.com/baker-laboratory/RoseTTAFold-All-Atom) was deployed on our local servers to predict the protein structures. The root-mean-square deviation (RMSD) of global structures and local residues within 5Å around the binding ligands were visualized and calculated via PyMol and in-house scripts.

Retrospective molecular docking against known ligands

We initially redocked and docked these co-bound inhibitors to the mixed target library, including 12 experimental and AI-predicted structures of four targets. The molecular docking was performed by QVina2 (https://github.com/QVina/qvina), setting the exhaustiveness value at 12 for ATM and PARP1, at 24 for ATR and DNA-PKcs. Interactions between ligands and targets were further analyzed by PLIP[41].

We prepared the known active ligands from ChEMBL databases[30], with inclusion criteria of affinities (IC50, EC50, Ki, Kd) less than 1µM and molecular weight less than 600 Da. Based on these active ligands, we generated property-matched decoys using the tldr web-service[42]. Thus, we constructed the active-decoys library for target proteins, including 16,972 ligands for ATM, 14,316 ligands for ATR, and 35,540 ligands for DNA-PKcs. Of note, for the PARP1, we retrieved its active-decoys library using the previously prepared

one in the DUDE-Z database[43], with 30,543 ligands. We evaluated the ability of each structure to distinguish actives from decoys by calculating the Area Under Curve (AUC) values. Furthermore, we utilized log-adjusted AUC ($logAUC$) to demonstrate the capability of enriching actives with higher rankings. The $logAUC$ is defined as:

$$logAUC = logAUC_{model} - logAUC\_rando \tag{1}$$

Where equation (1) $logAUC\_model$ is the area under the semilogarithmic ROC curve for the model predictions. Equation (1) $logAUC\_random$ is the area under the semilogarithmic ROC curve for a random guess.

Prospective molecular docking against the Natural Products Database

In the discovery stage, 79,263 molecules acquired from ZINC20 (https://zinc20.docking.org/) natural products (NP) subset. We prospectively docked the database against the mixed target library of ATM, ATR, DNA-PKcs, and PARP1 using QVina2. We used the affinity score of the 1000th-ranked compound as the cutoff, selecting approximately 1,000 compounds for each structure. These top-ranked compounds were clustered using the 1,024-length binary Morgan fingerprint with a Tanimoto coefficient (Tc)=0.5. Molecules with the highest affinity value in each cluster were retained as the representative compound. The druglikeness, medicinal chemistry property, and pharmacokinetics of representative compounds were predicted using the SwissADME[44]. Five rules from Lipinski's Rule of Five, Veber's Rule, Ghose Filter, Muegge Filter, and Egan Filter were applied to check all representative compounds that achieved five "YES" were kept. Then, we kept compounds with predicted High Gastrointestinal (GI) absorption. Besides, the pan-assay interference compounds (PAINS) were excluded. We predicted their AMES toxicity and Carcinogenicity using the ADMETlab 3.0[45]. Compounds with values less than 0.3, labeled as AMES negative and non-carcinogens, were predicted to show excellent safety.

For each target protein, we merged candidates identified by both AI-predict and experimental structures. All candidates were further filtered for novelty by calculating their Morgan fingerprints similarity against corresponding known ligands in the ChEMBL database. Compounds with a Tc

value of more than 0.35 were removed from our candidate list.

Molecular Dynamics Simulations

We performed molecular dynamics simulations (MDS) with GROMACS (version 2024.1)[46]. The protein PDB files were repaired and prepared using PDBFixer[47], including adding missing atoms, residues, and addressing improper formatting. The protonation states(pH=7.4) of ionizable residues in proteins were defined by their pKa (negative logarithm of the acid dissociation constant), which were predicted by PROPKA (v3.5.1)[48]. All ligands were prepared for topology files generation by adding hydrogens(pH=7.4).

The topologies of proteins were prepared by the 'pdb2gmx' function via the CHARMM36m all-atom force filed[49], and all ligands topology files were generated through the web server of CHARMM General Force Field (CGenFF)[50]. We then rebuilt the protein-ligand complex for following simulations. The complex conformations were placed into a cubic shape box with water molecules, where the distance between the solute and the box edge was set at 1.5nm. To minimize the system, the steepest descent algorithm was used. The minimization process terminated until the maximum force under 1000 $kJ\ mol^{-1}nm^{-1}$ or reached the maximum minimization steps of 50,000. During the equilibration simulations, each protein-ligand complex was position restrained and equilibrated with NVT and NPT ensembles for 100ps, respectively. The MDS runs were performed for 100ns in each complex, and energies were saved per 10ps. The temperature and pressure coupling were controlled by the V-rescale[51] and Parrinello-Rahman[52] methods, with a time constant of 0.1ps and 2ps, reference temperature at 300K and pressure at 1bar, respectively. The short-range van der Waals cutoff was set at 1.2nm and calculated under the Verlet cutoff-scheme. The long-range electrostatics were calculated using Particle Mesh Ewald (PME) method[53]. The hydrogen-bonds were constrained via LINCS algorithm[54]. After the production of MDS runs, the trajectories were used to calculate the RMSD of protein backbones and ligands to evaluate the convergence of the simulation system, and root mean square fluctuation (RMSF) of protein residues to check the flexibility of the protein structure.

MMPBSA Calculations

The Molecular mechanics/Poisson–Boltzmann (Generalized-Born) surface area (MM/PB(GB)SA) is a popular method to calculate binding free energies (BFE) of protein-ligand complexes. Here, the MMPBSA model was used for our complexes using gmx_MMPBSA(version 1.6.3)[55]. The BFE is normally calculated as the equation below:

$$\Delta G_{bind} = \Delta H - T\Delta S \tag{1}$$

Where equation (1) ΔH represents the enthalpy of the binding, while $-T\Delta S$ corresponds to the conformational entropy change upon ligand binding. Since we concerned the relative BFE of ligands towards repeated protein structures, the entropy term was dismissed. The terms of enthalpy are given by:

$$\Delta H = \Delta E_{MM} + \Delta G_{SOLV} \tag{2}$$

Where equation (2) the gas phase contributions of ΔE_{MM} and the solvation energy of ΔG_{SOLV} can be further decomposed into multiple terms:

$$\Delta E_{MM} = \left(\Delta E_{bond} + \Delta E_{angle} + \Delta E_{dihedral}\right) + \left(\Delta E_{ele} + \Delta E_{vdW}\right) \tag{3}$$

$$\Delta G_{SOLV} = \Delta G_{EPB} + \Delta G_{ENPOLAR}$$
(4)

The average BFE calculation for each complex was performed using the last 40ns of trajectory.

Cell Culture

A549 cells were purchased from Chinese Academy of Sciences Cell Bank (Identifier: SCSP-503) and grown in RPMI1640 media (Gibco) supplemented with 10% fetal bovine serum (Gibco) and 1% penicillin-streptomycin Solution (Beyotime, China). The cell line was incubated at 37℃ with 5% CO_2 environment (HCP-168, Haier, China).

Chemicals and Reagents

According to the vendor list on the ZINC20, the compound '1966, Fmoc-Tic-OH (ZINC ID:621966, purity≥97%, F117108) was purchased from

Aladdin Scientific (Shanghai, China). While the compound '7417, L-365260(ZINC ID:3797417, purity=99.75%, T22895) was purchased from TargetMol Chemicals. During the cell experiments, two compounds were dissolved in dimethyl sulfoxide (DMSO) at 100mM and diluted to $6.25\,\mu M$, $12.5\mu M$, $25\mu M$, $50\mu M$ and $100\mu M$ by the complex medium.

CCK-8 Assays and Radiation Methods

The cell viability was checked by the CCK-8 assay kit (Beyotime, China). A549 cells were seeded with 5,000 cells per well in the 96-well flat-bottom plates (Beyotime, China) for 24 hours. The medium was then replaced by compounds of '1966 and '7417 diluted at five different concentrations. After incubation with '1966 and '7417 for another 24 hours, the medium was replaced by 100μL medium with 10μL CCK-8 solution and incubated at 37℃ for 1h. The plates were measured at 450nm using a microplate reader (Spark, Tecan, Switzerland). Together, suitable concentrations of '1966 and '7417 were determined by the CCK-8 assays.

To determine the radiation dose for the cell culture, we plated 5,000 cells in each well of the 96-well plates for incubating 24 hours. On the second day, cells were treated by $12.5\mu M$ '7417 and $25\mu M$ '1966 medium, following by the treatment of radiation doses of $2\,Gy$, $4\,Gy$, $6\,Gy$, and $8\,Gy$ using the radiator (RS2000PRO, Rad Source Technologies Inc., USA). After another 24 hours incubation, the medium of radiated cells was replaced by CCK-8 medium. The CCK-8 assay was used to evaluate the radiosensitivity of compounds at different radiation dosages.

Colony formation experiment

The number of 1,000 A549 cells were seeded into each well in 6-well plates. We set up four groups: the control group, the drug-positive group, the IR-positive group, and the drug-IR-positive group. After the cell adhesion was formed, we gave drugs and the $4\,Gy$ dose of IR to corresponding labeled groups for incubating 24 hours. The medium in each well was replaced by fresh complete medium. After 8 days of incubation, cells were washed with PBS for twice, were fixed by 1 mL 4% paraformaldehyde fix solution (Beyotime, China) for 45 min, and stained by crystal violet (Beyotime, China) for 20 min.

Flow Cytometry Assay

The cell apoptosis in four groups were measured using the Annexin V-FITC/Propidium Iodide (PI) Apoptosis Detection Kit (Beyotime, China). After treatment of drugs and IR doses, cells were collected and resuspended with the binding buffer, and stained with FITC and PI following the manufacturer instructions. The proportions of apoptotic cells were counted by the flow cytometry (Attune NxT, Thermo Scientific, USA).

Proteomics Sample Preprocessing

Five bio-replicates were set for cells treated under different conditions. The cell samples were washed by the cold PBS and centrifuged (300g, 5 min, 4℃) for three times. The following steps, including cell lysis, reduction and alkylation, and proteolytic digestion, were processed by the pressure cycling technology (PCT)[56] using the Barocycler 2320EXT (Pressure Biosciences, Inc., USA). After remove the supernatant, $30\,\mu L$ lysis buffer ($8M$ Urea and $2M$ Thiourea, Sigma Aldrich) was added and the sample was transferred into PCT-Micro Tubes (Pressure Biosciences, Inc., USA) for cell lysis (25℃, 60 cycles, 45 kpsi). The reduction and alkylation were conducted by adding 100 mM Tris(2-carboxyethyl) phosphine (TCEP, Sigma Aldrich) and 800mM Iodoacetamide (IAA, Sigma Aldrich) stock. Meanwhile, the standard curve of protein concentration was determined using the Pierce BCA Protein Assay Kit (Thermo Scientific, USA) and Nanodrop (Thermo Scientific, USA). The protein amount in experimental samples was calculated by comparing to the standard curve. Two-steps proteolytic digestion was performed with Lys-C (enzyme-to-substrate ratio of 1:40, Hualishi Scientific) for a PCT condition (33℃, 50 cycles, 45 $kpsi$), and Trypsin (enzyme-to-substrate ratio of 1:50, Hualishi Scientific) for a PCT condition (33℃, 90 cycles, 45$kpsi$). All reagents used in the PCT steps were dissolved and diluted with 100 mM Ammonium bicarbonate (Sigma Aldrich) solution. The digestion reaction was quenched by 20% Trifluoroacetic acid (TFA, Sigma Aldrich) when the pH at 2. The peptide desalting procedure was conducted using the hydrophilic-lipophilic balanced (HLB) cartridges (Oasis, Waters), resulting peptides were in 200 μL elution buffer (70% ACN, Sigma Aldrich). The desalted peptides were dried using the SpeedVac SPD1030 (Thermo Scientific, USA) at 35℃, and redissolved in the Buffer (0.1% formic acid (FA), 4% ACN). Finally, the peptide concentrations

were measured by Nanodrop and adjusted to $0.1mg/mL$.

DDA and DIA Proteomics

A nanoflow DIONEX UltiMate 3000 RSLCnano System connected to an Orbitrap Exploris 480 mass spectrometer equipped with Nanospray Flex ion source and FAIMS Pro Duo system (Thermo Scientific, USA) were used for both data-dependent acquisition (DDA) and data-independent acquisition (DIA) proteomics. Buffer A was H_2O containing 0.1% FA, and Buffer B was ACN containing 0.1% FA. For the DDA mode, $200ng$ peptides were injected into the LC system and separated by a 120min LC gradient with a flow rate of $300nL/min$. The initial gradient was 3% Buffer B and increased to 8% by 3 min. Buffer B was gradually raised to 35% over 103 min, and reached at 90% in the following 5 min. The column was washed by maintaining a high rate of 90% Buffer B for 5 min. The global parameters of mass spectrometer were set as follows: the FAIMS Mode was standard resolution; the Carrier Gas Flow was at $4.8L/min$; two FAIMS CVs were used, both -65V and -45V. During the MS1 phase, the scan range of m/z was from 350 to 1,500 with a resolution of 60,000, the RF lens was set at 50%, the normalized AGC target was 300%, and the maximum injection time was $20ms$. The time between Masters Scans was set at 3 seconds. Precursors for MS/MS experiments had a minimum signal intensity of $2.0e^4$ and charge states from +2 to +6. For MS/MS scans, the isolation window was set at $1.6m/z$, the normalized HCD collision energy was 30%, the resolution was at 15,000, the normalized AGC target was 75%, and the maximum injection time was 22ms. In DIA mode, we used the same LC gradient in DDA proteomics. The parameters for MS1 scans were similar to the DDA experiment, except the scan range was 400-1200m/z. The 64 variable windows were used in the DIA experiment. The MS/MS spectra were acquired with parameters: the resolution was at 30,000, RF lens was at 50%, the normalized HCD collision energy was 32%, the normalized AGC target was 2,000%, and the maximum injection time was 54ms.

Protein identification and quantification

The resultant DIA MS data were processed by FragPipe(version 22.0)[57] using the DIA_SpecLib_Quant workflow, which builds a spectral library via MSFragger-DIA (version 4.1) and quantifies with DIA-NN (version 1.8.2)[58].

The DIA MS data comprised of two FAIMS CVs (-45V and -65V) were separated into two individual files per CV channel using the FreeStyle (version 1.6, Thermo Scientific), following the recommendations of DIA-NN developers. All DIA and DDA MS data were converted into mzML format files by MSConvert (version 3.0)[59] for the FragePipe workflow. Both DDA and DIA mzML files were used for the spectral library generation. The reviewed human protein database downloaded from Uniport on 25 Sep 2024, containing 40,936 entries (includes 20469 decoys: 50%). The spectral library generated by FragPipe was passed into DIA-NN for the DIA quantification. Of note, the file with only one CV channel was regarded as a separate experiment in DIA-NN, so two CVs reports were combined and merged as one file as described by developers on the DIA-NN Github repository (issue 67).

Statistical Analysis

During the proteomics data pre-processing, we first established a threshold for missing data, where any protein with more than 50% missing values across all samples in a group was excluded from further analysis. For the remaining proteins with missing values, these were imputed by replacing them with the median value calculated from the non-missing data points. To reduce systematic variation between samples, we applied quantile normalization upon the data using the normalizeBetweenArrays function from the limma package[60] in R software (v4.3.2). Following normalization, the data were $log2$-transformed to stabilize variance and make the data more normally distributed.

The principal component analysis (PCA) was conducted on the normalized data by Python (v3.9.13). For subsequent differential expression analysis, we then applied linear models and empirical Bayes methods on the data provided by the limma package. Adjusted p-values were calculated using the Benjamini-Hochberg method to control the false discovery rate (FDR), and the fold change were $log2$-transformed ($log2FC$). Differentially expressed proteins (DEPs) were identified based on an adjusted p-value threshold of less than 0.05 and a $log2FC$ >2 or <-2. The enrichment analysis of DEPs was performed using the DAVID online server[61], annotated by the GO terms, KEGG pathways, and Reactome pathways. The sankey diagram with pathway enrichment dots was plotted by an online platform (https://www.bioinformatics.com.cn).

Protein-Protein Interaction Analysis

Protein-protein interaction (PPI) network analysis was performed using the STRING database (v12.0)[62] and Cytoscape software (v3.10.2)[63]. We first uploaded DEPs to the STRING platform, with *Homo sapiens* selected as the organism and a high-confidence interaction score threshold set to ≥0.7. Disconnected nodes were excluded to refine the network. The resulting PPI data were exported and imported into Cytoscape for visualization and topological analysis. To identify hub genes, the CytoNCA plugin was employed to select hub genes based on Betweenness (BC), Closeness (CC), and Degree (DC) centralities.

The EnrichmentMap plugin (v3.5.0)[64] in Cytoscape was employed to visualize functional relationships among enriched terms. DAVID output files, enrichment results of DEPs, were directly imported into EnrichmentMap. The edge cutoff was set at 0.375 and the node cutoff was set at p-value < 0.05. The results from the EnrichmentMap were subsequently utilized to construct mechanism of action diagrams for the studied drugs.

References

1. Buckley, A.M., et al., *Targeting hallmarks of cancer to enhance radiosensitivity in gastrointestinal cancers.* Nat Rev Gastroenterol Hepatol, 2020. **17**(5): p. 298-313.
2. Huang, R.X. and P.K. Zhou, *DNA damage response signaling pathways and targets for radiotherapy sensitization in cancer.* Signal Transduct Target Ther, 2020. **5**(1): p. 60.
3. Ray Chaudhuri, A. and A. Nussenzweig, *The multifaceted roles of PARP1 in DNA repair and chromatin remodelling.* Nat Rev Mol Cell Biol, 2017. **18**(10): p. 610-621.
4. Blackford, A.N. and S.P. Jackson, *ATM, ATR, and DNA-PK: the trinity at the heart of the DNA damage response.* Molecular cell, 2017. **66**(6): p. 801-817.
5. Hopkins, J.L., L. Lan, and L. Zou, *DNA repair defects in cancer and therapeutic opportunities.* Genes & development, 2022. **36**(5-6): p. 278-293.
6. Villaruz, L.C., et al., *ATM protein is deficient in over 40% of lung adenocarcinomas.* Oncotarget, 2016. **7**(36): p. 57714.
7. Wang, C., et al., *Genetic vulnerabilities upon inhibition of DNA damage response.* Nucleic acids research, 2021. **49**(14): p. 8214-8231.
8. Gurley, K.E. and C.J. Kemp, *Synthetic lethality between mutation in Atm and DNA-PKcs during murine embryogenesis.* Current Biology, 2001. **11**(3): p.

191-194.

9. Fok, J.H., et al., *AZD7648 is a potent and selective DNA-PK inhibitor that enhances radiation, chemotherapy and olaparib activity.* Nature communications, 2019. **10**(1): p. 5065.

10. Yi, J., et al., *Potential of natural products as radioprotectors and radiosensitizers: Opportunities and challenges.* Food & Function, 2021. **12**(12): p. 5204-5218.

11. Komorowska, D., et al., *Natural radiosensitizers in radiotherapy: Cancer treatment by combining ionizing radiation with resveratrol.* International Journal of Molecular Sciences, 2022. **23**(18): p. 10627.

12. Eberhardt, J., et al., *AutoDock Vina 1.2. 0: New docking methods, expanded force field, and python bindings.* Journal of chemical information and modeling, 2021. **61**(8): p. 3891-3898.

13. Alhossary, A., et al., *Fast, accurate, and reliable molecular docking with QuickVina 2.* Bioinformatics, 2015. **31**(13): p. 2214-2216.

14. Coleman, R.G., et al., *Ligand pose and orientational sampling in molecular docking.* PloS one, 2013. **8**(10): p. e75992.

15. Díaz-Holguín, A., et al., *AlphaFold accelerated discovery of psychotropic agonists targeting the trace amine–associated receptor 1.* Science Advances, 2024. **10**(32): p. eadn1524.

16. Jumper, J., et al., *Highly accurate protein structure prediction with AlphaFold.* nature, 2021. **596**(7873): p. 583-589.

17. Baek, M., et al., *Accurate prediction of protein structures and interactions using a three-track neural network.* Science, 2021. **373**(6557): p. 871-876.

18. Díaz-Rovira, A.M., et al., *Are deep learning structural models sufficiently accurate for virtual screening? Application of docking algorithms to AlphaFold2 predicted structures.* Journal of Chemical Information and Modeling, 2023. **63**(6): p. 1668-1674.

19. Karelina, M., J.J. Noh, and R.O. Dror, *How accurately can one predict drug binding modes using AlphaFold models?* Elife, 2023. **12**: p. RP89386.

20. Lyu, J., et al., *AlphaFold2 structures guide prospective ligand discovery.* Science, 2024. **384**(6702): p. eadn6354.

21. Abramson, J., et al., *Accurate structure prediction of biomolecular interactions with AlphaFold 3.* Nature, 2024. **630**(8016): p. 493-500.

22. Krishna, R., et al., *Generalized biomolecular modeling and design with RoseTTAFold All-Atom.* Science, 2024. **384**(6693): p. eadl2528.

23. Lyu, J., et al., *Ultra-large library docking for discovering new chemotypes.* Nature, 2019. **566**(7743): p. 224-229.

24. Alon, A., et al., *Structures of the σ2 receptor enable docking for bioactive ligand discovery.* Nature, 2021. **600**(7890): p. 759-764.

25. Liu, F., et al., *Structure-based discovery of CFTR potentiators and inhibitors.* Cell, 2024. **187**(14): p. 3712-3725. e34.

26. Liu, F., et al., *Large library docking identifies positive allosteric modulators of the calcium-sensing receptor.* Science, 2024. **385**(6715): p. eado1868.

27. Lyu, J., J.J. Irwin, and B.K.J.N.c.b. Shoichet, *Modeling the expansion of virtual screening libraries.* 2023. **19**(6): p. 712-718.

28. Sorokina, M. and C. Steinbeck, *Review on natural products databases: where to find data in 2020.* Journal of cheminformatics, 2020. **12**(1): p. 20.

29. Irwin, J.J., et al., *ZINC20—a free ultralarge-scale chemical database for ligand discovery.* Journal of chemical information and modeling, 2020. **60**(12): p. 6065-6073.

30. Zdrazil, B., et al., *The ChEMBL Database in 2023: a drug discovery platform spanning multiple bioactivity data types and time periods.* Nucleic acids research, 2024. **52**(D1): p. D1180-D1192.

31. Fuchss, T., et al., *Highly potent and selective ATM kinase inhibitor M4076: A clinical candidate drug with strong anti-tumor activity in combination therapies.* Cancer Research, 2019. **79**(13_Supplement): p. 3500-3500.

32. Yap, T.A., et al., *Phase I trial of first-in-class ATR inhibitor M6620 (VX-970) as monotherapy or in combination with carboplatin in patients with advanced solid tumors.* Journal of Clinical Oncology, 2020. **38**(27): p. 3195-3204.

33. van Bussel, M.T., et al., *A first-in-man phase 1 study of the DNA-dependent protein kinase inhibitor peposertib (formerly M3814) in patients with advanced solid tumours.* British Journal of Cancer, 2021. **124**(4): p. 728-735.

34. Coleman, R.L., et al., *Veliparib with first-line chemotherapy and as maintenance therapy in ovarian cancer.* New England Journal of Medicine, 2019. **381**(25): p. 2403-2415.

35. Liang, S., et al., *Structural insights into inhibitor regulation of the DNA repair protein DNA-PKcs.* Nature, 2022. **601**(7894): p. 643-648.

36. Kumar, V. and J.V. Aldrich, *A solid-phase synthetic strategy for labeled peptides: synthesis of a biotinylated derivative of the δ opioid receptor antagonist TIPP (Tyr-Tic-Phe-Phe-OH).* Organic Letters, 2003. **5**(5): p. 613-616.

37. Bock, M.G., et al., *Benzodiazepine gastrin and brain cholecystokinin receptor ligands; L-365,260.* Journal of medicinal chemistry, 1989. **32**(1): p. 13-16.

38. Lotti, V.J. and R.S. Chang, *A new potent and selective non-peptide gastrin antagonist and brain cholecystokinin receptor (CCK-B) ligand: L-365,260.* European journal of pharmacology, 1989. **162**(2): p. 273-280.

39. Yang, Z., et al., *miR-328-3p enhances the radiosensitivity of osteosarcoma and regulates apoptosis and cell viability via H2AX.* Oncology reports, 2018. **39**(2): p. 545-553.

40. Liu, T.-T., et al., *Discovery of a meisoindigo-derived protac as the ATM degrader: Revolutionizing colorectal cancer therapy via synthetic lethality with ATR inhibitors.* Journal of Medicinal Chemistry, 2024. **67**(9): p. 7620-7634.

41. Adasme, M.F., et al., *PLIP 2021: expanding the scope of the protein–ligand interaction profiler to DNA and RNA.* Nucleic acids research, 2021. **49**(W1): p. W530-W534.

42. Irwin, J.J., et al., *Automated docking screens: a feasibility study.* Journal of medicinal chemistry, 2009. **52**(18): p. 5712-5720.

43. Stein, R.M., et al., *Property-unmatched decoys in docking benchmarks.* Journal of chemical information and modeling, 2021. **61**(2): p. 699-714.

44. Daina, A., O. Michielin, and V. Zoete, *SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules.* Scientific reports, 2017. **7**(1): p. 42717.

45. Fu, L., et al., *ADMETlab 3.0: an updated comprehensive online ADMET prediction platform enhanced with broader coverage, improved performance, API functionality and decision support.* Nucleic acids research, 2024. **52**(W1): p. W422-W431.

46. Abraham, M.J., et al., *GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers.* SoftwareX, 2015. **1**: p. 19-25.

47. Eastman, P., et al., *OpenMM 7: Rapid development of high performance algorithms for molecular dynamics.* PLoS computational biology, 2017. **13**(7): p. e1005659.

48. Olsson, M.H., et al., *PROPKA3: consistent treatment of internal and surface residues in empirical p K a predictions.* Journal of chemical theory and computation, 2011. **7**(2): p. 525-537.

49. Huang, J., et al., *CHARMM36m: an improved force field for folded and intrinsically disordered proteins.* Nature methods, 2017. **14**(1): p. 71-73.

50. Vanommeslaeghe, K., et al., *CHARMM general force field: A force field for drug-like molecules compatible with the CHARMM all-atom additive biological force fields.* Journal of computational chemistry, 2010. **31**(4): p. 671-690.

51. Berendsen, H.J., et al., *Molecular dynamics with coupling to an external bath.* The Journal of chemical physics, 1984. **81**(8): p. 3684-3690.

52. Parrinello, M. and A. Rahman, *Polymorphic transitions in single crystals: A new molecular dynamics method.* Journal of Applied physics, 1981. **52**(12): p. 7182-7190.

53. Darden, T., D. York, and L. Pedersen, *Particle mesh Ewald: An N log (N) method for Ewald sums in large systems.* Journal of chemical physics, 1993. **98**: p. 10089-10089.

54. Hess, B., et al., *LINCS: A linear constraint solver for molecular simulations.* Journal of computational chemistry, 1997. **18**(12): p. 1463-1472.

55. Valdés-Tresanco, M.S., et al., *gmx_MMPBSA: a new tool to perform end-state free energy calculations with GROMACS.* Journal of chemical theory and computation, 2021. **17**(10): p. 6281-6291.

56. Cai, X., et al., *High-throughput proteomic sample preparation using pressure cycling technology.* Nature Protocols, 2022. **17**(10): p. 2307-2325.

57. Yu, F., et al., *Analysis of DIA proteomics data using MSFragger-DIA and FragPipe computational platform.* Nature communications, 2023. **14**(1): p. 4154.

58. Demichev, V., et al., *DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput.* Nature methods, 2020. **17**(1): p. 41-44.

59. Chambers, M.C., et al., *A cross-platform toolkit for mass spectrometry and proteomics.* Nature biotechnology, 2012. **30**(10): p. 918-920.

60. Ritchie, M.E., et al., *limma powers differential expression analyses for RNA-sequencing and microarray studies.* Nucleic acids research, 2015. **43**(7): p. e47-e47.

61. Sherman, B.T., et al., *DAVID: a web server for functional enrichment analysis and functional annotation of gene lists (2021 update).* Nucleic acids research, 2022. **50**(W1): p. W216-W221.

62. Szklarczyk, D., et al., *The STRING database in 2021: customizable protein–protein networks, and functional characterization of user-uploaded gene/measurement sets.* Nucleic acids research, 2021. **49**(D1): p. D605-D612.

63. Shannon, P., et al., *Cytoscape: a software environment for integrated models of biomolecular interaction networks.* Genome research, 2003. **13**(11): p. 2498-2504.

64. Merico, D., et al., *Enrichment map: a network-based method for gene-set enrichment visualization and interpretation.* PloS one, 2010. **5**(11): p. e13984.

Acknowledgements

This work was supported by the Hainan Province Academician Workstation (Changbin Yu), the Specific Research Fund of The Innovation Platform for Academicians of Hainan Province (YSPTZX202407); the Natural Science Foundation of Shandong Province (2022HWYQ-081 and ZR2023LZY009); Scientific Research Cultivation Project in Emerging Strategic Fields (202402); the Academic promotion project of Shandong First Medical University, and funding from Jinan City.

Author contributions

J.T. and C.Y. designed the work; J.T., X.W., M.L. conducted a research and investigation process; J.T., T.W., J.C., J.C. carried out the proteomics experiments; J.T., X.W., M.L. analyzed the data; J.T., X.W., Y.L., C.X., and Y.F. reviewed the manuscript; J.T., M.L., and C.Y. supervised the study; J.T. wrote the paper.

Competing interest

The authors declare no competing interests.

Corresponding authors

Correspondence to Changbin Yu.