In Situ Amplification and Spatial Mapping of α-Synuclein Seeding Activity

Hengxu Mao[1, #, *], Yaoyun Kuang[1, #], Wei Dai[1], Hao Lin[1], Jinkun Huang[1, *] and Ping-Yi Xu[1, *]

Author affiliations:

1. Dept. of Neurology, the First Affiliated Hospital of Guangzhou Medical University, Guangzhou, 510120, China.

[#]**These authors are equally contributing to this work.**

*Corresponding to:

1. Hengxu Mao, Dept. of Neurology, the First Affiliated Hospital of Guangzhou Medical University, Guangzhou, 510120, China. Email address: hengxu2010@qq.com.

2. Jinkun Huang, Dept. of Neurology, the First Affiliated Hospital of Guangzhou Medical University, Guangzhou, 510120, China. Email address: 2009681002@gzhmu.edu.cn.

3. Ping-Yi Xu, Dept. of Neurology, the First Affiliated Hospital of Guangzhou Medical University, Guangzhou, 510120, China. Email address: pingyixu@sina.com.

Abstract

The prion-like propagation of proteopathic α-synuclein (αSyn) "seeds" is a central pathogenic mechanism in synucleinopathies. However, methods to directly visualize this "seeding activity" within intact tissue are lacking. Conventional antibody-based techniques provide only static snapshots of pathological deposits, while sensitive seed amplification assays lose spatial information via mechanical fragmentation. Here, we introduce the αSyn Quiescent Seed Amplification Assay (QSAA), a method for specific, robust, in situ amplification of

αSyn seeds directly within tissue sections. QSAA demonstrates superior sensitivity and earlier detection than pS129 and ThS staining, revealing widespread seeding in a PD mouse model, and successfully identifies endogenous seeds in archival human brains. By integrating with immunofluorescence, IF-QSAA uncovered microglial seeding activity and established that pS129 phosphorylation and seeding activity represent distinct pathological states. This offers unprecedented spatial insights into synucleinopathy pathogenesis, holding significant promises for early diagnosis and therapeutic evaluation.

Keywords: α-Synuclein, Seed amplification assay, Quiescent SAA, Parkinson's disease, Seeding activity

Introduction

Synucleinopathies, including Parkinson's disease (PD), dementia with Lewy bodies (DLB), and multiple system atrophy (MSA), are progressive neurodegenerative disorders defined by the pathological accumulation of α-synuclein (αSyn) [1,2]. In these diseases, αSyn misfolds into proteopathic seeds capable of self-propagation through a prion-like mechanism, templating the conversion of native αSyn into insoluble aggregates[3,4]. This functional property, termed seeding activity, is a primary driver of pathology progression, leading to widespread neuronal dysfunction and neurodegeneration[5,6]. Consequently, visualizing the spatiotemporal distribution and propagation pathways of αSyn seeding activity is critical for understanding synucleinopathy pathogenesis and developing targeted therapies.

Conventional antibody-based techniques, which often target pathological markers like αSyn phosphorylated at Serine 129 (pS129), have been invaluable in localizing αSyn aggregates in diseased brains[7,8]. However, these methods provide only a static snapshot of aggregate presence and fail to capture the dynamic seeding activity that underpins pathology[9-12]. To bridge this gap, seed amplification assays (SAAs) have been developed, exploiting the

inherent ability of misfolded αSyn seeds to induce amplification in vitro by recruiting exogenous monomeric substrates[13,14]. Despite their high sensitivity, traditional SAAs depend on tissue homogenization and mechanical fragmentation (e.g., shaking or sonication) to sustain the amplification reaction[15,16]. This process disrupts tissue architecture and abolishes spatial information about seed localization.

To address these challenges, we developed the αSyn Quiescent Seed Amplification Assay (QSAA), a method that enables in situ amplification of αSyn seeds within tissue sections without mechanical fragmentation. By optimizing reaction conditions, including the introduction of sulfate ions (SO_4^{2-}), elevating temperature to 70°C, and using mouse-derived αSyn monomers as the substrate, QSAA achieves seed-dependent fibril elongation under quiescent conditions. This approach preserves spatial context and generates high-resolution propagation maps, revealing pathological spread patterns inaccessible by antibody staining alone. QSAA is compatible with a variety of samples, demonstrating broad applicability by successfully detecting endogenous seeds in human brain sections, including long-term formalin-fixed archival specimens. Furthermore, by integrating QSAA with immunofluorescence (IF), IF-QSAA can simultaneously visualize amplified seeds and specific cellular markers. This dual-labeling approach provides a comprehensive, cellular-level view of αSyn seeding activity, offering new insights into the mechanisms of synucleinopathies. Together, QSAA and IF-QSAA represent powerful tools to advance our understanding of αSyn propagation, holding great promise for improving diagnosis and therapeutic development.

Results

Development of a quiescent system for αSyn seed amplification

A key prerequisite for in situ amplification of αSyn seeds is the ability to achieve robust and

specific amplification under quiescent conditions. Traditional SAAs primarily rely on cyclic

physical fragmentation to continuously generate new growth ends from fibrils that have

entered the reaction plateau[17,18], thereby sustaining rapid amplification (**Fig. 1a, solid line**).

In contrast, a quiescent system necessitates an alternative mechanism to maintain high

amplification efficiency (**Fig. 1a, dashed line**). To address this, our strategy focused on

optimizing the biochemical environment to intrinsically accelerate fibril amplification in a

seed-dependent manner, thereby circumventing the need for physical fragmentation (**Fig. 1b**).

We systematically examined several factors known to affect αSyn fibril growth,

including the αSyn monomer substrate[13,19,20], chemical additives[21-24], and incubation

temperature[25,26]. Initially, we constructed expression plasmids for human wild-type (hWT),

human A53T mutant (hA53T), and mouse wild-type (mWT) αSyn, confirming their

sequences (**Fig. 1c**). Notably, mWT αSyn naturally contains a threonine at position 53,

analogous to the pathogenic A53T mutation in humans, which is known to accelerate αSyn

aggregation[27]. Consistent with this, we demonstrated a progressively faster aggregation rate

from hWT to hA53T and then to mWT, confirming that mWT αSyn serves as a more

aggregation-prone substrate suitable for rapid amplification. Next, we explored the impact of

additives and temperature on fibril amplification. Previous studies have shown that sulfate

ions (SO_4^{2-}) can promote αSyn fibril elongation by electrostatically stabilizing misfolded

aggregates[21,22] and that an elevated temperature accelerates aggregation kinetics[25,26]. We

evaluated these parameters individually to establish their baseline effects in the presence of

αSyn pre-formed fibrils (PFFs) under cyclic fragmentation conditions (**Fig. 1d**). Each

parameter (mWT αSyn monomer, SO_4^{2-}, 70°C) independently shortened the lag phase of

αSyn seed amplification (**Fig. 1e**). However, combining these factors under cyclic

fragmentation resulted in rapid spontaneous amplification, leading to poor specificity (**Fig.**

1f). Remarkably, applying the same combination (mWT αSyn monomer, SO_4^{2-}, 70°C) under

quiescent conditions yielded rapid and specific amplification with a high fold separation of

7.547 (**Fig. 1g**). These results underscore that the optimized chemical environment and

elevated temperature in QSAA effectively compensate for the lack of mechanical

fragmentation, enabling continuous and robust fibril growth from existing seeds while

minimizing spontaneous nucleation.

To further characterize the amplification products, we performed Transmission Electron

Microscopy (TEM) on samples generated by QSAA (**Fig. 1h**). TEM analysis revealed

fibrillar morphologies characteristic of αSyn aggregates, including both densely packed and

loosely associated fibrillar assemblies. In contrast, no fibrillar products were observed in the

unseeded group, further demonstrating the specificity of the assay (**Fig. 1h**, Inset).

QSAA Enables In Situ Amplification and Mapping of αSyn Seeding Activity in Brain

Sections

Having established a quiescent amplification system, we sought to expand its application for

in situ amplification on brain tissue sections. We developed a standardized operating

procedure (SOP) involving tissue fixation, dehydration, sectioning, post-fixation, and

incubation with the QSAA reaction buffer (**Fig. 2a**). To minimize artifacts caused by

microscale fluid movement, a semi-solid agarose membrane was overlaid on the tissue

sections during incubation[28].

Brain sections from a synucleinopathy mouse model[29,30] at 180 days post-injection (dpi)

of mouse αSyn pre-formed fibrils (mPFFs) revealed widespread αSyn seeding activity after 5

hours of incubation (**Fig. 2b**). The signal propagated along neuroanatomical tracts, such as

the bilateral corpus callosum (CC) and anterior commissure (AC), into the contralateral

hemisphere, whereas control sections from PBS-injected mice were negative (**Fig. 2b, Inset**).

This pattern, consistently observed across coronal, sagittal, and horizontal sections,

effectively mapped pathology routes directly within the tissue, with sagittal views clearly

showing dissemination along the anterior commissure's olfactory limb (ACO) towards the

olfactory bulb (OB) (**Fig. 2b, sagittal**).

To confirm the Thioflavin T (ThT) signal represented seed-dependent fibril formation,

we used TEM on QSAA-treated tissue, which revealed typical αSyn amyloid fibrils within

the tissue interstitium and on its surface (**Fig. 2c**). Brightfield microscopy further showed that

fluorescent signals precisely co-localized with macroscopic aggregates, confirming that

tangible amplification products were generated (**Fig. 2d**). These observations validate that

QSAA amplifies αSyn seeds within intact tissue architecture.

QSAA Provides Superior Sensitivity and Early Detection of αSyn Pathology

By leveraging the ability of QSAA to preserve spatial information, we generated a brain-wide

map of αSyn seeding activity in mPFF-injected mice (180 dpi) and compared it with pS129-

αSyn immunofluorescence on consecutive sections (**Fig. 3a**). QSAA and pS129 signals

showed broad concordance across multiple brain regions, including the striatum (STR),

primary and secondary motor cortex (M1/M2), secondary somatosensory cortex (S2), and

entorhinal cortex (ECT), confirming that QSAA faithfully localizes pathological seeds.

Importantly, QSAA revealed a much wider pathological landscape, particularly within white

matter tracts such as the bilateral corpus callosum (CC), anterior commissure (ACA/ACP),

ipsilateral internal capsule (IC), external capsule (EC), medial forebrain bundle (MFB),

fasciculus retroflexus (FR), and nigrostriatal tract (NST). In these regions, pS129

immunoreactivity was relatively sparse, likely reflecting the lower density of neuronal soma.

To further demonstrate the sensitivity of QSAA, we compared QSAA, pS129-IF, and

Thioflavin S (ThS) staining on sections from mice injected with one-tenth the standard dose

of mPFFs and analyzed at 30 dpi (**Fig. 3b**). QSAA revealed widespread fluorescent signals

across the cortex, corpus callosum, and striatum of both hemispheres, whereas pS129

detected only sparse, neuritic-like structures, and ThS staining showed even weaker signals

(**Fig. 3c**).

We next examined the ability of QSAA to detect early-stage pathology. Because QSAA

directly amplifies functional seeding activity rather than depending on downstream

phosphorylation events, it confers a mechanistic advantage over pS129 immunostaining.

Remarkably, QSAA detected distinct seeding activity as early as 7 dpi, a stage at which

pS129 signals were undetectable (**Fig. 3d**). By 14 dpi, QSAA revealed widespread

dissemination of αSyn seeding throughout the brain, whereas pS129 immunoreactivity was

only beginning to emerge. Whole-brain QSAA mapping at 14 dpi confirmed broad

propagation of seeding activity beyond the injection site, with no signal observed in PBS-

injected controls (**Fig. 3e**). These findings demonstrate that extensive αSyn seeding precedes

the detectable onset of pS129 pathology in the exogenous seeding model.

QSAA is Applicable to Endogenous Seeds in Human Brain Tissue

We next evaluated the applicability of QSAA to detect endogenous αSyn seeds in postmortem

human brain tissue. Striatal sections from patients with PD, progressive supranuclear palsy

(PSP), and healthy controls (HC) were analyzed. QSAA generated ThT fluorescence

selectively in PD brains, with no signal observed in HC or PSP samples, demonstrating high

specificity (**Fig. 4a**). Notably, QSAA was able to detect seeding activity in PD specimens that

had been fixed in formalin for more than 20 years, in which pS129-IF failed to detect

pathology (**Fig. 4a, FF-PD**). This finding indicates the stability of seeding-competent αSyn

aggregates compared with antibody epitopes that deteriorate during prolonged fixation. To

validate these results independently, we performed Real-Time Quaking-Induced Conversion

(RT-QuIC) on lysates from the same brain specimens, which confirmed the seeding activity

in both frozen and formalin-fixed PD samples (**Fig. 4b-c**).

QSAA further revealed distinct spatial distributions of seeding activity in human PD

brains (**Fig. 4d**). Seeding was detected predominantly in gray matter regions such as the

striatum and cortex, whereas the corpus callosum showed no detectable activity (**Fig. 4e**).

This contrasts with the mPFF-inoculated mouse model, where seeding also occurs within

white matter tracts, suggesting differences in propagation dynamics between exogenously

induced pathology and endogenously developed diseases. Finally, a time-course analysis of

QSAA on PD brain sections revealed a clear, time-dependent increase in ThT fluorescence

intensity at defined sites of αSyn seeding (**Fig. 4f**). Signals appeared as early as 1 h and

continued to amplify through 5 h, highlighting both the sensitivity of the method and its

ability to dynamically monitor in situ seeding activity.

IF-QSAA Reveals Distinct Cellular Localization and Complementary Pathological

States of Seeding Activity and pS129

We developed an integrated immunofluorescence-QSAA (IF-QSAA) protocol to determine

the cellular localization of αSyn seeding activity. First, we confirmed the spectral

compatibility of ThT with common IF fluorophores, finding that its excitation spectrum is

distinct from DAPI, Cy3, and Cy5, but overlaps significantly with FITC (**Fig. 5a**). Next, a

sequential optimization revealed that the QSAA reaction conditions ($70°C$, SO_4^{2-}, 5h)

degraded antibody epitopes, necessitating an "IF-first" workflow to preserve both signals

(**Fig. 5b**). Applying this protocol with confocal microscopy on brain tissue from mPFF-

injected mice (180 dpi), we visualized intraneuronal QSAA products (**Fig. 5c, arrows**) and

larger products extending into the extracellular space (**Fig. 5c, arrowheads**). These results

validated the capability of IF-QSAA to map seeding activity with subcellular precision,

leading to our standardized IF-QSAA protocol (**Fig. 5d**).

To identify the specific cell types involved in αSyn seeding, we co-labeled sections from

mPFF-injected mice (30 dpi) with markers for neurons (NeuN), microglia (IBA1), and

astrocytes (GFAP). Strikingly, in addition to neuronal signals (**Fig. 5e**), we observed

extensive co-localization between QSAA signals and IBA1-positive microglia (**Fig. 5f**), but

not with GFAP-positive astrocytes (**Fig. 5g**). This finding contrasted with the cellular

distribution of pS129-αSyn pathology[31], which was predominantly found within neurons

(**Fig. 5h**) and showed minimal co-localization with microglia (**Fig. 5i**). These findings

suggest a previously underappreciated role for microglia in the uptake, processing, or

propagation of αSyn seed species, differing from the traditional view that Lewy pathology

primarily accumulates within neurons.

To further investigate the relationship between αSyn seeding activity and pS129

pathology, we applied IF-QSAA to brain tissue sections from the mPFF mouse model.

Through simultaneous imaging of the QSAA signal and pS129, we observed a heterogeneity

among αSyn aggregates, with multiple subtypes coexisting in the same microspatial

environment (**Fig. 5j**). Initially, we identified a large population of aggregates positive for

both signals (QSAA+/pS129+), which exhibited a characteristic "core-halo" structure,

comprising a dense, pS129-positive core enveloped by a more diffuse QSAA signal halo.

This finding suggested that phosphorylated αSyn with seeding capacity is a significant

component of the pathological landscape. However, this observation raises a critical question:

is pS129 a reliable surrogate for αSyn seeding activity? A closer examination of the

magnified regions in Figure 5j revealed that, in addition to the dual-positive aggregates, we

could also identify: 1) aggregates that were exclusively pS129-positive but QSAA-negative

(QSAA−/pS129+), appearing as discrete red puncta; and 2) aggregates that were exclusively

QSAA-positive but pS129-negative (QSAA+/pS129−), appearing as distinct green puncta.

This direct visual evidence demonstrates that αSyn seeding activity and its pS129

phosphorylation state are not strictly concordant or co-localized (**Fig. 5k**).

We therefore propose a "four-quadrant model" to classify αSyn aggregates (**Fig. 5l**):

QSAA+/pS129+ (Co-existent): Phosphorylated aggregates with seeding capacity, forming the

characteristic core-halo structure. QSAA+/pS129− (Seeding-Dominant): Aggregates

possessing seeding activity but with pS129 levels below the detection threshold or with a

masked epitope. QSAA−/pS129+ (Phosphorylation-Dominant): Phosphorylated aggregates

that have either lost or inherently lack seeding competency. QSAA−/pS129− (Pathology-

Free): Regions devoid of detectable seeding activity and pS129 pathology. This model

emphasizes that αSyn seeding activity and pS129 immunoreactivity represent two distinct yet

complementary pathological states, rather than being equivalent disease markers. It provides

a more precise framework for understanding the molecular heterogeneity and

microenvironmental complexity of αSyn pathology in synucleinopathies.

Discussion

In this study, we established QSAA, a method that enables the in situ detection of αSyn

seeding activity. Unlike antibody-based methods that capture pre-formed aggregates[32], or

conventional SAAs that require tissue homogenization[33,34], QSAA achieves rapid, seed-

dependent amplification under quiescent conditions. This methodology allows for the direct

visualization of functional seeding activity within intact tissue sections.

QSAA provided several key insights that extend beyond the capabilities of existing

techniques. First, its amplification-based nature provided markedly superior sensitivity,

enabling the detection of nascent αSyn seeds. Second, QSAA revealed a broader pathological

landscape, particularly within white matter tracts and microglial populations, where antibody-

based methods are often less effective. This is likely because conventional antibodies

targeting post-translational modifications (PTMs) like pS129 rely on late-stage pathological

events[9,11], giving QSAA a distinct advantage in detecting seeding species that are either

unmodified or at very early stages of aggregation. Together with the IF-QSAA protocol, these

findings led us to propose a conceptual "four-quadrant model," in which seeding activity and

pS129 phosphorylation define both overlapping and distinct dimensions of αSyn pathology.

This framework highlights early-stage QSAA+/pS129− seeds as a potentially critical driver

of disease progression that is not captured by conventional markers. Importantly, QSAA also

demonstrated strong translational potential. It successfully amplified endogenous seeds from

postmortem PD brains, including archival tissue fixed for over 20 years, where pS129

immunoreactivity had deteriorated. This underscores the remarkable stability of seeding-

competent αSyn structures and positions QSAA as a promising tool for both retrospective

neuropathological studies and diagnostic applications.

While QSAA offers the unique advantage of in situ assessment, it currently cannot

differentiate between distinct αSyn strains, a property reported for some conventional

SAAs[35-37], nor can it directly identify PTMs as antibody-based methods can[38,39]. Future

methodological refinements, such as fluorescence-based conformational fingerprinting of

amplified fibrils[40] or the engineering of strain-specific monomer substrates, may broaden its

utility.

Although the high sensitivity of QSAA arises from controlled, seed-dependent

amplification rather than from non-specific background signals, the risk of experimental false

positives still requires rigorous control. Despite multiple layers of mitigation strategies

including an agarose overlay, strict incubation time, parallel negative samples, and in situ

TEM validation, further standardization will be essential for multi-center implementation.

Uniform parameters such as section thickness, fixation protocols, and data normalization will

be critical for ensuring inter-laboratory reproducibility[17].

In conclusion, QSAA provides a powerful platform for mapping αSyn seeding activity in

situ with high sensitivity and specificity. By extending beyond static aggregate detection to

capture the functional and spatial dimensions of pathogenic seeds, QSAA deepens our

understanding of synucleinopathy pathogenesis and holds promise for clinical applications in

early diagnosis and therapeutic evaluation. Moreover, the principle of quiescent amplification

may be adaptable to other misfolded proteins, such as Tau and TDP-43, particularly those that

have already demonstrated cyclic fragmentation-based amplification in vitro[41,42], further

broadening its relevance to neurodegenerative disease research.

Methods

Ethics and Oversight

All animal procedures were conducted in strict accordance with the guidelines established by the Institutional Animal Care and Use Committee of Guangzhou Medical University. Human brain tissues were obtained from the Department of Neurology, First Affiliated Hospital of Guangzhou Medical University, with written informed consent from the next of kin and approval from the hospital Ethics Committee (MRE. No. 155). All experiments complied with relevant ethical regulations.

Animals and Stereotaxic Injections

Male C57BL/6J mice (10-12 weeks old, RRID: IMSR_JAX:000664) were purchased from the Experimental Animal Center of Guangzhou Medical University. Animals were maintained in a SPF facility on a 12-h light/dark cycle with ad libitum access to food and water. Sample size was chosen based on previous studies using similar models to ensure sufficient power for detecting inter-group differences. Animals were randomly assigned to experimental groups. All injections, data acquisition, and subsequent analyses were performed by experimenters blinded to the treatment conditions

For stereotaxic surgery, mice were anesthetized with isoflurane (2.5% for induction, 1.5% for maintenance) and placed in a stereotaxic frame (Kopf Instruments, Model 940). Recombinant mouse αSyn pre-formed fibrils (mPFFs; 5 μg in 1 μL) or an equal volume of sterile PBS were injected unilaterally into the right striatum. Coordinates were determined relative to bregma according to the Paxinos & Franklin mouse brain atlas (3rd edition): AP: +0.8 mm, ML: +2.0 mm, DV: -3.5 mm. Injections were performed using a 10-μL Hamilton syringe with a 33-gauge needle connected to a microinfusion pump at a rate of 0.2 μL/min. The needle was left in place for 10 min post-injection to minimize backflow before being

slowly withdrawn. For the low-dose sensitivity comparison, a one-tenth dose of mPFFs (0.5

µg in 1 µL) was used. At specified time points (7, 10, 14, 30, or 180 days post-injection),

mice were deeply anesthetized and transcardially perfused with ice-cold PBS followed by 4%

(w/v) paraformaldehyde (PFA) in 0.1 M phosphate buffer (PB, pH 7.4). Brains were post-

fixed in the same fixative for 24 h at 4°C, then cryoprotected in 30% (w/v) sucrose in PBS

until sectioning.

Human Brain Specimens

Postmortem human brain tissues (collected 2005-2022) were obtained from the First

Affiliated Hospital of Guangzhou Medical University. The cohort included pathologically

confirmed cases of Parkinson's disease (PD; n=3), progressive supranuclear palsy (PSP; n=3),

and age-matched healthy controls with no history of neurological disease (HC; n=3).

Diagnoses were established postmortem based on consensus neuropathological criteria,

including Braak staging for Lewy body pathology. The postmortem interval (PMI) for all

fresh-frozen and standard-fixed samples was <12 h. For each case, the left hemisphere was

fixed in 10% neutral buffered formalin (NBF) for 2-4 weeks, while the right hemisphere was

coronally sliced and snap-frozen at -80°C. Along-term fixed PD specimen (FF-PD) had been

stored in 10% NBF for over 20 years at room temperature.

α-Synuclein Monomer Expression and PFF Preparation/QC

Recombinant full-length mouse wild-type α-synuclein (mWT, UniProt ID: P58992) was

expressed in *E. coli* BL21 (DE3) cells (TransGen Biotech, CD601) and purified as previously

described with modifications[43,44]. Following IPTG induction (1 mM, 4 h), cells were lysed by

sonication in high-salt buffer (10 mM Tris-HCl, pH 7.5, 500 mM NaCl, 1 mM EDTA, 1 mM

PMSF, and 1x Protease inhibitor). The lysate was boiled for 15 min, followed by 45%

ammonium sulfate precipitation and pH 3.0 acid precipitation. Endotoxin was removed using

a high-capacity endotoxin removal resin (Sangon Biotech, B641718). The protein was further

purified by anion exchange chromatography (HiTrap Q HP column, Cytiva, 17115401). The

final monomer was dialyzed against TBS (10mM Tris-HCl pH 7.5, 50mM NaCl), filtered

through a 50-kDa cutoff filter (Amicon Ultra), and stored at -80°C.

To generate mPFFs, monomeric mouse αSyn (5 mg/mL in PBS) was incubated at 37°C

with continuous agitation (1000 rpm) in an Eppendorf ThermoMixer C for 7 days. The

resulting fibril suspension was fragmented into seeds using a non-contact water-bath

sonicator (SCIENTZ, SCIENTZ08-III) for a total of 5 min using 300 pulses of 1-s on/1-s off

at 90% power. PFF quality control was performed for each batch[45]. TEM confirmed fibrillar

morphology. Dynamic light scattering (DLS) confirmed an average hydrodynamic diameter

of 50-150 nm. Functional activity was validated using a standard SAA, with each batch

required to produce a lag time within ±15% of a reference standard batch when seeded with 1

pg of PFFs.

Plate-based Seed Amplification Assays

Agitated SAA: Reactions were prepared in a 96-well black, clear-bottom plate (Thermo

Scientific, 165305) in a final volume of 100 μL. Each well contained 1 mg/mL m-αSyn

monomer, 40 μM Thioflavin T (ThT), and 1 pg PFFs (or PBS for no-seed controls) in

Agitation Buffer (10 mM NaCl, 10 mM Tris-HCl, pH 7.5). Where indicated, 800 mM Na_2SO_4

was added. The plate was sealed with an optical adhesive film (Applied Biosystems,

4311971) and incubated in a plate reader (SpectraMax iD5, Molecular Devices) at 37°C. A

cycle of 1 min orbital shaking (700 rpm, double orbital) followed by 14 min of rest was

repeated. ThT fluorescence was measured every 15 min (Ex: 450 nm, Em: 490 nm, bottom-

read).

Quiescent SAA: Reactions were prepared in a 96-well polypropylene PCR plate

(Thermo Scientific, AB0700) in a final volume of 20 μL. Each reaction contained 1 mg/mL

mWT-αSyn monomer and 40 μM ThT in QSAA Reaction Buffer (800 mM Na_2SO_4, 25 mM

NaCl, 10mM Tris-HCl, pH 6.0), seeded with 1 pg mPFFs or TBS. Each reaction was overlaid

with 20 μL of mineral oil (Biosharp, BS927) to prevent evaporation. The plate was sealed and

incubated in a real-time PCR instrument (Bio-Rad, CFX96) at 70°C without agitation. ThT

fluorescence was monitored every 30 min via the FAM channel (Ex: 452-486 nm, Em: 505-

545 nm). Lag phase was defined as the time to cross a threshold of the mean baseline

fluorescence plus five standard deviations.

In Situ QSAA on Tissue Sections

Brains were sectioned at 14 μm thickness onto high-adhesion glass slides (CITOTEST,

188105W). Sections were air-dried for 30 min, then post-fixed in ice-cold acetone for 10 min

at -20°C and air-dried again. For FF-PD tissues, 95°C PBS treatment for 1 h was performed

prior to sectioning to reverse formalin cross-linking and restore seeding activity. To perform

QSAA, a reaction mix was prepared containing 1 mg/mL m-αSyn monomer, 40 μM ThT in

Reaction Buffer (800 mM Na_2SO_4, 25 mM NaCl, 10 mM Tris-HCl, pH 6.0). To prevent

micro-convection and ensure even reagent distribution, a pre-warmed (~50 °C) 1% (w/v)

agarose solution (in ddH_2O) was overlaid onto the tissue section to form a gel approximately

1 mm thick. After gelling at room temperature (~5min), the agarose was carefully trimmed

flush with the tissue edges using a scalpel. Subsequently, 300 μL of the reaction mix was

added to cover each section. The slide was then placed in a humidified, sealed chamber on a

temperature-controlled stage (Ruicheng Instrument Co., TS400) and incubated at 70°C for 0-

12 h unless otherwise specified. Following incubation, the agarose film was gently removed

by immersing the slide in heated (95 °C) PBS, and the slide was washed 3 times for 1 min

each in PBS before mounting with a coverslip.

IF-QSAA Workflow

An "IF-first" protocol was implemented to preserve antibody epitopes, which were found to be sensitive to the 70°C incubation. Sections were first processed for standard immunofluorescence. Briefly, sections were permeabilized with 0.5% Triton X-100 in PBS for 15 min and blocked with 5% normal goat serum in PBS for 1 h at room temperature. Primary antibodies (see Table 1 for details) were diluted in blocking buffer and incubated overnight at 4°C. After washing, sections were incubated with corresponding Alexa Fluor-conjugated secondary antibodies (1:1000, Invitrogen) for 1 h at room temperature. Following IF, sections were washed thoroughly in PBS and then subjected to the In Situ QSAA protocol as described above.

Thioflavin S Staining

Sections were incubated in 0.05% ThS in 50% ethanol for 8 min, differentiated in 80% ethanol, and treated with TrueBlack® Lipofuscin Autofluorescence Quencher (Biotium, 23007) to suppress background fluorescence. After rinsing in PBS, slides were coverslipped with aqueous mounting medium and imaged.

Microscopy and Image Quantification

Fluorescence images were acquired using a NIKON ECLIPSE Ti2-E microscope equipped with a 4×, 10×, and 20× objectives. High-resolution confocal images were acquired on a Leica SP8 scanning confocal microscope using a 40× (NA 0.75) or a 63× oil-immersion (NA 1.40) objective. To prevent spectral bleed-through, a sequential scanning strategy was employed, acquiring signals in descending order of emission wavelength: Cy5/Alexa Fluor 647 (Ex: 633 nm, Em: 650-720 nm) → Cy3/Alexa Fluor555 (Ex: 561 nm, Em: 570-630 nm) → ThT/QSAA (Ex: 488 nm, Em: 495-550 nm) →DAPI (Ex: 405 nm, Em: 415-470 nm). Notably, because ThT emission (not excitation spectra) partially overlaps with DAPI,

subtraction or spectral unmixing was applied during analysis. Z-stacks were acquired with a 2

µm step size over a 50 µm range.

All images intended for comparison were acquired using identical microscope settings

(laser power, gain, offset) and processed with the same linear adjustments for brightness and

contrast. Image analysis was performed using Fiji/ImageJ (v2.3.0). For quantification of

fluorescence intensity, regions of interest were manually drawn, and the mean gray value was

measured. All experiments were performed with at least three independent biological

replicates.

Transmission Electron Microscopy

For solution-state products, 10 µL of QSAA reaction mixture was adsorbed onto glow-

discharged, 400-mesh carbon-coated copper grids for 2 min, washed with ddH$_2$O, and

negatively stained with 2% (w/v) uranyl acetate. For in situ products, QSAA-treated tissue

sections were scraped from the slide, fixed in 2.5% glutaraldehyde/2% PFA in 0.1 M

cacodylate buffer, post-fixed with 1% osmium tetroxide, dehydrated in ethanol, and

embedded in EMBed 812 resin. Ultrathin sections (70 nm) were cut, placed on grids, and

stained with uranyl acetate and lead citrate. All grids were imaged on a FEI Talos L120C

TEM at 120 kV.

RT-QuIC Validation

Brain homogenates (10% w/v) from human samples were prepared in ice-cold PBS using a

bead beater. Homogenates were serially diluted from 10^{-2} to 10^{-5} in PBS. RT-QuIC reactions

were set up in a 96-well plate with each well containing 100 µL of reaction mix: 0.1 mg/mL

human αSyn (1-140), 10 µM ThT, 10 mM Tris (pH 7.5), and 50 mM NaCl. 2 µL of diluted

brain homogenate was added to seed the reactions. The plate was incubated in a BMG

FLUOstar Omega plate reader at 37°C with cycles of 1 min shaking (700 rpm, double orbital)

and 1 min rest. ThT fluorescence (Ex: 450 nm, Em: 490 nm) was recorded every 15 min. A

sample was considered positive if the fluorescence of at least two of three technical replicates

crossed a threshold calculated as the mean of negative control wells plus five standard

deviations.

Statistical Analysis

All statistical analyses were performed using GraphPad Prism (v9.0). Data are presented as

mean ± SD unless otherwise noted. Before applying parametric tests, data were assessed for

normality using the Shapiro-Wilk test and for homogeneity of variances using Levene's test.

Comparisons between two groups were performed using a two-tailed, unpaired Student's t-

test. Comparisons among three or more groups were performed using one-way ANOVA

followed by Tukey's post hoc test for multiple comparisons. For data involving two

independent variables, a two-way ANOVA with Tukey's multiple comparisons test was used.

A P value < 0.05 was considered statistically significant. Specific statistical details for each

experiment, including the exact value of n and what n represents, are provided in the figure

legends.

Data availability

433   Any additional information required to reanalyze the data reported in this report is available

434   from the corresponding author upon reasonable request. All materials used in this study will

435   be made available subject to a materials transfer agreement (MTA). This paper does not

436   report the original code.

Acknowledgments

438   This work was supported by grants from the National Natural Science Foundation of China

439   (grant numbers 81870856, 81870992, and 32171236).

Author contributions

441 H.M. designed the study, performed experiments, and wrote the manuscript. Y.K. supervised

442 experiments, analyzed data, and contributed to manuscript preparation. D.W. and H.L.

443 assisted with experimental procedures and data collection. H.M., J.H., and P-Y.X. provided

444 project oversight, funding, and critical revisions to the manuscript.

Competing interests

446 The authors declare that they have no competing interests.

References

Schweighauser, M. *et al.* Structures of alpha-synuclein filaments from multiple system atrophy. *Nature* **585**, 464–469 (2020). https://doi.org/10.1038/s41586-020-2317-6

Yang, Y. *et al.* Structures of alpha-synuclein filaments from human brains with Lewy pathology. *Nature* **610**, 791–795 (2022). https://doi.org/10.1038/s41586-022-05319-3

Rodriguez, J. A. *et al.* Structure of the toxic core of alpha-synuclein from invisible crystals. *Nature* **525**, 486–490 (2015). https://doi.org/10.1038/nature15368

Theillet, F. X. *et al.* Structural disorder of monomeric alpha-synuclein persists in mammalian cells. *Nature* **530**, 45–50 (2016). https://doi.org/10.1038/nature16531

Volpicelli-Daley, L. A. *et al.* Exogenous alpha-synuclein fibrils induce Lewy body pathology leading to synaptic dysfunction and neuron death. *Neuron* **72**, 57–71 (2011). https://doi.org/10.1016/j.neuron.2011.08.033

Thomzig, A. *et al.* Transmissible alpha-synuclein seeding activity in brain and stomach of patients with Parkinson's disease. *Acta Neuropathol* **141**, 861–879 (2021). https://doi.org/10.1007/s00401-021-02312-4

Anderson, J. P. *et al.* Phosphorylation of Ser-129 is the dominant pathological modification of alpha-synuclein in familial and sporadic Lewy body disease. *J Biol Chem* **281**, 29739–29752 (2006). https://doi.org/10.1074/jbc.M600933200

Fujiwara, H. *et al.* alpha-Synuclein is phosphorylated in synucleinopathy lesions. *Nat Cell Biol* **4**, 160–164 (2002). https://doi.org/10.1038/ncb748

Mahul-Mellier, A. L. *et al.* The process of Lewy body formation, rather than simply alpha-synuclein fibrillization, is one of the major drivers of neurodegeneration. *Proc Natl Acad Sci U S A* **117**, 4971–4982 (2020). https://doi.org/10.1073/pnas.1913904117

Lashuel, H. A. *et al.* Revisiting the specificity and ability of phospho-S129 antibodies to capture alpha-synuclein biochemical and pathological diversity. *NPJ Parkinsons Dis* **8**, 136 (2022). https://doi.org/10.1038/s41531-022-00388-7

Ghanem, S. S. *et al.* alpha-Synuclein phosphorylation at serine 129 occurs after initial protein deposition and inhibits seeded fibril formation and toxicity. *Proc Natl Acad Sci U S A* **119**, e2109617119 (2022). https://doi.org/10.1073/pnas.2109617119

Ramalingam, N. *et al.* Dynamic physiological alpha-synuclein S129 phosphorylation is driven by neuronal activity. *NPJ Parkinsons Dis* **9**, 4 (2023). https://doi.org/10.1038/s41531-023-00444-w

Groveman, B. R. *et al.* Rapid and ultra-sensitive quantitation of disease-associated alpha-synuclein seeds in brain and cerebrospinal fluid by alphaSyn RT-QuIC. *Acta Neuropathol Commun* **6**, 7 (2018). https://doi.org/10.1186/s40478-018-0508-2

Shahnawaz, M. *et al.* Development of a Biochemical Diagnosis of Parkinson Disease by Detection of alpha-Synuclein Misfolded Aggregates in Cerebrospinal Fluid. *JAMA Neurol* **74**, 163–172 (2017). https://doi.org/10.1001/jamaneurol.2016.4547

15   Soto, C., Saborio, G. P. & Anderes, L. Cyclic amplification of protein misfolding:
application to prion-related disorders and beyond. *Trends Neurosci* **25**, 390–394 (2002).
https://doi.org/10.1016/s0166-2236(02)02195-1

16   Paciotti, S., Bellomo, G., Gatticchi, L. & Parnetti, L. Are We Ready for Detecting alpha-
Synuclein Prone to Aggregation in Patients? The Case of "Protein-Misfolding Cyclic
Amplification" and "Real-Time Quaking-Induced Conversion" as Diagnostic Tools. *Front
Neurol* **9**, 415 (2018). https://doi.org/10.3389/fneur.2018.00415

17   Concha-Marambio, L., Pritzkow, S., Shahnawaz, M., Farris, C. M. & Soto, C. Seed
amplification assay for the detection of pathologic alpha-synuclein aggregates in
cerebrospinal fluid. *Nat Protoc* **18**, 1179–1196 (2023). https://doi.org/10.1038/s41596-
022-00787-3

18   Rossi, M. *et al.* Ultrasensitive RT-QuIC assay with high sensitivity and specificity for Lewy
body-associated synucleinopathies. *Acta Neuropathol* **140**, 49–62 (2020).
https://doi.org/10.1007/s00401-020-02160-8

19   Sorrentino, Z. A. *et al.* Physiological C-terminal truncation of alpha-synuclein
potentiates the prion-like formation of pathological inclusions. *J Biol Chem* **293**, 18914–
18932 (2018). https://doi.org/10.1074/jbc.RA118.005603

20   Ulamec, S. M. *et al.* Single residue modulators of amyloid formation in the N-terminal
P1-region of alpha-synuclein. *Nat Commun* **13**, 4986 (2022).
https://doi.org/10.1038/s41467-022-32687-1

21   Munishkina, L. A., Henriques, J., Uversky, V. N. & Fink, A. L. Role of protein-water
interactions and electrostatics in alpha-synuclein fibril formation. *Biochemistry* **43**,
3289–3300 (2004). https://doi.org/10.1021/bi034938r

22   Metrick, M. A., 2nd *et al.* Million-fold sensitivity enhancement in proteopathic seed
amplification assays for biospecimens by Hofmeister ion comparisons. *Proc Natl Acad
Sci U S A* **116**, 23029–23039 (2019). https://doi.org/10.1073/pnas.1909322116

23   Giehm, L., Oliveira, C. L., Christiansen, G., Pedersen, J. S. & Otzen, D. E. SDS-induced
fibrillation of alpha-synuclein: an alternative fibrillation pathway. *J Mol Biol* **401**, 115–
133 (2010). https://doi.org/10.1016/j.jmb.2010.05.060

24   Giehm, L. & Otzen, D. E. Strategies to increase the reproducibility of protein fibrillization
in plate reader assays. *Anal Biochem* **400**, 270–281 (2010).
https://doi.org/10.1016/j.ab.2010.02.001

25   Orru, C. D. *et al.* Rapid and sensitive RT-QuIC detection of human Creutzfeldt-Jakob
disease using cerebrospinal fluid. *mBio* **6** (2015). https://doi.org/10.1128/mBio.02451-
14

26   Bhumkar, A. *et al.* Single-Molecule Counting Coupled to Rapid Amplification Enables
Detection of alpha-Synuclein Aggregates in Cerebrospinal Fluid of Parkinson's Disease
Patients. *Angew Chem Int Ed Engl* **60**, 11874–11883 (2021).
https://doi.org/10.1002/anie.202014898

27   Ohgita, T., Namba, N., Kono, H., Shimanouchi, T. & Saito, H. Mechanisms of enhanced

aggregation and fibril formation of Parkinson's disease-related variants of alpha-
synuclein. *Sci Rep* **12**, 6770 (2022). https://doi.org/10.1038/s41598-022-10789-6

28   Lira, R. B., Steinkuhler, J., Knorr, R. L., Dimova, R. & Riske, K. A. Posing for a picture:
vesicle immobilization in agarose gel. *Sci Rep* **6**, 25254 (2016).
https://doi.org/10.1038/srep25254

29   Uemura, N. *et al.* alpha-Synuclein aggregates amplified from patient-derived Lewy
bodies recapitulate Lewy body diseases in mice. *Nat Commun* **14**, 6892 (2023).
https://doi.org/10.1038/s41467-023-42705-5

30   Luk, K. C. *et al.* Pathological alpha-synuclein transmission initiates Parkinson-like
neurodegeneration in nontransgenic mice. *Science* **338**, 949–953 (2012).
https://doi.org/10.1126/science.1227157

31   Tanriover, G. *et al.* Prominent microglial inclusions in transgenic mouse models of
alpha-synucleinopathy that are distinct from neuronal lesions. *Acta Neuropathol*
Commun **8**, 133 (2020). https://doi.org/10.1186/s40478-020-00993-8

32   Kumar, S. T. *et al.* How specific are the conformation-specific alpha-synuclein
antibodies? Characterization and validation of 16 alpha-synuclein conformation-specific
antibodies using well-characterized preparations of alpha-synuclein monomers, fibrils
and oligomers with distinct structures and morphology. *Neurobiol Dis* **146**, 105086
(2020). https://doi.org/10.1016/j.nbd.2020.105086

33   Soto, C. alpha-Synuclein seed amplification technology for Parkinson's disease and
related synucleinopathies. *Trends Biotechnol* **42**, 829–841 (2024).
https://doi.org/10.1016/j.tibtech.2024.01.007

34   Vascellari, S., Orru, C. D. & Caughey, B. Real-Time Quaking- Induced Conversion Assays
for Prion Diseases, Synucleinopathies, and Tauopathies. *Front Aging Neurosci* **14**,
853050 (2022). https://doi.org/10.3389/fnagi.2022.853050

35   Peng, C. *et al.* Cellular milieu imparts distinct pathological alpha-synuclein strains in
alpha-synucleinopathies. *Nature* **557**, 558–563 (2018). https://doi.org/10.1038/s41586-
018-0104-4

36   Peelaerts, W. *et al.* alpha-Synuclein strains cause distinct synucleinopathies after local
and systemic administration. *Nature* **522**, 340–344 (2015).
https://doi.org/10.1038/nature14547

37   Shahnawaz, M. *et al.* Discriminating alpha-synuclein strains in Parkinson's disease and
multiple system atrophy. *Nature* **578**, 273–277 (2020). https://doi.org/10.1038/s41586-
020-1984-7

38   Zhang, S. *et al.* Post-translational modifications of soluble alpha-synuclein regulate the
amplification of pathological alpha-synuclein. *Nat Neurosci* **26**, 213–225 (2023).
https://doi.org/10.1038/s41593-022-01239-7

39   Altay, M. F., Liu, A. K. L., Holton, J. L., Parkkinen, L. & Lashuel, H. A. Prominent astrocytic
alpha-synuclein pathology with unique post-translational modification signatures
unveiled across Lewy body disorders. *Acta Neuropathol Commun* **10**, 163 (2022).

https://doi.org/10.1186/s40478-022-01468-8

40   Okuzumi, A. *et al.* Propagative alpha-synuclein seeds as serum biomarkers for
synucleinopathies. *Nat Med* **29**, 1448–1455 (2023). https://doi.org/10.1038/s41591-
023-02358-9

41   Scialo, C. *et al.* TDP-43 real-time quaking induced conversion reaction optimization and
detection of seeding activity in CSF of amyotrophic lateral sclerosis and frontotemporal
dementia patients. *Brain Commun* **2**, fcaa142 (2020).
https://doi.org/10.1093/braincomms/fcaa142

42   Kraus, A. *et al.* Seeding selectivity and ultrasensitive detection of tau aggregate
conformers of Alzheimer disease. *Acta Neuropathol* **137**, 585–598 (2019).
https://doi.org/10.1007/s00401-018-1947-3

43   Volpicelli-Daley, L. A., Luk, K. C. & Lee, V. M. Addition of exogenous alpha-synuclein
preformed fibrils to primary neuronal cultures to seed recruitment of endogenous
alpha-synuclein to Lewy body and Lewy neurite-like aggregates. *Nat Protoc* **9**, 2135–
2146 (2014). https://doi.org/10.1038/nprot.2014.143

44   Mao, H. *et al.* Optimal Preservation of PFFs in Glycerol Enhances Suitability for Modeling
Parkinson's Disease. *Small* **20**, e2401552 (2024). https://doi.org/10.1002/smll.202401552

45   Polinski, N. K. *et al.* Best Practices for Generating and Using Alpha-Synuclein Pre-
Formed Fibrils to Model Parkinson's Disease in Rodents. *J Parkinsons Dis* **8**, 303–322
(2018). https://doi.org/10.3233/JPD-171248

Figure 1

Fig. 1 | Development of a quiescent system for αSyn seed amplification. a, Schematic

illustrating the kinetics of ThT assays. Conventional SAAs rely on cyclic fragmentation (solid

line) to sustain rapid amplification. In each quiescent phase without fragmentation,

elongation would typically plateau. In contrast, QSAA is designed to maintain continuous

amplification under quiescent conditions (dashed line). **b,** Schematic comparing the

mechanisms of SAA (cyclic fragmentation) and QSAA (quiescent amplification). **c,** Amino

acid sequence alignment of human wild-type (hWT), human A53T, and mouse wild-type

(mWT) αSyn. The bar chart quantifies the lag phase. **d,** ThT fluorescence kinetics of αSyn

aggregation under cyclic fragmentation, assessing the effect of different monomers (MM),

sulfate ions (SO_4^{2-}), and temperature (70°C). Reactions were seeded with 1 pg of PFFs (blue)

or unseeded (orange). **e,** Quantification of the lag phase from experiments in **d**. **f-g,**

Aggregation kinetics and specificity under cyclic fragmentation (**f**) versus quiescent

conditions (**g**); bar charts show PAR (1 pg PFFs / no seed). **h,** Representative TEM images of

the fibrillar products generated by QSAA under optimized quiescent conditions. Insert,

unseeded control. Data are mean ± SD (n=3). *P < 0.05, ***P < 0.001, ****P < 0.0001; one-

way ANOVA with Tukey's post hoc test. Scale bars are indicated.

Figure 2

Fig. 2 | QSAA enables in situ amplification and mapping of αSyn seeding activity in mouse brain sections. a, Schematic of the standardized operating procedure for in situ QSAA on brain tissue, including tissue preparation, reagent preparation, and incubation on a

temperature-controlled stage with an agarose overlay. **b,** Representative images of in situ

QSAA on coronal, sagittal, and horizontal sections from a mouse brain 180 dpi with mPFFs.

Green fluorescence (ThT) indicates αSyn seeding activity. Corresponding anatomical atlas

plates show the signal distribution (red dots). Inset: a PBS-injected control. **c,** TEM of a

QSAA-treated tissue showing fibrillar structures; arrowheads indicate fibrils adjacent to

nuclei. **d,** Coronal sections imaged by fluorescence and brightfield after QSAA incubation;

macroscopic aggregates are shown alongside ThT signal. Images are representative of at least

three independent experiments. Scale bars are indicated.

Figure 3

Fig.3 | QSAA provides superior sensitivity and earlier detection of αSyn pathology in a mouse model. a, Brain-wide maps comparing the distribution of QSAA signal with pS129-IF on consecutive sections from mPFF-injected mouse brain (180 dpi). **b,** Direct comparison of QSAA, pS129-IF, and ThS staining on consecutive coronal sections from a mouse brain

injected with one-tenth the standard mPFFs dose and analyzed at 30 dpi. Colored boxes

indicate magnified regions. **c,** Quantification of relative fluorescence units (RFU) from the

cortical regions shown in **b**. Data are mean ± SD (n=3). **P < 0.01, ****P < 0.0001; one-way

ANOVA with Tukey's post hoc test. **d,** Comparison of QSAA and pS129-IF in the mPFF-

injected mouse model at 7, 10, and 14 dpi. **e,** Brain-wide map of αSyn seeding activity

revealed by QSAA at 14 dpi in mPFF-injected mice compared to PBS-injected controls.

Images are representative of at least three independent experiments. Scale bars are indicated.

Figure 4

Fig. 4 | QSAA detects endogenous seeding activity in human postmortem brain tissue. a, Representative pS129-IF and QSAA results on striatal sections from HC, PSP, PD, and long-term formalin-fixed PD (>20 years; FF-PD). **b,** RT-QuIC of brain lysates from the same cases. **c,** Quantification of the PAR from RT-QuIC data in **b**. Data are mean ± SD (n=3). ****P < 0.0001; one-way ANOVA with Tukey's post hoc test. **d,** Schematic of brain section indicating the regions analyzed in **e**. **e,** Representative QSAA images of the corpus callosum, striatum, and cortex in HC and PD brains. L, left; R, right. **f,** Time-lapse QSAA imaging of a PD brain section during incubation (0-5 h). Scale bars are indicated.

Figure 5

Fig. 5 | IF-QSAA reveals cellular localization of seeding activity and its relationship with pS129 pathology. a, Normalized excitation spectra of ThT and common fluorophores (DAPI, FITC, Cy3, Cy5). **b,** Workflow comparison showing QSAA-first versus IF-first sequence with NeuN staining. **c,** Confocal image of IF-QSAA on striatal sections from mPFF-inoculated mice (180 dpi) co-labeled for NeuN (red) and QSAA (green). Arrow and

arrowhead indicate selected cellular and extracellular structures. **d,** Schematic of the

optimized IF-QSAA protocol. **e, g,** IF-QSAA images of mPFF-injected mouse S2 cortex

sections (30 dpi) co-labeled with NeuN (**e**), IBA1 (**f**), and GFAP (**g**). Insets show magnified

cells. **h, i,** Consecutive S2 cortex sections (30 dpi) stained for pS129 and either NeuN (**h**) or

IBA1 (**i**); DAPI in blue. Arrows in **i** indicate selected pS129-positive neurons. **j,** High-

resolution image showing QSAA (green) with pS129 (red) in the mPFF-model at the injected

site. **k,** Schematic of core-halo arrangement of QSAA and pS129 signals in **j**. **l,** Conceptual

four-quadrant classification model of αSyn aggregates. Images are representative of at least

three independent experiments. Scale bars are indicated.

 Table 1: Key Reagents and Resources

Reagent/Resource	Source	Catalog #	Application/Dilution
Antibodies			
Rabbit anti-pS129-αSyn	Abcam	ab51253	IF (1:1000)
Mouse anti-pS129-αSyn (clone 81A)	Abcam	ab184674	IF (1:1000)
Mouse anti-NeuN (clone A60)	Millipore	MAB377	IF (1:1000)
Rabbit anti-IBA1	FUJIFILM Wako	019-19741	IF (1:1000)
Mouse anti-GFAP	Cell Signaling Technology	3670	IF (1:1000)
Goat anti-Rabbit IgG (Alexa Fluor 488)	Invitrogen	A-11008	IF (1:1000)
Goat anti-Mouse IgG (Alexa Fluor 488)	Invitrogen	A-11001	IF (1:1000)
Goat anti-Rabbit IgG (Alexa Fluor 555)	Invitrogen	A-21428	IF (1:1000)
Goat anti-Mouse IgG (Alexa Fluor 555)	Invitrogen	A-21422	IF (1:1000)
Goat anti-Rabbit IgG (Alexa Fluor 647)	Invitrogen	A-21245	IF (1:1000)
Goat anti-Mouse IgG (Alexa Fluor 647)	Invitrogen	A-21235	IF (1:1000)
Chemicals, Buffers and Reagents			
Thioflavin T (ThT)	Sigma-Aldrich	T3516	40 μM (assays)
Thioflavin S (ThS)	Sigma-Aldrich	T1892	Staining
Na_2SO_4 (Sodium sulfate)	Sigma-Aldrich	238597	800 mM (QSAA buffer)

(NH$_4$)$_2$SO$_4$ (Ammonium sulfate)	Sigma-Aldrich	A4915	Purification
TureBlack Autofluorescence Quencher	Biotium	23007	Staining
Mineral oil	Biosharp	BS927	Overlay (quiescent SAA)
Protease inhibitor	Roche	04693116001	Protein expression/lysis
Endotoxin removal resin	Sangon Biotech	B641718	Endotoxin removal
Plastics and Consumables			
High-adhesion glass slides	CITOTEST	188105W	Section mounting
96-well black, clear-bottom plate	Nunc	165305	Agitated SAA
96-well PP PCR plate	Thermo Scientific	AB0700	Quiescent SAA
Optical adhesive film	Applied Biosystems	4311971	Plate sealing
HiTrap Q HP anion-exchange column	Cytiva	17115401	Protein purification
Amicon Ultra 50-kDa cutoff filter	Millipore	UFC905096	Filtration
10-µL Hamilton syringe, 33-g needle	Hamilton	65460-05	Stereotaxic injection
Instruments and Equipment			
Stereotaxic frame	Kopf Instruments	Model 940	Mouse surgery
ThermoMixer C	Eppendorf	5382	Fibril formation
Non-contact water-bath sonicator	SCIENTZ	SCIENTZ08-III	PFF fragmentation

Temperature-controlled stage	Ruicheng Instrument	TS400	In situ QSAA
Plate reader	Molecular Devices	SpectraMax iD5	ThT kinetics
Real-time PCR instrument	Bio-Rad	CFX96	Quiescent SAA
Plate reader (RT-QuIC)	BMG LABTECH	FLUOstar Omega	RT-QuIC
Inverted microscope	Nikon	ECLIPSE Ti2-E	Imaging
Confocal microscope	Leica Microsystems	SP8	Imaging
Transmission electron microscope	Thermo Fisher/FEI	Talos L120C	TEM
Software and Algorithms			
GraphPad Prism	GraphPad Software	v9.0	Statistics
Fiji/ImageJ	NIH	v2.3.0	Image analysis
NIS-Elements Advanced Research	Nikon	v5.30	Microscope image acquisition
Leica Application Suite X (LAS X)	Leica Microsystems	v3.5.7	Confocal image acquisition

664