Seasonally differential migration strategies, detour, and bottleneck in Common Crane *Grus*

***grus* wintering in western India**

Harindra L. Baraiya[1*], Gaurav Sirola[1], Anju Baroth[1], R. Suresh Kumar[1]

[1]Wildlife Institute of India, Dehradun, India

[*]Corresponding author: hareendra3994@gmail.com

Abstract

Background: Understanding migratory route selection and seasonal migration strategies helps

better plan transboundary conservation. We investigated the seasonal dynamics of migration

behavior in Common Cranes *Grus grus*, a potential sentinel species for understanding and

conserving the Central Asian Flyway. We tracked five individuals from Gujarat, western India, to

breeding sites in south-western Siberia using solar-powered GPS-GSM transmitters. We analysed

18 migration trajectories to investigate the seasonally-explicit migration behaviour, to quantify the

migratory detour and bottleneck, and to identify high-use areas along the migration route.

Results: The results highlighted seasonally differential migration strategies in Common Crane. GLM

analyses confirmed significant seasonal differences: Outward migration covered greater distances,

lasted longer, and was slower, with increased stopover duration and reduced straightness. Detours,

likely undertaken to avoid the Hindukush Himalaya and capitalize resources, increased migration

distances by 23.3% (inward) and 36.8% (outward). Cranes followed distinct seasonal migration

routes that converged in Turkmenistan, creating a migratory bottleneck between 33°N and 35°N.

Conclusion: These findings reveal the adaptive strategies of Common Cranes in response to

seasonal and topographical challenges, emphasizing the urgency of protecting key stopover sites

and migratory corridors.

Keywords: Central Asian Flyway, Movement Ecology, GPS telemetry, Flyway connectivity, Stop-

over site

Background

Avian migration is a periodic, repetitive, and large-scale movement, taking place at the same season

each year, with individuals commuting between breeding and non-breeding areas (Rappole and

Jones 2002; Newton 2011; Åkesson and Helm 2020). Migratory species are exposed to the highest

risk during their migratory journeys (Sillett and Holmes 2002; Rockwell et al. 2017) encountering a

spectrum of natural and anthropogenic challenges during migration. These challenges include

extreme weather changes, habitat alterations, hunting and poisoning along with many other (Kirby

et al. 2008; Gallo-Cajiao et al. 2020; Newth et al. 2022). Certain avian migratory species allocate more

than half of their annual life cycle to traversing between breeding and wintering habitats (Newton

2011). Adversities encountered during these transitions significantly contribute to a considerable

proportion of annual mortality across various species(Sillett and Holmes 2002; Klaassen et al. 2014).

Considering these, extensive studies have been undertaken to understand the patterns, drivers, and

consequences of migratory movements (Vansteelant et al. 2017; Norevik et al. 2020; Gregory et al.

2023), including the evolution of partial migration (B. B. Chapman et al. 2011), stop-over site

ecology (Cai, Huettmann, and Guo 2014; Kölzsch et al. 2016; Arizaga et al. 2018; Elmore et al. 2024),

and the influence of environmental changes on the shifting migratory timings and movements

(Brisson-Curadeau, Elliott, and Côté 2019; Haig et al. 2019; Remisiewicz and Underhill 2020; Davies

et al. 2023; Ozsanlav-Harris et al. 2023).

A flyway concept provides a spatial framework for the management and conservation of migratory

avian species across political boundaries as it encompasses the multiple migratory routes spanning

countries along the entire geographic range (Boere 2006). One such flyway is the Central Asian

Flyway spread across a large continental area of Eurasia between the Arctic and Indian Oceans and

the associated island chains (cms.int). This flyway falls in the Asia-Pacific region where the highest

number of threatened soaring bird species have been reported (Kirby et al. 2008). However, this is

also the flyway that is the least known among all the flyways (Mundkur et al. 2023). The Asian

Flyways are experiencing large-scale threats, including hunting migrant birds (Ahmad et al. 2021;

Hussain and Khan 2021; Mundkur et all. 2023), habitat loss, and urbanization, which will likely have

population-scale impacts on birds (Clark et al. 2013; Malakoff et al. 2016). Understanding a species'

movements and migratory connectivity can help understand the species-specific and route-specific

threats including the spread of infectious diseases that may have population-level consequences

(Webster et al. 2002; Takekawa et al. 2023).

Among the numerous migratory species that travel along the Central Asian Flyway, Common Crane

Grus grus is one of the widespread species that winters in tens of thousands in India. Across the

world, four subspecies of the Common Crane have been recognized: the Western common Crane

G. g. Grus in Europe; the Transcaucasian subspecies *G. g. archibaldi* in the Inner

Caucasus/borderland of Turkey, Georgia, and Iran (V. Y. Ilyashenko et al. 2007; V. Yu Ilyashenko

2008); the Tibetan subspecies *G. g. korelovi* in Tibet, Kyrgyzstan, and Kazakhstan (V. Yu Ilyashenko

and Belyalov 2011); and the Eastern Common Crane *G. g. lilfordi* in Asia from the Ural Mountains

eastward to Northeastern Siberia (Haase and Ilyashenko 2012; Ilyashenko, V.Y. 2013). The Western

and Eastern Common cranes are separated by the Ural Mountains (Mudrik et al. 2015). The

population size of the eastern subspecies is estimated at 125,000–130,000 individuals (Prange 2015),

of which more than 100,000 migrate from Western Siberia and Kazakhstan to the wintering

grounds in India (Higuchi et al. 1994 & 2008; Ram et al. 2023) and Central Asia in the Amudarya

River Valley (Bulyuk and Shamuradov 1995; Rustamov et al. 2011; Sorokin et al. 2011).

The arid plains in the States of Gujarat and Rajasthan in western India support large populations of

the Common Crane that arrive from different parts of their breeding range (Higuchi et al. 1994 &

2008; Ram et al. 2023). Few of the wintering sites witness huge congregations of the cranes and

these are often species-specific flocks numbering in thousands (HB unpublished count data).

Previous studies on the migratory behaviour of the Common Crane have provided important, yet

fragmented, insights into their movement patterns along the Central Asian Flyway. The first

attempt to document Common Crane migration routes was made by Higuchi et al. (1994), who

tracked two individuals from Rajasthan, western India, to southwestern Siberia. A subsequent study

by Higuchi et al. (2008) expanded on this by tracking three additional cranes from Gujarat, western

India, to Siberia. However, both studies were constrained to outward (spring) migration data only,

limited by the technology available at the time, thus offering an incomplete picture of the species'

annual migratory dynamics. Ram et al. (2023) later employed improved tracking technology to

follow two Common cranes from the Saurashtra region of Gujarat, documenting detailed migratory

routes during both inward and outward migrations, as well as movement on wintering grounds.

Nevertheless, these prior studies were limited by single season or single year data, restricting their

ability to characterize migration patterns comprehensively across seasons or to delineate primary

migration corridors and critical stopover sites. Further, these studies did not quantify migratory

detours and bottleneck, crucial parameters to understand the migration ecology and conservation

priorities.

To address these limitations, our study applied state-of-the-art solar-powered GPS-GSM

transmitters to track five Common Cranes from Gujarat across multiple years, capturing fine-scale,

seasonal migratory movements from western India to the Siberian steppes in Russia and

Kazakhstan. This dataset enabled an in-depth characterization of migration parameters, allowing

us to analyze and compare inward and outward migrations, and thus identify any significant

seasonal variation in migratory strategies. Additionally, we investigated migratory detours and

bottlenecks along the migratory path, and mapped high-use areas critical for the conservation of

this species along the Central Asian Flyway. By expanding both the scope and resolution of tracking

data, this study addresses key gaps in our understanding of Common Crane migration ecology and

provides foundational data for prioritizing conservation efforts in high-use regions.

Methods

Study population and GPS tracking

During the winter seasons of 2019-20 and 2021-22, efforts to tag Common cranes were

concentrated in Nal Sarovar Bird Sanctuary (22.8183 N; 72.0439 E, area: 12,000 ha) and Thol Lake

Wildlife Sanctuary (23.1339 N; 72.4001 E, area: 699 ha) in the western Indian state of Gujarat. Both

sanctuaries are designated as Ramsar Sites under the Ramsar Convention and harbor a substantial

population of Common Cranes, estimated to be as high as 30,000 (based on unpublished data from

regular, systematic roost counts by the first author).

We tagged five adult Common cranes with solar-powered GPS-GSM leg-mount transmitters

(OrniTela Inc., Lithuania). The cranes were captured during their arrival at the roosting wetland and

while foraging during daylight hours. In the roosting wetland, mist nets were strategically deployed

at the appropriate angle to the direction of Crane arrival during dusk, and in foraging fields, Cranes

were captured using slip-knot leg nooses. Upon capture, each Crane was immediately extracted

and blindfolded to minimize stress. Subsequently, a 40 g L-40 transmitter was fitted to the tibia of

each Crane, adding approximately 0.93% to the average weight of the tagged cranes. The

deployment of transmitters was completed within a minute, morphological measurements were

taken and the tagged cranes were released within 17–30 minutes after capture. All transmitters

were programmed to record GPS locations at 10-minute intervals, which were then transmitted via

a 2G GSM network.

Data processing, migration parameters, and high-use areas

The transmitter data was downloaded in CSV format and was subsequently visualized in QGIS

3.34.5 to look for any errors in GPS locations during which duplicate and unusually deviated data

points were removed. Trajectories between the breeding and wintering grounds were extracted in

QGIS. The breeding and wintering areas were identified through visual inspection, observing when

the cranes began their characteristic routine movements between the roosting wetlands and

foraging fields within their respective breeding and wintering ranges. In this study, various

parameters were defined and calculated to precisely characterize the migration patterns. The

"departure date" was defined as the date of the first recorded position when a Crane left its

wintering or breeding sites and stopped showing regular movements between the foraging fields

and roosting wetland. Conversely, the "arrival date" was identified as the date when a bird was determined to have reached its wintering or breeding sites and started showing the regular movements between the foraging patches and roosting wetland.

The "migration duration" was calculated as the total time a Crane took to travel, including any stopovers, between the last breeding site and the first wintering site for inward migration, or between the last wintering site and the first breeding site for outward migration. The "migration distance" was measured as the cumulative travel distance in km between all consecutive GPS locations during migration. The "migration speed" was defined as the total migration distance divided by the total duration of the migration, expressed in km/day. The "travel speed" was calculated as the daily distance traveled during the travel days, where the total migration distance was divided by the total travel duration and expressed as km/day. The "travel duration" was defined as the total number of travel days during migration, subtracting any days spent at stopover sites.

The "stopover duration" was the aggregate of all days spent at various stopover sites during each migration season. Further, the "straightness index" was calculated to measure orientation efficiency, defined as the ratio of the straight-line distance 'D' from the migration starting point to the ending point to the actual path length 'L' traveled by the bird. For this, the straight lines were simulated for each migration trajectory and the ratio D/L was calculated to determine an empirical measure of the directness of the migration path.

To assess the extent of longitudinal detours adopted by each individual, we computed the linear distance of each location along the actual migration path from the simulated straight line (D. Liu et al. 2018). The mean longitudinal detours were calculated for each latitude to pinpoint the latitudes exhibiting the highest degree of detour. We identified the migratory bottleneck, characterized by the convergence of all migratory trajectories through a very narrow passage. To determine this, we aggregated all trajectories and calculated the distance between the easternmost and westernmost GPS locations (Supplementary Information) for each latitude. It was anticipated that at the bottleneck, the distance between these points would be minimal. The latitude with the shortest distance was designated as the migratory bottleneck.

Additionally, we utilized Kernel Density Estimates (KDE) (Worton 1989) via ArcGIS Pro to map utilization distribution and identify high-occurrence areas along the migratory flyway. This method allowed us to delineate regions of high utilization during both stopovers and migratory corridors. The final maps were produced using QGIS 3.34.5, while the graphs were generated using R and Microsoft Excel.

Statistical analysis

To assess the impact of migration direction on key migration parameters, a General Linear Model (GLM) was utilized, employing ordinary least squares (OLS) regression with robust methodology. Bootstrapped confidence intervals were generated using 1000 samples and the bias-corrected and accelerated (BCa) method to ensure accurate parameter estimates and address potential deviations from normality in the residuals. Heteroscedasticity-consistent standard errors (HC3) were applied to account for potential violations of homoscedasticity. The model fit was evaluated using R^2. The summary statistics for each migration parameter were calculated and data is presented as mean ± SE and range. The visual comparison of different parameters between outbound and inbound migrations were presented using dot whisker plots where whiskers show 95% CI.

Results

A total of 40,336 GPS fixes were collected attributed to the migratory journeys of five Common Crane individuals across three years (Supplementary table S1). One of the individuals tagged in 2020 completed a full migration cycle (defined as complete inward and outward migration) before data transmission ceased, while the other four individuals tagged in 2022 completed two full migration cycles each, resulting in a total nine complete migration cycles containing 18 trajectories.

Migration timing, routes and stop-over sites

During their outward migration, the birds left India for their breeding grounds between March 18 and April 11, with an average departure date of March 29. They arrived at the breeding sites between April 9 and April 26, averaging an arrival date of April 19. For their return journey, the birds began migrating back to India from southwestern Siberia (55°N - 58°N) between September 28 and October 12, with an average departure date of October 5. They reached Gujarat, in western India (22°N - 23°N), between October 6 and October 25, with an average arrival date of October 16.

The migration route of all tagged Common cranes passed through Pakistan, Afghanistan, northwest Iran, Turkmenistan, Uzbekistan, and Kazakhstan, ending in southwestern Siberia and northern Kazakhstan (Figure 1 and 2). During both migrations, the cranes never crossed the Hindukush Himalayas. Along the route, they also traversed the vast Registan, Margo, and Karakum deserts. On the outward journey, the cranes initially flew in a northwest direction towards the Aral Sea at 45°N latitude. From there, they turned northeast to reach their breeding grounds. However, on two occasions, the cranes maintained a northwest direction until reaching 50°N before heading northeast. Notably, the cranes never crossed 56°E, the western limit of the Ural Mountains of Russia.

During their return migration, the cranes generally avoided flying over the Aral Sea, except on two

occasions, and took a more eastern route, still steering clear of the Hindukush Himalayas. The

tracking data revealed two migration corridors: the outward corridor, which is relatively more

western and crosses the Aral Sea, and the inward corridor, which is more eastern and largely avoids

the Aral Sea. Both corridors converge in Turkmenistan and then merge into a single major corridor

through Afghanistan and Pakistan, leading to western India.

The key stop-over sites identified by KDE include arid deserts, river floodplains, and steppe habitats.

Notable desert stop-over locations include the Karakum Desert near Ashgabat, Mary, and Kerpichli

(Turkmenistan), as well as Tejen and the Hanhowuz Reservoir. Floodplain habitats, particularly the

Amu Darya floodplains was also used by multiple individuals. In the northern steppe regions, cranes

utilize stop-over areas in the northern Aktobe and Kostanay regions of Kazakhstan, as well as in

North Kazakhstan. Additional stop-over sites are situated in the Adamovsky district, Orenburg

Oblast (Russia), and near Taybad (Iran) (Figure 3).

Seasonal differences in migration parameters

The results of the General Linear Model (GLM) analyses highlight significant seasonal differences in

migration parameters between inward and outward migration directions. The model for migration

distance exhibited a strong fit (R^2 = 0.78, $\hat{\beta}$ = 652.67; SE = 90.91, 95% CI = 490.30 - 817.78, p =

<0.0001), indicating that outward migration covers significantly longer distances. Similarly, the

model for migration duration revealed a robust fit (R^2 = 0.68, $\hat{\beta}$ = 10.11; SE = 1.86, 95% CI = 7.05 -

13.72, p = <0.0001), reflecting a notable increase in migration duration for outward migration.

For migration speed, the analysis indicated a strong fit and a significant difference (R^2 = 0.64, $\hat{\beta}$ = -

147.75, SE = 29.47, 95% CI = -97.13 - -5.01, p = 0.00013) where the outward migration was associated

with a marked decrease in speed. The stopover duration also demonstrated a strong model fit (R^2

= 0.61, $\hat{\beta}$ = 6.33, SE = 1.34, 95% CI = 3.91 − 8.65, p = 0.00024), indicating increased resting periods

during outward migration. Furthermore, the analysis of the straightness index (SI) revealed a low

model fit (R^2 = 0.48, $\hat{\beta}$ = -0.05, SE = 0.01 95% CI = -0.07 - -0.02, p = 0.00231), suggesting that outward

migration exhibited a slight reduction in straightness, indicating less direct routes. The travel speed

showed the weakest model fit and no significant difference between the inward and outward

migration (R^2 = 0.20, $\hat{\beta}$ = -60.61, SE = 31.73, 95% CI = -124.03 − 2.81, p = 0.07). The summary statistics

of the key migration parameters are given in Table 1 and supplementary table S2, and are

graphically compared in Figure 4.

Migratory detour and bottleneck

During inward migration, detours commence from the highest latitude of 55°N, where minimal

detour distances (11.44) are observed, and progressively increase as latitude decreases toward the

southern latitudes and peaks at 34°N with the detour of 829.56 km. Beyond this peak, detour

distances decrease at lower latitudes, as the trajectories approach wintering ground. During the

outward migration, the detour distances initially increase gradually, showing moderate detours

from 22°N to 29°N. A more pronounced increase in detour distances is observed between latitudes

29°N and 35°N, with the highest detour distance of 969.32 recorded at 35°N. Post-peak, outward

detours decline toward higher latitudes, as the trajectories approach the breeding ground (Figure

5). The detour resulted in 23.3 % (SE = 1.29, 95% CI = 25.3 – 31.2) and 36.8 % (SE = 1.85, 95% CI =

32.5 – 41.0) increase in the actual distance traveled by the cranes during inward and outward

migrations respectively.

The spatial spread of crane migration trajectories showed a distinct migratory bottleneck between

latitudes 33°N and 35°N, where the longitudinal spread narrows to its lowest values (3.60° at 34°N).

Beyond 35°N, the spread increases, indicating a divergence of migratory paths at higher latitudes

(Figure 6).

Discussion

Migration timing, routes and stop-over sites

The migration timings and routes observed in our study, based on multi-year tracking of five tagged

Common cranes, offer a more comprehensive understanding compared to previous studies, which

tracked fewer cranes over a single season or year. The outward migration timing aligns with prior

research by Higuchi et al. (1994, 2008) and Ram et al. (2023), where cranes departed western India

in March and early April. The inward departure date is consistent with one of the two individuals

tracked in Ram et al. 2023 where the Crane departed from Russia in the last week of September.

However, the second individual from Ram et al. 2023 study departed on 14th August, which is 52

days earlier than the mean departure date recorded in our study, suggesting the individual

variability in departure decision. The multi-year data in our study reveals more consistent patterns

in departure and arrival dates, suggesting stable temporal patterns across years. Further, the

migration timings recorded in our study establish a crucial baseline for Common cranes along the

Central Asian Flyway, providing a foundation for detecting climate-driven phenological shifts. Long-

term studies in Europe and Russia have demonstrated significant advances in Common Crane

migration timing in response to rising temperatures (Filippi-Codaccioni et al. 2011; Kondrakova et
al., 2025), suggesting the role of climate change in altering avian movement patterns. Given the
accelerating impacts of global warming, continued monitoring of migration phenology is essential
for understanding species-level responses and informing conservation strategies.

The broad migration route along crossing Pakistan, Afghanistan, northern Iran, Turkmenistan,
Uzbekistan and finally reaching the breeding ground in south-western Siberia is consistent with
Higuchi et al. 2008 and Ram et al. 2023. While the breeding region of the cranes is also consistent
with Higuchi et al. (1994), the outward migration route differs slightly, as their cranes did not go all
the way to northern Iran and the Aral Sea, flying through northern Afghanistan instead of southern
Afghanistan. This variation stems from the northern Indian origin of their Crane population. The
dual migration corridors observed in our study—western during outward migration and eastern
during return migration—offer new insights into seasonal route flexibility. These differences, likely
arising from the broader temporal and spatial dataset, emphasize the dynamic nature of Common
Crane migrations and the need for long-term tracking to understand these patterns. The migration
distances recorded in our study are consistent with those reported by Higuchi et al. (2008) and one
individual tracked by Ram et al. (2023). However, another individual in Ram et al. (2023) was
recorded traveling 12,982 km during outward migration and 14,873 km during the inward journey.
These distances are 2.5 times (outward) and 3.3 times (inward) greater than those recorded in our
multi-year study. Given the similarity in breeding and wintering sites between studies, this
discrepancy is likely due to differences in analytical approaches.

The identification of key stop-over sites along the Central Asian Flyway highlights the ecological
significance of arid deserts, river floodplains, and steppe habitats in sustaining migratory
populations of Common cranes. The spatial distribution of these stop-over locations suggests a
strong dependence on landscape heterogeneity, wherein cranes utilize resource-rich environments
that provide both foraging opportunities and safe roosting sites. The Karakum Desert, including
sites near Ashgabat, Mary, Kerpichli, and Tejen, serves as a crucial staging area despite its arid
conditions, emphasizing the species' ability to exploit ephemeral water sources and localized
productivity within desert ecosystems (Meine and Archibald 1996). Similarly, the Amu Darya
floodplains function as essential stop-over sites, offering stable wetland as well as food resources in
form of food crops in an otherwise arid landscape, reinforcing the well-documented importance of
riparian corridors in Crane migration (Harris and Mirande 2013). Steppe habitats in northern
Kazakhstan, including the Aktobe and Kostanay regions, as well as in North Kazakhstan, represent
key staging areas that likely support extensive refuelling due to the availability of agricultural fields

and natural grasslands. The prolonged use of these sites aligns with previous studies demonstrating

that Common cranes frequently exploit anthropogenically modified landscapes during migration

(Alonso, Alonso, and Nowald 2008; Li et al. 2013; Erdenechimeg et al. 2023). Furthermore, the

inclusion of Adamovsky district, Orenburg Oblast (Russia), and Taybad (Iran) as stop-over locations

indicates the species' reliance on a broad network of habitat patches to mitigate the energetic

demands of long-distance migration.

The presence of cranes at artificial reservoirs, such as Hanhowuz Reservoir, highlights the potential

role of human-managed water bodies in supplementing natural wetlands, a trend observed in other

migratory waterbirds (Li et al. 2013). Given the increasing anthropogenic pressures on natural

habitats within the Central Asian Flyway (Mundkur et al. 2023), the conservation of these critical

stop-over sites requires a multi-scalar approach integrating habitat protection, sustainable land-use

practices, and transboundary conservation initiatives. Future research should focus on quantifying

habitat selection at these stop-over locations using high-resolution tracking data and remote

sensing to refine conservation strategies and predict the potential impacts of environmental

change on migratory connectivity. Further, the identified stopover sites likely host high densities

of migratory birds, creating potential hotspots for disease transmission. Recent outbreaks of highly

pathogenic avian influenza (H5N1) have caused mass mortality in Common cranes (Braverman

2024; Djurdjević et al. 2024), highlighting the vulnerability of migratory populations to emerging

infectious diseases. Given the risks associated with high-density aggregations, regular disease

monitoring and screening at key stopover sites are essential for early detection and mitigation.

Integrating such measures into transboundary flyway conservation and management plans will be

critical for safeguarding both migratory bird populations and ecosystem health.

Seasonally explicit migration duration

The analysis of migration parameters revealed that outward migration in Common cranes was

significantly slower than inward migration, with longer stopovers contributing to the extended

duration. This contrasts with findings from the Eastern Mongolian Common Crane population

(Erdenechimeg et al. 2023), where outward (spring) migration was faster. Outward migration found

to be faster in the case of many other long-distance migratory species (Nilsson, Klaassen, and

Alerstam 2013), however, these species are predominantly small in size. Faster outward migration

is typically linked to securing better territories, enhanced reproductive success, and improved

readiness for subsequent migrations (Van Noordwijk, McCleery, and Perrins 1995; Kokko 1999; R

2005). Like in our case, the slower outward migration is also reported in many other long-distance

migrants with large body seizes such as White Stork (Shamoun-Baranes et al. 2003), Common Eider
(Mosbech et al. 2006), Black-backed Gull (Bustnes et al. 2013), Bewick's swan (Nuijten et al. 2014),
Booted Eagle (Mellone et al. 2014), Egyptian Vulture (López-López, García-Ripollés, and Urios 2014)
and White-fronted Goose (Kölzsch et al. 2016; Deng et al. 2019). This suggests that the seasonally
differential migration duration is a function of body size and environmental factors.

Given the cranes are long-lived capital-income breeders, the slower outward migration observed in
our cranes may reflect the females' need to accumulate energy reserves for reproduction, a strategy
documented in geese species which is also a large grazing species like cranes (Prop, Black, and
Shimmings 2003; Deng et al. 2019). Additionally, the Cranes may delay migration to coincide with
snowmelt at their northern breeding grounds ensuring that breeding areas are accessible and
favorable upon arrival (Nuijten et al. 2014). This behavior helps cranes optimize their reproductive
success, as arriving too early could mean facing harsh conditions on breeding grounds. The
proximate factors governing stopover duration during outward migration in Central Asian Flyway
Common Crane warrant further investigation.

Migratory detour and bottleneck

This study presents the first quantitative assessment of detours undertaken by Common cranes
along the Central Asian Flyway (CAF), revealing a substantial deviation from the shortest possible
migration route. Earlier, migratory detour has been examined in White-naped Crane *Grus vipio*
where the cranes took detours to capitalise on high-quality habitat (Fujita et al. 2004). Conversely,
Black-necked Crane, an Asian Crane species did not exhibit detour while migrating within the
Tibetan Plateau (Q. Liu et al. 2012). While long-distance migration is inherently constrained by the
energetic demands of sustained flight (Alerstam 2011), migratory birds frequently exhibit route
adjustments to optimize refuelling opportunities and minimize physiological stress and the risk of
predation (Ydenberg, Butler, and Lank 2007; Aharon-Rotman et al. 2016; Norevik et al. 2020) . Our
findings demonstrate that detours increased the total migration distance by up to 41% compared
to the shortest route. The significant increase in migration length observed across latitudes during
both spring and autumn migrations suggests that environmental and topographical factors
strongly influence route selection.

A shortest straight migratory trajectory would necessitate crossing the Hindukush and Kashmir
Himalaya. The complex topography of these high-altitude regions is characterized by turbulent
wind regimes, low atmospheric pressure, and unpredictable thermal availability (Ohlmann 2017;
Galtbalt et al. 2022), conditions that are largely unsuitable for facultative soaring species like

Common Crane. Given that Common cranes also rely on thermal uplift to sustain long-distance flights with minimal energy expenditure, the scarcity of stable thermals in this region likely renders the shortest route energetically expensive. Consequently, the observed detours reflect a strategic avoidance of suboptimal flight conditions, favoring alternative routes that facilitate soaring flight and reduce reliance on energetically costly flapping flight.

The westward deviation observed in both spring and autumn migrations directs the cranes through the arid landscapes of Pakistan, Afghanistan, and Turkmenistan, where extensive desert expanses generate strong and consistent thermal uplift and also provide wind assistance (Galtbalt et al. 2022; Mi et al. 2022). The availability of such atmospheric conditions likely enables Common cranes to employ mixed flying strategy, mitigating the energetic costs associated with increased detour distances. Beyond the benefits conferred by thermal availability, the presence of critical stopover habitats along this route appears to further justify the observed deviations from the shortest path. The agricultural landscapes near the Iran-Afghanistan border, wetland complexes and desert oases in Turkmenistan, and the highly productive agricultural fields of the Amu Darya River delta provide key restring and foraging opportunities, enabling cranes to replenish energy reserves at strategically spaced intervals. The pronounced detours and prolonged stop-over period observed during outward migration further highlight the role of habitat quality in influencing route selection, as cranes must accumulate substantial energy reserves before continuing into breeding grounds. By integrating thermal soaring opportunities with access to high-quality stopover habitats, the observed flyway effectively offsets the energetic costs associated with increased travel distance.

The identified migratory bottleneck along the Iran-Afghanistan border, particularly between latitudes 33°N and 35°N, highlights a critical conservation priority for the Common Crane. This bottleneck possibly results from geographic constraints such as the Aladagh, Binalud and East Iranian Mountain ranges in the West and the high reaches of the Hindukush Himalaya of Afghanistan in the East, forcing cranes to funnel through a narrow corridor. Such natural barriers shape migratory routes by limiting alternative flight paths, concentrating individuals into specific regions (La Sorte et al. 2016). However, this convergence makes the species highly vulnerable to habitat degradation, land-use changes, and direct threats such as hunting even if the birds are not resting or foraging in these areas (Mostafawi, Naseri, and Ostrowski 2017). The consequences of migratory bottlenecks extend beyond localized threats. Because such areas channel a significant portion of a species' population, disturbances can have disproportionate impacts on overall population dynamics (Iwamura et al. 2013). Recognizing bottlenecks as ecological priorities can help integrate their protection into broader conservation strategies, benefiting not only cranes but

also other migratory species that rely on these corridors. By addressing threats at these critical
junctures, conservationists can enhance the resilience of migratory populations and maintain the
ecological integrity of these essential flyways.

Conclusion

Our multi-year tracking study of Common cranes along the Central Asian Flyway (CAF) provides
critical insights into the species' migration timing, routes, and stop-over areas. Notably, this study
is the first quantitative assessment of migratory detours and the identification of a migratory
bottleneck of Common Crane along the CAF. The findings reveal that Common cranes deviate
substantially from the shortest possible migration route, increasing their travel distance by up to
41% to optimize energy expenditure and access high-quality stop-over habitats. This detour
behavior is driven by the avoidance of physiologically demanding ecological barriers, the Hindukush
and Kashmir Himalaya, and the strategic use of thermal soaring conditions in arid landscapes.
Additionally, a migratory bottleneck recorded at the Iran-Afghanistan border, where geographic
constraints funnel all the Crane migration routes, highlights a critical vulnerability in the migratory
flyway of Common Crane. Given the

The study also reinforces the importance of long-term, multi-individual tracking in capturing the
full spectrum of migratory behaviors. The observed temporal consistency in migration schedules,
coupled with individual variability in departure decisions, underlines the dynamic nature of
Common Crane migrations. The identification of dual migration corridors—western during
outward migration and eastern during return migration—further demonstrates seasonal flexibility
in route selection, likely driven by environmental and topographical constraints and breeding
location. Furthermore, the reliance on arid landscapes, river floodplains, and steppe habitats for
stop-over sites highlights the ecological importance of these regions in sustaining migratory
populations.

Conservation implications

Key stopover habitats and a migratory bottleneck along the Iran-Afghanistan border has crucial
conservation implications for Common cranes and other species using the Central Asian Flyway.
Stopover sites such as the Karakum Desert, Amu Darya floodplains, and northern Kazakhstan's
steppe habitats provide essential resources for refueling and resting. Anthropogenic modifications
in the habitats in these areas could have cascading effects on migratory population of not only

Common Crane but other species using the same flyway, necessitating habitat protection and integration into transboundary conservation initiatives.

Beyond habitat threats, disease outbreaks pose an increasing risk to migratory birds. The recent mass mortality of Common cranes due to highly pathogenic avian influenza (H5N1) highlights the vulnerability of birds congregating in large numbers. Given that the stopover sites identified in this study likely host a high abundance of migratory birds, they could serve as hotspots for disease transmission. Regular monitoring and screening at these key locations are essential for early detection and mitigation of emerging infectious diseases. Integrating disease surveillance into conservation strategies will be crucial for maintaining population health and ensuring the stability of the broader migratory network.

Importantly, the Common Crane can serve as a sentinel species for monitoring the Central Asian Flyway, particularly along the desert route passing through Pakistan, Afghanistan, Turkmenistan, Uzbekistan, and Kazakhstan. Its dependence on diverse habitats—arid deserts, river floodplains, agricultural fields, steppe grasslands, and human-managed reservoirs—demonstrates its adaptability to both natural and modified landscapes. The identification of key stopover sites, alongside the quantification of detours and a critical bottleneck, underscores its sensitivity to environmental and topographical constraints. These traits, combined with its wide distribution and long-distance migrations, make the Common Crane an ideal indicator for assessing flyway health and the impacts of environmental change. By prioritizing its conservation, we can safeguard not only this species but also the broader ecological network dependent on this vital corridor.

Acknowledgments

We thank the Director and Dean of the Wildlife Institute of India for their support during the study. We thank the Mr. Shwetank Pandit (Rtd. IFS) and and Mr. V.J. Rana (Rtd. IFS), Sh. P. Purushothama (IFS), Dr. Chaudhary (IFS), Mr. Dipak Solanki (GFS), Mr. Swapnil Patel (GFS) of Gujarat State Forest Department for their support in conducting fieldwork. We thank Mr. Gani Sama and late Mr. Karshan Padhar the frontline staff of the Nal Sarovar Ramsar Site, for their expert help in trapping birds. We thank interns and field support staff for their support in field work. We thank the people of Vadla village for providing logistics while field work. Mention of trade names or commercial products in this publication is solely to provide specific information and does not imply recommendation or endorsement by the authors or the affiliated institution.

Author contributions

HLB, RSK, and AB conceived the study. RSK and AB secured funding. HLB, RSK and GS arranged fieldwork logistics. HLB, GS and RSK participated in fieldwork. HLB conducted data analyses. HLB

445 wrote the original draft of the manuscript. All authors reviewed, edited and approved the final
446 manuscript.

Funding

448 This research was funded by the PowerGrid Corporation of India Ltd. under its Corporate Social
449 Responsibility initiative. The funding agency had no role in study design, data collection, analysis,
450 interpretation, or conclusions.

Conflict of interest

452 The authors declare no conflict of interest.

Availability of data and materials

454 The datasets used and/or analyzed during the current study are uploaded in Move Bank and are
455 kept unavailable for download. The data may be made available from the corresponding author on
456 reasonable written request.

References

Aharon-Rotman, Yaara, Ken Gosbell, Clive Minton, and Marcel Klaassen. 2016. "Why Fly the Extra
Mile? Latitudinal Trend in Migratory Fuel Deposition Rate as Driver of Trans-Equatorial
Long-Distance Migration." *Ecology and Evolution* 6 (18): 6616–24.
https://doi.org/10.1002/ece3.2388.
Ahmad, Tariq, Gul Saba, Nadeem Munawar, Tariq Mahmood, Faraz Akram, and Ammara Baig. 2021.
"Hunting Pressure on Two Migratory Species, Common Crane (*Grus grus*) and Demoiselle
Crane (*Anthropoides virgo*) in Khyber Pakhtunkhwa Province, Pakistan."
Åkesson, Susanne, and Barbara Helm. 2020. "Endogenous Programs and Flexibility in Bird
Migration." *Frontiers in Ecology and Evolution* 8 (March).
https://doi.org/10.3389/fevo.2020.00078.
Alerstam, Thomas. 2011. "Optimal Bird Migration Revisited." *Journal of Ornithology* 152 (S1): 5–23.
https://doi.org/10.1007/s10336-011-0694-1.
Alonso, Javier A., Juan C. Alonso, and Günter Nowald. 2008. "Migration and Wintering Patterns of a
Central European Population of Common Cranes *Grus grus*." *Bird Study* 55 (1): 1–7.
https://doi.org/10.1080/00063650809461499.
Arizaga, Juan, Jaime Resano-Mayor, Diego Villanúa, Daniel Alonso, Juan M. Barbarin, Alfredo
Herrero, Jesús M. Lekuona, and Ricardo Rodríguez. 2018. "Importance of Artificial Stopover
Sites through Avian Migration Flyways: A Landfill-Based Assessment with the White Stork
Ciconia Ciconia." *Ibis* 160 (3): 542–53. https://doi.org/10.1111/ibi.12566.
Boere, G. C. 2006. "The Flyway Concept: What It Is and What It Isn't In Boere GC, Editor;, Galbraith
CA, Editor;, & Stroud DA, Editor.(Eds.), Waterbirds around the World (Pp. 40–47)."
Edinburgh, UK: The Stationery Office.[Google Scholar].
Braverman, Irus. 2024. "Settler Ecologies and More-than-One Health: From Malaria to Avian Flu in
the Hula Valley, Palestine-Israel." *Environment and Planning E: Nature and Space* 7 (3): 977–
93. https://doi.org/10.1177/25148486241232525.

Brisson-Curadeau, Émile, Kyle H Elliott, and Pascal Côté. 2019. "Factors Influencing Fall Departure
Phenology in Migratory Birds That Bred in Northeastern North America." *The Auk*,
November, ukz064. https://doi.org/10.1093/auk/ukz064.

Bulyuk, Viktor N., and Allamurad K. Shamuradov. 1995. "The Migration of the Common Crane, *Grus*
grus, in Southern Turkmenistan." *Zoology in the Middle East* 11 (1): 21–30.
https://doi.org/10.1080/09397140.1995.10637667.

Bustnes, Jan O., Børge Moe, Morten Helberg, and Richard A. Phillips. 2013. "Rapid Long-distance
Migration in N Orwegian L Esser B Lack-backed G Ulls *L Arus Fuscus Fuscus* along Their
Eastern Flyway." Edited by Morten Frederiksen. *Ibis* 155 (2): 402–6.
https://doi.org/10.1111/ibi.12022.

Cai, Tianlong, Falk Huettmann, and Yumin Guo. 2014. "Using Stochastic Gradient Boosting to Infer
Stopover Habitat Selection and Distribution of Hooded Cranes *Grus* Monacha during
Spring Migration in Lindian, Northeast China." Edited by Cédric Sueur. *PLoS ONE* 9 (2):
e89913. https://doi.org/10.1371/journal.pone.0089913.

Chapman, Ben B., Christer Brönmark, Jan-Åke Nilsson, and Lars-Anders Hansson. 2011. "The Ecology
and Evolution of Partial Migration." *Oikos* 120 (12): 1764–75.
https://doi.org/10.1111/j.1600-0706.2011.20131.x.

Clark, Natalie E., Elizabeth H. Boakes, Philip J. K. McGowan, Georgina M. Mace, and Richard A. Fuller.
2013. "Protected Areas in South Asia Have Not Prevented Habitat Loss: A Study Using
Historical Models of Land-Use Change." *PLOS ONE* 8 (5): e65298.
https://doi.org/10.1371/journal.pone.0065298.

Davies, Jacob G., Máire Kirkland, Mark G. R. Miller, James W. Pearce-Higgins, Philip W. Atkinson, and
Chris M. Hewson. 2023. "Spring Arrival of the Common Cuckoo at Breeding Grounds Is
Strongly Determined by Environmental Conditions in Tropical Africa." *Proceedings of the*
Royal Society B: Biological Sciences 290 (2001): 20230580.
https://doi.org/10.1098/rspb.2023.0580.

Deng, Xueqin, Qingshan Zhao, Lei Fang, Zhenggang Xu, Xin Wang, Haoren He, Lei Cao, and Anthony
David Fox. 2019. "Spring Migration Duration Exceeds That of Autumn Migration in Far East
Asian Greater White-Fronted Geese (Anser Albifrons)." *Avian Research* 10 (1): 19.
https://doi.org/10.1186/s40657-019-0157-6.

Djurdjević, Biljana, Tamaš Petrović, Vladimir Gajdov, Dejan Vidanović, Ivana Vučićević, Milena
Samojlović, and Marko Pajić. 2024. "Natural Infection of Common Cranes (*Grus grus*) with
Highly Pathogenic Avian Influenza H5N1 in Serbia." *Frontiers in Veterinary Science* 11
(December):1462546. https://doi.org/10.3389/fvets.2024.1462546.

Elmore, Jared A., David W. Londe, Craig A. Davis, Samuel D. Fuhlendorf, and Scott R. Loss. 2024.
"Associations with Landscape and Local-scale Wetland Habitat Conditions Vary among
Migratory Shorebird Species during Stopovers." *Wildlife Biology* 2024 (2): e01132.
https://doi.org/10.1002/wlb3.01132.

Erdenechimeg, Baasansuren, Gankhuyag Purev-Ochir, Amarkhuu Gungaa, Oyunchimeg Terbish,
Yajie Zhao, and Yumin Guo. 2023. "Migration Pattern, Habitat Use, and Conservation Status
of the Eastern Common Crane (*Grus grus* Lilfordi) from Eastern Mongolia." *Animals* 13 (14):
2287. https://doi.org/10.3390/ani13142287.

Filippi-Codaccioni, Ondine, Jean-Pierre Moussus, Jean-Paul Urcun, and Frédéric Jiguet. 2011.
"Advanced Autumn Migration of the Common Crane *Grus grus* Over Western Pyrenean
Passes." *Acta Ornithologica* 46 (1): 37–45. https://doi.org/10.3161/000164511X589901.
Fujita, Go, Guan Hong-Liang, Mutsuyuki Ueta, Oleg Goroshko, Vladimir Krever, Kiyoaki Ozaki,
Nagahisa Mita, and Hiroyoshi Higuchi. 2004. "Comparing Areas of Suitable Habitats along
Travelled and Possible Shortest Routes in Migration of White-naped Cranes *Grus Vipio* in
East Asia." *Ibis* 146 (3): 461–74. https://doi.org/10.1111/j.1474-919X.2004.00286.x.
Gallo-Cajiao, Eduardo, Tiffany H. Morrison, Bradley K. Woodworth, Alexander C. Lees, Liliana C.
Naves, Ding Li Yong, Chi-Yeung Choi, et al. 2020. "Extent and Potential Impact of Hunting
on Migratory Shorebirds in the Asia-Pacific." *Biological Conservation* 246 (June):108582.
https://doi.org/10.1016/j.biocon.2020.108582.
Galtbalt, Batbayar, Nyambayar Batbayar, Tuvshintugs Sukhbaatar, Bernd Vorneweg, Georg Heine,
Uschi Müller, Martin Wikelski, and Marcel Klaassen. 2022. "Differences in On-Ground and
Aloft Conditions Explain Seasonally Different Migration Paths in Demoiselle Crane."
Movement Ecology 10 (1): 4. https://doi.org/10.1186/s40462-022-00302-z.
Gregory, Killian A., Charlotte Francesiaz, Frédéric Jiguet, and Aurélien Besnard. 2023. "A Synthesis of
Recent Tools and Perspectives in Migratory Connectivity Studies." *Movement Ecology* 11 (1):
69. https://doi.org/10.1186/s40462-023-00388-z.
Haase, Martin, and Valentin Ilyashenko. 2012. "A Glimpse on Mitochondrial Differentiation among
Four Currently Recognized Subspecies of the Common Crane *Grus grus*." *Ardeola* 59 (1):
131–35. https://doi.org/10.13157/arla.59.1.2012.131.
Hafid, Hinda, Moussa Houhamdi, Menouar Saheb, Seddik Sihem, Bouzegag Abdelaziz, Yassine
Nouidjem, Tayeb Bensaci, and Mohamed-Chérif Maazi. 2010. *Cranes Wintering (Grus grus)*
in the High Plains Complex, East Algeria.
Haig, Susan M., Sean P. Murphy, John H. Matthews, Ivan Arismendi, and Mohammad Safeeq. 2019.
"Climate-Altered Wetlands Challenge Waterbird Use and Migratory Connectivity in Arid
Landscapes." *Scientific Reports* 9 (1): 4666. https://doi.org/10.1038/s41598-019-41135-y.
Harris, James, and Claire Mirande. 2013. "A Global Overview of Cranes: Status, Threats and
Conservation Priorities." *Chinese Birds* 4 (3): 189–209.
https://doi.org/10.5122/cbirds.2013.0025.
Higuchi, Hiroyoshi, Sàlim Javed, Meenakshi Nagendran, and Masaki Fujita. n.d. "Spring Migration of
Eurasian Cranes *Grus grus* from Gujarat, India to Their Northern Breeding Grounds."
Hussain, A., and A. A. Khan. 2021. "Wild Birds Trade in Dera Ismael Khan and Bannu Divisions of
Khyber PakhtunKhwa (KPK) Province, Pakistan." *Brazilian Journal of Biology* 83
(September):e247915. https://doi.org/10.1590/1519-6984.247915.
Ilyashenko, V. Y., M. G. Ghasabyan, Y. M. Markin, E. I. Ilyashenko, A. F. Kovshar, and S. V. Winter.
2007. "The Morphology Variability of the Common Crane–*Grus grus* (Linnaeus, 1758)(Aves:
Gruidae)." In *Proceedings of the International Conference "Cranes of Eurasia: Biology and*
Conservation", Moscow, Russia, 1–4.
Ilyashenko, V. Yu. 2008. "The New Form of Crane from Trans-Caucasus." *Russ. J. Ornithol* 17:559–62.
Ilyashenko, V. Yu, and O. V. Belyalov. 2011. "The New Subspecies of the Eurasian Crane–*Grus grus*
Korelovi Ssp. n." *Aves: Gruidae), from Central and Eastern Tien-Shan. Russian Ornithological*
Journal 20:1803–11.

Iwamura, Takuya, Hugh P. Possingham, Iadine Chadès, Clive Minton, Nicholas J. Murray, Danny I. Rogers, Eric A. Treml, and Richard A. Fuller. 2013. "Migratory Connectivity Magnifies the Consequences of Habitat Loss from Sea-Level Rise for Shorebird Populations." *Proceedings of the Royal Society B: Biological Sciences* 280 (1761): 20130325. https://doi.org/10.1098/rspb.2013.0325.

Kirby, Jeff S., Alison J. Stattersfield, Stuart H. M. Butchart, Michael I. Evans, Richard F. A. Grimmett, Victoria R. Jones, John O'Sullivan, Graham M. Tucker, and Ian Newton. 2008. "Key Conservation Issues for Migratory Land- and Waterbird Species on the World's Major Flyways." *Bird Conservation International* 18 (S1): S49–73. https://doi.org/10.1017/S0959270908000439.

Klaassen, Raymond H. G., Mikael Hake, Roine Strandberg, Ben J. Koks, Christiane Trierweiler, Klaus-Michael Exo, Franz Bairlein, and Thomas Alerstam. 2014. "When and Where Does Mortality Occur in Migratory Birds? Direct Evidence from Long-Term Satellite Tracking of Raptors." *Journal of Animal Ecology* 83 (1): 176–84. https://doi.org/10.1111/1365-2656.12135.

Kokko, Hanna. 1999. "Competition for Early Arrival in Migratory Birds." *Journal of Animal Ecology* 68 (5): 940–50. https://doi.org/10.1046/j.1365-2656.1999.00343.x.

Kölzsch, Andrea, Gerhard J. D. M. Müskens, Helmut Kruckenberg, Peter Glazov, Rolf Weinzierl, Bart A. Nolet, and Martin Wikelski. 2016. "Towards a New Understanding of Migration Timing: Slower Spring than Autumn Migration in Geese Reflects Different Decision Rules for Stopover Use and Departure." *Oikos* 125 (10): 1496–1507. https://doi.org/10.1111/oik.03121.

Kondrakova, Kristina D, Sergey N Spiridonov, Alexander V Sharikov, Olga S Grinchenko, Yuri M Markin, Tatyana S Kovinka, and Sergey V Volkov. n.d. "LONG-TERM MIGRATORY CHANGES IN *GRUS GRUS* (GRUIFORMES, AVES) ARRIVAL AND DEPARTURE ON BREEDING GROUNDS IN CENTRAL RUSSIA."

La Sorte, Frank A., Daniel Fink, Wesley M. Hochachka, and Steve Kelling. 2016. "Convergence of Broad-Scale Migration Strategies in Terrestrial Birds." *Proceedings of the Royal Society B: Biological Sciences* 283 (1823): 20152588. https://doi.org/10.1098/rspb.2015.2588.

Li, Donglai, Sihang Chen, Huw Lloyd, Shuyu Zhu, Kai Shan, and Zhengwang Zhang. 2013. "The Importance of Artificial Habitats to Migratory Waterbirds within a Natural/Artificial Wetland Mosaic, Yellow River Delta, China." *Bird Conservation International* 23 (2): 184–98. https://doi.org/10.1017/S0959270913000099.

Liu, Dongping, Guogang Zhang, Hongxing Jiang, and Jun Lu. 2018. "Detours in Long-Distance Migration across the Qinghai-Tibetan Plateau: Individual Consistency and Habitat Associations." *PeerJ* 6 (January):e4304. https://doi.org/10.7717/peerj.4304.

Liu, Qiang, Feng-Shan Li, Paul Buzzard, Fa-Wen Qian, Fan Zhang, Jian-Lin Zhao, Jun-Xing Yang, and Xiao-Jun Yang. 2012. "Migration Routes and New Breeding Areas of Black-Necked Cranes." *The Wilson Journal of Ornithology* 124 (4): 704–12. https://doi.org/10.1676/1559-4491-124.4.704.

López-López, Pascual, Clara García-Ripollés, and Vicente Urios. 2014. "Individual Repeatability in Timing and Spatial Flexibility of Migration Routes of Trans-Saharan Migratory Raptors." *Current Zoology* 60 (5): 642–52.

Malakoff, D., Wigginton, N. S., Fahrenkamp-Uppenbrink, J. & Wible, B. 2016. "Rise of the urban planet." Science 80, 272 (2016) Accessed February 25, 2025. https://www.science.org/content/article/use-our-infographics-explore-rise-urban-planet.

Meine, Curt, and George Archibald. 1996. *The Cranes: Status Survey and Conservation Action Plan.* IUCN/SSC Action Plans for the Conservation of Biological Diversity. Gland, Switzerland: IUCN.

Mellone, Ugo, Javier De la Puente, Pascual López-López, Ruben Limiñana, Ana Bermejo-Bermejo, and Vicente Urios. 2014. "Seasonal Differences in Migration Patterns of a Soaring Bird in Relation to Environmental Conditions: A Multi-Scale Approach." *Behavioral Ecology and Sociobiology*, October. https://doi.org/10.1007/s00265-014-1818-4.

Mi, Chunrong, Xinhai Li, Falk Huettmann, Oleg Goroshko, and Yumin Guo. 2022. "Time and Energy Minimization Strategy Codetermine the Loop Migration of Demoiselle Cranes around the Himalayas." *Integrative Zoology* 17 (5): 715–30. https://doi.org/10.1111/1749-4877.12632.

Mosbech, Anders, Grant Gilchrist, Flemming Merkel, Christian Sonne, Annette Flagstad, and Helene Nyegaard. 2006. "Year-Round Movements of Northern Common Eiders Somateria Mollissima Borealis Breeding in Arctic Canada and West Greenland Followed by Satellite Telemetry." *ARDEA-WAGENINGEN-* 94 (3): 651.

Mostafawi, Sayed Naqibullah, Jalaludin Naseri, and Stephane Ostrowski. 2017. "Hunting for the Wealthiest Threatens Migrating Cranes in Afghanistan." *Oryx* 51 (4): 583–84. https://doi.org/10.1017/S003060531700134X.

Mudrik, E. A., T. A. Kashentseva, P. S. Redchuk, and D. V. Politov. 2015. "Microsatellite Variability Data Confirm Low Genetic Differentiation of Western and Eastern Subspecies of Common Crane *Grus grus* L. (Gruidae, Aves)." *Molecular Biology* 49 (2): 260–66. https://doi.org/10.1134/S0026893315020090.

Mundkur, T., Azhar A., Chaudhary, A., Evans, M., Jia, Y., Koshkina, A., Kumar, R., Nergui, J., Niven, R., Rao, M., Scott, T. and Taq, M.A.. 2023. Central Asian Flyway–Situation Analysis: The status of migratory birds and their habitats and recommendations for the observation. Report by BirdLife International, Cambridge.

Newth, Julia L., Robbie A. McDonald, Kevin A. Wood, Eileen C. Rees, Igor Semenov, Anton Chistyakov, Galina Mikhaylova, et al. 2022. "Predicting Intention to Hunt Protected Wildlife: A Case Study of Bewick's Swan in the European Russian Arctic." *Oryx* 56 (2): 228–40. https://doi.org/10.1017/S0030605320000435.

Newton, Ian. 2011. *The Migration Ecology of Birds.* San Diego: Elsevier Science & Technology.

Nilsson, Cecilia, Raymond H. G. Klaassen, and Thomas Alerstam. 2013. "Differences in Speed and Duration of Bird Migration between Spring and Autumn." *The American Naturalist* 181 (6): 837–45. https://doi.org/10.1086/670335.

Norevik, Gabriel, Susanne Åkesson, Tom Artois, Natalie Beenaerts, Greg Conway, Brian Cresswell, Ruben Evens, Ian Henderson, Frédéric Jiguet, and Anders Hedenström. 2020. "Wind-Associated Detours Promote Seasonal Migratory Connectivity in a Flapping Flying Long-Distance Avian Migrant." *Journal of Animal Ecology* 89 (2): 635–46. https://doi.org/10.1111/1365-2656.13112.

Nuijten, Rascha JM, Andrea Kölzsch, Jan A. Van Gils, Bethany J. Hoye, Kees Oosterbeek, Peter P. De Vries, Marcel Klaassen, and Bart A. Nolet. 2014. "The Exception to the Rule: Retreating Ice

Front Makes Bewick's Swans Cygnus Columbianus Bewickii Migrate Slower in Spring than in Autumn." *Journal of Avian Biology* 45 (2): 113–22.

Ohlmann, Klaus. 2017. "The Wind System in the Himalayas: From a Bird's-Eye View." In *Bird Migration across the Himalayas*, edited by Herbert H. T. Prins and Tsewang Namgail, 1st ed., 219–28. Cambridge University Press. https://doi.org/10.1017/9781316335420.017.

Ozsanlav-Harris, Luke, Malcolm D. Burgess, Geoff M. Hilton, and Stuart Bearhop. 2023. "Changes in the Spatial Patterns of Avian Migrations: Evidence, Mechanisms and Causes." *Diversity and Distributions* 29 (12): 1527–45. https://doi.org/10.1111/ddi.13785.

Prange, H. 2015. "Distribution and Migration of the Common Crane on the West European Flyway." *Cranes Eurasia Biol. Distrib. Captiv. Breed* 5:304–12.

Prop, Jouke, Jeffrey M. Black, and Paul Shimmings. 2003. "Travel Schedules to the High Arctic: Barnacle Geese Trade-off the Timing of Migration with Accumulation of Fat Deposits." *Oikos* 103 (2): 403–14. https://doi.org/10.1034/j.1600-0706.2003.12042.x.

R, MOORE F. 2005. "Stopover Ecology of Intercontinental Migrants: En Route Problems and Consequences for Reproductive Performance." *Birds of Two Worlds: The Ecology and Evolution of Migration*, 251–61.

Ram, Mohan, Devesh Gadhavi, Aradhana Sahu, Nityanand Srivastava, Tahir Ali Rather, Lahar Jhala, Priyank Kapadi, et al. 2023. "Satellite Telemetry Insights into the Winter Habitat Use and Movement Ecology of Common and Demoiselle Cranes." *Birds* 4 (4): 337–58. https://doi.org/10.3390/birds4040029.

Rappole, John H, and Peter Jones. n.d. "EVOLUTION OF OLD AND NEW WORLD MIGRATION SYSTEMS."

Remisiewicz, Magdalena, and Les G. Underhill. 2020. "Climatic Variation in Africa and Europe Has Combined Effects on Timing of Spring Migration in a Long-Distance Migrant Willow Warbler *Phylloscopus Trochilus*." *PeerJ* 8 (March):e8770. https://doi.org/10.7717/peerj.8770.

Rockwell, Sarah M., Joseph M. Wunderle, T. Scott Sillett, Carol I. Bocetti, David N. Ewert, Dave Currie, Jennifer D. White, and Peter P. Marra. 2017. "Seasonal Survival Estimation for a Long-Distance Migratory Bird and the Influence of Winter Precipitation." *Oecologia* 183 (3): 715–26. https://doi.org/10.1007/s00442-016-3788-x.

Rustamov, E. A., N. N. Efimenko, V. V. Marochkina, and A. A. Amanov. 2011. "Dynamics of Common Crane Numbers on Main Flyways and Wintering Grounds in Turkmenistan." *Cranes Eurasia Biol. Distrib. Migr. Manag* 4:364–70.

Shamoun-Baranes, Judy, Anat Baharad, Pinhas Alpert, Peter Berthold, Yoram Yom-Tov, Yoav Dvir, and Yossi Leshem. 2003. "The Effect of Wind, Season and Latitude on the Migration Speed of White Storks Ciconia Ciconia, along the Eastern Migration Route." *Journal of Avian Biology* 34 (1): 97–104.

Sillett, T. Scott, and Richard T. Holmes. 2002. "Variation in Survivorship of a Migratory Songbird throughout Its Annual Cycle." *Journal of Animal Ecology* 71 (2): 296–308. https://doi.org/10.1046/j.1365-2656.2002.00599.x.

Sorokin, A. G., E. N. Lanovenko, Yu M. Markin, M. G. Mitropolski, A. P. Shilina, and E. A. Filatova. 2011. "Common Crane Count at the Wintering Grounds in Uzbekistan in 2011." In *Cranes of Eurasia (Biology, Distribution, Migrations, Management) 4. Proceedings of the International*

Conference *"Cranes of Palearctic: Biology, Conservation, Management (in Memory of Academician PS Pallas)*, 11–16.

Takekawa, John Y., Diann J. Prosser, Jeffery D. Sullivan, Shenglai Yin, Xinxin Wang, Geli Zhang, and Xiangming Xiao. 2023. "Potential Effects of Habitat Change on Migratory Bird Movements and Avian Influenza Transmission in the East Asian-Australasian Flyway." *Diversity* 15 (5): 601. https://doi.org/10.3390/d15050601.

Van Noordwijk, A.J., R.H. McCleery, and C.M. Perrins. 1995. "Selection for the Timing of Great Tit Breeding in Relation to Caterpillar Growth and Temperature." *Journal of Animal Ecology* 64 (4): 451–58. https://doi.org/10.2307/5648.

Vansteelant, Wouter M. G., Judy Shamoun-Baranes, Willem Van Manen, Jan Van Diermen, and Willem Bouten. 2017. "Seasonal Detours by Soaring Migrants Shaped by Wind Regimes along the East Atlantic Flyway." Edited by Jason Chapman. *Journal of Animal Ecology* 86 (2): 179–91. https://doi.org/10.1111/1365-2656.12593.

Webster, Michael S., Peter P. Marra, Susan M. Haig, Staffan Bensch, and Richard T. Holmes. 2002. "Links between Worlds: Unraveling Migratory Connectivity." *Trends in Ecology & Evolution* 17 (2): 76–83. https://doi.org/10.1016/S0169-5347(01)02380-1.

Worton, B. J. 1989. "Kernel Methods for Estimating the Utilization Distribution in Home-Range Studies." *Ecology* 70 (1): 164–68. https://doi.org/10.2307/1938423.

Ydenberg, Ronald C., Robert W. Butler, and David B. Lank. 2007. "Effects of Predator Landscapes on the Evolutionary Ecology of Routing, Timing and Molt by Long-Distance Migrants." *Journal of Avian Biology* 38 (5): 523–29. https://doi.org/10.1111/j.0908-8857.2007.04202.x.

717 **Figures:**

719 Figure 1. A map showing migration trajectories of five tagged Common Cranes between 2020 –

720 2023. (Base map: ESRI Physical)

722 Figure 2. A line graph showing the latitudinal distribution of Common Crane movement

723 trajectories. The different colour shows different tagged individuals.

Figure 3. A map showing high occurrence areas of Common Cranes derived through Kernel Density

Estimation. (Base map: ESRI Physical)

728   Table 1. A table showing the summary statistics of key migration parameters of the five tagged

729   Cranes during inward and outward migration journeys

Parameter	Migration Direction	Mean	Median	SE	Range
Migration Distance (km)	Inward	4378.78	4340.00	51.04	4165.00 - 4593.00
	Outward	5031.44	5046.00	68.86	4638.00 - 5278.00
Migration Duration (days)	Inward	11.56	12.00	0.65	9.00 - 14.00
	Outward	21.67	22.00	1.62	15.00 - 31.00
Straightness Index	Inward	0.78	0.78	0.01	0.74 - 0.81
	Outward	0.73	0.73	0.01	0.69 - 0.78
Stop-over Duration (days)	Inward	0.22	0.00	0.15	0.00 - 1.00
	Outward	6.56	8.00	1.26	0.00 - 12.00
Migration Speed (km/day)	Inward	388.93	381.17	23.15	310.00 - 500.78
	Outward	241.19	239.91	15.37	169.00 - 320.53
Travel Days	Inward	11.33	12.00	0.67	8.00 - 14.00
	Outward	15.11	15.00	0.70	12.00 - 19.00
Travel Speed (km/day)	Inward	398.39	381.17	26.55	324.15 - 551.38
	Outward	337.78	330.87	13.80	275.79 - 394.69

Figure 4. Dot-whisker plots comparing the key migration parameters of the five tagged Common

Cranes. The whiskers show 95% confidence interval.

Figure 5. A graph showing the seasonally explicit spatial patterns in the migratory detour, visualised

by fitting smoothed LOESS lines. The values of detour are in km.

Figure 6. A figure showing spatial spread (depicted as a longitudinal difference (in km) between the

west-most and the east-most location in each latitude) in Common Crane migration trajectories.

The lowest spread is observed at 34°N, making it a migratory bottleneck.

741 **Supplementary material**

742 Table S1: A table showing basic information of tagged Common cranes

Sr. No.	Name of the individual	Date of tagging	Capture and Tagging location	Weight (kg)	No. of locations used in the study
1	Vadla	12-03-2020	Vadla Wetland (22.922671 N; 71.958432 E)	4.7	3649
2	Gani	10-02-2022	Thol Ramsar Site (23.139794 N; 72.405048 E)	4.3	8283
3	Sanand	22-02-2022	Thol Ramsar Site (23.135603 N; 72.401336 E)	4.5	7759
4	Nal	04-03-2022	Vadla Wetland (22.918991 N; 71.96749 E)	4.7	9289
5	Bhal	05-03-2022	Vadla Wetland (22.910345 N; 71.984059 E)	4.6	11356

744 Table S2: A table showing the migration parameters of the tagged Common cranes

Migration Direction	Individual	Migration Distance (km)	Migration Duration (days)	Straightness Index	Stop-over Duration (Days)	Migration Speed (km/day)	Travel Days	Travel Speed (km/day)
Inward	Bhal_1	4593.00	14.00	0.81	0.00	328.07	14.00	328.07
	Bhal_2	4340.00	14.00	0.80	1.00	310.00	13.00	333.85
	Gani_1	4295.00	11.00	0.80	0.00	390.45	11.00	390.45
	Gani_2	4165.00	10.00	0.78	0.00	416.50	10.00	416.50
	Nal_1	4507.00	9.00	0.78	0.00	500.78	9.00	500.78
	Nal_2	4214.00	13.00	0.79	0.00	324.15	13.00	324.15
	Sanand_1	4574.00	12.00	0.76	0.00	381.17	12.00	381.17
	Sanand_2	4411.00	9.00	0.76	1.00	490.11	8.00	551.38
	Vadla	4310.00	12.00	0.74	0.00	359.17	12.00	359.17
Outward	Bhal_1	5046.00	26.00	0.78	10.00	194.08	16.00	315.38
	Bhal_2	5240.00	31.00	0.77	12.00	169.03	19.00	275.79
	Gani_1	5016.00	19.00	0.74	2.00	264.00	17.00	295.06
	Gani_2	5278.00	22.00	0.74	8.00	239.91	14.00	377.00
	Nal_1	4963.00	23.00	0.73	8.00	215.78	15.00	330.87
	Nal_2	5163.00	23.00	0.72	8.00	224.48	15.00	344.20
	Sanand_1	4638.00	17.00	0.72	5.00	272.82	12.00	386.50
	Sanand_2	5131.00	19.00	0.70	6.00	270.05	13.00	394.69
	Vadla	4808.00	15.00	0.69	0.00	320.53	15.00	320.53

745