Root exudation facilitates water infiltration and rewetting of dry soil

Emma Gomez Peral [a], Andrew Mair [a], Iker Martin Sanchez [a], Gloria de las Heras
Martinez [a], Mariya Ptashnyk [b], Lionel X Dupuy [a c †]

[a] Department of Conservation of Natural Resources, Neiker, Derio, 48160,
Spain

[b] Department of Mathematics, School of Mathematical and Computer Sciences,
Heriot-Watt University, Scotland, United Kingdom

[c] Ikerbasque, Basque Foundation for Science, Bilbao, 48009, Spain

[†] Corresponding author: ldupuy@neiker.eus

Keywords

Root architecture, exudates, water infiltration, drought

Abstract

The way plant roots facilitate water infiltration in soil may be just as important as
the efficiency with which the root system in turn extracts it from the soil. Here
we studied the mechanisms through which the root system facilitates water
infiltration through a dry soil layer. Dye tracing experiments were conducted in
model soil microcosms to characterise how root growth and exudation affects
the permeability of dry layers of the model soil. Results showed that dissolved
root exudates may be the primary facilitator of water infiltration, which may be
linked to water surface tension. We conclude that in dry soil, root architecture
and root exudation may combine to facilitate the infiltration of water and
decrease the water lost by evaporation. These findings could enhance our
understanding of the traits that provide drought resistance in crops.

Introduction

Hydrologists have long acknowledged the significant role of vegetation in
influencing water movement in soil, and as Clothier and Green aptly expressed,
the roots are the "big movers" (Clothier and Green 1997). A primary driver is the
extraction of water from the soil through root uptake and subsequent
transpiration, which not only reduces soil water content but also causes water to
move along gradients of the water potential. However, prior to uptake, the roots
also influence water transport through modifications of the soil hydraulic
properties. The presence of plant roots in soil has been linked to preferential
flow of water in dye-tracing experiments. For example, Jačka et al. (2021)
observed increased dye concentration around plant roots in a tracer
experiment. This finding was corroborated by measurements of saturated
hydraulic conductivity in soil column experiments, which showed that the soil
conductivity increased due to the presence of roots (Leung et al. 2018). Various
studies have also demonstrated that the hydraulic conductivity increases with
the root length density (Xiao et al. 2024). In field experiments, it was also
observed that most types of vegetation increase the infiltration rate of bare soil
(Marshall et al. 2014; Song et al. 2017).

Changes in soil hydraulic properties can be attributed to physical modifications in the soil's pore structure. As roots grow, they displace the surrounding soil, creating a cavity and rearranging the distribution of the surrounding soil pores (Dexter 1987). Because the size and connectivity of soil pores determine the resistance of the flow of water, these changes in soil porosity in turn affect the permeability of the soil (Schulz et al. 2019). In the study of (Anselmucci et al. 2021) roots grown in sand created an increase in porosity because of dilation. In more structured soil however, the expansion of the root cavity compacted the surrounding soil, and this can lead to a reduction in the soil porosity (Bruand et al. 1996; Helliwell et al. 2017; Koebernick et al. 2019), although this usually occurs in the bulk soils which initial porosity is low (Lucas et al. 2019). Other factors such as the shrinking or decay of the root can also create macropores (Ni et al. 2019; Duddek et al. 2022) where the flow of water can occur with least resistance.

Chemical modifications of the soil solution caused by rhizodeposition can influence water movement in soils as well. High molecular weight polymers such as mucilage increase the viscosity of the soil solution (Read et al. 1999) which in turns reduces the flow speed and increases the resistance to water movements. Mucilages behave as hydrogels because they contain polymers that are hydrophilic, crosslinked, and possess the ability to absorb and retain a significant amount of water (Naveed et al. 2019). This property is critical to maintain moisture around the root when the soil is drying (Carminati et al. 2010). Once dry however, root rhizodeposits can become water repellent and may slow down water infiltration during rewetting (Ahmed et al. 2016). Rhizodeposits also contain surfactants, and this likely facilitates water movement during infiltration and the rewetting of dry and water repellent soils (Read et al. 1999, 2003).

This study aims to determine whether physical or chemical modifications of the soil substrate have a greater impact on soil water infiltration. We carried out dye tracing experiments in model transparent soil microcosms containing dry hydrophobic layers and combined image analysis and mathematical models to quantify the respective role of root exudation and root growth on the permeability of the dry transparent soil layer.

Material and methods

Plant material and germination

Seeds of winter wheat (*Triticum aestum var. Filon*) were first hydrated for one hour in sterilised water and then sterilised for 15 minutes in a 1% calcium hypoclorite solution (042548.30, Thermo Scientific, Spain). After numerous washes, seeds were placed in distilled water for two more hours before transfer to agar plates in an incubator at 23 ºC. The incubation time was three days when running microcosm experiments and five days for root exudate extraction experiments.

Extraction of root exudates

Seedlings whose radicles had reached 3-5 cm were placed on a 3D printed
punched plate (5 mm diameter holes arranged on a regular grid with 2 cm
spacing). Each plate hosted approximately 50 seeds and was placed on top of a
15×21×7 glass dish (Ikea, Spain) with roots dipping into 1 L of Hoagland
nutrient solution (092621822, MP Biomedicals, Spain) adjusted to a pH of 7
using potassium hydroxide (KOH). Plants grew in a growth chamber (23 ºC,
60% humidity with 14/10 h day/night cycle) with the nutrient solution aerated
using a pneumatic pump (Souslow, Spain) during seven days. To suppress the
growth of contaminants and avoid root responses to light, the glass dish was
painted in black on the outside. After six days of growth, the nutrient solution
was replaced with 0.5 L of sterile distilled water. Roots grew 24 h in distilled
water before three 50 ml falcon tubes (EP0030122232, Sigma-Aldrich, Spain)
were used to collect 120 mL of solution from each glass dish. The root exudate
solutions were then frozen at -80 ºC. To adjust the concentration of root
exudates, falcon tubes containing root exudate solutions were freeze-dried and
2 mL of distilled water were subsequently added before pooling all the root
exudates from one glass dish into a 15 mL falcon tube (EP0030122194, Sigma-
Aldrich, Spain). The root exudate solution obtained was freeze-dried a second
time for weighing, and distilled water was added to adjust the concentration of
the root exudate solution to 1 mg mL^{-1}. Root exudates were then stored frozen
at -20 ºC. We ran 4 experiments with a total of 8 plates.

Transparent soil

Transparent soil was prepared as described earlier (Downie et al. 2012). Briefly
Nafion pellets (NR50 1100, Ion Power Inc, USA) were fractured to a texture
similar to sand (0.25 to 1.25 mm) using a freezer mill (6850 Freezer/Mill, SPEX
CertiPrep, UK) and a series of sieves. The pH of the substrate was then
adjusted by washing with Hoagland nutrient solution. With a density for Nafion
of approximately 1.9 g cm^{-3} and a porosity of 0.3, the estimated substrate
density is 1.58 g cm^{-3}.

Microcosm preparation

Microcosm chambers were assembled following the protocol described by Liu et
al. (2025) from microscope glass slides (76 × 26 × 1 mm^3, VWR, Spain). Glass
slides were bonded by 4 mm polydimethylsiloxane spacer (PDMS, SYLGARD
184, Sigma-Aldrich, Spain) following surface treatment by oxygen plasma for 15
s at 100 W (HPT-100, Henniker Plasma, UK). After surface treatment, the glass
and PDMS surfaces were brought together, and pressure was applied to ensure
a strong bond between all surfaces in contact. Following the assembly of the
different part, the space available for placing plants and soil is 68 × 18 × 4 mm^3.
The microcosm was then filled with transparent soil layers with different water
content (Figure 1A). The bottom 1.5 cm layer (approximate dry weight of 1.7 g)
of the microcosm chamber was filled with transparent soil fully saturated with
water. Then a 0.8 cm layer of dry transparent soil was added (approximate dry
weight of 0.9 g). The dry transparent soil was obtained by heating in an oven at

105 ºC for 24 hours. The top layer was 1 cm thick (approximate dry weight of
1.1 g) of transparent soil fully saturated with water.

Dye tracing experiment

First, we studied how the presence of plant roots affected the permeability of
the dry transparent soil layer. Control samples consisted of microcosm
chambers prepared as described in the previous section but without seedlings
transferred into them. Wheat seedlings were transferred when the root length
was 0.5 to 0.8 mm in length. All microcosm chambers were closed with parafilm
tape (HS234526B, Sigma-Aldrich, Spain) and placed in a growth chamber at 23
ºC, 60% humidity with 14/10 h day/night cycle. Distilled water was added to the
samples using a syringe to maintain the top layer of transparent soil saturated.
The tracer dye was added 8 days after the transfer of the root seedlings. Most
plants had at least one root in the bottom of the microcosm chamber at that
time. A food dye (red food colorant, Vahiné, Spain) was diluted to 0.1 mL mL^{-1} to
obtain the tracer dye solution. 0.2 mL of that solution was then added on the top
of the microcosm chamber. Half an hour later, the first image was taken (t=0
hour). The following images were taken 24, 48 and 72 hours after the first
image. The experiment was repeated 3 times with a total of 15 samples with
roots and 13 controls samples without roots. Two root samples were discarded
because of leakage of the microcosm chamber.

In the second experiment, we quantified the effect of root exudates on the
infiltration of water. In this case the tracer dye solution was diluted with the root
exudate solution so that resulting root exudate concentration was approximately
0.9 mg mL. Then 0.2 mL of that solution was then added at the top of the
microcosm (concentration of root exudate approximately 0.16 mg g^{-1}). All
microcosm chambers were sealed with parafilm tape and placed in the growth
chamber under the previously described conditions. The first images were
obtained half an hour after inoculation (t=0 hour). The following images were
obtained 72, 96 and 120 hours after the first image. The experiment was
repeated 3 times with a total of 19 samples containing root exudates and 16
control samples containing only water. One control sample was discarded due
to a leak.

In a third experiment, we studied the effect of the presence of root exudates in
the dry transparent soil layer. To prepare dry transparent soil containing root
exudates, 5 g of wet transparent soil was mixed 1 mL of the root exudate
solution. The wet transparent soil was then left to dry at 105 ºC for 48 hours in
an oven. Therefore, the resulting root exudate concentration in the dry
transparent soil was similar to that of the top layer, 0.633 mg g^{-1}. Four types of
samples were prepared. Namely, samples that did not contain root exudates
(NN), samples that contained root exudates in the middle layer (NE), samples
that contained root exudates in the top layer EN), and samples that contained
root exudates in both layers (EE). All microcosm chambers were sealed with
parafilm tape and placed in the growth chamber. The first image was acquired
half an hour after introduction of the tracer dye (t=0 hour). The following images

were taken 72, 96 and 120 hours after the first image. The experiment was
repeated 3 times with a total of 8 samples for each combination.

Finally, in the fourth experiment, we studied the effect of the rearrangement of
transparent soil particles induced by penetration of the dry middle layer. A
hypodermic needle (12684536, Fisher Scientific, Spain) was pushed
approximately 2.5 cm into the transparent soil. The needle was 0.8 mm in
diameter which is similar to the diameter of wheat roots (Hallam et al. 2021).
Control samples were not penetrated by the hypodermic needle. 0.2 ml of the
tracer dye solution was added at the top of the microcosm chamber. All
microcosm chambers were sealed with parafilm tape and placed in the growth
chamber. Half an hour after inoculation of the tracer dye, the first image was
taken (t=0 hour). The following images were taken 24, 48 and 120 hours after
the first image. The experiment was repeated 3 times with a total of 15 samples
and 15 controls. One sample was discarded due to a leak.

Quantitative image analysis

Images of microcosm chambers were acquired using a Canon EOS 2000D
camera (Canon, Spain) with an exposure time of 1/1000 s, ISO-400 speed, and
focal length was 39 mm and a f-stop of f/5.6. Colour images were stored
uncompressed in colour format (sRGB) at resolution of 6000 × 4000 pixels. To
quantify the tracer dye concentration from image data, calibration samples with
dye concentration of 0, 0.01, 0.05, 0.1, 0.5 and 1 mL mL^{-1} were prepared in
triplicate. Images were acquired for each sample and relationships were
established between the dye concentration and the image brightness, and
between dye concentration and the image saturation value (Figure 1B). To
quantify dye concentration in the microcosm chambers containing transparent
soil layers, regions of the image corresponding to the top, middle and bottom
layers of transparent soil were cropped from the raw image data. Images were
transformed to hue, saturation and brightness (HSB) format and the saturation
value was used to determine the dye concentration (Figure 1C). All image
processing tasks were performed using the ImageJ software (Schmid et al.
2010).

The permeability of the dry transparent soil layer k [s^{-1}] was calculated using the
following formula
205

$$k = \frac{(C_2 - C_1) \ / \ (C_0 - C_1)}{T_1 - T_0}. \tag{1}$$

Here, C_0 and C_1 are the dye concentrations at time T_0 in the top and bottom
layer respectively. C_2 is the dye concentration in the bottom layer at time T_1.

Statistical analysis

Statistical models were constructed to analyse the effects of plant roots, root
exudates or to analyse the effect of the needle on both the permeability *k* and
the tracer dye concentration in the dry transparent soil layer. We used
generalized linear models to correlate the permeability of the dry transparent

soil layer with both time and treatment effects. We tried different variance
functions and found that heterogeneity was best corrected using variance
covarying with treatments rather than with time. Model selection was based on
the Akaike information criterion (AIC). Statistical analyses were performed with
R (R Core Team 2021) using the glm package.

Model of water infiltration and rewetting

Water transport in the transparent soil microcosms was represented using the
model of Mair (unpublished). The model solves Richards equations (Richards,
1931), on a spatial domain which here is in 1D, $x \in [-3.3, 0]$, $t \in (0, T]$. Final
time was T=5 days (120 hours).
To match the experimental conditions, the pressure head was adjusted to -0.01
kPa -0.1 cm at the top -100 kPa in the middle layer and -0.01 kPa at the
bottom. The water content and hydraulic conductivity functions were expressed
from the soil pressure head using the formulations of (van Genuchten 1980)
and (Mualem 1976):

$$\theta(h, c_W, c_D) = \theta_r + (\theta_s - \theta_r)S_e(h, c_W, c_D),$$ (2)

$$S_e(h, c_W, c_D) = \frac{1}{(1 + |\alpha(c_W, c_D)h|^n)^m},$$

$$K(h, c_W, c_D) = K^* + \alpha_K K_s K_r(h, c_W, c_D).$$ (3)

Both the Nafion particles and root exudates are known to have different
behaviour during wetting and drying (hysteresis in both the hydraulic
conductivity and water retention functions), and this was modelled represented
using the model of Kool and Parker (1987) for water content θ ($\mathrm{cm^3 cm^{-3}}$) and
the model of Vogel and Zhang (1996) for the hydraulic conductivity K ($\mathrm{cm\,d^{-1}}$).
Hence, we focused primarily on the influence of the concentration of root
exudates suspended in the solution c_W ($\mathrm{mg\,mL^{-1}}$) and the concentration of root
exudates dried and attached to the surface of the particles c_D ($\mathrm{mg\,g^{-1}}$). These
parameters affect the inverse air entry pressure head α ($\mathrm{cm^{-1}}$), as described in
(Karagunduz et al. 2001), through the surface tension of water γ ($\mathrm{mN\,m^{-1}}$) and
water contact angle ω (rad)

$$\alpha^W = \alpha_0^W \times \frac{\gamma(c_W)}{\gamma_0} \times \frac{\cos(\omega_0)}{\cos(\omega(c_D))},$$ (4)

$$\alpha^D = \alpha_0^D \times \frac{\gamma_0}{\gamma(c_W)},$$

239 Here α^W and α^D are the inverse air entry pressure head for wetting and drying
240 respectively and α_0^W and α_0^D are the default inverse air entry pressures if no
241 exudates are present in either form. The effect of root exudates is also
242 incorporated into the saturated hydraulic conductivity as

$$K_s^Z(c_W, c_D) = K_{s,0}^Z \times \left(\frac{\gamma_0}{\gamma(c_W)}\right)^\beta \times \frac{\cos(\omega(c_D))}{\cos(\omega_0)},$$ (5)

where $Z = W, D$ for wetting or drying. Here $K_{s,0}^Z$ are the default saturated
hydraulic conductivities during wetting and drying if no exudates are present,

and β determines the extent to which an exudate-induced reduction in surface
tension increases hydraulic conductivity. The terms K^* and α_K in (3) are
included to effectively incorporate hysteresis. The expressions for surface
tension and contact angle as functions of the concentrations of suspended and
dried root exudates are fitted to the data of Read et al. (2003) and Zickenrott et
al., (2016) respectively,

$$\gamma(c_W) = 47.5 + \frac{72.86-47.5}{\left(1+e^{18.79(c_W-0.36)}\right)}, \tag{6}$$

$$\omega(c_D) = \frac{3.529}{18}\pi(1 + e^{-1879.14c_D})^{-4.39}.$$

251 The dynamics of the root exudates in solution and the dried root exudates are
252 given by the following differential equations

$$\partial_t(\theta c_W) - \partial_x(\theta D_W \partial_x c_W - \vec{q}c_W) = \rho\kappa_W c_D - \theta\kappa_D c_W, \tag{7}$$

$$\rho\partial_t c_D = \theta\kappa_D c_W - \rho\kappa_W c_D,$$

where D_W $(\text{cm}^2\,\text{d}^{-1})$ is the coefficient of diffusion of root exudates through soil
water, $\vec{q}$ $(\text{cm}\,\text{d}^{-1})$ is the advective flux of soil water and ρ $(\text{mg}\,\text{cm}^{-3})$ is the bulk
density of Nafion. The terms κ_W and $\kappa_D\,(\text{d}^{-1})$ are the rates at which dried root
exudate joins the soil water solution and root exudate in solution dries to the
pore surface respectively. Finally, the movement of dye within the chambers is
modelled using an additional differential equation

$$\partial_t(\theta u) - \partial_x(\theta D_u \partial_x u - \vec{q}u) = 0. \tag{8}$$

Here u $(\text{mL}\,\text{mL}^{-1})$ is the concentration of dye in solution and D_u $(\text{cm}^2\text{d}^{-1})$ is the
diffusion coefficient of the dye in solution. The dye concentration at the start of
the experiment is 0.04 $\text{mL}\,\text{mL}^{-1}$ in the upper layer where root exudates have
been introduced, 0 otherwise.

A finite element scheme, with an implicit Euler discretisation in time, was used
to approximate solutions to the coupled system of equations.

Model parameterisation

Model parameters were taken from previously published literature assuming
that coarse sand would be the best approximation of the transparent soil.
However, the reference inverse air-entry pressure heads α_0^D, α_0^W and saturated
hydraulic conductivities $K_{s,0}^D, K_{s,0}^W$ were altered considerably from the values cited
in the literature in order to reflect the hydrophobicity of dry Nafion that was
observed in the control infiltration experiments. The full list of parameters can
be found in Table 1.

Parameter	Value	Source
$\boldsymbol{\theta_{r,0}}$	1×10^{-10} $\text{cm}^3\text{cm}^{-3}$	$N\backslash A$
$\boldsymbol{\theta_{s,0}}$	0.404 $\text{cm}^3\text{cm}^{-3}$	(Benson et al. 2014)
$\boldsymbol{\alpha_0^D}$	0.049 cm^{-1}	(Benson et al. 2014)

α_0^W	$0.164\ \mathrm{cm^{-1}}$	(Benson et al. 2014)
n	3.16	(Benson et al. 2014)
m	0.684	(Benson et al. 2014)
$K_{s,0}^D$	$8.196 \times 10^{-4}\ \mathrm{cm\,d^{-1}}$	(Carsel and Parrish 1988)
$K_{s,0}^W$	$4.917 \times 10^{-4}\ \mathrm{cm\,d^{-1}}$	(Carsel and Parrish 1988)
D_W	$0.65\ \mathrm{cm^2 d^{-1}}$	(Scott et al. 1995), average taken
ρ	$1.9\ \mathrm{g\,cm^{-3}}$	(Oberbroeckling et al. 2002)
D_u	$0.5\ \mathrm{cm^2 d^{-1}}$	(Syms 2017)

Table 1: Model parameter values dervied from the scientific litterature

Two types of critical parameters could not be determined from the literature.
First, the coefficient linking surface tension to hydraulic conductivity β is both
very specific to the model system and could not be found in the literature. To
calibrate the value of β, data was used from the second experiment and
compared observations to simulated dye concentration between depths $-3.3\ \mathrm{cm}$
and $-1.8\mathrm{cm}$ at times $t = 3, 4, 5\ \mathrm{d}$. Using Bayesian optimisation (Brochu et al.
2010) we obtained the values β that most accurately matched experimental
results. The rates at which root exudates dissolve or reverse to a dry state are
also very specific to our model system and were obtained in a similar way. Here
we used the observation from the third experiment and Bayesian optimisation to
determine the values of κ_W and κ_D that best predicted dye concentrations in the
lower third of the domain.

Figure 1: Model system for quantifying the impact of root exudates on water transport (A) Diagram of the microcosm system. At the start of the experiment, the microcosm is comprised of three layers of transparent soil (left). The top layer of transparent soil is saturated with water to allow fast diffusion of the dye. The middle layer is oven dried to produce a hydrophobic barrier. The bottom layer is saturated with water for precise determination of the quantity of dye moving through the dry layer. During the experiment (right) the dye was introduced at T0 at the top of the microcosm chamber and diffused rapidly through the top layer of transparent soil. At time T1 the tracer dye became visible, and a permeability coefficient could be calculated. (B) The dye concentration in the liquid can be quantified from the image saturation value (red) at low dye concentration, and with the image brightness value (green) at high concentration. (C) Example of time lapse data of the saturation value captured in the bottom transparent soil layer.

Results

Plant roots increase the infiltration of water through dry transparent soil in a microcosm system

Following introduction into the microcosm, the dye was observed spreading non homogeneously but rapidly within the top transparent soil layer (Figure 2A), but there was no evidence of the dye being absorbed by the particles. The dry

transparent soil layer blocked further spreading of the tracer dye into the lower layers of transparent soil. There was no obvious effect of the roots on the transport of the dye at this stage of the experiment. The spatial distribution of the tracer dye was analysed over a 72-hour period following the start of the experiment (Figure 2B). In the control samples, the wetting front advanced downward by 1 to 2 mm during the first 48 hours. We did not observe further progress of the wetting front during the rest of the experiment, even in the case where the tracer dye was seen in the bottom transparent soil layer (Figure 2B). 24 hours after the introduction of the tracer dye, no coloration was observed in the bottom transparent soil layer of the control samples. Only 5 samples out of 13 showed some coloration at the end of the experiment. In the samples containing roots, we observed a greater movement of the wetting front. 48 hours after the experiment began, the wetting front had advanced downward by 3 to 4 mm. When the root was visible, we observed the dye both attached to the root and in the surrounding transparent soil (Figure 2B). The tracer dye began to appear in the lower transparent soil layer 24 hours after the introduction of the tracer dye. The coloration increased in intensity over time until the end of the experiments. All samples showed some coloration at the end of the experiment.

The quantitative analysis of the image data confirmed visual observations. The dye concentration in the dry transparent soil layer increased with time (Figure 2C). In the case of the control samples, the dye concentration increased up to a mean value of 3.5×10^{-4} mL mL^{-1}. In the case of the samples containing plants, an increase in dye concentration was observed during the entire experiment with a final dye concentration recorded at 2.1×10^{-2} mL mL^{-1}. Quantitatively, the estimated dye concentration in the dry transparent soil layer increased when a root was present, but not in control treatments (Figure 2C). We did not measure a significant change over time in the permeability of the dry transparent soil layer in the control samples (Figure 2D). Statistical analysis of the data using a linear model showed that in the absence of the root, the permeability of the dry transparent soil layer was not significantly different from 0 during the entire experiment (p=0.23 and p=0.69 for slope and intercept). However, the presence of the root had a significant effect on the rate of increase in the permeability of the dry transparent soil layer (p<0.001).

Figure 2: Effect of the presence of winter wheat (*Triticum aestum*) roots on the infiltration of a tracer dye through a dry transparent soil layer. **(A)** Microcosm samples just after introduction of the tracer dye into the top transparent soil layer. Sample without root (Control, left) and sample with root (right). **(B)** Microcosm samples 72 hours after introduction of the tracer dye. **(C)** Concentration of the tracer dye in the dry transparent soil layer measured at different time points. **(D)** Permeability of the dry transparent soil layer measured at different time points. C&D, Green control samples and pink samples with roots.

Root exudates increase water infiltration through the dry transparent soil layer

Following introduction into the microcosm, the dye was observed spreading non homogeneously but rapidly within the top transparent soil layer (Figure 3A), but there was no evidence of the dye being absorbed by the transparent soil particles. The dry transparent soil layer blocked further spreading of the tracer dye into the lower layers of transparent soil. There was no obvious effect of the presence of root exudates on the transport of the dye at this stage of the experiment. The spatial distribution of the tracer dye was analysed over a 120-hour period following the start of the experiment (Figure 3A&B). In control samples, the wetting front did not advance during the first 72 hours but had advanced by 1-2 mm by the end of the experiment (Figure 3A&B). 96 hours after the introduction of the tracer dye, red coloration began visible in the bottom

transparent soil layer in all samples. The coloration increased in intensity with
time until the end of the experiments. In the samples with root exudates, the
wetting front advanced irregularly but continuously into the dry layer throughout
the experiment. At the end of the experiment, the dry transparent soil layer was
completely wet (Figure 3A&B).

Figure 3: Effect of the presence or absence of root exudates from winter wheat
(*Triticum aestum*) on the infiltration of a tracer dye through a dry transparent soil
layer. (A) Microcosm samples just after introduction of the tracer dye into the top
transparent soil layer. Sample without root exudates (left) and sample with root
exudates (right). (B) Microcosm samples 120 hours after introduction the tracer
dye. (C) Concentration of the tracer dye in the dry soil layer measured at
different time points. (D) Permeability of the dry transparent soil layer measured
at different time points. C&D green control samples and pink samples with
exudates.

Quantitatively, the mean dye concentration in the dry transparent soil layer
increased with time (Figure 3C) for both control samples and samples
containing root exudates. When samples contained root exudates, the increase
in dye concentration in the bottom layer was sustained during the experiment.
We did not measure a significant change with time in the permeability of the dry
transparent soil layer in the control samples (Figure 3D). For samples with root
exudates, permeability was maximal 120 hours after introduction of the tracer

dye. Statistical analysis of the data using a linear model showed that in the
absence of root exudates, the permeability of the dry transparent soil layer was
not significantly different from 0 during the entire experiment (p=0.54 and
p=0.45 for slope and intercept). However, the presence of root exudates had a
significant effect on the rate of increase in the permeability of the dry
transparent soil layer (p<0.001).

The presence of root exudates in the dry layers has limited effect on water
infiltration and rewetting of the dry transparent soil layer

Following introduction into the microcosm, the dye spread following a similar
pattern as reported in the previous experiments (Figure 4A). There was no
obvious effect of the presence of root exudates (in the top or dry layer) on the
transport of the dye at this stage of the experiment. The spatial distribution of
the tracer dye was analysed over a 120-hour period following the start of the
experiment (Figure 4A&B). In samples, that did not contain root exudates, the
wetting front did not advance during the first 72 hours but had advanced by 1 to
1.5 mm both in the NN treatment and in the NE treatments (Figure 3A&B). 96
hours after the introduction of the tracer dye, red coloration began visible in the
bottom transparent soil layer in all samples. The coloration increased in
intensity with time until the end of the experiments. No colouration was
observed at the bottom of the microcosm chambers at any stage of the
experiment. There were no strong visual differences between samples with and
without root exudates in the dry transparent soil layer. Samples containing root
exudates in the tracer dye (EN and EE) exhibited qualitatively similar behaviour
to samples containing root exudates in the tracer dye in experiment 2.

Quantitative analysis of the image data confirmed the visual observations. The
dye concentration in the dry transparent soil layer increased with time (Figure
4C). For samples containing no exudates in the tracer dye, the increase was
gradual, but for samples containing root exudates in the tracer dye, a sharp
increase in dye concentration was observed within the first 72 hours and this
was followed by a more moderate increase in dye concentration. We did not
measure a significant change over time in the permeability of the dry
transparent soil layer in samples containing water in the tracer dye (Figure 4D).
In the case of samples with root exudates in the top layer, permeability steadily
increased during the experiment. Statistical analysis of the data using a linear
model showed that in the absence of root exudates, the permeability of the dry
transparent soil layer was not significantly different from 0 during the entire
experiment but the presence of root exudates in the top transparent soil layer
had both a significant effect on the slope and intercept of the relationship
(p<0.001 and p=0.01 respectively). The presence of root exudates in the middle
layer did not have a significant effect on the permeability of the dry transparent
soil layer.

Figure 4 Effect of location of root exudates of winter wheat (*Triticum aestum*) on the infiltration of a tracer dye through a dry transparent soil layer. (A) Microcosm samples just after introduction of the tracer dye into the top transparent soil layer. (B) Microcosm samples 120 hours after the start of the experiment. (C) Concentration of the tracer dye in the dry transparent soil layer measured at different time points. (D) Permeability of the dry transparent soil layer measured at different time points. In C&D green NN, blue NE pink EN and yellow EE.

Physical modifications due to penetration of the dry transparent soil layer do not affect the infiltration and rewetting

Following introduction into the microcosm, the dye spread following a similar pattern than experiments where the tracer dye did not contain root exudates (Figure 5A&B). There was no obvious effect of the presence of the needle on the transport of the dye at this stage of the experiment. The spatial distribution of the tracer dye was analysed over a 120-hour period following the start of the experiment. No difference was observed between treatments during the entire experiment. No dye was observed at the bottom transparent soil layer, nor in the dry transparent soil layer. At the end of the experiment, we observed that the wetting front had advanced about 1 mm downward for all treatments.

Quantitative analysis of the image data confirmed the visual observations. The dye concentration in the dry transparent soil layer did not increase with time in

both conditions (Figure 5C). Calculation of the permeabilities of the dry transparent soil layer revealed that that when the needle was present, there was much higher variability and mean value for the permeability (Figure 5D), which is likely due to the needle having an effect on the saturation of the image. Statistical analyses however showed that the presence of a needle did not have an effect on the calculated permeability (P = 0.0073).

Figure 5 Effect of the presence or absence of a needle on the infiltration of a tracer dye through a dry transparent soil layer. (A) Microcosm samples just after introduction of the tracer dye into the top transparent soil layer. Sample without needle (Control, left) and sample with needle (right). (B) Microcosm samples 120 hours after introduction of the tracer dye. (C) Concentration of the tracer dye in the dry transparent soil layer measured at different time points. (D) Permeability of the dry transparent soil layer measured at different time points. C&D, green control samples, pink samples with a needle.

The hydraulic conductivity is critically sensitive to surface tension and root exudate concentration

Model calibration completed successfully with optimal parameters values for
$\beta^*, \kappa_D^*, \kappa_W^*$ determined from experimental data. Following calibration, the
predicted dye concentration was plotted against experimental data (Figure 6).
Results showed good agreements between the model and experimental
observations. Discrepancies were observed in the control treatment where no
exudates were introduced (NN). Experiments showed an abrupt stop in the
permeability of the dry transparent soil layer, which the model could not
represent. The estimated model parameters gave also interesting insights into
the effects of root exudates on water transport in transparent soil. The exponent
of the power law relating surface tension to hydraulic conductivity was $\beta = 8.8$,
indicating that the concentration of root exudate in solution had a strong effect
on the hydraulic conductivity of transparent soil. The parameter for the
solubilisation or sorption of root exudates $\kappa_D^* = 9.9 \times 10^{-2}\ d^{-1}$ and $\kappa_W^* = 1.8 \times$
$10^{-2}d^{-1}$ also provided useful insight. They showed that although the rate of
solubilisation may not be limiting, because permeability is very small at the start
of the experiments, the fraction of the transparent soil exposed to water limits
the capacity of the dry root exudates to meaningfully increase the infiltration of
water.

Figure 6 Model predictions against experimental data for dye concentration in
the bottom transparent soil layer. (A) Comparison between model and
experimental data for experiments when root exudates are introduced in the top
layer in the tracer dye solution. EN is shown in pink and NN is shown in green.
(B) Comparison between model and experimental data for experiments when
root exudates are introduced dry in the middle layer. NE is shown in pink and
NN is shown in green. In all plots, dotted lines represent model predictions data
whereas plain lines indicate experimental observations.

Discussion

Water infiltration and loss of water from the rhizosphere

Plant roots absorb water stored in the soil's pore spaces, which is subsequently
replenished through irrigation, rainfall, or capillary rise. However, a substantial
portion of the water input is lost before it can be absorbed by the plant. Losses
are due to evaporation, deep percolation, runoff, and lateral flow and can be
particularly high before canopy closure (Chen et al. 2020). In a cropping
system, 20% of the evapotranspiration is typically due to bare soil evaporation,
but it can exceed 50% of the total influx of water. For example, a review of the
seasonal losses of water by evaporation in rain fed wheat in Australia showed
that between 27% and 68% of the evapotranspiration is due to evaporation
(Unkovich et al. 2018). The percentage of evaporation was between 15% and
40% in Maize in a range of water and N regimes (Hernández et al. 2015). In
sandy soils, deep percolation can become the dominant source of water loss.
Nassah et al. (2018) reported nearly 50% loss of water inputs lost due to water
leaving the rooted zone through deep percolation.

The root system architecture, the spatial and topological arrangement of roots
into a soil, is therefore a critical trait to determine the ability of a crop to extract
water from the soil. First because it determines the regions of the soil where
water can be extracted. When a root system is shallow, the plant can more
readily acquire immobile nutrients that are strongly absorbed to soil particles
(Lynch and Brown 2002). When the root system is deep, it can reach the water
table and extract groundwater during drought (Lynch 2013). However, water
balance in soil is very dynamic and another important role of root systems may
be to intercept water, that is to distribute water into soil following a rainfall, to
minimize losses from the rhizosphere. Although roots are known to facilitate
preferential water flow through the soil, this function of root systems has been
largely overlooked as a valuable trait to improve water management in cropping
systems. Recently, mathematical models that combined water uptake with
preferential flow induced by plant roots showed that root system architectures
may have very different abilities to intercept rainfall water, and that root system
ideotypes may vary as a function of soil type (Mair et al. 2023).

The experiments described in this study aimed to investigate the impact of root
traits on water infiltration in dry soil. We used a dye tracing approach rather than
infiltrometers to miniaturise and shorten the duration of the microcosm
experiments. Observation of the dye also gave insights into the mechanisms of
infiltration. The use of an artificial soil improved the accuracy of measurements
since all the dye below the dry layer was considered in the measurement of the
fluxes. The artificial soils also made experiments more manageable. Nafion^{TM},
the material used as substrate, is mildly hydrophobic when dry (water contact
angle of 105º (Goswami et al. 2008)), and resulted in low infiltration rates but
with measurable effects after a few days of experiments. Agricultural soils which
have low organic matter content usually exhibit higher infiltration rates (Basset
et al. 2023), and organic matter rich soils such as peat are too hydrophobic
when dry (water contact angle of up to 120º (Valat et al. 1991)).

Physical processes involved in preferential flow induced by plant roots

We observed that root exudation was a primary factor influencing the infiltration
of water through dry soil. This result suggests that surfactants found in root
exudates may play a crucial role in affecting water movement within the soil.
Not only root exudates are known to contain surfactants (Read et al. 2003), but
measurements of water retention in rhizosphere and bulk soil have also
indicated that rhizosphere solutions may exhibit lower surface tension (Whalley
et al. 2005). Surfactants are also used for the irrigation of plants growing in
hydrophobic soils and arid environments (Baratella and Trinchera 2018;
Ogunmokun et al. 2020), which confirms that the exudation of surfactants may
be a beneficial trait in certain soils.

Experiments using syringe needles indicated that changes in porosity induced
by root growth may not affect significantly the infiltration of water. In a coarse
and loosely packed, sand, similar to our artificial soil, root growth was shown to
induce dilation of <4% and increase the porosity of about 10% over a layer of
<2 mm around the root (Anselmucci et al. 2021). The Kozeny–Carman equation
$K \propto \text{porosity}^3/(1 - \text{porosity})^2$ predicts that such changes can result in an
increase of 50% of the hydraulic conductivity (Schulz et al. 2019). But a 1 mm
thick layer represents about 10% of the cross section of the soil. Also, dilation of
the rhizosphere soil is associated with reduction of porosity further away from
the root and collectively, this would likely limit the changes in the hydraulic
conductivity to less than 5%. In the Young-Laplace equation, pore size and
surface tension exert equal effects on capillary entry pressure (Hallett 2008).
Given that root exudates have been shown to reduce surface tension by 27%,
reducing it from 72 mN m^{-1} to 50 mN m^{-1} (Read et al. 2003), it is plausible for
root exudation to have a stronger effect than dilation in some conditions.

Root exudation as a mechanism to influence water availability in soil

During drought conditions, evidence suggests that both the quantity and
composition of root exudates undergo significant changes (Karlowsky et al.
2018; Williams and de Vries 2020). However, it remains uncertain whether
these changes reflect a plant's response to modify the physical properties of the
soil. It was suggested that for a large part, changes in root exudation result from
shift in plants metabolic activity, with root exudates reflecting the most abundant
root metabolites (Gargallo-Garriga et al. 2018). The lifespan of root exudates in
soils is likely to be highly variable. Many of the compounds deposited in the soil
by plant roots are consumed by soil microbes (Sasse et al., 2018), and it is
challenging to determine the extent to which changes in soil physical properties
are directly attributable to rhizodeposition, as opposed to being secondary
effects arising from microbial activity. For example, soil bacteria are
extraordinary producers of biosurfactants (Karanth et al. 1999; Domingues et al.
2020). They are also able to synthesize extracellular polymeric substances
(EPS) which enhance soil water retention (Zheng et al. 2018; Benard et al.
2023). The reduction of surface tension typically necessitates small amounts of
a single surfactant molecule, such as Lecithin found in soybeans (Read et al.
2003). Consequently, even subtle changes in the composition of root exudates
could significantly impact soil properties.

It is also far from clear what the desired physical properties of the rhizosphere solution should be. While surfactant production may be desirable in low soil water content when water is trapped in small pores (Whalley et al. 2005) or in soils rich in organic matter (strong hysteresis between wetting and drying), the broader benefits are less clear. The level of organic matter in agricultural soils is minimal (Loveland and Webb 2003; Rawls et al. 2004) and heterogeneously distributed (Lehmann et al. 2008). Also, losses due to evaporation and deep percolation are extremely dependent on rainfall or irrigation patterns. Answering such questions will be challenging and is likely to require further experimental characterisation, together with the integration of mathematical modelling and simulation studies.

Acknowledgements

We acknowledge the funding from the Spanish Ministry of Science, Innovation and Universities under the grant agreements No. PID2020-112950RR-I00 and No. PID2023-149435OR-I00 (projects MICROCROWD and BIOFLOW). This work was also supported by the European Union's Horizon Europe under the grant agreement No.101060124 (Project Root2Res).

References

Ahmed MA, Kroener E, Benard P, et al (2016) Drying of mucilage causes water repellency in the rhizosphere of maize: measurements and modelling. Plant Soil 407:161–171

Anselmucci F, Andò E, Viggiani G, et al (2021) Imaging local soil kinematics during the first days of maize root growth in sand. Sci Rep 11:22262

Baratella V, Trinchera A (2018) Organosilicone surfactants as innovative irrigation adjuvants: Can they improve water use efficiency and nutrient uptake in crop production? Agric Water Manag 204:149–161

Basset C, Abou Najm M, Ghezzehei T, et al (2023) How does soil structure affect water infiltration? A meta-data systematic review. Soil Tillage Res 226:105577

Benard P, Bickel S, Kaestner A, et al (2023) Extracellular polymeric substances from soil-grown bacteria delay evaporative drying. Adv Water Resour 172:104364

Benson CH, Chiang I, Chalermyanont T, Sawangsuriya A (2014) Estimating van Genuchten parameters α and n for clean sands from particle size distribution data. In: From soil behavior fundamentals to innovations in geotechnical engineering: Honoring Roy E. Olson. pp 410–427

Berry JD, Neeson MJ, Dagastine RR, et al (2015) Measurement of surface and interfacial tension using pendant drop tensiometry. J Colloid Interface Sci 454:226–237

Brochu E, Cora VM, De Freitas N (2010) A tutorial on Bayesian optimization of expensive cost functions, with application to active user modeling and hierarchical reinforcement learning. arXiv preprint arXiv:10122599

Bruand A, Cousin I, Nicoullaud B, et al (1996) Backscattered electron scanning images
of soil porosity for analyzing soil compaction around roots. Soil Science Society of
America Journal 60:895–901

Carminati A, Moradi AB, Vetterlein D, et al (2010) Dynamics of soil water content in the
rhizosphere. Plant Soil 332:163–176. https://doi.org/10.1007/s11104-010-0283-8

Carsel RF, Parrish RS (1988) Developing joint probability distributions of soil water
retention characteristics. Water Resour Res 24:755–769

Chen S, Du T, Wang S, et al (2020) Evaluation and simulation of spatial variability of
soil property effects on deep percolation and nitrate leaching within a large-scale
field in arid Northwest China. Science of the Total Environment 732:139324

Clothier BE, Green SR (1997) Roots: the big movers of water and chemical in soil. Soil
Sci 162:534–543

Dexter AR (1987) Compression of soil around roots. Plant Soil 97:401–406

Domingues VS, de Souza Monteiro A, Júlio ADL, et al (2020) Diversity of metal-
resistant and tensoactive-producing culturable heterotrophic bacteria isolated from
a copper mine in Brazilian Amazonia. Sci Rep 10:1–12

Downie HF, Holden N, Otten W, et al (2012) Transparent Soil for Imaging the
Rhizosphere. PLoS One 7:. https://doi.org/10.1371/journal.pone.0044276

Duddek P, Carminati A, Koebernick N, et al (2022) The impact of drought-induced root
and root hair shrinkage on root–soil contact. Plant Physiol 189:1232–1236

Gargallo-Garriga A, Preece C, Sardans J, et al (2018) Root exudate metabolomes
change under drought and show limited capacity for recovery. Sci Rep 8:1–15

Goswami S, Klaus S, Benziger J (2008) Wetting and absorption of water drops on
Nafion films. Langmuir 24:8627–8633

Hallam J, Holden J, Robinson DA, Hodson ME (2021) Effects of winter wheat and
endogeic earthworms on soil physical and hydraulic properties. Geoderma 400:.
https://doi.org/10.1016/j.geoderma.2021.115126

Hallett PD (2008) A brief overview of the causes, impacts and amelioration of soil water
repellency–a review. Soil and Water Research 3:521–528

Helliwell JR, Sturrock CJ, Mairhofer S, et al (2017) The emergent rhizosphere: imaging
the development of the porous architecture at the root-soil interface. Sci Rep
7:14875

Hernández M, Echarte L, Della Maggiora A, et al (2015) Maize water use efficiency and
evapotranspiration response to N supply under contrasting soil water availability.
Field Crops Res 178:8–15

Jačka L, Walmsley A, Kovář M, Frouz J (2021) Effects of different tree species on
infiltration and preferential flow in soils developing at a clayey spoil heap.
Geoderma 403:115372

Karagunduz A, Pennell KD, Young MH (2001) Influence of a Nonionic Surfactant on the
Water Retention Properties of Unsaturated Soils. Soil Science Society of America
Journal 65:1392–1399. https://doi.org/10.2136/sssaj2001.6551392x

Karanth NGK, Deo PG, Veenanadig NK (1999) Microbial production of biosurfactants and their importance. Curr Sci 116–126

Karlowsky S, Augusti A, Ingrisch J, et al (2018) Drought-induced accumulation of root exudates supports post-drought recovery of microbes in mountain grassland. Front Plant Sci 9:1593

Koebernick N, Daly KR, Keyes SD, et al (2019) Imaging microstructure of the barley rhizosphere: particle packing and root hair influences. New Phytologist 221:1878–1889

Kool JB, Parker JC (1987) Development and evaluation of closed-form expressions for hysteretic soil hydraulic properties. Water Resour Res 23:105–114. https://doi.org/10.1029/WR023i001p00105

Lehmann J, Solomon D, Kinyangi J, et al (2008) Spatial complexity of soil organic matter forms at nanometre scales. Nat Geosci 1:238–242

Leung AK, Boldrin D, Liang T, et al (2018) Plant age effects on soil infiltration rate during early plant establishment. Géotechnique 68:646–652

Liu Y, Patko D, de la Mata AL, et al (2025) Microcosm fabrication platform for live microscopy of plant-soil systems. Biosyst Eng 252:105–114

Loveland P, Webb J (2003) Is there a critical level of organic matter in the agricultural soils of temperate regions: a review. Soil Tillage Res 70:1–18

Lucas M, Schlüter S, Vogel H-J, Vetterlein D (2019) Roots compact the surrounding soil depending on the structures they encounter. Sci Rep 9:16236

Lynch JP (2013) Steep, cheap and deep: an ideotype to optimize water and N acquisition by maize root systems. Ann Bot 112:347–357

Lynch JP, Brown KM (2002) Topsoil foraging - an architectural adaptation of plants to low phosphorus availability. Plant Soil 237:225–237

Mair A, Dupuy L, Ptashnyk M (2023) Can root systems redistribute soil water to mitigate the effects of drought? Field Crops Res 300:109006. https://doi.org/https://doi.org/10.1016/j.fcr.2023.109006

Marshall MR, Ballard CE, Frogbrook ZL, et al (2014) The impact of rural land management changes on soil hydraulic properties and runoff processes: results from experimental plots in upland UK. Hydrol Process 28:2617–2629

Mualem Y (1976) A new model for predicting the hydraulic conductivity of unsaturated porous media. Water Resour Res 12:513–522. https://doi.org/10.1029/WR012i003p00513

Nassah H, Er-Raki S, Khabba S, et al (2018) Evaluation and analysis of deep percolation losses of drip irrigated citrus crops under non-saline and saline conditions in a semi-arid area. Biosyst Eng 165:10–24

Naveed M, Ahmed MA, Benard P, et al (2019) Surface tension, rheology and hydrophobicity of rhizodeposits and seed mucilage influence soil water retention and hysteresis. Plant Soil 437:65–81

Ni JJ, Leung AK, Ng CWW (2019) Modelling effects of root growth and decay on soil
water retention and permeability. Canadian Geotechnical Journal 56:1049–1055

Oberbroeckling KJ, Dunwoody DC, Minteer SD, Leddy J (2002) Density of Nafion
exchanged with transition metal complexes and tetramethyl ammonium, ferrous,
and hydrogen ions: commercial and recast films. Anal Chem 74:4794–4799

Ogunmokun FA, Liu Z, Wallach R (2020) The influence of surfactant-application
method on the effectiveness of water-repellent soil remediation. Geoderma
362:114081

R Core Team (2021) R Core Team (2021). R: A language and environment for
statistical computing

Rawls WJ, Nemes A, Pachepsky YA (2004) Effect of soil organic carbon on soil
hydraulic properties. Developments in soil science 30:95–114

Read DB, Bengough AG, Gregory PJ, et al (2003) Plant roots release phospholipid
surfactants that modify the physical and chemical properties of soil. New
phytologist 157:315–326

Read DB, Gregory PJ, Bell AE (1999) Physical properties of axenic maize root
mucilage. Plant Soil 211:87–91

Schmid B, Schindelin J, Cardona A, et al (2010) A high-level 3D visualization API for
Java and ImageJ. BMC Bioinformatics 11:274

Schulz R, Ray N, Zech S, et al (2019) Beyond Kozeny–Carman: predicting the
permeability in porous media. Transp Porous Media 130:487–512

Scott EM, Rattray EAS, Prosser JI, et al (1995) A mathematical model for dispersal of
bacterial inoculants colonizing the wheat rhizosphere. Soil Biol Biochem 27:1307–
1318

Song L, Li JH, Zhou T, Fredlund DG (2017) Experimental study on unsaturated
hydraulic properties of vegetated soil. Ecol Eng 103:207–216

Syms R (2017) Rapid evaporation-driven chemical pre-concentration and separation
on paper. Biomicrofluidics 11:

Unkovich M, Baldock J, Farquharson R (2018) Field measurements of bare soil
evaporation and crop transpiration, and transpiration efficiency, for rainfed grain
crops in Australia–A review. Agric Water Manag 205:72–80

Valat B, Jouany C, Riviere LM (1991) Characterization of the wetting properties of air-
dried peats and composts. Soil Sci 152:100–107

van Genuchten MTh (1980) A Closed-form Equation for Predicting the Hydraulic
Conductivity of Unsaturated Soils. Soil Science Society of America Journal
44:892–898. https://doi.org/10.2136/sssaj1980.03615995004400050002x

Vogel T, Zhang R (1996) The HYDRUS Code for Simulating One-Dimensional Water
Flow, Solute Transport, and Heat Movement in Variably-Saturated Media. Version
5.0. https://doi.org/10.13140/RG.2.1.3456.7525

Whalley WR, Riseley B, Leeds-Harrison PB, et al (2005) Structural differences
between bulk and rhizosphere soil. Eur J Soil Sci 56:353–360

Williams A, de Vries FT (2020) Plant root exudation under drought: implications for
ecosystem functioning. New Phytologist 225:1899–1905

Xiao T, Li P, Fei W, Wang J (2024) Effects of vegetation roots on the structure and
hydraulic properties of soils: A perspective review. Science of The Total
Environment 906:167524

Zheng W, Zeng S, Bais H, et al (2018) Plant growth-promoting rhizobacteria (PGPR)
reduce evaporation and increase soil water retention. Water Resour Res 54:3673–
3687

Zickenrott IM, Woche SK, Bachmann J, et al (2016) An efficient method for the
collection of root mucilage from different plant species—A case study on the
effect of mucilage on soil water repellency. Journal of Plant Nutrition and Soil
Science 179:294–302. https://doi.org/10.1002/jpln.201500511