1 **Vocal mimicry in corvids**

2 Claudia A.F. Wascher[1]*, Gemini Waterhouse[1] & Bret A. Beheim[2]

3 [1]Behavioural Ecology Research Group, School of Life Sciences, Anglia Ruskin University, United Kingdom
4 [2] Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6,
5 04103 Leipzig, Germany

30 *Corresponding author: Claudia A.F. Wascher: School of Life Sciences, Anglia Ruskin University, East

31 Road, Cambridge, CB1 1PT United Kingdom; Phone: +44 (0) 1223 698270;

32 e-mail: claudia.wascher@gmail.com

Abstract

Vocal mimicry, the copying of sounds produced by another species or the environment, is commonly described in vocal learners, such as songbirds. Understanding the functions of vocal mimicry can help to uncover the evolutionary drivers of vocal learning. Different adaptive functions like interspecific and intraspecific communication have been suggested, as well as the possibility of vocal mimicry to be a mistake during vocal learning. In the present study, we review the occurrence of mimicry in the family of corvids and investigate the socio-ecological factors driving the evolution of vocal mimicry in this group of birds. We recorded evidence of vocal mimicry from primary (xeno-canto recordings) as well as secondary sources (published literature) and found evidence for vocal mimicry in 31 out of 128 corvid species (24%). Smaller species were described as mimics more compared to larger species. Other socio-ecological factors like breeding status, habitat and trophic niche did not have a significant effect. More mimicry was recorded during the non-breeding season compared to the breeding season. We used Bayesian modelling based on existing data from primary and secondary sources to estimate the occurrence of mimicry, suggesting that vocal mimicry may be more widespread among corvids than currently documented, with many species potentially being 'hidden mimics'. Our study for the first time systematically reviews the occurrence of vocal mimicry across the family of corvids and investigates a range of socio-ecological factors driving the behaviour. Our analysis suggests many corvid species likely to be 'hidden' mimics and we provide specific suggestions for species which according to our statistical model are most likely mimicking corvid species, hopefully inspiring future field work on those species.

Introduction

Vocal mimicry, the copying of sounds produced by another species or the environment, is commonly described in songbirds (Kelley et al. 2008; Dalziell et al. 2015) and also has been reported in mammals including an Asian elephant, *Elephas maximus* (Stoeger et al. 2012), a white whale, *Delphinapterus leucas* (Ridgway et al. 2012), a harbor seal, *Phoca vitulina* (Duengen et al. 2023), and greater mouse-eared bats, *Myotis myotis* (Ancillotto et al. 2022). Mimics usually acquire sounds through vocal production learning, the ability to produce novel signals through imitation or modification of existing vocalisations (Janik and Knörnschild 2021). Vocal production learning in animals has parallels with human vocal ontogeny (Ter Haar et al. 2021), as it requires an individual to process acoustic information from an appropriate model, engage in vocal practice and evaluation through auditory feedback, and requires enhanced forebrain control of the vocal organ (Nieder and Mooney 2020).

According to Dalziell et al. (2015), vocal mimicry is defined as vocalisations resembling heterospecific or environmental sounds which change the behaviour of the receiver after perceiving the acoustic resemblance between the mimic and the model, and the behavioural change confers a selective advantage on the species mimicking. This definition excludes other forms for vocal resemblance, for example as a result of chance or common ancestry. Vocal mimicry may be a result of mistakes during vocal learning and have no function (Kelley et al., 2008). In contrast to this, a number of different potential functions of vocal mimicry have been previously suggested, broadly falling into two categories (a) interspecific communication, to avoid threats, dissuade competitors, or imitate threats to allow access to food resources and (b) intraspecific communication, in the context of sexual selection or for social affiliation (reviewed by Kelley et al., 2008). For example, mimics are suggested to produce vocalisations from predatory or agonistic species to deter competitors or potential predators from their territories (Dobkin 1979). Australian magpies, *Gymnorhina tibicen*, mimic several potential nest predators, for example the barking owl, *Ninox connivens*, and the boobook owl, *Ninox novaeseelandiae* (Kaplan 1999). Fork-tailed drongos, *Dicrurus adsimilis,* mimic alarm calls of other species to deter heterospecifics from food sources, in order to steal the food (Flower et al. 2014). Individuals might also benefit from vocal mimicry when used to attract other individuals. For example, greater racket-tailed drongos, *Dicrurus paradiseus,* mimic songs and contact calls to attract other species, which increases foraging efficiency (Goodale and Kotagama 2006a) and alarm calls of other species alongside their own mobbing calls, potentially attracting other species to support them when mobbing predators (Goodale and Kotagama 2006b). This shows that some species flexibly use vocal mimicry, and this may be context dependent. Finally, mimicked calls are sometimes incorporated into song or sexual displays and might play a role in sexual selection (Kelley et al., 2008). Overall, surprisingly little empirical research on the functional explanations of vocal mimicry is available.

Phylogenetic analysis has previously shown that vocal mimicry is unlikely to be an ancestral state in songbirds, meaning that it may have evolved independently on multiple occasions (Goller and Shizuka 2018). A review of European passerines estimated approximately 40% of songbird species to mimic (Garamszegi et al., 2007). Sounds mimicked by songbirds include birdsong (northern mockingbird, *Mimus polyglottos*: Gammon and Altizer 2011), alarm calls of other species (spotted bowerbird, *Ptilonorhynchus maculatus*: Kelley and Healy 2011; red-capped robin-chat, *Cossypha natalensis*: Oatley 1969; thick-billed euphonia, *Euphonia laniirostris*: Morton 1976), calls of birds of prey (blue jay, *Cyanocitta cristata*: Hailman 1990), and human voices (European starlings, *Sturnus vulgaris*: West et al. 1983). As suggested by Gammon (2013), the 'model selection', *i.e.* type of sound mimicked, can provide valuable insight into the potential functions of vocal mimicry. For example, northern mockingbirds, *Mimus polyglottos,* preferentially mimic species whose sounds are acoustically similar to their non-imitative song, supporting the acoustic similarity hypothesis, and suggesting a non-adaptive explanation of vocal mimicry (Gammon, 2013). Mimicry of predator calls by Steller's jays, *Cyanocitta stelleri,* is also linked to mate selection, with mimicry occurring more frequently during the breeding season and when in the presence of their mates (Tippin 2017).

Corvids *(Aves: Corvidae),* are a large and diverse group with over 120 species split between 21 genera, an array of which can be found on all continents except Antarctica (Madge and Burn 1994). Many corvid species have demonstrated advanced cognitive abilities in the social and physical domain (Taylor 2014). Corvids belong to the suborder of oscine passerine birds, where the vocal organ (syrinx) is highly developed to produce vocalizations (Suthers and Zollinger 2004). Vocalisations in corvids present a wide range of ecological and social adaptations, such as communicating information about predators such as predator type or level of risk or individual recognition (Wascher and Reynolds 2025). Corvids are open-end vocal learners, which means they can acquire new vocalisations throughout their lives and not only in a specific sensitive period (Brenowitz et al. 1997). Carrion crows are capable of volitional control of vocalisations, *i.e.* can be trained to vocalise and withhold vocalisations in response to arbitrary stimuli (Brecht et al. 2019). Vocalisations in New Caledonian crows, *Corvus moneduloides* (Bluff et al. 2010) and common ravens, *Corvus corax* (Enggist-Dueblin and Pfister 2002) are socially learned from conspecifics. Several species of corvids have demonstrated the ability to mimic human speech when hand-raised and kept in close proximity to humans (Bluff et al., 2010). Mimicry of predator calls has been documented in several corvid species including Steller's jays (Hope 1980; Tippin 2017), Sri Lanka blue-magpies, *Urocissa ornata* (Ratnayake et al. 2010), Florida scrub-Jays, *Aphelocoma coerulescens* (Woolfenden and Fitzpatrick 2020), blue jays, *Cyanocitta cristata* (Hailman 1990), and Eurasian jays, *Garrulus glandarius* (Goodwin 1951). Tippin (2017) conducted an in-depth study of vocal

mimicry in Steller's jays, which found significant differences in age and sex between mimicking and

non-mimicking individuals, where males were less likely to mimic compared to females, and the

likelihood of an individual mimicking decreased with age.

Missing data presents a significant challenge in the study of animal behaviour, limiting our

understanding of key phenomena (Garamszegi and Møller 2012; Nakagawa 2015). Factors such as

habitat inaccessibility (Evans et al. 2018), species rarity (Roberts et al. 2016), human-centric biases in

study design (Archer et al. 2014; Ellison et al. 2021; Bowler et al. 2025; Daw et al. 2025), and

technological constraints (Hebblewhite and Haydon 2010) contribute to missingness. In evolutionary

biology, missing data can skew our understanding of trait evolution, obscuring the selective pressures

and socio-ecological contexts that drive behavioural adaptations. The problem of missing data can be

addressed with more systematic data collection, e.g., planned missing data design (Noble and

Nakagawa 2018), application of emerging technologies and statistical approaches, such as imputation

procedures (Augustine et al. 2014), and a recognition that the absence of evidence is not evidence of

absence in behavioural research.

The aim of the present study is to describe the occurrence of mimicry in corvids. We investigated the

evidence for mimicry from primary, i.e. recordings on Xeno-canto, and secondary sources, i.e.

published literature. Further, we investigate empirical support for mimicry in corvids to serve an

interspecific or intraspecific function or whether evidence base hints more towards mimicry as an error

during vocal learning. If vocal mimicry in corvids serves a function of interspecific communication, we

expect smaller corvid species to mimic more, as they are more at risk of predation (Dierschke 2003).

Further, if mimicry in corvids functions to avoid threats, we would expect them to mainly mimic birds

of prey. If vocal mimicry serves intraspecific communication, such as sexual selection, we expect

mimicry to mostly occur during the breeding season. We further investigate the relationship between

size of the vocal repertoire and the occurrence of vocal mimicry, as we expect species with larger

repertoires to mimic more. In addition to biological factors driving vocal mimicry in corvids, species

differ significantly in terms of study effort (Wascher & Reynolds, 2025). We investigate how differences

in recording effort affect the detection of mimicry in corvids and evaluate the likelihood of 'hidden'

mimicry.

Methods

Data on vocal mimicry

The present study was approved by the School of Life Sciences departmental ethics panel at Anglia Ruskin University (ETH2223-3747). This is a desk-based study, and no animals have been used for the research. Data were compiled from a variety of primary (Xeno-canto) and secondary sources (Birds of the World and Handbook of the Birds of Europe the Middle East and North Africa). Scientific and common names were standardized using a combination of the IOC Bird list version 14.1. (Gill et al. 2024) and the 'Clements checklist' (Clements et al. 2024) and 128 species of corvids were included. A Xeno-canto advanced search was conducted between the 27th of April and the 1st of May 2023 (searches have been updated on the 22nd of June 2023 and 5th May 2024). The Xeno-canto search yielded a total of 19,566 audio recordings of corvid calls from 128 different species (mean±SD per species: 153 ± 360), belonging to 21 genera (mean±SD per genus: $889 \pm 1,767$). The number of recordings per species ranged from zero to 2,272 recordings. All recordings with mentions of 'mimicry', 'imitation' or 'production of heterospecific calls' in the comments or labels section were identified. For the purpose of this study, we defined vocal mimicry as the occurrence of any type of non-conspecific sound. We noted the duration of the recording, date, whether the recording was taken during the breeding season or not, latitude and longitude and type of sound mimicked. Recordings were then listened to by GW and CAFW, and if both observers agreed that they contained convincing evidence for production of mimicry, the type of sound was noted (*e.g.*, species mimicked) and the observation was considered as evidence of mimicry. A total of 447 recordings containing mimicry have been identified. For 297 recordings it was indicated that the vocalising bird was seen by the recorder, for 106 recordings the bird was not observed and for 44 recordings this information was unknown. Additionally, secondary literature was searched for additional reports, resulting in additional records of mimicry from 22 species (Cramp and Perrins 1994; Poole and Gill 2020). For each species, the number of entries in the corvids literature database (Droege and Töpfer 2016) plus one entry in Birds of the World (Poole & Gill, 2020) was used as an estimate of the volume of secondary literature available (range: 1-423 entries). For each piece of evidence of vocal mimicry, CAFW allocated a 'reliability score' of high, moderate or low. Sixty-five percent of recordings have been rated as 'high', 23% as 'moderate' and 12% as 'low' reliability.

For each of the 128 species, we recorded body mass (information available for 121 species), vocal repertoire (98 species), habitat (coastal, desert, forest, grassland, human modified, rock, shrubland, woodland; reported for 124 species) and trophic niche (frugivore, granivore, invertivore, omnivore; reported for 124 species).

Statistical analysis

Description of the dataset

All analyses were conducted in R version 4.3.1. (R Core Team 2019). We constructed a phylogenetic

tree for the Corvidae family using data from the Open Tree of Life (OTL). The tree was retrieved from

the Open Tree of Life database (OpenTreeOfLife et al. 2019) using the rotl R package (Michonneau et

al. 2016). We employed Bayesian generalized linear mixed-effects models built with the MCMCglmm

package (Hadfield 2010) to determine effects of body size, vocal repertoire size, habitat and trophic

niche on vocal mimicry in corvids. Genus was included as a random factor and the OTL tree served as

the foundational phylogeny for the MCMCglmm. We ran Markov chain Monte Carlo (MCMC) chains for

3,000,000 iterations, thinned by 2000, and employed a burn-in of 2,000,000. All main effects were

given normal(0, 0.5) log-odds prior, with a normal(0, 1) log-odds intercept prior. Random effects by

genus were given an inverse Wishart prior with V = 1 and nu = 10.

To investigate whether corvids mimicked birds of prey more often compared to other types of sound

and whether more recordings occurred during the breeding season compared to the non-breeding

season, we conducted Wilcoxon signed rank tests. Data and code of the present study are published as

online supplementary materials (Supplementary File 2; supplementary File 3).

Estimating the occurrence of mimicry

In order to estimate the occurrence of mimicry, we have developed a Bayesian model of evidence

updating, assuming that for any specific kind of evidence (database entries, audio recordings), there is

a corresponding rate ρ at which mimics are detected per-unit evidence. The model enables the

calculation of the posterior probability a species in which no mimicry is detected is, in fact, a mimic. Full

details of the model are presented in the supplementary materials.

<u>Results</u>

Description of the dataset

We recorded evidence of vocal mimicry in 32 species of *Corvidae*, belonging to twelve genera (25% of

species; Figure 1). Twenty-two species of corvids are described as mimics in the literature and

recordings from 18 species have been identified on xeno-canto, including nine species not previously

described as mimics (Indochinese green-magpie, *Cissa hypoleuca;* pied crow, *Corvus albus;* hooded

crow, *Corvus cornix;* carrion crow, *Corvus corone;* black-chested jay, *Cyanocorax affinis;* plush-crested

jay, *Cyanocorax chrysops;* white-naped jay, *Cyanocorax cyanopogon;* Yucatan jay, *Cyanocorax*

yucatanicus; clark's nutcracker*, Nucifraga Columbiana;* Table 1). One to 71 different types of sounds

were mimicked (mean ± standard deviation: 5.687 ± 12.568). Sounds have been mimicked from a wide

range or birds (Accipitriformes, Anseriformes, Charadriiformes, Cuculiformes, Falconiformes,

Galliformes, Gaviiformes, Gruiformes, Pelecaniformes, Passeriformes, Piciformes, Strigiformes),

mammals (Artiodactyla, Carnivora, Perissodactyla, Primates (human voice), Rodentia) amphibian

(Anura) and insect species (Hemiptera) as well as anthropogenic sounds, such as bells, lawnmower,

motorbike (see supplementary materials for full list of mimicked species). Corvids did not mimic

predator sounds more often compared to sounds of non-predatory species (Wilcoxon signed-rank test:

V=215, n=32, p=0.792). On xeno-canto mimicry more files containing mimicry were recorded outside of

the breeding season compared to during the breeding season (Wilcoxon signed-rank test: V=16.5,

n=18, p=0.002).

Figure 1: Vocal mimicry in 128 species of corvids. Grey points indicate species with no mimicry

reported, red points indicate species with recordings of mimicry on xeno-canto and reported as mimics

in the literature, blue points illustrate species with recordings of mimicry on xeno-canto but no records

in the literature and yellow dots indicate species only reported as mimics in the literature but no

recordings of mimicry are available on xeno-canto.

Table 1: Cross-table indicating the number of species detected as mimics in the literature (2°) and on

xeno-canto (1°). Records per species indicate number of records in the covid literature

database (2°) and xeno-canto (1°)

mimicry category	n.species	2° records per species	1° records per species
no mimicry seen	97	15	60
2° evidence only	13	40	192
1° evidence only	9	45	433
1° & 2° evidence	9	115	825

261 Body mass significantly affected the occurrence of mimicry, with smaller species being detected as
262 mimics more compared to larger species (Table 2, Figure 2). Repertoire size, habitat, breeding system
263 and trophic niche did not affect mimicry status (Table 2).

265 **Table 2.** Results of Markov chain Monte Carlo generalized linear mixed-effect model (MCMCglmm).
266 Factors with significant effects ($p \leq 0.05$) are shown in bold.

parameters	post.mean	l-95% CI	u-95% CI	eff.samp	pMCMC
(intercept)	-0.109	-1.783	1.998	500	0.916
body mass	**-0.229**	**-0.541**	**-0.004**	**405.738**	**<0.002**
repertoire	0.237	-1.141	1.283	617.836	0.704
habitat breadth (generalist relative to hypergeneralist)	0.011	-1.179	1.54	598.69	1
habitat breadth (generalist relative to specialist)	-0.054	-1.279	1.443	500	0.908
breeding system (cooperative breeding relative to groups)	-0.013	-1.583	1.221	500	0.988
breeding system (cooperative breeding relative to territorial pairs)	-0.021	-1.322	1.274	500	0.968
trophic niche (frugivore relative to granivore)	0.024	-1.278	1.538	500	0.972
trophic niche (frugivore relative to invertivore)	-0.042	-1.535	1.214	500	0.928
trophic niche (frugivore relative to omnivore)	0.056	-1.228	1.502	500	0.94

Figure 2: Body mass (g) in species with no mimicry detected (0) and mimicry detected (1)

271 *Estimating the occurrence of mimicry*

272 Parameter estimates for estimates of occurrence of mimicry from primary sources and secondary

273 sources are presented in supplementary materials. Figure 3 shows how different information sources

274 update the detection of occurrence of mimicry, both by increasing the average but also decreasing the

275 posterior uncertainty about that average.

278
279 Figure 3: Prior and posterior estimates (means and 89% HPDI) of the frequency of occurrence of
280 mimicry across corvid species conditioning on different information evidence (primary sources,
281 secondary sources, and both combined).
282

283 Using this model, we can calculate a composite evidence score, in practice converting a count of

284 recordings on xeno-canto into the equivalent number of corvid database entries (or vis versa), which is

285 useful to understand how the hidden mimicry probabilities for each species relates to research

286 attention (Figure 4).

288
Figure 4: Composite evidence score related to the probability of hidden mimicry.

Looking at posterior probabilities of hidden mimicry for each species individually. The ten species which

are predicted as least likely to be mimics and ten species nominated as most likely to be mimic are

presented in the supplementary materials.

<u>Discussion</u>

In the present study, we describe vocal mimicry in 32 species of corvids belonging to twelve genera,

showing that vocal mimicry is widespread across this family of birds. With 24% of corvids known to

mimic, it is more prevalent in this group, as opposed to the 8.9% observed in the entire suborder of

songbirds (Goller and Shizuka 2018). Additionally, because mimicry detection is strongly associated

with the amount of research effort directed at any specific corvid species, there is reason to believe

that the occurrence of vocal mimicry is starkly underestimated by the current empirical evidence and

many species of corvids are likely to be 'hidden mimics', meaning they are undetected as mimics by the

currently available literature and data.

Our study underscores the importance of crowd-sourced databases, such as xeno-canto, revealing

potentially overlooked or underreported behaviours, such as vocal mimicry. We have identified nine

species of corvids as mimics based on xeno-canto recordings, which have not previously been reported

as mimics. Xeno-canto has previously been used scientifically to develop machine learning methods for

automatic detection and classification (Ovaskainen et al. 2018), describing geographic variation (Deng

et al. 2019), and interspecific variation (Benedetti et al. 2018). Our analysis also highlighted the utility

of combining primary and secondary data sources in assessing the occurrence of mimicry. By using a

Bayesian model to integrate these data, we were able to generate more robust estimates of mimicry

occurrence across species, reducing uncertainty in our findings. The composite evidence score, which

converts xeno-canto recordings into a standardized measure comparable to traditional literature-based

records, provides a valuable tool for assessing the hidden probabilities of mimicry within under-studied

species. This model also offers insights into how research attention and available data influence our

understanding of mimicry, suggesting that species with more extensive documentation are more likely

to be perceived as mimics, while those with less attention may still exhibit mimicry but remain

underreported. These calculations are useful starting places for future fieldwork focusing on vocal

mimicry and can already be subjected to an empirical test. White-necked Ravens (*Corvus albicollis*) are

not described as mimics in either primary or secondary data sources used here, though based on only

three entries in the corvid literature database and 57 Xeno-Canto recordings. Given this, the model

predicts white-necked ravens to have a 71 % chance of being a hidden mimic, among the more likely

candidates, and indeed vocal mimicry of human speech has been documented on social media (Bucalo

2015).

Another novel aspect the Bayesian modelling allowed us to do is performing several robustness checks

to our interpretation of data. For example, supplying the empirical estimates and evidence distribution

as a simulation input for 1,280 corvid species, which presents ten times the real number of species in

the group, the posterior estimates of occurrence of mimicry consistently approach the true value of

each data sources (primary only, secondary only, primary and secondary). We have also scored the

reliability of each report of vocal mimicry as high, moderate or low. Most recordings were rated as high

reliability; however mimicry was only identified on a low number of low reliability recordings, hence

increasing the likelihood of type I error. To check for sensitivity of false positives, we built models under

the assumption that all low reliability records are false and another model that only high reliability

records are true. Changes in the model did not substantially shift the estimate of occurrence, with still

being above 70% in the most conservative false positive models. This means we can assume the

evidence presented in the current study to be robust.

Corvids mimicked a wide range of different sounds, including calls from birds of prey or other non-

predatory bird species, human speech, and mammalian vocalisations. The fact that corvids are capable

of mimicking such a wide range of sounds further highlights their cognitive flexibility and auditory

learning abilities. The type of sound mimicked can provide insights regarding potential functions of

vocal mimicry (Gammon 2013). Interestingly, contrary to our expectations, we found no evidence that

predator-related sounds were mimicked more often than non-predator sounds. This result contrasts

with some studies in other avian species, where mimicry of predator calls has been hypothesized to

serve as a defensive or deceptive strategy (Igic et al. 2015). On the other hand, we found smaller

species more often to be identified as mimics compared to larger corvid species, with body size being

linked to risk of predation, this could suggest that vocal mimicry could serve a 'deterrent', e.g. anti-

predator and decrease competition, function in corvids. This is supported by anecdotal reports in the

literature, for example Steller's jays mimicked hawk calls while being mobbed by hummingbirds, as

reported in the notes on xeno-canto and green jays are reported to disperse a foraging group of plain

chachalacas (*Ortalis vetula*), whereupon jay moved in to consume seeds being eaten by chachalaca

(Gayou 2020) and in Canada jays most sounds mimicked are known or potential predators of adults or

nests, hence mimicking predators is assumed to serve as a warning call to other group members, to

confuse the predator itself, or merely to signal in a general way that a threat is present (Strickland and

Ouellet 2020). Blue jays, *Cyanocitta cristata,* have been observed to mimic several species of birds of

prey and other bird species, in order to make other species to leave food sources, *e.g.*, feeder and

therefore reduce competition (Hailman 2009). This suggests vocal mimicry in corvids to have an

adaptive evolutionary function and heterospecific sounds might not, as previously suggested, be

learned by mistake (Kelley et al. 2008), however in order to investigate functional contexts of vocal

mimicry behavioural observations of context or experimental tests on behavioural responses of

receivers are necessary. An overabundance of mimicry of sounds from alarm and aggressive contexts

has been also described for example in spotted bowerbirds, *Ptilonorhynchus maculatus* (Kelley and

Healy 2011). Fork-tailed drongos have been described to mimic in order to deceive heterospecific, for

example to lead pied babblers, *Turdoides bicolor,* and meerkats, *Suricata suricatta,* away from food

sources in order to steal the food (Flower et al. 2014). Mimicry by female birds has been observed to

deter nest predators and protect eggs and nestlings (Igic et al., 2015).

Interestingly, there were more recordings of mimicry on Xeno-Canto recorded outside of the breeding

season compared to the breeding season. Sexual selection has been proposed as one of the potential

selective drivers of vocal mimicry, especially in birdsong, where mimicked vocalisations could provide

an honest indicator of male quality (Dalziell et al. 2015). We did not find empirical support of this

hypothesis, although we have to acknowledge that our categorization of breeding season was done per

species and based on rough estimates, ignoring potential geographical variation between species.

However, the fact that recordings of vocal mimicry existed year-round on Xeno-Canto is encouraging to

support our claim that vocal mimicry is not sexually selected in corvids.

Vocal mimicry has previously been described to be driven by social factors. Parrots for example

preferentially mimic socially relevant vocalisations (Bradbury and Balsby 2016) and budgerigars,

Melopsittacus undulatus, converge contact calls from group members (Hile and Striedter 2000).

Choughs, *Pyrrhocorax pyrrhocoraxc* and *Pyrrhocorax graculus*, have been shown to imitate calls of

individuals they are closely socially bonded with (Sitasuwan and Thaler 1985). In our study, we showed

that the social system (territoriality, family groups and cooperative breeders) does not seem to affect

the occurrence of mimicry. This does not mean that social factors do not affect vocal mimicry in

general. Corvids generally are a group of highly social birds (Wascher, 2018) forming strong social

relationships within their groups (Fraser and Bugnyar 2010) and memorizing vocalisations of social

partners for several years (Boeckle and Bugnyar 2012).

To summarize, our study contributed to a broader understanding of vocal mimicry. Vocal mimicry in

birds has been described to have evolved several times in different families and independently of vocal

learning (Goller and Shizuka 2018). A better understanding of the socio-ecological drivers of the

behaviour offers valuable insights into the evolution of communication systems in birds and other

animals. By investigating mimicry in corvids, we gain an opportunity to explore how complex vocal

behaviours evolve, including vocal learning.

References

Ancillotto L, Pafundi D, Cappa F, et al (2022) Bats mimic hymenopteran insect sounds to deter
predators. Curr Biol 32:R408–R409. https://doi.org/10.1016/j.cub.2022.03.052

Archer CR, Pirk CWW, Carvalheiro LG, Nicolson SW (2014) Economic and ecological implications of
geographic bias in pollinator ecology in the light of pollinator declines. Oikos 123:401–407.
https://doi.org/10.1111/j.1600-0706.2013.00949.x

Augustine BC, Tredick CA, Bonner SJ (2014) Accounting for behavioural response to capture when
estimating population size from hair snare studies with missing data. Methods Ecol Evol
5:1154–1161. https://doi.org/10.1111/2041-210X.12289

Benedetti Y, Slezak K, Møller AP, et al (2018) Number of syllables in cuckoo Cuculus canorus calls: A test
using a citizen science project. Sci Rep 8:12872. https://doi.org/10.1038/s41598-018-31329-1

Bluff LA, Kacelnik A, Rutz C (2010) Vocal culture in New Caledonian crows, Corvus moneduloides. Biol J
Linn Soc Lond 101:767–776. https://doi.org/10.1111/j.1095-8312.2010.01527.x

Boeckle M, Bugnyar T (2012) Long-Term memory for affiliates in ravens. Curr Biol 22:801–806.
https://doi.org/10.1016/j.cub.2012.03.023

Bowler DE, Boyd RJ, Callaghan CT, et al (2025) Treating gaps and biases in biodiversity data as a missing
data problem. Biological Reviews 100:50–67. https://doi.org/10.1111/brv.13127

Bradbury JW, Balsby TJS (2016) The functions of vocal learning in parrots. Behav Ecol Sociobiol 70:293–
312. https://doi.org/10.1007/s00265-016-2068-4

Brecht KF, Hage SR, Gavrilov N, Nieder A (2019) Volitional control of vocalizations in corvid songbirds.
PLoS Biol 17:e3000375. https://doi.org/10.1371/journal.pbio.3000375

Brenowitz EA, Margoliash D, Nordeen KW (1997) An introduction to birdsong and the avian song
system. J Neurobiol 33:495–500

Bucalo P, (2015) Ravens can talk! https://www.youtube.com/watch?v=AfsnHVaScjg, accessed: 2025-
03-22

Clements JF, Rasmussen PC, Schulenberg TS, et al (2024) The eBird/Clements checklist of Birds of the
World: v2024.

Cramp S, Perrins CM (1994) Handbook of the Birds of Europe the Middle East and North Africa. Oxford
University Press, Oxford

Dalziell AH, Welbergen JA, Igic B, Magrath RD (2015) Avian vocal mimicry: a unified conceptual
framework. Biol Rev 90:643–668. https://doi.org/10.1111/brv.12129

Daw F, Beheim BA, Wascher CAF (2025) Evolutionary drivers of caching behaviour in corvids. Anim
Cogn 28:17. https://doi.org/10.1007/s10071-025-01938-1

Deng Z, Lloyd H, Xia C, et al (2019) Components of variation in female common cuckoo calls.
Behavioural Processes 158:106–112. https://doi.org/10.1016/j.beproc.2018.10.007

Dierschke V (2003) Predation hazard during migratory stopover : are light or heavy birds under risk?
Journal of Avian Biology 34:24–29. https://doi.org/10.1034/j.1600-048X.2003.03049.x

Dobkin DS (1979) Functional and evolutionary relationships of vocal copying phenomena in birds. Z
Tierpsychol 50:348–363. https://doi.org/10.1111/j.1439-0310.1979.tb01037.x

Droege G, Töpfer T (2016) The Corvids Literature Database: 500 years of ornithological research from a
crow's perspective.

Duengen D, Fitch WT, Ravignani A (2023) Hoover the talking seal. Curr Biol 33:R50–R52.
https://doi.org/10.1016/j.cub.2022.12.023

Ellison G, Jones M, Cain B, Bettridge CM (2021) Taxonomic and geographic bias in 50 years of research
on the behaviour and ecology of galagids. PLoS ONE 16:e0261379.
https://doi.org/10.1371/journal.pone.0261379

Enggist-Dueblin P, Pfister U (2002) Cultural transmission of vocalizations in ravens, Corvus corax. Anim
Behav 64:831–841. https://doi.org/10.1006/anbe.2002.2016

Evans T, Pigot A, Kumschick S, et al (2018) Determinants of data deficiency in the impacts of alien bird
species. Ecography 41:1401–1410. https://doi.org/10.1111/ecog.03232

Flower TP, Gribble M, Ridley AR (2014) Deception by flexible alarm mimicry in an african bird. Science
344:513–516. https://doi.org/10.1126/science.1249723

Fraser ON, Bugnyar T (2010) Do ravens show consolation? Responses to distressed others. PLoS ONE
5:e10605. https://doi.org/10.1371/journal.pone.0010605

Gammon DE (2013) How is model selection determined in a vocal mimic?: Tests of five hypotheses.
Behav 150:1375–1397. https://doi.org/10.1163/1568539X-00003101

Gammon DE, Altizer CE (2011) Northern mockingbirds produce syntactical patterns of vocal mimicry
that reflect taxonomy of imitated species. J Field Ornithol 82:158–164.
https://doi.org/10.1111/j.1557-9263.2011.00318.x

Garamszegi LZ, Møller AP (2012) Untested assumptions about within-species sample size and missing
data in interspecific studies. Behav Ecol Sociobiol 66:1363–1373.
https://doi.org/10.1007/s00265-012-1370-z

Gayou DC (2020) Green Jay (Cyanocorax yncas), version 1.0. Cornell University Press, Ithaca, NY, USA

Gill F, Donsker D, Rasmussen P (2024) IOC World Bird List (v14.1). https://doi.org/doi :
10.14344/IOC.ML.14.1.

Goller M, Shizuka D (2018) Evolutionary origins of vocal mimicry in songbirds. Evol Lett 2:417–426.
https://doi.org/10.1002/evl3.62

Goodale E, Kotagama SW (2006a) Context-dependent vocal mimicry in a passerine bird. Proc R Soc B
273:875–880. https://doi.org/10.1098/rspb.2005.3392

Goodale E, Kotagama SW (2006b) Vocal mimicry by a passerine bird attracts other species involved in
mixed-species flocks. Anim Behav 72:471–477. https://doi.org/10.1016/j.anbehav.2006.02.004

Goodwin D (1951) A comparative study of the voice and some aspects of behaviour in two old-world
jays. Behav 4:293–316. https://doi.org/10.1163/156853951X00188

Hadfield JD (2010) MCMC methods for multi-response generalized linear mixed models: The
MCMCglmm R package. J Stat Softw 33:1–22

Hailman JP (1990) Blue jay mimics osprey. Fla Field Nat 18:81–82

Hailman JP (2009) Context of Blue jay (Cyanocitta cristata) mimicking Cooper's hawk (Accipiter
cooperii) cackle. Fla Field Nat 37:94

Hebblewhite M, Haydon DT (2010) Distinguishing technology from biology: a critical review of the use
of GPS telemetry data in ecology. Phil Trans R Soc B 365:2303–2312.
https://doi.org/10.1098/rstb.2010.0087

Hile AG, Striedter GF (2000) Call convergence within groups of female budgerigars (Melopsittacus
undulatus). Ethology 106:1105–1114. https://doi.org/10.1046/j.1439-0310.2000.00637.x

Hope S (1980) Call form in relation to function in the Steller's jay. Am Nat 116:788–820.
https://doi.org/10.1086/283670

Igic B, McLachlan J, Lehtinen I, Magrath RD (2015) Crying wolf to a predator: deceptive vocal mimicry
by a bird protecting young. Proc R Soc B 282:20150798.
https://doi.org/10.1098/rspb.2015.0798

Janik VM, Knörnschild M (2021) Vocal production learning in mammals revisited. Phil Trans R Soc B
376:20200244. https://doi.org/10.1098/rstb.2020.0244

Kaplan G (1999) Song structure and function of mimicry in the Australian magpie (Gymnorhina tibicen):
Compared to lyrebird (Menura ssp.). Int J Comp Psych 12:. https://doi.org/10.46867/C4J30H

Kelley LA, Coe RL, Madden JR, Healy SD (2008) Vocal mimicry in songbirds. Anim Behav 76:521–528.
https://doi.org/10.1016/j.anbehav.2008.04.012

Kelley LA, Healy SD (2011) The mimetic repertoire of the spotted bowerbird Ptilonorhynchus
maculatus. Naturwissenschaften 98:501–507. https://doi.org/10.1007/s00114-011-0794-z

Madge S, Burn H (1994) Crows and jays: a guide to the crows, jays and magpies of the world. A&C Black

Michonneau F, Brown JW, Winter DJ (2016) rotl: an R package to interact with the Open Tree of Life
data. Methods Ecol Evol 7:1476–1481. https://doi.org/10.1111/2041-210X.12593

Morton ES (1976) Vocal mimicry in the thick-billed euphonia. Wilson Bull 88:485–487

Nakagawa S (2015) Missing Data: Mechanisms, Methods and Messages. In: Ecological statistics:
Contemporary Theory and Application. Oxford University Press, Oxford

Nieder A, Mooney R (2020) The neurobiology of innate, volitional and learned vocalizations in
mammals and birds. Phil Trans R Soc B 375:20190054. https://doi.org/10.1098/rstb.2019.0054

Noble DWA, Nakagawa S (2018) Planned missing data design: stronger inferences, increased research
efficiency and improved animal welfare in ecology and evolution

Oatley TB (1969) The functions of vocal imitation by African cossyphas. Ostrich 40:85–89.
https://doi.org/10.1080/00306525.1969.9639112

OpenTreeOfLife, Redelings B, Sanchez Reyes LL, et al (2019) Open tree of life synthetic tree

Ovaskainen O, Moliterno De Camargo U, Somervuo P (2018) Animal Sound Identifier (ASI): software for
automated identification of vocal animals. Ecology Letters 21:1244–1254.
https://doi.org/10.1111/ele.13092

Poole AF, Gill FB (2020) Birds of the World. Cornell Lab of Ornithology, Ithaca, NY, USA

R Core Team (2019) R: a language and environment for statistical computing.

Ratnayake CP, Goodale E, Kotagama SW (2010) Two sympatric species of passerine birds imitate the
same raptor calls in alarm contexts. Naturwissenschaften 97:103–108.
https://doi.org/10.1007/s00114-009-0617-7

Ridgway S, Carder D, Jeffries M, Todd M (2012) Spontaneous human speech mimicry by a cetacean.
Curr Biol 22:R860–R861. https://doi.org/10.1016/j.cub.2012.08.044

Roberts DL, Taylor L, Joppa LN (2016) Threatened or data deficient: assessing the conservation status of
poorly known species. Diversity and Distributions 22:558–565.
https://doi.org/10.1111/ddi.12418

Sitasuwan N, Thaler E (1985) Vocal inventory and communication in the Chough (Pyrrhocorax
pyrrhocorax), in the Alpine Chough (P. graculus) and in their hybrids. J Ornithol 126:181–193

Stoeger AS, Mietchen D, Oh S, et al (2012) An asian elephant imitates human speech. Curr Biol
22:2144–2148. https://doi.org/10.1016/j.cub.2012.09.022

Strickland D, Ouellet HR (2020) Canada Jay (Perisoreus canadensis). In: Billerman SM, Keeney BK,
Rodewald PG, Schulenberg TS (eds) Birds of the World. Cornell Lab of Ornithology

Suthers RA, Zollinger SA (2004) Producing song: The vocal apparatus. Ann N Y Acad Sci 1016:109–129.
https://doi.org/10.1196/annals.1298.041

Taylor AH (2014) Corvid cognition. WIREs Cogn Sci 5:361–372. https://doi.org/10.1002/wcs.1286

Ter Haar SM, Fernandez AA, Gratier M, et al (2021) Cross-species parallels in babbling: animals and
algorithms. Phil Trans R Soc B 376:20200239. https://doi.org/10.1098/rstb.2020.0239

Tippin TC (2017) Propensity of predator mimicry in wild Steller's jays. The Faculty of Humboldt State
University

Wascher CAF, Reynolds S (2025) Vocal communication in corvids: a systematic review. Animal
Behaviour 221:123073. https://doi.org/10.1016/j.anbehav.2024.123073

West MJ, Stroud N, King AP (1983) Mimicry of the human voice by European starlings: The role of social
interaction. Wilson Bull 95:635–640

Woolfenden GE, Fitzpatrick JW (2020) Florida Scrub-Jay (Aphelocoma coerulescens). In: Birds of the
World. Cornell Lab of Ornithology, Ithaca, NY, USA

541