1 **Vocal mimicry in corvids**

2 Claudia A.F. Wascher[1]*, Gemini Waterhouse[1] & Bret A. Beheim[2]

3 [1]Behavioural Ecology Research Group, School of Life Sciences, Anglia Ruskin University, United Kingdom
4 [2] Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6,
5 04103 Leipzig, Germany

*Corresponding author: Claudia A.F. Wascher: School of Life Sciences, Anglia Ruskin University, East

Road, Cambridge, CB1 1PT United Kingdom; Phone: +44 (0) 1223 698270;

e-mail: claudia.wascher@gmail.com

Abstract

Vocal mimicry, the copying of sounds produced by another species or the environment, is commonly

described in vocal learners, such as songbirds. Understanding the functions of vocal mimicry can help

to uncover the evolutionary drivers of vocal learning. Different adaptive functions like interspecific and

intraspecific communication have been suggested, as well as the possibility of vocal mimicry to be a

mistake during vocal learning. In the present study, we review the occurrence of mimicry in the family

of corvids and investigate the socio-ecological factors driving the evolution of vocal mimicry in this

group of birds. We recorded evidence of vocal mimicry from primary (xeno-canto recordings) as well as

secondary sources (published literature) and found evidence for vocal mimicry in 39 out of 128 corvid

species (30%). Socio-ecological factors like breeding status, habitat and trophic niche did not have a

significant effect. We used Bayesian modelling based on existing data from primary and secondary

sources to estimate the occurrence of mimicry, suggesting that vocal mimicry may be more widespread

among corvids than currently documented, with many species potentially being 'hidden mimics'. Our

study for the first time systematically reviews the occurrence of vocal mimicry across the family of

corvids and investigates a range of socio-ecological factors driving the behaviour, hopefully inspiring

future field work on those species.

56 **Key words:** *Corvidae*, deterrent mimicry, heterospecific vocal mimicry, vocal communication,

Introduction

Vocal mimicry, the copying of sounds produced by another species or the environment, is commonly described in songbirds (Kelley et al. 2008; Dalziell et al. 2015; Goller and Shizuka 2018) and also has been reported in mammals including an Asian elephant, *Elephas maximus* (Stoeger et al. 2012), a white whale, *Delphinapterus leucas* (Ridgway et al. 2012), a harbor seal, *Phoca vitulina* (Duengen et al. 2023), and greater mouse-eared bats, *Myotis myotis* (Ancillotto et al. 2022). Mimics usually acquire sounds through vocal production learning, the ability to produce novel signals through imitation or modification of existing vocalisations (Janik and Knörnschild 2021). Vocal production learning in animals has parallels with human vocal ontogeny (Ter Haar et al. 2021), as it requires an individual to process acoustic information from an appropriate model, engage in vocal practice and evaluation through auditory feedback, and requires enhanced forebrain control of the vocal organ (Nieder and Mooney 2020).

According to Dalziell et al. (2015), vocal mimicry is defined as production of vocalisations resembling heterospecific or environmental sounds which change the behaviour of the receiver after perceiving the acoustic resemblance between the mimic and the model, and the behavioural change confers a selective advantage on the species mimicking. This definition excludes other forms for vocal resemblance, for example as a result of chance, common ancestry (Goller and Shizuka 2018) or social learning (Bluff et al. 2010) . Vocal mimicry may be a result of mistakes during vocal learning and have no function (Kelley et al., 2008). In contrast to this, a number of different potential functions of vocal mimicry have been previously suggested, broadly falling into two categories (a) interspecific communication, to avoid threats, dissuade competitors, or imitate threats to allow access to food resources and (b) intraspecific communication, in the context of sexual selection or for social affiliation (reviewed by Kelley et al., 2008). For example, mimics are suggested to produce vocalisations from predatory or agonistic species to deter competitors or potential predators from their territories (Dobkin 1979). Australian magpies, *Gymnorhina tibicen*, mimic several potential nest predators, for example the barking owl, *Ninox connivens*, and the boobook owl, *Ninox novaeseelandiae* (Kaplan 1999). Fork-tailed drongos, *Dicrurus adsimilis,* mimic alarm calls of other species to deter heterospecifics from food sources, in order to steal the food (Flower et al. 2014). Individuals might also benefit from vocal mimicry when used to attract other individuals. For example, greater racket-tailed drongos, *Dicrurus paradiseus,* mimic songs and contact calls to attract other species, which increases foraging efficiency (Goodale and Kotagama 2006a) and alarm calls of other species alongside their own mobbing calls, potentially attracting other species to support them when mobbing predators (Goodale and Kotagama 2006b). This shows that some species flexibly use vocal mimicry, and this may be context dependent. Finally, mimicked calls are sometimes incorporated into song or sexual displays and might play a role in

sexual selection (Kelley et al., 2008). Overall, surprisingly little empirical research on the functional

explanations of vocal mimicry is available.

Phylogenetic analysis has previously shown that vocal mimicry is unlikely to be an ancestral state in

songbirds, meaning that it may have evolved independently on multiple occasions (Goller and Shizuka

2018). A review of European passerines estimated approximately 40% of songbird species to mimic

(Garamszegi et al., 2007). Sounds mimicked by songbirds include birdsong (northern mockingbird,

Mimus polyglottos: Gammon and Altizer 2011), alarm calls of other species (spotted bowerbird,

Ptilonorhynchus maculatus: Kelley and Healy 2011; red-capped robin-chat, *Cossypha natalensis*: Oatley

1969; thick-billed euphonia, *Euphonia laniirostris*: Morton 1976), calls of birds of prey (blue jay,

Cyanocitta cristata: Hailman 1990), and human voices (European starlings, *Sturnus vulgaris*: West et al.

1983). As suggested by Gammon (2013), the 'model selection', *i.e.* type of sound mimicked, can provide

valuable insight into the potential functions of vocal mimicry. For example, northern mockingbirds,

Mimus polyglottos, preferentially mimic species whose sounds are acoustically similar to their non-

imitative song, supporting the acoustic similarity hypothesis, and suggesting a non-adaptive

explanation of vocal mimicry (Gammon, 2013). Mimicry of predator calls by Steller's jays, *Cyanocitta*

stelleri, is also linked to mate selection, with mimicry occurring more frequently during the breeding

season and when in the presence of their mates (Tippin 2017).

Corvids *(Aves: Corvidae),* are a large and diverse group with over 120 species split between 21 genera,

an array of which can be found on all continents except Antarctica (Madge and Burn 1994). Many

corvid species have demonstrated advanced cognitive abilities in the social and physical domain (Taylor

2014). Corvids belong to the suborder of oscine passerine birds, where the vocal organ (syrinx) is highly

developed to produce vocalisations (Suthers and Zollinger 2004). Vocalisations in corvids present a

wide range of ecological and social adaptations, such as communicating information about predators

such as predator type or level of risk or individual recognition (Wascher and Reynolds 2025). Corvids

are open-ended vocal learners, which means they can acquire new vocalisations throughout their lives

and not only in a specific sensitive period (Brenowitz et al. 1997). Carrion crows are capable of

volitional control of vocalisations, *i.e.* can be trained to vocalise and withhold vocalisations in response

to arbitrary stimuli (Brecht et al. 2019). Vocalisations in New Caledonian crows, *Corvus moneduloides*

(Bluff et al. 2010) and common ravens, *Corvus corax* (Enggist-Dueblin and Pfister 2002) are socially

learned from conspecifics. Several species of corvids have demonstrated the ability to mimic human

speech when hand-raised and kept in close proximity to humans (Bluff et al., 2010). Mimicry of

predator calls has been documented in several corvid species including Steller's jays (Hope 1980; Tippin

2017), Sri Lanka blue-magpies, *Urocissa ornata* (Ratnayake et al. 2010), Florida scrub-Jays, *Aphelocoma*

coerulescens (Woolfenden and Fitzpatrick 2020), blue jays, *Cyanocitta cristata* (Hailman 1990), and

Eurasian jays, *Garrulus glandarius* (Goodwin 1951). Tippin (2017) conducted an in-depth study of vocal

mimicry in Steller's jays, which found significant differences in age and sex between mimicking and

non-mimicking individuals, where males were less likely to mimic compared to females, and the

likelihood of an individual mimicking decreased with age.

Missing data present a significant challenge in the study of animal behaviour, limiting our

understanding of key phenomena (Garamszegi and Møller 2012; Nakagawa 2015). Factors such as

habitat inaccessibility (Evans et al. 2018), species rarity (Roberts et al. 2016), human-centric biases in

study design (Archer et al. 2014; Ellison et al. 2021; Bowler et al. 2025; Daw et al. 2025), and

technological constraints (Hebblewhite and Haydon 2010) contribute to missingness. In evolutionary

biology, missing data can skew our understanding of trait evolution, obscuring the selective pressures

and socio-ecological contexts that drive behavioural adaptations. The problem of missing data can be

addressed with more systematic data collection, e.g., planned missing data design (Noble and

Nakagawa 2018), application of emerging technologies and statistical approaches, such as imputation

procedures (Augustine et al. 2014), and a recognition that the absence of evidence is not evidence of

absence in behavioural research.

The aim of the present study is to describe the occurrence of mimicry in corvids. We investigated the

evidence for mimicry from primary, i.e. recordings on xeno-canto, and secondary sources, i.e. published

literature. Further, we investigate socio-ecological factors affecting the occurrence of vocal mimicry in

corvids. Vocal mimicry in corvids might serve a function of interspecific communication, such as

deterring a potential predator and competitors, by copying the vocalisations of animals that are

predatory or agonistic (Dobkin 1979). If this is the case, we expect smaller corvid species to mimic

more, as they are more at risk of predation (Dierschke 2003) and we expect them to mainly mimic birds

of prey. If vocal mimicry serves intraspecific communication, such as sexual selection (Kelley et al.

2008; Dalziell et al. 2015), we expect a positive correlation between the size of the vocal repertoire and

the occurrence of vocal mimicry. Additionally, we investigate whether habitat, trophic niche and

breeding system affect the occurrence of mimicry in corvids, although we do not have specific

predictions in this regard and consider our analysis exploratory. In addition to biological factors driving

vocal mimicry in corvids, species differ significantly in terms of study effort (Wascher & Reynolds,

2025). We investigate how differences in recording effort affect the detection of mimicry in corvids and

evaluate the likelihood of 'hidden' mimicry.

163
164

Methods

Data on vocal mimicry

The present study was approved by the School of Life Sciences departmental ethics panel at Anglia Ruskin University (ETH2223-3747). This is a desk-based study, and no animals have been used for the research. Data were compiled from a variety of primary (xeno-canto) and secondary sources (peer-reviewed journal articles and books). Scientific and common names were standardized using a combination of the IOC Bird list version 14.1. (Gill et al. 2024) and the 'Clements checklist' (Clements et al. 2024) and 128 species of corvids were included. A xeno-canto advanced search was conducted between the 27th of April and the 1st of May 2023 (searches have been updated on the 22nd of June 2023 and 5th May 2024). Xeno-canto is a citizen science project and repository in which volunteers record, upload and annotate recordings of bird calls (www.xeno-canto.org, last accessed on 29th July 2025). The xeno-canto search yielded a total of 19,566 audio recordings of corvid calls from 128 different species (mean±SD per species: 153 ± 360), belonging to 21 genera (mean±SD per genus: 889 ± 1,767). The number of recordings per species ranged from zero to 2,272 recordings. All recordings with mentions of 'mimicry', 'imitation' or 'production of heterospecific calls' in the comments or labels section were identified. For the purpose of this study, we defined vocal mimicry as the occurrence of any type of non-conspecific sound and a resemblance between the mimicked and model sound (Kelley et al. 2008; Goller and Shizuka 2018). Our definition does not require the receivers to respond behaviourally to mimicked sound in a way that benefits them, which is the case for the definition of vocal mimicry in Dalziell et al. (2015). We noted the duration of the recording, date, latitude and longitude and type of sound mimicked. Recordings were then listened to by GW and CAFW, and if both observers agreed that they contained convincing evidence for production of mimicry the observation was considered as evidence of mimicry. We fully acknowledge that this process involves a high level of subjectivity and a risk of false positives, i.e., the recordings not being a case of mimicry in corvids. In order to account for this, we have rated the perceived 'reliability' of the recordings, indicating whether credibility of the recording was high, moderate or low. A total of 466 records of mimicry have been identified. For 350 records it was indicated that the vocalising bird was seen by the recorder, for 86 records the bird was not observed and for 30 records this information was unknown. The majority of records (325) have been classed as 'high' reliability, meaning that the rater (CAFW) had high confidence in the recording containing mimicry by a corvid, 53 records were classed as 'moderate' reliability and 88 records as 'low' reliability, meaning the record is more likely to be a false positive.

Data were compiled from a range of primary and secondary literature sources. A search was conducted using Google Scholar, combining search terms such as 'mimic', 'imitate', and 'copy' with both English and Latin species names. In addition, we consulted authoritative field guides and handbooks, including

for example the *Handbook of the Birds of the World* and *Birds of the World*, which were manually

reviewed for references to vocal mimicry or imitation. Only sources that explicitly used terms such as

"mimic" or "imitate" and clearly identified the mimicking species and the target of mimicry (e.g.,

species, family, order, or group such as 'environmental sounds') were included. We excluded any

instances where vague descriptors like 'similar to' or 'reminiscent of' were used. We have included

reports of mimicry from six books and field-guides (non-peer reviewed sources) and 11 peer-reviewed

journal articles. A full list of sources is provided in the online supplementary material.

For each species, the number of entries in the corvids literature database (Droege and Töpfer 2016)

plus one entry in Birds of the World (Poole & Gill, 2020) was used as an estimate of the volume of

secondary literature available (range: 1-423 entries). For each piece of evidence of vocal mimicry,

CAFW allocated a 'reliability score' of high, moderate or low. Sixty-five percent of recordings have been

rated as 'high', 23% as 'moderate' and 12% as 'low' reliability.

For each of the 128 species, we recorded body mass (information available for 121 species), vocal

repertoire (98 species), breeding system (cooperative breeding, groups, territorial pairs; reported for

121 species), habitat (coastal, desert, forest, grassland, human modified, rock, shrubland, woodland;

reported for 124 species) and trophic niche (frugivore, granivore, invertivore, omnivore; reported for

124 species).

Statistical analysis

All analyses were conducted in R version 4.3.1. (R Core Team 2019). We constructed a phylogenetic

tree for the Corvidae family using data from the Open Tree of Life (OTL). The tree was retrieved from

the Open Tree of Life database (OpenTreeOfLife et al. 2019) using the rotl R package (Michonneau et

al. 2016). We employed Bayesian generalized linear mixed-effects models built with the MCMCglmm

package (Hadfield 2010) to determine effects of body size, vocal repertoire size, habitat and trophic

niche on vocal mimicry in corvids. Genus was included as a random factor and the OTL tree served as

the foundational phylogeny for the MCMCglmm. We ran Markov chain Monte Carlo (MCMC) chains for

3,000,000 iterations, thinned by 2000, and employed a burn-in of 2,000,000. All main effects were

given normal(0, 0.5) log-odds prior, with a normal(0, 1) log-odds intercept prior. Random effects by

genus were given an inverse Wishart prior with V = 1 and nu = 10. Data and code of the present study

are published as online supplementary materials (Supplementary File 2; supplementary File 3).

In order to estimate the occurrence of mimicry, we have developed a Bayesian model of evidence

updating, assuming that for any specific kind of evidence (citation database entries, audio recordings),

there is a corresponding rate at which mimics are detected per-unit evidence. The model enables the

calculation of the posterior probability a species in which no mimicry is detected is, in fact, a mimic,

which is a more principled way to handle the presence of zeros in such datasets than taking them at

face value (Beheim et al. 2021; Sigmundson et al. 2025). Full details of the model are presented in the

supplementary materials, including several robustness checks for variable reporting rate and quality

across species, insensitivity to priors, and the effects of reliability scores on evidence accumulation.

Results of these checks are presented in the online supplementary materials. The present manuscript

was posted as pre-print and full statistical analysis code is available in the accompanying materials

repository (Wascher et al. 2025; https://github.com/babeheim/corvid-mimicry).

<u>Results</u>

Description of the dataset

We recorded evidence of vocal mimicry in 39 species of *Corvidae*, belonging to thirteen genera (30% of

species; Figure 1). Thirty-three species of corvids are described as mimics in the literature and

recordings from 17 species have been identified on xeno-canto, including six species not previously

described as mimics (Indochinese green-magpie, *Cissa hypoleuca*; carrion crow, *Corvus corone*; black-

chested jay, *Cyanocorax affinis;* Yucatan jay, *Cyanocorax yucatanicus;* clark's nutcracker*, Nucifraga*

Columbiana; rook, *Corvus frugilegus;* Table 1). One to 71 different types of sounds were mimicked

(mean ± standard deviation: 5.512 ± 11.568). Sounds have been mimicked from a wide range or birds

(Accipitriformes, Anseriformes, Charadriiformes, Cuculiformes, Falconiformes, Galliformes,

Gaviiformes, Gruiformes, Pelecaniformes, Passeriformes, Piciformes, Strigiformes), mammals

(Artiodactyla, Carnivora, Perissodactyla, Primates (human voice), Rodentia) amphibian (Anura) and

insect species (Hemiptera) as well as anthropogenic sounds, such as bells, lawnmower, motorbike (see

supplementary materials for full list of mimicked species). Body mass, repertoire size, habitat, breeding

system and trophic niche did not affect mimicry status (Table 2).

Figure 1: Vocal mimicry in 128 species of corvids. Grey points indicate species with no mimicry

in the literature, blue points illustrate species with recordings of mimicry on xeno-canto but no records

in the literature and yellow dots indicate species only reported as mimics in the literature, but no

recordings of mimicry are available on xeno-canto.

Table 1: Cross-table indicating the number of species detected as mimics in the literature and on xeno-

canto. Records per species indicate the number of records in the corvid literature database and xeno-

canto.

mimicry category	n.species	number of records in the corvid literature database	number of records on xeno-canto
no mimicry seen	89	11	51
number of species detected as mimics in the literature only	22	29	203
number of species detected as mimics on xeno-canto only	6	126	477
number of species detected as mimics both in the literature and on xeno-canto	11	94	697

277 **Table 2.** Results of Markov chain Monte Carlo generalized linear mixed-effect model (MCMCglmm).

278 Factors with significant effects ($p \leq 0.05$) are shown in bold.

parameters	post.mean	l-95% CI	u-95% CI	eff.samp	pMCMC
(intercept)	-0.014	-1.935	2.178	500	0.928
body mass	-0.137	-0.373	-0.026	500	0.068
repertoire	0.161	-0.917	1.426	429.547	0.768
habitat breadth (generalist relative to hypergeneralist)	0.011	-1.374	1.501	500	0.996
habitat breadth (generalist relative to specialist)	-0.022	-1.41	1.263	500	0.996
breeding system (cooperative breeding relative to groups)	-0.012	-1.356	1.361	500	0.996
breeding system	-0.034	-1.391	1.217	628.116	0.952

(cooperative breeding relative to territorial pairs)					
trophic niche (frugivore relative to granivore)	-0.018	-1.397	1.298	500	0.984
trophic niche (frugivore relative to invertivore)	-0.007	-1.473	1.504	356.051	0.964
trophic niche (frugivore relative to omnivore)	0.003	-1.459	1.453	500	0.976

Estimating the occurrence of mimicry

Because species with more evidence are more likely to be reported as mimics, the observed rate of occurrence of mimicry in the published literature (33/128 species) and in the xeno-Canto recordings (17/128 species) are likely to undercount the true occurrence of mimicry, as species that are capable of mimicry have not (yet) been documented as such. In order to formalize this intuition, we developed a Bayesian model of evidence accumulation that can accurately estimate the true rate of occurrence of a phenomenon with variable rates of reporting, detection and source quality (see the supplementary information for full specification). Figure 2 shows how different information sources update prior assumptions about the occurrence of mimicry across corvids. When both primary and secondary sources are incorporated into the model, each with their own detection rates, we estimate that mimicry occurs in 82% (+/- 7%) of *Corvidae*.

296 Figure 2: Prior and posterior estimates (means and 89% HPDI) of the frequency of occurrence of
297 mimicry across corvid species conditioning on different information evidence (primary sources,
298 secondary sources, and both combined).
299

Using this model, we can also calculate a composite evidence score, in practice converting a count of

recordings on xeno-canto into the equivalent number of corvid database entries (or vice versa), which

is useful to understand how the hidden mimicry probabilities for each species relates to research

attention (Figure 3). When very little is known about a particular species (e.g. no uploaded recordings

or entries in the citation database), our model assumes that the best guess about their mimicry ability

is the corvid-wide intercept in the model, q. As information accumulates with no positive detection of

mimicry, the species-specific posterior estimate declines according to the mathematical relationship

described in the supplement. Thus, although most species are not detected as mimics, only a handful

actually have sufficient evidence to update species-specific posteriors near zero probability (Figure 3).

310
311 Figure 3: Composite evidence score related to the probability of hidden mimicry.

313 Looking at posterior probabilities of hidden mimicry for each species individually. The ten species which

314 are predicted as least likely to be mimics and ten species nominated as most likely to be mimic are

315 presented in the supplementary materials.

Discussion

In the present study, we describe cases of vocal mimicry in 39 species of corvids belonging to thirteen genera, showing that vocal mimicry is widespread across this family of birds. With 30% of corvids known to mimic, it is more prevalent in this group, as opposed to the 8.9% observed in the entire suborder of songbirds (Goller and Shizuka 2018). Additionally, because mimicry detection is strongly associated with the amount of research effort directed at any specific corvid species, there is reason to believe that the occurrence of vocal mimicry is starkly underestimated by the current empirical evidence and many species of corvids are likely to be 'hidden mimics', meaning they are undetected as mimics by the currently available literature and data.

Our study underscores the importance of crowd-sourced databases, such as xeno-canto, revealing potentially overlooked or underreported behaviours, such as vocal mimicry. We have identified five species of corvids as mimics based on xeno-canto recordings, which have not previously been reported as mimics. Xeno-canto has previously been used scientifically to develop machine learning methods for automatic detection and classification (Ovaskainen et al. 2018), describing geographic variation (Deng et al. 2019), and interspecific variation (Benedetti et al. 2018). Our analysis also highlighted the utility of combining primary and secondary data sources in assessing the occurrence of mimicry. By using a Bayesian model to integrate these data, we were able to generate more probabilistic estimates of mimicry occurrence across species, reducing uncertainty in our findings. The composite evidence score, which converts xeno-canto recordings into a standardized measure comparable to traditional literature-based records, provides a valuable tool for assessing the hidden probabilities of mimicry within under-studied species. This model also offers insights into how research attention and available data influence our understanding of mimicry, suggesting that species with more extensive documentation are more likely to be perceived as mimics, while those with less attention may still exhibit mimicry but remain underreported. These calculations are useful starting places for future fieldwork focusing on vocal mimicry and can already be subjected to an empirical test.

Another novel aspect the Bayesian modelling allowed us to do is performing several robustness checks to our interpretation of data. For example, supplying the empirical estimates and evidence distribution as a simulation input for 1,280 corvid species, which presents ten times the real number of species in the group, the posterior estimates of occurrence of mimicry consistently approach the true value of each data sources (primary only, secondary only, primary and secondary). Potentially low data quality, is a persisting challenge in citizen science projects (Kosmala et al. 2016), including the use of data from xeno-canto. For example, misclassification and 'noisy' data can significantly hinder application of machine learning on citizen science datasets (Gupta and Gupta 2019). However, depending on type of

application, unscreened crowd contributors also can be as effective as experts in data annotation

(Cocos et al. 2017). In order to evaluate the impact of potentially low data quality, potentially increase

the number of false positives and hence us over-estimating the occurrence of mimicry in corvids, we

have scored the perceived reliability of each report of vocal mimicry as high, moderate or low. Most

recordings were rated as high reliability; however, in three species only low reliability recordings were

available, hence increasing the likelihood of type I error. To check for sensitivity of false positives, we

built models under the assumption that all low reliability records are false and another model that only

high reliability records are true. Changes in the model did not substantially shift the estimate of

occurrence, which remained above 80% in the most conservative false positive models. This means we

can assume the evidence presented in the current study to be robust.

We did not find body mass or other socio-ecological factors such as breeding system or habitat to

affect mimicry status, hence we cannot draw conclusions about potential functions of vocal mimicry in

corvids. Anecdotal reports in the literature suggest a potential 'deterrent' function, e.g. anti-predator

or to decrease competition. For example Steller's jays mimicked hawk calls while being mobbed by

hummingbirds, as reported in the notes on xeno-canto. Similarly, green jays, *Cyanocorax yncas*, were

reported to disperse a foraging group of plain chachalacas, *Ortalis vetula*, whereupon jays moved in to

consume seeds being eaten by chachalaca (Gayou 2020). In Canada jays most sounds mimicked are

known or potential predators of adults or nests, hence mimicking predators is assumed to serve as a

warning call to other group members, to confuse the predator itself, or merely to signal in a general

way that a threat is present (Strickland and Ouellet 2020). Blue jays, *Cyanocitta cristata,* have been

observed to mimic several species of birds of prey and other bird species, in order to make other

species to leave food sources, *e.g.*, feeder and therefore reduce competition (Hailman 2009). This

suggests vocal mimicry in corvids to have an adaptive evolutionary function and heterospecific sounds

might not, as previously suggested, be learned by mistake (Kelley et al. 2008). However, in order to

investigate functional contexts of vocal mimicry behavioural observations of context or experimental

tests on behavioural responses of receivers are necessary. An overabundance of mimicry of sounds

from alarm and aggressive contexts has been also described for example in spotted bowerbirds,

Ptilonorhynchus maculatus (Kelley and Healy 2011). Fork-tailed drongos have been described to mimic

in order to deceive heterospecifics, for example to lead pied babblers, *Turdoides bicolor,* and meerkats,

Suricata suricatta, away from food sources in order to steal the food (Flower et al. 2014). Mimicry by

female birds has been observed to deter nest predators and protect eggs and nestlings (Igic et al.,

2015).

Corvids mimicked a wide range of different sounds, including calls from birds of prey or other non-
predatory bird species, human speech, and mammalian vocalisations. The fact that corvids are capable
of mimicking such a wide range of sounds further highlights their cognitive flexibility and auditory
learning abilities. Interestingly, in a number of species corvids have been reported to mimic human
speech in captivity (Landsborough Thomson 1964; Lorenz 2002), whereas such reports are rare from
wild birds. This further supports the high level of flexibility in mimicry. We suggest that while many, if
not all corvid species possess a general capacity for vocal mimicry, the extent to which this ability is
expressed in the wild likely depends on socio-ecological factors such as predation pressure and
competition.
395
Our study was of explorative nature, our capacity to test specific hypotheses regarding the evolutionary
drivers of mimicry in corvids was limited. Vocal mimicry has previously been described to be driven by
social factors. Parrots for example preferentially mimic socially relevant vocalisations (Bradbury and
Balsby 2016). In our study, we showed that the breeding system (territoriality, family groups and
cooperative breeders) does not seem to affect the occurrence of mimicry. This does not mean that
social factors do not affect vocal mimicry in general. Corvids generally are a group of highly social birds
(Wascher, 2018) forming strong social relationships within their groups (Fraser and Bugnyar 2010) and
memorizing vocalisations of social partners for several years (Boeckle and Bugnyar 2012). Because of its
flexibility, the social system of corvids is difficult to characterize, for example a number of species are
facultative cooperative breeders, where the occurrence of cooperative breeding depends on
environmental factors. Additionally, the social system depends on life-history factors, for example
individuals form large flocks outside of the breeding season or when they are young and become
territorial, when they mate and start to reproduce (Wascher 2018). We suggest that because we did
not find an effect of the breeding system on the occurrence of vocal mimicry, this might indicate sexual
selection not to be a main evolutionary driver of vocal mimicry in corvids. Sexual selection has been
proposed as one of the potential selective drivers of vocal mimicry, especially in birdsong, where
mimicked vocalisations could provide an honest indicator of male quality (Dalziell et al. 2015).
413
To summarize, our study contributed to a broader understanding of vocal mimicry. Vocal mimicry in
birds has been described to have evolved several times in different families and independently of vocal
learning (Goller and Shizuka 2018). A better understanding of the socio-ecological drivers of the
behaviour offers valuable insights into the evolution of communication systems in birds and other
animals. By investigating mimicry in corvids, we gain an opportunity to explore how complex vocal
behaviours evolve, including vocal learning.
420

Acknowledgement

We thank all xeno-canto members for sharing their recordings and two anonymous referees for their

constructive advice and helpful comments on our manuscript.

Competing interest

At the time of writing, C.A.F.W. was a member of the Editorial Board of Animal Cognition, but had no

involvement in the review or assessment of the paper. C.A.F.W. received funding from Anglia Ruskin

University, and B.A.B. received funding from the Max Planck Institute for Evolutionary Anthropology.

G.W. did not receive any funding.

References

Ancillotto L, Pafundi D, Cappa F, et al (2022) Bats mimic hymenopteran insect sounds to deter
        predators. Curr Biol 32:R408–R409. https://doi.org/10.1016/j.cub.2022.03.052

Archer CR, Pirk CWW, Carvalheiro LG, Nicolson SW (2014) Economic and ecological
        implications of geographic bias in pollinator ecology in the light of pollinator declines.
        Oikos 123:401–407. https://doi.org/10.1111/j.1600-0706.2013.00949.x

Augustine BC, Tredick CA, Bonner SJ (2014) Accounting for behavioural response to capture
        when estimating population size from hair snare studies with missing data. Methods
        Ecol Evol 5:1154–1161. https://doi.org/10.1111/2041-210X.12289

Beheim B, Atkinson QD, Bulbulia J, et al (2021) Treatment of missing data determined
        conclusions regarding moralizing gods. Nature 595:E29–E34.
        https://doi.org/10.1038/s41586-021-03655-4

Benedetti Y, Slezak K, Møller AP, et al (2018) Number of syllables in cuckoo Cuculus canorus
        calls: A test using a citizen science project. Sci Rep 8:12872.
        https://doi.org/10.1038/s41598-018-31329-1

Bluff LA, Kacelnik A, Rutz C (2010) Vocal culture in New Caledonian crows, Corvus
        moneduloides. Biol J Linn Soc Lond 101:767–776. https://doi.org/10.1111/j.1095-
        8312.2010.01527.x

Boeckle M, Bugnyar T (2012) Long-Term memory for affiliates in ravens. Curr Biol 22:801–
        806. https://doi.org/10.1016/j.cub.2012.03.023

Bowler DE, Boyd RJ, Callaghan CT, et al (2025) Treating gaps and biases in biodiversity data
as a missing data problem. Biological Reviews 100:50–67.
https://doi.org/10.1111/brv.13127

Bradbury JW, Balsby TJS (2016) The functions of vocal learning in parrots. Behav Ecol
Sociobiol 70:293–312. https://doi.org/10.1007/s00265-016-2068-4

Brecht KF, Hage SR, Gavrilov N, Nieder A (2019) Volitional control of vocalizations in corvid
songbirds. PLoS Biol 17:e3000375. https://doi.org/10.1371/journal.pbio.3000375

Brenowitz EA, Margoliash D, Nordeen KW (1997) An introduction to birdsong and the avian
song system. J Neurobiol 33:495–500

Clements JF, Rasmussen PC, Schulenberg TS, et al (2024) The eBird/Clements checklist of
Birds of the World: v2024.

Cocos A, Qian T, Callison-Burch C, Masino AJ (2017) Crowd control: Effectively utilizing
unscreened crowd workers for biomedical data annotation. Journal of Biomedical
Informatics 69:86–92. https://doi.org/10.1016/j.jbi.2017.04.003

Dalziell AH, Welbergen JA, Igic B, Magrath RD (2015) Avian vocal mimicry: a unified
conceptual framework. Biol Rev 90:643–668. https://doi.org/10.1111/brv.12129

Daw F, Beheim BA, Wascher CAF (2025) Evolutionary drivers of caching behaviour in
corvids. Anim Cogn 28:17. https://doi.org/10.1007/s10071-025-01938-1

Deng Z, Lloyd H, Xia C, et al (2019) Components of variation in female common cuckoo calls.
Behavioural Processes 158:106–112. https://doi.org/10.1016/j.beproc.2018.10.007

Dierschke V (2003) Predation hazard during migratory stopover : are light or heavy birds under
risk? Journal of Avian Biology 34:24–29. https://doi.org/10.1034/j.1600-
048X.2003.03049.x

Dobkin DS (1979) Functional and evolutionary relationships of vocal copying phenomena in
birds. Z Tierpsychol 50:348–363. https://doi.org/10.1111/j.1439-0310.1979.tb01037.x

Droege G, Töpfer T (2016) The Corvids Literature Database: 500 years of ornithological
research from a crow's perspective.

Duengen D, Fitch WT, Ravignani A (2023) Hoover the talking seal. Curr Biol 33:R50–R52.
https://doi.org/10.1016/j.cub.2022.12.023

Ellison G, Jones M, Cain B, Bettridge CM (2021) Taxonomic and geographic bias in 50 years
of research on the behaviour and ecology of galagids. PLoS ONE 16:e0261379.
https://doi.org/10.1371/journal.pone.0261379

Enggist-Dueblin P, Pfister U (2002) Cultural transmission of vocalizations in ravens, Corvus
corax. Anim Behav 64:831–841. https://doi.org/10.1006/anbe.2002.2016

Evans T, Pigot A, Kumschick S, et al (2018) Determinants of data deficiency in the impacts of
alien bird species. Ecography 41:1401–1410. https://doi.org/10.1111/ecog.03232

Flower TP, Gribble M, Ridley AR (2014) Deception by flexible alarm mimicry in an african
bird. Science 344:513–516. https://doi.org/10.1126/science.1249723

Fraser ON, Bugnyar T (2010) Do ravens show consolation? Responses to distressed others.
PLoS ONE 5:e10605. https://doi.org/10.1371/journal.pone.0010605

Gammon DE (2013) How is model selection determined in a vocal mimic?: Tests of five
hypotheses. Behav 150:1375–1397. https://doi.org/10.1163/1568539X-00003101

Gammon DE, Altizer CE (2011) Northern mockingbirds produce syntactical patterns of vocal
mimicry that reflect taxonomy of imitated species. J Field Ornithol 82:158–164.
https://doi.org/10.1111/j.1557-9263.2011.00318.x

Garamszegi LZ, Møller AP (2012) Untested assumptions about within-species sample size and
missing data in interspecific studies. Behav Ecol Sociobiol 66:1363–1373.
https://doi.org/10.1007/s00265-012-1370-z

Gayou DC (2020) Green Jay (Cyanocorax yncas), version 1.0. Cornell University Press, Ithaca,
NY, USA

Gill F, Donsker D, Rasmussen P (2024) IOC World Bird List (v14.1). https://doi.org/doi :
10.14344/IOC.ML.14.1.

Goller M, Shizuka D (2018) Evolutionary origins of vocal mimicry in songbirds. Evol Lett
2:417–426. https://doi.org/10.1002/evl3.62

Goodale E, Kotagama SW (2006a) Context-dependent vocal mimicry in a passerine bird. Proc
R Soc B 273:875–880. https://doi.org/10.1098/rspb.2005.3392

Goodale E, Kotagama SW (2006b) Vocal mimicry by a passerine bird attracts other species
involved in mixed-species flocks. Anim Behav 72:471–477.
https://doi.org/10.1016/j.anbehav.2006.02.004

Goodwin D (1951) A comparative study of the voice and some aspects of behaviour in two old-
world jays. Behav 4:293–316. https://doi.org/10.1163/156853951X00188

Gupta S, Gupta A (2019) Dealing with noise problem in machine learning data-sets: A
systematic review. Procedia Computer Science 161:466–474.
https://doi.org/10.1016/j.procs.2019.11.146

Hadfield JD (2010) MCMC methods for multi-response generalized linear mixed models: The
MCMCglmm R package. J Stat Softw 33:1–22

Hailman JP (1990) Blue jay mimics osprey. Fla Field Nat 18:81–82

Hailman JP (2009) Context of Blue jay (Cyanocitta cristata) mimicking Cooper's hawk
(Accipiter cooperii) cackle. Fla Field Nat 37:94

Hebblewhite M, Haydon DT (2010) Distinguishing technology from biology: a critical review
of the use of GPS telemetry data in ecology. Phil Trans R Soc B 365:2303–2312.
https://doi.org/10.1098/rstb.2010.0087

Hope S (1980) Call form in relation to function in the Steller's jay. Am Nat 116:788–820.
https://doi.org/10.1086/283670

Janik VM, Knörnschild M (2021) Vocal production learning in mammals revisited. Phil Trans
R Soc B 376:20200244. https://doi.org/10.1098/rstb.2020.0244

Kaplan G (1999) Song structure and function of mimicry in the Australian magpie (Gymnorhina
tibicen): Compared to lyrebird (Menura ssp.). Int J Comp Psych 12:.
https://doi.org/10.46867/C4J30H

Kelley LA, Coe RL, Madden JR, Healy SD (2008) Vocal mimicry in songbirds. Anim Behav
76:521–528. https://doi.org/10.1016/j.anbehav.2008.04.012

Kelley LA, Healy SD (2011) The mimetic repertoire of the spotted bowerbird Ptilonorhynchus
maculatus. Naturwissenschaften 98:501–507. https://doi.org/10.1007/s00114-011-0794-
z

Kosmala M, Wiggins A, Swanson A, Simmons B (2016) Assessing data quality in citizen
science. Frontiers in Ecol & Environ 14:551–560. https://doi.org/10.1002/fee.1436

Landsborough Thomson A (1964) A New Dictionary of Birds. Nelson, London

Lorenz K (2002) King Solomon's ring: new light on animal ways. Psychology Press, London

Madge S, Burn H (1994) Crows and jays: a guide to the crows, jays and magpies of the world.
A&C Black

Michonneau F, Brown JW, Winter DJ (2016) rotl: an R package to interact with the Open Tree
of Life data. Methods Ecol Evol 7:1476–1481. https://doi.org/10.1111/2041-
210X.12593

Morton ES (1976) Vocal mimicry in the thick-billed euphonia. Wilson Bull 88:485–487

Nakagawa S (2015) Missing Data: Mechanisms, Methods and Messages. In: Ecological
statistics: Contemporary Theory and Application. Oxford University Press, Oxford

Nieder A, Mooney R (2020) The neurobiology of innate, volitional and learned vocalizations in
mammals and birds. Phil Trans R Soc B 375:20190054.
https://doi.org/10.1098/rstb.2019.0054

Noble DWA, Nakagawa S (2018) Planned missing data design: stronger inferences, increased
research efficiency and improved animal welfare in ecology and evolution

Oatley TB (1969) The functions of vocal imitation by African cossyphas. Ostrich 40:85–89.
https://doi.org/10.1080/00306525.1969.9639112

OpenTreeOfLife, Redelings B, Sanchez Reyes LL, et al (2019) Open tree of life synthetic tree

Ovaskainen O, Moliterno De Camargo U, Somervuo P (2018) Animal Sound Identifier (ASI):
software for automated identification of vocal animals. Ecology Letters 21:1244–1254.
https://doi.org/10.1111/ele.13092

R Core Team (2019) R: a language and environment for statistical computing.

Ratnayake CP, Goodale E, Kotagama SW (2010) Two sympatric species of passerine birds
imitate the same raptor calls in alarm contexts. Naturwissenschaften 97:103–108.
https://doi.org/10.1007/s00114-009-0617-7

Ridgway S, Carder D, Jeffries M, Todd M (2012) Spontaneous human speech mimicry by a
cetacean. Curr Biol 22:R860–R861. https://doi.org/10.1016/j.cub.2012.08.044

Roberts DL, Taylor L, Joppa LN (2016) Threatened or data deficient: assessing the
conservation status of poorly known species. Diversity and Distributions 22:558–565.
https://doi.org/10.1111/ddi.12418

Sigmundson R, Girard-Buttoz C, Le Floch A, et al (2025) Vocal sequence diversity and length
remain stable across ontogeny in a catarrhine monkey (Cercocebus atys). Commun Biol
8:465. https://doi.org/10.1038/s42003-025-07922-2

Stoeger AS, Mietchen D, Oh S, et al (2012) An asian elephant imitates human speech. Curr Biol
22:2144–2148. https://doi.org/10.1016/j.cub.2012.09.022

Strickland D, Ouellet HR (2020) Canada Jay (Perisoreus canadensis). In: Billerman SM,
Keeney BK, Rodewald PG, Schulenberg TS (eds) Birds of the World. Cornell Lab of
Ornithology

Suthers RA, Zollinger SA (2004) Producing song: The vocal apparatus. Ann N Y Acad Sci
1016:109–129. https://doi.org/10.1196/annals.1298.041

Taylor AH (2014) Corvid cognition. WIREs Cogn Sci 5:361–372.
https://doi.org/10.1002/wcs.1286

Ter Haar SM, Fernandez AA, Gratier M, et al (2021) Cross-species parallels in babbling:
animals and algorithms. Phil Trans R Soc B 376:20200239.
https://doi.org/10.1098/rstb.2020.0239

Tippin TC (2017) Propensity of predator mimicry in wild Steller's jays. The Faculty of
Humboldt State University

Wascher CAF (2018) Corvids. In: Vonk J, Shackelford T (eds) Encyclopedia of Animal
Cognition and Behavior. Springer International Publishing, Cham, pp 1–12

Wascher CAF, Reynolds S (2025) Vocal communication in corvids: a systematic review.
Animal Behaviour 221:123073. https://doi.org/10.1016/j.anbehav.2024.123073

Wascher CAF, Waterhouse G, Beheim BA (2025) Vocal mimicry in corvids. BioRxiv.
https://doi.org/10.1101/2025.03.26.645457

West MJ, Stroud N, King AP (1983) Mimicry of the human voice by European starlings: The
role of social interaction. Wilson Bull 95:635–640

Woolfenden GE, Fitzpatrick JW (2020) Florida Scrub-Jay (Aphelocoma coerulescens). In:
Birds of the World. Cornell Lab of Ornithology, Ithaca, NY, USA

597