Impact of thermal stress during pupal development in a key pollinator

Sandra Laußer[1] and Christoph Kurze[1]*

[1] Institute for Zoology and Evolutionary Biology, University of Regensburg, Germany

* Corresponding author, email: Christoph.Kurze@ur.de

Abstract

The increasing frequency and intensity of extreme thermal events like heatwaves, driven

by climate change, threaten global biodiversity and entire ecosystems. Despite alarming

projections of population declines for key pollinators like bumblebees, little is known about

the direct impact of varying elevated temperatures during their pupal development. To

gain a better understanding of the dose-response to thermal stress, we experimentally

exposed *Bombus terrestris* L. (Hymenoptera: Apidae) pupae *in vitro* to varying thermal

stress lengths (3-7 days) and intensities (36-38°C). This mechanistic approach allowed us

to uniquely isolate the direct physiological effects of temperature on pupal development

from complex colony dynamics and assess individual emergence success, developmental

duration, body size, relative lipid content, and wing geometrics morphometrics. At the

critical threshold of 38°C, pupal emergence was drastically reduced (2-5 fold) and wing

deformities increased (4-7 fold). Even excluding wing deformities, we found thermal

stress-induced effects on emerged worker wing shape and directional asymmetry. Hence,

our findings suggest that increased extreme thermal stress negatively impacts bumblebee

pupal development. Given the relative heat tolerance of *B. terrestris*, we expect even

greater vulnerability in more cold-adapted bumblebee species.

Keywords social insects, wild bees, extreme heat, acute stress, metamorphosis,

malformations

1. Introduction

Global warming is one of the most pressing challenges of our time [1, 2]. Particularly the increase in frequencies and intensities of heatwaves, defined as periods of abnormally hot temperatures sustained over at least three consecutive days, is concerning [3-6]. In fact, extreme thermal stress can severely impact a wide range of terrestrial animals such as insects, birds and mammals [4, 7, 8]. Despite recent advances in understanding the impact of thermal stress, our knowledge of the dose-response of extreme thermal stress (i.e. exposure variable temperatures and durations) on insect development and their physiological responses is still limited [7, 8].

Gaining a better understanding of the dose-response to thermal stress on insect physiology and fitness might be particularly important in threatened keystone species such as bumblebees (*Bombus* sp., Hymenoptera: Apidae). Bumblebees provide crucial pollination services in ecosystems and agriculture in the Northern hemisphere [9, 10]. As a cold-adapted taxon, bumblebees appear to be particularly vulnerable to climate change, which is reflected in their alarming population declines in some species [11-18]. However, heat tolerance varies significantly across species, with alpine and polar species being especially vulnerable to thermal stress [15, 19]. While ambient temperature fluctuations are buffered in ground nesting bumblebee species [20], those nesting above ground (e.g. *B. hypnorum* and *B. pascuorum* [15, 19]) and commercially managed species like *Bombus terrestris* L., widely used as crucial crop pollinators in greenhouses [21], face an increased risk of direct exposure to extreme thermal stress. Previous studies have shown that exposure to elevated temperatures impair cognition [22] and scent perception [23] in worker bumblebees. Under extreme thermal stress, the survival is significantly affected [24, 25]. Additionally, increasing temperatures negatively correlate with colony growth, which in turn negatively affects gyne production [26].

Colony fitness may not only be influenced by the quantity of offsprings [26], but also by their quality, expressed in their phenotypic traits. Wing morphology (shape, size

55 and asymmetry) and storage reserves (e.g. fat body size) have been suggested as

56 potential markers for monitoring bee health and pollination service, as they correlate with

57 reproductive success, immunity, resilience and foraging efficiency [27]. While detailed

58 dose-response data to environmental stressors at different life stages of bumblebees are

59 still limited [28, 29], evidence suggest that offspring from colonies under constant thermal

60 stress exhibit alterations in wing morphology, reduced body size and smaller antennae

61 [30-33]. As a natural colony response to elevated nest temperatures worker spend

62 significantly more time fanning to cool the brood than under optimal thermal conditions

63 [34, 35]. As complex behaviroual responses can vary significantly between colonies,

64 making it challenging to conduct detailed dose-response experiments under controlled

65 conditions, *in vitro* experiments offer an alternative approach. These controlled

66 experiments are particularly helpful to gain a better mechanistic understanding of the

67 effects of environmental stressors at the precise developmental stages by uniquely

68 isolating the direct physiological responses of individuals from the colony context at the

69 cost of losing natural realism [28, 29].

70  While different developmental stages are likely differentially affected by thermal

71 stress, it remains unclear which stages are more vulnerable and if short periods of thermal

72 stress, like heatwaves, are sufficient to impair their development. To study the direct

73 impact of thermal stress length and sevirity on pupal development in bumblebees, we

74 used an *in vitro* rearing approach [28, 29]. This allowed us to precisely control the

75 temperature by individual pupae, thus isolating their direct physiological and

76 developmental responses from complex colony-level influences. We hypothesised that an

77 increase in rearing temperatures and prolonged exposure would adversely affect the

78 emergence rate, developmental duration, adult body size, fat body size, and wing

79 development.

2. Methods

(a) Experimental overview

To study the impact of thermal stress length and severity on pupal development in bumblebees, we collected 887 fourth instar larvae from 25 *Bombus terrestris* colonies to rear them *in vitro* following previous protocols [28, 29]. While our *in vitro* rearing temperature of 34°C [28, 29] exceeds typical nest temperatures [20], it aligns with brood temperatures, which are typically 2°C warmer than ambient nest temperatures [36, 37]. At the beginning of the pupal stage, pupae were pseudo-randomly assigned to one of the seven treatment groups (detailed in Table 1). We chose temperatures of 36°C, 37°C, and 38°C and durations of 3, 5, and 7 days to reflect potentially relevant extreme heatwave conditions observed in Europe [38-40]. Decision on thermal stress durations were based on our pilot study (unpublished data), which tested only 36°C for 3 days alongside controls and confirmed that an average pupation period would last about 8 days [29]. While we decided to expose pupae to 36°C for 5 and 7 days, we opted for shorter 3- and 5-day exposure durations for the treatments for 37°C and 38°C to avoid excessive pupal mortality at these more extreme temperatures and to focus on sub-lethal effects.

Table 1. Overview of the experimental design. This table details the rearing conditions for all experimental groups. Controls (Ctrl/34C) were reared in vitro at a constant temperature of 34°C until emergence as adults. Pupae of the thermal stress treatments experienced elevated temperatures (36°C, 37°C, and 38°C) for 3, 5 or 7 days at the start of pupation, followed by rearing at 34°C until emergence. Sample sizes (n) reflect the total number of pupae per treatment groups at the start of the experiment, but actual sample size for subsequent measurements were lower due to incomplete emergence.

Treatment groups	Temperature	Duration	Sample size (n)
34C/Ctrl	34°C	until emergence	127
36C5d	36°C	5 d	79
36C7d	36°C	7 d	80
37C3d	37°C	3 d	75
37C5d	37°C	5 d	73
38C3d	38°C	3 d	64
38C5d	38°C	5 d	106

We aimed to distribute pupae from the same collection date and source colony as

broadly as possible among treatment groups (see Supplementary Table S1 for detailed

information about collection cohorts, colony ID, and treatment assignment). Although a

direct comparison between all treatment groups for each cohort was not feasible as we

had access to only three incubators for four different rearing temperatures, our

experimental design relied on a robust concurrent control group. These controls represent

all collection cohorts and colonies as we continuously kept in one incubator at 34°C

throughout the entire experiment. The other two incubators were rotated among 36°C,

37°C, and 38°C across experimental replications. This meant that not all treatment

combinations/temperatures were run concurrently and did not derive from the same

collection cohort (see Supplementary Table S1). To account for variability among colonies

and collection cohorts, we included 'colony ID' and 'collection date' as a random effect into

the statistical models whenever possible.

We checked their survival until adult bees emerged and measured their body

mass. If adult bees emerged, we sampled them at age of 2 days and stored them at -20°C

until subsequent morphometric measurements of their head width, intertegular distance

(ITD), dry mass, and wings.

(b) Colony husbandry

Larvae were collected from 25 queen-right colonies of *Bombus terrestris* (Natupol

Research Hives, Koppert B.V., Netherlands) at various colony phases across multiple

days between 19th of June and 31st of October 2023 to ensure a continuous supply and

broad representation of colony cohorts (see Supplementary Table S1 for detailed

collection dates, number of larvae collected per day, and their colony origin). Colonies

were maintained under similar conditions to previous studies [28, 41]. In short, colonies

were housed in the standard Natupol nest boxes (23 (l) x 25 (w) x 15 (h) cm), which were

connected to foraging arenas (59 (l) x 39 (w) x 26 (h) cm) under a 14:10 h light:dark cycle.

Bumblebees had *ad libitum* access to 70% (w/v) sucrose solution. Depending on colony

size, each colony received 3-11 g of pollen candy daily, which consisted of 67% organic

pollen (naturwaren-niederrhein GmbH, Germany), 25% sucrose (Südzucker AG,

Germany), and 8% tap water. The room temperature was maintained at 25 °C ± 1 °C and

30-50% relative humidity.

(c) Larvae collection and *in vitro* rearing

To collect 4th instar larvae (L4), all adult bees from a colony were temporarily transferred

to a separate cage. Then, L4 larvae were carefully removed from individual, spherical

brood cells using soft tweezers. L4 larvae can be easily identified by the small opening for

food provisioning (figure S1*a*). To minimize disturbance, colonies were only resampled

after minimum period of at least two weeks.

Depending on the size of L4 larvae, they were carefully transferred into either 0.6

mL or 0.96 mL large 3D-printed polylactide (PLA) artificial brood cells (diameter of either 8

mm or 10 mm respectively, figure S1*b*). Those were then placed in 24- or 48-well clear flat

bottom plates (Falcon/Corning, USA) (figure S1*c*) and kept inside a ventilated plastic

container (18.5 x 18.5 x 11.5 cm) at 34°C (KB115, BINDER GmbH, Germany). A 120 mL

cup of saturated sodium chloride solution was included within each container to maintain a

relative humidity (RH) of 65 ± 10 %. The use of this setup facilitated the measurement of

larval and pupal weight gain without the need for repeated handling them directly and

allowed for the easy removal of any dead individuals.

L4 larvae were fed a pollen medium twice daily (morning and evening) until

pupation and the start of the experiment. The pollen medium consisted of 50% w/v

sucrose solution (Südzucker AG, Germany), 40% honeybee-collected organic pollen (Bio-

Blütenpollen, naturwaren-niederrhein GmbH, Germany), 10% Bacto yeast extract

(Bacto™, BD, USA), and 1% casein sodium salt from bovine milk (Sigma-Aldrich,

Germany) [29]. The pollen medium was aliquoted and stored at -20°C until use. Before

hand-feeding all L4 larvae, aliquots were warmed to 34°C and vortexed. For each 20-

minute feeding session, well-plates containing the L4 larvae were placed on a heated

plate at 35°C (Medax model 12801, Medax Nagel GmbH, Germany). Each larva initially

received about 6 µL droplet (7.1 ± 1.6 mg, 10 µL micropipette tips had to be cut off at the

1 µL mark) of pollen medium applied to its ventral abdomen (figure S1*d*). Larvae that

consumed all initial food received a second droplet. Satiation was determined when a

larva curled up and stopped movement, while active larvae were considered hungry

(Supplementary video 1). At the end of each feeding session, any remaining food was

carefully removed to prevent tracheal blockage. Larvae were no longer fed when entering

the pupal stage.

169   At the start of pupal stage, individuals were weighed (d = 0.1 mg, Sartorius

AC120S, Sartorius AG, Germany) and pseudo-randomly assigned to one of seven

treatment groups (detailed in Table 1), ensuring a balanced distribution across treatments

and natal colonies (Table S1). Pupae were transferred to new humidity-controlled

containers (65 ± 10 % RH), each containing a cup of saturated sodium chloride solution,

inside an incubator set to the designated temperature. After the treatment period, pupae

were returned to the control incubator at 34°C. We recorded the proportion of larvae that

successfully reached adulthood.

(d) Emerged adult bees

For each bee reaching adulthood, the emergence date was recorded to calculate the

developmental time required to complete metamorphosis (i.e. pupal phase). Their body

weight was directly measured using a precision scale (d = 0.1 mg, Sartorius AC120S,

Sartorius AG, Germany). Adult bees were then housed individually in 25 mL snap-cap vial

containing a cardboard bottom (approximately 5.5 x 1.5 cm) and mesh lid. Bees had *ad*

libitum access to 60% (w/v) sugar solution via simple 1.5 mL feeder and kept at 25 °C ± 1

°C and 30-50% RH. Two-day-old adults, with fully developed and hardened wings, were

frozen and stored at -20°C until further analysis.

(e) Body size and lipid measurements

As a proxy for adult body size, the intertegular distance (ITD) and head width were

measured using a digital microscope (VHX-500F, Keyence GmbH, Germany) [41]. In

addition, their sex was determined by counting flagellomeres (workers have 10, males 11)

[42]. We determined adult dry mass and lipid content following previous protocols [28, 41].

First, each specimen was carefully dissected by one cut at ventral abdominal segments

from the stinger to the fourth sternite, then dried at 60°C for three days (U40, Memmert

GmbH & Co. KG, Germany), and weighed (d = 0.1 mg, analytic balance M-Pact AX224,

Sartorius GmbH, Germany). Subsequently, the body lipids were extracted using

petroleum ether for five days. The ether was discarded, and the specimen were rinsed

with fresh ether before being dried for an additional three days. The lipid content was

calculated as the difference between the initial and post-extraction dry masses. The

relative lipid content, serving as a proxy for the fat body size, was calculated as the

percentage of lipid contents relative to the initial dry masses.

(e) Wings

Wings were first detached at the tegulae and embedded side-by-side on a microscope

slide (Roti®Mount, Carl Roth GmbH + Co. KG, Germany). Then, wing pairs were

photographed at 20x magnification (Keyence VHX-500F). Two-dimensional coordinates of

twenty landmarks [43] for each wing (figure S1*e*) were obtained using TpsDig2 (v2.32)

[44]. While individuals with visibly undeveloped or fully deformed wings (figure S1*f*) were

excluded from geometric morphometric analysis, we compared the proportion of

individuals with developed, undeformed wings among treatment groups.

(g) Statistical analyses

All statistical analyses and data visualizations were performed using R Statistical Software

in R version 4.4.2 [45]. The complete code as RMD (R Markdown) output and datasets

are provided in Dryad repository: https://doi.org/10.5061/dryad.n02v6wx7z.

We followed a hypothesis driven statistical approach. Whenever possible, we used

generalized linear mixed effect models (GLMMs) using the *glmmTMB* package [46] to

analyse effects of elevated temperatures and exposure duration, and their interactions, as

fixed factors on emergence rates (i.e. pupae reaching adulthood), developmental time (i.e.

days to emergence), adult mass, head width, ITD, relative lipid content, and wing

development (i.e. either fully developed or deformed) with binomial (logit link), beta

regression (logit link) or gamma (log link) data structures (see Supplementary RMD output

for detailed model structure and selection procedure). We included 'colony ID' and

'collection cohort' (see Table S1) as random effects to account for colony and temporal-

specific variability. Due to limited sample sizes for male specific data and unbalanced

sample size distribution, we excluded male data and analysed the effects of thermal

stress on adult mass, dry mass, ITD, and head width for worker data separately (the entire

data are presented in figure S3). This approach was chosen to avoid sex-specific biases

that could potentially mask treatment effects. In fact, we found sex-specific effects on

adult mass (Kruskal-Wallis: χ^2 = 99.723, df = 1, p < 0.0001), adult head width (χ^2 =

88.556, df = 1, p < 0.0001), ITD (χ^2 = 59.588, df = 1, p < 0.0001), and adult relative lipid

content (χ^2 = 6.748, df = 1, p < 0.01) (see Supplementary RMD output for a detailed

analysis of the entire data).

234   For each response variable, a candidate set of GLMMs including potential

explanatory variables was constructed to identify the most parsimonious and best-fitting

model, based on the Akaike information criterion (AIC), with lower AIC values indicating a

better fit (see Table S2 of a simplified overview selection procedure for models meeting

model assumptions). The best models were compared with their respective null-models,

including only random effects, using likelihood ratio tests (LRTs) to determine statistical

significance of added terms. Model assumptions and dispersion of the data were checked

using the *DHARMa* package [47]. Significance levels (p < 0.05) were determined using

the *Anova* function when model assumptions were met. Due to our experimental design

limitations described above, we specifically tested the effects of each discrete treatment

combination compared to the controls using the function *emmeans* [48], adjusted with

Holm-Bonferroni correction for multiple comparisons. When model assumptions of

GLMMs were not met, treatment effects were tested with a non-parametric Kruskal-Wallis

rank sum test, followed by post-hoc comparisons between each treated group with the

controls using Dunn's test with Holm-Bonferroni correction.

Geometric morphometric analyses of wing size, shape, and asymmetry on fully

developed left and right forewings (i.e. excluding all deformed wings) of workers on the

obtained TPS data using geomorph package [49], following a similar approach as

previous studies [31, 43]. Due to sample size limitation of males, we also excluded male

data from the geometric morphometric analyses of wings.

First, right wings were first mirrored to ensure landmark correspondence. Missing

landmark data, representing morphological variation, were accounted for using the

function *estimate.missing*. To remove non-shape variation, a Generalized Procrustes

Analysis (GPA) was performed using *gpagen*. A Principal Component Analysis (PCA) was

conducted on the Procrustes coordinates using *gm.prcomp*, PCA scores (PC1 and PC2)

were calculated, and data distribution was visually assessed. Wings with PC1 z-scores

exceeding $\pm$ 2 standard deviations were removed as outliers, where PC1 explained 78%

of the data's variation. Treatment groups with fewer than six individuals were also

excluded from further analysis, which was only the case for workers from the 5-day-

exposure at 38°C (38C5d).

Then, centroid size and mean centroid size were calculated for each wing and

each bee, respectively, as a measure of wing size. To analyse shape variation, a

permutational multivariate analysis of variance (PERMANOVA) was performed using the

adonis2 function. Homoscedasticity was evaluated using *betadisper*, and pairwise

comparisons were conducted using *pairwise.adonis* with Holm-Bonferroni correction.

Directional asymmetry (DA) was calculated as the difference between left and right wing

Procrustes coordinates, and DA magnitude was calculated as the mean Euclidean

distance between corresponding landmarks. To assess fluctuating asymmetry (FA), a

bilateral symmetry analysis was performed using *bilat.symmetry* and calculating the FA

magnitude as the Euclidean norm of the FA component. The effects of treatment on

centroid size, DA and FA magnitude were tested using linear regression models (LM)

when assumptions were met, followed by contrast pairwise comparisons for significant

results as described above. Kruskal-Wallis tests were used as a non-parametric

alternative, followed by Dunn's post-hoc test to compare treatment groups to the control

by applying Holm-Bonferroni corrections.

Results

We found a strong interaction effect between thermal stress temperature and duration on

the proportion of pupae reaching adulthood (GLMM: χ^2 = 25.813, df = 1, p < 0.0001; figure

1a). This effect was mainly driven by increasing temperatures (χ^2 = 24.549, df = 1, p <

0.0001). Pairwise comparisons revealed that pupae exposed to 38°C for 3 days were 43%

(38C3d: z = 3.655, p < 0.01) less likely to reach adulthood compared to the controls.

When exposed to 38°C for 5 days the emergence rate dropped by 83 % (38C5d: z =

7.236, p < 0.0001) compared to the controls.

Thermal stress intensity also significantly affected pupal developmental duration

(Kruskal-Wallis: χ^2 = 69.584, df = 6, p < 0.0001; figure 1b; worker and male data are

presented separately in figure S2). While an exposure to 36°C shortened development

compared to controls (Dunn's test: 36C5d, z = 3.253, p < 0.01; 36C7d, z = 3.142, p <

0.01), an exposure to 38°C prolonged pupal development (38C3d, z = -4.393, p < 0.0001;

38C3d, z = -2.739, p < 0.01). There was no significant effect found for the 37°C exposure

compared to controls (37C3d, z = -1.361, p > 0.5; 37C5d, z = -1.164, p > 0.5).

Figure 1. Direct effects of varying intensity of thermal stress on *(a)* emergence rate and *(b)*

developmental duration in *B. terrestris* pupae. Controls (Ctrl, reared at constant 34°C, blue) were

compared to individuals exposed to increased temperatures: of 36°C for 5 days (light warm yellow) and 7 days

(warm yellow), 37°C for 3 days (light reddish-orange) and 5 days (reddish orange), and 38°C for 3 days (light

red) and 5 days (deep red). Sample sizes are indicated by white numbers and significant differences are

marked by asterisks (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

We found no significant effects of thermal stress treatments on worker body size

proxies, including adult mass (Kruskal-Wallis: $x^2 = 9.125$, df $= 6$, $p > 0.05$; figure 2*a*) and

ITD (Kruskal-Wallis: $x^2 = 9.571$, df $= 6$, $p > 0.05$; figure 2*c*). While there was a weak

significant effect on worker head width (Kruskal-Wallis: $x^2 = 16.131$, df $= 6$, $p < 0.05$, figure

2*b*), we found no significant differences between treatment groups and controls (Dunn's

tests: $p > 0.05$). However, female head width varied with temperature (Kruskal-Wallis: $x^2 =$

12.311, df $= 3$, $p < 0.01$), where workers exposed to 38°C as pupae had slightly wider

heads than the controls ($z = -2.227$, $p < 0.05$).

In contrast, we found that thermal stress temperature significantly influenced

relative lipid content in adult workers (Kruskal-Wallis: χ^2 = 25.847, df = 6, p < 0.001, figure

2*d*). Pairwise comparisons indicated significant differences between 37C5d (Dunn's test: z

= -2.592, p < 0.05), 38C3d (z = -2.844, p < 0.05), and 38C5d (z = -3.012, p < 0.01) and

the controls (figure 2*d*).

Figure 2. Effects thermal stress exposure during pupal development on adult worker morphometrics

and lipid content in *B. terrestris*: *(a)* adult mass, *(b)* head width, *(c)* intertegular distance (ITD), *(d)* and

relative lipid content (2-day-old adults). Controls (34°C, blue) were compared to individuals exposed to

increased temperatures: of 36°C for 5 days (light warm yellow) and 7 days (warm yellow), 37°C for 3 days

(light reddish-orange) and 5 days (reddish orange), and 38°C for 3 days (light red) and 5 days (deep red).

Sample sizes (*n*) are provided in the legend. Asterisks denote significant differences (* p < 0.05, ** p < 0.01).

We found that thermal stress temperature impacted wing development (GLMM:

x^2 = 9.410, df = 1, p < 0.01, figure 3*a*), but there was no significant effect of exposure

duration (x^2 = 0.280, df = 1, p > 0.05) and their interactions (x^2 = 3.491, df = 1, p = 0.062).

Specifically, 38C3d pupae were four times more likely to develop deformed wings

compared to the controls (z = -3.367, p < 0.01), while 38C5d pupae were nearly seven

times more likely of developing deformed wings (z = -3.920, p < 0.001). Pupae of the

36C7d treatment showed a marginally significant, twofold increase in deformed wings (z =

-2.482, p = 0.052). Pairwise comparisons between the other treatment groups and

controls were not significant (36C5d: z = 0.656, p > 0.05; 37C3d: z = -2.001, p > 0.05; and

37C5d: z = -2.279, p > 0.05).

Our geometric morphometrics wing analysis, excluding deformed wings, indicated

a significant effect on worker wing shape (PERMANOVA: F = 11.677, R^2 = 0.197, df = 5,

p < 0.001, figure 3*b*). Pairwise comparisons revealed significant differences between all

treatment groups and the controls (36C5d, F = 7.259, p < 0.05; 36C7d, F = 5.005, p <

0.05; 37C3d, F = 21.125, p < 0.05; 37C5d, F = 29.494, p < 0.05; 38C3d, F = 25.168, p <

0.05). Similarly, we found that exposure affected centroid wing size in workers (LM: F =

4.205, df = 5, p < 0.01, figure S3*c*). Pairwise comparisons indicated a marginally

significant increased worker wing size of the 37C5d (t = -2.568, p = 0.058) and 38C3d (t =

-2.394, p = 0.073) treatment groups compared to the controls.

Furthermore, our thermal stress treatment significantly affected the DA magnitude

in worker wings (LM: F = 9.466, df = 5, p < 0.0001, figure 3*d*), but not FA magnitude

(Kruskal-Wallis: x^2 = 3.903, df = 5, p > 0.05, figure 3*e*). Workers exposed to 36°C showed

higher DA (36C5d, t = -2.520, p < 0.05; 36C7d, t = -2.536, p < 0.05), while workers

exposed to 37°C (37C3d, t = 2.655, p < 0.05; 37C5d, t = 2.932, p < 0.05) and 38°C

(38C3d, t = 2.500, p < 0.05) showed lower DA compared to controls.

351
352

Figure 3. Effects of thermal stress on wing development in *B. terrestris.* *(a)* Proportion of deformed wings

in emerged adult bees (including both males and workers) for each treatment group (sample sizes are

indicated by white numbers). Deformed and male wings presented in *(a)* were excluded for detailed wing

geometric morphometric analyses *(b-e)*, therefore resulting in lower sample sizes. *(b)* Principal component

analysis (PCA) of wing morphology in emerged workers with fully developed wings across thermal stress

treatments (see legend, n = samples size). *(c)* Influence of thermal stress treatments on wing size (centroid

size) and the magnitude of *(d)* directional and *(e)* fluctuating asymmetry in fully developed worker wings.

Controls (34°C, blue) were compared to individuals exposed to increased temperatures: of 36°C for 5 days

(light warm yellow) and 7 days (warm yellow), 37°C for 3 days (light reddish-orange) and 5 days (reddish

orange), and 38°C for 3 days (light red). Asterisks denote significant differences (* $p < 0.05$, ** $p < 0.01$, *** $p <$

0.001).

3. Discussion

Our study provides compelling evidence that thermal stress temperature and duration

significantly impact pupal development in *B. terrestris* with notable effects on emergence

rate (figure 1*a*), developmental duration (figure 1*b*), lipid stores (figure 2*d*), and wing

development (figure 3). These findings highlight the extreme vulnerability of the pupal

stage to thermal stress, particularly at 38°C, which appears to be a critical threshold for

development disruption.

Although our treatments are extreme, they reflect potential heatwave scenarios in

Europe with temperatures exceeding 38°C for several days as reported in parts of the

British Isles, Mediterranean and Eastern Europe [38-40]. We are aware that our *in vitro*

experiment oversimplifies the thermal stress that bumblebee pupae would experience

naturally, especially in ground-nesting species such as *B. terrestris*, where ambient

temperatures are buffered [20]. However, bumblebee species that typically nest above

ground (e.g. *B. hypnorum* and *B. pascuorum*) or commercially managed colonies (e.g. *B.*

terrestris), widely used for pollination services in greenhouses [21], are potentially more

directly exposed to extreme thermal stress over prolonged periods. Furthermore, we

expect more cold-adapted bumblebee species such as *B. lapidarius*, *B. alpinus*, and *B.*

poralris to be even more vulnerable to thermal stress than *B. terrestris*, given its relative

high heat tolerance [15, 19]. While natural colony responses to thermal stress, such as

fanning, aim to cool nest temperatures, these thermoregulation efforts may negatively

affect colony fitness and could be overwhelmed or fail under prolonged periods of stress

[34, 35]. Here, we used *B. terrestris* as a model species to gain a mechanistic inside into

the dose-response to thermal stress on pupal development by isolating them from colony.

Most notably, our findings show that emergence rates decreased by almost twofold

and fivefold in pupae exposed to 38°C for 3 and 5 days, respectively, compared to

controls (figure 1*a*). Despite *in vitro* rearing of L4 larvae and pupae deviates from natural

nest conditions, the high emergence rate of controls suggests that our experimental setup

allowed for normal development. While 38°C exposure slightly prolonged pupal

development, a 36°C exposure slightly accelerated it (figure 1*b*), suggesting a potential

developmental benefit in pupae to temperature elevation up to 37°C, followed by severe

disruption at 38°C.

We observed no clear effects on worker body size proxies (figures 2*a-c*), except

for an increased worker head width at 38°C. These results contrast with previous studies

showing that colonies exposed to prolonged thermal stress would produce smaller

offspring, indicated by smaller ITD [30, 32] and body mass [33], potentially reflecting

resource allocation strategies such as reduced larval feeding. Alternatively, thermally

stressed L4 larvae may have a decreased body weight gain due to reduced food

consumption [28]. Despite potential limitations in statistical power due to the relative low

emergence rates and resulting small sample sizes (figure 1*a*), particularly for the 38°C

treatments, we speculate that larger pupae have a selective advantage under thermal

stress (figure 1*b,* figure S3*a*). Although small insect body sizes should generally be

favoured under warmer environmental conditions over the long term according to

Bergmann's rule [8], this rule does not apply to bumblebees [31].

In addition, we found that individuals exposed to thermal stress during pupal

development (37°C for 5 days, 38°C for 3 and 5 days) had increased levels of the relative

lipid content (fat body) as adult workers (figure 2*d*). This may indicate a potentially higher

thermal stress resilience due to increased energy reserves, immunocompetence, and heat

shock protein (HSP) synthesis [27, 50-53]. In fact, the insect fat body is the primary site

for HSP synthesis in response to thermal challenges by activating stress-response

pathways that upregulate the production of these chaperones to mitigate protein damage

[50, 53]. Interestingly, the relative lipid content was not affected in workers that were

exposed to thermal stress as L4 larvae, further underscoring the differential vulnerability of

developmental stages [28].

Wing development was severely impaired by thermal stress temperature, notably

with a fourfold and sevenfold increase in wing deformities in emerged adult bees from

pupae exposed to 38°C for 3 and 5 days, respectively (figure 3*a*). Even excluding

individuals with these deformities, workers wing shape altered for all treatments (figure

3*b*), which appears to be a natural response in colonies under thermal and parasitic stress

[31, 54]. However, our data shows only marginal effects on worker wing size (figure 3*c*)

with a similar pattern as for the other worker body size proxies (figure 2*a-c*). DA was

affected by temperature in workers, with increasing DA at 36°C and decreasing DA at

37°C and 38°C (figure 3*d*), potentially also reflecting the different selection pressures

among treatments. While FA remained unaffected in workers (figure 3*e*), confirming

previous studies in honeybees [55] and bumblebees [31].

In conclusion, our *in vitro* study provides novel insights into the direct impact of

varying thermal stress scenarios on *B. terrestris* pupal development. While our

experimental design sacrifices ecological realism, it enabled precise control over timing

and temperature. Our findings highlight a critical thermal stress threshold at 38°C, where

severe disruptions in bumblebee pupal development manifest by particularly impairing

emergence and wing development. Given the heat tolerance of *B. terrestris*, we expect

more cold-adapted and above ground nesting bumblebee species are potentially more

vulnerable to increasingly more frequent and severe heatwaves as climate change

continues.

Acknowledgments

Author contributions

[Anonymous]

Data accessibility

Datasets and R script used for statistical analysis and producing all figures have archived

in Dryad repository:

http://datadryad.org/share/vpCETsQuLDiGFIHSwioKRfRjwKhWotnYFRwZ72ovPC8

Ethics statement

This study was conducted in accordance with the ethical regulations of the German

Animal Welfare Act (TierSchG) for conducting experiments with insects.

Funding

This study was carried out without any third-party funding.

Competing interests

The authors have no competing interests.

455
References

1.   IPCC. Summary for Policymakers. Climate Change 2022 – Impacts, Adaptation
and Vulnerability: Working Group II Contribution to the Sixth Assessment Report of the
Intergovernmental Panel on Climate Change. Cambridge: Cambridge University Press;
2023. p. 3-34.

2.   Garcia RA, Cabeza M, Rahbek C, Araújo MB. Multiple dimensions of climate
change and their implications for biodiversity. Science. 2014;344(6183):1247579.

3.   Lhotka O, Kyselý J, Farda A. Climate change scenarios of heat waves in Central
Europe and their uncertainties. Theoretical and applied climatology. 2018;131(3):1043-54.

4.   Stillman JH. Heat waves, the new normal: summertime temperature extremes will
impact animals, ecosystems, and human communities. Physiology. 2019;34(2):86-100.

5.   Perkins-Kirkpatrick SE, Lewis SC. Increasing trends in regional heatwaves. Nature
communications. 2020;11(1):3357.

6.   Meehl GA, Tebaldi C. More intense, more frequent, and longer lasting heat waves
in the 21st century. Science. 2004;305(5686):994-7.

7.   Ma C-S, Ma G, Pincebourde S. Survive a warming climate: insect responses to
extreme high temperatures. Annual Review of Entomology. 2021;66:163-84.

8.   González-Tokman D, Córdoba-Aguilar A, Dáttilo W, Lira-Noriega A,
Sánchez-Guillén RA, Villalobos F. Insect responses to heat: physiological mechanisms,
evolution and ecological implications in a warming world. Biological Reviews.
2020;95(3):802-21.

9.   Klein A-M, Vaissiere BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen
C, et al. Importance of pollinators in changing landscapes for world crops. P Roy Soc B.
2007;274:303-13. doi: 10.1098/rspb.2006.3721. PubMed PMID: WOS:000243354100001.

10.   Goulson D, Lye GC, Darvill B. Decline and conservation of bumble bees. Annual
Review of Entomology. 2008;53(1):191-208.

11.   Ghisbain G, Thiery W, Massonnet F, Erazo D, Rasmont P, Michez D, et al.
Projected decline in European bumblebee populations in the twenty-first century. Nature.
2024;628(8007):337-41.

12.   Soroye P, Newbold T, Kerr J. Climate change contributes to widespread declines
among bumble bees across continents. Science. 2020;367(6478):685-8.

13.   Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. Global
pollinator declines: trends, impacts and drivers. Trends Ecology and Evolution.
2010;25:345-53. doi: 10.1016/j.tree.2010.01.007.

14.   Maebe K, Hart AF, Marshall L, Vandamme P, Vereecken NJ, Michez D, et al.
Bumblebee resilience to climate change, through plastic and adaptive responses. Global
Change Biology. 2021;27(18):4223-37. doi: https://doi.org/10.1111/gcb.15751.

15.   Martinet B, Dellicour S, Ghisbain G, Przybyla K, Zambra E, Lecocq T, et al. Global
effects of extreme temperatures on wild bumblebees. Conservation Biology.
2021;35(5):1507-18.

16.   Cameron SA, Sadd BM. Global trends in bumble bee health. Annual Review of
Entomology. 2020;65:209-32.

17.   Cameron SA, Lozier JD, Strange JP, Koch JB, Cordes N, Solter LF, et al. Patterns
of widespread decline in North American bumble bees. Proceedings of the National
Academy of Sciences. 2011;108(2):662-7.

18.   Kerr JT, Pindar A, Galpern P, Packer L, Potts SG, Roberts SM, et al. Climate
change   impacts   on   bumblebees   converge   across   continents.   Science.
2015;349(6244):177-80.

19.   Zambra E, Martinet B, Brasero N, Michez D, Rasmont P. Hyperthermic stress
resistance of bumblebee males: test case of Belgian species. Apidologie. 2020;51:911-20.

20.   Kevan PG, Rasmont P, Martinet B. Thermodynamics, thermal performance and
climate change: temperature regimes for bumblebee (*Bombus* spp.) colonies as examples
of   superorganisms.   Frontiers   in   Bee   Science.   2024;   2-2024.   doi:
10.3389/frbee.2024.1351616.

21.   Potts SG, Imperatriz-Fonseca V, Ngo HT, Aizen MA, Biesmeijer JC, Breeze TD, et
al. Safeguarding pollinators and their values to human well-being. Nature. 2016;540:220-
9.

22.   Gérard M, Amiri A, Cariou B, Baird E. Short-term exposure to heatwave-like
temperatures affects learning and memory in bumblebees. Global Change Biology.
2022;28(14):4251-9.

23.   Nooten SS, Korten H, Schmitt T, Kárpáti Z. The heat is on: reduced detection of
floral scents after heatwaves in bumblebees. Proceedings of the Royal Society B.
2024;291(2029):20240352.

24.   Quinlan GM, Feuerborn C, Hines HM, Grozinger CM. Beat the heat: thermal
respites and access to food associated with increased bumble bee heat tolerance. Journal
of Experimental Biology. 2023;226(17):jeb245924.

25.   Kuo Y, Lu Y-H, Lin Y-H, Lin Y-C, Wu Y-L. Elevated temperature affects energy
metabolism and behavior of bumblebees. Insect Biochemistry and Molecular Biology.
2023;155:103932.

26.   Theodorou P, Kühn O, Baltz LM, Wild C, Rasti SL, Bucksch CR, et al. Bumble bee
colony health and performance vary widely across the urban ecosystem. Journal of
Animal Ecology. 2022;91(10):2135-48. doi: https://doi.org/10.1111/1365-2656.13797.

27.   Vanderplanck M, Breeze T, Cini E, Potts SG, Senapathi D. Monitoring bee health
in European agroecosystems using wing morphology and fat bodies. One Ecosystem.
2021;6.

28.   Wögler L, Kurze C. Experimental short-term heatwaves negatively impact body
weight gain and survival during larval development in *Bombus terrestris* L. (Hymenoptera:
Apidae). Biology Open. 2025;14(4). doi: 10.1242/bio.061781.

29.   Kato Y, Kikuta S, Barribeau SM, Inoue MN. In vitro larval rearing method of
eusocial bumblebee Bombus terrestris for toxicity test. 2022;12(1):15783.

30.   Guiraud M, Cariou B, Henrion M, Baird E, Gérard M. Higher developmental
temperature increases queen production and decreases worker body size in the
bumblebee *Bombus terrestris*. Journal of Hymenoptera Research. 2021;88:39-49.

31.   Gérard M, Michez D, Debat V, Fullgrabe L, Meeus I, Piot N, et al. Stressful
conditions reveal decrease in size, modification of shape but relatively stable asymmetry
in bumblebee wings. Sci Rep. 2018;8(1):15169.

32.   Gérard M, Guiraud M, Cariou B, Henrion M, Baird E. Elevated developmental
temperatures impact the size and allometry of morphological traits of the bumblebee
Bombus terrestris. Journal of Experimental Biology. 2023;226(8):jeb245728.

33.   Perl CD, Johansen ZB, Moradinour Z, Guiraud M, Restrepo CE, Wen Jie V, et al.
Heatwave-Like Events During Development Are Sufficient to Impair Bumblebee Worker
Responses to Sensory Stimuli. Frontiers in Ecology and Evolution. 2022;9. doi:
10.3389/fevo.2021.776830.

34.   Sepúlveda Y, Nicholls E, Schuett W, Goulson D. Heatwave-like events affect
drone production and brood-care behaviour in bumblebees. PeerJ. 2024;12:e17135.

35.   Bretzlaff T, Kerr JT, Darveau C-A. Handling heatwaves: balancing
thermoregulation, foraging and bumblebee colony success. Conservation Physiology.
2024;12(1). doi: 10.1093/conphys/coae006.

36.   Weidenmüller A, Kleineidam C, Tautz J. Collective control of nest climate
parameters in bumblebee colonies. Animal Behaviour. 2002;63(6):1065-71.

37.   Vogt FD. Thermoregulation in bumblebee colonies. I. Thermoregulatory versus
brood-maintenance behaviors during acute changes in ambient temperature.
Physiological zoology. 1986;59(1):55-9.

38.   Barriopedro D, Fischer EM, Luterbacher J, Trigo RM, García-Herrera R. The hot
summer of 2010: redrawing the temperature record map of Europe. Science.
2011;332(6026):220-4.

39.   Lhotka O, Kyselý J. Three-dimensional analysis reveals diverse heat wave types in
Europe. Communications Earth & Environment. 2024;5(1):323.

40.   Rita A, Camarero JJ, Nolè A, Borghetti M, Brunetti M, Pergola N, et al. The impact
of drought spells on forests depends on site conditions: The case of 2017 summer heat
wave in southern Europe. Global change biology. 2020;26(2):851-63.

41.   Gilgenreiner M, Kurze C. Age dominates flight distance and duration, while body
size shapes flight speed in *Bombus terrestris* L.(Hymenoptera: Apidae). Proceedings of
the Royal Society B. 2024;291(2027):20241001.

42.   Wcislo WT. Sensilla numbers and antennal morphology of parasitic and non-
parasitic bees (Hymenoptera : Apoidea). 1995;24(1):63-81.

43.   Aytekin MA, Terzo M, Rasmont P, Çağatay N. Landmark based geometric
morphometric analysis of wing shape in *Sibiricobombus* Vogt (Hymenoptera: Apidae:
Bombus Latreille). Annales de la Société entomologique de France (NS). 2007;43(1):95-
102.

44.   Rohlf FJ. The tps series of software. Hystrix. 2015;26(1):9-12.

45.   R Core Team. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/;
2024.

46.   Brooks ME, Kristensen K, Van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et
al. glmmTMB balances speed and flexibility among packages for zero-inflated generalized
linear mixed modeling. The R journal. 2017;9(2):378-400.

47.   Hartig F. DHARMa: residual diagnostics for hierarchical (multi-level/mixed)
regression models. R package version 03. 2020;3(5).

48.   Lenth R, Lenth MR. Package 'lsmeans'. The American Statistician.
2018;34(4):216-21.

49.   Adams D, Collyer M, Kaliontzopoulou A, Baken E. Geometric morphometric
analyses of 2D and 3D landmark data, version 4.0. 8. R Foundation for Statistical
Computing. Retrieved from https://cran. r-proje …; 2024.

50.   Arrese EL, Soulages JL. Insect fat body: energy, metabolism, and regulation.
Annual Review of Entomology. 2010;55:207-25.

51.   Korner P, Schmid-Hempel P. In vivo dynamics of an immune response in the
bumble bee *Bombus terrestris.* 2004;87(1):59-66.

52.   Vesterlund S-R, Lilley T, Van Ooik T, Sorvari J. The effect of overwintering
temperature on the body energy reserves and phenoloxidase activity of bumblebee
Bombus lucorum queens. Insectes sociaux. 2014;61:265-72.
53.   King AM, MacRae TH. Insect heat shock proteins during stress and diapause.
Annual Review of Entomology. 2015;60(1):59-75.
54.   Gérard M, Cariou B, Henrion M, Descamps C, Baird E. Exposure to elevated
temperature during development affects bumblebee foraging behavior. Behavioral
Ecology. 2022;33(4):816-24.
55.   Jones JC, Helliwell P, Beekman M, Maleszka R, Oldroyd BP. The effects of rearing
temperature on developmental stability and learning and memory in the honey bee, *Apis*
mellifera. Journal of comparative physiology A. 2005;191:1121-9.

606