Heatwave length and severity drive development disruption in a key pollinator

Sandra Laußer[1] and Christoph Kurze[1]*

[1] Institute for Zoology and Evolutionary Biology, University of Regensburg, Germany

* Corresponding author, email: Christoph.Kurze@ur.de

Abstract

Heatwaves are becoming more frequent and severe due to climate change, posing a

significant threat to global biodiversity and entire ecosystems. Despite alarming

projections of population declines for key pollinators like bumblebees, crucial for crop

pollination and ecosystem services, surprisingly little is known about the implications of

heatwaves on their larval and pupal development. Here, we experimentally exposed

Bombus terrestris L. (Hymenoptera: Apidae) pupae *in vitro* to varying heatwave scenarios

in terms of length (3, 5, or 7 days) and intensity (36°C, 37°C, and 38°C) to study the

impact on their development at this critical life stage. We assessed their emergence rate,

developmental duration, lipid content, sex-specific body size and wing geometrics

morphometrics. Our major findings revealed a critical heat stress threshold at 38°C, where

pupal emergence was drastically reduced by almost twofold and fivefold for 3 and 5 days,

respectively. Adults emerged from these heatwave treatments exhibited a fourfold and

sevenfold increase in wing deformities. Even excluding these extreme cases, we found

clear evidence for heatwave-induced effects on female wing shape and directional

asymmetry (DA) among treatments. Given the relative heat tolerance of our model

species *B. terrestris*, we expect even greater vulnerability in more cold-adapted

bumblebee species. Our findings underscore the profound ecological consequences of

heatwaves on bumblebee pupal development, with potentially devastating cascading

effects on colony fitness, population viability, crop pollination, and ultimately, ecosystem

services essential for human well-being.

Keywords social insect, wild bees, extreme heat, acute stress, metamorphosis,

malformations

1. Introduction

Global warming is undoubtedly one of the most pressing challenges of our time (Garcia, Cabeza, Rahbek, & Araújo, 2014; IPCC, 2023). Particularly the increase in frequencies and intensities of extreme weather events such as heatwaves are concerning (Lhotka, Kyselý, & Farda, 2018; Meehl & Tebaldi, 2004; Perkins-Kirkpatrick & Lewis, 2020; Stillman, 2019). Heatwaves, defined as periods of abnormally hot temperatures of at least three consecutive days, can severely affect the life of wide range of terrestrial animals such as insects, birds and mammals (González-Tokman et al., 2020; Ma, Ma, & Pincebourde, 2021; Stillman, 2019). Despite recent advances in understanding the impact of heatwaves, our knowledge of their impact on animal development, their fitness and entire populations is still limited (González-Tokman et al., 2020; Jentsch, Kreyling, & Beierkuhnlein, 2007; Ma et al., 2021; Stillman, 2019).

Gaining a better understanding of how the length and severity heatwaves impact animal physiology and fitness might be particularly important in threatened key-stone species such as bumblebees (*Bombus* sp., Hymenoptera: Apidae). Bumblebees provide crucial pollination services in ecosystems and agriculture in the Northern hemisphere (Goulson, Lye, & Darvill, 2008; Klein et al., 2007). As cold-adapted taxon, bumblebees appear to be particularly vulnerable to climate change, which is reflected in their alarming population declines in some species (Cameron et al., 2011; Cameron & Sadd, 2020; Ghisbain et al., 2024; Kerr et al., 2015; Maebe et al., 2021; Martinet et al., 2021; Potts et al., 2010; Soroye, Newbold, & Kerr, 2020). However, heat tolerance varies significantly across species, with alpine and polar species being especially vulnerable to heat stress (Martinet et al., 2021a; Zambra et al., 2020). In contrast, commercially used bumblebees like *Bombus terrestris* L. are commonly kept aboveground in greenhouses as crucial crop pollinators (Potts et al., 2016), which puts them at increased risk of exposure to extreme heat. Previous studies have shown that exposure to elevated heatwave-like temperatures impairs cognition (Gérard, Amiri, Cariou, & Baird, 2022) and scent perception (Nooten, Korten, Schmitt, & Kárpáti, 2024) in worker bumblebees. Under extreme heat stress, the

survival is significantly affected (Kuo, Lu, Lin, Lin, & Wu, 2023; Quinlan, Feuerborn, Hines,

& Grozinger, 2023). Additionally, increasing temperatures negatively correlate with colony

growth, which in turn negatively affects gyne production (Theodorou et al., 2022).

Colony fitness may not only be influenced by quantity of offsprings (Theodorou et

al., 2022), but also by their quality, expressed in their phenotypic traits. In particular, wing

morphology (shape, size and asymmetry) and storage reserves (e.g. fat body size) have

been suggested as potential markers for monitoring bee health and pollination service, as

they correlate with reproductive success, immunity, resilience and foraging efficiency

(Vanderplanck, Breeze, Cini, Potts, & Senapathi, 2021). Indeed, experimental exposure of

bumblebee colonies to prolonged constant heat led morphological changes in their

offspring, including alterations in wing morphology, reduced body size and smaller

antennae (Guiraud, Cariou, Henrion, Baird, & Gérard, 2021; Gérard, Cariou, Henrion,

Descamps, & Baird, 2022; Gérard, Guiraud, Cariou, Henrion, & Baird, 2023; Gérard et al.,

2018; Perl et al., 2022). Moreover, elevated nest temperatures during their larval

development seem to have carry-over effects on adult behaviour (Gérard, Cariou, et al.,

2022; Perl et al., 2022). To prevent or limit the negative consequences of elevated nest

temperatures workers spend significantly more time fanning to cool the brood (Bretzlaff,

Kerr, & Darveau, 2024; Sepúlveda, Nicholls, Schuett, & Goulson, 2024). While different

developmental stages are likely differentially affected by heat stress, it remains unclear

which stages are most vulnerable and if short-term heatwaves are sufficient to impair their

development.

To study the impact of heatwave length and sevirity on pupal development in

bumblebees, we used an *in vitro* rearing protocol (Kato, Kikuta, Barribeau, & Inoue, 2022;

Woegler & Kurze, 2024), isolating confounding influences of the worker behaviour in the

colony. We hypothesised that an increase in rearing temperatures and prolonged

exposure would adversely affect the emergence rate, developmental duration, adult body

size, fat body size, and wing development.

2. fMethods

(a) Experimental overview

To study the impact of heatwave length and sevirity on pupal development in bumblebees, we collected 887 fourth instar larvae from 25 commercial *Bombus terrestris* colonies to rear them *in vitro* following previous protocols (Kato et al., 2022; Woegler & Kurze, 2024). While our *in vitro* rearing temperature of 34°C (Kato et al., 2022) exceeds typical nest temperatures, it aligns with brood temperatures, which are typically 2°C warmer than ambient nest temperatures (Vogt, 1986; Weidenmüller, Kleineidam, & Tautz, 2002). With the beginning of the pupal stage, they were pseudo-randomly assigned to one of the seven treatment groups to ensure relative equal distribution among treatments and colonies. Pupae were exposed to either 36°C for 5 day (36C5d, n = 79), 36°C for 7 days (36C7d, n = 80), 37°C for 3 day (37C3d, n = 75), 37°C for 5 days (37C5d, n = 73), 38°C for 3 day (38C3d, n = 64), and 38°C for 5 days (38C5d, n = 106), while the controls (Ctrl/34C, n = 127) were kept at a constant 34°C. We checked their survival until adult bees emerged and measured their body mass. If adult bees emerged, we freeze killed them at age of 2 days and kept them at -20°C until subsequent morphometric measurements of their head width, intertegular distance (ITD), dry mass, and wings.

(b) Colony husbandry

Larvae were collected from 25 queen-right colonies of *Bombus terrestris* (Natupol Research Hives, Koppert B.V., Netherlands) at various colony phases. Colonies were maintained under conditions similar to our previous studies (Gilgenreiner & Kurze, 2024; Woegler & Kurze, 2024). In short, they were housed in the standard Natupol nest boxes (23 (l) x 25 (w) x 15 (h) cm), which were connected to foraging arenas (59 (l) x 39 (w) x 26 (h) cm) under a 14:10 h light:dark cycle. Bumblebees had *ad libitum* access to 70% (w/v) sucrose solution. Depending on colony size, each colony received 3-11 g of pollen candy daily, which consisted of 67% organic pollen (naturwaren-niederrhein GmbH, Germany),

25% sucrose (Südzucker AG, Germany), and 8% tap water. The room temperature was

maintained at 25 °C ± 1 °C and 30-50% relative humidity.

(c) Larvae collection and *in vitro* rearing

To collect 4[th] instar larvae (L4), all adult bees from a colony were temporarily transferred

to a separate cage. Then, L4 larvae were carefully removed from individual, spherical

brood cells using soft tweezers. L4 larvae can be easily identified by the small opening for

food provisioning (figure S1a) (Woegler & Kurze, 2024). To minimize disturbance,

colonies were only resampled after minimum period of at least two weeks.

Depending on the size of L4 larvae, they were carefully transferred into smaller or

larger 3D-printed polylactide (PLA) artificial brood cells (capacity of either 0.6 mL or 0.96

mL, diameter of either 8 mm or 10 mm respectively, figure S1b). Those were then placed

in 24- or 48-well clear flat bottom plates (Falcon/Corning, USA) (figure S1c) and kept

inside a ventilated plastic container (18.5 x 18.5 x 11.5 cm) at 34°C (KB115, BINDER

GmbH, Germany). A 120 mL cup of saturated sodium chloride solution was included

within each container to maintain a relative humidity of 65 ± 10 %. The use of this setup

facilitated the measurement of larval and pupal weight gain without the need for repeated

handling them directly and allowed for the easy removal of any dead individuals.

L4 larvae were fed a pollen medium twice daily (morning and evening) until

pupation and the start of the experiment. The pollen medium consisted of 50% w/v

sucrose solution (Südzucker AG, Germany), 40% honeybee-collected organic pollen (Bio-

Blütenpollen, naturwaren-niederrhein GmbH, Germany), 10% Bacto yeast extract

(Bacto™, BD, USA), and 1% casein sodium salt from bovine milk (Sigma-Aldrich,

Germany) (Kato et al., 2022). The pollen medium was aliquoted and stored at -20°C until

use. Before hand-feeding all L4 larvae, aliquots were warmed to 34°C and vortexed. For

each 20-minute feeding session, well-plates containing the L4 larvae were placed on a

heated plate at 35°C (Medax model 12801, Medax Nagel GmbH, Germany). Each larva

initially received about 6 μL droplet (7.1 ± 1.6 mg, 10 μL micropipette tips had to be cut off

at the 1 μL mark) of pollen medium applied to its ventral abdomen (figure S1d). Larvae

that consumed all initial food received a second droplet. Satiation was determined when a

larva curled up and stopped movement, while active larvae were considered hungry

(Supplementary video 1). At the end of each feeding session, any remaining food was

carefully removed to prevent tracheal blockage. Larvae were no longer fed when entering

the pupal stage.

At the start of pupal stage, individuals were weighed (d = 0.1 mg, Sartorius

AC120S, Sartorius AG, Germany) and pseudo-randomly assigned to one of seven

treatment groups, ensuring a balanced distribution across treatments and natal colonies.

Pupae were exposed to one of the following heatwave scenarios: 36°C for 5 day (36C5d,

n = 79), 36°C for 7 days (36C7d, n = 80), 37°C for 3 day (37C3d, n = 75), 37°C for 5 days

(37C5d, n = 73), 38°C for 3 day (38C3d, n = 64), or 38°C for 5 days (38C5d, n = 106).

Controls (Ctrl/34C, n = 127) were reared at a constant 34°C. Pupae were transferred to

new humidity-controlled containers inside an incubator set to the designated temperature.

After the treatment period, they were returned to the control incubator at 34°C. Although

pupal survival was recorded daily, these data are not presented due to the difficulty in

accurately determining viability in the pupae. Instead, we decided to report the proportion

of larvae that successfully reached adulthood.

(d) Emerged adult bees

For each bee reaching adulthood, the emergence date was recorded to calculate the

developmental requited to complete metamorphosis (i.e. pupal phase). Their body weight

was directly measured using a fine scale (d = 0.1 mg, Sartorius AC120S, Sartorius AG,

Germany). Adult bees were then housed individually in 25 mL snap-cap vial containing a

cardboard bottom (approximately 5.5 x 1.5 cm) and mesh lid. Bees had *ad libitum* access

to 60% (w/v) sugar solution via simple 1.5 mL feeder and kept within the same rearing

room where their natal colonies were maintained. Two-day-old adults, with fully developed

and hardened wings, were then euthanized by freezing and stored at -20°C until further

analysis.

(e) Body size and lipid measurements

As a proxy for adult body size, the intertegular distance (ITD) and head width were

measured using a digital microscope (VHX-500F, Keyence GmbH, Germany)

(Gilgenreiner & Kurze, 2024). In addition, their sex was determined by counting antennal

segments (females have 12, males 13). To determine adult dry mass, each corpse was

carefully dissected by one cut at ventral abdominal segments, then dried at 60°C for three

days (U40, Memmert GmbH & Co. KG, Germany), and weighed (d = 0.1 mg, analytic

balance M-Pact AX224, Sartorius GmbH, Germany). Subsequently, the body lipids were

extracted using petroleum ether for five days. The ether was discarded, and the corpses

were rinsed with fresh ether before being dried for an additional three days. The lipid

content was calculated as the difference between the initial and post-extraction dry

masses. The relative lipid content, serving as a proxy for the fat body size, was calculated

as the percentage of lipid contents relative to the initial dry masses.

(e) Wings

Wings were first detached at the tegulae and embedded side-by-side on a microscope

slide (Roti®Mount, Carl Roth GmbH + Co. KG, Germany). Then, wing pairs were

photographed at 20x magnification (Keyence VHX-500F). Two-dimensional coordinates of

twenty landmarks for each wing (figure S1e) were obtained using TpsDig2 (v2.32) (Rohlf,

2015). While individuals with visibly undeveloped or fully deformed wings (figure S1f) were

excluded from geometric morphometric analysis, we compared the proportion of

individuals with developed, undeformed wings among treatment groups.

(g) Statistical analyses

All statistical analyses and data visualizations were performed using R Statistical Software

in R version 4.4.2 (R Core Team, 2024). The complete code and output are provided in

the electronic supplementary material.

Generalized linear mixed effect models (GLMMs) using the *glmmTMB* package

(Brooks et al., 2017) were used to analyse effects of the treatment length and intensity

(temperature), and their interactions, as fixed factors on emergence rates (i.e. pupae

reaching adulthood), developmental time (i.e. days to emergence), adult mass, dry mass,

ITD, head width, and wing development (i.e. either fully developed or deformed) with

binomial (logit link) or gamma (log) data structures. We included 'colony ID' and 'initial

pupae mass' as random effects to account for colony and larval-specific variability.

Additionally, we analysed the effects on adult mass, dry mass, ITD, and head width for

female (i.e. worker) and male data separately, to account for any sex-specific biases.

Model selection was performed based on the Akaike information criterion (AIC)

and likelihood ratio tests. The final models were compared with their respective null-

models. Model assumptions and dispersion of the data were checked using the *DHARMa*

package (Hartig, 2020). Significance levels ($p < 0.05$) were determined using the *Anova*

function when assumptions were met. Contrast pairwise comparisons between each

treated group with the controls were conducted using the function *emmeans* (Lenth &

Lenth, 2018) adjusted with Holm-Bonferroni correction. When model assumptions were

not met, treatment effects were tested with a Kruskal-Wallis rank sum test, followed by

post-hoc comparisons between each treated group with the controls using Dunn's test

with Holm-Bonferroni correction.

Geometric morphometric analyses of wing size, shape, and asymmetry on fully

developed left and right forewings from females and male were performed separately on

the obtained TPS data using geomorph package (Adams, Collyer, Kaliontzopoulou, &

Baken, 2024), following a similar approach as previous studies (Aytekin, Terzo, Rasmont,

& Çağatay, 2007; Gérard et al., 2018). Briefly, right wings were first mirrored to ensure

landmark correspondence. Missing landmark data, representing morphological variation,

were accounted for using the function *estimate.missing*. To remove non-shape variation, a

Generalized Procrustes Analysis (GPA) was performed using *gpagen*. A Principal

Component Analysis (PCA) was conducted on the Procrustes coordinates using

gm.prcomp and PCA scores (PC1 and PC2) calculated and data distribution was visually

assessed. Wing with PC1 z-scores exceeding ± 2 standard deviations were removed as

outliers, where PC1 explained 71% of the data's variation. Treatment groups with fewer

than six individuals were also excluded from further analysis. Then, centroid size and

mean centroid size were calculated for each wing and each bee, respectively, as a

measure of wing size. To analyse shape variation, a permutational multivariate analysis of

variance (PERMANOVA) was performed using the *adonis2* function. Homoscedasticity

was evaluated using *betadisper*, and pairwise comparisons were conducted using

pairwise.adonis with Holm-Bonferroni correction. Directional asymmetry (DA) was

calculated as the difference between left and right wing Procrustes coordinates, and DA

magnitude was calculated as the mean Euclidean distance between corresponding

landmarks. To assess fluctuating asymmetry (FA), a bilateral symmetry analysis was

performed using *bilat.symmetry* and calculating the FA magnitude as the Euclidean norm

of the FA component. The effects of treatment on centroid size, DA and FA magnitude

were tested using linear regression models (LM) when assumptions were met, followed by

contrast pairwise comparisons for significant results as described above. Kruskal-Wallis

tests were used as non-parametric alternative, followed by Dunn's post-hoc test to

compare treatment groups to the control by applying Holm-Bonferroni corrections.

3. Results

We found a strong interaction effect of heatwave temperature and length on the proportion

of pupae reaching adulthood (GLMM: χ^2 = 23.056, df = 1, p < 0.0001; figure 1a). Pupae of

treatment 38C3d and 38C5d were 43% (z = 3.523, p < 0.01) and 83 % (z = 6.563, p <

0.0001) less probable to reach adulthood compared to controls, respectively. Heatwave

intensity also significantly affected pupal developmental duration (Kruskal-Wallis: χ^2 =

69.584, df = 6, p < 0.0001; figure 1b; female and male data presented separately in figure

S2). While an exposure to 36°C shortened development compared to controls (Dunn's

test: 36C5d, z = 3.253, p < 0.01; 36C7d, z = 3.142, p < 0.01), an exposure to 38°C

prolonged development (38C3d, z = -4.393, p < 0.0001; 38C3d, z = -2.739, p < 0.01).

There was no significant effect found for the 37°C exposure compared to controls (37C3d,

z = -1.361, p > 0.5; 37C5d, z = -1.164, p > 0.5).

Figure 1. Direct effects of heatwaves of varying intensity on (a) emergence rate and (b) developmental

duration in *B. terrestris* pupae. Controls (Ctrl, reared at constant 34°C, blue) were compared to individuals

exposed to increased temperatures: of 36°C for 5 days (light warm yellow) and 7 days (warm yellow), 37°C for

3 days (light reddish-orange) and 5 days (reddish orange), and 38°C for 3 days (light red) and 5 days (deep

red). Sample sizes are indicated by white numbers and significant differences are marked by asterisks (* p <

0.05, ** p < 0.01, *** p < 0.001).

When testing the effects of heatwave treatments on adult mass, we found a

significant effect (Kruskal-Wallis: χ^2 = 30.552, df = 6, p < 0.0001) with 36C7d pupae

resulting in significantly lighter adults compared to controls (Dunn's test: z = 2.576, p <

0.05) and 38C3d pupae resulting in heavier adults (z = -2.679, p < 0.05). However, sex-

specific analyses revealed no significant treatment effects on adult mass (Kruskal-Wallis:

females, χ^2 = 9.125, df = 6, p > 0.05, figure 2a; males, χ^2 = 6.526, df = 6, p > 0.05, figure

S3a). While there was a significant effect on female head width (Kruskal-Wallis: χ^2 =

16.131, df = 6, p < 0.05, figure 2b), we found no significant differences between treatment

groups and controls (Dunn's tests: p > 0.05). However, female head width varied with

temperature (Kruskal-Wallis: χ^2 = 12.311, df = 3, p < 0.01), where adults from pupae

exposed to 38°C had slightly wider heads than controls (z = -2.227, p < 0.05). We found

no significant effect on female ITD (Kruskal-Wallis: χ^2 = 9.571, df = 6, p > 0.05, figure 2c).

Similarly, we also found no significant effects on male body size proxies (Kruskal-Wallis:

head width, χ^2 = 9.532, df = 6, p > 0.05, figure S3b; ITD, χ^2 = 4.743, df = 6, p > 0.05, figure

S3c).

In contrast, we found that heatwave intensity significantly influenced relative lipid

content in adult females (Kruskal-Wallis: χ^2 = 25.069, df = 6, p < 0.0001, figure 2d), but

not in males (Kruskal-Wallis: χ^2 = 8.287, df = 6, p > 0.05, figure S3d). We found an

increase in relative lipid content in adult females of 37C5d (Dunn's test: z = -2.592, p <

0.05), 38C3d (z = -2.844, p < 0.05), and 38C5d (z = -3.012, p < 0.01) compared to

controls.

Figure 2. Effects of pupal heatwave exposure on adult female morphometrics and lipid content in *B.*

***terrestris*:** (a) adult mass, (b) head width, (c) intertegular distance (ITD), (d) and relative lipid content (2-day-

old adults). Controls (34°C, blue) were compared to individuals exposed to increased temperatures: of 36°C

for 5 days (light warm yellow) and 7 days (warm yellow), 37°C for 3 days (light reddish-orange) and 5 days

(reddish orange), and 38°C for 3 days (light red) and 5 days (deep red). Asterisks denote significant

differences (* $p < 0.05$, ** $p < 0.01$).

We found that heatwave intensity impacted wing development (GLMM: $\chi^2 = 9.992$,

df = 1, $p < 0.01$, figure 3a). Specifically, 38C3d pupae were four times more likely to

develop deformed wings compared to the controls (z = -3.367, $p < 0.01$), while 38C5d

pupae were nearly seven times more likely of developing deformed wings (z = -3.920, $p <$

0.001). Pupae of the 36C7d treatment showed a marginally significant, twofold increase in

deformed wings (z = -2.482, p = 0.052).

Our geometric morphometrics analysis without deformed wings indicated a

significant effect on female wing shape (PERMANOVA: F = 12.484, R^2 = 0.21, df = 5, p <

0.001, figure 3b), but not on male wing shape (F = 0.269, R^2 = 0.01, df = 2, p > 0.05,

figure S4b). Pairwise comparisons revealed significant differences between all treatment

groups and the controls (36C5d, F = 9.860, p < 0.05; 36C7d, F = 7.599, p < 0.05; 37C3d,

F = 25.643, p < 0.05; 37C5d, F = 31.365, p < 0.05; 38C3d, F = 29.824, p < 0.05).

Similarly, we found that heatwave exposure affected centroid wing size in females

(Kruskal-Wallis: χ^2 = 18.633, df = 5, p < 0.01, figure S3c), but not in males (χ^2 = 0.657, df

= 2, p > 0.05, figure S4b). Pairwise comparisons indicated marginally significant decrease

in wing size for females of the 36C7d treatment (z = -2.279, p = 0.057).

Furthermore, our heatwave treatment significantly affected the DA in female wings

(LM: F = 9.466, df = 5, p < 0.0001, figure 3d), but not FA (Kruskal-Wallis: χ^2 = 3.903, df =

5, p > 0.05, figure 3e). Females exposed to 36°C showed higher DA (36C5d, t = -2.520, p

< 0.05; 36C7d, t = -2.536, p < 0.05), while females exposed to 37°C (37C3d, t = 2.655, p

< 0.05; 37C5d, t = 2.932, p < 0.05) and 38°C (38C3d, t = 2.500, p < 0.05) showed lower

DA compared to controls. In males, we found no effect on DA (Kruskal-Wallis: χ^2 = 1.318,

df = 2, p > 0.05, figure S4d), but on FA (χ^2 = 6.932, df = 2, p < 0.05, figure S4e). Males of

37C3d (z = -1.884, p < 0.05) and 37C5d (z = -2.547, p < 0.05) showed slight increase in

FA compared to controls.

321
322
Figure 3. Heatwave effects on wing development in *B. terrestris*. (a) Proportion of deformed wings in

emerged adult bees for each treatment group. (b) Principal component analysis (PCA) of wing morphology in

emerged females with fully developed wings across heatwave treatments (see legend; 95% confidence

intervals are shown in figure S4a). (c) Influence of heatwave treatments on wing size (centroid size) and the

magnitude of (d) directional and (e) fluctuating asymmetry in fully developed female wings. Controls (34°C,

blue) were compared to individuals exposed to increased temperatures: of 36°C for 5 days (light warm yellow)

and 7 days (warm yellow), 37°C for 3 days (light reddish-orange) and 5 days (reddish orange), and 38°C for 3

days (light red). Asterisks denote significant differences (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

4. Discussion

Our study provides compelling evidence that heatwave length and severity significantly impact pupal development in the key pollinator *B. terrestris* with notable effects on emergence rate (figure 1a), developmental duration (figure 1b), lipid stores (figure 2d), and wing development (figure 3). These findings highlight the extreme vulnerability of the pupal stage to heatwaves, particularly at 38°C, which appears to be a critical threshold for development disruption, aligning with our previous study on L4 larvae (Woegler & Kurze, 2024). Although our treatments are extreme, they reflect potential heatwave scenarios in Europe (Lhotka & Kyselý, 2024; Lhotka et al., 2018; Perkins-Kirkpatrick & Lewis, 2020; Simolo & Corti, 2025; Stillman, 2019). Recent heatwaves reaching or even exceeding temperatures of 38°C were, for example, reported in parts of the British Isles, Mediterranean and Eastern Europe (Barriopedro, Fischer, Luterbacher, Trigo, & García-Herrera, 2011; Lhotka & Kyselý, 2024; Rita et al., 2020).

Most notably, our findings show that emergence rates decreased by almost twofold and fivefold in pupae exposed to 38°C for 3 and 5 days, respectively, compared to controls (figure 1a). Despite *in vitro* rearing of L4 larvae and pupae deviates from natural nest conditions, the high emergence rate of controls suggests that our experimental setup allowed for normal development. While 38°C exposure slightly prolonged pupal development, a 36°C exposure slightly accelerated it (figure 1b), suggesting a potential developmental benefit in pupae to temperature elevation up to 37°C, followed by severe disruption at 38°C. Given the relative heat tolerance of our model species, *B. terrestris*, we expect cold-adapted bumblebee species such as *B. lapidarius*, *B. alpinus*, and *B. poralris* to be even more vulnerable to heatwaves (Martinet et al., 2021; Zambra, Martinet, Brasero, Michez, & Rasmont, 2020). Therefore, extreme heatwave events could have a devastating impact on bumblebee populations, which could lead to local extinctions and a decline in species diversity, with significant implications for crop pollination and ecosystem services (Ghisbain et al., 2024).

We observed no clear heatwave effects on sex-specific body size proxies (figures

2 and S3), except for an increased female head width at 38°C. These results contrast with

previous studies showing that colonies exposed to prolonged heat stress would produce

smaller offspring, indicated by smaller ITD (Guiraud et al., 2021; Gérard et al., 2023) and

body mass (Perl et al., 2022), potentially reflecting resource allocation strategies such as

reduced larval feeding. Alternatively, heat stressed larvae may have a decreased body

weight gain due to reduced food consumption (Woegler & Kurze, 2024). We speculate

that our morphometric data potentially reflect a selective advantage of larger pupae under

heat stress. Although small insect body sizes should generally be favoured under warmer

environmental conditions over the long term (González-Tokman et al., 2020), aligning with

Bergmann's rule, larger workers in numerous ant species have been described to be more

heat resistant than smaller workers (Perez & Aron, 2020). In addition, we found increased

relative lipid content (fat body) in females with increasing heatwave intensity (figure 2d),

indicating a potentially higher heat stress resilience due to increased energy reserves,

immunocompetence, and heat shock protein (HSP) synthesis (Arrese & Soulages, 2010;

King & MacRae, 2015; Korner & Schmid-Hempel, 2004; Vanderplanck et al., 2021;

Vesterlund, Lilley, Van Ooik, & Sorvari, 2014). Interestingly, the relative lipid content was

not affected in heat stressed L4 larvae, further underscoring the differential vulnerability of

developmental stages (Woegler & Kurze, 2024).

Wing development was severely impaired by heatwave intensity, notably with a

fourfold and sevenfold increase in wing deformities in emerged adult bees from pupae

exposed to 38°C for 3 and 5 days, respectively (figure 3a). Even excluding individuals with

these deformities, females wing shape altered for all treatments (figure 3b), which appears

to be a natural response in colonies under heat and parasitic stress (Gérard, Cariou, et

al., 2022; Gérard et al., 2018). However, our data shows only marginal effects on wing

size (figure 3c) with a similar pattern as for the other body size proxies (figure 2). DA was

affected by temperature in females, with increasing DA at 36°C and decreasing DA at

37°C and 38°C (figure 3d), potentially also reflecting the different selection pressures

among treatments. While FA remained unaffected in females (figure 3e), confirming

previous studies in honeybees (Jones, Helliwell, Beekman, Maleszka, & Oldroyd, 2005)

and bumblebees (Gérard et al., 2018), males showed a slight increase in FA at 37°C

(figure S4e).

In conclusion, our study provides novel insights into the direct impacts of varying

heatwaves scenarios on *B. terrestris* pupal development, eliminating confounding colony-

level effects. While our experimental design sacrifices some ecological realism, it enabled

precise control over timing and temperature to match potential heatwave scenarios,

making our study relevant for predicting future pollinator declines. Our findings highlight a

critical heat stress threshold at 38°C, where severe disruptions in bumblebee pupal

development manifest by particularly impairing emergence and wing development. Given

the heat tolerance of *B. terrestris*, we expect other cold-adapted bumblebee species to be

even more vulnerable to increasingly more frequent and severe heatwaves as climate

change continues to cause more extreme weather events. Therefore, our findings

underscore the profound fitness implications for bumblebee colonies with cascading

consequences for whole populations, biodiversity, crop pollination and ecosystem

services of bumblebees as key pollinators.

Data accessibility

All data and statistical code output supporting the findings of this study are provided in the

electronic supplementary material.

Acknowledgments

We thank Nadine Wildemann, Franziska Altmann, and Erik Köhler for their assistance in

the laboratory. We would like to extend our gratitude to Dr. Tomer J. Czaczkes for

sponsoring the bumblebee colonies.

Author's contributions

SL and CK collected the data. CK conceived, designed, and supervised the study;

analysed and interpreted the data; and wrote the manuscript. SL gave approval for

publication.

Ethics statement

This study was conducted in accordance with the ethical regulations of the German

Animal Welfare Act (TierSchG) for conducting experiments with insects.

Funding

This study was carried out without any third-party funding.

Competing interests

The authors have no competing interests.

References

Adams, D., Collyer, M., Kaliontzopoulou, A., & Baken, E. (2024). Geometric morphometric analyses of 2D and 3D landmark data, version 4.0. 8. In: R Foundation for Statistical Computing.

Arrese, E. L., & Soulages, J. L. (2010). Insect fat body: energy, metabolism, and regulation. *Annual review of entomology, 55*, 207-225.

Aytekin, M. A., Terzo, M., Rasmont, P., & Çağatay, N. (2007). *Landmark based geometric morphometric analysis of wing shape in Sibiricobombus Vogt (Hymenoptera: Apidae: Bombus Latreille).* Paper presented at the Annales de la Société entomologique de France.

Barriopedro, D., Fischer, E. M., Luterbacher, J., Trigo, R. M., & García-Herrera, R. (2011). The hot summer of 2010: redrawing the temperature record map of Europe. *Science, 332*(6026), 220-224.

Bretzlaff, T., Kerr, J. T., & Darveau, C.-A. (2024). Handling heatwaves: balancing thermoregulation, foraging and bumblebee colony success. *Conservation Physiology, 12*(1). doi:10.1093/conphys/coae006

Brooks, M. E., Kristensen, K., Van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., . . . Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R journal, 9*(2), 378-400.

Cameron, S. A., Lozier, J. D., Strange, J. P., Koch, J. B., Cordes, N., Solter, L. F., & Griswold, T. L. (2011). Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences, 108*(2), 662-667.

Cameron, S. A., & Sadd, B. M. (2020). Global trends in bumble bee health. *Annual review*
of entomology, 65, 209-232.
Garcia, R. A., Cabeza, M., Rahbek, C., & Araújo, M. B. (2014). Multiple dimensions of
climate change and their implications for biodiversity. *Science, 344*(6183),
1247579.
Ghisbain, G., Thiery, W., Massonnet, F., Erazo, D., Rasmont, P., Michez, D., & Dellicour,
S. (2024). Projected decline in European bumblebee populations in the twenty-first
century. *Nature, 628*(8007), 337-341.
Gilgenreiner, M., & Kurze, C. (2024). Age dominates flight distance and duration, while
body size shapes flight speed in *Bombus terrestris* L.(Hymenoptera: Apidae).
Proceedings of the Royal Society B, 291(2027), 20241001.
González-Tokman, D., Córdoba-Aguilar, A., Dáttilo, W., Lira-Noriega, A., Sánchez-
Guillén, R. A., & Villalobos, F. (2020). Insect responses to heat: physiological
mechanisms, evolution and ecological implications in a warming world. *Biological*
Reviews, 95(3), 802-821.
Goulson, D., Lye, G. C., & Darvill, B. (2008). Decline and conservation of bumble bees.
Annual Review Entomology, 53(1), 191-208.
Guiraud, M., Cariou, B., Henrion, M., Baird, E., & Gérard, M. (2021). Higher
developmental temperature increases queen production and decreases worker
body size in the bumblebee *Bombus terrestris. Journal of Hymenoptera Research,*
88, 39-49.
Gérard, M., Amiri, A., Cariou, B., & Baird, E. (2022). Short-term exposure to heatwave-like
temperatures affects learning and memory in bumblebees. *Global Change Biology,*
28(14), 4251-4259.
Gérard, M., Cariou, B., Henrion, M., Descamps, C., & Baird, E. (2022). Exposure to
elevated temperature during development affects bumblebee foraging behavior.
Behavioral Ecology, 33(4), 816-824.
Gérard, M., Guiraud, M., Cariou, B., Henrion, M., & Baird, E. (2023). Elevated
developmental temperatures impact the size and allometry of morphological traits
of the bumblebee *Bombus terrestris. Journal of Experimental Biology, 226*(8),
jeb245728.
Gérard, M., Michez, D., Debat, V., Fullgrabe, L., Meeus, I., Piot, N., . . . Vanderplanck, M.
(2018). Stressful conditions reveal decrease in size, modification of shape but
relatively stable asymmetry in bumblebee wings. *Scientific Reports, 8*(1), 15169.
Hartig, F. (2020). DHARMa: residual diagnostics for hierarchical (multi-level/mixed)
regression models. *R package version 0.3, 3*(5).
IPCC. (2023). Summary for Policymakers. In *Climate Change 2022 – Impacts, Adaptation*
and Vulnerability: Working Group II Contribution to the Sixth Assessment Report of

the Intergovernmental Panel on Climate Change (pp. 3-34). Cambridge:

Cambridge University Press.

Jentsch, A., Kreyling, J., & Beierkuhnlein, C. (2007). A new generation of climate-change

experiments: events, not trends. *Frontiers in Ecology and the Environment, 5*(7),

365-374.

Jones, J. C., Helliwell, P., Beekman, M., Maleszka, R., & Oldroyd, B. P. (2005). The

effects of rearing temperature on developmental stability and learning and memory

in the honey bee, *Apis mellifera*. *Journal of comparative physiology A, 191*, 1121-

1129.

Kato, Y., Kikuta, S., Barribeau, S. M., & Inoue, M. N. (2022). In vitro larval rearing method

of eusocial bumblebee *Bombus terrestris* for toxicity test. *Scientific Reports 12*(1),

15783.

Kerr, J. T., Pindar, A., Galpern, P., Packer, L., Potts, S. G., Roberts, S. M., . . .

Richardson, L. L. (2015). Climate change impacts on bumblebees converge

across continents. *Science, 349*(6244), 177-180.

King, A. M., & MacRae, T. H. (2015). Insect heat shock proteins during stress and

diapause. *Annual review of entomology, 60*(1), 59-75.

Klein, A.-M., Vaissiere, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A.,

Kremen, C., & Tscharntke, T. (2007). Importance of pollinators in changing

landscapes for world crops. *Proceedings of the Royal Society B, 274*, 303-313.

doi:10.1098/rspb.2006.3721

Korner, P., & Schmid-Hempel, P. (2004). In vivo dynamics of an immune response in the

bumble bee *Bombus terrestris*. *87*(1), 59-66.

Kuo, Y., Lu, Y.-H., Lin, Y.-H., Lin, Y.-C., & Wu, Y.-L. (2023). Elevated temperature affects

energy metabolism and behavior of bumblebees. *Insect Biochemistry and*

Molecular Biology, 155, 103932.

Lenth, R., & Lenth, M. R. (2018). Package 'lsmeans'. *The American Statistician, 34*(4),

216-221.

Lhotka, O., & Kyselý, J. (2024). Three-dimensional analysis reveals diverse heat wave

types in Europe. *Communications Earth & Environment, 5*(1), 323.

Lhotka, O., Kyselý, J., & Farda, A. (2018). Climate change scenarios of heat waves in

Central Europe and their uncertainties. *Theoretical and applied climatology,*

131(3), 1043-1054.

Ma, C.-S., Ma, G., & Pincebourde, S. (2021). Survive a warming climate: insect responses

to extreme high temperatures. *Annual Review of Entomology, 66*, 163-184.

Maebe, K., Hart, A. F., Marshall, L., Vandamme, P., Vereecken, N. J., Michez, D., &

Smagghe, G. (2021). Bumblebee resilience to climate change, through plastic and

adaptive responses. *Global Change Biology, 27*(18), 4223-4237. doi:https://doi.org/10.1111/gcb.15751

Martinet, B., Dellicour, S., Ghisbain, G., Przybyla, K., Zambra, E., Lecocq, T., . . . Rasmont, P. (2021). Global effects of extreme temperatures on wild bumblebees. *Conservation Biology, 35*(5), 1507-1518.

Meehl, G. A., & Tebaldi, C. (2004). More intense, more frequent, and longer lasting heat waves in the 21st century. *Science, 305*(5686), 994-997.

Nooten, S. S., Korten, H., Schmitt, T., & Kárpáti, Z. (2024). The heat is on: reduced detection of floral scents after heatwaves in bumblebees. *Proceedings of the Royal Society B, 291*(2029), 20240352.

Perez, R., & Aron, S. (2020). Adaptations to thermal stress in social insects: recent advances and future directions. *Biological Reviews, 95*(6), 1535-1553.

Perkins-Kirkpatrick, S. E., & Lewis, S. C. (2020). Increasing trends in regional heatwaves. *11*(1), 3357.

Perl, C. D., Johansen, Z. B., Moradinour, Z., Guiraud, M., Restrepo, C. E., Wen Jie, V., . . . Baird, E. (2022). Heatwave-Like Events During Development Are Sufficient to Impair Bumblebee Worker Responses to Sensory Stimuli. *Frontiers in Ecology and Evolution, 9.* doi:10.3389/fevo.2021.776830

Potts, S. G., Biesmeijer, J. C., Kremen, C., Neumann, P., Schweiger, O., & Kunin, W. E. (2010). Global pollinator declines: trends, impacts and drivers. *Trends in Ecology and Evolution, 25*, 345-353. doi:10.1016/j.tree.2010.01.007

Potts, S. G., Imperatriz-Fonseca, V., Ngo, H. T., Aizen, M. A., Biesmeijer, J. C., Breeze, T. D., . . . Settele, J. (2016). Safeguarding pollinators and their values to human well-being. *Nature, 540*, 220-229.

Quinlan, G. M., Feuerborn, C., Hines, H. M., & Grozinger, C. M. (2023). Beat the heat: thermal respites and access to food associated with increased bumble bee heat tolerance. *Journal of Experimental Biology, 226*(17), jeb245924.

Rita, A., Camarero, J. J., Nolè, A., Borghetti, M., Brunetti, M., Pergola, N., . . . Ripullone, F. (2020). The impact of drought spells on forests depends on site conditions: The case of 2017 summer heat wave in southern Europe. *Global change biology, 26*(2), 851-863.

Rohlf, F. J. (2015). The tps series of software. *Hystrix, 26*(1), 9-12.

R Core Team. (2024). R: A Language and Environment for Statistical Computing. In: R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Sepúlveda, Y., Nicholls, E., Schuett, W., & Goulson, D. (2024). Heatwave-like events affect drone production and brood-care behaviour in bumblebees. *PeerJ, 12*, e17135.

Simolo, C., & Corti, S. (2025). Enhanced risk of hot extremes revealed by observation-
constrained model projections. *Communications Earth & Environment, 6*(1), 165.

Soroye, P., Newbold, T., & Kerr, J. (2020). Climate change contributes to widespread
declines among bumble bees across continents. *Science, 367*(6478), 685-688.

Stillman, J. H. (2019). Heat waves, the new normal: summertime temperature extremes
will impact animals, ecosystems, and human communities. *Physiology, 34*(2), 86-
100.

Theodorou, P., Kühn, O., Baltz, L. M., Wild, C., Rasti, S. L., Bucksch, C. R., . . . Kurze, C.
(2022). Bumble bee colony health and performance vary widely across the urban
ecosystem. *Journal of Animal Ecology, 91*(10), 2135-2148.
doi:https://doi.org/10.1111/1365-2656.13797

Vanderplanck, M., Breeze, T., Cini, E., Potts, S. G., & Senapathi, D. (2021). Monitoring
bee health in European agroecosystems using wing morphology and fat bodies.
One Ecosystem, 6.

Vesterlund, S.-R., Lilley, T., Van Ooik, T., & Sorvari, J. (2014). The effect of overwintering
temperature on the body energy reserves and phenoloxidase activity of
bumblebee *Bombus lucorum* queens. *Insectes sociaux, 61*, 265-272.

Vogt, F. D. (1986). Thermoregulation in bumblebee colonies. I. Thermoregulatory versus
brood-maintenance behaviors during acute changes in ambient temperature.
Physiological zoology, 59(1), 55-59.

Weidenmüller, A., Kleineidam, C., & Tautz, J. (2002). Collective control of nest climate
parameters in bumblebee colonies. *Animal Behaviour, 63*(6), 1065-1071.

Woegler, L., & Kurze, C. (2024). Experimental short-term heatwaves negatively impact
body weight gain and survival during larval development in a wild pollinator.
bioRxiv, 2024.2010.2009.617416. doi:10.1101/2024.10.09.617416

Zambra, E., Martinet, B., Brasero, N., Michez, D., & Rasmont, P. (2020). Hyperthermic
stress resistance of bumblebee males: test case of Belgian species. *Apidologie,*
51, 911-920.

Supplementary information

Heatwave length and severity drive development disruption in a key pollinator

Sandra Laußer[1] and Christoph Kurze[1]*

[1] Institute for Zoology and Evolutionary Biology, University of Regensburg, Germany

* Corresponding author, email: Christoph.Kurze@ur.de

Figure S1. *In vitro* larval rearing of L4 (a-d) and wing morphology analysis (e, f). (a) L4 larvae were identified by the small opening for food provisioning (black arrows) and carefully collected. (b) Design of 3D-printed polylactide (PLA) artificial brood cells (capacity: 0.6 ml or 0.96 ml). (c) L4 larvae inside artificial brood cells in a 24-well plate. (d) Observation of L4 feeding on medium (see also supplementary video 1). (e) Fully developed right forewing showing the positions of the 20 landmarks (red dots) used for the geometric morphometric analysis, including centroid size, wing shape, directional asymmetry (DA), and fluctuating asymmetry (FA). (f) Representative examples of left and right forewings from two emerged adults with slight (top) and more severe (bottom) wing deformities. Individuals with deformed wings were excluded from further geometric morphometric analyses. Scale bar = 1mm.

Figure S2. Direct effects of heatwaves of varying intensity on (a) female and (b) male developmental duration in *B. terrestris* pupae. Controls (Ctrl, reared at constant 34°C, blue) were compared to individuals exposed to increased temperatures: of 36°C for 5 days (light warm yellow) and 7 days (warm yellow), 37°C for 3 days (light reddish-orange) and 5 days (reddish orange), and 38°C for 3 days (light red) and 5 days (deep red). Significant differences are marked by asterisks (* p < 0.05, ** p < 0.01, *** p < 0.001).

Figure S3. Effects of heatwaves intensity and duration during *B. terrestris* pupal development on adult male morphometrics and lipid content: (a) adult mass, (b) head width, (c) intertegular distance (ITD), (d) and relative lipid content (2-day-old adults). Controls (34°C, blue) were compared to individuals exposed to increased temperatures: of 36°C for 5 days (light warm yellow) and 7 days (warm yellow), 37°C for 3 days (light reddish-orange) and 5 days (reddish orange), and 38°C for 3 days (light red) and 5 days (deep red). Asterisks denote significant differences (* p < 0.05, ** p < 0.01).

Figure S4. Heatwave effects on wing development in male *B. terrestris*. (a, b) General wing morphology variation visualised by principal component analysis (PCA) for females (a) and males (b) with fully developed wings across heatwave treatments (see legend). 95% confidence intervals are shown as ellipses. (c) Influence of heatwave treatments on wing size (centroid size) and the magnitude of (d) directional (DA) and (e) fluctuating asymmetry (FA) in fully developed male wings. Controls (34°C, blue) were compared to individuals exposed to increased temperatures of 37°C for 3 days (light reddish-orange) and 5 days (reddish orange). Asterisks denote significant differences (* p < 0.05).