A breath of fresh air: comparative evaluation of passive versus active airborne eDNA sampling strategies

Hugo Jager[1], Krijn B. Trimbos[1], Jan-Maarten Luursema[2], Adrianus G.C.L. Speksnijder[3,4], Kathryn A. Stewart[1*]

[1]Institute of Environmental Sciences, Leiden University, The Netherlands

[2] Department of Surgery, Radboud University Medical Center, Nijmegen, the Netherlands

[3] Leiden University of Applied Sciences, Leiden, the Netherlands

[4] Naturalis Biodiversity Center, Leiden, the Netherlands

* Corresponding author: k.a.stewart@cml.leidenuniv.nl

Abstract

Terrestrial biodiversity's rapid decline worldwide demands for expansion of high-resolution biomonitoring to support science-based policy. Environmental DNA (eDNA) analysis has proven effective at biomonitoring aquatic systems but remains underexplored for terrestrial habitats, partly due to point-source sampling bias and difficulty in upscaling. Air has recently emerged as a promising substrate for terrestrial biomonitoring yet most studies use active samplers which tend to be expensive, bulky, and require power. We used a newly developed, inexpensive, reusable, and easy-to-use passive airborne eDNA sampler (DNAir Sampler) to capture eDNA suspended in air over 96 hours within Rotterdam Zoo, the Netherlands. Its performance was compared to two commonly used active airborne eDNA samplers for vertebrate diversity detection. In total, 88 species were detected, including 24 zoo residents. The DNAir Sampler was the most effective at detecting zoo residents, surpassing active samplers in species richness within 48 hours and continuing to accumulate new species beyond 96 hours. It also detected the furthest species signal (515m). Zoo airborne eDNA further demonstrated a positive correlation to species total biomass. Our findings indicate that for long, unsupervised biomonitoring, passive airborne eDNA sampling outperforms active

sampling for assessing vertebrate communities and potentially reduces detection noise in

stochastic air eDNA signals.

Keywords

biomonitoring, environmental DNA, airborne DNA, passive samplers, DNAir Sampler, verte-

brate biodiversity, bioaerosols

Synopsis

Passive airborne eDNA sampling outperforms active methods, accumulates species over

time, correlates with biomass, and enables scalable biodiversity monitoring across land-

scapes.

Introduction

Global biodiversity is essential for ecosystem resilience and critical services such as food security and medicine [1,2]. However, anthropogenic pressures, including climate change, habitat destruction, and globalisation threaten biodiversity, necessitating high-resolution biomonitoring to guide conservation efforts [3,4].

Traditional biomonitoring methods are often time-consuming, require expert taxonomic knowledge, can be invasive to the species and fail to document rare species [5]. Environmental DNA (eDNA) approaches, which collect genetic material directly from the environment rather than individuals [5-7], have gained widespread use [4]. These approaches are non-invasive, highly sensitive, cost effective, and capable of detecting taxa across the tree of life while reducing observer bias and enabling large-scale monitoring [8,9]. In many cases, eDNA has outperformed traditional methods [10,11]. While eDNA sampling from water is well-established [12-14], terrestrial applications remain underdeveloped partly due to challenges in identifying optimal substrates [15]. Various terrestrial substrates, including soil, plants and spider webs (among others) have been explored but often yield limited taxonomic coverage and rely on point-source samples that may fail to give a comprehensive and representative estimate of biodiversity [16-19].

Airborne eDNA sampling offers a novel approach for terrestrial biodiversity monitoring. Air contains bioaerosols—genetic material from microorganisms, pollen, and animal tissue fragments [20]. [21] first demonstrated animal DNA detection in air using active (powered) air pumps, with subsequent studies confirming its efficacy across various environments, including zoos and natural habitats [22-24]. Recent research has detected insects, birds, frogs, and mammals in field conditions, highlighting the potential of airborne eDNA for large-scale biodiversity monitoring [7,20,25].

Most air eDNA studies rely on active samplers, which, despite their effectiveness, are costly, bulky, and require power sources, limiting their use in remote areas [26]. Passive air eDNA sampling provides a cheaper and more mobile alternative, with promising results for

detecting diverse taxa [27-29]. However, questions remain regarding its sensitivity and spa-

tial resolution compared to active methods, particularly given evidence of long-range airborne

eDNA dispersal [22] and continuous accumulation over time [29]. Understanding airborne

eDNA's spatial and temporal dynamics is crucial for optimizing sampling strategies [30].

Our study evaluated the efficacy and sensitivity of a newly-designed passive airborne

eDNA sampler (the DNAir Sampler) against two, commonly used, active air eDNA samplers

(Pollensniffer and Coriolus Micro) in Rotterdam Zoo, Netherlands. The DNAir sampler, de-

signed to enhance DNA capture efficiency by directing airflow over a protected DNA-binding

filter, aims to mitigate the limitations of previous passive methods that rely on surface settle-

ment [27, 29, 31]. We assessed species detection rates, expected zoo vertebrate recovery,

species dispersion distances, accuracy in detecting known zoo residents, and the correlation

between species biomass and airborne eDNA abundance. Additionally, we compared spe-

cies accumulation curves to determine optimal sampling strategies across the three eDNA

samplers.

MATERIALS AND METHODS

Study site

99 We collected air samples at Rotterdam Zoo (Diergaarde Blijdorp, Netherlands; Figure 1), a

100 32-acre zoo wherein sampling took place at five outdoor points and one indoor point. Sur-

101 rounded by residential areas and water bodies, the zoo houses diverse avian and mammal-

102 ian species in spacious outdoor enclosures. Besides listed zoo animals, expected eDNA

103 sources include food, visitors, their pets, and nearby wildlife.

FIGURE 1 Map of Rotterdam Zoo (the Netherlands) including the 6 sampling locations (numbered circles). Location 6 represents an indoor site. Photo insert depicts sampling setup of the Pollensniffer (left bottom), Coriolis Micro (left top), and the passive DNAir Samplers (right).

Airborne eDNA sampling

Between March 25 and April 12 2024, we collected air samples at six locations over three 5-day sampling campaigns (Table S1). Locations were selected based on 1) secure placement of passive airborne eDNA samplers out of visitor's reach, 2) sufficient distance to avoid re-dundant detections, and 3) proximity to at least four enclosures.

Three air sampling devices were used, each varying in mobility, cost and flow-rate: two active samplers - the Pollensniffer (~8 L/min) [32] and the Coriolis Micro (~100 L/min) [33], and the passive DNAir sampler [34] which relied on external air movement. On the first day of each sampling campaign, we deployed 5 passive samplers at ~2.5m height in trees at each location (including the indoor location) and took 1 m-high air samples with both active samplers. To minimize disruption (in consultation with zoo staff) and in accordance with sug-

gested sampling practice by the manufacturers, both active samplers were operated simulta-

neously for 10 minutes per sampling event. Passive samplers were collected at 6, 24, 48, 72,

and 96 hours after deployment. All equipment was sterilized between uses with 30% bleach

solution followed by ultrapure water rinsing. Passive DNAir samples were collected on 0.7µm

glass fibre filters, stored in 2 ml Eppendorf tubes. The Coriolis Micro samples were collected

in 4 mL PBS-filled cones, sealed immediately after sampling. Samples were handled with

facemasks and nitrile gloves and transported on ice before storage the same day at -20°C

until DNA extraction.

For each campaign, field negative controls were collected: glass fibre filters placed

inside without being hung (DNAir Sampler), PBS solution placed within the sampling cup

(Coriolis Micro), and a Vaseline-coated slide (Pollensniffer). In total, 90 DNAir samples, 18

Pollensniffer samples, 18 Coriolis Micro samples, and three field negatives per device across

the three weeks were obtained.

DNA extraction

For extraction, we fit each sample in a 2mL Eppendorf tube: DNAir filters were cut

into small pieces, the Pollensniffer slides were removed from the cassettes and

folded with the exposed side facing inwards, the 4mL Coriolis Micro PBL samples

were transferred to an Eppendorf tube, centrifuged for two minutes at 10,000rpm,

thereafter supernatant removed and pellets retained. We extracted DNA for all sam-

ples following the protocol of [35]. Samples were incubated overnight in a 56°C lysis buffer

for 14.5 hours. QIAshredder (QIAGEN) was then used to release lysed cellular content from

the sample substrates, which after centrifuging for three minutes at 11,000 rpm resulted in

two 390 µL tubes per sample. We vortexed samples and added 700µL AL:EtOH (1:1) to each

tube. Finally, we eluted DNA in 100µL AE buffer. All samples were stored at -20°C until me-

tabarcoding.

DNA metabarcoding

We used two vertebrate primer sets for metabarcoding: 16S rRNA primer

(16Smam1/16Smam2) targeting mammals (~95bp) [36], and 12S rRNA (12S05 forward

/12S05 reverse) targeting vertebrates (~97bp) [37]. All primers had 5'-overhang adapters for

Nextera XT (Illumina) indexing. A 1:100 dilution of harbor porpoise (*Phocoena phocoena*)

DNA (Naturalis Biodiversity Center, Netherlands) served as a positive PCR control, with mul-

tiple negative controls to detect contamination.

For the 16S primer set, 20µL reactions consisted of 4µL DNA template, 10µL

TaqMan™ Environmental Master Mix 2.0, 1µL of each primer (10µM), 3µL RNase-free water,

and 1µL human blocker (60µM) [38]. The thermal cycling profile was 95°C for 10 minutes,

followed by 40 cycles of 95°C for 12 s, 59°C for 30 s, 70°C for 25 s, and a final extension

time of 72°C for 7 minutes. The thermal cycling profile was 95°C for 10 min, followed by 40

cycles of 95°C for 12 s, 59°C for 30 s, 70°C for 25 s, and a final extension at 72°C for 7 min.

The 12S reactions followed the same protocol but omitted the human blocker [39] and had a

cycling profile of 94°C for 30 s, 59°C for 45 s, and 72°C for 60 s. Three PCR replicates were

performed for all 131 DNA extracts, controls, and both primer sets.

PCR products were visualized on 2% agarose gels (Invitrogen) with SYBR™ Safe

alongside a low-range quantitative DNA ladder. Three technical replicates (10µL each) were

pooled per DNA extract. Amplicon pools were purified using NucleoMag NGS Clean-up and

Size Select beads (Mackerey-Nagel) at a 1.6× bead-to-amplicon ratio. Nextera XT (Illumina)

indices were ligated in a second PCR (20µL), consisting of 2µL amplicon pool, 10µL

TaqMan™ Environmental Master Mix 2.0, 1.5µL of both N- and S-indices, and 5µL RNase-

free water. The thermal profile was 95°C for 10 min, 10 cycles of 95°C for 15 s, 55°C for 30

s, 72°C for 1 min, and a final extension at 72°C for 5 min.

Amplicon pool concentrations were measured using the 4200 Tapestation System

(Agilent Technologies) and pooled equimolarly using the OT-2 Robot (Opentrons). The final

pool was purified using a 1.2× bead-to-amplicon ratio and quantified on the 4200 Tapesta-

tion. Libraries were sequenced at Macrogen Europe (Netherlands) on an Illumina MiSeq (v3

chemistry, 150bp paired-end), targeting 50,000 reads per sample.

Data preparation

Quality assurance was performed separately for each primer set. Illumina adaptors and

primers were trimmed using Cutadapt v4.9 [40], and reads were processed into amplicon se-

quence variants (ASVs) using DADA2 [41] in R [42]. Filtering parameters included maxEE =

2 and minLen = 50, removing reads with >2 expected errors or <50 bp. Dereplicated and er-

ror estimation followed, using DADA2 to calculate true ASV counts. We merged reads, re-

moved chimeras, retaining only ASVs between 80 -106 bp for 12S, and 83 - 118 bp for 16S.

Taxonomic identification was performed using BLASTn against GenBank 12S and

16S databases via a custom pipeline on Naturalis Biodiversity Center's OpenStack GALAXY

platform [43]. A weighted LCA algorithm applied cutoffs of 85% query coverage, 98% iden-

tity, and a minimum score of 170 [44]. ASVs identified as human, unidentified, or detected in

controls were removed. To minimize contamination, negative control read counts were sub-

tracted from DNAir sampler reads for their respective weeks and from all reads for Coriolis

Micro and Pollensniffer. ASVs of the same species were grouped for both primer sets.

Afterward, we manually validated all species-level identifications and removed ASVs

that were identified as human, could not be identified, and/or were detected in the positive

and negative PCR controls. To control for potential contamination, we removed the number

of reads present in our negative controls from the read counts in their respective weeks for

the DNAir sampler, and from all read counts for both the Coriolis Micro and Pollensniffer All

ASVs belonging to the same species were grouped together for both primer sets.

Read counts from both primers were combined into an ASV table summarizing total

species read counts per sample. Separate datasets were created for each eDNA sampler,

summing species-specific reads across all samples for each method. Identified species were

cross-referenced with a detailed Rotterdam Zoo inventory (N=266), including those housed

off-exhibit.

Quantification and statistical analyses

We assessed normality using Shapiro-Wilk and KS tests in R 4.4.0, finding all data normal

except zoo species sequencing read and total biomass (p>0.05), which were log-

transformed. To compare species richness detected by the three airborne eDNA samplers,

we performed ANOVA and Tukey's HSD tests.

Species richness accumulation over time was visualized using *vegan* v.2.6.6.1 and

ggplot2 [45,46]. DNAir samples were analyzed across 75 outdoor sites (three replicate cam-

paigns in five locations, each with five samplers collected at 6, 24, 48, 72, and 96 hours).

Two accumulation curves were generated: one for all species and one for listed zoo species.

These were averaged per campaign and modeled using a Generalized Linear Mixed Model

using time as a fixed effect and location as a random effect. For active samplers (Pollensnif-

fer and Coriolis Micro), total and average detected species were plotted per replication. We

report static values instead of curves for the active samplers since the active samplers were

only used once for each location/replication.

Zoo species detection distances were estimated using Google Earth, calculating the

mean detection distance per sampling method. Using these radii, we determined the per-

centage of expected zoo species detected per sampler over three weeks. Maximum detec-

tion distances were also averaged, omitting zeros to prevent overdispersion.

To examine the relationship between species biomass and detection probability

(sensu [22,24], we multiplied published mean biomass values (Table S4) by the number of

individuals housed in the zoo. A Pearson correlation tested the relationship between total

biomass and read counts.

RESULTS

Sequencing results

To detect airborne eDNA from vertebrates and mammals, we targeted mitochondrial regions using 12S and 16S markers. For the 12S dataset, we reduced our original 3,862,876 sequencing reads to 783,731 reads across 293 ASVs after filtering, removing unidentifiable ASVs and correcting for negative controls. Subsequently, questionable species identifications were adjusted using the weighted LCA algorithm in Galaxy. Despite our use of human blocking primers, 299,733 reads were identified as human (*Homo sapiens*), which we subsequently removed. This resulted in 483,998 reads across 161 ASVs, which we identified to 65 taxa,15 of which were expected resident species of the Rotterdam Zoo (total read count of 48,735 across 26 ASVs).

For 16S, we filtered 5,871,156 sequencing reads down to 651,690 reads after filtering, removing unidentifiable ASVs and correcting for negative controls. Of the remaining reads, 293,267 were identified as *Homo sapiens*. After removal, the 16S dataset consisted of 358,249 reads across 92 ASVs that we identified to 47 taxa. Of these taxa, 23 were not identified using the 12S primer set, and 15 ASVs were identified as expected resident species of Rotterdam Zoo (total read count of 146,019 across 23 ASVs).

Cumulatively, we detected airborne eDNA from 88 distinct species, of which 24 were found in both primer sets (Table S2). Of these 24 species, six were expected zoo species: the Bactrian camel (*Camelus bactrianus*), the black rhinoceros (*Diceros bicornis*), the giraffe (*Giraffa camelopardalis*), the European otter (*Lutra lutra*), the Indian rhinoceros (*Rhinoceros unicornis*), and the Arctic fox (*Vulpes lagopus*).

Species detection

Across all sampling weeks at the Rotterdam Zoo, we detected a total of 88 species: 34 mammals, 29 birds, 15 fish, and 10 amphibians (Figure 2). The passive DNAir Sampler accounted for 62.17% of all reads and detected 76 species, whereas the active samplers, the

Pollensniffer and Coriolis Micro, detected 50 species (21.66% of the reads) and 25 species

(16.17% of reads), respectively. Of the species diversity detected, the majority (30) were ex-

clusively collected by the DNAir Sampler, and 17 were detected by all three samplers (Figure

2a). This pattern was similar when looking specifically at zoo residents (Fig. 3a). Overall, 24

of the total 88 species detected using these eDNA samplers were listed as Rotterdam Zoo

residents: 15 mammals and 9 birds (Figure 3), accounting for 9% of the total inventory at

Rotterdam Zoo (including those not on exhibit). Of these zoo species, the DNAir Sampler de-

tected 20 species, 9 of which were exclusive to this passive sampler (Fig. 2a; Fig. 2b),

whereas the Pollensniffer collected 14 zoo species, of which 4 were exclusive to the Pol-

lensniffer, and the Coriolis Micro detected 4 zoo species.

a)

b)

FIGURE 2 a) Venn diagram of total species detection across different air eDNA samplers. b) Non-

target species detected (icons per family) across three weeks sampling in Rotterdam Zoo using differ-

ent air eDNA samplers: DNAir Sampler (passive sampler), Coriolis Micro, and Pollensniffer. Asterisk

denotes Netherlands non-native species. Total sequencing reads across all samples per sampler indi-

cated for each species by the corresponding circle colour (Legend bottom right).

a)

296 b)

FIGURE 3 a) Venn diagram of total species detection distribution across different air eDNA sam-

plers b) Rotterdam Zoo resident species (icons per family) detected across three weeks of airborne

eDNA sampling using different samplers: DNAir Sampler (passive sampler), Coriolis Micro, and Pol-

lensniffer. total sequencing reads across all samples per sampler indicated for each species by the

corresponding circle colour (Legend bottom right).

While the majority of the non-target species (N=64) were those native to the Nether-

lands, including mammals, birds, amphibians and fish, 10 non-native species were also iden-

tified (Fig. 2b). At the indoor sampling site, four species were detected that were not detected

outdoors.

Accumulating species richness detection over time

Comparing the three air eDNA samplers, the average total detected species richness by the

DNAir Sampler per outdoor sampling location exceeded that of the Coriolis Micro (100L/min)

after only 6 hours (Fig. 4a), whereas the Pollensniffer at 8L/min was exceeded by the DNAir

Sampler between 24 and 48 hours of passive sampling. The DNAir Sampler detected on av-

erage 15.07 (SD ± 2.62) species per outdoor site after 96 hours of sampling (Fig. 4a), with

site 2 detecting the most species diversity, on average 19 species (SD ± 5.29), and site 3 de-

tecting the least number of species (11.67 SD ± 3.21), after 96 hours. Comparatively, the

Pollensniffer detected an average of 6.6 species (SD ± 1.38), whereas the Coriolis Micro de-

tected an average of 3.47 species (SD ± 1.15) per site after 10 minutes of active sampling

Excluding all non-target species, we detected an average of 2.87 zoo species (SD ±

1.57) per outdoor sampling site after 96 hours of sampling using the DNAir Sampler. By

comparison, the Pollensniffer and the Coriolis Micro detected an average of 1.67 (SD ± 0.82)

and 0.4 (SD ± 0.43) zoo species at each outdoor sampling site, respectively. The average

detected zoo species richness by the Coriolis Micro was exceeded after 24 hours of sam-

pling using the DNAir Sampler, and after approximately 72 hours using the Pollensniffer.

Our indoor sampling site (location 6) demonstrated the mean total species richness

and zoo species richness obtained by the Coriolis Micro were both exceeded by the DNAir

Sampler within the first 24 hours, similar to the outdoor sampling locations (Figure S2). In

contrast, the DNAir Samplers only exceeded the species richness obtained by the Pollensnif-

fer after 72 to 96 hours (Figure S2).

FIGURE 4 Species richness accumulation curves for outdoor sampling locations only. Grey dotted

lines represent species richness accumulation curves for the outdoor sampling locations (sites 1-5) as

detected by the DNAir Sampler, black line is a GLMM of best fit. (a) accumulation of the total detected

species richness by the DNAir Sampler for the outdoor sampling locations over 96 hours across the

three weeks plotted against the mean total detected species richness by the Coriolis Micro and Pol-

lensniffer for the outdoor sampling locations across the three weeks, with (b) showing only zoo resi-

dent species.

The total detected species richness by the DNAir Sampler per outdoor sampling location ex-

ceeded the mean total detected species richness by the Coriolis Micro within the first 24

hours, while it exceeded the mean total detected species richness by the Pollensniffer be-

tween 24 and 48 hours of sampling, with the exception of location 3. Our results indicate the

DNAir Sampler detected significantly higher total species richness, followed by the Pol-

lensniffer, and the Coriolis Micro detected the least ($F_{(2,12)} = 43.21$, $p < 0.001$) (Figure S3a).

The Coriolis Micro detected a total of 0.8 zoo species on average across three measure-

ments per outdoor sampling location (SD ± 0.84), whereas the Pollensniffer detected a total

of 3.6 (SD ± 1.67) zoo species on average across three measurements per outdoor sampling

location.

We also found total detected zoo species per outdoor sampling location to be signifi-

cantly different among all three samplers ($F_{(2,12)} = 6.80$, $p = 0.011$), with the highest total

species richness detected by the DNAir Sampler (Figure S3b).

Accuracy, dispersion, and biomass

The DNAir Sampler detected zoo species at the greatest average distance (188.67 ± 61.33

m), followed by the Pollensniffer (172.58 ± 91.29 m), and the Coriolis Micro (68 ± 50.21 m),

though these differences were not statistically significant ($F_{(2,10)} = 2.8$, $p = 0.11$) (Figure 5a;

Table S3). Excluding the indoor site, the DNAir Sampler also recorded the furthest single de-

tection (373 ± 124.37 m), significantly outperforming the Coriolis Micro ($p = 0.036$), with a

maximum detection of an American bison (*Bison bison*) at 515 m (location 2) (Figure 5b).

The furthest detections by the Pollensniffer and Coriolis Micro were an Arctic fox (*Vulpes*

lagopus, 491 m) and a black-crowned night heron (*Nycticorax nycticorax*, 128 m), both at the

indoor location (location 6). Outdoors, the farthest detections were the Asian elephant (*Ele-*

phas maximus, 466 m, Pollensniffer) and the night heron again (116 m, Coriolis Micro).

FIGURE 5 Boxplots showing the average (a) and maximum (b) distance in meters between the de-

tected zoo species and the sampling location at which they were detected for the different outdoor

sampling locations per air eDNA sampler across three weeks of sampling.

In terms of detection accuracy, the DNAir Sampler was highest, identifying 19.11%

(±10.87%) of zoo species within its average range, followed by the Pollensniffer (12.25% ±

5.49%) and the Coriolis Micro (10% ± 13.69%). Using the lowest average detection distance

(68 m, Coriolis Micro) as a baseline, both the DNAir Sampler (23.71% ± 22.94%) and the

Pollensniffer (21.86% ± 18.31%) showed slightly higher detection accuracy across five out-

door locations, indicating effective identification of nearby zoo species (Fig. 6). Additionally,

there was a significant positive correlation between total sequencing reads and the log-

transformed biomass of detected zoo species ($r(24) = 0.44$, $p = 0.031$; Fig. 7). This trend re-

mained, though weaker, after removing four large outliers ($r(20) = 0.38$, $p = 0.097$; Fig. S4).

 FIGURE 6 Mean % accuracy dispersion curve across distance (meters ± SD) from sampling loca-

 tion. Sampler type is denoted by colour (legend to the right).

396 Correlation between total sequencing reads (read counts) and Log transformed total

397 biomass (kg) of detected zoo species.

399 DISCUSSION

This study aimed to assess airborne eDNA sampling strategies by comparing the effi-

ciency of a newly designed passive approach and two commonly used active methods, while

exploring the spatial and temporal dynamics of airborne eDNA. Our results show that the

DNAir Sampler detected a wide range of zoo and non-target species and was more effective

at capturing vertebrate biodiversity over 96 hours than either active sampler. Although mean

detection distances did not differ significantly, the DNAir Sampler identified taxa at the fur-

thest range and achieved the highest accuracy within its average detection radius. Addition-

ally, we found a positive correlation between species biomass and total sequencing reads

obtained via airborne eDNA.

410 Species detection within Rotterdam Zoo

We detected airborne eDNA from 88 vertebrate species, spanning both terrestrial (mammals,

birds) and aquatic (fish, amphibians) habitats. Of these, 76 species were detected by the

passive DNAir Sampler within 96 hours, including 30 not detected by either active sampler—

highlighting the superior detection capacity of passive sampling. While the Pollensniffer

showed similar diversity to previous active air eDNA studies in zoos [22,24], 64 of the 88 taxa

were non-target species, likely local wildlife or food (e.g., livestock, fish).

Currently, little is known about the movement of airborne eDNA, and the impact of

sampling strategies. While some species were detected across multiple samples and sam-

pling sites (e.g. *Cyprinus caprio* in 58 samples; *Elephas maximus* in 28 samples), 32 species

were present in only a single sample, highlighting the likely stochastic nature in detecting air-

borne eDNA in general. The greater detection capacity of the DNAir Sampler may stem from

factors like sampling height (~2.5 m vs. ≤1 m for active samplers), extended sampling dura-

tion (96 hours vs. short active bursts), and a higher cumulative air volume. This longer, pas-

sive sampling may integrate effects tied to eDNA release variability due to species behavior

or environmental factors (e.g., wind, rain) increase species detection over other methods.

At the indoor site (location 6), however, active sampling by the Pollensniffer outper-

formed passive methods possibly due to limited airflow indoors which hampers passive col-

lection but promotes airborne eDNA accumulation [7, 47]. In contrast, the Coriolis Micro de-

tected the lowest diversity despite its higher flow rate. Notably, the Pollensniffer consistently

detected more species per location (Fig. S3a), despite sampling less air per minute. This

may reflect differences in sampling mechanics—its impingement method might better cap-

ture fine aerosols than the vortex mechanism of the Coriolis Micro, where high airflow could

reduce DNA particle retention. Further studies using matched air volumes but varying flow

rates are needed to test this hypothesis.

Accumulation patterns of airborne eDNA

The DNAir Sampler outperformed the Coriolis Micro in species richness within the first 24

hours, both overall and for zoo species. Compared to the Pollensniffer, it surpassed species

richness after 24–48 hours (overall) and 48–72 hours (zoo species). Given that active sam-

plers are noisy, labour-intensive, costly, and require trained personnel, passive sampling—by

simply hanging a device and retrieving it after a few days—offers a low-cost, minimally intru-

sive alternative.

Consistent with [29], who showed airborne eDNA accumulates over 24 hours using

water-based passive samplers, we observed no plateau in species detection even after 96

hours. Continuous detection of new species suggests that longer sampling durations could

further increase richness. Unlike aquatic eDNA, which may be more homogenous, airborne

eDNA might be released in pulses or be influenced by weather, as suggested by [27] and

[48]. We observed variation in both species richness and composition over time, highlighting

the potential impact of pulsed release, weather, and eDNA persistence on airborne eDNA

signatures. Untangling these effects experimentally and devising optimized sampling strate-

gies is a challenge we look forward to addressing.

Spatial distribution of target species

Out of all airborne eDNA samplers, the DNAir Sampler had the highest average detection

distance and accuracy, identifying nearly 1 in 5 target species within its 188 m range. Both

the DNAir Sampler and Pollensniffer were more effective at detecting nearby species than

the Coriolis Micro, aligning with findings from [22] and [24], who observed stronger detection

closer to source animals. While airborne eDNA typically concentrates around its origin, it can

also travel considerable distances—such as the American bison (*Bison bison*), detected

515m from its enclosure by the DNAir Sampler within 24 hours. Notably, the furthest detec-

tions by the active samplers occurred indoors, underscoring the challenge of tracing eDNA

back to its source in natural settings [7]. Similar difficulties have occasionally been reported

in aquatic systems for localizing sources of eDNA (e.g., [49]), highlighting challenges regard-

less of the environmental media sampled.

Conversely, many nearby target species went undetected, with contributing factors

likely being heterogeneity, wind patterns, UV degradation, or low species-specific shedding

rates [7, 50]. Echoing findings from aquatic studies [51], zoo-based airborne eDNA research

has shown a correlation between species biomass and detection likelihood [22,24]. Our re-

sults similarly revealed a strong positive relationship between species biomass and sequenc-

ing reads, corroborating what is becoming a salient characteristic of eDNA.

Future directions and recommendations

This study is the first to apply the passive DNAir Sampler for airborne eDNA-based verte-

brate detection, though its optimal setup remains untested. We used 0.7 μm glass fiber fil-

ters, but other materials may enhance detection. For instance, [24] used F8 pleated filters,

and [22] used Sterivex-HV filters (0.22–0.45 µm), finding that pore size did not significantly

affect community composition. Since the DNAir Sampler passively collects particles without

forcing air through pores, pore size may be less critical than with active samplers.

Several variables remain unexplored—filtering duration, seasonal effects, sampler

height, and air movement. Wind direction, in particular, is thought to influence detection more

during passive sampling [52], though we did not track it in this study. Despite this, the DNAir

Sampler effectively detected both zoo and local species and continued to register new detec-

tions over time.

Overall, the DNAir Sampler offers a low-cost, non-invasive, and efficient method for

vertebrate biodiversity assessment compared to its active sampler counterparts. Its ease of

deployment and wide detection range position it a valuable tool, especially for expanding ter-

restrial biomonitoring. Future research should refine passive sampling protocols and evaluate

their performance in natural habitats. While terrestrial eDNA research lags behind aquatic

systems, airborne eDNA has transformative potential for large-scale biodiversity monitoring.

Given its strong detection performance across taxa, passive air eDNA is an underappreci-

ated yet powerful method for biodiversity assessments and may emerge as the preferred

strategy for future terrestrial biomonitoring, particularly for scaling up efforts.

Resource availability

493   Upon acceptance, all data will be uploaded to Figshare and raw sequence files will

494   be made available upon request.

Acknowledgements

496   This research was made possible through the Aspasia Grant 0.15.021.046 (K.A.S.),

497   as well as the Impact Fund through the Institute of Environmental Sciences at Leiden

498   University (K.A.S & K.B.T). We would also like to thank Blijdorp Zoo for access to

499 their facilities, for enabling feedback and assistance during the sampling campaigns,

500 as well as access to their resident zoo species records.

Author contributions

K.A.S., K.T., A.G. C. L. S. and H.J. conceived and planned the experiments, JM.L.

built the samplers. H.J. and JM.L. conducted sample collections. H.J. and K.A.S. car-

ried out the laboratory work. H.J., K.A.S. and K.T. planned and carried out the data

analysis. All authors contributed to the interpretation of the results. H.J. and K.A.S.

took the lead in writing the manuscript but all authors provided feedback, edited, and

helped shape the research, analysis and manuscript.

Declaration of interest

509 The authors declare no competing interests.

REFERENCES

1. Tong, S.; Bambrick, H.; Beggs, P. J.; Chen, L.; Hu, Y.; Ma, W.; Steffen, W.; Tan, J.
Current and Future Threats to Human Health in the Anthropocene. Environ. Int.
2022, 158, 106892. https://doi.org/10.1016/j.envint.2021.106892.
2. Zimmerer, K. S.; de Haan, S.; Jones, A. D.; Creed-Kanashiro, H.; Tello, M.;
Carrasco, M.; Meza, K.; Plasencia Amaya, F.; Cruz-Garcia, G. S.; Tubbeh, R.; et
al. The Biodiversity of Food and Agriculture (Agrobiodiversity) in the Anthropo-
cene: Research Advances and Conceptual Framework. Anthropocene 2019, 25,
100192. https://doi.org/10.1016/j.ancene.2019.100192.
3. Baird, D. J.; Hajibabaei, M. Biomonitoring 2.0: A New Paradigm in Ecosystem As-
sessment Made Possible by Next-Generation DNA Sequencing. Mol. Ecol. 2012,
21(8), 2039–2044. https://doi.org/10.1111/j.1365-294X.2012.05519.x.
4. Rodríguez-Ezpeleta, N., Zinger, L., Kinziger, A., Bik, H. M., Bonin, A., Coissac, E.,
Emerson, B. C., Lopes, C. M., Pelletier, T. A., Taberlet, P., & Narum, S. (2021).
Biodiversity monitoring using environmental DNA. *Molecular Ecology Resources*,
21(5), 1405–1409. https://doi.org/10.1111/1755-0998.13399
5. Taberlet, P.; Coissac, E.; Pompanon, F.; Brochmann, C.; Willerslev, E. Towards
Next-Generation Biodiversity Assessment Using DNA Metabarcoding. Mol. Ecol.
2012, 21(8), 2045–2050. https://doi.org/10.1111/j.1365-294X.2012.05470.x.
6. Didaskalou, E. A.; Trimbos, K. B.; Stewart, K. A. Environmental DNA. Curr. Biol.
2022, 32(22), R1250–R1252. https://doi.org/10.1016/j.cub.2022.09.052.
7. Garrett, N. R., Watkins, J., Francis, C. M., Simmons, N. B., Ivanova, N., Naaum,
A., Briscoe, A., Drinkwater, R., & Clare, E. L. (2023). Out of thin air: surveying

tropical bat roosts through air sampling of eDNA. *PeerJ*, *11*, e14772.
https://doi.org/10.7717/peerj.14772

8. Kelly, R. P., Port, J. A., Yamahara, K. M., Martone, R. G., Lowell, N., Thomsen, P.
F., Mach, M. E., Bennett, M., Prahler, E., Caldwell, M. R., & Crowder, L. B. (2014).
Harnessing DNA to improve environmental management. *Science*, *344*(6191),
1455–1456. https://doi.org/10.1126/science.1251156

9. Qu, C., & Stewart, K. A. (2019). Evaluating monitoring options for conservation:
comparing traditional and environmental DNA tools for a critically endangered
mammal. *The Science of Nature*, *106*(3–4), 9. https://doi.org/10.1007/s00114-
019-1605-1

10. Hallam, J., Clare, E. L., Jones, J. I., & Day, J. J. (2021). Biodiversity assessment
across a dynamic riverine system: A comparison of eDNA metabarcoding versus
traditional fish surveying methods. *Environmental DNA*, *3*(6), 1247–1266.
https://doi.org/10.1002/edn3.241

11. Fediajeviate, J., Priestly, V., Arnold, R., Savolainen, V. 2021. Meta-analysis shows
that environmental DNA outperforms traditional surveys but warrants better re-
porting standards. Ecology and Evolution. 11(9), 4803-4815.
https://doi.org/10.1002/ece3.7382

12. Ruppert, K. M., Kline, R. J., & Rahman, M. S. Past, present, and future perspec-
tives of environmental DNA (eDNA) metabarcoding: A systematic review in meth-
ods, monitoring, and applications of global eDNA. Global Ecology and Conserva-
tion (2019), *17*, e00547. https://doi.org/10.1016/j.gecco.2019.e00547

13. Gold, Z., Sprague, J., Kushner, D. J., Zerecero Marin, E., & Barber, P. H. (2021).
eDNA metabarcoding as a biomonitoring tool for marine protected areas. *PLOS*
ONE, *16*(2), e0238557. https://doi.org/10.1371/journal.pone.0238557

14. Thomsen, P. F.; Kielgast, J.; Iversen, L. L.; Møller, P. R.; Rasmussen, M.; Willer-
slev, E. Detection of a Diverse Marine Fish Fauna Using Environmental DNA from
Seawater Samples. PLoS ONE 2012, 7(8), e41732.
https://doi.org/10.1371/journal.pone.0041732.

15. Beng, K. C.; Corlett, R. T. Applications of Environmental DNA (eDNA) in Ecology
and Conservation: Opportunities, Challenges, and Prospects. Biodivers. Conserv.
2020, 29, 2089–2121. https://doi.org/10.1007/s10531-020-01980-0.

16. Kinoshita, G., Yonezawa, S., Murakami, S., & Isagi, Y. (2019). Environmental DNA
Collected from Snow Tracks is Useful for Identification of Mammalian Species.
Zoological Science, *36*(3), 198. https://doi.org/10.2108/zs180172

17. Lyman, J. A., Sanchez, D. E., Hershauer, S. N., Sobek, C. J., Chambers, C. L.,
Zahratka, J., & Walker, F. M. (2022). Mammalian eDNA on herbaceous vegeta-
tion? Validating a qPCR assay for detection of an endangered rodent. *Environ-*
mental DNA, *4*(5), 1187–1197. https://doi.org/10.1002/edn3.331

18. Thomsen, P. F.; Sigsgaard, E. E. Environmental DNA Metabarcoding of Wild
Flowers Reveals Diverse Communities of Terrestrial Arthropods. Ecol. Evol. 2019,
9(4), 1665–1679. https://doi.org/10.1002/ece3.4809.

19. Sigsgaard, E. E., Olsen, K., Hansen, M. D. D., Hansen, O. L. P., Høye, T. T.,
Svenning, J., & Thomsen, P. F. Environmental DNA metabarcoding of cow dung
reveals taxonomic and functional diversity of invertebrate assemblages. *Molecular*
Ecology (2021), *30*(13), 3374–3389. https://doi.org/10.1111/mec.15734

20. Roger, F., Ghanavi, H. R., Danielsson, N., Wahlberg, N., Löndahl, J., Pettersson,
L. B., Andersson, G. K. S., Boke Olén, N., & Clough, Y. (2022). Airborne environ-
mental DNA metabarcoding for the monitoring of terrestrial insects—A proof of

concept from the field. Environmental DNA 2014, 4(4), 790–807. https://doi.org/10.1002/edn3.290

21. Clare, E. L., Economou, C. K., Faulkes, C. G., Gilbert, J. D., Bennett, F., Drinkwater, R., & Littlefair, J. E. (2021). eDNAir: proof of concept that animal DNA can be collected from air sampling. *PeerJ*, *9*, e11030. https://doi.org/10.7717/peerj.11030

22. Clare, E. L., Economou, C. K., Bennett, F. J., Dyer, C. E., Adams, K., McRobie, B., Drinkwater, R., & Littlefair, J. E. (2022). Measuring biodiversity from DNA in the air. *Current Biology*, *32*(3), 693-700.e5. https://doi.org/10.1016/j.cub.2021.11.064

23. Gusareva, E. S., Acerbi, E., Lau, K. J. X., Luhung, I., Premkrishnan, B. N. V., Kolundžija, S., Purbojati, R. W., Wong, A., Houghton, J. N. I., Miller, D., Gaultier, N. E., Heinle, C. E., Clare, M. E., Vettath, V. K., Kee, C., Lim, S. B. Y., Chénard, C., Phung, W. J., Kushwaha, K. K., … Schuster, S. C. (2019). Microbial communities in the tropical air ecosystem follow a precise diel cycle. *Proceedings of the National Academy of Sciences of the United States of America*, **116**(46), 23299–23308. https://doi.org/10.1073/pnas.1908493116

24. Lynggaard, C., Bertelsen, M. F., Jensen, C. V., Johnson, M. S., Frøslev, T. G., Olsen, M. T., & Bohmann, K. (2022). Airborne environmental DNA for terrestrial vertebrate community monitoring. *Current Biology*, *32*(3), 701-707.e5. https://doi.org/10.1016/j.cub.2021.12.014

25. Polling, M., Buij, R., Laros, I., & de Groot, A. (2024). Continuous daily sampling of airborne eDNA detects all vertebrate species identified by camera traps. *Environmental DNA*, *6*(4), Article e591. https://doi.org/10.1002/edn3.591

26. Garrett, N. R., Watkins, J., Simmons, N. B., Fenton, B., Maeda-Obregon, A., Sanchez, D. E., Froehlich, E. M., Walker, F. M., Littlefair, J. E., & Clare, E. L. (2022). Airborne eDNA documents a diverse and ecologically complex tropical bat and other mammal community. *Environmental DNA*, *5*(2), 350–362. https://doi.org/10.1002/edn3.385

27. Johnson, M. D.; Barnes, M. A.; Garrett, N. R.; Clare, E. L. Answers Blowing in the Wind: Detection of Birds, Mammals, and Amphibians with Airborne Environmental DNA in a Natural Environment Over a Yearlong Survey. Environ. DNA (2023), 5(2), 375–387. https://doi.org/10.1002/edn3.388.

28. Johnson, M. D., Fokar, M., Cox, R. D., & Barnes, M. A. (2021). Airborne environmental DNA metabarcoding detects more diversity, with less sampling effort, than a traditional plant community survey. *BMC Ecology and Evolution*, *21*(1), 218. https://doi.org/10.1186/s12862-021-01947-x

29. Klepke, M. J., Sigsgaard, E. E., Jensen, M. R., Olsen, K., & Thomsen, P. F. (2022). Accumulation and diversity of airborne, eukaryotic environmental DNA. *Environmental DNA*, *4*(6), 1323–1339. https://doi.org/10.1002/edn3.340

30. Lynggaard, C., Guldberg Frøslev, T., Johnson, M.S., Tange Olsen, M., Bohmann K. 2023. Airborne environmental DNA captures terrestrial vertebrate diversity in nature. *Molecular Ecology Resources* 24(1). E13840. https://doi.org/10.1111/1755-0998.13840

31. Johnson M, Barnes MA. (2023). Macrobial airborne environmental DNA analysis: A review of progress, challenges, and recommendations for an emerging application. Mol Ecol Resour. 2024 Oct;24(7):e13998. doi: 10.1111/1755-0998.13998.

32. de Weger, L. A.; Molster, F.; de Raat, K.; den Haan, J.; Romein, J.; van Leeuwen, W.; de Groot, H.; Mostert, M.; Hiemstra, P. S. A New Portable Sampler to Monitor Pollen at Street Level in the Environment of Patients. Sci. Total Environ. 2020, 741, 140404. https://doi.org/10.1016/j.scitotenv.2020.140404.

33. Bertin Technologies SAS. Coriolis Micro – Microbial Air Sampler. Available at: https://www.bertin-technologies.com/product/air-samplers/coriolis-micro-air-sampler/ (accessed March 2, 2025).

34. Luursema, J. M. (2023). DNAir-sampler. Available at: https://github.com/J4n-M44rt3n/DNAir-sampler [accessed March 2, 2025].

35. Yates, M. C.; Glaser, D. M.; Post, J. R.; Cristescu, M. E.; Fraser, D. J.; Derry, A. M. The Relationship between eDNA Particle Concentration and Organism Abundance in Nature is Strengthened by Allometric Scaling. Mol. Ecol. 2021, 30(13), 3068–3082. https://doi.org/10.1111/mec.15543.

36. Taylor, P. G. Reproducibility of Ancient DNA Sequences from Extinct Pleistocene Fauna. Mol. Biol. Evol. 1996, 13(1), 283–285. https://doi.org/10.1093/oxfordjournals.molbev.a025566.

37. Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P., & Coissac, E. (2011). ecoPrimers: inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic Acids Research*, *39*(21), e145–e145. https://doi.org/10.1093/nar/gkr732

38. Vestheim, H.; Jarman, S. N. Blocking Primers to Enhance PCR Amplification of Rare Sequences in Mixed Samples – A Case Study on Prey DNA in Antarctic Krill Stomachs. Front. Zool. 2008, 5(1), 12. https://doi.org/10.1186/1742-9994-5-12.

39. Calvignac-Spencer, S., Merkel, K., Kutzner, N., Kühl, H., Boesch, C., Kappeler, P. M., Metzger, S., Schubert, G., & Leendertz, F. H. (2013). Carrion fly-derived DNA as a tool for comprehensive and cost-effective assessment of mammalian biodiversity. *Molecular Ecology*, *22*(4), 915–924. https://doi.org/10.1111/mec.12183

40. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.Journal*, *17*(1), 10. https://doi.org/10.14806/ej.17.1.200

41. Callahan, B. J.; McMurdie, P. J.; Rosen, M. J.; Han, A. W.; Johnson, A. J. A.; Holmes, S. P. DADA2: High-Resolution Sample Inference from Illumina Amplicon Data. Nat. Methods 2016, 13(7), 581–583. https://doi.org/10.1038/nmeth.3869.

42. RStudio Team (2020). RStudio: integrated development for R. http://www.rstudio.com/.

43. Afgan, E.; Baker, D.; Batut, B.; van den Beek, M.; Bouvier, D.; Čech, M.; Chilton, J.; Clements, D.; Coraor, N.; Grüning, B. A.; Guerler, A.; Hillman-Jackson, J.; Hiltemann, S.; Jalili, V.; Rasche, H.; Soranzo, N.; Goecks, J.; Taylor, J.; Nekrutenko, A.; Blankenberg, D. The Galaxy Platform for Accessible, Reproducible, and Collaborative Biomedical Analyses: 2018 Update. Nucleic Acids Res. 2018, 46(W1), W537–W544. https://doi.org/10.1093/nar/gky379.

44. Beentjes, K. K.; Barmentlo, S. H.; Cieraad, E.; Schilthuizen, M.; van der Hoorn, B. B.; Speksnijder, A. G. C. L.; et al. Environmental DNA Metabarcoding Reveals Comparable Responses to Agricultural Stressors on Different Trophic Levels of a Freshwater Community. Mol. Ecol. 2021, 31, 1430–1443. https://doi.org/10.1111/MEC.16326.

45. Oksanen, A.J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., Mcglin, D., Minchin, P.R., Hara, R. B. O., Simpson, G.L., Solymos, P., Stevens, M. H. H., & Szoecs, E. (2016). Package 'vegan' January.

46. Wickham, H. ggplot2: Elegant Graphics for Data Analysis; Springer-Verlag: New York, 2016. https://ggplot2.tidyverse.org.

47. Serrao, N. R., Weckworth, J. K., McKelvey, K. S., Dysthe, J. C., & Schwartz, M. K. Molecular genetic analysis of air, water, and soil to detect big brown bats in North

America. *Biological Conservation* (2021), *261*, 109252.
https://doi.org/10.1016/j.biocon.2021.109252

48. Bessey, C.; Gao, Y.; Truong, Y. B.; Miller, H.; Jarman, S. N.; Berry, O. Comparison of Materials for Rapid Passive Collection of Environmental DNA. Mol. Ecol. Resour. 2022, 22(7), 2559–2572. https://doi.org/10.1111/1755-0998.13640.

49. Jane, S. F., Wilcox, T. M., McKelvey, K. S., Young, M. K., Schwartz, M. K., Lowe, W. H., Letcher, B. H., & Whiteley, A. R. (2015). Distance, flow and PCR inhibition: eDNA dynamics in two headwater streams. *Molecular Ecology Resources*, *15*(1), 216–227. https://doi.org/10.1111/1755-0998.12285

50. Stewart, K. A. Understanding the Effects of Biotic and Abiotic Factors on Sources of Aquatic Environmental DNA. Biodivers. Conserv. 2019, 28(5), 983–1001. https://doi.org/10.1007/s10531-019-01709-8.

51. Takahara, T.; Minamoto, T.; Yamanaka, H.; Doi, H.; Kawabata, Z. Estimation of Fish Biomass Using Environmental DNA. PLoS ONE 2012, 7(4), e35868. https://doi.org/10.1371/journal.pone.0035868.

52. Waza, A.; Schneiders, K.; May, J.; Rodríguez, S.; Epple, B.; Kandler, K. Field Comparison of Dry Deposition Samplers for Collection of Atmospheric Mineral Dust: Results from Single-Particle Characterization. Atmos. Meas. Tech. 2019, 12(12), 6647–6665. https://doi.org/10.5194/amt-12-6647-2019.