1 **Title:** Goal-directed visual information processing with GABAergic inhibition in parietal
2 cortex
3

4 **Authors:** Zhiyan Wang, Sinah Wiborg, Antonia Wittmann, Nina Beck, Susanna Hirschle,
5 Dominik Aschenbrenner, Markus Becker, Sebastian M. Frank
6
7 **Affiliation:** Institute for Experimental Psychology, University of Regensburg, 93053
8 Regensburg, Germany.
9
10 **Correspondence:** sebastian1.frank@psychologie.uni-regensburg.de

12 **Funding:** This research was supported by the Deutsche Forschungsgemeinschaft (DFG,
13 German Research Foundation; Emmy Noether grant – project number 491290285) and the
14 Julitta und Richard Müller Stiftung.

15
16 **Declaration of Conflicting Interests:** The author declares that there is no conflict of interest.
17

Abstract

Goal-directed visual information processing involves tracking relevant visual signals (targets)
over space and time. However, goal-irrelevant visual signals (distractors) can interfere with the
tracking of targets. The neural mechanisms, which promote the tracking of targets in the
challenge of interference from distractors remain elusive. Here, we used time-resolved
functional magnetic resonance spectroscopy (fMRS) to measure concentrations of γ-
aminobutyric acid (GABA), a chief inhibitory neurotransmitter, and glutamate, a chief
excitatory neurotransmitter, in parietal and visual cortex while participants performed a visual
tracking task for targets among distractors. We found that the more targets had to be tracked,
the greater the concentrations of GABA and glutamate in parietal cortex. In visual cortex only
the concentration of glutamate increased with the number of targets. Concentration changes of
GABA and glutamate only in parietal cortex were differentially associated with tracking
performance: Better target tracking was associated with greater increase in GABA
concentration and smaller increase in glutamate concentration. The results of control
experiments showed that this differential association between tracking performance and
metabolite changes in parietal cortex reflected individual differences in distractor suppression:
participants with better distractor suppression during tracking tended to have greater increase

in GABA concentration and smaller increase in glutamate concentration. This suggests that target-distractor interference during tracking is minimized by GABAergic suppression of goal-irrelevant distractors in parietal cortex, thereby promoting goal-directed visual information processing.

Keywords: Distractor Suppression, GABA, Glutamate, Goal-Directed Information Processing, Interference, Parietal Cortex, Tracking, Visual Cortex

Introduction

Tracking visual signals relevant to current goals across space and time is critical for adaptive behavior (Carrasco, 2011; Cavanagh & Alvarez, 2005; Chun et al., 2011; Egeth & Yantis, 1997; Gazzaley & Nobre, 2012; Scholl, 2001; Van Ede & Nobre, 2023; Xu & Chun, 2009). However, in a dynamically changing environment goal-irrelevant distractor signals compete for processing and can interfere with tracking of goal-relevant target signals. How is the interference between targets and distractors resolved? Previous results showed that there are distinct neural representations of targets and distractors in visual cortex (Christophel et al., 2012; Ester et al., 2013; Harrison & Tong, 2009; Rademaker et al., 2019; Serences, 2016; Sprague et al., 2014, 2015) and parietal cortex (Bracci et al., 2017; Buschman & Miller, 2007; Vaziri-Pashkam & Xu, 2017). Especially the representation of targets in parietal cortex was found to be robust against interference from distractors (Bettencourt & Xu, 2016; Lorenc et al., 2018; Xu, 2024). However, it is unclear which neural mechanisms promote the emergence of target representations in parietal cortex that are robust to distractor interference.

A crucial neural mechanism for minimizing target-distractor interference could be inhibitory processing with γ-aminobutyric acid (GABA). GABA is a chief inhibitory neurotransmitter (Isaacson & Scanziani, 2011; Petroff, 2002; Schmidt-Wilcke et al., 2018; Tremblay et al., 2016; Tritsch et al., 2016) and critical for suppressing spontaneous activity related to distraction (Bang et al., 2018; De La Vega et al., 2014; Frangou et al., 2019; Frank et al., 2022; Shibata et al., 2017; H. R. Snyder et al., 2010; Tamaki et al., 2020; Yamada et al., 2024). While reduced GABAergic function can impair the selection of goal-relevant signals (H. R. Snyder et al., 2010), increased GABAergic activity is associated with enhanced efficiency in resolving competitions between task-relevant and task-irrelevant signals (De La Vega et al., 2014). On a cellular level, GABAergic inhibition minimizes interneuron correlations and promotes goal-relevant sensory signal processing (Cohen & Maunsell, 2009; Harris & Thiele, 2011; A. C. Snyder et al., 2016; Srivastava et al., 2023).

Here we investigated whether and, if so, how GABAergic inhibition in parietal cortex
minimizes target-distractor interference. We used a recently advanced imaging method referred
to as time-resolved functional magnetic resonance spectroscopy (fMRS) (Apšvalka et al., 2015;
Craven et al., 2024; Ip & Bridge, 2022; Koolschijn et al., 2023; Lally et al., 2014; Mullins,
2018, 2024; Stanley & Raz, 2018) to measure concentrations of GABA and Glx (corresponding
to the composite measure of glutamate and its precursor glutamine) in parietal and visual cortex
while participants performed a multiple object tracking (MOT) task (Cavanagh & Alvarez,
2005; Meyerhoff et al., 2017; Pylyshyn & Storm, 1988). In different conditions of MOT
participants tracked two or four targets among multiple distractors (henceforth referred to as
low and high load conditions), to increase target-distractor interference with increasing load.
Importantly, the total number of stimuli (sum of targets and distractors) remained identical
across conditions to maintain constant visual input (Alvarez & Franconeri, 2007; Cavanagh &
Alvarez, 2005).

We found that GABA and glutamate concentrations in parietal cortex increased in MOT
with high compared to low load. Critically, a greater increase in GABA concentration and lower
increase in glutamate concentration in parietal cortex predicted better tracking performance in
MOT with high load. In visual cortex only the concentration of glutamate increased with load
and no association with tracking performance was found.

This suggests that GABAergic inhibition in parietal cortex is crucial for tracking
performance under conditions of high target-distractor interference. However, minimizing
target-distractor interference in MOT with high load could occur through enhancing targets
(Brefczynski & DeYoe, 1999; Rorden, 2008), or suppressing distractors (Serences et al., 2004;
Wöstmann et al., 2022), or a combination of both (Föcker et al., 2023). To further elucidate the
mechanisms underlying GABAergic inhibition in parietal cortex in MOT with high load, we
conducted control experiments in which we separately varied the need to enhance targets and
suppress distractors during tracking (Bettencourt & Somers, 2009). The results of the control
experiments showed a significant association between parietal GABA and distractor
suppression, such that a greater increase in GABA concentration predicted better tracking
performance with increasing need to suppress distractors during MOT. No significant
association between parietal GABA and target enhancement was found.

Together, the results of this study suggest that GABAergic inhibition in parietal cortex
minimizes target-distractor interference through distractor suppression, which may promote the
emergence of robust target representations. This effect of GABA was found in parietal but not

in visual cortex, indicating an important role of higher order areas for goal-directed visual
information processing.
104
105 **Methods**
106
Participants
A combined total of 72 participants [49 females, 23 males, mean ± standard-error-of-
the-mean (SEM) age = 24.0 ± 0.60 years old; 70 right-handed, 2 left-handed] with normal or
corrected-to-normal vision was recruited. Participants gave informed written consent prior to
participation. The study was approved by the internal review board of the University of
Regensburg. Thirty-six participants were recruited for the main imaging experiment, among
which thirty participants completed the GABA measurement and a subset of twenty-four
participants also volunteered to complete the Glx measurement. Six additional new participants
were recruited for the Glx measurement. Twenty participants completed the control behavioral
experiments. Twenty-one participants including five participants from the main imaging
experiment were recruited for the control imaging experiments, among which nineteen
participants completed both the target enhancement and distractor suppression experiments.
One participant was only available for the target enhancement experiment. Another participant
was only available for the distractor suppression experiment.
121
MOT Task in Main Imaging Experiment
For the main imaging experiment, a MOT task with two attentional load conditions
(tracking two targets = low load; tracking four targets = high load) was used (Cavanagh &
Alvarez, 2005; Meyerhoff et al., 2017; Pylyshyn & Storm, 1988) (Figure 1). Each trial consisted
of cueing, tracking and response phases. The tracking phase was always 12 s-long. The cueing
and response phases were each 3 s-long in the GABA measurement and 2 s-long in the Glx
measurement. During the cueing phase a total of twelve disks (diameter: 0.5°) was arranged
circularly on a black screen (24 x 18° in size). Disks remained stationary during the cueing
phase. A subset of disks was highlighted in green as targets while the other disks (i.e., the
distractors) were displayed in white. During the tracking phase, the highlighting was turned off
and all disks moved independently with a speed of 4.5°/s in different and unpredictable
directions across the screen. The motion trajectory of each disk was constrained to avoid
collisions with other disks and overlap with the central fixation spot. Disks were repelled from
the borders of the screen. For each trial and session trajectories were calculated in advance and

136 randomly drawn (without replacement) from a predefined pool of unique trajectories that was

137 used for each participant. Participants were instructed to track the targets with their attention

138 and maintain central fixation. Each trial terminated with a response phase. During the response

139 phase all disks remained stationary at their final position at the end of the tracking phase and

140 one disk was highlighted in green. Participants indicated whether the highlighted disk was a

141 target or a distractor by pressing one of two buttons on the MRI-safe buttonbox. In half of all

142 trials, the highlighted disk was a target, resulting in a chance level of 50% correct response

143 accuracy. Participants were instructed to respond as quickly and accurately as possible. No

144 feedback about participants' response accuracy was provided.

146 *Main Imaging Experiment*

147 The main imaging experiment included separate fMRS runs for GABA and Glx

148 measurements in the parietal and visual cortex during MOT. A checklist with all fMRS

149 acquisition parameters, analyses and quality checks, as proposed by Choi et al. (2021), is

150 provided in Table 1. In each fMRS run participants performed MOT with low and high load

151 conditions while a time-series of fMRS scans was collected (see Figure 1). Low and high load

152 conditions were presented in random order within each fMRS run. For each load condition there

153 was a total of 18 trials in the GABA measurement and 9 trials in the Glx measurement. For

154 fMRS a mixed design was used in which each trial with MOT was followed by a jittered

155 intertrial interval. This intertrial interval ranged between 9 s, 18 s, and 27 s in the GABA

156 measurement and between 8 s, 16 s, and 24 s in the Glx measurement. During the intertrial

157 interval a central fixation spot on an otherwise blank screen was displayed. Each fMRS run

158 commenced with an on-ramp and terminated with an off-ramp. During ramps a central fixation

159 spot on an otherwise blank screen was displayed. Ramps were 60 s-long in the GABA

160 measurement and 64 s-long in the Glx measurement.

162 *MOT Tasks in Control Experiments*

163 For the control behavioral and imaging experiments, two modified versions of the MOT

164 task were used, which were based on previously proposed designs to separately manipulate

165 levels of target enhancement and distractor suppression during tracking (Bettencourt & Somers,

166 2009) (Figure 5). Using these modified designs, Bettencourt and Somers (2009) found that

167 higher levels of target enhancement and distractor suppression each led to a decrease in tracking

168 performance. Moreover, they showed that this change in behavioral performance did not result

169 from stronger crowding during tracking with high than low target enhancement and distractor

suppression (Bettencourt & Somers, 2009). All parameters in the modified versions of the MOT task including motion parameters, trial duration and number of trials with target vs. distractor highlighted in the response phase were identical with the MOT task in the main imaging experiment except for the following aspects: For the target enhancement experiment, a MOT task with two different sizes of all disks was used (large disk size = low enhancement; small disk size = high enhancement). A total of eight disks were presented in each trial. Four disks were highlighted as targets during the cueing phase. The remaining four disks were distractors. For the low and high enhancement conditions, the disk sizes were varied between 1.25° and 0.25°, respectively. For the distractor suppression experiment, a MOT task with two different numbers of distractor disks was used (low number of distractors = low suppression; high number of distractors = high suppression). For the low suppression condition, four out of eight disks were highlighted as targets during the cueing phase yielding a total of four distractors. For the high suppression condition, four out of twelve disks were highlighted as targets yielding a total of eight distractors.

Control Behavioral Experiments

Two control behavioral experiments were carried out to establish behavioral measurements of target enhancement and distractor suppression during MOT as in Bettencourt & Somers (2009). The results were also used to determine the number of participants required for the control imaging experiments. The control behavioral experiments were conducted in a single session in a psychophysics-dedicated room with room lights turned off. Participants used a chin rest and responded by pressing one of two keys on the keyboard to indicate whether the highlighted disk in the response phase was a target or a distractor. Participants completed two runs for each of the target enhancement and distractor suppression experiments using the same number of trials as in the GABA and Glx measurements in the main imaging experiment. For the control behavioral experiments, the intertrial interval was shortened to 2 s and on-/off-ramps were omitted.

Control Imaging Experiments

Two control imaging experiments were carried out to measure GABA and Glx concentrations in parietal cortex during MOT with low and high levels of target enhancement and distractor suppression, respectively. The MOT designs were identical with the control behavioral experiments. The number of trials, intertrial intervals and on-/off-ramps for GABA and Glx measurements were identical with the main imaging experiment. The results of the

control behavioral experiments were used to calculate the number of participants necessary to achieve a power of 0.95 at alpha = 0.001 (two-tailed) using a paired-sample t-test to detect a minimum effect size of interest for a difference in inverse efficiency score between high and low levels of target enhancement and distractor suppression, respectively. The results of the power analysis (using G*Power; Faul et al., 2007) showed that a total of fifteen participants was necessary to detect a minimum effect size of interest ($d = 1.65$, calculated from the control behavioral experiment) for a difference between high and low target enhancement. A total of thirteen participants was necessary to detect a minimum effect size of interest ($d = 1.85$, calculated from the control behavioral experiment) for a difference between high and low distractor suppression. Based on these results, we decided to run the same number of participants in the control imaging experiments as in the control behavioral experiments (i.e., twenty participants).

Scanning Parameters

In the main imaging experiment, single-voxel proton (^{1}H) fMRS was conducted for two types of fMRS scans (MEGAPRESS for GABA measurement and PRESS for Glx measurement) and volumes-of-interest (VOIs; located in the parietal and occipital lobes, respectively). In the control imaging experiments, MEGAPRESS and PRESS scans were collected for the parietal VOI.

MEGAPRESS scans (Mescher et al., 1996, 1998) were acquired with alternating Edit On and Edit Off scans with the following parameters: time-to-repeat (TR) = 1.5 s; time-to-echo (TE) = 68 ms; flip angle (FA) = 90°. The total number of scans (Edit On and Edit Off combined) for each MEGAPRESS fMRS run was 472. A frequency selective, single band Gauss pulse was utilized to saturate the β-CH$_2$ signal at 1.94 ppm and to refocus the J evolution of the triplet γ-CH$_2$ resonance of GABA at 3 ppm ('Edit On'). The very same Gauss pulse was used to irradiate the opposite part of the spectrum at 7.46 ppm ('Edit Off'). The 'Edit Off' spectrum was subtracted from the 'Edit On' spectrum to produce a difference spectrum. For each load condition in the main imaging experiment and for each level of target enhancement and distractor suppression in the control imaging experiments, there were a total of 18 scans for the cueing phase, 72 scans for the tracking phase, and 18 scans for the response phase. The remaining scans corresponded to the intertrial interval (216 scans) and on-/off-ramps (20 scans for each ramp). The total number of scans collected during the tracking phase in each condition was based on previous studies (Floyer-Lea et al., 2006; Frank et al., 2022), which used 64 scans to estimate GABA from a MEGAPRESS difference spectrum.

PRESS scans (Bottomley et al., 1984; Bottomley & Hardy, 1987) were acquired with

the following parameters: TR = 2 s; TE = 30 ms; FA = 90°. The total number of scans for each

PRESS fMRS run was 352. For each load condition in the main imaging experiment and for

each level of target enhancement and distractor suppression in the control imaging experiments,

there were a total of 9 scans for the cueing phase, 54 scans for the tracking phase, and 9 scans

for the response phase. The remaining scans corresponded to the intertrial interval (144 scans)

and on-/off-ramps (32 scans for each ramp).

WET water suppression was used (Ogg et al., 1994). In the main imaging experiment, a

total of four fMRS runs were conducted corresponding to MEGAPRESS for the parietal VOI,

MEGAPRESS for the visual VOI, PRESS for the parietal VOI, and PRESS for the visual VOI.

In the control imaging experiments, a total of four fMRS runs were conducted for the parietal

VOI corresponding to MEGAPRESS for target enhancement, MEGAPRESS for distractor

suppression, PRESS for target enhancement, and PRESS for distractor suppression. The parietal

VOI had a size of 2 x 2 x 4 cm and was placed approximately perpendicular to the intraparietal

sulcus. The visual VOI had a size of 2 x 2 x 2 cm and was placed perpendicular to the calcarine

sulcus. Proximity to the dura was avoided during VOI placement to minimize macromolecule

contamination. See Figure 3 for VOI locations in a representative participant. VOIs were

centered between the two hemispheres. Non-neural tissue containing lipids was avoided during

VOI placement. For the purpose of VOI placement, three high-resolution anatomical scans were

acquired for the coronal, sagittal and transverse planes prior to fMRS. These anatomical scans

had the following parameters: coronal plane: TR = 0.25 s, TE = 2.46 ms, FA = 70°, acquisition

matrix (AM) = 288 x 288, 35 slices, voxel size = 0.8 x 0.8 x 4.0 mm, inter-slice gap = 1.20 mm;

sagittal plane: TR = 0.19 s, TE = 2.46 ms, FA = 70°, AM = 288 x 288, 25 slices, voxel size =

0.8 x 0.8 x 4.0 mm, inter-slice gap = 1.20 mm; transverse plane: TR = 0.19 s, TE = 2.46 ms, FA

= 70°, AM = 288 x 288, 27 slices, voxel size = 0.8 x 0.8 x 4.0 mm, inter-slice gap = 1.20 mm.

After each fMRS run, a water reference scan without water suppression was acquired for the

same VOI using the following parameters: MEGAPRESS: TR = 1.5 s; TE = 30 ms; FA = 90°;

8 scans; PRESS: TR = 2 s; TE = 30 ms; FA = 90°; 16 scans.

Automatic shimming (gradient field adjustments to increase the homogeneity of the

magnet field B_0) was conducted prior to each fMRS run. For each fMRS run and participant the

shim values were kept below a full-width at half-maximum of 20 Hz. In the main imaging

experiment, the mean ± SEM shim values across participants were as follows: MEGAPRESS:

13.6 ± 0.26 Hz (parietal VOI) and 13.8 ± 0.18 Hz (visual VOI); PRESS: 13.6 ± 0.22 Hz (parietal

VOI) and 14.5 ± 0.26 Hz (visual VOI); water reference scan in MEGAPRESS: 13.9 ± 0.31 Hz

(parietal VOI) and 13.9 ± 0.21 Hz (visual VOI); water reference scan in PRESS: 13.7 ± 0.28

Hz (parietal VOI) and 14.5 ± 0.24 Hz (visual VOI). In the control imaging experiments, the

values for the parietal VOI were as follows: target enhancement: MEGAPRESS: 14.1 ± 0.39

Hz; PRESS: 14.0 ± 0.32 Hz; water reference scan in MEGAPRESS: 14.9 ± 0.55 Hz; water

reference scan in PRESS: 14.2 ± 0.34 Hz; distractor suppression: MEGAPRESS: 15.2 ± 0.46

Hz; PRESS: 14.8 ± 0.47 Hz; water reference scan in MEGAPRESS: 15.0 ± 0.42 Hz; water

reference scan in PRESS: 14.8 ± 0.45 Hz.

In order to calculate volume fractions for each VOI (see *Imaging Analysis* below) a high-

resolution anatomical scan of the brain was collected using a magnetization prepared rapid

gradient echo sequence with the following scanning parameters: TR= 2.3 s, TE = 2.32 ms, FA

= 8°, AM = 256 x 256, 192 sagittal slices, voxel size = 0.9 x 0.9 x 0.9 mm, interslice gap = 0.45

mm. For a subset of five participants, this scan could not be collected due to time constraints

and the sagittal scan acquired for VOI placement was used instead.

Some studies reported that the concentration of GABA is modulated by the menstrual

cycle (De Bondt et al., 2015; Duncan et al., 2014). Therefore, in female participants not using

contraception, the experiments were only conducted during the follicular phase.

Stimulus Presentation

Stimuli were generated using Psychtoolbox (Brainard, 1997; Pelli, 1997) running in

Matlab (The Mathworks, Natick, MA, USA). In the scanner, stimuli were presented using a

projector with gamma-correction onto a translucent screen located at the back of the scanner

bore. Participants viewed the screen with a headcoil-mounted mirror. Outside the scanner,

stimuli were presented on an LCD-screen with gamma-correction.

Behavioral Analysis

Response accuracy and median response time (in seconds) were combined into the

inverse efficiency score for each run and participant (Townsend & Ashby, 1978, 1983). To this

purpose median response time was divided by response accuracy (in proportion of one) and the

result was log-transformed (Bruyer & Brysbaert, 2011). The inverse efficiency score was then

combined across runs by averaging for each participant in each experiment. For correlational

analyses between fMRS and behavioral results, the inverse efficiency score of the

corresponding fMRS run was used. For one participant in the main imaging experiment,

response accuracy was at chance level in the GABA measurement. The behavioral and GABA

results from this participant were excluded from all further analyses. For one participant in the

control imaging experiments, response accuracy was at chance level in all fMRS runs. The behavioral and fMRS results from this participant were excluded from all further analyses.

Imaging Analysis

The fMRS data were analyzed using the LC Model (Provencher, 1993, 2001). Analysis steps as in previous studies were applied (Frank et al., 2023). Details are reported in Table 1 (Choi et al., 2021; Lin et al., 2021). Before running the analysis, spectra from different scans in the fMRS time-series were averaged across trials and conditions as follows. Average spectra were produced for the tracking phase in each condition for trials with correct participant response (Supplementary Figures 1-4). The mean ± SEM number of spectra used for averaging across participants in the main imaging experiment was as follows: MEGAPRESS: 69.7 ± 0.81 (parietal VOI) and 70.5 ± 0.46 (visual VOI) in low load; 63.3 ± 1.45 (parietal VOI) and 61.7 ± 1.58 (visual VOI) in high load; PRESS: 50.0 ± 1.01 (parietal VOI) and 51.4 ± 0.80 (visual VOI) in low load; 47.8 ± 1.33 (parietal VOI) and 48.2 ± 0.97 (visual VOI) in high load. The number of spectra used in the control imaging experiments was as follows: MEGAPRESS: 68.8 ± 1.25 (low enhancement) and 65.7 ± 2.08 (high enhancement); 65.5 ± 1.65 (low suppression) and 61.9 ± 2.23 (high suppression); PRESS: 50.5 ± 1.24 (low enhancement) and 47.1 ± 1.40 (high enhancement); 49.9 ± 1.66 (low suppression) and 47.7 ± 1.34 (high suppression).

The averaged MEGAPRESS difference spectra were analyzed in the chemical shift range between 1.95 and 4.2 ppm. The averaged PRESS spectra were analyzed in the chemical shift range between 0.2 and 4.0 ppm. Water scaling and eddy current correction were performed. The metabolite intensities were fitted to a linear combination of spectra of individual metabolites derived from imported metabolite basis sets (Provencher, 1993, 2001; Table 1).

The following metabolite concentrations estimated for the tracking phase in each condition and fMRS run were further analyzed: MEGAPRESS: GABA and N-acetyl-aspartate (NAA). NAA is a marker of neuronal density and mitochondrial function (Schuff et al., 2006) and is commonly used as a control metabolite for normalization due to its stability over experimental conditions and scan time (Duncan et al., 2014; Frank et al., 2023; Mullins et al., 2014; Stagg, 2014). PRESS: Glx and NAA. Prior to normalization, metabolite concentrations were corrected for each participant's volume fractions inside the fMRS VOIs. Volume fractions within the fMRS VOIs corresponding to gray matter, white matter and other types of volume including cerebrospinal fluid were calculated using the Freesurfer segmentation of each participant's reconstructed high-resolution anatomical scan of the brain (Dale et al., 1999; Fischl et al., 1999). Across participants and experiments the mean ± SEM volume fractions

were as follows: Gray matter: $61.7 \pm 0.96\%$ (parietal VOI) and $60.1 \pm 0.90\%$ (visual VOI); white matter: $15.9 \pm 0.65\%$ (parietal VOI) and $34.1 \pm 0.89\%$ (visual VOI); other types of volume: $22.4 \pm 1.17\%$ (parietal VOI) and $5.85 \pm 0.59\%$ (visual VOI). Volume fraction correction was performed using the approach proposed Kolasinski et al. (2017). The concentrations of GABA and Glx were corrected for the proportion of gray matter within each VOI by dividing by [gray matter / (gray matter + white matter + other)]. The concentration of NAA was corrected for the proportion of total brain volume within each VOI by dividing by [(gray matter + white matter) / (gray matter + white matter + other)]. Thereafter, the corrected concentration of GABA was normalized by dividing it by the corrected concentration of NAA extracted from the same MEGAPRESS difference spectrum. The corrected concentration of Glx was normalized by dividing it by the corrected concentration of NAA extracted from the same PRESS spectrum.

Post-hoc analyses were performed to calculate which brain regions were included in the gray matter of each VOI. The PALS segmentation (Van Essen, 2005) was remapped from the Freesurfer standard brain to each participant's individual brain and the percentage of Brodmann areas in the gray matter of each VOI was calculated for each participant. Across participants and experiments, approximately seventy percent of gray matter within the parietal VOI corresponded to Brodmann areas 1, 2, 3, 5, 7 and 31 (mean $\pm$ SEM: areas 1-3: $7.33 \pm 0.69\%$; area 5: $22.1 \pm 1.08\%$; area 7: $20.2 \pm 1.25\%$; area 31: $16.9 \pm 1.31\%$). The remaining gray matter corresponded mostly to the medial part of Brodmann area 4 ($32.0 \pm 2.39\%$). Within the gray matter of the visual VOI, $60.4 \pm 2.10\%$ corresponded to Brodmann area 17 and $37.5 \pm 1.95\%$ corresponded to Brodmann area 18 across participants. Gray matter corresponding to other areas was negligible. The mean $\pm$ SEM MNI coordinates of the parietal and visual VOIs across participants and experiments were as follows: Parietal VOI: $X = 0.04 \pm 0.23$, $Y = -47.9 \pm 0.76$, $Z = 52.7 \pm 0.54$; Visual VOI: $X = 0.39 \pm 0.31$, $Y = -81.4 \pm 0.42$, $Z = 4.39 \pm 0.54$.

We only included spectral results for which the signal-to-noise (SN) ratio as estimated by the LC model was higher than 5% (Wilson et al., 2019). None of a participant's spectra had to be excluded. The mean SN ratios ($\pm$ SEM) across participants' spectra in the main imaging experiment were as follows: MEGAPRESS: $24.8 \pm 1.42\%$ (parietal VOI) and $16.2 \pm 0.54\%$ (visual VOI) in low load; $23.1 \pm 1.14\%$ (parietal VOI) and $15.2 \pm 0.51\%$ (visual VOI) in high load; PRESS: $30.2 \pm 1.36\%$ (parietal VOI) and $28.7 \pm 0.83\%$ (visual VOI) in low load; $29.7 \pm 1.39\%$ (parietal VOI) and $28.1 \pm 0.85\%$ (visual VOI) in high load. The values in the control imaging experiments for the parietal VOI were as follows: MEGAPRESS: $22.3 \pm 1.84\%$ (low enhancement) and $21.5 \pm 1.91\%$ (high enhancement); $24.2 \pm 2.01\%$ (low suppression) and 24.2

± 1.97% (high suppression); PRESS: 30.5 ± 2.30% (low enhancement) and 30.4 ± 2.24% (high enhancement); 29.9 ± 2.23% (low suppression) and 29.4 ± 2.17% (high suppression).

The Cramer-Rao Lower Bounds (CRLB) indicate the reliability of quantification of GABA, Glx, and NAA. A criterion of 25% was used to reject low-quality results (following the approach in Frank et al., 2022). In the main imaging experiment, GABA results for the parietal VOI in three participants and for the visual VOI in four participants exceeded the inclusion criterion. GABA and NAA results from these participants in MEGAPRESS were excluded. Therefore, out of twenty-nine participants in MEGAPRESS, GABA and NAA results in the parietal VOI for twenty-six participants and for twenty-five participants in the visual VOI were further analyzed. None of the Glx and NAA results in any VOI in PRESS exceeded the CRLB inclusion criterion. The mean CRLB (± SEM) across included participants in the main imaging experiment were as follows: GABA (MEGAPRESS): 16.8 ± 0.58% (parietal VOI) and 19.4 ± 0.57% (visual VOI) in low load; 15.4 ± 0.42% (parietal VOI) and 20.1 ± 0.49% (visual VOI) in high load; NAA (MEGAPRESS): 3.50 ± 0.16% (parietal VOI) and 3.40 ± 0.10% (visual VOI) in low load; 3.69 ± 0.15% (parietal VOI) and 3.72 ± 0.12% (visual VOI) in high load; Glx (PRESS): 6.47 ± 0.15% (parietal VOI) and 6.73 ± 0.17% (visual VOI) in low load; 6.10 ± 0.14% (parietal VOI) and 6.70 ± 0.13% (visual VOI) in high load; NAA (PRESS): 2.60 ± 0.10% (parietal VOI) and 2.57 ± 0.11% (visual VOI) in low load; 2.67 ± 0.11% (parietal VOI) and 2.70 ± 0.11% (visual VOI) in high load.

In the control imaging experiments for the parietal VOI, GABA results from one participant exceeded the CRLB inclusion criterion. GABA and NAA results from this participant in MEGAPRESS were excluded. Therefore, out of nineteen participants in MEGAPRESS, GABA and NAA results for eighteen participants in MEGAPRESS for target enhancement and distractor suppression were further analyzed. None of the Glx and NAA results in PRESS exceeded the CRLB inclusion criterion. The mean CRLB (± SEM) across included participants in the control imaging experiments were as follows: GABA (MEGAPRESS): 16.1 ± 0.80% (low enhancement) and 16.7 ± 0.87% (high enhancement); 17.1 ± 0.71% (low suppression) and 18.3 ± 0.75% (high suppression); NAA (MEGAPRESS): 4.58 ± 0.59% (low enhancement) and 4.84 ± 0.67% (high enhancement); 3.63 ± 0.29% (low suppression) and 3.52 ± 0.22% (high suppression); Glx (PRESS): 6.68 ± 0.29% (low enhancement) and 6.68 ± 0.32% (high enhancement); 6.84 ± 0.36% (low suppression) and 6.84 ± 0.41% (high suppression); NAA (PRESS): 2.94 ± 0.18% (low enhancement) and 2.94 ± 0.16% (high enhancement); 3.05 ± 0.34% (low suppression) and 3.05 ± 0.25% (high suppression).

 Statistics

The results were analyzed using *t*-tests and Pearson correlations. The two-tailed alpha-

level was set to 0.05. Cohen's *d* and Pearson's *r* were calculated as measures of effect size for

t-test and Pearson correlation, respectively. Cohen's *d* for repeated measures was calculated

using the formula implemented in G*Power (Faul et al., 2007).

Results

Main Imaging Experiment

The behavioral results showed that the inverse efficiency score was significantly greater

in MOT with high load than with low load [$t(34) = 5.79$, $p < 0.001$, $d = 0.98$], suggesting that

participants responded slower and were less accurate with increasing tracking load (Figure 2

and Supplementary Figure 5). The fMRS results showed that the concentration of GABA was

significantly greater in MOT with high load than with low load in the parietal VOI [$t(25) =$

3.25, $p = 0.003$, $d = 0.64$] but not in the visual VOI [$t(24) = -0.84$, $p = 0.41$] (Figure 3). The

concentration of Glx was significantly greater in MOT with high load than with low load in

both VOIs [parietal VOI: $t(29) = 3.27$, $p = 0.003$, $d = 0.60$; visual VOI: $t(29) = 2.37$, $p = 0.02$,

$d = 0.43$] (Figure 3).

The fMRS and behavioral results in the parietal VOI were correlated (Figure 4): The

greater was the increase in GABA concentration from low to high load, the smaller was the

difference in inverse efficiency score between load conditions (i.e., the smaller was the decrease

in response accuracy and increase in response time with increasing tracking load) across

participants ($r = -0.47$, $p = 0.02$). A different association was found between Glx concentration

in the parietal VOI and tracking performance (Figure 4): The greater the increase in Glx

concentration from low to high load, the greater was the increase in inverse efficiency score

with load (i.e., the worse was response accuracy and the slower was response time with

increasing tracking load) across participants ($r = 0.49$, $p = 0.006$). There was no significant

correlation with tracking performance in the visual VOI (GABA: $r = -0.17$, $p = 0.43$; Glx: $r =$

0.24, $p = 0.21$) (Figure 4).

Control Behavioral Experiments

The inverse efficiency score was significantly greater in MOT with high than with low

target enhancement [$t(19) = 7.36$, $p < 0.001$, $d = 1.65$], suggesting that participants responded

less accurately and more slowly when the disk size was small during tracking (Figure 6A and

Supplementary Figure 6A,B). Similarly, the inverse efficiency score was significantly greater

in MOT with high than with low distractor suppression [$t(19) = 8.29$, $p < 0.001$, $d = 1.85$],

suggesting that participants were less accurate and responded more slowly when the number of

distractors was high during tracking (Figure 6B and Supplementary Figure 6C,D).

Control Imaging Experiments

Results for tracking performance in the control imaging experiments replicated results

in the control behavioral experiments as follows: The inverse efficiency score was significantly

greater in MOT with high than with low enhancement [$t(18) = 6.52$, $p < 0.001$, $d = 1.50$] and in

MOT with high than with low suppression [$t(18) = 3.61$, $p = 0.002$, $d = 0.83$] (Figure 7 and

Supplementary Figure 7). The fMRS results showed no significant differences in metabolite

concentrations for the between-condition comparison in the target enhancement [GABA: $t(17)$

$= 0.58$, $p = 0.57$; Glx: $t(18) = -0.69$, $p = 0.50$] and distractor suppression [GABA: $t(17) = -1.50$,

$p = 0.15$; Glx: $t(18) = -0.14$, $p = 0.89$] experiments (Figure 8). However, the fMRS and

behavioral results were correlated (Figure 9). In the target enhancement experiment, the greater

was the increase in Glx concentration from low to high enhancement, the smaller was the

difference in inverse efficiency score between the two target enhancement conditions (i.e., the

smaller was the decrease in response accuracy and increase in response time from low to high

enhancement) across participants ($r = -0.49$, $p = 0.03$) (Figure 9). There was no significant

correlation between changes in GABA concentration and inverse efficiency score from low to

high enhancement ($r = -0.15$, $p = 0.56$) (Figure 9). In the distractor suppression experiment,

metabolite concentrations and tracking performance were correlated in similar directions as in

the main imaging experiment: The greater was the increase in GABA concentration from low

to high suppression, the smaller was the difference in inverse efficiency score between the two

distractor suppression conditions across participants ($r = -0.52$, $p = 0.03$) (Figure 9). Glx

concentration and tracking performance was correlated in the opposite direction: The greater

was the increase in Glx concentration from low to high suppression, the larger was the

difference in inverse efficiency score between the two distractor suppression conditions across

participants ($r = 0.46$, $p = 0.049$) (Figure 9).

Discussion

We examined how visual signals relevant to current goals can be processed while being

interfered with by goal-irrelevant distractor signals. For this purpose, we used a MOT task in

which participants tracked goal-relevant targets among goal-irrelevant distractors. fMRS
measurements were conducted in parietal and visual cortex while participants performed MOT
with increasing tracking load to increase target-distractor interference. The results showed that
GABA and Glx concentrations both increased in parietal cortex during tracking with high load
compared to low load. In visual cortex only the concentration of Glx increased with load. A
greater increase in GABA concentration and lower increase in Glx concentration in parietal
cortex predicted better tracking performance with increasing load across participants. Control
experiments showed similar correlational results between changes in parietal GABA and Glx
concentrations and tracking performance when the need to suppress distractors increased during
MOT. Taken together, our results suggest that GABAergic suppression of distractors in parietal
cortex is critical for minimizing target-distractor interference during tracking.

Our results suggest a key role of GABAergic inhibition in parietal cortex for goal-
directed visual information processing. GABA is a chief inhibitory neurotransmitter involved
in refining neural responses for goal-relevant signals (De La Vega et al., 2014; Isaacson &
Scanziani, 2011; Koolschijn et al., 2019; Okun & Lampl, 2008; Schmidt-Wilcke et al., 2018;
H. R. Snyder et al., 2010; Tremblay et al., 2016; Tritsch et al., 2016; Wehr & Zador, 2003). The
results of our main imaging experiment agree with this role of GABA as they show that
GABAergic inhibition in parietal cortex increased as the need to minimize interference between
targets and distractors increased in MOT with high load. Our correlational results further
support the association between parietal GABA and tracking performance during pronounced
target-distractor interference, as a greater increase in GABA concentration predicted better
tracking performance in high load across participants. To further establish the role of
GABAergic inhibition in parietal cortex for minimizing target-distractor interference, we
conducted control imaging experiments in which the need to enhance targets and suppress
distractors during MOT was varied separately (Bettencourt & Somers, 2009). The results
showed that parietal GABA was correlated with distractor suppression, such that a greater
increase in GABA concentration predicted better tracking performance when the need to
suppress distractors increased. This suggests that GABA may suppress competition from
spontaneous signals evoked by distractors during MOT, leading to refined processing and
robust representations of goal-relevant targets in parietal cortex (Lavie, 1995; Pinsk et al., 2004;
Torralbo et al., 2016). This is comparable to findings observed with other types of cognitive
tasks including semantic memory and selection during language processing, which showed that
increased levels of GABA sharpened the representation of relevant information for a more
efficient and precise selection among competitive semantic or linguistic information (De La

Vega et al., 2014; Koolschijn et al., 2019; Williams et al., 2023). It is possible that increased levels of GABA reflect a general mechanism underlying the brain's ability to resist interference from irrelevant and unwanted signals and is universally used as a mechanism in different types of cognitive tasks (Isaacson & Scanziani, 2011; Tremblay et al., 2016; Tritsch et al., 2016). The important role of GABAergic inhibition in parietal cortex to minimize target-distractor interference is consistent with previous results from functional magnetic resonance imaging (fMRI). These previous fMRI studies showed that the representation of goal-relevant information in parietal cortex, rather than visual cortex, is resilient to interference from goal-irrelevant signals (Bettencourt & Xu, 2016; Lorenc et al., 2018; Xu, 2024).

Although our results suggest that GABAergic inhibition plays a key role in minimizing target-distractor interference, glutamatergic excitation may still be relevant to other aspects of goal-directed visual information processing. In the main imaging experiment, the concentration of Glx increased with load in parietal and visual cortex. This load-dependent increase in Glx concentration may reflect target enhancement, as our control imaging experiments showed that participants with greater increase in Glx concentration in parietal cortex also tended to perform better when the need to enhance targets during MOT increased. No such association was found between GABA concentration and target enhancement. Although we did not test this directly in our study, glutamatergic target enhancement might occur at both early and late stages of cortical visual processing as electroencephalographic recordings showed increased stimulus-evoked potentials for targets compared to distractors already in (early) visual cortex (Adamian & Andersen, 2022; Störmer et al., 2013, 2014).

Changes in GABA and Glx concentrations in parietal cortex were differentially related to changes in tracking performance with increasing load in the main imaging experiment. Higher tracking performance with increasing load was associated with greater GABA increase and lower Glx increase across participants. This suggests that excitatory processing may be most necessary when the task proves challenging to participants, whereas high-performing participants engage inhibitory processing to a greater extent. Together with the results of our control imaging experiments, this suggests that effective minimization of target-distractor interference requires GABAergic suppression of distractors to a greater extent than glutamatergic enhancement of targets.

GABA and Glx concentrations in the control imaging experiments were not significantly different between low and high target enhancement and distractor suppression conditions. Significant results were only found when changes in GABA and Glx concentrations were correlated with changes in tracking performance between conditions across participants.

Contrary to the main imaging experiment, the visual stimulation was varied between conditions in the control imaging experiments (large vs. small disks in target enhancement and a total of eight vs. twelve disks in distractor suppression), following the approach of a previous study (Bettencourt & Somers, 2009). We could speculate that significant results only in the correlational analyses indicate that changes in metabolite concentrations between conditions reflected individual differences in tracking performance rather than variations in visual stimulation.

Using fMRI, it is difficult to separate the contributions of inhibitory and excitatory processing to the blood-oxygenation-level-dependent (BOLD) response. Therefore, it is uncertain whether increased BOLD responses with increasing tracking load as reported in previous studies (Alnaes et al., 2014; Culham et al., 1998; Frank et al., 2016; Jahn et al., 2012; Jovicich et al., 2001; Maechler et al., 2025; Nummenmaa et al., 2017; Shim et al., 2010) reflect increased inhibitory or excitatory processing or a combination of both. fMRS is a novel imaging approach that overcomes this limitation of fMRI (Apšvalka et al., 2015; Craven et al., 2024; Ip & Bridge, 2022; Koolschijn et al., 2023; Lally et al., 2014; Mullins, 2018, 2024; Stanley & Raz, 2018). In addition, with fMRS it is possible to detect changes in GABA and Glx concentrations with little delay compared to fMRI (Mullins, 2018). However, fMRS also has limitations that should be considered when interpreting the results of this study. Due to the large VOI size required for fMRS, the measured concentrations reflect a combination of extracellular and intracellular neurotransmitter concentrations across large portions of the parietal and occipital lobes. Although previous studies found extended activations in the parietal and visual cortex during MOT (Adamian & Andersen, 2022; Alnaes et al., 2014; Culham et al., 2001; Frank et al., 2016; Jahn et al., 2012; Jovicich et al., 2001; Maechler et al., 2025; Nummenmaa et al., 2017; Shim et al., 2010; Störmer et al., 2013, 2014), possible differences between subregions cannot be measured with fMRS. The time-resolved fMRS design used in this study makes it likely that changes in metabolite concentrations between conditions result from neurotransmission rather than slower metabolic processes related to neurotransmitter production. However, it is still possible that both processes contributed to the measured concentration changes.

In conclusion, the results of this study suggest that GABAergic inhibition in parietal cortex is crucial for goal-directed visual information processing. By minimizing interference between target and distractors through distractor suppression, GABAergic inhibition promotes the emergence of goal-relevant target representations in parietal cortex that are robust to interference from goal-irrelevant distractors.

579 **Author Contribution:** Z.W. and S.M.F. designed the study. Z.W., S.W., A.W., N.B., S.H., D.A.,

580 M.B., and S.M.F. collected data. Z.W. and S.M.F. analyzed data. Z.W. and S.M.F. interpreted

581 results. Z.W. and S.M.F. wrote the first draft of the manuscript. Z.W., M.B., and S.M.F. edited

582 and revised manuscript. S.M.F. approved final version of manuscript. S.M.F. acquired funding.

References

Adamian, N., & Andersen, S. K. (2022). Attentional Enhancement of Tracked Stimuli in Early Visual Cortex Has Limited Capacity. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *42*(46), 8709–8715. https://doi.org/10.1523/JNEUROSCI.0605-22.2022

Alnaes, D., Sneve, M. H., Espeseth, T., Endestad, T., Van De Pavert, S. H. P., & Laeng, B. (2014). Pupil size signals mental effort deployed during multiple object tracking and predicts brain activity in the dorsal attention network and the locus coeruleus. *Journal of Vision*, *14*(4), 1–1. https://doi.org/10.1167/14.4.1

Alvarez, G. A., & Franconeri, S. L. (2007). How many objects can you track?: Evidence for a resource-limited attentive tracking mechanism. *Journal of Vision*, *7*(13), 14. https://doi.org/10.1167/7.13.14

Apšvalka, D., Gadie, A., Clemence, M., & Mullins, P. G. (2015). Event-related dynamics of glutamate and BOLD effects measured using functional magnetic resonance spectroscopy (fMRS) at 3T in a repetition suppression paradigm. *NeuroImage*, *118*, 292–300. https://doi.org/10.1016/j.neuroimage.2015.06.015

Bang, J. W., Shibata, K., Frank, S. M., Walsh, E. G., Greenlee, M. W., Watanabe, T., & Sasaki, Y. (2018). Consolidation and reconsolidation share behavioural and neurochemical mechanisms. *Nature Human Behaviour*, *2*(7), 507–513. https://doi.org/10.1038/s41562-018-0366-8

Bettencourt, K. C., & Somers, D. C. (2009). Effects of target enhancement and distractor suppression on multiple object tracking capacity. *Journal of Vision*, *9*(7), 9–9. https://doi.org/10.1167/9.7.9

Bettencourt, K. C., & Xu, Y. (2016). Decoding the content of visual short-term memory under distraction in occipital and parietal areas. *Nature Neuroscience*, *19*(1), 150–157. https://doi.org/10.1038/nn.4174

Bottomley, P. A., Foster, T. B., & Darrow, R. D. (1984). Depth-resolved surface-coil spectroscopy (DRESS) for in Vivo 1H, 31P, and 13C NMR. *Journal of Magnetic Resonance (1969)*, *59*(2), 338–342. https://doi.org/10.1016/0022-2364(84)90179-3

Bottomley, P. A., & Hardy, C. J. (1987). PROGRESS in efficient three-dimensional spatially localized in vivo31P NMR spectroscopy using multidimensional spatially selective (ρ) pulses. *Journal of Magnetic Resonance (1969)*, *74*(3), 550–556. https://doi.org/10.1016/0022-2364(87)90276-9

Bracci, S., Daniels, N., & Op de Beeck, H. (2017). Task Context Overrules Object- and

Category-Related Representational Content in the Human Parietal Cortex. *Cerebral*

Cortex (New York, N.Y.: 1991), *27*(1), 310–321.

https://doi.org/10.1093/cercor/bhw419

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

Brefczynski, J. A., & DeYoe, E. A. (1999). A physiological correlate of the „spotlight" of

visual attention. *Nature Neuroscience*, *2*(4), 370–374. https://doi.org/10.1038/7280

Bruyer, R., & Brysbaert, M. (2011). Combining Speed and Accuracy in Cognitive

Psychology: Is the Inverse Efficiency Score (IES) a Better Dependent Variable than

the Mean Reaction Time (RT) and the Percentage Of Errors (PE)? *Psychologica*

Belgica, *51*(1), 5. https://doi.org/10.5334/pb-51-1-5

Buschman, T. J., & Miller, E. K. (2007). Top-Down Versus Bottom-up Control of Attention

in the Prefrontal and Posterior Parietal Cortices. *Science*, *315*(Ci), 1860–1862.

https://doi.org/10.1126/science.1138071

Carrasco, M. (2011). Visual attention: The past 25 years. *Vision Research*, *51*, 1484–1525.

https://doi.org/10.1016/j.visres.2011.04.012

Cavanagh, P., & Alvarez, G. (2005). Tracking multiple targets with multifocal attention.

Trends in Cognitive Sciences, *9*(7), 349–354.

https://doi.org/10.1016/j.tics.2005.05.009

Choi, I., Andronesi, O. C., Barker, P., Bogner, W., Edden, R. A. E., Kaiser, L. G., Lee, P.,

Marjańska, M., Terpstra, M., & De Graaf, R. A. (2021). Spectral editing in[1] H

magnetic resonance spectroscopy: Experts' consensus recommendations. *NMR in*

Biomedicine, *34*(5), e4411. https://doi.org/10.1002/nbm.4411

Christophel, T. B., Hebart, M. N., & Haynes, J.-D. (2012). Decoding the Contents of Visual

Short-Term Memory from Human Visual and Parietal Cortex. *The Journal of*

Neuroscience, *32*(38), 12983–12989. https://doi.org/10.1523/JNEUROSCI.0184-

12.2012

Chun, M. M., Golomb, J. D., & Turk-Browne, N. B. (2011). A Taxonomy of External and

Internal Attention. *Annual Review of Psychology*, *62*(1), 73–101.

https://doi.org/10.1146/annurev.psych.093008.100427

Cohen, M. R., & Maunsell, J. H. R. (2009). Attention improves performance primarily by

reducing interneuronal correlations. *Nature Neuroscience*, *12*(12), 1594–1600.

https://doi.org/10.1038/nn.2439

Craven, A. R., Dwyer, G., Ersland, L., Kazimierczak, K., Noeske, R., Sandøy, L. B., Johnsen,
E., & Hugdahl, K. (2024). GABA, glutamatergic dynamics and BOLD contrast
assessed concurrently using functional MRS during a cognitive task. *NMR in
Biomedicine*, *37*(3), e5065. https://doi.org/10.1002/nbm.5065

Culham, J. C., Brandt, S. A., Cavanagh, P., Kanwisher, N. G., Dale, A. M., & Tootell, R. B.
H. (1998). Cortical fMRI Activation Produced by Attentive Tracking of Moving
Targets. *Journal of Neurophysiology*, *80*(5), 2657–2670.
https://doi.org/10.1152/jn.1998.80.5.2657

Culham, J. C., Cavanagh, P., & Kanwisher, N. G. (2001). Attention Response Functions.
Neuron, *32*(4), 737–745. https://doi.org/10.1016/S0896-6273(01)00499-8

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis. I.
Segmentation and surface reconstruction. *NeuroImage*, *9*(2), 179–194.
https://doi.org/10.1006/nimg.1998.0395

De Bondt, T., De Belder, F., Vanhevel, F., Jacquemyn, Y., & Parizel, P. M. (2015). Prefrontal
GABA concentration changes in women-Influence of menstrual cycle phase,
hormonal contraceptive use, and correlation with premenstrual symptoms. *Brain
Research*, *1597*, 129–138. https://doi.org/10.1016/j.brainres.2014.11.051

De La Vega, A., Brown, M. S., Snyder, H. R., Singel, D., Munakata, Y., & Banich, M. T.
(2014). Individual Differences in the Balance of GABA to Glutamate in pFC Predict
the Ability to Select among Competing Options. *Journal of Cognitive Neuroscience*,
26(11), 2490–2502. https://doi.org/10.1162/jocn_a_00655

Duncan, N. W., Wiebking, C., & Northoff, G. (2014). Associations of regional GABA and
glutamate with intrinsic and extrinsic neural activity in humans—A review of
multimodal imaging studies. *Neuroscience & Biobehavioral Reviews*, *47*, 36–52.
https://doi.org/10.1016/j.neubiorev.2014.07.016

Egeth, H. E., & Yantis, S. (1997). VISUAL ATTENTION: Control, Representation, and Time
Course. *Annual Review of Psychology*, *48*(1), 269–297.
https://doi.org/10.1146/annurev.psych.48.1.269

Ester, E. F., Anderson, D. E., Serences, J. T., & Awh, E. (2013). A Neural Measure of
Precision in Visual Working Memory. *Journal of Cognitive Neuroscience*, *25*(5), 754–
761. https://doi.org/10.1162/jocn_a_00357

Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical
power analysis program for the social, behavioral, and biomedical sciences. *Behavior
Research Methods*, *39*(2), 175–191. https://doi.org/10.3758/BF03193146

Fischl, B., Sereno, M. I., & Dale, A. M. (1999). Cortical surface-based analysis. II: Inflation,

flattening, and a surface-based coordinate system. *NeuroImage*, *9*(2), 195–207.

https://doi.org/10.1006/nimg.1998.0396

Floyer-Lea, A., Wylezinska, M., Kincses, T., & Matthews, P. M. (2006). Rapid modulation of

GABA concentration in human sensorimotor cortex during motor learning. *Journal of*

Neurophysiology, *95*(3), 1639–1644. https://doi.org/10.1152/jn.00346.2005

Föcker, J., Beer, A. L., & Bavelier, D. (2023). Evidence of target enhancement and distractor

suppression in early visual areas. *Attention, Perception, & Psychophysics*, *85*(3), 734–

748. https://doi.org/10.3758/s13414-023-02673-w

Frangou, P., Emir, U. E., Karlaftis, V. M., Nettekoven, C., Hinson, E. L., Larcombe, S.,

Bridge, H., Stagg, C. J., & Kourtzi, Z. (2019). Learning to optimize perceptual

decisions through suppressive interactions in the human brain. *Nature*

Communications, *10*(1), 474. https://doi.org/10.1038/s41467-019-08313-y

Frank, S. M., Becker, M., Malloni, W. M., Sasaki, Y., Greenlee, M. W., & Watanabe, T.

(2023). Protocol to conduct functional magnetic resonance spectroscopy in different

age groups of human participants. *STAR Protocols*, *4*(3), 102493.

https://doi.org/10.1016/j.xpro.2023.102493

Frank, S. M., Becker, M., Qi, A., Geiger, P., Frank, U. I., Rosedahl, L. A., Malloni, W. M.,

Sasaki, Y., Greenlee, M. W., & Watanabe, T. (2022). Efficient learning in children

with rapid GABA boosting during and after training. *Current Biology: CB*, *32*(23),

5022-5030.e7. https://doi.org/10.1016/j.cub.2022.10.021

Frank, S. M., Sun, L., Forster, L., Tse, P. U., & Greenlee, M. W. (2016). Cross-Modal

Attention Effects in the Vestibular Cortex during Attentive Tracking of Moving

Objects. *The Journal of Neuroscience: The Official Journal of the Society for*

Neuroscience, *36*(50), 12720–12728. https://doi.org/10.1523/JNEUROSCI.2480-

16.2016

Gazzaley, A., & Nobre, A. C. (2012). Top-down modulation: Bridging selective attention and

working memory. *Trends in Cognitive Sciences*, *16*(2), 129–135.

https://doi.org/10.1016/j.tics.2011.11.014

Harris, K. D., & Thiele, A. (2011). Cortical state and attention. *Nature Reviews Neuroscience*,

12(9), 509–523. https://doi.org/10.1038/nrn3084

Harrison, S. A., & Tong, F. (2009). Decoding reveals the contents of visual working memory

in early visual areas. *Nature*, *458*(7238), 632–635.

https://doi.org/10.1038/nature07832

Ip, I. B., & Bridge, H. (2022). Investigating the neurochemistry of the human visual system
using magnetic resonance spectroscopy. *Brain Structure & Function*, *227*(4), 1491–
1505. https://doi.org/10.1007/s00429-021-02273-0

Isaacson, J. S., & Scanziani, M. (2011). How inhibition shapes cortical activity. *Neuron*,
72(2), 231–243. https://doi.org/10.1016/j.neuron.2011.09.027

Jahn, G., Wendt, J., Lotze, M., Papenmeier, F., & Huff, M. (2012). Brain activation during
spatial updating and attentive tracking of moving targets. *Brain and Cognition*, *78*(2),
105–113. https://doi.org/10.1016/j.bandc.2011.12.001

Jovicich, J., Peters, R. J., Koch, C., Braun, J., Chang, L., & Ernst, T. (2001). Brain Areas
Specific for Attentional Load in a Motion-Tracking Task. *Journal of Cognitive*
Neuroscience, *13*(8), 1048–1058. https://doi.org/10.1162/089892901753294347

Kolasinski, J., Logan, J. P., Hinson, E. L., Manners, D., Divanbeighi Zand, A. P., Makin, T.
R., Emir, U. E., & Stagg, C. J. (2017). A Mechanistic Link from GABA to Cortical
Architecture and Perception. *Current Biology: CB*, *27*(11), 1685-1691.e3.
https://doi.org/10.1016/j.cub.2017.04.055

Koolschijn, R. S., Clarke, W. T., Ip, I. B., Emir, U. E., & Barron, H. C. (2023). Event-related
functional magnetic resonance spectroscopy. *NeuroImage*, *276*, 120194.
https://doi.org/10.1016/j.neuroimage.2023.120194

Koolschijn, R. S., Emir, U. E., Pantelides, A. C., Nili, H., Behrens, T. E. J., & Barron, H. C.
(2019). The Hippocampus and Neocortical Inhibitory Engrams Protect against
Memory Interference. *Neuron*, *101*(3), 528-541.e6.
https://doi.org/10.1016/j.neuron.2018.11.042

Lally, N., Mullins, P. G., Roberts, M. V., Price, D., Gruber, T., & Haenschel, C. (2014).
Glutamatergic correlates of gamma-band oscillatory activity during cognition: A
concurrent ER-MRS and EEG study. *NeuroImage*, *85 Pt 2*, 823–833.
https://doi.org/10.1016/j.neuroimage.2013.07.049

Lavie, N. (1995). Perceptual load as a necessary condition for selective attention. *Journal of*
Experimental Psychology. Human Perception and Performance, *21*(3), 451–468.
https://doi.org/10.1037//0096-1523.21.3.451

Lin, A., Andronesi, O., Bogner, W., Choi, I.-Y., Coello, E., Cudalbu, C., Juchem, C., Kemp,
G. J., Kreis, R., Krššák, M., Lee, P., Maudsley, A. A., Meyerspeer, M., Mlynarik, V.,
Near, J., Öz, G., Peek, A. L., Puts, N. A., Ratai, E.-M., … Experts' Working Group on
Reporting Standards for MR Spectroscopy. (2021). Minimum Reporting Standards for
in vivo Magnetic Resonance Spectroscopy (MRSinMRS): Experts' consensus

recommendations. *NMR in Biomedicine*, *34*(5), e4484.

https://doi.org/10.1002/nbm.4484

Lorenc, E. S., Sreenivasan, K. K., Nee, D. E., Vandenbroucke, A. R. E., & D'Esposito, M. (2018). Flexible Coding of Visual Working Memory Representations during Distraction. *The Journal of Neuroscience*, *38*(23), 5267–5276. https://doi.org/10.1523/JNEUROSCI.3061-17.2018

Maechler, M. R., Choe, E., Cavanagh, P., Kohler, P. J., & Tse, P. U. (2025). Hemifield Specificity of Attention Response Functions during Multiple-Object Tracking. *The Journal of Neuroscience*, *45*(19), e1340242025. https://doi.org/10.1523/JNEUROSCI.1340-24.2025

Mescher, M., Merkle, H., Kirsch, J., Garwood, M., & Gruetter, R. (1998). Simultaneousin vivo spectral editing and water suppression. *NMR in Biomedicine*, *11*(6), 266–272. https://doi.org/10.1002/(SICI)1099-1492(199810)11:6%253C266::AID-NBM530%253E3.0.CO;2-J

Mescher, M., Tannus, A., Johnson, M. O., & Garwood, M. (1996). Solvent Suppression Using Selective Echo Dephasing. *Journal of Magnetic Resonance, Series A*, *123*(2), 226–229. https://doi.org/10.1006/jmra.1996.0242

Meyerhoff, H. S., Papenmeier, F., & Huff, M. (2017). Studying visual attention using the multiple object tracking paradigm: A tutorial review. *Attention, Perception, & Psychophysics*, *79*(5), 1255–1274. https://doi.org/10.3758/s13414-017-1338-1

Mullins, P. G. (2018). Towards a theory of functional magnetic resonance spectroscopy (FMRS): A meta-analysis and discussion of using MRS to measure changes in neurotransmitters in real time. *Scandinavian Journal of Psychology*, *59*(1), 91–103. https://doi.org/10.1111/sjop.12411

Mullins, P. G. (2024). Considerations for event-related gamma-aminobutyric acid functional magnetic resonance spectroscopy. *NMR in Biomedicine*, *37*(11), e5215. https://doi.org/10.1002/nbm.5215

Mullins, P. G., McGonigle, D. J., O'Gorman, R. L., Puts, N. A. J., Vidyasagar, R., Evans, C. J., Cardiff Symposium on MRS of GABA, & Edden, R. A. E. (2014). Current practice in the use of MEGA-PRESS spectroscopy for the detection of GABA. *NeuroImage*, *86*, 43–52. https://doi.org/10.1016/j.neuroimage.2012.12.004

Nummenmaa, L., Oksama, L., Glerean, E., & Hyönä, J. (2017). Cortical Circuit for Binding Object Identity and Location During Multiple-Object Tracking. *Cerebral Cortex (New York, N.Y.: 1991)*, *27*(1), 162–172. https://doi.org/10.1093/cercor/bhw380

Ogg, R. J., Kingsley, P. B., & Taylor, J. S. (1994). WET, a T1- and B1-insensitive water-
suppression method for in vivo localized 1H NMR spectroscopy. *Journal of Magnetic*
Resonance. Series B, *104*(1), 1–10. https://doi.org/10.1006/jmrb.1994.1048

Okun, M., & Lampl, I. (2008). Instantaneous correlation of excitation and inhibition during
ongoing and sensory-evoked activities. *Nature Neuroscience*, *11*(5), 535–537.
https://doi.org/10.1038/nn.2105

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: Transforming
numbers into movies. *Spatial Vision*, *10*(4), 437–442.
https://doi.org/10.1163/156856897X00366

Petroff, O. A. C. (2002). Book Review: GABA and Glutamate in the Human Brain. *The*
Neuroscientist, *8*(6), 562–573. https://doi.org/10.1177/1073858402238515

Pinsk, M. A., Doniger, G. M., & Kastner, S. (2004). Push-pull mechanism of selective
attention in human extrastriate cortex. *Journal of Neurophysiology*, *92*(1), 622–629.
https://doi.org/10.1152/jn.00974.2003

Provencher, S. W. (1993). Estimation of metabolite concentrations from localized in vivo
proton NMR spectra. *Magnetic Resonance in Medicine*, *30*(6), 672–679.
https://doi.org/10.1002/mrm.1910300604

Provencher, S. W. (2001). Automatic quantitation of localized *in vivo*[1] H spectra with
LCModel. *NMR in Biomedicine*, *14*(4), 260–264. https://doi.org/10.1002/nbm.698

Pylyshyn, Z. W., & Storm, R. W. (1988). Tracking multiple independent targets: Evidence for
a parallel tracking mechanism*. *Spatial Vision*, *3*(3), 179–197.
https://doi.org/10.1163/156856888X00122

Rademaker, R. L., Chunharas, C., & Serences, J. T. (2019). Coexisting representations of
sensory and mnemonic information in human visual cortex. *Nature Neuroscience*,
22(8), 1336–1344. https://doi.org/10.1038/s41593-019-0428-x

Rorden, C. (2008). Event related potentials reveal that increasing perceptual load leads to
increased responses for target stimuli and decreased responses for irrelevant stimuli.
Frontiers in Human Neuroscience, *2*. https://doi.org/10.3389/neuro.09.004.2008

Schmidt-Wilcke, T., Fuchs, E., Funke, K., Vlachos, A., Müller-Dahlhaus, F., Puts, N. A. J.,
Harris, R. E., & Edden, R. A. E. (2018). GABA—from Inhibition to Cognition:
Emerging Concepts. *The Neuroscientist*, *24*(5), 501–515.
https://doi.org/10.1177/1073858417734530

Scholl, B. J. (2001). Objects and attention: The state of the art. *Cognition*, *80*(1–2), 1–46.
https://doi.org/10.1016/S0010-0277(00)00152-9

Schuff, N., Meyerhoff, D. J., Mueller, S., Chao, L., Sacrey, D. T., Laxer, K., & Weiner, M. W. (2006). N-Acetylaspartate as a Marker of Neuronal Injury in Neurodegenerative Disease. In J. R. Moffett, S. B. Tieman, D. R. Weinberger, J. T. Coyle, & A. M. A. Namboodiri (Hrsg.), *N-Acetylaspartate* (Bd. 576, S. 241–262). Springer US. https://doi.org/10.1007/0-387-30172-0_17

Serences, J. T. (2016). Neural mechanisms of information storage in visual short-term memory. *Vision Research*, *128*, 53–67. https://doi.org/10.1016/j.visres.2016.09.010

Serences, J. T., Yantis, S., Culberson, A., & Awh, E. (2004). Preparatory activity in visual cortex indexes distractor suppression during covert spatial orienting. *Journal of Neurophysiology*, *92*(6), 3538–3545. https://doi.org/10.1152/jn.00435.2004

Shibata, K., Sasaki, Y., Bang, J. W., Walsh, E. G., Machizawa, M. G., Tamaki, M., Chang, L.-H., & Watanabe, T. (2017). Overlearning hyperstabilizes a skill by rapidly making neurochemical processing inhibitory-dominant. *Nature Neuroscience*, *20*(3), 470–475. https://doi.org/10.1038/nn.4490

Shim, W. M., Alvarez, G. A., Vickery, T. J., & Jiang, Y. V. (2010). The Number of Attentional Foci and Their Precision Are Dissociated in the Posterior Parietal Cortex. *Cerebral Cortex*, *20*(6), 1341–1349. https://doi.org/10.1093/cercor/bhp197

Snyder, A. C., Morais, M. J., & Smith, M. A. (2016). Dynamics of excitatory and inhibitory networks are differentially altered by selective attention. *Journal of Neurophysiology*, *116*(4), 1807–1820. https://doi.org/10.1152/jn.00343.2016

Snyder, H. R., Hutchison, N., Nyhus, E., Curran, T., Banich, M. T., O'Reilly, R. C., & Munakata, Y. (2010). Neural inhibition enables selection during language processing. *Proceedings of the National Academy of Sciences*, *107*(38), 16483–16488. https://doi.org/10.1073/pnas.1002291107

Sprague, T. C., Ester, E. F., & Serences, J. T. (2014). Reconstructions of Information in Visual Spatial Working Memory Degrade with Memory Load. *Current Biology*, *24*(18), 2174–2180. https://doi.org/10.1016/j.cub.2014.07.066

Sprague, T. C., Saproo, S., & Serences, J. T. (2015). *Visual attention mitigates information loss in small- and large-scale neural codes*. *19*(4), 215–226.

Srivastava, S., Wang, W. Y., & Eckstein, M. P. (2023). *Bridging Neurons and Behavior in a Convolutional Neural Network with Emergent Human-like Covert Attention*. Neuroscience. https://doi.org/10.1101/2023.09.17.558171

Stagg, C. J. (2014). Magnetic Resonance Spectroscopy as a tool to study the role of GABA in
motor-cortical plasticity. *NeuroImage*, *86*, 19–27.
https://doi.org/10.1016/j.neuroimage.2013.01.009

Stanley, J. A., & Raz, N. (2018). Functional Magnetic Resonance Spectroscopy: The „New"
MRS for Cognitive Neuroscience and Psychiatry Research. *Frontiers in Psychiatry*, *9*,
76. https://doi.org/10.3389/fpsyt.2018.00076

Störmer, V. S., Alvarez, G. A., & Cavanagh, P. (2014). Within-Hemifield Competition in
Early Visual Areas Limits the Ability to Track Multiple Objects with Attention. *The*
Journal of Neuroscience, *34*(35), 11526–11533.
https://doi.org/10.1523/JNEUROSCI.0980-14.2014

Störmer, V. S., Winther, G. N., Li, S.-C., & Andersen, S. K. (2013). Sustained Multifocal
Attentional Enhancement of Stimulus Processing in Early Visual Areas Predicts
Tracking Performance. *The Journal of Neuroscience*, *33*(12), 5346–5351.
https://doi.org/10.1523/JNEUROSCI.4015-12.2013

Tamaki, M., Wang, Z., Barnes-Diana, T., Guo, D., Berard, A. V., Walsh, E., Watanabe, T., &
Sasaki, Y. (2020). Complementary contributions of non-REM and REM sleep to
visual learning. *Nature Neuroscience*, *23*(9), 1150–1156.
https://doi.org/10.1038/s41593-020-0666-y

Torralbo, A., Kelley, T. A., Rees, G., & Lavie, N. (2016). Attention induced neural response
trade-off in retinotopic cortex under load. *Scientific Reports*, *6*(1), 33041.
https://doi.org/10.1038/srep33041

Townsend, J. T., & Ashby, F. G. (1978). Methods of modeling capacity in simple processing
systems. In *Cognitive theory* (S. 199–239). Psychology Press.

Townsend, J. T., & Ashby, F. G. (viaf)94267600. (1983). *The stochastic modeling of*
elementary psychological processes. Cambridge : Cambridge university press.
http://lib.ugent.be/catalog/rug01:000725686

Tremblay, R., Lee, S., & Rudy, B. (2016). GABAergic Interneurons in the Neocortex: From
Cellular Properties to Circuits. *Neuron*, *91*(2), 260–292.
https://doi.org/10.1016/j.neuron.2016.06.033

Tritsch, N. X., Granger, A. J., & Sabatini, B. L. (2016). Mechanisms and functions of GABA
co-release. *Nature Reviews Neuroscience*, *17*(3), 139–145.
https://doi.org/10.1038/nrn.2015.21

Van Ede, F., & Nobre, A. C. (2023). Turning Attention Inside Out: How Working Memory

Serves Behavior. *Annual Review of Psychology*, *74*(1), 137–165.

https://doi.org/10.1146/annurev-psych-021422-041757

Van Essen, D. C. (2005). A Population-Average, Landmark- and Surface-based (PALS) atlas

of human cerebral cortex. *NeuroImage*, *28*(3), 635–662.

https://doi.org/10.1016/j.neuroimage.2005.06.058

Vaziri-Pashkam, M., & Xu, Y. (2017). Goal-Directed Visual Processing Differentially

Impacts Human Ventral and Dorsal Visual Representations. *The Journal of*

Neuroscience: The Official Journal of the Society for Neuroscience, *37*(36), 8767–

8782. https://doi.org/10.1523/JNEUROSCI.3392-16.2017

Wehr, M., & Zador, A. M. (2003). Balanced inhibition underlies tuning and sharpens spike

timing in auditory cortex. *Nature*, *426*(6965), 442–446.

https://doi.org/10.1038/nature02116

Williams, S., Ralph, M. L., & Jung, J. (2023). *The role of GABA in semantic memory and its*

neuroplasticity. https://doi.org/10.7554/eLife.91771.1

Wilson, M., Andronesi, O., Barker, P. B., Bartha, R., Bizzi, A., Bolan, P. J., Brindle, K. M.,

Choi, I.-Y., Cudalbu, C., Dydak, U., Emir, U. E., Gonzalez, R. G., Gruber, S.,

Gruetter, R., Gupta, R. K., Heerschap, A., Henning, A., Hetherington, H. P., Huppi, P.

S., … Howe, F. A. (2019). Methodological consensus on clinical proton MRS of the

brain: Review and recommendations. *Magnetic Resonance in Medicine*, *82*(2), 527–

550. https://doi.org/10.1002/mrm.27742

Wöstmann, M., Störmer, V. S., Obleser, J., Addleman, D. A., Andersen, S. K., Gaspelin, N.,

Geng, J. J., Luck, S. J., Noonan, M. P., Slagter, H. A., & Theeuwes, J. (2022). Ten

simple rules to study distractor suppression. *Progress in Neurobiology*, *213*, 102269.

https://doi.org/10.1016/j.pneurobio.2022.102269

Xu, Y. (2024). The human posterior parietal cortices orthogonalize the representation of

different streams of information concurrently coded in visual working memory. *PLOS*

Biology, *22*(11), e3002915. https://doi.org/10.1371/journal.pbio.3002915

Xu, Y., & Chun, M. M. (2009). Selecting and perceiving multiple visual objects. *Trends in*

Cognitive Sciences, *13*(4), 167–174. https://doi.org/10.1016/j.tics.2009.01.008

Yamada, T., Watanabe, T., & Sasaki, Y. (2024). Plasticity–stability dynamics during post-

training processing of learning. *Trends in Cognitive Sciences*, *28*(1), 72–83.

https://doi.org/10.1016/j.tics.2023.09.002

Table 1. fMRS checklist proposed by Choi et al. (2021).

Site (Name or Number): Brain Imaging Center at University of Regensburg / Bezirksklinikum Regensburg	
1. Hardware	
a. Field strength [T]	3 T
b. Manufacturer	Siemens
c. Model (software version if available)	Prisma
d. RF coils: nuclei (transmit/receive), number of channels, type, body part	64 channel head/neck coil
e. Additional hardware	N/A
2. Acquisition	
a. Pulse sequence	Siemens MEGAPRESS sequence; Siemens PRESS sequence
b. Volume of Interest (VOI) locations	Parietal lobes; occipital lobes
c. Nominal VOI size [cm^3, mm^3]	2 x 2 x 4 cm (parietal lobes); 2 x 2 x 2 cm (occipital lobes)
d. Repetition Time (TR), Echo Time (TE) [ms, s]	MEGAPRESS (for each of Edit On & Edit Off): TR 1500 ms; TE = 68 ms; PRESS: TR 2000 ms; TE = 30 ms
e. Total number of Excitations or acquisitions per spectrum	MEGAPRESS: 472 scans in total, which corresponded to the following experimental conditions (summed across all time-points): on-ramp: 20 scans; cueing phase low load/low enhancement/low suppression: 18 scans; tracking phase low load/low enhancement/low suppression: 72 scans; response phase low load/low enhancement/low suppression: 18 scans; cueing phase high load/high enhancement/ high suppression: 18 scans; tracking phase high load/high enhancement/high suppression: 72 scans; response phase high load/high enhancement/high suppression: 18 scans; intertrial interval 216 scans; off-ramp: 20 scans. For each participant spectra for the tracking phases in each condition were averaged across trials with correct participant response. PRESS: 352 scans in total, which corresponded to the following experimental conditions (summed across all time-points): on-ramp: 32 scans; cueing phase low load/low enhancement/low suppression: 9 scans; tracking phase low load/low enhancement/low suppression: 54 scans; response phase low load/low enhancement/low suppression: 9 scans;

	cueing phase high load/high enhancement/high suppression: 9 scans; tracking phase high load/high enhancement/high suppression: 54 scans; response phase high load/high enhancement/high suppression: 9 scans; intertrial interval 144 scans; off-ramp: 32 scans. For each participant spectra for the tracking phases in each condition were averaged across trials with correct participant response.
f. Additional sequence parameters (spectral width in Hz, number of spectral points, frequency offsets)	Spectral bandwidth: 1200 Hz; Spectral points: 1024
g. Water Suppression Method	WET
h. Shimming Method, reference peak, and thresholds for "acceptance of shim" chosen	Automated B_0 field mapping to < 20 Hz
i. Triggering or motion correction method	N/A
3. Data analysis methods and outputs	
a. Analysis software	LC Model Version 6.3-R; Osprey Version 2.4.0
b. Processing steps deviating from quoted reference or product	N/A
c. Output measure	Tissue-corrected concentrations relative to NAA from the same spectrum
d. Quantification references and assumptions, fitting model assumptions	Basis sets were imported from LC model software package and included the following spectra: MEGAPRESS difference: γ-Aminobutyric Acid (GABA), Glutamine (Gln), Glutamate (Glu), Glutathione (GSH), N-Acetylaspartate (NAA), N-Acetylaspartylglutamate (NAAG). PRESS: Aspartate (Asp), Creatine (Cr), Phosphocreatine (PCr), γ-Aminobutyric Acid (GABA), Glucose (Glc), Glutamine (Gln), Glutamate (Glu), Glycerophosphocholine (GPC), Glutathione (GSH), myo-Inositol (Ins), L-Lactate (Lac), N-Acetylaspartate (NAA), N-Acetylaspartylglutamate (NAAG), scyllo-Inositol (Scyllo), Taurine (Tau).
4. Data Quality	
a. Reported variables	CRLB, shimming, S/N
b. Data exclusion criteria	CRLB > 25%, shimming >= 20 Hz, S/N < 5
c. Quality measures of postprocessing Model fitting	CRLB, S/N

| d. Sample Spectrum | See Supplementary Figures 1-4 |

Figures

 Figure 1. Design of the main imaging experiment. Example trials of the multiple object tracking

 (MOT) task with low and high attentional load conditions corresponding to tracking two and

 four targets among distractors, respectively. γ-aminobutyric acid (GABA) was measured with

 MEGAPRESS scans. Glx (glutamate and its precursor glutamine) was measured with PRESS

 scans.

 Figure 2. Tracking performance across fMRS runs in the main imaging experiment. The thick

 line shows mean results across participants. Thin lines show results for each participant. The

 lower the inverse efficiency score, the faster and more accurate participants responded. The

 result of a paired-sample *t*-test between tracking with high load and low load is denoted by

 asterisks. *** $p < 0.001$.

Figure 3. Metabolite concentrations in the main imaging experiment. Thick lines show mean

results across participants for the parietal and visual volumes-of-interest (VOIs). Thin lines

show results for each participant. GABA and Glx concentrations are normalized to the

concentration of N-acetyl-aspartate (NAA). Results of paired-sample *t*-tests between tracking with high load and low load are denoted by asterisks. * $p < 0.05$, ** $p < 0.01$.

Figure 4. Correlational results in the main imaging experiment. Load-dependent changes in GABA and Glx concentrations during tracking were calculated by subtracting low load from high load concentrations for each participant. More positive values on the x-axis correspond to greater concentration increase from low to high load. Load-dependent changes in tracking performance were calculated by subtracting the inverse efficiency score in low load from high load for each participant. More positive values on the y-axis correspond to greater increase in inverse efficiency score from low to high load (i.e., greater decrease of response accuracy and greater increase of response time from low to high load). Each correlational analysis was carried out between metabolite concentrations and tracking performance from the same fMRS run. Each dot shows the result from a different participant.

Figure 5. Design of control behavioral and imaging experiments for the measurement of target enhancement and distractor suppression. In the control imaging experiments, GABA and Glx concentrations during MOT were measured in the parietal VOI exactly as in the main imaging experiment with MEGAPRESS and PRESS scans, respectively (see Figure 1). Cueing and response periods were 3 s-long each for MEGAPRESS scans and 2 s-long each for PRESS scans. A jittered intertrial interval was included after each MOT trial exactly as in the main imaging experiment.

 Figure 6. Tracking performance across runs in each control behavioral experiment. (**A**) Target
974 enhancement experiment. (**B**) Distractor suppression experiment. Otherwise same as Figure 2.
975 *** $p < 0.001$.
976

977
978 **Figure 7.** Tracking performance across fMRS runs in each control imaging experiment. (**A**)
979 Target enhancement experiment. (**B**) Distractor suppression experiment. Otherwise same as
980 Figure 6. ** $p < 0.01$, *** $p < 0.001$.
981

Figure 8. GABA and Glx measurements in the parietal VOI in the control imaging experiments. Otherwise same as Figure 3.

Figure 9. Correlational results in the control imaging experiments. More positive values on the x-axis correspond to greater concentration increase from low to high target enhancement and distractor suppression. More positive values on the y-axis correspond to greater increase in inverse efficiency score from low to high target enhancement and distractor suppression (i.e., greater decrease of response accuracy and greater increase of response time). Each correlational analysis was carried out between metabolite concentrations and tracking performance from the same fMRS run. Otherwise same as Figure 4.

 Supplementary Material

1000

1001 **Supplementary Figure 1.** Mean MEGAPRESS difference spectra across participants in the

1002 main imaging experiment. The gray shaded area shows standard-error-of-the mean. GABA at 3

1003 ppm and NAA at 2 ppm.

1004

1005

1006

1007 **Supplementary Figure 2.** Mean PRESS spectra across participants in the main imaging

1008 experiment. Glx between 2.1 and 2.5 ppm and NAA at 2 ppm. Otherwise same as

1009 Supplementary Figure 1.

1010

1011

1012 **Supplementary Figure 3.** Mean MEGAPRESS difference spectra across participants in the

1013 control imaging experiments. Otherwise same as Supplementary Figure 1.

1014

1015

1016 **Supplementary Figure 4.** Mean PRESS spectra across participants in the control imaging

1017 experiments. Otherwise same as Supplementary Figure 2.

1019

Supplementary Figure 5. Tracking performance across fMRS runs in the main imaging

experiment. (**A**) Response accuracy. Participants responded less accurately in tracking with

high load than with low load [paired-sample *t*-test on arcsin-square-root transformed response

accuracy; $t(34) = -4.40$, $p < 0.001$, $d = -0.74$]. Otherwise same as Figure 2. (**B**) Median response

time. Participants responded slower in tracking with high load than with low load [paired-

sample *t*-test on log-transformed response time; $t(34) = 5.42$, $p < 0.001$, $d = 0.92$]. Otherwise

same as (A). *** $p < 0.001$.

1027

1028

1029 **Supplementary Figure 6.** Tracking performance across runs in each control behavioral

1030 experiment. (**A**) Response accuracy in the target enhancement experiment. Participants

1031 responded less accurately in tracking with high than with low target enhancement [paired-

sample *t*-test on arcsin-square-root transformed response accuracy; $t(19)$ = -4.84, $p < 0.001$, d = -1.08]. Otherwise same as Supplementary Figure 5A. (**B**) Median response time in the target enhancement experiment. Participants responded slower in tracking with high than with low target enhancement [paired-sample *t*-test on log-transformed response time; $t(19)$ = 8.24, $p < 0.001$, d = 1.84]. Otherwise same as Supplementary Figure 5B. (**C**) Response accuracy in the distractor suppression experiment. Participants responded less accurately in tracking with high than with low distractor suppression [paired-sample *t*-test on arcsin-square-root transformed response accuracy; $t(19)$ = -6.05, $p < 0.001$, d = -1.35]. Otherwise same as (A). (**D**) Median response time in the distractor suppression experiment. Participants responded slower in tracking with high than with low distractor suppression [paired-sample *t*-test on log-transformed response time; $t(19)$ = 8.12, $p < 0.001$, d = 1.82]. Otherwise same as (B). *** $p < 0.001$.

Supplementary Figure 7. Tracking performance across fMRS runs in each control imaging experiment. (**A**) Response accuracy in the target enhancement experiment. Participants responded less accurately in tracking with high than with low target enhancement [paired-sample *t*-test on arcsin-square-root transformed response accuracy; $t(18)$ = -3.89, $p = 0.001$, d = -0.89]. Otherwise same as Supplementary Figure 6A. (**B**) Median response time in the target enhancement experiment. Participants responded slower in tracking with high than with low target enhancement [paired-sample *t*-test on log-transformed response time; $t(18)$ = 6.41, $p < 0.001$, d = 1.47]. Otherwise same as Supplementary Figure 6B. (**C**) Response accuracy in the distractor suppression experiment. Participants responded less accurately in tracking with high than with low distractor suppression [paired-sample *t*-test on arcsin-square-root transformed response accuracy; $t(18)$ = -3.50, $p = 0.003$, d = -0.80]. Otherwise same as (A). (**D**) Median response time in the distractor suppression experiment. Participants responded slower in

1058 tracking with high than with low distractor suppression [paired-sample t-test on log-
1059 transformed response time; $t(18) = 2.76$, $p = 0.01$, $d = 0.63$]. Otherwise same as (B). * $p < 0.05$,
1060 ** $p < 0.01$, *** $p < 0.001$.
1061