Asymmetric phosphoinositide lipid bilayers generated by spontaneous lipid insertion

Gwendal Guerin[1], Thi Lan Anh Nguyen[1], John Manzi[1], Christophe Le Clainche[2], Julien Heuvingh[3], Julien Berro[4,5,6], Nicolas Rodriguez[7], Sophie Cribier[8], and Feng-Ching Tsai[1]*

1. Institut Curie, Université PSL, Sorbonne Université, CNRS UMR168, Laboratoire Physique des Cellules et Cancer, Paris 75005, France

2. Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198 Gif-sur-Yvette, France

3. PMMH, ESPCI ParisPSL University, CNRS, Université Paris Cité, Sorbonne Université Paris France

4. Department of Molecular Biophysics and Biochemistry; Yale University, New Haven, CT, USA

5. Nanobiology Institute, Yale University; West Haven, CT 06516, USA

6. Department of Cell Biology, Yale University School of Medicine; New Haven, CT 06520, USA

7. Sorbonne Université, École Normale Supérieure, PSL University, CNRS, Laboratoire de Physique de l'ENS, Paris, France

8. Sorbonne Université, École Normale Supérieure, PSL University, CNRS, Laboratoire Chimie Physique et Chimie du Vivant, 75005, Paris, France

*Corresponding author: feng-ching.tsai@curie.fr

Phosphatidylinositol phosphate (PIP) lipids, enriched on the cytoplasmic leaflet of the plasma membrane, are key regulators of diverse cellular processes, often through interactions with partner proteins that regulate actin assembly. Supported lipid bilayers (SLBs) provide a powerful model system to study the interactions of PIP lipids with their partner proteins. However, despite advances in SLB preparation methods, it remains a challenge to robustly obtain fluid SLBs in which PIP lipids are both mobile and asymmetrically distributed. In this study, we report a simple and robust method to generate asymmetric SLBs containing tunable amounts of $PI(4,5)P_2$. By dissolving $PI(4,5)P_2$ below its critical micelle concentration (CMC), we enable its spontaneous insertion into the upper leaflet of SLBs exposed to bulk solution. The mobility of $PI(4,5)P_2$ is confirmed using fluorescence recovery after photobleaching (FRAP). Furthermore, we demonstrate that $PI(4,5)P_2$ incorporated using this method retains its functionality, recruiting binding partners, actin-membrane linker ezrin, and myosin 1 motors capable of sliding actin filaments on the SLBs. Our method offers a straightforward strategy to generate asymmetric $PI(4,5)P_2$-containing SLBs and is applicable to other lipid species with high CMC values.

Introduction

The asymmetric distribution of lipid species is a fundamental feature of the plasma membrane in eukaryotic cells[1]. Sphingomyelin (SM) is abundant in the exoplasmic leaflet, whereas phosphatidylethanolamine (PE), phosphatidylserine (PS), and notably phosphatidylinositol (PI) lipids are predominantly found in the cytoplasmic leaflet[1]. PI lipids account for $\sim 10\%$ of total lipids in mammalian cells[2]. Despite being a minor lipid species in the plasma membrane, PI and its phosphorylated derivatives, phosphatidylinositol phosphate (PIP) lipids, are recognized as key cellular regulators, affecting processes such as membrane trafficking and actin cytoskeleton assembly[3,4]. At neutral pH, PIP lipids exhibit a greater negative charge at the head group than PS, due to the presence of two negative charges per phosphate[5]. The phosphate modification enables PIP lipids to interact specifically with phosphoinositide-binding protein domains including pleckstrin homology (PH), Phox (PX), and BAR domains[6–8]. Among the PIP lipids, $PI(4,5)P_2$ (hereafter referred to as PIP_2) is the most abundant, accounting for 0.5 up to 4 mol% of the total plasma membrane lipids, estimated to be ~ 4000 up to $34,000$ molecules per μm^2 [5,6]. By interacting with hundreds of effector proteins, including actin-binding proteins, PIP_2 plays a central role in regulating numerous membrane-associated cellular processes[4,5,9,10]. PIP_2 has therefore been extensively studied, especially through in vitro reconstitution assays using model membranes. For instance, PIP_2 has been shown to form clusters by interacting with positively charged divalent ions such as Ca^{2+} and Mg^{2+} [11–14.] Moreover, interactions between PIP_2 and effector proteins, including members from the ezrin-radixin-moesin (ERM) protein family and actin nucleation promoting factors, have been implicated in the regulation of actin assembly[4,9,10,15–19]. However, despite these

significant findings, how the dynamic interactions between PIP lipids and their effector proteins give rise to the diverse membrane processes in cells remains to be fully resolved.

Supported lipid bilayers (SLBs) are widely used model membrane platforms for studying lipid-lipid and protein-lipid interactions. Thanks to the substrate support and the planar shape, SLBs are particularly well suited for experimental techniques that require the imaging plane to be close to the substrate, such as atomic force microscopy, total internal reflection fluorescence (TIRF) microscopy, fluorescence correlation spectroscopy (FCS), single molecule tracking, and super-resolution imaging[20–24]. Common methods for generating SLBs include Langmuir-Blodgett deposition, the fusion of small vesicles (of the order of 100 nm in diameter), the rupture of giant vesicles (of the order of 10 μm in diameter), and rapid solvent exchange to drive SLB formation[25–28]. Vesicle fusion and rupture can produce symmetric SLBs (sym-SLBs), in which the lipid composition is the same on both leaflets. However, a key limitation with sym-SLBs is the inherent and often undesirable interactions between the lipids and the substrates. These lipid-substate interactions are known to reduce lipid mobility, likely due to factors such as surface roughness and electrostatic interactions[29–32]. Such interactions are particularly pronounced for negatively charged lipids like PIP$_2$, which can become nearly immobilized in the bottom lipid leaflet that contacts the glass substrate[31,33–36]. Such immobilization compromises the quantitative analysis of lipid dynamics, and can also affect lipid lateral organization, such as phase separation, in the bilayer[37–39].

To mitigate substrate-induced lipid immobilization, one approach is to replace glass substrates with mica, which provides a smoother and more hydrophilic surface compared to glass[29]. Alternatively, introducing polymer cushions, such as polyethylene glycol (PEG), between the substrate and bilayer can reduce friction[40,41]. Moreover, suspended planar lipid bilayers generated

in micropores are elegant approaches that avoids the use of solid substrates[42–45]. However, these approaches often involve complex surface modifications or microfabrication[41–45].

To enable lipid-protein interactions while minimizing substrate-induced immobilization, an effective approach is to construct asymmetric SLBs (asym-SLBs) in which the lipid species of interest, such as PIP$_2$, are enriched in the top leaflet facing the bulk solution. Langmuir-Blodgett deposition can be used to generate asym-SLBs, but it requires sophisticated procedures[26]. Other methods include methyl-β-cyclodextrin-assisted lipid exchange to enrich sphingomyelin in the top lipid leaflet facing to the buffer[46,47], using negatively charged surfaces to promote a higher amount of negatively charged lipids in the top leaflet[48], transferring lipids to the top leaflet by the hemifusion of vesicles on the SLBs[49], and directly using asymmetric vesicles to form SLBs[50]. While these methods enable the generation of asym-SLBs, they are often limited to specific lipid types for asymmetric incorporation, and they do not offer precise control over the asymmetric distribution of lipids between the two leaflets. Recently, an elegant technique has been developed to generate asym-SLBs for PIP$_2$, in which additional vector small unilamellar vesicles (SUVs) containing PIP$_2$ were introduced to dope the top leaflet of pre-formed SLBs with PIP$_2$, although the underlying formation mechanism remains to be identified[35]. To date, there is a lack of robust methods to generate SLBs with asymmetric distribution of PIP lipids on the top leaflet of the SLBs, facilitating quantitative studies of lipid-protein interactions.

In this study, we propose a new method for preparing asym-SLBs enriched in PIP$_2$ exclusively in the top leaflet of the SLBs. Inspired by a recently developed method involving PIP$_2$-containing vector SUVs[35], our approach takes advantage of lipids with high critical micelle concentrations (CMC). At concentrations below the CMC, lipids do not form micelles but remain as monomers. Specifically, in our method, PIP$_2$ is introduced in solution at concentrations below its CMC,

enabling its spontaneous insertion into pre-formed symmetric SLBs. This strategy is analogous to previous reports of spontaneous insertions of short chain ceramides and single-chain lysolipids into the outer leaflet of liposomes[51,52].

Our method is simple and robustly reproducible. It allows control over the concentration of the lipids of interest incorporated into asym-SLBs. The asym-SLBs allow us to study the dynamics of the lipids of interest free from any interaction with the substrate. We quantitatively characterized membrane asymmetry and fluidity using fluorescently labelled PIP_2, TopFluor-PIP_2 (TF-PIP_2). Our method allows the amount of TF-PIP_2 incorporation to be fine-tuned. We further demonstrated the utility of asymmetrically incorporating PIP_2 into SLBs by recruiting the PIP_2-associated actin-membrane linker ezrin, and myosin I motor, myosin 1b, which glides single actin filaments on the PIP_2-asym-SLBs. Taken together, the asym-SLBs developed in this study provide a robust experimental model system for future quantitative studies of lipid dynamics and lipid-protein interactions in vitro.

Materials and Methods

Reagents

1,2-Dioleoyl-sn-glycero-3-Phosphatidylcholine (DOPC, 850375), 1,2-dioleoyl-sn-glycero-3-phospho-L-serine (DOPS, 840035), L-α-phosphatidylinositol-4,5-bisphosphate (PIP_2, Brain, Porcine, 840046P), and 1-oleoyl-2-[6-[4-(dipyrrometheneboron difluoride)butanoyl]amino]-hexanoyl-sn-glycero-3-phosphoinositol-4,5-bisphosphate (TF-PIP_2, 810184P) were purchased from Avanti Polar Lipids. Alexa Fluor 488 C5-Maleimide (AX488) and Alexa Fluor 647 (AX647) phalloidin were purchased from Invitrogen. Creatinine phosphate was purchased from Roche

(10621714001) and creatinine phosphate kinase from Sigma- Aldrich (C3755-3.5KU). Other reagents were purchased from Sigma-Aldrich.

Buffer

The buffer used to prepare PIP_2 solutions and to wash the SLBs, referred to as the working buffer, contains 0.8 mM Sodium Phosphate, 15.3 mM Sodium Chloride, 0.3 mM Potassium Chloride, 0.2 mM Potassium Phosphate and 1 mM EDTA, pH 7.5. For PIP_2 clustering experiments, the 1 mM EDTA is replaced by 1 mM $CaCl_2$. MilliQ (MQ) water was used in all buffer preparations.

Preparation of SUVs

The desired amounts of lipids were mixed in $CHCl_3$ and then dried with N_2, followed by drying in a vacuum chamber overnight to ensure that the $CHCl_3$ was completely evaporated. The dried lipid film was hydrated with MQ H_2O for lipid mixtures without PIP_2, or the working buffer for lipid mixtures containing PIP_2, at a lipid concentration of 1.5 mg/mL and incubated at room temperature for 30 min to form vesicles. The resulting vesicle suspension was vortexed, followed by sonication in a bath sonicator (Fisher Scientific, ref. 10611983, at maximum power), twice for 30 min each, at room temperature, to generate SUVs. The resulting SUVs were then centrifuged at 20000g for 1 hour at 4°C to remove any lipid debris. The SUV supernatant was collected and stored at 4°C for a maximum of two weeks.

Preparation of PIP_2 solutions

The desired amount of PIP_2 in $CHCl_3$ was dried with N_2, followed by drying in a vacuum chamber for a few hours or overnight to ensure that the $CHCl_3$ was completely evaporated. The resulting lipid film was resuspended in the working buffer to obtain a final lipid concentration of

10 µM for PIP_2 and TF-PIP_2. The lipid solution was either stored at 4°C for a maximum of two weeks or at -20°C for several months.

Preparation of PIP_2-asym-SLBs

Glass coverslips were first cleaned by sonication in the following solutions for 20 min each: H_2O, Acetone, H_2O, 1M KOH and 5 times with H_2O. The coverslips were dried with N_2 and stored under vacuum before use. The experimental chamber was assembled from two cleaned coverslips glued together using double sided tapes, with the bottom coverslip plasma cleaned for 3 min to make it hydrophilic. SUVs composed of DOPC:DOPS at 9:1 molar ratio were prepared following the above-mentioned method. The chamber was filled with SUVs at 0.5 mg/mL, supplemented with 1 mM $CaCl_2$, followed by incubation at room temperature for 30 min to form SLBs. The resulting SLB was rinsed with 200 µL working buffer (~10 times the chamber volume). Then, the chamber was filled with PIP_2 or TF- PIP_2 solution for 10 min, followed by rinsing with 100 µL working buffer. The asym-SLB was used immediately for experiments.

Preparation of sym-SLBs containing TF-PIP_2

SUVs composed of DOPC:DOPS:TF-PIP_2 at 85:10:5 molar ratio were prepared following the above procedure, using the working buffer to rehydrate lipid films. An experimental chamber was filled with SUVs at 0.5 mg/mL, supplemented with 3 mM $CaCl_2$, followed by incubation at room temperature for 30 min to form SLBs. The resulting SLB was rinsed with 200 µL working buffer (~10 times the chamber volume). The SLB was used immediately for experiments.

FRAP experiments and data analysis

Confocal imaging was performed on an inverted microscope (Eclipse Ti-E, Nikon) equipped with a spinning disk CSU-X1 (Yokogawa), a sCmos camera Prime 95B (Photometrics) and a 60X

objective (NA 1.40, oil, Nikon). A laser bench of four lines (405nm, 491nm, 561nm, 642nm; Gataca Systems) was used for FRAP and acquisition at 491 nm. The fluorescent intensities at each time points were obtained using Metamorph software (Gataca Systems). For each image, the intensity of the bleached area was corrected with the intensity of an unbleached area, far from the bleached region, to account for photobleaching and potential variation of the intensity overtime. For each FRAP experiment, we performed FRAP with four bleaching areas with diameters of 9.2 μm, 16.5 μm, 23.8 μm and 31.2 μm. We performed three FRAP measurements for each area sizes. The analysis of the FRAP curves and the extraction of characteristic time and immobile fraction were performed as previously described[53]. The diffusion coefficient was obtained by fitting the diffusion coefficient using the following equation: $D = \frac{d^2}{16\tau}$, where the values of d and τ obtained from each individual FRAP measurement, and D is the diffusion coefficient, d is the diameter of the bleached area, and τ is the characteristic time of the FRAP recovery curve. (See supplementary figure 1)

Quantification of TF-PIP$_2$ incorporation into asym-SLBs

To quantify the incorporation of TF-PIP$_2$ into the top leaflet of asym-SLBs, we compared their fluorescent intensity with that of sym-SLBs. Sym-SLBs served as a reference because their TF-PIP$_2$ content is directly determined by the lipid composition of the SUVs used for their preparation. Specifically, sym-SLBs were formed from SUVs containing DOPC supplemented with 10 mol% DOPS, and with 0.1, 1, 2, 3 or 4 mol% TF-PIP$_2$. For each confocal image of SLBs, the mean fluorescent intensity was calculated within a 200 x 200 pixel region. These data were used to generate a calibration curve, yielding the slope $m_{sym-SLB}$ (intensity units per mol%) (Fig. S2). All imaging and analysis were performed under identical microscope settings, including laser power

and detector gain, for both sym-SLBs and asym-SLBs. The mole fration of TF-PIP$_2$ in the top leaflet of aysm-SLBs was then calculated as $f_{top} = 2 \times \frac{I-b}{m_{sym-SLB}}$, where I is the measured intensity of asym-SLBs and b is the background intensity. The factor of 2 accounts for the fact that in sym-SLBs TF-PIP$_2$ is present in both leaflets, whereas in asym-SLBs it is present only in the top leaflet.

Determination of the critical micelle concentration (CMC) of TF-PIP$_2$

Solutions of TF-PIP$_2$ in the working buffer are prepared at TF-PIP$_2$ concentrations ranging from 1 µM to 2 mM. The principle of the CMC determination is that monomers of TF-PIP$_2$ are not quenched and are thus visible in fluorescence spectrometry, whereas TF-PIP$_2$ in micelles is self-quenched. When TF-PIP$_2$ concentrations are increased below the CMC, the fluorescence intensities are thus expected to increase proportionally to the concentration but the fluorescence intensity will plateau as soon as the CMC is reached and additional TF-PIP$_2$ is incorporated in micelles. The fluorescence intensities of TF-PIP$_2$ were measured with a Tecan microplate spectrophotometer.

Thermodynamic model of TF-PIP$_2$ incorporation in asym-SLBs

The principle of the thermodynamic model is the thermal equilibrium between the three different forms of TF-PIP$_2$ (monomer in solution, in a micelle in solution, inserted in the SLB):

$$\mu_S^0 + k_B T ln \left(\frac{c_S}{c_{H2O}} \right) = \mu_M^0 + \frac{kT}{N} ln \left(\frac{c_M}{N} \frac{1}{c_{H2O}} \right) = \mu_B^0 + kT ln \left(\frac{s_B}{s_L} \right),$$

μ_S^0, μ_M^0, and μ_B^0 are the standard chemical potentials of TF-PIP$_2$ in solution (s), micelles (M), and asym-SLBs (B), respectively. c_S and c_M are the bulk concentrations of TF-PIP$_2$ in solution and in micelles, $c_{H2O} = 55.6\ M$ is the water concentration. s_B and s_L are the surface densities of TF-PIP$_2$ and lipids in asym-SLBs, respectively ($s_L = 1.3 \times 10^{18}$ molecules/m^2 if the surface per lipid is

0.7 nm^2). N is the number of TF-PIP$_2$ molecules per micelle; this number is not known but its exact value has no impact here provided that the monomer concentration remains below CMC and that N>~10.

The total TF-PIP$_2$ concentration is computed by taking into account the TF-PIP$_2$ fraction in the SLB, knowing that our sample has a volume of 20 µL and our SLB covers 2.6 cm^2.

The model must reproduce: (1) the proportionality between the proportion of TF-PIP$_2$ in asym-SLB (%) and the total TF-PIP$_2$ concentration (µM) when the proportion is below 5% (no self-quenching), and (2) the TF-PIP$_2$ CMC = 290 µM. A good fit is obtained with: $\mu_M^0 = 0 \, kT$ (our reference), $\mu_B^0 = -0.6 \, kT$ and $\mu_S^0 = 12.2 \, kT$ (Fig. 3, blue line). From these standard chemical potentials, the TF-PIP$_2$ partition coefficient between the membrane and the solution can be

calculated as $K = \dfrac{\frac{s_B}{s_L}}{c_S} = \dfrac{1}{c_{H2O}} \times e^{\frac{\mu_S^0 - \mu_B^0}{kT}} = 5.9 \times 10^3 M^{-1}$.

TF-PIP$_2$ as monomer in solution (µM)	TF-PIP$_2$ in micelles in solution (µM)	Proportion of TF-PIP$_2$ in asym-SLB (%)	Total TF-PIP$_2$ concentration (µM)
2	$< 10^{-3}$	1.2	2.4
4.7	$< 10^{-3}$	2.8	5.6
6.5	$< 10^{-3}$	3.8	7.7
8.5	$< 10^{-3}$	5.0	10.1
15	$< 10^{-3}$	8.9*	17.8

Table 1. Thermodynamic model prediction of TF-PIP$_2$ in asym-SLBs. *at the bulk concentration of 15 µM, TF-PIP$_2$ signal is quenched.

Measurement of TF-PIP₂ self-quenching

To monitor the conditions of TF-PIP$_2$ self-quenching, we prepared liposomes in water at 1 mM POPC final concentration with various TF-PIP$_2$ concentrations, so that the proportion of TF-PIP$_2$ in liposome membranes varies from 0% to 10%. To obtain the right TF-PIP$_2$ proportion, we have taken into account the partition of TF-PIP$_2$ between membranes and solution according to the partition coefficient we have determined ($K = 5.9 \times 10^3 M^{-1}$), which leads to adding ~20% more TF-PIP$_2$ than if it is fully embedded in liposome membranes. The fluorescence intensities of the liposome samples were measured with a Tecan microplate spectrophotometer.

As shown in Fig. S3, we observed that the fluorescence intensities of TF-PIP$_2$ increase proportionally to the TF-PIP$_2$ percentage in the liposome membrane up to 5% TF-PIP$_2$. Above 5%, the fluorescence intensity remains approximately constant, which can be explained by TF-PIP$_2$ self-quenching.

Myosin 1b, actin and ezrin purification

Human myosin 1b was purified from HEK293-Flp-In cells following the method described previously[54]. Myo1b was stored in the following buffer: 30 mM Hepes pH 7.5, 150 mM KCl, 4 mM MgCl$_2$, 1 mM EGTA, 0.1% Methylcellulose, 4 mM Na$_2$ATP, and 1 mM DTT. Muscle actin was purified from rabbit muscle and isolated in monomeric form in actin buffer (5 mM Tris-HCl, pH 7.8, 0.1 mM CaCl$_2$, 0.2 mM ATP, 1 mM DTT and 0.01% NaN$_3$)[55]. Phosphomimetic ezrin T567D (ezrin TD) was purified and labelled with AX488 as previously described[56]. EzrinTD was stored in 20 mM Tris pH 7.4, 50 mM KCl, 0.1 mM EDTA, 2 mM β-Mercaptoethanol, and supplemented with 0.1% methylcellulose.

Preparation and analysis of ezrin-actin binding assays

To evaluate ezrin binding and actin recruitment on PIP$_2$-asym-SLBs, base SLBs were prepared by the fusion of SUVs composed of DOPC: DOPS at a 9:1 molar ratio and supplemented with 0.5 mol% Texas Red-DHPE for membrane visualization. SUVs were prepared as described above and introduced into the chamber at a final lipid concentration of 0.5 mg/mL. The chamber was incubated with the SUVs for 30 min at room temperature, protected from light. SLBs were then gently washed with 200 µL of the working buffer to remove excess vesicles.

After the formation of the base SLB, 20 µL of a 10 µM PIP$_2$ solution was added directly to the chamber, followed by incubating with the SLB for at least 10 min. The chamber was subsequently washed with ~100 µL of the working buffer. For ezrin recruitment assays, phosphomimetic ezrin T567D (ezrin TD) labeled with AX488 was prepared at 10 µM in 20 mM HEPES pH 7.4, 50 mM KCl, and 0.1 mM EDTA. The protein solution (20 µL) was added to the chamber and incubated for 10 min. Next, pre-polymerized actin filaments labeled with AX647-tagged phalloidin were introduced in the chamber, followed by incubating for 10 min at room temperature. The chamber was sealed and prepared for imaging. Control experiments were performed using base SLBs, without PIP$_2$ incorporation.

Actin filaments were prepared by polymerizing globular actin monomers at 5 µM, then diluted to 1 µM in the actin polymerization buffer (5 mM Tris HCl pH 8.0, 3 mM MgCl$_2$, 0.2 mM EGTA, 1 mM DABCO, 10 mM DTT, 100 mM KCl, 2 mM ATP), supplemented with AX647-taggd phalloidin at 330 nM.

Samples were imaged using a spinning-disk confocal microscope: Inverted Eclipse Ti-E (Nikon) equipped with a CSU-X1 spinning disk (Yokogawa), controlled by Metamorph software (Gataca Systems), and a 100x oil objective. Imaging was performed using appropriate laser lines and filters

for Texas Red, AX488, and AX647. The pixel-averaged intensity of ezrin fluorescence was calculated from a 500×500 pixel-sizes for each image.

Preparation and analysis of motor motility assays

The actin motility assays were performed and analyzed following the method described previously[57]. Briefly, myo1b at 600 nM was incubated with PIP$_2$-asym-SLBs for 30 min at room temperature. Then, a 50 nM solution of pre-polymerized actin filaments (in the actin polymerization buffer, supplemented with 0.3% methylcellulose and the ATP regeneration system) stabilized with AX647 phalloidin at 165 nM was introduced into the experimental chambers. The ATP regeneration system is composed of 2 mM Na$_2$ATP, 2 mM MgCl$_2$, 20 mM Creatinine phosphate and 3.5 U/mL of Creatinine phosphate kinase.

The samples were observed using TIRF microscopy (Eclipse Ti inverted microscope, 100X TIRF objective, Quantem512SC camera). Kymographs of gliding actin filaments were generated using the ImageJ plug-in KymoToolBox.

Results and Discussions

The generation of asymmetric SLBs with (PIP$_2$-asym-SLBs)

A wide range of CMC values of PIP$_2$ has been reported. For unlabeled PIP$_2$, the CMC is on the order of a few μM in 100 mM NaCl, 30 μM in 50 mM PIPES or Tris pH 7, and 200 μM in water[58–60]. For BODIPY® TMR PIP$_2$ in water, the CMC value was estimated to be up to 500 μM[58–60]. We hypothesized that dissolving PIP$_2$ at a relatively high concentration in solution, but below its CMC, could facilitate its insertion into the top leaflet of pre-formed SLBs. This spontaneous lipid insertion is due to the less favorable hydrophobic interaction of the fatty acid chains of PIP$_2$ with the solution than with the lipid membrane. We first determined the CMC of TF-PIP$_2$ in the

Working buffer (0.8 mM Na_3PO_4, 15.3 mM NaCl, 0.3 mM KCl, 0.2 mM KH_2PO_4, and 1 mM EDTA, pH 7.5), as used previously[35]. We prepared TF-PIP$_2$ solutions at concentrations ranging from 1 µM to 2 mM. The principle of CMC determination is that monomeric TF-PIP$_2$ is not quenched and its fluorescent signal is detectable in fluorescence spectrometry, while TF-PIP$_2$ in micelles is self-quenched. Thus, TF-PIP$_2$ fluorescence intensity increases linearly with concentrations below the CMC, then plateaus once the CMC is reached due to micelle formation. As shown in Fig. 1A, the CMC of TF-PIP$_2$ was determined to be 290 µM in the working buffer.

To test our hypothesis of TF-PIP$_2$ spontaneous insertion into SLBs, we first prepared a SLB containing DOPC and DOPS (9:1 molar ratio) on a glass substrate using the well-established SUV fusion method[61]. Then, a solution of TF-PIP$_2$ at a concentration below its CMC, at 5 µM, was introduced into the experimental chamber. We incubated the base SLB with TF-PIP$_2$ for 10 min, allowing it to insert into the top leaflet of the base SLB (Fig. 1B). The resulting asym-SLB was rinsed with the working buffer to remove excess TF-PIP$_2$ remaining in the solution. Using 5 µM of TF-PIP$_2$ in solution produced asym-SLBs with relatively homogenous fluorescence, demonstrating the proper incorporation of TF-PIP$_2$ on the SLBs (Fig. 2A *Top panel*).

Figure 1. Proposed mechanism for PIP₂ asym-SLB formation. *(A) Determination of the CMC of TF-PIP₂ based on fluorescence intensities measurements (blue symbols. See Materials and Methods for details.). Dotted line is the linear fit of the measurements. Dashed line represents the plateau value of the intensities at high TF-PIP₂ concentration. The CMC, 290 μM, is indicated in red. (B) Schematic representation of PIP₂ asym-SLB formation. Step 1: Formation of a base SLBs composed of DOPC and DOPS via SUV fusion on a glass substrate. Step 2: Introduction of a PIP₂ solution at a concentration below CMC, enabling its spontaneous insertion into the top leaflet of the SLB and resulting in an asymmetric TF-PIP₂ distribution. Green-headed lipids represent PIP₂.*

Quantitative characterization of PIP₂-asym-SLBs

To quantitatively characterize the resulting TF-PIP₂-asym-SLBs, we performed Fluorescence Recovery After Photobleaching (FRAP) experiments to measure the diffusion coefficients of TF-PIP₂ and its immobile fractions within the asym-SLBs (Fig. 2A, Asym-SLB). For freshly prepared asym-SLBs, within 1 hour after preparation, we found the immobile fractions of 2.8 ± 1.7 % (Fig.

2B, Asym-SLB, 1h). By analyzing the FRAP curves (Fig. S1), we found that the diffusion coefficients of TF-PIP$_2$ in these asym-SLBs were around 2 ± 0.4 μm^2/sec (Fig. 2C, Asym-SLB, 1h), which are comparable to previously reported values for SLBs[34–36]. The initial residual immobile fraction in TF-PIP$_2$ asym-SLBs may be due to the presence of unfused SUVs in the pre-formed SLBs, as previously shown by scanning electron microscopy[62]. Notably, by unzipping lipid monolayers from SLBs, a recent study demonstrated that PIP$_2$ can translocate from the top to the bottom leaflet at the edges of bilayer defects in sym-SLBs, contributing to the immobilized fraction[36]. Moreover, it was reported that the translocation of PIP$_2$ from the top to the bottom leaflet at bilayer defects sites results in a progressive increase in the PIP$_2$ immobile fraction over time[36]. Here, we compared the immobile fraction of a freshly prepared asym-SLB with those of asym-SLBs a few hours after formation. Indeed, we found the mean immobile fractions increased after formation, up to 9% after 5 hours (Fig. 2B, Asym-SLB, 2h-3h and 3h-5h), while there is no significant difference for the diffusion coefficients, ranging between $1.5 - 3$ μm^2/sec (Fig. 2C, Asym-SLB, pink bars, p=0.1 one way ANOVA test). These results are consistent with the previous report[36], and suggest that, under our experimental conditions, TF-PIP$_2$ in asym-SLBs undergoes a modest degree of translocation over the timescale of the experiments.

We compared the immobile fractions and diffusion coefficients of TF-PIP$_2$-asym-SLBs with those measured in sym-SLBs containing TF-PIP$_2$. We prepared the symmetric SLBs using SUVs composed of DOPC:DOPS:TF-PIP$_2$ at 88:10:2 molar ratio. Given the known tendency of PIP$_2$ to remain immobile when in contact with a glass substrate[31,35,36], the measured immobile fraction of PIP$_2$ serves as an indicator of its asymmetric distribution in the two leaflets of the SLB. The time series of the FRAP images showed a clear difference between the sym- and asym-SLBs (Fig. 2A).

While the asym-SLB exhibited an immobile fraction of less than 20% (Fig. 2B, Asym-SLB, pink data points), the sym-SLB showed much less fluorescence recovery, reflecting a higher immobile fraction of 75% on average (Fig. 2B, Sym-SLB, blue data points). In the sym-SLB, the higher immobile fraction of TF-PIP$_2$ can be partly attributed to the fact that 50% of the TF-PIP is in direct contact with the glass substrate, resulting in its immobilization, as previously observed[33–36,63]. For the mobile fraction of TF-PIP$_2$, we found similar diffusion coefficients in both asym- and sym-SLBs (Fig. 2C, compare pink and blue bars). This similarity probably arises from the fact that mobile PIP$_2$ lipids are only present in the top leaflet of both asym- and sym- SLBs. Overall, we showed here that the proposed method based on the spontaneous insertion of PIP$_2$ into pre-formed sym-SLBs allows us to generate asym-SLBs with preserved PIP$_2$ fluidity.

Divalent ions such as calcium and magnesium are known to cluster PIP$_2$ via electrostatic interactions, which can affect the immobile fraction and potentially alter the diffusion of PIP$_2$[12–14,64,65]. To evaluate how well PIP$_2$ in the asym-SLBs retains its expected properties, such as its ability to diffuse or form clusters in the presence Ca^{2+} ions, we quantified the effect of Ca^{2+} on the diffusion and immobile fractions of TF-PIP$_2$. By performing FRAP experiments, we observed an increase in the immobile fraction of TF-PIP$_2$ in the presence of 1 mM of Ca^{2+} ions compared to that without Ca^{2+} ions (Fig. 2B, green data points). The increased immobile fraction indicates PIP$_2$ molecules were clustered by Ca^{2+} ions. On the other hand, the diffusion coefficient of mobile TF-PIP$_2$ was only slightly affected by the presence of Ca^{2+} ions (Fig. 2C, green bars). This observation is consistent with a previous study using FCS reported a slight decrease in the diffusion of TF-PIP$_2$ monomers and TF-PIP$_2$ clustering in GUVs due to the presence of Ca^{2+} ions[11]. Our results indicated that in the presence of Ca^{2+}, the freely diffusing TF-PIP$_2$ was not significantly affected

by the presence of TF-PIP$_2$ clusters and possibly the kinetics of clustering and de-clustering is much slower than the timescale used for measuring the FRAP recoveries.

***Figure 2. Quantitative comparisons of sym- and asym-SLBs for TF-PIP$_2$, and the effect of Ca^{2+} in asym-SLBs. (A)** Left: Representative confocal images of FRAP experiments on TF-PIP$_2$ in sym- and asym-SLBs before bleaching (t<0), immediately after bleaching (t=0) and after recovery (t=245 sec). Scale bars, 10μm. Right: Representative relative fluorescent intensities of the bleached area overtime. By analyzing the FRAP curves, we found the recovery of ~98% and ~30% and the diffusion coefficient of 2 μm^2/sec and 1.5 μm^2/sec of TF-PIP$_2$ in the asym-SLB (Magenta*

*circles) and sym-SLB (Cyan circles), respectively. **(B)** Quantification of the immobile fractions of TF-PIP$_2$ in FRAP experiments performed within durations of one hour (1h), 2h-3h and 3h-5h after bilayer formation. N = 3 independent experiments, except "asym-SLB 2h-3h" with N = 6, for each condition, in which n = 34, 71, 36, 36, 36, 36 FRAP measurements (from left to right). The immobile fractions were calculated by data fitting to the FRAP curves. **(C)** Diffusion coefficients of TF-PIP$_2$ in asym- and sym-SLB. For the asym-SLBs, TF-PIP$_2$ bulk concentration was 5 μM. Each data point represents one independent FRAP experiment. Within each experiment, there were 4 different bleaching area sizes and 3 FRAP measurements were performed for each size. Thus, a total of 12 FRAP measurements in one FRAP experiment. Diffusion coefficients for each experiment were obtained from the linear fit of bleached areas to their recovery characteristic times (as shown in Fig. S1). The error bar represents the standard deviation. Statistic test, Mann-Whitney test, for conditions of Asym-SLB 3h-5h and Asym-SLB + Ca^{2+} 3h-5h, p = 0.1. In (B) and (C), for experiments with Ca^{2+}, Asym-SLB + Ca^{2+}, in the working buffer, 1 mM EDTA was replaced by 1 mM CaCl$_2$.*

Control of the PIP$_2$ amount incorporated into the top leaflet of pre-formed SLBs

To quantify the amount of PIP$_2$ incorporated into the top leaflet of the SLBs, we measured the fluorescent intensity of TF-PIP$_2$ in asym-SLBs. We calibrated our measurements using TF-PIP$_2$ fluorescence measurements on sym-SLBs since the concentrations of TF-PIP$_2$ incorporated into the SUVs used to generate the sym-SLBs is known (see Method section for details). We found that the amount of TF-PIP$_2$ incorporated into the top leaflet of asym-SLBs was directly proportional to the bulk concentration of TF-PIP$_2$, up to 10 μM of TF-PIP$_2$ (Fig. 3). The use of TF-PIP$_2$ at bulk concentrations ranging from 0 to 10 μM resulted in 0 to 5 mol% TF-PIP$_2$ in asym-SLBs (Fig. 3).

We observed that beyond 10 μM of TF-PIP$_2$, the intensity of TF-PIP$_2$ did not increase significantly. This observation suggests that either the asym-SLBs were saturated with TF-PIP$_2$, or the concentration of TF-PIP$_2$ was high enough for the TF dyes to quench each other[12,66]. To test for self-quenching, we prepared vesicles containing 0 mol% to 10 mol% TF-PIP$_2$. We observed self-quenching when the TF-PIP$_2$ fraction above 5 mol% (Fig. S3), supporting the hypothesis that fluorescence saturation at higher concentrations is due, at least in part, to quenching effects.

To quantitatively explain our observations of TF-PIP$_2$ incorporation into SLBs or micelles, we implemented a thermodynamic model of the equilibrium between monomeric and micellar TF-PIP$_2$ in solution and TF-PIP$_2$ inserted into the SLB (see method section). Using our experimentally observed proportionality between TF-PIP$_2$ incorporation in SLBs and bulk concentration below 5 mol% of TF-PIP$_2$, we used our thermodynamic model to calculate the TF-PIP$_2$ partition coefficient between the membrane and the solution to be $5.9 \times 10^3 M^{-1}$.

Figure 3. Quantification of TF-PIP₂ incorporation into asym-SLBs. *Quantification of the molar percentages of TF-PIP₂ in the top leaflet of asym-SLB generated by using different bulk concentrations of TF-PIP₂. The blue line is the prediction of the thermodynamic model for all the data points < 10µM, slope = 0.53. n = 568 measurements, N = 7 independent experiments.*

The activity of PIP₂-associated ezrin and molecular motor myosin 1b on PIP₂-asym-SLBs

Actin-membrane linkers often contain domains that specifically interact with PIP₂. To further validate the use of PIP₂-asym-SLBs, we examined the recruitment of ezrin, a well-characterized actin-membrane linker[18]. Ezrin contains an N-terminal FREM (band 4.1, ezrin, radixin, moesin) domain that binds specifically to PIP₂ and a C-terminal ERM associated domain that interacts with actin filaments. We assessed both ezrin recruitment to PIP₂-asym-SLBs and its ability of anchoring actin filaments to the bilayers. We incubated PIP₂-asym-SLBs with 10 µM AX488-labelled phosphomimetic ezrin T567D (ezrin TD), which adopts an open conformation that allows

simultaneous interactions with PIP_2 and actin. This was followed by the addition of 1 μM AX647-phalloidin-labelled actin filaments. In the absence of PIP_2, neither ezrinTD nor actin bound to the SLBs (Fig. 4A). However, when PIP_2 was present, both ezrin and actin filaments were clearly recruited to the membrane (Fig. 4, A and B). Additionally, we investigated the activity of the class I myosin motor, myo1b, which contains a C-terminal tail pleckstrin homology (PH) domain that binds PIP_2 and an N-termial motor domain that binds actin filaments and generates force[67]. We incubated PIP_2-asym-SLBs with 1 μM myo1b and 50 nM of AX647-phalloidin labelled actin filaments to perform motility assays[68]. We observed the gliding of the actin filaments on asym-SLB containing TF-PIP_2 or non-fluorescently labelled PIP_2, showing myo1b activity can be reconstituted in this system (Fig. 4C). The resulting actin gliding velocities are 57 nm/sec for TF-PIP_2 and 62 nm/sec for non-fluorescently labelled PIP_2, both comparable to those previously measured for myo1b on PIP_2-sym-SLBs (Fig. 4D)[57].

Figure 4. Actin filament recruitments on PIP₂-asym-SLBs by actin-membrane linkers, ezrin

and myo1 motors. *(A) Representative confocal images of AX488 labelled ezrinTD and AX647*

phalloidin-labelled actin filaments on SLBs without PIP₂ (Top) and PIP₂-asym-SLBs (Bottom).

From Left to Right panel: Membrane, Ezrin and Actin. Base SLBs were composed of DOPC:

DOPS at a 9:1 molar ratio and supplemented with 0.5 mol% Texas Red-DHPE for membrane

visualization. The asym-SLBs were generated by incubating the base SLBs with 10 μM PIP₂

solution. Scale bars, 10 μm. (B) Quantification of ezrin intensity on SLBs without PIP₂ and PIP₂-

asym-SLBs. N = 3 independent experiments. Analyzed images: 60 for SLBs without PIP$_2$ and 44 for PIP$_2$-asym- SLBs. **(C)** *Representative kymographs of actin filament gliding by myo1b bound on asym-SLBs with PIP$_2$ or TF-PIP$_2$. Base SLBs were incubated with 5 μM PIP$_2$ or TF-PIP$_2$ solutions. Horizontal scale bars, 5 μm, Vertical scale bars, 1 min. Fluorescent singles were AX647 phalloidin labelled actin filaments.* **(D)** *Velocities of single actin filament gliding by myo1b bound on asym-SLBs with TF-PIP$_2$ or PIP$_2$. The mean velocities and the corresponding standard deviations of myo1b-driven actin filament gliding are indicated in the figure. Welch's t test, p = 0.1119. For TF-PIP$_2$, n=62 filaments measured and N=3 independent experiments. For PIP$_2$, n=60 filaments measured, N=2 independent experiments.*

Conclusion

In this study, we introduced a simple two-step method to generate supported bilayers containing leaflets with different lipid compositions. We demonstrated the feasibility of the method to enrich one leaflet with PIP$_2$ and fluorescent TF-PIP$_2$. We also quantitatively characterized the fluidity and asymmetry of the SLBs. We demonstrated the ability to fine-tune the amount of the asymmetrically incorporated lipids by incubating the SLB with varying concentrations of TF-PIP$_2$.

Our results suggest the potential application of this method to incorporate other lipids with sufficiently high CMC values, such as the other members of the phosphoinositide family and fluorescently labelled lipids[69,70]. While our study focused on the incorporation of lipids into SLBs, the possibility of adapting the method to other model membranes such as giant unilamellar vesicles (GUVs), large unilamellar vesicles (LUVs), and suspended planar lipid bilayers, or to the plasma membrane of cells, could open the potential applications to studies involving asymmetric membranes.

Supporting information

Additional experimental data plots (PDF).

Acknowledgment

We thank Frederic Pincet (ENS, Paris) for his help with the FRAP analysis and careful reading of the manuscript. We thank Patricia Bassereau and Julien Pernier for fruitful discussions. J.H, N.R., S.C. and F.-C.T. is a member of the CNRS consortium Approches Quantitatives du Vivant (AQV), Labex Cell(n)Scale (ANR-11-889 LABX0038) and Paris Sciences et Lettres (ANR-10-IDEX-0001-02). G.G. and F.-C.T. are funded by ANR grant ANR-20-CE11-0010-01 (ActinFission). G.G. and F.-C.T., J.H., S.C., N.R., and J.B. are funded by a PSL-Qlife grant (MyoMemActin). JB is funded by grant R01GM115636 from NIH. T.L.A.N. is funded by École Doctorale Physique en Île-de-France (EDFPI) PhD fellowship. The authors greatly acknowledge the Cell and Tissue Imaging (PICT-IBiSA), Institut Curie, member of the French National Research Infractucture France-BioImaging (ANR10-INBS-04).

Reference

(1)   Lorent, J. H.; Levental, K. R.; Ganesan, L.; Rivera-Longsworth, G.; Sezgin, E.; Doktorova, M.; Lyman, E.; Levental, I. Plasma Membranes Are Asymmetric in Lipid Unsaturation, Packing and Protein Shape. *Nat. Chem. Biol.* **2020**, *16* (6), 644–652. https://doi.org/10.1038/s41589-020-0529-6.

(2)   Vance, J. E. Phospholipid Synthesis and Transport in Mammalian Cells. *Traffic* **2015**, *16* (1), 1–18. https://doi.org/10.1111/tra.12230.

(3)   Posor, Y.; Jang, W.; Haucke, V. Phosphoinositides as Membrane Organizers. *Nat. Rev. Mol. Cell Biol.* **2022**, *23* (12), 797–816. https://doi.org/10.1038/s41580-022-00490-x.

(4)   Janmey, P. A.; Bucki, R.; Radhakrishnan, R. Regulation of Actin Assembly by PI (4, 5) P2 and Other Inositol Phospholipids: An Update on Possible Mechanisms. *Biochem. Biophys. Res. Commun.* **2018**, *506* (2), 307–314.

(5)   Wills, R. C.; Hammond, G. R. V. PI(4,5)P2: Signaling the Plasma Membrane. **2022**, *0*, 2311–2325.

(6) Lemmon, M. A. Membrane Recognition by Phospholipid-Binding Domains. *Nat. Rev. Mol. Cell Biol.* **2008**, *9* (2), 99–111. https://doi.org/10.1038/nrm2328.

(7) Chandra, M.; Chin, Y. K.-Y.; Mas, C.; Feathers, J. R.; Paul, B.; Datta, S.; Chen, K.-E.; Jia, X.; Yang, Z.; Norwood, S. J.; Mohanty, B.; Bugarcic, A.; Teasdale, R. D.; Henne, W. M.; Mobli, M.; Collins, B. M. Classification of the Human Phox Homology (PX) Domains Based on Their Phosphoinositide Binding Specificities. *Nat. Commun.* **2019**, *10* (1), 1528. https://doi.org/10.1038/s41467-019-09355-y.

(8) Carman, P. J.; Dominguez, R. BAR Domain Proteins-a Linkage between Cellular Membranes, Signaling Pathways, and the Actin Cytoskeleton. *Biophys. Rev.* **2018**, *10* (6), 1587–1604. https://doi.org/10.1007/s12551-018-0467-7.

(9) Suetsugu, S.; Kurisu, S.; Takenawa, T. Dynamic Shaping of Cellular Membranes by Phospholipids and Membrane-Deforming Proteins. *Physiol. Rev.* **2014**, *94* (4), 1219–1248. https://doi.org/10.1152/physrev.00040.2013.

(10) Senju, Y.; Lappalainen, P. Regulation of Actin Dynamics by PI (4, 5) P2 in Cell Migration and Endocytosis. *Curr. Opin. Cell Biol.* **2019**, *56*, 7–13.

(11) Sarmento, M. J.; Coutinho, A.; Fedorov, A.; Prieto, M.; Fernandes, F. Ca2 + Induces PI(4,5)P2 Clusters on Lipid Bilayers at Physiological PI(4,5)P2 and Ca2 + Concentrations. *Biochim. Biophys. Acta - Biomembr.* **2014**, *1838* (3), 822–830. https://doi.org/10.1016/j.bbamem.2013.11.020.

(12) Wen, Y.; Vogt, V. M.; Feigenson, G. W. Multivalent Cation-Bridged PI(4,5)P2 Clusters Form at Very Low Concentrations. *Biophys. J.* **2018**, *114* (11), 2630–2639. https://doi.org/10.1016/j.bpj.2018.04.048.

(13) Wang, Y. H.; Collins, A.; Guo, L.; Smith-Dupont, K. B.; Gai, F.; Svitkina, T.; Janmey, P. A. Divalent Cation-Induced Cluster Formation by Polyphosphoinositides in Model Membranes. *J. Am. Chem. Soc.* **2012**, *134* (7), 3387–3395. https://doi.org/10.1021/ja208640t.

(14) Yu-Hsiu Wang, D. R. S. and P. A. J. Counterion-Mediated Cluster Formation by Polyphosphoinositides. *Chem Phys Lipids* **2014**, *182* (1), 38–51. https://doi.org/10.1016/j.chemphyslip.2014.01.001.Counterion-mediated.

(15) Bezanilla, M.; Gladfelter, A. S.; Kovar, D. R.; Lee, W.-L. Cytoskeletal Dynamics: A View from the Membrane. *J. Cell Biol.* **2015**, *209* (3), 329–337. https://doi.org/10.1083/jcb.201502062.

(16) Tsai, F.-C.; Guérin, G.; Pernier, J.; Bassereau, P. Actin-Membrane Linkers: Insights from Synthetic Reconstituted Systems. *Eur. J. Cell Biol.* **2024**, *103* (2), 151402. https://doi.org/10.1016/j.ejcb.2024.151402.

(17) Liebe, N. L.; Mey, I.; Vuong, L.; Shikho, F.; Geil, B.; Janshoff, A.; Steinem, C. Bioinspired Membrane Interfaces: Controlling Actomyosin Architecture and Contractility. *ACS Appl. Mater. Interfaces* **2023**, *15* (9), 11586–11598. https://doi.org/10.1021/acsami.3c00061.

(18) McClatchey, A. I. ERM Proteins at a Glance. *J. Cell Sci.* **2014**, *127* (15), 3199–3204. https://doi.org/10.1242/jcs.098343.

(19) Maniti, O.; Carvalho, K.; Picart, C. Model Membranes to Shed Light on the Biochemical and Physical Properties of Ezrin/Radixin/Moesin. *Biochimie* **2013**, *95* (1), 3–11. https://doi.org/10.1016/j.biochi.2012.09.033.

(20) Morandat, S.; Azouzi, S.; Beauvais, E.; Mastouri, A.; El Kirat, K. Atomic Force Microscopy of Model Lipid Membranes. *Anal. Bioanal. Chem.* **2013**, *405* (5), 1445–1461. https://doi.org/10.1007/s00216-012-6383-y.

(21) Oreopoulos, J.; Yip, C. M. Probing Membrane Order and Topography in Supported Lipid Bilayers by Combined Polarized Total Internal Reflection Fluorescence-Atomic Force Microscopy. *Biophys. J.* **2009**, *96* (5), 1970–1984. https://doi.org/10.1016/j.bpj.2008.11.041.

(22) Matysik, A.; Kraut, R. S. Preparation of Mica Supported Lipid Bilayers for High Resolution Optical Microscopy Imaging. *J. Vis. Exp. JoVE* **2014**, No. 88, 52054. https://doi.org/10.3791/52054.

(23) Niederauer, C.; Nguyen, C.; Wang-Henderson, M.; Stein, J.; Strauss, S.; Cumberworth, A.; Stehr, F.; Jungmann, R.; Schwille, P.; Ganzinger, K. A. Dual-Color DNA-PAINT Single-Particle Tracking Enables Extended Studies of Membrane Protein Interactions. *Nat. Commun.* **2023**, *14* (1), 4345. https://doi.org/10.1038/s41467-023-40065-8.

(24) Honigmann, A.; Mueller, V.; Hell, S. W.; Eggeling, C. STED Microscopy Detects and Quantifies Liquid Phase Separation in Lipid Membranes Using a New Far-Red Emitting Fluorescent Phosphoglycerolipid Analogue. *Faraday Discuss* **2013**, *161*, 77–89. https://doi.org/10.1039/C2FD20107K.

(25) Ferhan, A. R.; Yoon, B. K.; Park, S.; Sut, T. N.; Chin, H.; Park, J. H.; Jackman, J. A.; Cho, N.-J. Solvent-Assisted Preparation of Supported Lipid Bilayers. *Nat. Protoc.* **2019**, *14* (7), 2091–2118. https://doi.org/10.1038/s41596-019-0174-2.

(26) Kurniawan, J.; Ventrici de Souza, J. F.; Dang, A. T.; Liu, G.; Kuhl, T. L. Preparation and Characterization of Solid-Supported Lipid Bilayers Formed by Langmuir–Blodgett Deposition: A Tutorial. *Langmuir* **2018**, *34* (51), 15622–15639. https://doi.org/10.1021/acs.langmuir.8b03504.

(27) Lind, T. K.; Cárdenas, M. Understanding the Formation of Supported Lipid Bilayers via Vesicle Fusion-A Case That Exemplifies the Need for the Complementary Method Approach (Review). *Biointerphases* **2016**, *11* (2), 020801. https://doi.org/10.1116/1.4944830.

(28) Richter, R. P.; Bérat, R.; Brisson, A. R. Formation of Solid-Supported Lipid Bilayers: An Integrated View. *Langmuir* **2006**, *22* (8), 3497–3505. https://doi.org/10.1021/la052687c.

(29) Goodchild, J. A.; Walsh, D. L.; Connell, S. D. Nanoscale Substrate Roughness Hinders Domain Formation in Supported Lipid Bilayers. *Langmuir* **2019**, *35* (47), 15352–15363. https://doi.org/10.1021/acs.langmuir.9b01990.

(30) Scomparin, C.; Lecuyer, S.; Ferreira, M.; Charitat, T.; Tinland, B. Diffusion in Supported Lipid Bilayers: Influence of Substrate and Preparation Technique on the Internal Dynamics. *Eur. Phys. J. E* **2009**, *28* (2), 211–220. https://doi.org/10.1140/epje/i2008-10407-3.

(31) Schäfer, J.; Nehls, J.; Schön, M.; Mey, I.; Steinem, C. Leaflet-Dependent Distribution of PtdIns[4,5]P2 in Supported Model Membranes. *Langmuir* **2020**, *36* (5), 1320–1328. https://doi.org/10.1021/acs.langmuir.9b03793.

(32) Guo, L.; Har, J. Y.; Sankaran, J.; Hong, Y.; Kannan, B.; Wohland, T. Molecular Diffusion Measurement in Lipid Bilayers over Wide Concentration Ranges: A Comparative Study. *ChemPhysChem* **2008**, *9* (5), 721–728. https://doi.org/10.1002/cphc.200700611.

(33) Baumann, M. K.; Amstad, E.; Mashaghi, A.; Textor, M.; Reimhult, E. Characterization of Supported Lipid Bilayers Incorporating and Phosphoinositol-3,4,5-Triphosphate by Complementary Techniques. *Biointerphases* **2010**, *5* (4), 114–119. https://doi.org/10.1116/1.3516485.

(34) Braunger, J. A.; Kramer, C.; Morick, D.; Steinem, C. Solid Supported Membranes Doped with PIP2: Influence of Ionic Strength and pH on Bilayer Formation and Membrane Organization. *Langmuir ACS J. Surf. Colloids* **2013**, *29* (46), 14204–14213.

(35) Shi, X.; Kohram, M.; Zhuang, X.; Smith, A. W. Interactions and Translational Dynamics of Phosphatidylinositol Bisphosphate (PIP2) Lipids in Asymmetric Lipid Bilayers. *Langmuir* **2016**, *32* (7), 1732–1741. https://doi.org/10.1021/acs.langmuir.5b02814.

(36) Sun, S.; Liu, C.; Rodriguez Melendez, D.; Yang, T.; Cremer, P. S. Immobilization of Phosphatidylinositides Revealed by Bilayer Leaflet Decoupling. *J. Am. Chem. Soc.* **2020**, *142* (30), 13003–13010. https://doi.org/10.1021/jacs.0c03800.

(37) Blosser, M. C.; Honerkamp-Smith, A. R.; Han, T.; Haataja, M.; Keller, S. L. Transbilayer Colocalization of Lipid Domains Explained via Measurement of Strong Coupling Parameters. *Biophys. J.* **2015**, *109* (11), 2317–2327. https://doi.org/10.1016/j.bpj.2015.10.031.

(38) Stottrup, B. L.; Veatch, S. L.; Keller, S. L. Nonequilibrium Behavior in Supported Lipid Membranes Containing Cholesterol. *Biophys. J.* **2004**, *86* (5), 2942–2950. https://doi.org/10.1016/S0006-3495(04)74345-3.

(39) Tero, R. Substrate Effects on the Formation Process, Structure and Physicochemical Properties of Supported Lipid Bilayers. *Materials* **2012**, *5* (12), 2658–2680. https://doi.org/10.3390/ma5122658.

(40) Karedla, N.; Schneider, F.; Enderlein, J.; Chen, T. Leaflet-Specific Structure and Dynamics of Solid and Polymer Supported Lipid Bilayers. *Angew. Chem. Int. Ed Engl.* **2025**, *64* (21), e202423784. https://doi.org/10.1002/anie.202423784.

(41) Tanaka, M.; Sackmann, E. Polymer-Supported Membranes as Models of the Cell Surface. *Nature* **2005**, *437* (7059), 656–663.

(42) Sun, Y.; Zang, X.; Sun, Y.; Wang, L.; Gao, Z. Lipid Membranes Supported by Planar Porous Substrates. *Chem. Phys. Lipids* **2020**, *228*, 104893. https://doi.org/10.1016/j.chemphyslip.2020.104893.

(43) Bera, M.; Kalyana Sundaram, R. V.; Coleman, J.; Chatterjee, A.; Thoduvayil, S.; Pincet, F.; Ramakrishnan, S. Highly Stable Planar Asymmetric Suspended Membranes for Investigating Protein Dynamics and Membrane Fusion. *Nat. Protoc.* **2025**, 1–25. https://doi.org/10.1038/s41596-025-01192-2.

(44) Teske, N.; Sibold, J.; Schumacher, J.; Teiwes, N. K.; Gleisner, M.; Mey, I.; Steinem, C. Continuous Pore-Spanning Lipid Bilayers on Silicon Oxide-Coated Porous Substrates. *Langmuir* **2017**, *33* (49), 14175–14183. https://doi.org/10.1021/acs.langmuir.7b02727.

(45) Mion, D.; Bunel, L.; Ramakrishnan, S.; Heo, P.; Pincet, F. Biomimetic Membrane in a Microfluidic Chip for the Electrical and Optical Monitoring of Biological Reactions. *Nat. Protoc.* **2025**, 1–32. https://doi.org/10.1038/s41596-025-01171-7.

(46) Visco, I.; Chiantia, S.; Schwille, P. Asymmetric Supported Lipid Bilayer Formation via Methyl-β-Cyclodextrin Mediated Lipid Exchange: Influence of Asymmetry on Lipid Dynamics and Phase Behavior. *Langmuir* **2014**, *30* (25), 7475–7484. https://doi.org/10.1021/la500468r.

(47) Vázquez, R. F.; Ovalle-García, E.; Antillón, A.; Ortega-Blake, I.; Bakás, L. S.; Muñoz-Garay, C.; Maté, S. M. Asymmetric Bilayers Mimicking Membrane *Rafts* Prepared by Lipid Exchange: Nanoscale Characterization Using AFM-Force Spectroscopy. *Biochim. Biophys. Acta BBA - Biomembr.* **2021**, *1863* (1), 183467. https://doi.org/10.1016/j.bbamem.2020.183467.

(48) Stanglmaier, S.; Hertrich, S.; Fritz, K.; Moulin, J.-F.; Haese-Seiller, M.; Rädler, J. O.; Nickel, B. *Asymmetric Distribution of Anionic Phospholipids in Supported Lipid Bilayers*. ACS Publications. https://doi.org/10.1021/la3019887.

(49) Enoki, T. A.; Feigenson, G. W. Asymmetric Bilayers by Hemifusion: Method and Leaflet Behaviors. *Biophys. J.* **2019**, *117* (6), 1037–1050. https://doi.org/10.1016/j.bpj.2019.07.054.

(50) Doktorova, M.; Heberle, F. A.; Eicher, B.; Standaert, R. F.; Katsaras, J.; London, E.; Pabst, G.; Marquardt, D. Preparation of Asymmetric Phospholipid Vesicles for Use as Cell Membrane Models. *Nat. Protoc.* **2018**, *13* (9), 2086–2101. https://doi.org/10.1038/s41596-018-0033-6.

(51) López-Montero, I.; Rodriguez, N.; Cribier, S.; Pohl, A.; Vélez, M.; Devaux, P. F. Rapid Transbilayer Movement of Ceramides in Phospholipid Vesicles and inHumanErythrocytes *. *J. Biol. Chem.* **2005**, *280* (27), 25811–25819. https://doi.org/10.1074/jbc.M412052200.

(52) Hua, L.; Kaiser, M.; Carabadjac, I.; Meister, A.; Hause, G.; Heerklotz, H. Vesicle Budding Caused by Lysolipid-Induced Asymmetry Stress. *Biophys. J.* **2023**, *122* (20), 4011–4022. https://doi.org/10.1016/j.bpj.2023.08.023.

(53) Pincet, F.; Adrien, V.; Yang, R.; Delacotte, J.; Rothman, J. E.; Urbach, W.; Tareste, D. FRAP to Characterize Molecular Diffusion and Interaction in Various Membrane Environments. *PLoS ONE* **2016**, *11* (7), 1–19. https://doi.org/10.1371/journal.pone.0158457.

(54) Yamada, A.; Mamane, A.; Lee-Tin-Wah, J.; Di Cicco, A.; Prévost, C.; Lévy, D.; Joanny, J. F.; Coudrier, E.; Bassereau, P. Catch-Bond Behaviour Facilitates Membrane Tubulation by Non-Processive Myosin 1b. *Nat. Commun.* **2014**, *5*, 3624. https://doi.org/10.1038/ncomms4624.

(55) Spudich, J. A.; Watt, S. The Regulation of Rabbit Skeletal Muscle Contraction. *J. Biol. Chem.* **1971**, *246* (15), 4866–4871. https://doi.org/10.1016/s0021-9258(18)62016-2.

(56) Tsai, F.-C.; Bertin, A.; Bousquet, H.; Manzi, J.; Senju, Y.; Tsai, M.-C.; Picas, L.; Miserey-Lenkei, S.; Lappalainen, P.; Lemichez, E.; Coudrier, E.; Bassereau, P. Ezrin Enrichment on Curved Membranes Requires a Specific Conformation or Interaction with a Curvature-Sensitive Partner. *eLife* **2018**, *7*, e37262.

(57) Pernier, J.; Kusters, R.; Bousquet, H.; Lagny, T.; Morchain, A.; Joanny, J. F.; Bassereau, P.; Coudrier, E. Myosin 1b Is an Actin Depolymerase. *Nat. Commun.* **2019**, *10* (1), 1–7. https://doi.org/10.1038/s41467-019-13160-y.

(58) Moens, P. D. J.; Bagatolli, L. A. Profilin Binding to Sub-Micellar Concentrations of Phosphatidylinositol (4,5) Bisphosphate and Phosphatidylinositol (3,4,5) Trisphosphate. *Biochim. Biophys. Acta - Biomembr.* **2007**, *1768* (3), 439–449. https://doi.org/10.1016/j.bbamem.2006.12.012.

(59) Walsh, J. P.; Suen, R.; Glomset, J. A. Arachidonoyl-Diacylglycerol Kinase SPECIFIC IN VITRO INHIBITION BY POLYPHOSPHOINOSITIDES SUGGESTS A MECHANISM FOR REGULATION OF PHOSPHATIDYLINOSITOL BIOSYNTHESIS (*). *J. Biol. Chem.* **1995**, *270* (48), 28647–28653. https://doi.org/10.1074/jbc.270.48.28647.

(60) Palmer, F. B. The Phosphatidyl-Myo-Inositol-4,5-Biphosphate Phosphatase from Crithidia Fasciculata. *Can. J. Biochem.* **1981**, *59* (7), 469–476. https://doi.org/10.1139/o81-065.

(61) Richter, R. P.; Bérat, R.; Brisson, A. R. Formation of Solid-Supported Lipid Bilayers: An Integrated View. *Langmuir* **2006**, *22* (8), 3497–3505. https://doi.org/10.1021/la052687c.

(62) Beber, A.; Taveneau, C.; Nania, M.; Tsai, F.-C.; Di Cicco, A.; Bassereau, P.; Lévy, D.; Cabral, J. T.; Isambert, H.; Mangenot, S.; Aurélie, B. Membrane Reshaping by Micrometric Curvature Sensitive Septin Filaments. *Nat. Commun.* **2019**, *10* (1), 420.

(63) Schäfer, J.; Nehls, J.; Schön, M.; Mey, I.; Steinem, C. Leaflet-Dependent Distribution of PtdIns [4, 5] P2 in Supported Model Membranes. *Langmuir ACS J. Surf. Colloids* **2020**, *36* (5), 1320–1328.

(64) Wen, Y.; Vogt, V. M.; Feigenson, G. W. PI(4,5)P2 Clustering and Its Impact on Biological Functions. *Annu. Rev. Biochem.* **2021**.

(65) Ellenbroek, W. G.; Wang, Y. H.; Christian, D. A.; Discher, D. E.; Janmey, P. A.; Liu, A. J. Divalent Cation-Dependent Formation of Electrostatic PIP2 Clusters in Lipid Monolayers. *Biophys. J.* **2011**, *101* (9), 2178–2184. https://doi.org/10.1016/j.bpj.2011.09.039.

(66) Meredith, S. A.; Kusunoki, Y.; Connell, S. D.; Morigaki, K.; Evans, S. D.; Adams, P. G. Self-Quenching Behavior of a Fluorescent Probe Incorporated within Lipid Membranes Explored Using Electrophoresis and Fluorescence Lifetime Imaging Microscopy. *J. Phys. Chem. B* **2023**, *127* (8), 1715–1727. https://doi.org/10.1021/acs.jpcb.2c07652.

(67) McIntosh, B. B.; Ostap, E. M. Myosin-I Molecular Motors at a Glance. *J. Cell Sci.* **2016**, *129* (14), 2689–2695.

(68) Pyrpassopoulos, S.; Feeser, E. A.; Mazerik, J. N.; Tyska, M. J.; Ostap, E. M. Membrane-Bound Myo1c Powers Asymmetric Motility of Actin Filaments. *Curr. Biol.* **2012**, *22* (18), 1688–1692.

(69) Malinin, V. S.; Haque, Md. E.; Lentz, B. R. The Rate of Lipid Transfer during Fusion Depends on the Structure of Fluorescent Lipid Probes: A New Chain-Labeled Lipid Transfer Probe Pair. *Biochemistry* **2001**, *40* (28), 8292–8299. https://doi.org/10.1021/bi010570r.

(70) Collins, M. D.; Gordon, S. E. Short-Chain Phosphoinositide Partitioning into Plasma Membrane Models. *Biophys. J.* **2013**, *105* (11), 2485–2494. https://doi.org/10.1016/j.bpj.2013.09.035.