1 # Global Metagenomics Reveals Hidden Protist Diversity
2

3 Miguel F. Romero[1]*, Arianna I. Krinos[1,2], Xyrus Maurer-Alcala[3], John A. Burns[4], Ramunas
4 Stepanauskas[4], Tanja Woyke[1,5], Frederik Schulz[1]*
5
[1]DOE Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, California,
USA
[2]Department of Earth, Environmental, and Planetary Sciences, Brown University, Providence,
Rhode Island, USA
[3]American Museum of Natural History, New York, New York, USA
[4]Bigelow Laboratory for Ocean Sciences, East Boothbay, Maine, USA
[5]University of California Merced, Life and Environmental Sciences, Merced, CA 95343, USA
13
*Correspondence:
mfromerogutierrez@lbl.gov
fschulz@lbl.gov

Abstract

Protists, defined as eukaryotes distinct from animals, plants, and fungi, are a polyphyletic
group that dominates the eukaryotic tree of life, exhibiting significant phylogenetic
diversity and fulfilling critical ecological roles. Historically, research has prioritized protists
associated with animals and plants, particularly those of medical significance, thereby
overlooking the majority of protist diversity. Conventional molecular methods, such as 18S
rRNA gene amplicon sequencing, frequently encounter limitations, including primer
binding bias and PCR bias caused by gene length variations, resulting in a biased
representation of protistan diversity. Further, most protist lineages are notoriously difficult
to cultivate. Here, we analyzed over 30,000 metagenome assemblies and protist single-cell
genomes, together with 21 long-amplicon 18S datasets, spanning marine, freshwater, and
soil ecosystems. We recovered 157,956 18S rRNA gene sequences (≥800 bp), which
clustered into 103,338 operational taxonomic units (OTUs) at 97% sequence identity and
24,438 OTUs at 85% identity. Notably, 69% of 13,238 non-singleton clusters at 85% identity
consisted exclusively of environmental sequences, uncovering a wealth of novel,
uncultivated, and unclassified protist diversity. A comprehensive taxonomic framework of
eukaryotes based on concatenated 18S and 28S rRNA genes that incorporated most novel
lineages revealed substantial underrepresentation of Amoebozoa, Discoba, and Rhizaria in
reference databases, with many lacking isolate or genome sequence representation.
Further, we identified 13 lineages that likely represent deeply branching diversity,
including candidates at approximately class- to phylum-level depth, that lack
representation in public databases. The corresponding 85% OTUs were primarily affiliated

with Excavata, with some branching deeply in the eukaryotic tree. Comprehensive analysis
of the global distribution of eukaryotes revealed uneven microbial eukaryotic diversity
across supergroups and ecosystems, with notable novelty particularly in soil and marine
environments. We then examined co-occurrence between protists and prokaryotes,
predicting putative symbiotic or predator-prey relationships, particularly among
understudied protist groups with bacteria such as Verrucomicrobia and Rickettsiales. Our
results extend current reference coverage and provide a global, contig-based framework
for protistan diversity and distribution, highlighting major gaps in curated databases and
metabarcoding coverage and guiding targeted studies of these organisms' ecological
roles.

Introduction

The vast majority of eukaryotic diversity lies outside the well-established kingdoms of Animalia,
Fungi, and Plantae. These organisms have traditionally been referred to collectively as 'protists',
a term that encompasses a highly diverse and non-monophyletic assemblage of lineages. In
addition to their remarkable diversity, protists exhibit a wide array of cellular structures and
lifestyles (Foissner 2006; Adl et al. 2007). Previous research has largely focused on protists
associated with animals and plants, particularly those of medical relevance (Collins and Jeffery
2007; Horn 2022; Einarsson et al. 2016). Well-characterized lineages include the TSAR group,
which comprises Stramenopiles (e.g., diatoms and oomycetes), Alveolata (e.g., ciliates,
dinoflagellates, and apicomplexan parasites), Rhizaria (e.g., foraminifera and radiolarians), as
well as the Amoebozoa supergroup (sensu stricto amoebae) and the Discoba division (which
includes heteroloboseans and euglenoids, including kinetoplastids). Even less well studied
eukaryotic lineages include Metamonada, which encompass instances of symbiotic or parasitic
groups like diplomonads and parabasalids (Céza et al. 2022). Recently, advances in molecular
methods, such as cultivation-independent single-cell sequencing (Woyke et al. 2017), have
facilitated the exploration of phylogenetic associations among selected uncultivated groups,
resulting in the identification of new eukaryotic supergroups, such as Hemimastigophora
(Foissner et al. 1988; Lax et al. 2018) and Provora (Tikhonenkov et al. 2022), with additional
members continually being added as interest in these novel lineages expands (Eglit et al. 2024).
68
Molecular methods such as 18S rRNA gene amplicon sequencing have proven invaluable for
assessing eukaryotic diversity in various environmental contexts. In oceanic plankton
communities, studies revealed that heterotrophic protists represent the most diverse group,
despite metazoans dominating in terms of abundance (de Vargas et al. 2015). A protist-centric
survey in select soil environments indicated that Cercozoa are the most prevalent group,
revealing correlations between protistan and bacterial taxa, including evidence of potential
predator-prey interactions (Oliverio et al. 2020). At a global level, soil eukaryotic diversity has
primarily been evaluated with respect to specific groups such as Fungi and testate amoebae
(Fernández et al. 2016; Bahram et al. 2018). Diversity of eukaryotes in soil typically exhibits
higher values in the Simpson diversity index in equatorial regions and intermediate latitudes,

indicating that diversity peaks in temperate climates, likely linked to water-energy balance and
present similar patterns such as plants and animals (Fernández et al. 2016; Bahram et al. 2018).
A comprehensive meta-analysis that incorporated data from marine, freshwater, and topsoil
ecosystems demonstrated that topsoil harbors greater protistan diversity compared to aquatic
environments (Singer et al. 2021). Despite these extensive studies on eukaryotic diversity and
community structure, recent findings indicate that an overwhelming majority of protistan taxa
remain undocumented. For example, an analysis suggested that one-third of operational
taxonomic groups (OTUs) could not be assigned to known eukaryotic groups (de Vargas et al.
2015). Similarly, new lineages of protists, at both the phylum and supergroup levels, including
Hemimastigophora and Provora, have only recently been described (Foissner et al. 1988; Lax et
al. 2018; Tikhonenkov et al. 2022).
90
The 18S rRNA gene (also referred to as 18S rDNA) has emerged as the most frequently used
marker gene for identifying eukaryotic organisms in environmental studies. This gene encodes
the small subunit of the ribosomal RNA and is universally present in all nuclear genomes,
comprising regions of both high conservation and high variability (Wuyts et al. 2001). Public
databases, such as Silva (Quast et al. 2013) and the Protist Ribosomal Reference (PR2; Guillou
et al. 2013), have served as repositories for these sequences for decades. Polymerase chain
reaction (PCR)-based approaches (metabarcoding) are typically used to recover this marker
gene with high sensitivity in environmental samples. However, due to the constraints of short-
read DNA sequencing technologies, only a limited portion of the 18S rDNA is typically screened,
predominantly the V4 to V6 and V9 regions (Hugerth et al. 2014; de Vargas et al. 2015; Oliverio et
al. 2020). Unlike its prokaryotic counterpart (the 16S rRNA gene), the 18S rRNA gene exhibits
pronounced length variation, ranging from a minimum of 1.5 kb (in Diplomonadida) to a maximum
of 6.4 kb (in Amoebozoa), with the mature 18S rRNA molecule potentially reaching lengths of up
to 2.3 kb (Xie et al. 2011). This variability can be attributed to length-variable regions that
frequently contain introns, some exceeding 1 kb, especially in regions commonly amplified (Xie et
al. 2011). Such disparities in length, coupled with the rapid evolution of the 18S rRNA gene in
certain lineages (Meyer et al. 2010), may bias PCR-based metabarcoding studies, as only a
fraction of the eukaryotic diversity in the environment can be captured. Recently, long amplicon
sequencing has been deployed to recover multiple markers from the rRNA gene locus within
eukaryotic genomes (Jamy et al. 2022). Although this technique solves some of the challenges
associated with conventional methods (Overgaard et al. 2024), primer binding bias remains a
concern due to sequence divergence.
113
An alternative approach that is able to further mitigate limitations of amplicon-based methods is
shotgun metagenomics as it does not employ specific primers for amplifying target sequences.
Instead, it involves screening assembled sequences from the genomes of all taxa present in the
sample for rRNA genes. Depending on sequencing depth, genomic information from rarer taxa
can also be retrieved. Targeted marker gene recovery from metagenomic assemblies is more
sensitive than genome-resolved metagenomics, which relies on successful binning. This gene-
centric approach provides higher sensitivity by enabling the analysis of contigs from low-
abundance organisms or fragmented genomes that fall below the thresholds required for
accurate binning (Nayfach et al. 2021). However, this method remains susceptible to artifacts

inherent to the assembly process itself, including sequence fragmentation, chimerism, and the
collapse of strain-level diversity into consensus sequences (Sczyrba et al. 2017). In our study, we
employed a shotgun metagenomics approach combined with marker gene extraction from
assembled contigs. We leverage over 30,000 assembled metagenomes to conduct a global
census of eukaryotic diversity. Given the potential confounders introduced by metagenomic
sequence assembly, which can obscure fine-grained phylogenetic signals, we focused our
analysis primarily on higher taxonomic ranks. Our primary objective was to elucidate the
phylogenetic and ecosystem diversity of microbial eukaryotes on a global scale, thereby
advancing the understanding of protistan diversity and distribution, including the characterization
of previously unidentified lineages.

Results and discussion

Expansion of Known Eukaryotic Diversity

The screening of 32,520 assembled metagenomes yielded 157,956 18S rRNA genes from
18,444 sequencing datasets, all with a minimum length of 800 bp (a summary of the methods can
be found in Supplementary Figure 1). We clustered the recovered 18S sequences together with
curated reference sequences (SILVA, PR2, and recent long-amplicon datasets; Quast et al.
2013; Guillou et al. 2013; Jamy et al. 2022; Tikhonenkov et al. 2022). We used a two-step identity
threshold (97% and 85%) because 18S identity does not map uniformly to taxonomic rank across
eukaryotes. This produced 103,338 OTUs at 97% identity and 24,438 OTUs at 85% identity. We
discarded all 11,200 85% identity clusters containing only a single sequence, retaining 13,238
clusters with two or more sequences for further analysis. As reference databases such as Silva
and PR2 include sequences originating from both cultured isolates as well as environmental
sources, we were able to estimate the proportion of sequences corresponding to uncultivated
microbes within our dataset.

Our analysis revealed that 69% of the 85% identity clusters (n=9,146) were exclusively
comprised of environmental sequences, while 30% (n=4,006) included sequences from isolates
and less than 1% (n=86) contained sequences derived from genomes. Clusters that contained
sequences from isolates (18S rRNA genes) were generally larger, whereas the metagenomic-
only clusters tended to be smaller (Figure 1A). Clusters containing genome sequences had an
average of 1,334 members, greatly exceeding the average of 49 members for clusters containing
sequences from isolates. Notably, the mean size of clusters containing solely environmental
sequences was calculated at 10. Among the largest clusters containing only sequences from
reference databases were sequences from Aconoidasida (Apicomplexa, size=1,220) and
Conoidasida (Apicomplexa, size=910). Conversely, the largest clusters including metagenomic
sequences comprised sequences from Malasseziales (Fungi, size=18,608) and Colpodellida
(Chrompodellids, size=10,291). The largest cluster incorporating genome sequences was also
affiliated with Malasseziales, while the largest cluster containing isolate-derived sequences
without genome information was ascribed to Hexapoda (Metazoa, size=10,510). Notably, the
largest cluster exclusively composed of environmental sequences was designated to Syndiniales

Group II (Syndiniales, size=1,028). To further investigate the clusters with only reference
sequences, we estimated the number of distinct taxa found per cluster.
165
Our data-driven approach generated clusters containing, on average, 2.4 genera, 1.7 families,
and 1.32 orders (Supplementary Figure 2). Similarly, when considering only well-resolved orders
(excluding unclassified .U.order and unresolved _X lineages), each order encompassed an
average of 1.59 families and 2.4 genera. These data suggest that the 85% sequence identity
threshold used for clustering corresponds approximately to the taxonomic level of order or family.
Although metagenomic assembly is susceptible to inherent artifacts including sequence
fragmentation, chimerism, the collapse of strain-level diversity into consensus sequences
(Sczyrba et al. 2017), and exhibits low sensitivity for low-abundance members, we mitigated
these issues by focusing our analyses on higher taxonomic ranks.
175
Our results indicate that protists recognized as parasites of animals and humans, such as
Piroplasmorida (Apicomplexa, 1,220 sequences) and Cryptosporida (Apicomplexa, 910
sequences), are the predominant representatives within reference databases, revealing a
pronounced sampling bias in the representation of microbial eukaryotes (del Campo et al. 2014).
Some parasitic protists, such as apicomplexans, were also found to be dominant in soil
microbiomes (Mahé et al. 2017). Consequently, it is unsurprising to encounter sizable clusters of
sequences derived from parasitic protists, whether from isolates or environmental samples, with
the latter exhibiting a more balanced representation of the diversity of these organisms.
184
To enhance the taxonomic annotation of sequences and to better reflect the phylogenetic
relationships among eukaryotes, we constructed a eukaryotic reference tree utilizing 1,438 85%
OTU cluster representatives that also presented matching 28S rRNA genes. This subset of our
dataset excluded most of the highly populated clusters (Figure 1B). The resulting tree broadly
aligned with the established phylogeny of eukaryotes, as it effectively grouped the Amorphea
together, distinctly separate from the Diaphoretickes, in agreement with observations from both
rRNA and protein-based alignments (Jamy et al. 2022; Torruella et al. 2024). Further,
Metamonads were positioned near the root, consistent with phylogenomic analysis of the root of
the eukaryotic tree (Al Jewari and Baldauf 2023; Williamson et al. 2025) . Due to high mutation
rates and pronounced length variability, certain groups such as Microsporidia (Vossbrinck et al.
1987) were excluded from the reference tree, leaving their representatives unclassified within our
phylogenetic taxonomic annotation. Numerous clades grouped within Metamonada,
Stramenopiles, Alveolata, Amoebozoa, Metazoa, and Fungi divisions entirely lacked sequences
derived from isolates or genomic sources (Figure 1C). Clusters associated with these clades also
tended to exhibit representative sequences with low blast alignment identity to reference
sequences and increased average predicted primer binding bias (Figure 1C) compared to
sequences from other clusters, suggesting these represent taxonomic blindspots overlooked in
prior PCR amplification-based studies (Morey et al. 2024).
203
Our census of eukaryotic diversity revealed an uneven taxon richness across various
supergroups, with Obazoa (n=3,090), Amoebozoa (n=2,998), and TSAR (n=2,081) exhibiting the
highest number of OTUs. At the division level, Opisthokonta harbored the largest diversity,

predominantly characterized by sequences derived from isolates. A total of 36 clusters
incorporated sequences from genomes, while 2,039 sequences were drawn from isolates, and
1,015 clusters were comprised solely of environmental sequences (Figure 1D). The divisions
classified under Amoebozoa exhibited a high proportion of clusters exclusive to environmental
samples (95%), most of which remained unclassified at the division level (n=1,400). Similarly,
Discoba, the second most diverse division, demonstrated novelty in sequences presented as a
significantly larger proportion of clusters derived exclusively from metagenomic sequences
(n=1,312), with only 202 clusters containing sequences from isolates and a mere 6 clusters
encompassing genome-derived sequences. In the context of the TSAR supergroup, Rhizaria
emerged as the most diverse division, with 73.5% of the 937 clusters comprising entirely
environmental sequences. The predominance of Obazoa, which includes metazoans, may
reflect its evolutionary success due to complex multicellularity and diverse nutritional strategies
(Torruella et al. 2015), unlike other groups of multicellular eukaryotes such as brown algae.
Conversely, Amoebozoa's high diversity demonstrates its adaptive radiation across soil and
aquatic habitats through varied locomotion mechanisms and predatory strategies (Tekle et al.
2022), which could be true also for some groups of discobans. Within the TSAR supergroup,
Rhizaria's high OTU richness is consistent with rapid SSU divergence (Burki and Pawlowski
2006). The resulting diversification, which is particularly well-documented in marine systems, has
resulted in highly divergent SSU rRNA gene sequences even among closely related taxa
(Schuler et al. 2018). The 85% sequence identity threshold employed for our clustering was
specifically chosen to accommodate this substantial sequence divergence. This approach
successfully resolved numerous diverse rhizarian lineages, underscoring the degree to which
current cultivation-based methods fail to capture the true phylogenetic breadth of these major
eukaryotic groups.
231
Representative sequences that could not be assigned to established eukaryotic supergroups
were designated as 'Eukaryota.U'. Consequently, 'Eukaryota.U' does not represent a single
monophyletic group; rather, it is a polyphyletic assemblage of multiple, taxonomically distinct,
and unclassified eukaryotic lineages. Among these, 2,274 (78%) were exclusively derived from
metagenomic sequences, while 621 clusters contained sequences from isolates and 13 clusters
harbored sequences from genomes. Among the reference-only clusters that were unassigned to
any supergroup using the tree-placement method, most were represented in the reference tree
according to sequence annotation, such as Cnidaria (129 clusters), or had a sister group present
within the same supergroup or division, exemplified by Microsporidia (36 clusters). This finding is
somewhat expected for lineages such as Microsporidia, which are characterized by short yet
highly divergent 18S rRNA gene sequences (Vossbrinck et al. 1987). Challenges in supergroup
assignment using tree-placement methods often stem from incomplete reference databases or
rapid evolutionary rates. Despite sequence annotation indicating taxonomic placement, many
organisms remain unclassified at higher taxonomic levels while having clear affiliations at lower
ranks. This phenomenon reflects the evolutionary complexity of certain lineages that escape
traditional classification approaches, as has been previously observed (Berger et al. 2011).
248
We found differences in insertion number and length among eukaryotic supergroups, with
Centroplasthelida with the highest mean number of insertions (6.54), and Chlorophyta with the

longest mean insertion size (665.34; Figure 1E). Even with variation within lineages of eukaryotic
supergroups (Xie et al. 2011), we found differences between groups. Predicted binding bias
scores also varied among groups, with those with the highest proportion of phylogenetic novelty
(Figure 1D) showing also a high mean binding bias score, such as Metamonada (mean
score=3.83), Evosea (3.58), Discoba (3.02), and Eukaryota.U.division (unclassified at the
supergroup level, mean score=2.93). This observation could explain why some lineages within
these groups might have remained as taxonomic blindspots in metabarcoding studies.

258
259
Figure 1. Phylogenetic Diversity of Eukaryotes. (A) Histogram illustrating the distribution of cluster
sizes, color-coded based on the origin of the included sequences. The inset represents the number of
clustered sequences from each source. Data corresponds to 85% identity clusters after discarding
singletons. (B) Cluster size distribution of the OTUs that contained both 18S and 28S rRNA genes, which
were employed to construct the eukaryotic reference tree. (C) Reference tree inferred from a concatenated
alignment of the 18S and 28S rRNA genes derived from OTUs that had representative members with both
genes present. The heat map on the innermost track indicates the cluster-wide average of the accumulated
primer binding bias score of the corresponding 18S rRNA gene sequences utilizing a set of four commonly
used general eukaryotic primer pairs. A score of 4 (dark purple) suggests that the sequence would not bind
to any of the primer pairs, while a score of 0 (white) implies it would bind to all of them. The adjacent
heatmap delineates sequence identity compared to the best hit within our combined 18S rRNA gene
sequence database. The outermost track denotes the cluster composition corresponding to the legend
illustrated in panel A. Black circles at the tip of the leaves indicate that the respective clusters are
composed exclusively by environmental sequences extracted in this study. (D) Abundance of clusters for

each eukaryotic division within the analyzed environmental samples. (E) Distribution of insertion count and
cumulative insertion length among eukaryotic divisions. Each point shows the corresponding data for each
OTU representative sequence. Insertions are defined as sequences present in each 18S rRNA gene
sequence not represented in the reference covariance model. Data of insertions with length >= 10 bp is
shown. Box plots depict the interquartile range (IQR, 25th to 75th percentiles), with horizontal lines
representing median values. Whiskers extend to the most extreme data points within 1.5 × IQR from the
box boundaries, and data points beyond this range are displayed as individual circles, representing
statistical outliers. White circles indicate mean values. (F) Distribution of predicted primer binding bias
score for each OTU representative sequence ordered by its taxonomic classification at the division level.
Box plots show the interquartile range with median values (horizontal lines). Whiskers extend to data points
within 1.5 × IQR, with outliers represented as individual circles. Mean values are shown as white circles.

The Global Distribution Of Microbial Eukaryotes

The majority of 18S rRNA gene sequences recovered (present in non-singleton clusters) were
from the Integrated Microbial Genome/Metagenome (IMG/M) database (Chen et al. 2023),
representing samples from all continents, encompassing longitudes and latitudes from pole to
pole (Figure 2A). The main ecosystem type sampled was soil (n=5,487), followed by marine
(n=4,390) and freshwater (n=3,235). Among the frequently observed subtypes, lake samples
(n=1,588), peat (n=985), and oceanic samples (n=974) were predominant (Supplementary Table
1). In contrast, datasets without any identified 18S rRNA gene sequences were most frequently
associated with subtypes such as phylloplane/leaf (n=408), coalbed methane well (n=319), and
insect nests (n=248) (Supplementary Table 2), the presence of 16S rRNA genes in the same
samples indicates that these may represent a sample preparation bias or a harsher environment
for growth of microbial eukaryotes. Despite the broad spectrum from which these samples were
gathered, the majority originated from latitudes ranging from 60° to 30° North, particularly within
North America and Central Europe (Figure 2B). Key factors influencing protist diversity include
water availability, pH, and nutrient availability (Zhao et al. 2023). The captured protistan diversity
in freshwater and peat bogs is in agreement with what has been shown in previous studies (Lara
et al. 2011; Krinos et al. 2024; Oliverio et al. 2020). In contrast, the absence of 18S rRNA gene
sequences in habitats like coalbeds, that are characterized by low water availability and low
nutrient concentrations, reflect challenging growth conditions for eukaryotes.
304
Next, we investigated the overall distribution of predominant and ubiquitous microbial eukaryotic
orders and families. Relative abundance was defined as the average of the normalized mean
sequencing depth across all contigs pertaining to each taxonomic group, while ubiquity was
defined as the proportion of samples from comparable environments and latitudes in which each
taxonomic group was identified. Within soil ecosystems, well-studied groups such as
Exobasidiomycetes (Fungi; relative abundance = 2.73) and Insecta (Metazoa; relative
abundance = 5.78) were notably abundant, whereas the crustaceans Maxillopoda (Metazoa)
constituted the most abundant group within marine environments (relative abundance = 15.62).
Excluding Metazoa, Fungi, and Streptophyta (plants), the most abundant group detected in
marine ecosystems consisted of green algae from the family Bathycoccaceae (relative
abundance = 8.65; Figure 2C), followed by diatoms (family Thalassiosiraceae; relative
abundance = 8.06) and brown algae (family Phaeophyceae_XX; relative abundance = 6.37).

Additionally, parasitic dinoflagellates from Ichthyodinida Clade-1 and Dino-Group-II-Clade-7
were frequently observed with relatively high abundances (relative abundance = 3.55 and 2.76,
respectively). In freshwater systems, the most abundant groups included Thalassiosiraceae
(Bacillariophyta; relative abundance = 11.81) and green algae (Chlamydomonadales_X; relative
abundance = 5.61), with ciliates from the Oxytrichidae family being present across the largest
number of freshwater samples (ubiquity = 0.22). Bathycoccaceae (Chlorophyta) emerged as the
most abundant group within soil ecosystems (relative abundance = 2.85), closely followed by
flagellated bacteriotrophic protists from the family Sandonidae (Cercozoa; relative abundance =
1.11). Historically, diatoms and metazoans have been classified as dominant groups within
marine environments, alongside chlorophytes and parasitic dinoflagellates (de Vargas et al.
2015). Similarly, prior studies have indicated that lineages of Cercozoa hold dominant positions
within soil ecosystems, along with other groups of chlorophytes (Oliverio et al. 2020). Our
findings revealed Bathycoccaceae as highly abundant protists in soil, likely due to the inclusion of
diverse soil ecosystem subtypes, such as peatlands, glaciers, and wetlands, where
Bathycoccaceae were most frequently observed (Supplementary Table 3). This contrasts to
topsoils, which contain algae from various groups, such as Chlorophyta and Bacillariophyceae
(Zancan et al. 2006). The observed differences in abundance between our findings and previous
amplicon-based studies (Mahé et al. 2017; Oliverio et al. 2020) may stem from the heterogeneity
of samples, enhanced gene recovery of high-abundance eukaryotes in shotgun metagenomes,
and the absence of amplification and primer-binding bias. Taken together, our results provide a
more unbiased perspective of abundances of microeukaryotes in global ecosystems.
338
Our eukaryotic census also provided a more detailed perspective on the distribution of eukaryotic
groups across different latitudes (Figure 2D). Even at elevated taxonomic ranks, distinct patterns
emerged across latitudes and ecosystem types. The marine environment exhibited both high
ubiquity and abundance of the Stramenopiles, Alveolata, and Rhizaria (SAR) group. Notably, at
the arctic tundra around 90°, representatives from all three divisions were consistently present
(ubiquity = 1) and exhibited relative abundances of 5.8, 4.7, and 2 for Alveolata, Stramenopiles,
and Rhizaria, respectively. Opisthokonts were also found to be ubiquitous (ubiquity = 1) and
abundant (relative abundance = 7.6) in close proximity to the North Pole. Prior studies using
documentary sources, such as cell counts, showed the presence of alveolates, stramenopiles,
and unicellular opisthokonts within the pan-Arctic region; however, rhizarians were not observed
in this context (Poulin et al. 2011). In freshwater ecosystems, Opisthokonts, particularly Metazoa,
displayed the highest mean ubiquity (0.3) alongside high relative abundance at -10° and 40°
(relative abundance = 17.2 and 33.2, respectively), while groups such as Discoba were also
represented with mean ubiquity = 0.22 and high abundance at latitudes of 30° and 50° (relative
abundance = 60.2 and 16.7, respectively), with Discoba_X (primarily composed of
heteroloboseans) exhibiting a mean abundance of 5.5. Sequences lacking supergroup-level
classification were most ubiquitous in freshwater (mean ubiquity = 0.3), though their abundance
exhibited variability based on latitude, peaking at 64.6 at 50°. Within soil ecosystems,
opisthokonts and streptophytes were ubiquitous across all latitudes (mean ubiquity = 0.5 and 0.3,
respectively), whereas other groups were less frequently observed. Nevertheless, Cercozoa
persisted as the most predominant eukaryotic groups in soil, with classes such as Cercozoa.U.
leading with a mean relative abundance of 21.38, while Glissomonadida and

Thaumatomonadida displayed lower mean abundances of 1.1 and 0.2, respectively. This is in line with previous reports that showed that Cercozoans comprise a substantial proportion of microbial biomass in soil when evaluated using metatranscriptome sequencing data (Urich et al. 2008), often represented by highly divergent 18S rRNA gene sequences even among closely related lineages, as observed from isolates (Howe et al. 2011). Our results confirm that cercozoans represent a dominant eukaryotic group in soil environments with variable abundance as a function of latitude even within soil environments, and further expand the prevalence of this group through many newly identified lineages.

Our dataset of 18S rRNA gene sequences from thousands of samples presents a unique opportunity to estimate the taxonomic diversity of microbial eukaryotes on a global scale. Rarefaction curves of the number of observed OTUs as a function of number of sequencing datasets showed that microbial eukaryote OTU richness did not reach a plateau for any of the ecosystem types, indicating that more extensive sampling efforts could yield a substantial number of novel OTUs (Figure 2E). Partitioning the samples among ecosystem types according to their GOLD ecosystem subtype showed that the highest eukaryotic diversity within marine systems was observed in epipelagic, oceanic, and coastal subtypes (Figure 2F). The highest diversity in freshwater was found in river and peatland subtypes within soil. This contrasts with a prior amplicon sequencing-based study which reported soil as the most diverse ecosystem for eukaryotic taxa (Singer et al. 2021). However, this previous research focused exclusively on topsoil and planktonic samples, while our study highlights that sediment samples represent important contributors to eukaryotic diversity in aquatic environments (Supplementary Figure 3). Freshwater ecosystems might provide nutrients for trophic groups to grow and in turn allow growth of heterotrophic organisms. These ecosystem dynamics align with the crucial role of nutrient cycling in supporting diverse trophic interactions in freshwater systems (Cotner and Biddanda 2002). These results indicate that nutrient availability and a stable ecosystem can in turn produce microbial communities as diverse as others in complex ecosystems such as the soil as freshwater bacterial communities often rival soil communities in phylogenetic diversity (Newton et al. 2011). Further, environmental stability and resource availability are key drivers of microbial community complexity across ecosystem types, including both aquatic and terrestrial environments (Delgado-Baquerizo et al. 2018).

Further, we were interested in patterns of the latitudinal gradient of eukaryotic diversity, as these patterns have been studied for other microbial groups such as marine bacteria (Fuhrman et al. 2008) and is a recurrent question in ecology (Mittelbach et al. 2007). We assessed the correlation between absolute latitude and OTU diversity within each eukaryotic subdivision across the three most prevalent ecosystem types in our samples. We were able to identify distinct patterns in the correlation of the 18S rRNA diversity with absolute latitude. Most taxonomic groups displayed positive coefficients in the quadratic term, indicating that diversity peaks at the Equator and the poles (Figure 2G). This finding contrasts with previous studies, done on observation data based on richness data of observations of benthic diatoms, suggesting that protist diversity shows weak or no correlation with latitudinal gradients, in contrast to plants and animals, which exhibit pronounced responses influenced by body size (Hillebrand and Azovsky 2001). Other investigations focused on specific taxa, such as Fungi, which exhibited a higher diversity at

intermediate absolute latitudes (Bahram et al. 2018), in contrast to their metazoan counterparts.
This was also evident in our analysis. Further, fungal lineages have been shown to have an
increase in diversity near the equator, with tendencies towards reduced diversity in polar regions
(Azovsky and Mazei 2013; Lara et al. 2016). While this trend was also visible in our data, we
found that the Ciliophora subdivision displayed an even higher diversity in freshwater near the
equator, as did the Dinoflagellata subdivision in marine systems.
411

412
Figure 2. Global distribution of eukaryotes. A) World map illustrating the origin of the 17,371 samples
from which 18S rRNA genes were recovered and geolocalization data is available. B) Total number of

samples within each latitude range, partitioned by ecosystem type. C) Mean values of ubiquity and
normalized abundance for eukaryotic families that exhibited abundance and ubiquity above the 80%
quantile. Families belonging to Metazoa, Fungi, and Streptophyta have been excluded. D) Mean
abundance and ubiquity of each eukaryotic division across latitude ranges. Rows marked with cross marks
indicate no observed samples at that latitude and ecosystem type. Brown circles indicate normalized
abundance lower than 1. E) Rarefaction curves representing the mean number of observed OTUs
following 10,000 randomizations. Gray shades represent standard deviation values. F) Per-sample
diversity distribution for each ecosystem type with at least a 1% share of the datasets. Box plots display
medians (horizontal lines), interquartile ranges (boxes), and data distribution (whiskers), with outliers
shown as circles. G) Second-order polynomial fits for Simpson index values as a function of absolute
latitude across different ecosystem types; only significant regressions (adjusted p-value < 0.05) are
portrayed.

New Lineages Of Eukaryotes

Our comprehensive census of eukaryotic diversity has uncovered a substantial number of novel
eukaryotic groups (2,072 OTUs at 85% sequence similarity) that are solely represented by
metagenomic sequences, eluding classification within any known eukaryotic lineage. In order to
resolve these unclassified lineages, we expanded our robust species tree utilized for taxonomic
placement by incorporating these taxa.

The expanded tree preserved the majority of eukaryotic lineages as monophyletic, while also
accommodating several orphan sequences interspersed among them (black branches, n=1,670)
confirming them as polyphyletic (Figure 3A). The presence of long orphan branches grouping
within known eukaryotic phyla is noteworthy. A plausible explanation for these long branches is
accelerated evolutionary rates, analogous to those documented in obligate intracellular bacteria
(Ferla et al. 2013) and in eukaryotic parasites like microsporidians, which are known to exhibit
highly divergent 18S rRNA gene sequences (Vossbrinck et al. 1987). While such branches can
result from evolutionary processes, including rapid diversification or divergent molecular
evolution, this pattern warrants further investigation using complementary approaches. For
example, only through single-cell transcriptome sequencing was it possible to identify a novel
alveolate intracellular parasite closely related to perkinsids (Cooney et al. 2024). Cultivation of
numerous eukaryotic lineages is notoriously challenging, leaving a considerable proportion of
eukaryotic diversity unexplored on both genomic and phenotypic levels. In this study, we did not
identify previously documented orphan OTUs, as our work mitigated primer binding bias. Indeed,
the bias score exhibited significantly higher values (adjusted p value = 0) in orphan sequences
compared to reference sequences, irrespective of their origins from isolates or environmental
sequencing datasets (Figure 3B).

Notably, the majority of environmental orphan sequences (n=1,670) branched deeply in the
eukaryotic tree or in proximity to discoban sequences bridging excavates and the eukaryotic root.
We identified 13 clades entirely composed of orphan OTU representatives, likely indicative of
major, undescribed new lineages (NLs, colored ranges). Given that 28S rRNA gene sequences
were also detected within some metagenomic contigs containing orphan 18S OTU
representatives, we examined whether the inclusion of additional phylogenetic information

through adding the 28S rRNA gene sequences would impact their classification and placement
within the reference tree. In our results the placement remained consistent for sequences in
NL11, NL9, NL41, and several orphans within Fungi and TSAR (filled red circles). New lineages 5
and 11 branched outside of all eukaryotic sequences from the reference tree, suggesting they
may represent early branching eukaryotes. In contrast, several of these new clades were
affiliated with Excavata, specifically the Discoba division, such as NL9. Various clades branched
as sister groups to sequences from the discoban phylum Heterolobosea: NL37, NL33, NL12,
NL28, NL39, NL30, NL36, and NL41. The most phylogenetically diverse group within these new
lineages, NL41, formed a substantial clade that may represent a new discoban lineage closely
related to Heterolobosea.
468
We further validated NL41 as a distinct lineage using an independent dataset of single-amplified
genomes, derived from cells isolated from anoxic marine sediments in Maine. Phylogenomic
analyses using a set of 317 conserved eukaryotic proteins indicated that the SAGs formed a
novel clade , sister to *Percolomonas cosmopolitus* (Figure 3D). Within the novel SAG clade, we
found three SAGs containing 18S rRNA gene sequences, all of which group within the NL41
clade in the 18S tree (Figure 3A,D). This phylogenetic placement, together with the observation
that the best BLAST hit from the nucleotide database corresponding to these sequences was the
18S rRNA sequence of *Creneis carolina* (mean identity %=78.3), implies that NL41 belongs to
Heterolobosea, potentially related to Tetramitia.
478
We traced the ecosystem types from which sequencing datasets belonging to the new lineages
were obtained and observed that marine ecosystems contributed the largest proportion, followed
by freshwater and soil (Figure 3C). Upon assessing the environmental sources of these datasets,
we found that wetlands from both marine and freshwater ecosystems were the most frequently
represented ecosystem subtypes (Supplementary Figure 4; Supplementary Table 5). Wetlands
harbor diverse microbial communities, including the newly proposed bacterial phylum
Candidatus Cosmopoliota (Zhang et al. 2023), which was identified through metagenome-
assembled genomes (MAGs) recovered from this ecosystem.
487
We used an example lineage for which there was limited taxonomic confidence to demonstrate
that more than just the 18S rRNA gene could be extracted for these novel lineages from the
metagenomes. We chose the novel lineage NL5 because of its location in the tree among
discoban clades and other groups composed of unclassified sequences and better classify it in
the eukaryotic tree using recovered coding genes. Thus we applied the tax-aliquots clustering
approach (described in the Methods) to identify functionally annotated genes that clustered with
the non-18S rRNA gene portions of contigs containing the 18S rRNA gene established through
our discovery clustering pipeline (Krinos et al. 2024). This approach clusters sequences based
on their *k*-mer frequencies to identify sequences from similar taxonomic lineages that cannot be
aligned to existing reference sequences. We retained clustered contigs associated with protein
family annotations (Pfam database) with bitscores exceeding 100, thus stringently excluding
clustered sequences lacking robust protein annotations. The presence of functional genes within
nucleotide sequences which clustered with the 18S rRNA gene-containing contigs provides
additional evidence that the 18S rRNA genes are valid. Among sequences exclusively clustering

with NL5, we identified 24 distinct Pfam annotations on clustered contigs using the hierarchical
clustering approach. These Pfam annotation-containing contigs encompassed multiple
membrane proteins (amino acid permease and ABC membrane transporters), a stress/heat
shock-related protein (AHSA1), alpha/beta hydrolases, glycosyl transferases, carbohydrate
phosphorylases, and sulfur transport proteins. We aligned these functionally-annotated proteins
with proteins from the nr database from NCBI (Pruitt et al. 2006) to identify functionally-annotated
proteins with similar taxonomic alignment position to NL5. Of 52 unique Pfams annotated across
68 distinct proteins identified in nucleotide-clustered contigs, 29 of 68 (43%) were taxonomically
annotated as eukaryotes and 11 (38%) of those eukaryotes had a maximum percentage identity
of less than 80% assigned to the best hit to the NR database. 26 protein sequences (38.2%) did
not have any BLAST hit to the NR database at all. This result aligns with the novel diversity
identified via the 18S rRNA gene sequences and provides putative functional proteins that
appear to have derived from the new lineages.

Previous amplicon-based studies of eukaryotic diversity in environmental samples have
indicated that many protist groups contain representatives that remain virtually unknown (de
Vargas et al. 2015). The classification of these new clades poses challenges, often limited to
specific clades (Reboul et al. 2019; Fermani et al. 2021), primarily due to the conventional short
lengths of amplicon libraries, which cover only selected regions of the 18S rRNA gene (Hugerth
et al. 2014). To overcome some of these limitations, we recovered 18S rRNA genes (>1,200 bp)
from assembled contigs and, if co-located, also the 28S rRNA genes of novel eukaryotic
lineages, improving phylogenetic resolution. Our findings imply the presence of at least 13 major
clades of eukaryotes lacking both genomic representation and closely related sequences in
reference databases, representing taxonomic blind spots. For NL41, we provide independent
support from single-cell genomic data that corroborates the phylogenetic placement inferred from
our metagenome-derived rRNA sequences. The detailed information on sample origins provided
in this study, predominantly wetlands and sediment systems, will facilitate future targeted
sampling efforts aimed at further exploring this unexpected novelty within the eukaryotic tree of
life.

We mapped approximately 71,000 short-read V4 18S rRNA gene amplicon sequences against
our database of ~13,000 metagenomic OTU representative sequences. This comparison
revealed that the eukaryotic diversity captured by metabarcoding studies represents only a small
fraction of that contained within assembled metagenomic DNA (Supplementary Figure 5). The
cross-alignment was limited: only 14.3% of our metagenomic OTUs were matched by an
amplicon sequence, while only 72.2% of the short amplicon OTUs (short OTUs) aligned to our
metagenomic database. This disparity was particularly pronounced for novel lineages. For
instance, the only OTU from an orphan major clade (NL) that aligned with a short-read sequence
belonged to NL9; this corresponding amplicon sequence variant exhibited low abundance
(sum_reads_asv=233). Thus, the novel lineages we identified from metagenome assemblies
have been largely missed by the amplicon-based studies cataloged in the MetaPR2 database
(Vaulot et al. 2022). Further supporting this conclusion, we observed a small but significant
negative Spearman correlation (rho = -0.15, p << 0.05) between the weighted environmental
abundance of OTUs and their calculated mean primer-binding bias score. Our results highlight

the limitations of amplicon-centric surveys and underscore the necessity of using metagenome
assemblies to explore eukaryotic diversity, especially for the discovery and phylogenetic
placement of novel groups.
549
550

551
552
Figure 3. Distribution of new lineages of Eukaryotes. A) Maximum-likelihood tree derived from a
concatenated alignment of 18S rRNA and 28S rRNA genes representing reference sequences (branches
colored according to division-level taxonomy) and orphan 18S rRNA gene sequences (black branches).
Clades likely indicative of major new lineages (NL) are highlighted with colored segments. Dark purple
circles denote orphan contigs that encompass a 28S rRNA gene sequence. Filled circles indicate
conserved placements maintained upon adding the LSU sequence to the multiple sequence alignment.
Purple bars within NL41 represent SAG-derived 18S rRNA gene sequences. B) Distribution of primer
binding bias scores across groups of 18S rRNA gene sequences aggregated from our combined database.
All comparisons exhibited an adjusted p value = 0 in paired Mann-Whitney tests. Boxes show interquartile
ranges, lines indicate medians, whiskers represent data spread, and circles denote outliers. C) Diagram
illustrating the GOLD Ecosystem type category assigned to the sequencing samples containing sequences
from the newly identified lineages. D) Maximum likelihood tree based on a concatenated alignment of 341
protein sequences. Stars denote SAGs from which 18S rRNA gene sequences were extracted, branching
within NL41 in the phylogenetic tree represented in A.

Prediction of Associations between Microbes Across the Globe

Microbial eukaryotes are documented to coexist with other microbes, some serving as hosts for bacterial symbionts, which may function as either ecto- or endosymbionts. These relationships can provide nutrient benefits but may also confer defense mechanisms and/or environmental adaptability (Gast et al. 2009; Husnik et al. 2021). Our abundance correlation results predict associations among unclassified eukaryotes and bacteria annotated as Verrucomicrobia, ABY1, and unclassified bacteria in marine ecosystems (n=7, 5, and 5, respectively; Figure 4A). OTUs categorized within various families of the Amoebozoa supergroup demonstrated a relatively high incidence of positive correlations with prokaryotic OTUs identified as Vicinamibacteraceae (n=15), Ardenticatenales (n=25), and Bacteroidiota (n=13) in freshwater ecosystems (Figure 4B), as well as Rickettsiales (n=5) and Microgenomata (n=4) in soil environments (Figure 4C). Several network-predicted associations likely represent genuine symbioses. This is exemplified by interactions involving Rickettsiales bacteria (identified via BLAST: >80% identity, >700 bp alignment length), a lineage well-documented as endosymbionts of amoebozoans, algae, and ciliates (Schulz et al. 2025; Schulz et al. 2016; Lanzoni et al. 2019). The congruence between these network-inferred interactions and established ecological relationships supports the biological relevance of our predictions. Other associations likely reveal environmental or feeding preferences, as suggested by prior studies (Oliverio et al. 2020).

Correlations among various eukaryotic lineages within marine environments also revealed possible associations between unclassified eukaryotes at the supergroup level and unclassified sequences associated with groups such as Heterolobosea (n=6), NL39 (n=5), and NL41 (n=3; Figure 4D). Freshwater ecosystems exhibited the highest number of correlations overall (n=1,809), with sequences from Mastigamoeba and Heterolobosea representing 549 pairs (Figure 4E). The anaerobic growth preferences of many amoebae within these groups, particularly members of the Psalteriomonadidae family, coinciding with the anaerobic lifestyle of some heteroloboseans (Pánek et al. 2012) may account for this finding. We also identified 32 eukaryote-eukaryote co-occurrences across the soil samples, primarily involving unclassified groups of Heterolobosea and eukaryotes unclassified at the supergroup level (n=12; Figure 4F). The latter group was further correlated with opisthokonts, amoebozoans, and rotifers (n=4, 4, and 3, respectively). Some members of the new lineages identified in this work were found correlated with other eukaryotes, such as NL39 and NL41 correlated with unclassified eukaryotes in marine ecosystems, NL30 and NL41 with Mastigamoebidae in freshwater, and NL22 with heteroloboseans in soil. These associations could represent host-parasite interactions considering the highly divergent sequences of these new lineages, which is often a signature of an intracellular lifestyle (Vossbrinck et al. 1987). Predatory protists are known to prey upon both bacteria and other eukaryotes, and certain species of amoebozoans and heteroloboseans exhibit species-specific feeding habits (Berlinches de Gea et al. 2024). Our findings could facilitate the identification of more eukaryotic species-specific predatory relationships between members of these recognized groups and the newly identified eukaryotic lineages.

The proportion of correlated OTU pairs, whether classified to an existing supergroup, orphan, or assigned to a new lineage, varied across ecosystem types (Figure 4G). In marine and soil ecosystems, a predominance of associations occurred between classified and orphan OTUs

(0.79 and 0.82, respectively), while in freshwater systems, the majority of correlated pairs
comprised two successfully classified OTU representatives (0.82). This trend became even more
evident when comparing the proportions of taxon pairs with correlated pairs (Figure 4H), noting a
reduced ratio of taxon pairs classified at the supergroup level in freshwater (0.73). Likewise, we
found a reduced proportion of associations between orphan taxa and taxa classified at the
supergroup level (0.72). Within marine ecosystems, the relationships between orphan OTUs and
those classified as new lineages represented the second-largest proportion (0.21 and 0.2 for
correlated OTU pair count and taxa with correlated pairs, respectively). Established routine
sampling methods, often optimized for water quality monitoring and pathogen detection, seem to
have led to an increased effort but also bias in the exploration of protist taxa in freshwater
ecosystems (Dodds 2005). Soil has been characterized as the ecosystem type with the highest
novelty regarding documented 18S rRNA gene sequences and marine the ecosystem type with
even less novelty than freshwater ecosystems (Singer et al. 2021). Thus, the more frequently
recovered predicted associations between known taxa in freshwater can likely be attributed to the
bulk of previous work, alongside the fact that the taxa exhibiting the most associations
(amoebozoans and discobans) are found to be phylogenetically diverse and may establish
relationships with other eukaryotes. Conversely, the higher proportion of associations involving
known and orphaned taxa that were found in this study underscores the significance of
associations formed by understudied protist groups within marine and soil ecosystems.
630

631
Figure 4. Association between abundances of eukaryotes and prokaryotes across ecosystem
types. Heatmaps illustrating the count of significant positive correlations between 18S and 16S rRNA gene

sequences (A, B, and C) as well as between 18S and 16S rRNA sequences (D, E, and F) aggregated at the
family level. Values are represented for family pairs exhibiting a minimum of three significant (adjusted p-
value < 0.05) correlations. G. Proportions of the total number of correlated pairs (n) between each cluster
type according to their taxonomic classification (Classified, Orphan, New Lineage) for each ecosystem
type regarding abundance correlations among eukaryotic sequences. H. Proportion of the total number of
taxa manifesting correlated pairs between each cluster type across each ecosystem type.

Conclusion

Our comprehensive, metagenome-based survey of eukaryotic diversity substantially expands the known
breadth of protistan lineages and reveals previously unseen branches across the eukaryotic tree. By
leveraging assembled metagenomes to circumvent the inherent biases of amplicon-based approaches, we
captured novel lineages belonging to taxonomic blind spots that often lack cultured representatives and
inhabit a broad range of global environments. The identified protists include largely divergent groups within
Excavata and previously uncharacterized taxa branching deeply in the eukaryotic tree. We further
uncovered likely associations between understudied protists and bacteria, hinting at intricate symbiotic or
predator-prey relationships that shape community structure. Together, these findings set the stage for
more targeted cultivation efforts and future genomic characterization, and ecological studies aimed at
illuminating the roles of these enigmatic lineages and their partners. Together, these results advance and
refine our understanding of eukaryotic diversity by providing a global, contig-based census that
complements amplicon and single-cell surveys and prioritizes underrepresented lineages for targeted
follow-up.

Methods

rRNA Sequence Search

We compiled assembled contig files corresponding to 32,520 public datasets derived from
metagenome, metatranscriptome, and single-cell sort sequencing projects marked as both public
and unrestricted within the Integrated Microbial Genome/Metagenome (IMG/M) database. The
cmsearch program from the Infernal (Nawrocki and Eddy 2013) package was utilized to search
the contig files for sequence matches to each of the following small subunit rRNA (SSU rRNA)
models from the Rfam database (Kalvari et al. 2018): eukaryotic 18S rRNA (RF01960),
microsporidial 18S rRNA gene (RF02542), archaeal 16S rRNA gene (RF01959), and bacterial
16S rRNA gene (RF00177). Employing the --anytrunc option in cmsearch allowed for the
inclusion of long SSU rRNA sequences in our analysis, enabling us to parse the output for contigs
displaying sequential and multiple non-overlapping alignments to the covariance models
(Supplementary Methods). Furthermore, long subunit rRNA (LSU rRNA) sequences were
screened using the eukaryotic 28S rRNA gene (RF02543), archaeal 23S rRNA gene (RF02540),
and bacterial 23S rRNA gene (RF02541) covariance models from Rfam, utilizing cmsearch with
default parameters.

Preliminary Annotation

We searched the retrieved SSU rRNA sequences against a database that contained all SSU rRNA sequences from both Silva (Quast et al. 2013) and PR2 (Guillou et al. 2013) databases using the BLAST+ (Camacho et al. 2009) program with an e-value cutoff of 0.005 and considering hits with sequence identity % > 70 to account for potential divergent sequences. Sequences with no hits to the database were added to a reference SSU rRNA tree to confirm their phylogenetic affiliation at the domain level. The reference tree was built as follows: SSU rRNA from the Silva and PR2 databases were clustered at 75% identity to group higher groups and further reduce redundancy using Usearch (Edgar 2010) and the resulting centroids were aligned with Mafft (Katoh and Standley 2013) using the Linsi algorithm. Tree building was done with IQtree (Nguyen et al. 2015) with default parameters and model GTR+F+R10. Sequences that branched within a cluster that belonged to a different domain were considered misannotations and subsequently removed from the alignment and the tree then rebuilt with the filtered alignment. Sequences confirmed by tree building to belong to the domain Eukarya were added to the 18S rRNA gene sequence database together with those that had a best hit with 18S rRNA gene sequences in Silva and PR2. We retained sequences with a length >= 800 bp, as this threshold represents approximately half of the SSU rRNA gene and is sufficient to capture the conserved regions essential for robust phylogenetic analyses. This search and filtering protocol yielded 157,956 18S rRNA gene sequences derived from 18,444 distinct datasets.

rRNA SSU Sequence Clustering

We assembled a database inclusive of 18S rRNA gene sequences from the Silva database (Quast et al. 2013), the PR2 database (Guillou et al. 2013), sequences referenced in the description of the Provora supergroup (Tikhonenkov et al. 2022), long amplicon sequences from a recent study (Jamy et al. 2022), sequences extracted from eukaryotic genomes housed in GenBank (Sayers et al. 2022) represented within the EukProt database (Richter et al. 2022), as well as environmental sequences sourced from IMG/M in February 2022 (Chen et al. 2023). Two-step clustering was conducted using the Usearch program with the -cluster_fast option, initially targeting 97% identity and subsequently at 85% identity to account for the high variability of 18S rRNA gene sequences in related taxa. Following the removal of clusters containing only a single member, we obtained 13,238 sequence clusters (OTUs), wherein the largest cluster comprised 18,608 sequences.

Concurrently, SSU rRNA sequences extracted from the IMG/M database, those recognized as having a 16S rRNA gene as their best hit during BLAST searches, were clustered at 97% identity and subsequently at 85% together with the 16S rRNA gene sequences from Silva and PR2 databases. Clusters with only a single member were excluded from analysis. Taxonomic annotations for each sequence were inferred from the cluster representative's best hit and reformatted to be compatible with the NCBI Taxonomy structure. This procedure produced a total of 839,222 16S rRNA gene sequences divided among 23,233 clusters, representing Bacteria, Mitochondria, Archaea, and Plastids (Supplementary Figure 6).

Primer Binding Analysis

The software primer prospector was employed to search for four different primer pairs (Hugerth et al. 2014; Amaral-Zettler et al. 2009; Nolte et al. 2010; Stoeck et al. 2010) across each 18S rRNA gene sequence from the non-singleton 85% identity clusters. Sequences with an overall weighted score exceeding 1 for any forward or reverse primer were classified as undetected by respective primer pairs. We aggregated the weighted score of all four primer pairs to calculate the binding bias score and subsequently summarized the binding bias score at the cluster level, determining the mean bias score of all sequences within each cluster. We compared the distribution of primer binding bias scores across different dataset groups: reference sequences (originating from Silva, PR2, Provora, Genbank, and long amplicons), successfully classified environmental sequences, orphan environmental sequences (lacking classification at the supergroup level), and orphan sequences representing potential new lineages. To assess significance, we performed a Kruskal-Wallis test, followed by paired two-tailed Mann-Whitney tests across pairs, adjusting p-values via the Benjamini-Hochberg procedure using the scipy.stats module v1.16 (Virtanen et al. 2020). A p value < 0.05 was deemed statistically significant.

18S and 28S rRNA Gene Tree Construction

We selected a subset of 1,247 cluster representative sequences based on specific inclusion criteria: 1) both the 18S and 28S rRNA gene sequences could be extracted from the same contig; 2) 18S sequences exhibited lengths of >= 1,200 bp; 3) 28S sequences displayed lengths of >= 2,800 bp; and 4) centroids were sourced from environmental samples from IMG/M or from long amplicon sequencing (Jamy et al. 2022). Employing the cmalign program and the RF01960 model, we aligned 18S rRNA gene sequences, and utilized the RF02543 model to align 28S rRNA gene sequences, employing the --match-only option to restrict alignment sites to those represented within the model. We concatenated sequences from both rRNA genes utilizing AMAS (Borowiec 2016) and constructed a tree through IQTree applying the GTR+F+R10 model.

To enhance the tree, we removed terminal branches exhibiting lengths > 0.5. We then recovered the original versions of the remaining sequences and separately aligned the 28S and 28S rRNA gene sequences using mafft-linsi software. Subsequently, we concatenated the alignments of both genes and employed the trimAl program (Capella-Gutiérrez et al. 2009) with the -gt 0.1 option to exclude sites with significant indel content. Afterwards, we used the IQtree program with the GTR+F+R10 model along with 1,000 ultrafast bootstrap replicates to construct a phylogenetic tree. The resultant tree underwent additional manual inspection in which sequences branching within a cluster dominated by sequences from differing supergroups were excluded from the concatenated alignment. The aforementioned removal rogue taxa has the trade-off of masking true diversity that could result from misclassification of some groups, but improves tree stability for downstream analyses (Aberer et al. 2012). The final tree included 577 sequences.

Phylogenetic Placement Based Annotation

We selected an outgroup comprising 38 bacterial, archaeal, and organellar genomes from GenBank (Supplementary Table 4), aligning the respective SSU and LSU rRNA gene sequences with the RF01960 and RF02543 covariance models utilizing the cmalign program with the –match-only and --ifile options. A phylogenetic tree was developed using sequences from the eukaryotic 18S and 28S tree alongside outgroup SSU and LSU rRNA sequences, constructed by IQtree employing the GTR+F+R10 model and incorporating 1,000 ultrafast bootstrap replicates. All cluster representatives of 18S rRNA gene sequences were aligned with the RF01960 covariance model using the --match-only option, subsequently appending a tail of gaps (-) to each sequence to account for the absence of the corresponding 28S rRNA gene sequence. We utilized the EPA-NG software (Barbera et al. 2019) to place aligned centroid sequences into the reference tree with the inclusion of the added outgroup, assigning taxonomy to the placed sequences based on the reference tree taxonomy via the Gappa software (Czech et al. 2020). We implemented a hierarchical matching algorithm to harmonize Silva annotations with the PR2 taxonomic framework. This adopted framework designates clades of unconfirmed monophyly using an "_X" suffix (Guillou et al. 2013). The algorithm first queried the PR2 database for an exact match to each Silva annotation at its lowest taxonomic rank. If a match was identified, the complete higher-level PR2 classification was inherited. If no match was found at the lowest rank, the algorithm iteratively ascended the Silva taxonomic hierarchy, searching at each progressively higher rank until a corresponding entry was found in PR2, at which point that classification was assigned.

18S rRNA Insertion Distribution Analysis

Using the per-cluster 18S rRNA representative sequence insert information produced by the cmalign program with the RF01960 covariance model and the --ifile option, we recovered only those inserts with length >= 10 bp. Then we obtained and compared the distribution of both the number of inserts and length of inserts for each eukaryotic division. To assess significance, we performed a Kruskal-Wallis test, followed by paired two-tailed Mann-Whitney tests across pairs, adjusting p-values via the Benjamini-Hochberg procedure using the scipy.stats module v1.16 (Virtanen et al. 2020). A p value < 0.05 was considered statistically significant.

Global Distribution Analysis

We extracted longitude and latitude data from the IMG/M database metadata and visualized it through the geopandas module for Python3. Latitude data was digitized into a discrete scale with a step size of 10 degrees. Distribution was visualized using the geopandas module (Jordahl et al. 2022) for Python3. Abundance was defined as the average of the normalized mean sequencing depth across all contigs pertaining to each taxonomic group, while ubiquity was defined as the proportion of samples from comparable environments and latitudes in which each taxonomic group was identified. For the evaluation of eukaryotic family distribution, we focused exclusively on those groups attaining ubiquity and abundance at or above the 80th percentile to emphasize those groups that were highly abundant and ubiquitous.

Sequence Read Depth

We retrieved data on per-contig average read depths across metagenomic datasets whenever available from IMG/M, employing this information to estimate the sample-wide average read depth via the following expression:

$$Sample\,average\,depth = \frac{\sum_{i}^{n} \left(contig_i\,average\,read\,depth \times contig_i\,length \right)}{\Box}$$

The sample average depth was subsequently employed to compute the normalized average read depth for every contig harboring an SSU rRNA gene in the dataset.

Ecosystem Diversity Analysis

For this analysis, we exclusively considered samples from the three most prevalent GOLD ecosystem types represented in our dataset: Marine, Freshwater, and Soil. Given that the metrics utilized for analyzing diversity rely on count data, we simulated read counts based on the normalized average read depth data, employing the expression:

$$Simulated\,read\,count = \frac{Normalized\,average\,read\,depth}{Simulated\,read\,length}$$

With a simulated read depth of 100 bp, read counts were rounded to the nearest integer. Contigs with a simulated read count of 0 were subsequently removed. To account for sequencing disparity, we executed a random rarefaction employing the rrarefy function from R version (R Core Team, 2013) utilizing the vegan package (Oksanen et al. 2013) to align read counts with that of the smallest sample.

To visualize the 18S rRNA gene OTU sampling effort, we employed a per-sample basis rarefaction curve for each distinct ecosystem type, estimating the mean observed OTUs per new sample across 10,000 randomizations. We then fitted our accumulation curves to various linear and nonlinear models, as outlined below:

$$Simple\,linear: y = bX$$
$$Logarithmic: y = \log \left(X \right)$$
$$Power: y = X^b$$
$$Exponential: y = b^X$$
$$Rational: y = \left(a + bX \right) / \left(1 + cX \right)$$

We used the Python ols function from the statsmodels module (Seabold and Perktold 2010) and the curve_fit function from the scipy module v1.16 (Virtanen et al. 2020) for model fitting. The best-fitting model for the accumulation curve was determined using the Akaike Information

824   Criterion (AIC). The rarefaction curves for the Freshwater and Marine ecosystem types were
825   comparable, exhibiting overlaps. We fitted the rarefied count data to five distinct models,
826   inclusive of an asymptotic rational model. Across all assessments, the power function constituted
827   the best fit, confirming a non-asymptotic relationship.
828
Subsequently, we utilized the Python3 ecopy module (https://github.com/Auerilas/ecopy) to
quantitate Shannon H, Simpson, and Inverse Simpson diversity indices on the rarefied OTU
table. Second-order polynomial linear regressions were applied to model the relationship
between diversity (measured by the Simpson index) and absolute latitude. For this analysis, we
segmented the OTU table for each taxon across varying taxonomic levels: supergroup, division,
and subdivision, limiting our focus to taxonomic groups present in a minimum of 10 samples. To
account for data partitioning, we adjusted p-values for polynomial fits using the Benjamini-
Hochberg procedure.
837

Search For Potential New Lineages

We constructed a phylogenetic tree utilizing OTU representative sequences classified under the
eukaryotic domain and categorized unknown supergroups. These sequences were combined
with data from the reference tree employed for phylogenetic placement analysis. Using
concatenated alignments for the 18S and 28S rRNA genes based on the RF01960 and RF02543
covariance models, we constructed the tree employing the IQtree program with the GTR+F+R10
model and 1,000 ultrafast bootstrap replicates. Clades comprising at least five leaves classified
as eukaryotic with unknown supergroup status and belonging to clusters containing only
metagenomic sequences, along with maximum bootstrap support values, were defined as
candidates for potential new lineages. The environmental origins of samples harboring
sequences from these new lineages were visualized employing the plotting capabilities of the
plotly Python3 module (https://github.com/plotly/plotly.py).
850

Phylogenomic Analysis

To further validate the taxonomic affiliation of the potential new lineages, we used an
independent dataset of single-amplified genomes derived from cells isolated from anoxic marine
sediments in Maine. We performed a phylogenetic analysis expecting a robust classification
based on several marker genes. Initial filtering included the removal of SAGs with fewer than 100
putatively annotated eukaryotic proteins. The peptides from the remaining SAGs were
incorporated into broadly conserved gene families using PhyloToL (Cerón-Romero et al. 2019),
identifying 345 gene families well represented among the remaining SAGs. Subsequently, we
excluded SAGs with low representation within this pool of gene families (i.e., present in <80 gene
families), thereafter generating single-gene alignments, conducting homology assessments, and
selecting orthologs with PhyloToL. For each alignment, columns with more than 90% missing
data were removed with trimAl (Capella-Gutiérrez et al. 2009). Following alignment processing,
we concatenated the 341 single-gene alignments, utilizing IQTree2 (v2.2; Minh et al. 2019) for

phylogenomic reconstruction, applying the LG+C60+G+F model under the PMSF approximation
with 100 bootstrap iterations.
866

867 Tax-aliquots Clustering Method

The tax-aliquots clustering method (Krinos et al. 2024) was employed to revisit the original
metagenomes, recovering contigs akin to the non-18S rRNA gene portions of the 18S rRNA
gene-containing contigs used for identifying new lineages. We utilized mmseqs2 (version
13.45111) (Steinegger and Söding 2017) to precluster all contigs from samples from which a new
lineage was identified, establishing a threshold coverage value of 0.01 and a percent sequence
identity of 0.7, utilizing coverage mode 1. All contigs within clusters containing at least one
sequence inclusive of an 18S rRNA gene-containing contig were retained, thus minimizing the
total input size of sequence content while limiting sequence representatives to a manageable
number by restricting the search to contigs already exhibiting some degrees of similarity to the
rRNA gene-containing contigs. Subsequent to obtaining this original clustered assembly, we
differentiated sequences associated with the original 18S rRNA gene-containing contigs from
those that did not incorporate these genes. With the coordinates of the rRNA gene within the 18S
rRNA gene-containing contig, we masked the 18S rRNA gene sequences, recombining the files
as input for tax-aliquots. The reference content consisted of the masked 18S rRNA gene-
containing contigs, while query content comprised other contigs remaining subsequent to the
initial mmseqs2 filtering (employing SeqIO from Biopython for sequence masking; (Cock et al.
2009). We subsequently clustered this sequence content utilizing mmseqs2 (Steinegger and
Söding 2017) with a coverage of 0.2 and percent sequence identity of 0.2, retaining clusters that
encompassed at least one of the masked rRNA-gene containing contigs and at least one
additional contig that did not contain an 18S rRNA gene. The resulting assembly was subject to
further clustering with sequence information parsed into k-mers of length 3.

889 Short amplicon sequence mapping analysis

To understand the representation of our 18S rRNA gene-based OTUs in previous
metabarcoding-based studies, we collected a comprehensive amplicon sequence data set from
the MetaPR2 database (Vaulot, et al. 2022) belonging to the V4 region of the 18S rRNA gene
sequence from all available environments. Data from 71,043 amplicon sequence variants
(ASVs) was downloaded from the MetaPR2 web server and their sequences were clustered at
97% identity using vsearch --cluster_fast resulting in 15,532 short amplicon OTUs (short OTUs).
We used the bbmap v39.33 (https://sourceforge.net/projects/bbmap/) software to map the short
OTU representative sequences to the 13,238 metagenomic assembly OTU representative
sequences using ambig=all minid=0.80 options to capture multiple alignments per reference
sequence while also allowing sequence variation to account for the limitations of using short
sequences of the V4 region. The relationship between reference sequences and aligned
amplicon sequences was obtained using the bamtobed option in BEDTools v2.31 (Quinlan &
Hall, 2010). To get a measure of the diversity of sequence variants captured by each
metagenomic OTU in the sequence mapping we used the weighted environmental abundance
calculated using the following expression:
905

$$906 \qquad Normalized environmental abundance = \frac{Mapped shOTUs}{\log 10 \left(Average shOTU abundance \right)}$$

Short OTU abundance was obtained from the information provided in the MetaPR2 database and
used to normalize the number of aligned sequences to each OTU by the abundance of each short
OTU as more abundant short OTUs are expected to be more likely to be found and aligned to a
metagenomic OTU. To inspect the relationship between normalized environmental abundance
and predicted primer binding bias, we calculated the Spearman correlation coefficient between
the primer binding bias score and normalized environmental abundance using the spearmanr
function from scipy v 1.16 (Virtanen et al. 2020).

Correlation Between SSU rRNA Sequence Read Depth

For each of the most frequently represented ecosystem types in our dataset (Soil, Freshwater,
and Marine), we evaluated the sequences within each 18S rRNA gene cluster, retaining only
those with a minimum of three sequences across different samples from specific environments,
resulting in the acquisition of ecosystem-filtered 18S rRNA gene clusters. An analogous
procedure was conducted to retrieve ecosystem-filtered 16S clusters, from which we calculated
Spearman's rank correlation coefficients for each pair of 18S and 16S rRNA gene filtered clusters
whenever a minimum of three samples contained sequences from both clusters. Adjustments for
p-values were made using the Benjamini-Hochberg method for each GOLD ecosystem type
independently, with a replication of this process for correlations among 18S rRNA gene
sequence clusters, incorporating the additional step of excluding autocorrelations.

Acknowledgments

927 The work conducted by the U.S. Department of Energy Joint Genome Institute
928 (https://ror.org/04xm1d337), a DOE Office of Science User Facility, is supported by the Office of
929 Science of the U.S. Department of Energy operated under Contract No. DE-AC02-05CH11231.

Data availability

931 Sequence data including SSU rRNA sequences and SAG assemblies, together with tables
932 containing the full metadata from sequencing datasets from IMG/M and sequence cluster
933 information are available at: https://portal.nersc.gov/cfs/nelli/eukcensus/

Code availability

935 The pipeline for extracting long SSU rRNA sequences from metagenomic contigs developed for
936 this study is available at https://github.com/NeLLi-team/ssuextract.

Competing interests statement

F.S. serves as CEO of SampleX. This work was not funded by, nor does it benefit, SampleX. All other authors declare no competing interests.

References

Aberer, A. J., Krompass, D., & Stamatakis, A. (2013). Pruning rogue taxa improves phylogenetic accuracy: an efficient algorithm and webservice. *Systematic biology*, 62(1), 162-166.

Adl, S. M., Leander, B. S., Simpson, A. G. B., Archibald, J. M., Anderson, O. R., Bass, D., Bowser, S. S., Brugerolle, G., Farmer, M. A., Karpov, S., Kolisko, M., Lane, C. E., Lodge, D. J., Mann, D. G., Meisterfeld, R., Mendoza, L., Moestrup, Ø., Mozley-Standridge, S. E., Smirnov, A. V., & Spiegel, F. (2007). Diversity, nomenclature, and taxonomy of protists. *Systematic Biology*, *56*(4), 684–689.

Al Jewari, C., & Baldauf, S. L. (2023). An excavate root for the eukaryote tree of life. *Science Advances*, *9*(17), eade4973.

Amaral-Zettler, L. A., McCliment, E. A., Ducklow, H. W., & Huse, S. M. (2009). A method for studying protistan diversity using massively parallel sequencing of V9 hypervariable regions of small-subunit ribosomal RNA genes. *PloS One*, *4*(7), e6372.

Azovsky, A., & Mazei, Y. (2013). Do microbes have macroecology? Large-scale patterns in the diversity and distribution of marine benthic ciliates. *Global Ecology and Biogeography: A Journal of Macroecology*, *22*(2), 163–172.

Bahram, M., Hildebrand, F., Forslund, S. K., Anderson, J. L., Soudzilovskaia, N. A., Bodegom, P. M., Bengtsson-Palme, J., Anslan, S., Coelho, L. P., Harend, H., Huerta-Cepas, J., Medema, M. H., Maltz, M. R., Mundra, S., Olsson, P. A., Pent, M., Põlme, S., Sunagawa, S., Ryberg, M., … Bork, P. (2018). Structure and function of the global topsoil microbiome. *Nature*, *560*(7717), 233–237.

Barbera, P., Kozlov, A. M., Czech, L., Morel, B., Darriba, D., Flouri, T., & Stamatakis, A. (2019). EPA-ng: massively parallel evolutionary placement of genetic sequences. *Systematic biology*, 68(2), 365-369.

Berger, S. A., Krompass, D., & Stamatakis, A. (2011). Performance, accuracy, and web server for evolutionary placement of short sequence reads under maximum likelihood. *Systematic biology*, 60(3), 291-302.

Berlinches de Gea, A., Geisen, S., Grootjans, F., Wilschut, R. A., & Schwelm, A. (2024). Species-specific predation determines the feeding impacts of six soil protist species on bacterial and eukaryotic prey. *European Journal of Protistology*, *94*(126090), 126090.

Borowiec, M. L. (2016). AMAS: a fast tool for alignment manipulation and computing of
summary statistics. *PeerJ*, *4*, e1660.
Burki, F., & Pawlowski, J. (2006). Monophyly of Rhizaria and multigene phylogeny of
unicellular bikonts. *Molecular biology and evolution*, 23(10), 1922-1930.
Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T.
L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics*, *10*, 421.
Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: a tool for
automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* , *25*(15),
1972–1973.
Cerón-Romero, M. A., Maurer-Alcalá, X. X., Grattepanche, J.-D., Yan, Y., Fonseca, M. M., &
Katz, L. A. (2019). PhyloToL: A Taxon/Gene-Rich Phylogenomic Pipeline to Explore
Genome Evolution of Diverse Eukaryotes. *Molecular Biology and Evolution*, *36*(8), 1831–
1842.
Céza, V., Kotyk, M., Kubánková, A., Yubuki, N., Šťáhlavský, F., Silberman, J. D., & Čepička,
I. (2022). Free-living Trichomonads are Unexpectedly Diverse. *Protist*, *173*(4), 125883.
Chen, I.-M. A., Chu, K., Palaniappan, K., Ratner, A., Huang, J., Huntemann, M., Hajek, P.,
Ritter, S. J., Webb, C., Wu, D., Varghese, N. J., Reddy, T. B. K., Mukherjee, S.,
Ovchinnikova, G., Nolan, M., Seshadri, R., Roux, S., Visel, A., Woyke, T., … Ivanova, N. N.
(2023). The IMG/M data management and analysis system v.7: content updates and new
features. *Nucleic Acids Research*, *51*(D1), D723–D732.
Cock, P. J. A., Antao, T., Chang, J. T., Chapman, B. A., Cox, C. J., Dalke, A., Friedberg, I.,
Hamelryck, T., Kauff, F., Wilczynski, B., & de Hoon, M. J. L. (2009). Biopython: freely
available Python tools for computational molecular biology and bioinformatics.
Bioinformatics (Oxford, England), *25*(11), 1422–1423.
Collins, W. E., & Jeffery, G. M. (2007). Plasmodium malariae: parasite and disease. *Clinical
Microbiology Reviews*, *20*(4), 579–592.
Cooney, E. C., Leander, B. S., & Keeling, P. J. (2024). Hidden syndinian and perkinsid
infections in dinoflagellate hosts revealed by single-cell transcriptomics. *The ISME Journal*,
18(1).
Cotner, J. B., & Biddanda, B. A. (2002). Small players, large role: microbial influence on
biogeochemical processes in pelagic aquatic ecosystems. *Ecosystems*, 5, 105-121.
Jordahl, K., Van den Bossche, J., Fleischmann, M., Wasserman, J.,McBride, J., Gerard, J., …
Leblanc, F. (2020). *geopandas/geopandas*: v0.8.1 (Version v0.8.1). Zenodo.
http://doi.org/10.5281/zenodo.3946761
Czech, L., Barbera, P., & Stamatakis, A. (2020). Genesis and Gappa: processing, analyzing
and visualizing phylogenetic (placement) data. *Bioinformatics*, 36(10), 3263-3265.

del Campo, J., Sieracki, M. E., Molestina, R., Keeling, P., Massana, R., & Ruiz-Trillo, I.
(2014). The others: our biased perspective of eukaryotic genomes. *Trends in Ecology &*
Evolution, *29*(5), 252–259.

Delgado-Baquerizo, M., Oliverio, A. M., Brewer, T. E., Benavent-González, A., Eldridge, D. J.,
Bardgett, R. D., ... & Fierer, N. (2018). A global atlas of the dominant bacteria found in soil.
Science, 359(6373), 320-325.

de Vargas, C., Audic, S., Henry, N., Decelle, J., Mahé, F., Logares, R., Lara, E., Berney, C.,
Le Bescot, N., Probert, I., Carmichael, M., Poulain, J., Romac, S., Colin, S., Aury, J.-M.,
Bittner, L., Chaffron, S., Dunthorn, M., Engelen, S., … Karsenti, E. (2015). Ocean plankton.
Eukaryotic plankton diversity in the sunlit ocean. *Science*, *348*(6237), 1261605.

Dodds, W. K. (2005). Freshwater Microbiology: Biodiversity and Dynamic Interactions of
Microorganisms in the Aquatic Environment. By David C Sigee. Chichester (United
Kingdom) and Hoboken (New Jersey): John Wiley & Sons. 180.00 (hardcover); 65.00
(paper). xix + 524 p; ill.; index. ISBN: 0–471–48528–4 (hc); 0–471–48529–2 (pb). 2005. *The*
Quarterly Review of Biology, *80*(3), 355–356.

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST.
Bioinformatics , *26*(19), 2460–2461.

Eglit, Y., Shiratori, T., Jerlström-Hultqvist, J., Williamson, K., Roger, A. J., Ishida, K.-I., &
Simpson, A. G. B. (2024). Meteora sporadica, a protist with incredible cell architecture, is
related to Hemimastigophora. *Current Biology: CB*, *34*(2), 451–459.e6.

Einarsson, E., Ma'ayeh, S., & Svärd, S. G. (2016). An up-date on Giardia and giardiasis.
Current Opinion in Microbiology, *34*, 47–52.

Ferla, M. P., Thrash, J. C., Giovannoni, S. J., & Patrick, W. M. (2013). New rRNA gene-based
phylogenies of the Alphaproteobacteria provide perspective on major groups, mitochondrial
ancestry and phylogenetic instability. *PloS One*, *8*(12), e83383.

Fermani, P., Metz, S., Balagué, V., Descy, J. P., Morana, C., Logares, R., Massana, R., &
Sarmento, H. (2021). Microbial eukaryote assemblages and potential novel diversity in four
tropical East African Great Lakes. *FEMS Microbiology Ecology*, *97*(9).

Fernández, L. D., Fournier, B., Rivera, R., Lara, E., Mitchell, E. A. D., & Hernández, C. E.
(2016). Water–energy balance, past ecological perturbations and evolutionary constraints
shape the latitudinal diversity gradient of soil testate amoebae in south-western South
America. *Global Ecology and Biogeography: A Journal of Macroecology*, *25*(10), 1216–
1227.

Foissner, W. (2006). Biogeography and dispersal of micro-organisms: a review emphasizing
protists. *Acta Protozoologica*, *45*, 111–136.

Foissner, W., Blatterer, H., & Foissner, I. (1988). The hemimastigophora (Hemimastix
amphikineta nov. gen., nov. spec.), a new protistan phylum from gondwanian soils.
European Journal of Protistology, *23*(4), 361–383.
Fuhrman, J. A., Steele, J. A., Hewson, I., Schwalbach, M. S., Brown, M. V., Green, J. L., &
Brown, J. H. (2008). A latitudinal diversity gradient in planktonic marine bacteria.
Proceedings of the National Academy of Sciences, 105(22), 7774-7778.
Gast, R. J., Sanders, R. W., & Caron, D. A. (2009). Ecological strategies of protists and their
symbiotic relationships with prokaryotic microbes. *Trends in Microbiology*, *17*(12), 563–569.
Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., Boutte, C., Burgaud, G., de
Vargas, C., Decelle, J., Del Campo, J., Dolan, J. R., Dunthorn, M., Edvardsen, B.,
Holzmann, M., Kooistra, W. H. C. F., Lara, E., Le Bescot, N., Logares, R., … Christen, R.
(2013). The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote
small sub-unit rRNA sequences with curated taxonomy. *Nucleic Acids Research*,
41(Database issue), D597–D604.
Hillebrand, H., & Azovsky, A. I. (2001). Body size determines the strength of the latitudinal
diversity gradient. *Ecography*, *24*(3), 251–256.
Horn, D. (2022). A profile of research on the parasitic trypanosomatids and the diseases they
cause. *PLoS Neglected Tropical Diseases*, *16*(1), e0010040.
Howe, A. T., Bass, D., Scoble, J. M., Lewis, R., Vickerman, K., Arndt, H., & Cavalier-Smith, T.
(2011). Novel cultured protists identify deep-branching environmental DNA clades of
cercozoa: New Genera Tremula, Micrometopion, Minimassisteria, Nudifila, Peregrinia.
Protist, *162*(2), 332–372.
Hugerth, L. W., Muller, E. E. L., Hu, Y. O. O., Lebrun, L. A. M., Roume, H., Lundin, D.,
Wilmes, P., & Andersson, A. F. (2014). Systematic design of 18S rRNA gene primers for
determining eukaryotic diversity in microbial consortia. *PloS One*, *9*(4), e95567.
Husnik, F., Tashyreva, D., Boscaro, V., George, E. E., Lukeš, J., & Keeling, P. J. (2021).
Bacterial and archaeal symbioses with protists. *Current Biology: CB*, *31*(13), R862–R877.
Jamy, M., Biwer, C., Vaulot, D., Obiol, A., Jing, H., Peura, S., Massana, R., & Burki, F.
(2022). Global patterns and rates of habitat transitions across the eukaryotic tree of life.
Nature Ecology & Evolution, *6*(10), 1458–1470.
Kalvari, I., Nawrocki, E. P., Argasinska, J., Quinones-Olvera, N., Finn, R. D., Bateman, A., &
Petrov, A. I. (2018). Non-Coding RNA Analysis Using the Rfam Database. *Current
Protocols in Bioinformatics / Editoral Board, Andreas D. Baxevanis ... [et Al.]*, *62*(1), e51.
Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4), 772–
780.

Krinos, A. I., Bowers, R. M., Rohwer, R. R., McMahon, K. D., Woyke, T., & Schulz, F. (2024).

Time-series metagenomics reveals changing protistan ecology of a temperate dimictic lake.

Microbiome, *12*(1), 133.

Krinos, A. I., Mars Brisbin, M., Hu, S. K., & Cohen, N. R. (2024). Missing microbial eukaryotes

and misleading meta-omic conclusions. *Nature Communications*.

Lanzoni, O., Sabaneyeva, E., Modeo, L., Castelli, M., Lebedeva, N., Verni, F.,

Schrallhammer, M., Potekhin, A., & Petroni, G. (2019). Diversity and environmental

distribution of the cosmopolitan endosymbiont "Candidatus Megaira." *Scientific Reports*, *9*.

Lara, E., Mitchell, E. A. D., Moreira, D., & López García, P. (2011). Highly diverse and

seasonally dynamic protist community in a pristine peat bog. *Protist*, *162*(1), 14–32.

Lara, E., Roussel-Delif, L., Fournier, B., Wilkinson, D. M., & Mitchell, E. A. D. (2016). Soil

microorganisms behave like macroscopic organisms: patterns in the global distribution of

soil euglyphid testate amoebae. *Journal of Biogeography*, *43*(3), 520–532.

Lax, G., Eglit, Y., Eme, L., Bertrand, E. M., Roger, A. J., & Simpson, A. G. B. (2018).

Hemimastigophora is a novel supra-kingdom-level lineage of eukaryotes. *Nature*,

564(7736), 410–414.

Mahé, F., de Vargas, C., Bass, D., Czech, L., Stamatakis, A., Lara, E., Singer, D., Mayor, J.,

Bunge, J., Sernaker, S., Siemensmeyer, T., Trautmann, I., Romac, S., Berney, C., Kozlov,

A., Mitchell, E. A. D., Seppey, C. V. W., Egge, E., Lentendu, G., … Dunthorn, M. (2017).

Parasites dominate hyperdiverse soil protist communities in Neotropical rainforests. *Nature*

Ecology & Evolution, *1*(4), 91.

Meyer, A., Todt, C., Mikkelsen, N. T., & Lieb, B. (2010). Fast evolving 18S rRNA sequences

from Solenogastres (Mollusca) resist standard PCR amplification and give new insights into

mollusk substitution rate heterogeneity. *BMC Evolutionary Biology*, *10*, 70.

Minh, B. Q., Trifinopoulos, J., Schrempf, D., Schmidt, H. A., & Lanfear, R. (2019). IQ-TREE

version 2.0: tutorials and Manual Phylogenomic software by maximum likelihood. *URL*

Http://www. Iqtree. Org.

Mittelbach, G. G., Schemske, D. W., Cornell, H. V., Allen, A. P., Brown, J. M., Bush, M. B., ...

& Turelli, M. (2007). Evolution and the latitudinal diversity gradient: speciation, extinction

and biogeography. *Ecology letters*, 10(4), 315-331.

Morey, K. C., Myler, E., Hanner, R., & Tetreault, G. (2024). Taxonomic blind spots: A

limitation of environmental DNA metabarcoding-based detection for Canadian freshwater

fishes. *Environmental DNA (Hoboken, N.J.)*, *6*(6), e70054.

Nawrocki, E. P., & Eddy, S. R. (2013). Infernal 1.1: 100-fold faster RNA homology searches.

Bioinformatics , *29*(22), 2933–2935.

Newton, R. J., Jones, S. E., Eiler, A., McMahon, K. D., & Bertilsson, S. (2011). A guide to the
natural history of freshwater lake bacteria. *Microbiology and molecular biology reviews*,
75(1), 14-49.
Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular*
Biology and Evolution, *32*(1), 268–274.
Nolte, V., Pandey, R. V., Jost, S., Medinger, R., Ottenwälder, B., Boenigk, J., & Schlötterer,
C. (2010). Contrasting seasonal niche separation between rare and abundant taxa conceals
the extent of protist diversity. *Molecular Ecology*, *19*(14), 2908–2915.
Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'hara, R. B., &
Oksanen M. (2013). Package "vegan." *Community Ecology Package, Version*, *2*(9), 1–295.
Oliverio, A. M., Geisen, S., Delgado-Baquerizo, M., Maestre, F. T., Turner, B. L., & Fierer, N.
(2020). The global-scale distributions of soil protists and their contributions to belowground
systems. *Science Advances*, *6*(4), eaax8787.
Overgaard, C. K., Jamy, M., Radutoiu, S., Burki, F., & Dueholm, M. K. D. (2024).
Benchmarking long-read sequencing strategies for obtaining ASV-resolved rRNA operons
from environmental microeukaryotes. *Molecular Ecology Resources*, e13991.
Pánek, T., Silberman, J. D., Yubuki, N., Leander, B. S., & Cepicka, I. (2012). Diversity,
evolution and molecular systematics of the Psalteriomonadidae, the main lineage of
anaerobic/microaerophilic heteroloboseans (excavata: discoba). *Protist*, *163*(6), 807–831.
Poulin, M., Daugbjerg, N., Gradinger, R., Ilyash, L., Ratkova, T., & von Quillfeldt, C. (2011).
The pan-Arctic biodiversity of marine pelagic and sea-ice unicellular eukaryotes: a first-
attempt assessment. *Marine Biodiversity: A Journal of the Senckenberg Research Institute /*
Senckenberg Forschungsinstitut Und Naturmudeum, *41*(1), 13–28.
Pruitt, K., Tatusova, T., & Maglott, D. (2006). NCBI reference sequences (RefSeq): a curated
non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids*
Research, *35*, D61–D65.
Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., &
Glöckner, F. O. (2013). The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Research*, *41*(Database issue), D590–
D596.
Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics*, 26(6), 841-842.
Reboul, G., Moreira, D., Bertolino, P., Hillebrand-Voiculescu, A. M., & López-García, P.
(2019). Microbial eukaryotes in the suboxic chemosynthetic ecosystem of Movile Cave,
Romania. *Environmental Microbiology Reports*, *11*(3), 464–473.

Richter, D. J., Berney, C., Strassert, J. F. H., Poh, Y.-P., Herman, E. K., Muñoz-Gómez, S. A.,
Wideman, J. G., Burki, F., & de Vargas, C. (2022). EukProt: A database of genome-scale
predicted proteins across the diversity of eukaryotes. *Peer Community Journal*, *2*(e56).

Sayers, E. W., Bolton, E. E., Brister, J. R., Canese, K., Chan, J., Comeau, D. C., Connor, R.,
Funk, K., Kelly, C., Kim, S., Madej, T., Marchler-Bauer, A., Lanczycki, C., Lathrop, S., Lu, Z.,
Thibaud-Nissen, F., Murphy, T., Phan, L., Skripchenko, Y., … Sherry, S. T. (2022).
Database resources of the national center for biotechnology information. *Nucleic Acids
Research*, *50*(D1), D20–D26.

Schuler, G. A., Tice, A. K., Pearce, R. A., Foreman, E., Stone, J., Gammill, S., ... & Brown, M.
W. (2018). Phylogeny and classification of novel diversity in Sainouroidea (Cercozoa,
Rhizaria) sheds light on a highly diverse and divergent clade. *Protist*, 169(6), 853-874.

Schulz, F., Martijn, J., Wascher, F., Lagkouvardos, I., Kostanjšek, R., Ettema, T. J. G., &
Horn, M. (2016). A Rickettsiales symbiont of amoebae with ancient features. *Environmental
Microbiology*, *18*(8), 2326–2342.

Schulz, F., Yan, Y., Weiner, A. K., Ahsan, R., Katz, L. A., & Woyke, T. (2025). Single-cell
genomics reveals complex microbial and viral associations in ciliates and testate amoebae.
Nature Communications, 16(1), 10336.

Seabold, S., & Perktold, J. (2010). Statsmodels: econometric and statistical modeling with
python. *SciPy*, *7*(1).

Singer, D., Seppey, C. V. W., Lentendu, G., Dunthorn, M., Bass, D., Belbahri, L., Blandenier,
Q., Debroas, D., de Groot, G. A., de Vargas, C., Domaizon, I., Duckert, C., Izaguirre, I.,
Koenig, I., Mataloni, G., Schiaffino, M. R., Mitchell, E. A. D., Geisen, S., & Lara, E. (2021).
Protist taxonomic and functional diversity in soil, freshwater and marine ecosystems.
Environment International, *146*, 106262.

Steinegger, M., & Söding, J. (2017). MMseqs2 enables sensitive protein sequence searching
for the analysis of massive data sets. *Nature Biotechnology*, *35*(11), 1026–1028.

Stoeck, T., Bass, D., Nebel, M., Christen, R., Jones, M. D. M., Breiner, H.-W., & Richards, T.
A. (2010). Multiple marker parallel tag environmental DNA sequencing reveals a highly
complex eukaryotic community in marine anoxic water. *Molecular Ecology*, *19 Suppl 1*, 21–
31.

Team, R. C., & Others. (2013). R: A language and environment for statistical computing.
Foundation for Statistical Computing, Vienna, Austria.

Tekle, Y. I., Wang, F., Wood, F. C., Anderson, O. R., & Smirnov, A. (2022). New insights on
the evolutionary relationships between the major lineages of Amoebozoa. *Scientific reports*,
12(1), 11173.

Tikhonenkov, D. V., Mikhailov, K. V., Gawryluk, R. M. R., Belyaev, A. O., Mathur, V., Karpov,
S. A., Zagumyonnyi, D. G., Borodina, A. S., Prokina, K. I., Mylnikov, A. P., Aleoshin, V. V., &

Keeling, P. J. (2022). Microbial predators form a new supergroup of eukaryotes. *Nature*,
612(7941), 714–719.

Torruella, G., De Mendoza, A., Grau-Bové, X., Antó, M., Chaplin, M. A., Del Campo, J., ... &
Ruiz-Trillo, I. (2015). Phylogenomics reveals convergent evolution of lifestyles in close
relatives of animals and fungi. *Current Biology*, 25(18), 2404-2410.

Torruella, G., Galindo, L. J., Moreira, D., & López-García, P. (2025). Phylogenomics of
neglected flagellated protists supports a revised eukaryotic tree of life. *Current Biology*,
35(1), 198-207.

Urich, T., Lanzén, A., Qi, J., Huson, D. H., Schleper, C., & Schuster, S. C. (2008).
Simultaneous assessment of soil microbial community structure and function through
analysis of the meta-transcriptome. *PloS One*, *3*(6), e2527.

Vaulot, D., Sim, C. W. H., Ong, D., Teo, B., Biwer, C., Jamy, M., & Lopes dos Santos, A.
(2022). metaPR2: a database of eukaryotic 18S rRNA metabarcodes with an emphasis on
protists. *Molecular Ecology Resources*, 22(8), 3188-3201.

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,
Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson,
J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., … SciPy
1.0 Contributors. (2020). SciPy 1.0: fundamental algorithms for scientific computing in
Python. *Nature Methods*, *17*(3), 261–272.

Vossbrinck, C. R., Maddox, J. V., Friedman, S., Debrunner-Vossbrinck, B. A., & Woese, C. R.
(1987). Ribosomal RNA sequence suggests microsporidia are extremely ancient
eukaryotes. *Nature*, *326*(6111), 411–414.

Williamson, K., Eme, L., Baños, H., McCarthy, C. G., Susko, E., Kamikawa, R., ... & Roger, A.
J. (2025). A robustly rooted tree of eukaryotes reveals their excavate ancestry. *Nature*, 1-8.

Woyke, T., Doud, D. F. R., & Schulz, F. (2017). The trajectory of microbial single-cell
sequencing. *Nature Methods*, *14*(11), 1045–1054.

Wuyts, J., Van de Peer, Y., & De Wachter, R. (2001). Distribution of substitution rates and
location of insertion sites in the tertiary structure of ribosomal RNA. *Nucleic Acids Research*,
29(24), 5017–5028.

Xie, Q., Lin, J., Qin, Y., Zhou, J., & Bu, W. (2011). Structural diversity of eukaryotic 18S rRNA
and its impact on alignment and phylogenetic reconstruction. *Protein & Cell*, *2*(2), 161–170.

Zancan, S., Trevisan, R., & Paoletti, M. G. (2006). Soil algae composition under different
agro-ecosystems in North-Eastern Italy. *Agriculture, Ecosystems & Environment*, *112*(1), 1–
12.

Zhang, Z.-F., Liu, L.-R., Pan, Y.-P., Pan, J., & Li, M. (2023). Long-read assembled
metagenomic approaches improve our understanding on metabolic potentials of microbial
community in mangrove sediments. *Microbiome*, *11*(1), 188.

Zhao, J., Fan, D., Guo, W., Wu, J., Zhang, X., Zhuang, X., & Kong, W. (2023). Precipitation
drives soil protist diversity and community structure in dry grasslands. *Microbial Ecology*,
86(4), 2293–2304.

Supplementary Figures

Supplementary figure 1. Summary of the methods used in this work. Cylinders correspond to sequence
data, while the ovals indicate phylogenies, stars indicate metadata, and squares represent analyses. Some
final deliverables include an F and one or two numbers in parenthesis, which indicate the figure of the main
text in which they are included. "Iterative refinement" refers to the process of manually inspecting the tree
and removing sequences representing rogue taxa and long branches that could produce artifacts in the
topology from the seed alignment and rebuilding the tree.

1234

1235
1236   Supplementary figure 2. Distribution of clusters comprising any given number of different taxa within a
1237   single reference sequence-only cluster.

1238

1239
1240 Supplementary figure 3. Per-sample diversity distribution for each specific ecosystem with at least 1%
1241 share of the datasets separated by ecosystem types. Box plots display medians (horizontal lines),
1242 interquartile ranges (boxes), and data distribution (whiskers), with outliers shown as circles.
1243

1244
1245   Supplementary figure 4. Contribution of each ecosystem subtype and ecosystem type to the observed
1246   pool of 18S rRNA gene sequences attributable to the 13 new lineages.
1247

1248
Supplementary figure 5. 18S rRNA gene V4 region short amplicon sequence mapping to long
metagenomic assembly OTU representative sequences. A. Number of metagenomic OTU representatives
for each eukaryotic division colored by their alignment status to short amplicon OTU (shOTU)
representatives. B. Number of metagenomic OTU representatives for each of the new lineages presented
in this work colored by their alignment status to shOTU representative sequences. C. Number of
metagenomic OTU representatives that were aligned at least once with shOTU representative sequences
colored by the cluster type regarding the source of the sequences belonging to each OTU. D. Number of
metagenomic OTU   representatives that were not aligned at least once with shOTU representative
sequences colored by the cluster type. E. Environmental abundance for each eukaryotic division
expressed as the total number of alignments of metagenomic OTU representative sequences and shOTU
representatives. F. Relationship between mean primer binding bias score and the weighted environmental
abundance for each metagenomic OTU with alignments to short read sequences.

Supplementary figure 6. Abundance of clusters for each prokaryotic taxonomic group representing at least
0.5% of the total number of clusters within the analyzed environmental samples. Bars are partitioned
according to their share of cluster types as defined by the origin of the member sequences. REF_IMG:
includes both sequences from IMG/M and from reference databases; REF_only: includes sequences from
reference databases exclusively.

1268 Supplementary Tables

Supplementary Table 1. Number of sequencing datasets with observed 18S rRNA sequences by
Ecosystem Subtype.
Supplementary Table 2. Counts of samples with no observed 18S rRNA sequences by
Ecosystem Subtype.
Supplementary Table 3. Mean ubiquity, abundance and weighted abundance data of 18s rRNA
sequences among ecosystem types at the family level.
Supplementary Table 4. Information on selected genomes of archaea, bacteria, mitochondria
and plastids used as outgroup for the tree built for phylogenetic placement-based taxonomic
classification.
Supplementary Table 5. Metadata and frequency of 18S rRNA genes for samples in which new
lineages were found.

Supplementary Methods

1280

To search for rRNA sequences taking into account the extreme variations in length due to introns
and other additional sequences in eukaryotic genomes, we developed a workflow named
"ssuextract". This workflow uses the Infernal software (Nawrocki and Eddy 2013) to search a
covariance model (CM) to nucleic acid sequencing contigs and search homologs using blastn
(Camacho et al. 2009) against the Silva (Quast et al. 2013) and PR2 (Guillou et al. 2013)
databases as a first approach to get a taxonomic classification for each sequence. It was
designed with the 18S rRNA gene in mind, but it can also be used to extract 28S, 16S, 23S rRNA
gene sequences, and practically any nucleic acid sequence for which a covariance model is
available.

The key component of this workflow is the detection of long sequences represented by the
covariance model through multiple sequential hits along a single contig reported in the cmsearch
output table produced using the –anytrunc option (i.e. a fragmented alignment). Full alignments
to the model in which the start and end position of the covariance model are found within the
subject contig are returned instantly, while repetitions of the covariance model in the contig (i.e.
paralog sequences) are also considered and extracted individually. This process is implemented
in Python3 as illustrated with the following block of pseudocode:

1297

1298   1. Open the cmsearch output table for each combination of FASTA file and CM
1299   2. For each contig:
1300   1. Get all the alignments found by cmsearch
1301   2. If number of alignments > 1:
1302   1. Adjust alignment coordinates based on the strand
1303   2. If a full alignment is found:
1304   1. Return the coordinates in the contig to extract*
1305   3. Else:
1306   1. For each alignment:
1307   1. Get the coordinates in the CM
1308   2. Get the coordinates in the contig
1309   2. If order(coordinates in the CM) != order(coordinates in the contig):
1310   1. Search for blocks of sequential alignments and treat them as repetitions
1311   2. For each repetition:
1312   1. Return coordinates to extract*
1313   3. Else:
1314   1. Treat alignments as a single, fragmented hit
1315   2. Return coordinates to extract*
1316   3. Else:
1317   1. Return coordinates to extract*

1318

1319 Asterisks denote the different final outcomes for the alignment information on each contig. Only
1320 sequences representing an alignment to the CM with length >= 800 bp are extracted for further
1321 analyses. The workflow uses snakemake (Mölder et al. 2021) to manage sub processes in an
1322 ordered and scalable manner, and anaconda (https://anaconda.com/) to install and load

dependencies such as Infernal and Blast+. The code is publicly available at GitHub (https://github.com/NeLLi-team/ssuextract) and has been successfully tested in SUSE Linux Enterprise Server 15 SP5 and Ubuntu 24.04 LTS in an WSL instance for Windows 11.

References

Anaconda Software Distribution. Computer software. Vers. 2-2.4.0. Anaconda, Nov. 2016. Web. https://anaconda.com.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics*, *10*, 421.

Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., Boutte, C., Burgaud, G., de Vargas, C., Decelle, J., Del Campo, J., Dolan, J. R., Dunthorn, M., Edvardsen, B., Holzmann, M., Kooistra, W. H. C. F., Lara, E., Le Bescot, N., Logares, R., … Christen, R. (2013). The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote small sub-unit rRNA sequences with curated taxonomy. *Nucleic Acids Research*, *41*(Database issue), D597–D604.

Mölder, F., Jablonski, K. P., Letcher, B., Hall, M. B., Tomkins-Tinch, C. H., Sochat, V., ... & Köster, J. (2021). Sustainable data analysis with Snakemake. *F1000Research*, 10, 33.

Nawrocki, E. P., & Eddy, S. R. (2013). Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics* , *29*(22), 2933–2935.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O. (2013). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research*, *41*(Database issue), D590–D596.