Organ-level gene regulatory network models enable the identification of
central transcription factors in *Solanum lycopersicum*

<u>Running title</u>: Organ-level gene regulatory networks in *Solanum lycopersicum.*

José D. Fernández[1,2,3], David Navarro-Payá[4], Antonio Santiago[4], Jonathan
Canan[1,2,5], Sebastián Contreras-Riquelme[2,7], Ariel Cerda[2,3,7], Tomás C.
Moyano[2,3,7], Lorena Melet[1,2,3], Nathan R. Johnson[1,2], Javier Canales[2,6], José M.
Álvarez[2,3,7,*], José Tomás Matus[4,*], Elena A. Vidal[1,2,3,*]

[1]Centro de Genómica y Bioinformática, Universidad Mayor, 8580745 Santiago,
Chile.
[2]Agencia Nacional de Investigación y Desarrollo-Millennium Science Initiative
Program-Millennium Institute for Integrative Biology (iBio), 8331150 Santiago,
Chile.
[3]Agencia Nacional de Investigación y Desarrollo-Millennium Nucleus in Data
Science for Plant Resilience (Phytolearning), 8370186 Santiago, Chile.
[4]Institute for Integrative Systems Biology (I^{2}SysBio), Universitat de València-
CSIC 46908 Paterna, Valencia, Spain.
[5]Instituto de Ciencias de la Ingeniería, Universidad de O'Higgins, 2820000
Rancagua, Chile.
[6]Instituto de Bioquímica y Microbiología, Facultad de Ciencias, Universidad
Austral de Chile, 5110566 Valdivia, Chile.
[7]Centro de Biotecnología Vegetal, Facultad de Ciencias de la Vida, Universidad
Andrés Bello, 8370186 Santiago, Chile.

*Corresponding authors:
José Miguel Alvarez (jose.alvarez.h@unab.cl)
José Tomás Matus (tomas.matus@uv.es)
Elena A. Vidal (elena.vidal@umayor.cl)

Date of submission: 26-03-2025
Number of tables and figures:
1 Table, 6 Figures, 18 Supplementary Tables, 9 Supplementary Figures.
Word count: 7,071

Highlight

We developed organ-level gene regulatory networks for tomato using 10,000+ RNA-seq libraries, validated predictions and identified new regulators of fruit ripening and ABA response. These networks are available at https://plantaeviz.tomsbiolab.com/tomviz.

Abstract

Tomato (*Solanum lycopersicum*) is a globally important crop, yet the gene regulatory networks (GRNs) controlling gene expression remain poorly understood. In this study, we constructed GRNs for roots, leaves, flowers, fruits, and seeds by inferring transcription factor (TF)–target interactions from over 10,000 RNA-seq libraries using the GENIE3 algorithm. We refined these networks with gene co-expression data and computational predictions of TF binding sequences in open chromatin sites. Our networks confirmed key TFs, including TOMATO AGAMOUS LIKE 1 and RIPENING INHIBITOR in fruit ripening, as well as ABF3 and ABF5 in abscisic acid response in leaves. Additionally, we identified novel candidate regulators, including AUXIN RESPONSE FACTOR 2A and ETHYLENE RESPONSE FACTOR.E2 in fruit ripening and G-BOX BINDING FACTOR 3 (*Sl*GBF3) in ABA-related and drought pathways. To further validate the GRNs, we used DNA Affinity Purification Sequencing (DAP-seq) for *Sl*GBF3*,* confirming the accuracy of our GRNs. This study provides a valuable resource for dissecting transcriptional regulation in tomato, with potential applications in crop improvement. The GRNs are publicly accessible through a user-friendly web platform at https://plantaeviz.tomsbiolab.com/tomviz.

Keywords

Organ-level Gene Regulatory Network, GENIE3, Tomato, DAP-seq, ABA, ripening.

Abbreviations

ABA: Abscisic acid

ABF: ABA response element binding factor

AUPR: Area under the precision-recall

AUROC: Area under the receiver operating characteristic

DAP-seq: DNA affinity purification sequencing

DEG: Differentially Expressed Gene

GENIE3: GEne Network Inference with Ensemble of Trees

GO: Gene Ontology

GCN: Gene coexpression network

GRN: Gene regulatory network

GSEA: Gene Set Enrichment Analysis

IVI: Integrated Value of Influence

OCS: Open Chromatin Sites

PP2C: Protein Phosphatase class 2 C.

PWM: Position Weight Matrix

RIN: RIPENING INHIBITOR

SnRK2: SNF1-related protein kinase 2.

TAGL1: TOMATO AGAMOUS-LIKE 1

TF: Transcription factor

TPM: Transcripts per Million

Introduction

Tomato *(Solanum lycopersicum L.)* is one of the most widely cultivated and consumed crops globally. It also serves as a key model organism for research on fleshy fruit development, ripening, and plant defense responses (Kimura and Sinha, 2008; Gascuel et al., 2017). Despite its significance, the regulatory networks controlling tomato responses to internal and external signals remain largely unexplored.

Various approaches exist to help in the identification of TF-target interactions and generate gene regulatory networks. Chromatin immunoprecipitation followed by sequencing (ChIP-seq) and DNA affinity purification sequencing (DAP-seq) can directly assess events of TF binding in genomic regions; however, these methods are technically challenging, particularly in non-model organisms (Park, 2009). In *Solanum lycopersicum*, only a limited number of TFs have been studied using ChIP-seq (Fujisawa *et al.*, 2011; Ricardi *et al.*, 2014; Du *et al.*, 2017; Lü *et al.*, 2018; Gao *et al.*, 2019; Lira *et al.*, 2020; Liu *et al.*, 2020; Ding *et al.*, 2022; Tu *et al.*, 2022; Yang *et al.*, 2022; Jiang *et al.*, 2023). Similarly, just a limited number of TFs have been studied using DAPseq (López-Vidriero *et al.*, 2021; Chong *et al.*, 2022; Huang *et al.*, 2023; Zhu *et al.*, 2023). While these approaches have identified TF binding sites and regulatory targets of TFs, they are typically limited to individual TFs. Alternatively, studies addressing accessible chromatin sites to derive binding of multiple TFs at a genome-wide scale in tomato are scarce. Most ATAC-seq analyses are focused on fruit responses to abiotic stress (Maher *et al.*, 2018; Reynoso *et al.*, 2019; Hendelman *et al.*, 2021; Kajala *et al.*, 2021), while DNase-seq studies also remain limited (Qiu *et al.*, 2016; Lü *et al.*, 2018).

The existing biological network models in tomato have focused primarily on protein-protein interactions or gene co-expression networks (GCNs), often derived from small datasets or limited to condition-specific analyses (Ozaki *et al.*, 2010; Fukushima *et al.*, 2012; Gao *et al.*, 2013; Koenig *et al.*, 2013; Pan *et al.*, 2013; Ichihashi *et al.*, 2014; Arhondakis *et al.*, 2016; Kim *et al.*, 2017; Zouine *et al.*, 2017; Xie *et al.*, 2019; Bae *et al.*, 2021; Bizouerne *et al.*, 2021; Kusano *et al.*, 2022; Wang *et al.*, 2023*a*; Li *et al.*, 2024). Some studies have integrated larger scale transcriptomic datasets from microarrays and RNA-Seq to generate GCNs. Fukushima et al., (2012) analyzed 307 microarrays from 17 experiments,

encompassing diverse conditions in a single GCN that was separated into organ-specific submatrices. Similarly, Kim et al., (2017) compiled 1,473 expression samples from 12 studies of microarrays and mRNA-Seq data to generate organ-independent coexpression networks. Zouine et al., (2017) integrated 29 RNA-seq studies to generate a global GCN. However, these networks are either organ-independent or focus only on fruits. Additionally, they rely on outdated tomato genome annotations (ITAG2.4 or ITAG3.0) (Fukushima *et al.*, 2012; Kim *et al.*, 2017; Zouine *et al.*, 2017). Moreover, co-expression network approaches infer associations based on expression similarity rather than directed regulatory interactions (Babu *et al.*, 2004; Chai *et al.*, 2014; Swift and Coruzzi, 2017). Gene regulatory networks (GRNs) offer a more comprehensive framework to study TF-target interactions at a genome-wide scale. Unlike GCNs, GRNs infer directed regulatory interactions, identifying key regulatory hubs that orchestrate biological processes (Doidy *et al.*, 2016; Vidal *et al.*, 2020; Escorcia-Rodríguez *et al.*, 2023). Therefore, developing genome-wide GRN models that integrate updated genome assemblies, diverse transcriptomic datasets, and organ-level regulation is crucial to uncovering transcriptional regulatory mechanisms in tomato.

Machine learning based approaches, such as GENIE3 (GEne Network Inference with Ensemble of Trees) have been widely used to infer GRNs from large-scale transcriptomic datasets. GENIE3 applies an ensemble of regression trees to predict TF-target interactions and has demonstrated a high performance in the DREAM4 and DREAM5 Network Inference challenges (Huynh-Thu *et al.*, 2010; Huynh-Thu and Geurts, 2019). This approach has been successfully applied to generate GRNs in various plant species, including *Arabidopsis thaliana*, wheat, and maize (Huang *et al.*, 2018; Harrington *et al.*, 2020; Tu *et al.*, 2020; De Clercq *et al.*, 2021; Chen *et al.*, 2023; Ranjan *et al.*, 2024).In this study, we integrate large-scale tomato datasets to generate, validate, and refine organ-level GRNs. Using an extensive set of transcriptomic data and the GENIE3 algorithm, we generated five reference GRNs representing different tomato organs (roots, leaves, flowers, fruits and seeds). These networks were further enhanced with TF-target interactions derived from accessible chromatin data, predicted TF binding events, and genome-wide GCNs. The resulting organ-level GRNs provide a robust foundation for exploring diverse biological contexts, from developmental processes to stress responses, offering a valuable resource for

164 addressing unresolved questions in tomato biology. The organ-level GRNs are
165 available at https://plantaeviz.tomsbiolab.com/tomviz.
166

Materials and Methods
167

Tomato gene annotation update
169

Gene models from tomato annotation iTAG 4.2 beta (provided by Dr. Surya Saha,
SolGenomics) were merged with gene models from iTAG4.0. New gene models
reported in iTAG4.2 beta were incorporated into the iTAG4.0 annotation when
their genome coordinates did not overlap existing entries, resulting in the final
annotation file utilized in this work (hereafter iTAG4-merge). To expand the
functional annotations for tomato, ITAG 4.0c protein sequences were analyzed
using eggNOG-mapper (Cantalapiedra *et al.*, 2021) to predict Gene Ontology
(GO) terms, functional categories, and orthologs relationships. Additionally,
functional annotations for all genes in ITAG 4.0c were generated using
InterProScan v5.57-90 (Jones *et al.*, 2014) under default parameters and
complemented with GO information for *Solanum lycopersicum* from PLAZA 5.0
(Van Bel *et al.*, 2022) database. The resulting GO annotations for molecular
functions and biological processes were consolidated and used to create an
updated GO set.
To update the list of TFs in ITAG 4.0c, we integrated evidence from multiple
sources: 1. Gene descriptions for iTAG4.0 and iTAG4.1 available at SolGenomics
(description contains the term "transcription factor") (Fernandez-Pozo *et al.*,
2015; Hosmani *et al.*, 2019); 2. TF catalogs for *S. lycopersicum* obtained from
PlantTFDB (Jin *et al.*, 2017) and ITAK (Zheng *et al.*, 2016); 3. ITAG 4.1c GO
annotations (description contains the terms "transcription factor", "DNA-binding");
The output of an OrthoFinder v3.0 (Emms and Kelly, 2019) analysis of
Arabidopsis TFs from PlantTFDB; and 4. A search of proteins that we annotated
with an IPR code associated to eukaryote TFs (list of IPR codes obtained from
InterPro and manually curated to exclude IPR codes not representing TFs)
(Jones *et al.*, 2014). Genes supported by at least three independent lines of
evidence were selected and a manual curation step was conducted to exclude
proteins associated with enzymatic activities, non-transcriptional molecular

processes (e.g., DNA replication, repair, splicing, translation), transcriptional
regulators other than TFs (e.g., basal transcription factors, RNA polyadenylation
factors), and chromatin remodeling complex subunits. We associated Position
Weight Matrices (PWMs) to the TFs by combining information obtained from
CisBP v.2 (Weirauch *et al.*, 2014) with PWM inference from protein sequences
using the JASPAR profile inference tool "infer_profile.py" (Castro-Mondragon *et*
al., 2022).

Processing of RNA-seq data

To obtain tomato RNA-seq datasets, we queried the NCBI SRA database using
("*Solanum lycopersicum*"[Organism] AND ILLUMINA[Platform]) NOT (RIP-
Seq[Strategy]   OR   OTHER[Strategy]   OR   ChIP-Seq[Source]   OR
METATRANSCRIPTOMIC[Source]   OR   Bisulfite-Seq[Strategy]   OR
GENOMIC[Source]   OR   METAGENOMIC[Source]   OR   DNase-
Hypersensitivity[Strategy] OR WGS[Strategy] OR ncRNA-Seq[Strategy] OR
WCS[Strategy] OR degradome OR miRNA-Seq[Strategy] OR small RNA[Title]
OR sRNA[Title]) (Leinonen *et al.*, 2011). The metadata was classified by organ
of origin following the protocol in Santiago *et al.* (2024). The libraries were
downloaded using SRAtools (Kans, 2010). Adapters were trimmed, and low-
quality reads (reads with average quality inferior to q<30 and shorter than 20
bases) were filtered out using fastp v.0.20.0 (Chen *et al.*, 2018). Reads were
aligned to the SL4.0 *S. lycopersicum* genome assembly (Hosmani *et al.*, 2019)
using STAR v.2.7.3 (Dobin *et al.*, 2013). After mapping, a total of 10,618 mRNA-
seq libraries were retained. Gene counts were obtained with FeatureCounts
v.2.0.0 (Liao *et al.*, 2014) using the ITAG4-merge annotation. Total counts were
normalized to transcripts per million (TPM). Finally, genes with ≥ 5 TPM in at least
10% of the total libraries for each organ were classified as expressed, as
described in Huang et al., (2018).

Processing of ChIP-seq data

The   following   query   (("*Solanum   lycopersicum*"[Organism]   AND
ILLUMINA[Platform]) AND ChIP-Seq[Source]) was used to obtain tomato TF-
binding (ChiP-seq) datasets from the NCBI SRA website (Leinonen et al., 2011).

ChIP-seq libraries were processed using the methods described by ENCODE
(Hitz *et al.*, 2023). Briefly, the libraries were downloaded from the NCBI SRA
using SRAtools (Kans, 2010). Adapters were trimmed, and low-quality reads
(reads with average quality q<30 and shorter than 20 bases) were filtered using
Cutadapt v.4.9 (Martin, 2011). Each file was mapped with Bowtie2 v.2.54
(Langmead and Salzberg, 2012) to the SL4.0 assembly (Hosmani *et al.*, 2019).
Alignment files were sorted and filtered with Samtools v.1.21 (Li *et al.*, 2009) and
peaks were identified with MACS2 v.2.2.9.1 (Zhang *et al.*, 2008). Only libraries
with ≥80% mapping efficiency and over 1,000 peaks assigned to annotated genes
were retained as high-quality datasets for downstream analysis.
240

Processing of ATAC-seq and DNAse-seq data

The following query (("*Solanum lycopersicum*"[Organism] AND
ILLUMINA[Platform]) AND ATAC-Seq[Source] AND DNAse-Seq[Source]) was
used to obtain tomato open chromatin sites (OCSs) datasets from the NCBI SRA
website (Leinonen *et al.*, 2011). A total of 183 open chromatin experiments
(DNase-seq, ATAC-seq libraries) were downloaded using SRAtools (Kans,
2010). Reads were trimmed of adapters, and low-quality reads (reads with
average quality q<30 and shorter than 20 bases) were filtered using Cutadapt
v.4.9 (Martin, 2011). The ATAC-seq libraries were processed following the
protocol described in Reynoso et al., (2019), while the DNase-seq libraries were
processed following the protocol described in Moyano et al., (2021). Briefly, reads
were mapped to the SL4.0 genome assembly (Hosmani *et al.*, 2019) using
Bowtie2 v.2.54 (Langmead and Salzberg, 2012). The ATAC-seq alignments were
sorted and filtered with Samtools v.1.21 (Li *et al.*, 2009) and peaks were identified
with HOMER v.4.11 (Heinz *et al.*, 2010). The DNase-seq regions were mapped
into DNase hypersensitive sites using HOTSPOT (Meuleman *et al.*, 2020). OCS
peak files were then merged by experiment and converted into FASTA
sequences with BedTools v.2.31.1 (Quinlan and Hall, 2010).

Determination of TF binding sites using FIMO

TF-target datasets were generated by mapping TF DNA binding preferences,
represented as Position Weight Matrices (PWMs) to genomic regions of the
Solanum lycopersicum SL4.0 genome assembly using the Find Individual Motif

Occurrences (FIMO) search tool, with default parameters (p-value < $1\times10e^{-4}$)

(Grant *et al.*, 2011). As query sequences, we used all promoter sequences

represented as two kilobases upstream of the transcription start site (TSS) of

each gene, or organ-level OCS sequences. For OCS sequences, the results were

assigned to genes using the BedTools command ClosestGene (Quinlan and Hall,

2010).

GENIE3 inference of regulatory interactions

Raw gene counts for each organ, as well as the updated list of TFs, were given

as input to the GENIE3 algorithm (Huynh-Thu *et al.*, 2010). The GENIE3 tool was

run with standard parameters, using 2,000 decision trees and a seed of 122 for

reproducibility. The output scores were used to create subnetworks based on the

top 1%, 2%, 5%, 8%, and 10% scores of TF-target pairs. These thresholds were

used in previously published GENIE3 networks (Cuesta-Astroz et al., 2021;

J.Huang et al., 2018; Olivares-Yañez et al., 2021).

Co-expression network generation

To generate co-expression networks for tomato organs, we used the pipeline

described in Orduña, et al. (2023). Briefly, the gene count tables from the

transcriptome of each organ were used to calculate the Pearson correlation

values for each pair of genes. The results were ordered by gene rank in

descending order and computed in a highest reciprocal ranking (HRR) matrix per

organ following the formula: HRR (A,B) = max (rank (A,B), rank (B,A)). Finally, to

avoid noise and low confidence pairings, the top 1% of the highest frequency

HRR per gene were chosen to make the tomato organ-specific GCNs.

Evaluation of network performance

To evaluate the performance of the predicted GRNs in capturing experimentally

validated regulatory interactions, we followed the protocol in Contreras-López et

al., (2022) to compute the area under the receiver operating characteristic

(AUROC) and precision-recall (AUPR) curves. Briefly, these analyses were

performed using organ-level GRNs and tested against validated regulatory

interactions derived from ChIP-seq datasets. Gene interactions were filtered to

retain only regulatory genes present in both the GENIE3-inferred and ChIP-seq

networks, with edges assigned as binary labels indicating the validation status.

True and false positive rates were calculated using the *precrec* v.0.14.4 package

in R. To assess statistical significance, the AUROC and AUPR values were

compared against 1,000 randomized networks generated by shuffling edge

weights. Percentile-based confidence intervals (2.5–97.5%) were used to

benchmark the performance of the GENIE3 network, and significance was

determined via a permutation test.

Visualization of GRNs and network analysis

Network visualizations were generated using Cytoscape v.3.10.1 (Shannon *et al.*,

2003) and network topology analyses were conducted using the Cytoscape

NetworkAnalyzer tool. The R package Influential v. 2.2.9 (Salavaty *et al.*, 2020)

identified the TF hubs with the highest integrated value of influence (IVI).

Gene Set Enrichment Analysis (GSEA)

Gene Set Enrichment Analysis (GSEA) was conducted to identify

overrepresented biological process GO terms using a hypergeometric test with

Benjamini and Hochberg false discovery rate (FDR) correction (threshold < 0.05).

The analysis was performed using the BinGO v.3.0.5 tool (Maere *et al.*, 2005)

within Cytoscape, with input from the updated tomato 4.1c GO terms catalog. The

REVIGO v. 1.8.1 (Supek *et al.*, 2011) web application was used to improve and

narrow down the grouping of GO terms, and GO terms within levels 5–7 were

selected to focus on more specific terms.

Generation of DAP-seq libraries

Genomic DNA (gDNA) was extracted from fully expanded mature leaves of 3-

month-old *Solanum lycopersicum* cv. Moneymaker plants using the Wizard

Genomic DNA Purification Kit (Promega), according to the manufacturer's

instructions. gDNA was fragmented to an average size of ~200 bp using a M220

sonicator (Covaris, Woburn, MA, USA). The fragmented gDNA underwent end repair, A-tailing and ligation of Illumina adapters.

The full-length coding sequence of *SlGBF3* (*Solyc03g120460*) was amplified from cDNA obtained from *S. lycopersicum* cv. Moneymaker leaves using primers SlGBF3_Fw (5'-CACCATGGGAAATAGTGAGGATGGGAAATCATGTAAGC-3') and SlGBF3_Rv (5'-TCACCCAGCTGCTACTGCATCA-3'). PCR products were cloned into the pFN19K_HaloTag-T7-SP6 Flexi expression vector (Promega, Madison, WI, USA), with the Halo-Tag at the N-terminal. Expression of the Halo-tagged fusion protein was performed using the TNT SP6 Coupled Wheat Germ Extract System (Promega). HaloTag-ligand conjugated magnetic beads (Promega) were used to pull-down the Halo-tagged TF. Pulled-down TFs were exposed to adapter-ligated gDNA libraries. Bound DNA was eluted and sequencing libraries were generated by PCR amplification with Illumina TruSeq Universal and Index primers. An empty expression vector was used as a negative control to account for non-specific DNA binding (input library). Libraries were sequenced on an Illumina NovaSeq 6000 platform (150 pb PE, approximately 22 million reads per library). Two replicates were used for all experiments.

Processing of DAP-seq data

Raw sequence data were processed using the pipeline described by *Hutin et al., (2023)*. Briefly, adapter sequences were trimmed, and low-quality reads (reads with average quality q<30 and shorter than 20 bases) were filtered using Cutadapt v.4.9 (Martin, 2011). Filtered reads were mapped to the *Solanum lycopersicum* genome assembly SL4.0 (Hosmani *et al.*, 2019) using Bowtie2 v2.5.4 (Langmead and Salzberg, 2012). Alignment files were sorted, filtered, and deduplicated using Samtools v1.21 (Li *et al.*, 2009). Peaks were identified using MACS2 v2.2.9.1 (Zhang *et al.*, 2008) with the parameters -f BAMPE -q 0.00001 --call-summits. Final peak-associated regions were annotated using BedTools closestGene function (Quinlan and Hall, 2010). Binding motifs were identified using the MEME-ChIP tool v5.1.1 (Machanick and Bailey, 2011).

Results

Refining tomato gene annotation and transcription factor catalog for GRN construction

As a first step to generate organ-level GRNs for tomato, we obtained a revised gene annotation for genome SL4.0 from SolGenomics (iTAG4.2 beta). To include all the genes, present in iTAG4.0 and include new genes, we merged the iTAG4.0 and iTAG4.2 beta versions, into the iTAG4-merge version. This version contains a total of 37,467 genes (Supplementary Datasets 1,2,3). Considering that only 15.53% of tomato genes had a functional annotation in ITAG4.0, we generated an updated annotation for protein coding genes by integrating different sources of evidence. These included obtaining GO terms assigned by EggNOG-mapper (Cantalapiedra *et al.*, 2021) and InterProScan (Jones *et al.*, 2014) and gathering gene functional annotations compiled in PLAZA 5.0 for tomato (Van Bel *et al.*, 2022). The GO annotation in ITAG 4.0c covers 70.47% (26,403 out of 37,467) of protein-coding genes, each associated with at least one GO term. This dataset includes 9,379 unique GO terms, providing biological function annotations for 21,630 genes and molecular function annotations for 23,809 genes, for a total of 982,370 annotations (Supplementary Table S1 and S2).

To generate an updated list of TFs, we retrieved TF catalogs from various repositories, we conducted orthologous TF analyses using the set of *Arabidopsis thaliana* TFs and we obtained information from the GO annotations. Only TFs supported by at least three lines of evidence were selected, resulting in a set of 1,840 TFs for further analyses (Supplementary Table S3). DNA-binding preferences were determined by retrieving position weight matrices (PWMs) from CisBP v.2 (Weirauch *et al.*, 2014) or assigning PWMs from Arabidopsis and Maize orthologs in JASPAR (Castro-Mondragon *et al.*, 2022), yielding 846 TFs with assigned PWMs (Supplementary Table S3). This updated annotation and TF list served as the foundation for constructing the GRNs in the subsequent steps.

Tomato genes exhibit widespread expression, yet their expression levels vary across different organs

We started our analysis by compiling a comprehensive dataset of publicly available tomato RNA-Seq libraries (Supplementary Table S4). The

transcriptomic libraries were categorized into five main organs: roots (1,840

libraries from 124 studies), leaves (3,778 libraries from 279 studies), flowers (568

libraries from 55 studies), fruits (4,149 libraries from 147 studies), and seeds (270

libraries from 13 studies) for a total of 10,605 libraries that surpassed quality filters

(Supplementary Fig. S1A). These libraries encompass a diverse range of tomato

genotypes and growth conditions. Additionally, they represent multiple

experimental contexts, including abiotic stress (3,379 runs), biotic interactions

(2,085 runs), developmental studies (2,872 runs) and genetic modification

studies (2,035 runs) (Supplementary Fig. S1B). This comprehensive dataset

provides a robust foundation for constructing reference GRNs to address multiple

research questions.

Reads were mapped to gene models using the ITAG4-merge annotation, and

genes with expression levels above 5 TPM in more than 10% of the total libraries

for a given organ were considered as expressed in that organ. We identified

26,910 genes (71.82%) as expressed in at least one organ, from which 19,361

(71.95%) genes are expressed in all organs, while a minor fraction (15.9/% of

expressed genes) exhibit organ-level expression (Fig. 1A, Supplementary Table

S5). Similar expression patterns have been reported in maize, flaxseed and

wheat (Huang et al., 2018; Qi et al., 2023; Ramírez-González et al., 2018).

Examples of specific genes include *Solyc06g051770* and *Solyc10g047720* in

seeds, whose Arabidopsis homologs, *Oleosin 1* and *2 (OLEO1-2)*, are involved

in seed oil body formation (Siloto *et al.*, 2006). In roots, we identified *SULTR1;1*

(*Solyc10g047170*), which encodes a sulfate transporter associated with external

sulfate uptake (Takahashi *et al.*, 2000). In flowers, *Tapetum Determinant 1-like*

(TPD1-like) genes such as *Solyc11g005500*, *Solyc12g009850*, *Solyc05g010190*

and *Solyc04g071640* were specifically expressed, consistent with their role in

tapetal cell development and gametogenesis (Ezura *et al.*, 2017). In leaves, we

found *Longifolia* 1 (*SILNG1, Solyc02g089030)*, whose Arabidopsis homolog

LNG1 (AT5G15580) regulates leaf morphology by promoting longitudinal cell

expansion (Lee *et al.*, 2006) (Fig. 1A).

Concerning TFs, 1612 (87.7%) were expressed in at least one organ. Among

them, 1014 (62.9%) were expressed across all organs, while a smaller subset

(16.46% of expressed TFs) exhibited organ-level expression (Fig. 1B,

Supplementary Table S5). This latter group includes *SlBRC1a* and *SlBRC1b*

(*Solyc03g119770* and *Solyc06g069240*), paralogs of Arabidopsis *BRANCHED1*,
involved in leaf and axillary bud development (Martín-Trillo *et al.*, 2011), *SlFER*
(*Solyc06g051550*) a key regulator of root iron uptake (Aviña-Padilla *et al.*, 2023),
SlWUSCHEL (WUS, Solyc02g083950), which controls floral meristem identity
and development (Hawar *et al.*, 2022), *SlSHINE2* (*SHN2*, *Solyc12g009490*),
encoding a TF that controls epidermal growth in developing fruits (Bres *et al.*,
2022), and two Arabidopsis *ABI4* paralogs
(*Solyc03g095977* and *Solyc03g095973*) that are exclusively expressed in seeds,
consistent with their role in seed vigor (Bizouerne *et al.*, 2021). Importantly, while
most TFs and other genes are expressed across all organs, their expression
levels varied substantially depending on the organ analyzed (Fig. 1C, D). These
quantitative differences suggest organ-level regulatory mechanisms, where
distinct expression patterns contribute to the specialized functions and
characteristics of each organ.

To assess how gene expression in tomato organs relates to the occurrence of
organ-specific and widely expressed biological processes, we performed a Gene
Set Enrichment Analysis (GSEA) for each organ. The analysis revealed that most
enriched biological processes (151 GO terms) were common across all organs
(adjusted p-value < 0.05), including gene expression regulation, circadian rhythm,
vacuole and vesicle organization, immune responses, mRNA methylation and
response to abscisic acid (Figure 1E, Supplementary Table S6). In contrast, only
22 enriched GO terms were identified as unique for each organ. These include
processes related to fruit ripening in fruits, root meristem identity and response
to reactive oxygen species in roots, phototropism and photoperiodism in leaves,
shoot apical meristem identity, brassinosteroid signaling, and pollen tube
guidance in flowers as well as lipid storage and seed development in seeds
(Figure 1E, Supplementary Fig. S2, Supplementary Table S6). These findings
demonstrate that while fundamental biological processes are conserved across
all tomato organs, a subset of organ-level processes support their unique
functions, emphasizing a complex regulatory mechanism that defines organ
identity in tomato.

Organ-level GENIE3 networks recapitulate experimentally obtained TF-target interactions

To generate organ-level GRNs, we employed the GENIE3 algorithm using the transcriptomic count tables separated by organ and the updated TF list as input. GENIE3 generated a ranked list of putative TF-target interactions, from which we selected the top 1%, 2%, 5%, 8% and 10% of the highest-scoring interactions to evaluate network accuracy. The inferred organ-level networks were benchmarked against ChIP-seq datasets for tomato (Supplementary Table S7). These included data for GLK1 and 2 (Solyc07g053630, Solyc10g008160) (Tu *et al.*, 2022), MYC2 (Solyc08g076930) (Du *et al.*, 2017), JMJ4 (Solyc08g076390) (Ding *et al.*, 2022), WOX13 (Solyc02g082670) (Jiang *et al.*, 2023), EIL4 (Solyc06g073730), TAGL1 (Solyc07g055920) and RIN (Solyc05g012020) (Fujisawa *et al.*, 2011; Gao *et al.*, 2019) (Supplementary Table S7). Each GENIE3 network was benchmarked against a ChIP-seq gold-standard network through Fisher's exact test enrichment analysis. The analysis included the TFs GLK1 and 2, MYC2, EIL4, JMJ4, and WOX13 to evaluate all organs, while RIN and TAGL1 were assessed exclusively in reproductive organs (excluding roots and leaves networks), as their expression is restricted to these tissues. The top 2% networks showed the highest enrichment (indicated by higher $\log_2$-transformed Fisher odds ratios) and statistical significance (lower $\log_{10}$-transformed adjusted p-values) in the overlap between TF-target pairs obtained by GENIE3 and TF-target pairs obtained by ChIP-Seq (Table 1, Supplementary Table S8). Furthermore, comparison with existing tomato gene networks from PlantRegmap (Tian *et al.*, 2020) and TomatoNet (Kim *et al.*, 2017) demonstrated that the GENIE3-derived GRNs exhibited greater enrichment and overlap with the gold-standard dataset, indicating a better performance in predicting TF-target interactions obtained experimentally (Supplementary Table S9).

To further evaluate the performance of the top 2% GRNs, we calculated the AUROC (Area under the receiver Operating Characteristic) and AUPR (Area under the precision-recall) curves for each organ-level network and compared these values against a network consisting of TF-target interactions obtained from the gold-standard dataset. The GRNs derived from tomato roots, leaves, flowers, fruits and seeds showed statistically higher AUROC and AUPR values than

randomly generated TF-target pairs (Supplementary Fig. S3). These results confirm that the GRNs successfully recapitulate experimentally validated TF-target interactions, underscoring their utility in predicting regulatory interactions for TFs lacking experimental validation.

To provide further support to our networks, we integrated additional layers of information from complementary approaches to the GENIE3-inferred edges. Gene co-expression is widely used to infer biologically relevant relationships between genes (Wolfe *et al.*, 2005; Yin *et al.*, 2021). Using the same RNA-seq datasets, we generated aggregated gene co-expression networks (GCNs) following the protocol in Orduña et al., (2023). The TF-target pairs from each GCN were extracted as additional evidence for the GENIE3-predicted interactions. While the GENIE3 algorithm predicts regulatory interactions based on expression patterns, additional evidence is necessary to determine whether these interactions occur via direct TF binding to regulatory sequences. To integrate TF-binding information into the GENIE3 networks, we extracted upstream sequences (2 Kb from the transcription start site) for each annotated gene on ITAG4-merge and performed TF-binding motif prediction using the FIMO tool (Grant *et al.*, 2011). Additionally, we conducted the same analysis on sequences within open chromatin sites (OCSs) identified in tomato fruit, flower, leaf, and root organs, using data from DNase-seq and ATAC-seq experiments (Supplementary Table S10). The results from all evidence layers were compiled, ensuring that our analysis remained constrained to TF-target pairs identified by GENIE3. This approach maintained the predefined network structure, with additional regulatory evidence mapped onto it rather than introducing new interactions. We found that between 51% and 61% of GENIE3-predicted interactions were supported by at least one additional evidence, with most edges validated by one or two different approaches (Fig. 3A). Furthermore, a substantial portion of GENIE3 edges were validated by the presence of cis-binding motifs detected by FIMO within promoter sequences or OCSs, suggesting a direct regulation of TFs over their inferred targets.

As previously mentioned, most genes and TFs were expressed across all tomato organs, albeit with varying expression levels. To assess how these expression patterns influence regulatory interactions, we examined the distribution of TF-target gene pairs across the five organ-level GRNs. Over 75% of these pairs were

unique to a single organ (Fig. 3B), indicating that while TFs and targets are broadly expressed, the existence of regulatory pairs is lar |gely   organ specific. This has previously been described for other plant GRNs (Huang *et al.*, 2018; Ranjan *et al.*, 2024).

To further evaluate how specific TF-target pairs distribute across organs and how target conservation correlates with TF connectivity, we calculated the percentage of conserved targets for each TF across all organs. As expected, most TFs showed low target conservation due to the organ-level nature of TF-target interactions. Interestingly, we observed a significant correlation between TF connectivity and target conservation ($R^2 = 0.36$, $p < 2.2e^{-16}$), indicating that highly connected TFs tend to regulate conserved target genes across organs (Fig. 3C; Supplementary Table S11). Gene Set Enrichment Analysis (GSEA) of the top five most connected and conserved TFs revealed strong enrichment for biological process GO terms related to fundamental cellular processes, including nucleic acid metabolism, vesicle transport, and RNA metabolism (FDR-adjusted p-value < 0.05) (Supplementary Fig. S4). These results suggest that TFs with high connectivity and conserved targets function as global regulators of essential cellular pathways. In contrast, TFs with lower connectivity and limited target conservation appear to mediate more specific functions. For instance, despite their ubiquitous expression, TFs such as *LOB27* (*Solyc06g062630*) and *bHLH17* (*Solyc02g093280*) show enriched functions in flowers, *WRKY47* (*Solyc01g058540*) in roots, *B3* (*Solyc05g004000*) in flowers and seeds, and *MYB104* (*Solyc01g090530*) exhibits distinct functions across multiple organs (Supplementary Fig. S5). In sum, organ-level GENIE3 networks present topological features expected for GRNs and can recapitulate TF-target interactions experimentally obtained by complementary approaches.

The fruit GRN captures known regulatory interactions and identifies novel central controllers of ripening

To evaluate the ability of the organ-level GRNs to capture biologically relevant regulatory interactions, we focused on ripening, one the most extensively studied processes in tomato, linked to hormonal signaling pathways such as ethylene, cell wall remodeling, pigments biosynthesis and other processes (Karlova *et al.*,

2014; Li *et al.*, 2019; Zhu *et al.*, 2022). Tomato fruit ripening is governed by a
complex regulatory cascade, involving interactions between well-characterized
TFs including APETALA2a (AP2a), NON-RIPENING (NOR), FRUITFULL
(FUL1/TDR4 and FUL2/MBP7), TOMATO AGAMOUS-LIKE 1 (TAGL1),
RIPENING INHIBITOR (RIN) and COLORLESS NON-RIPENING (CNR) (Li *et*
al., 2021*a*; Zhu *et al.*, 2022). Among these, TAGL1 and RIN are recognized as
central regulators of ripening (Gao *et al.*, 2019; Li *et al.*, 2021*b*).
To determine whether the fruit GRN reproduced regulatory interactions of known
TFs, we compared the targets of TAGL1 and RIN from the fruit GRN with gene
lists compiled from previous omics studies. These included differentially
expressed genes (DEGs) identified in TAGL1 and RIN knockout and RNAi plants
(Gao et al., 2019; Ito et al., 2020; Li et al., 2018), as well as direct binding targets
identified via RIN ChIP-chip and ChIP-seq (Fujisawa et al., 2013; Gao et al.,
2019; Zhong et al., 2013), and TAGL1 ChIP-seq (Gao *et al.*, 2019) experiments.
For RIN, we observed a statistically significant overlap of the targets determined
in the fruit GRN with targets obtained in all the experiments, including ChIP-
binding targets and regulatory targets identified in RIN-deficient plants (Fig. 4A).
Similarly, for TAGL1, the fruit GRN was able to identify targets validated by ChIP-
seq and/or knock-out experiments (Fig. 4B). This finding highlights the potential
of the GENIE3 GRN to capture experimentally validated regulatory interactions,
with many of these corresponding to direct binding of a TF to a target promoter
(45% of RIN GRN targets (411/857) and 89% of TAGL1 GRN targets (740/827)
are validated by ChIP binding evidence) (Figure 4A-B).
To further investigate the regulatory roles of RIN and TAGL1 in fruit ripening, we
generated RIN and TAGL1 subnetworks focusing on genes involved in fruit
ripening described in Li *et al.*, (2019) and Zhu *et al.*, (2022). We found that the
GENIE3 network predicts interactions for both RIN and TAGL1 with an important
number of ripening genes (Supplementary Table S12). Furthermore, most of the
GENIE3 edges connecting RIN and TAGL1 to these target genes (90% for RIN
and 100% for TAGL1) are supported by ChIP evidence (Fig. 4C, D). Additionally,
the subnetworks highlight the regulatory influence of RIN and TAGL1 across
diverse biological processes including central or secondary metabolism, cell wall
modification, and their interactions with other known TFs in fruit ripening including
CNR, NOR, and AP2a (Fig. 4C, D).

To identify novel ripening regulators, we used the fruit GENIE3 network to generate a subnetwork including the list of ripening-associated genes (Li *et al.*, 2019; Zhu *et al.*, 2022). We applied the Integrated Value of Influence (IVI), a metric that combines centrality measures such as degree centrality, cluster rank, neighborhood connectivity, betweenness centrality, and collective influence into a single value to quantify network hub influence (Salavaty *et al.*, 2020). We found that among TFs with higher IVI are CNR, NOR, FUL1, AP2a, RIN, and TAGL1, confirming their roles as central regulators of fruit ripening genes. Notably, two additional TFs, *Sl*ARF2A (Solyc03g118290) and *Sl*ERF.E2 (Solyc06g063070), emerged as TFs with high IVI, revealing a potentially relevant role in controlling ripening-related genes (Supplementary Table S13). *Sl*ARF2A TF has been identified as a regulator of axillary shoot development and is predominantly expressed in the late stages of ripening (Xu *et al.*, 2016). RNAi lines targeting *Sl*ARF2A exhibit ripening defects and ethylene insensitivity, while overexpression lines show accelerated and uneven ripening (Hao *et al.*, 2015; Breitel *et al.*, 2016). On the other hand, *Sl*ERF.E2 expression is downregulated in *cnr*, *nor*, and *rin* mutants, however its role in ripening has not been elucidated yet (M. Liu et al., 2016). Interestingly, we found that *Sl*ARF2A and *Sl*ERF.E2 may act upstream of important TFs such as AP2, NOR, and CNR, in addition to multiple ripening-relevant genes (Fig. 4E).

Tomato organ-level GRNs validate the role of ABF TFs on abscisic acid (ABA) regulatory cascades and identify SlGBF3 as a new regulator of ABA-related genes

ABA is a key plant hormone involved in the regulation of seed dormancy, germination, seedling development, root growth, flowering, and responses to abiotic and biotic stresses (Vishwakarma *et al.*, 2017; Krukowski *et al.*, 2023). Notably, the GO term "response to ABA" was consistently enriched across all organs (Supplementary Fig. S2). This category includes 730 genes, 714 of which are ubiquitously expressed in tomato organs (Supplementary Table S14, hereafter referred as "ABA-related genes"). Among known TFs participating in ABA responses are the members of the AREB/ABF (ABA response element binding/ABA response element binding factor) family of bZIP TFs (Uno *et al.*,

2000; Krukowski *et al.*, 2023). This family has ten members in tomato (Pan *et*
al., 2023), however the relative contribution of each ABF TF at the organ-level
remains unexplored. To address this, we constructed organ-level networks for
the ten ABFs, focusing specifically on their regulation of ABA-related genes.
While most ABF family members are expressed at similar levels across organs,
-except for *Sl*ABF6 (not expressed in flowers) and *Sl*ABF7 (not expressed in fruits
and leaves), their regulatory potential differs depending on the organ. *Sl*ABF1
and *Sl*ABF4 appear to play key roles in regulating ABA-related genes in fruits,
whereas *Sl*ABF2, *Sl*ABF3, *Sl*ABF5, and *Sl*ABF10 regulate more genes in the leaf
network. *Sl*ABF5, *Sl*ABF9, and SlABF10 are involved in ABA response regulation
in roots, while *Sl*ABF2 seems to play a more important role in the flower GRN.
For seeds, *Sl*ABF6 and *Sl*ABF7 exhibit the highest regulatory activity on ABA-
related genes (Figure 5A, Supplementary Table S15). These findings point out
that the ABF TFs have different relative contributions on the regulation of ABA-
related genes across organs.

Plant response to drought is closely linked to ABA signaling (Kang *et al.*,
2002; Krukowski *et al.*, 2023). To validate the regulatory predictions from our
GRNs, we focused on two TFs, *Sl*ABF3 and *Sl*ABF5, which have been identified
as relevant regulators of drought responses in tomato leaves (Kang *et al.*, 2002;
Hsieh *et al.*, 2010). Using the leaf GRN, we extracted the predicted targets of
ABA-related genes from both TFs and compared them to the DEGs found in a
transcriptome analysis of leaves from drought-exposed plants (Wang et al.,
2023). Our analysis revealed a significant enrichment of drought-responsive
genes among the targets of both TFs (Fig. 5B, Supplementary Table S16). We
found that 284 out of 410 (~69%) ABF5 targets and 63 out of 168 (~38%) ABF3
targets overlapped with drought-responsive DEGs. Additionally, 14 out of 22
(~64%) ABF3 targets and 35 out of 45 (~78%) ABA-relevant genes were also
present among the drought DEGs. The GRN also indicated that both TFs regulate
ABA signaling components encoded by *protein phosphatase class 2 C (PP2C)*
genes (Kang *et al.*, 2002; Fujii *et al.*, 2009; Krukowski *et al.*, 2023), including
Solyc03g121880 and *Solyc05g052980* (regulated by both TFs), and
Solyc03g096670 and *Solyc06g076400* (regulated specifically by *Sl*ABF5).
Furthermore, we observed a potential feedback regulatory mechanism between
*Sl*ABF3 and *Sl*ABF5, as previously reported for Arabidopsis (Chang *et al.*, 2019)

(Fig. 5B). These findings confirm the key roles of *Sl*ABF3 and *Sl*ABF5 in the ABA-

mediated drought regulatory cascades (Kang *et al.*, 2002; Hsieh *et al.*, 2010;

Krukowski *et al.*, 2023; Pan *et al.*, 2023).

To identify novel regulators of ABA responses beyond the ABF family of TFs

across all organs, we filtered the five organ-level GRNs to retain TFs with

regulatory connections to ABA-related genes. A network analysis calculating the

Integrated Value of Influence (IVI) (Salavaty *et al.*, 2020) of the network hubs

identified the *Sl*GBF3 TF (Solyc01g095460) as one of the top ten most influential

TFs in the ABA-related networks across all organs, and the most influential in the

leaf GRN (Supplementary Table S17). The *Sl*GBF3 has recently been found to

be co-expressed with drought-responsive genes in tomato leaves (Bortolami *et*

al., 2024). To further explore the role of *Sl*GBF3, we performed a GSEA on its

target genes in each organ-level GRN. We found a significant enrichment of

genes belonging to the "response to abscisic acid stimulus" biological process

shared across all organ networks, indicating a potential conserved role in the

regulation of ABA-related genes (Fig. 5C).

To further validate the potential regulatory interactions of *Sl*GBF3 identified in the

GRNs, we performed DNA Affinity Purification Sequencing (DAP-seq). The DAP-

seq analysis identified approximately 46,705 binding events (peaks) associated

with genomic regions. These peaks were linked to a total of 11,381 genes

annotated in ITAG4-merge in both experimental replicates (Supplementary Table

S18, Supplementary Fig. S6A). Using the identified peak regions, we predicted

the DNA-binding motif of *Sl GBF3* through a MEME-chip analysis (Machanick and

Bailey, 2011). Our analysis revealed that *Sl*GBF3 predominantly binds to a

consensus sequence "AYGTGGCA" (e-value = 5.5 e^{-132} from 28,413 sites) (Fig.

S6B). Comparative analysis with existing TF binding motifs in Cis-BP (Weirauch

et al., 2014) and JASPAR (Castro-Mondragon *et al.*, 2022) databases for

Arabidopsis thaliana GBF3 (*At*GBF3) showed that it has three binding motifs

(MA1351.1-3) with an average consensus sequence of "CGTGGCA" (Fig. S6C).

Notably, the binding motif of *Sl*GBF3 exhibited a significant correlation (Pearson

correlation coefficient between 0.7-0.865) against *At*GBF3 motifs (Fig. S6D),

indicating a strong evolutionary conservation between these TFs and validating

our experimental results.

To evaluate the functional relevance of the identified binding events, we
compared the predicted targets in the tomato organ-specific GRNs with those
identified through DAP-seq. Genes that overlapped between these datasets were
classified as high-confidence targets (HCTs). Our analysis revealed that 42–58%
of the GRN-predicted targets of *SlGBF3* were supported by DAP-seq binding
evidence, with this overlap being significantly enriched (Supplementary Fig. S7).
A GSEA of the complete network of HCTs according to DAP-seq revealed a
significant enrichment in functions related to water deprivation, abiotic stimuli,
and hormone responses, with ABA as a central component (Supplementary Fig.
S8). To further explore *Sl*GBF3 role in ABA-related regulatory networks, we
constructed GRNs specifically for ABA-responsive genes across different organs.
Among these, the leaf GRN exhibited the highest representation of ABA-related
genes. Notably, over 60% of the targets of *Sl*GBF3 in the leaf network were
associated with drought stress responses, including multiple *PP2C* genes
(*Solyc03g121880*, *Solyc03g096670*, *Solyc05g052980*, *Solyc06g076400*) and
two *SNF1-related protein kinase 2 (SnRK2)* genes (*Solyc08g077780*,
Solyc04g012160) (Fig. 5D). Additionally, *Sl*GBF3 appears to function as an
upstream regulator of key TFs controlling ABA-related genes, including *SlABF2*,
SlABF3, and *SlABF5*. In the leaf network, over 50% of ABA-related targets were
classified as high-confidence targets (HCTs) based on DAP-seq evidence (Fig.
5D). In ABA-related GRNs from other organs, *Sl*GBF3 was found to regulate
smaller subsets of ABA-responsive genes, including *PP2C* genes, ABF TFs, and
key regulators such as MYB1 (Solyc12g099120), a TF implicated in ABA-
mediated pathogen susceptibility (Abuqamar *et al.*, 2009) (Supplementary Fig.
S10). Although the leaf-specific GRN contained the largest number of ABA-
related targets, all organ-specific ABA networks maintained a consistent
proportion of HCTs (~50%) (Supplementary Fig. S9). This result suggests a
conserved regulatory role of *Sl*GBF3 across different organs, reinforcing its
significance in ABA-mediated stress responses.
723

724 *TomViz- GRN App: An online tool for the visualization of tomato GRNs*

726 To provide the scientific community with a comprehensive framework of tomato
727 organ-level GRNs and a user-friendly resource, we have developed a public web

platform featuring an interactive interface that allows users to explore the results of this study. Our GRN apps within the TomViz module of the PlantaeViz platform (https://plantaeviz.tomsbiolab.com/tomviz) (Santiago *et al.*, 2024) adhere to the Findability, Accessibility, Interoperability, and Reusability (FAIR) principles. Through the website app, users can interact with organ-level GRNs, select and subset network data for download (Fig. 6A, C). The TomViz-GRNs app provides various features for data analysis. In the Regulatory Targets Tab, users can query individual TFs or genes to explore central regulatory TFs and their validation layers (Fig. 6B). The D3 Subnetwork Tab allows users to upload gene lists and generate GRNs based on specific queries. It categorizes data and enables an organ-level study of stress responses, helping to detect novel regulatory pathways and TFs involved in multiple regulatory cascades (Fig. 6C). The TomViz-GRNs app thus provides an intuitive platform for studying tomato gene regulation and investigating stress responses across different organs.

Discussion

Bioinformatic validation of GRNs

To generate tomato GRNs, our first step was to generate an updated resource of gene models, TFs and functional annotations. Using well-established gene functional annotation pipelines (Jones *et al.*, 2014; Cantalapiedra *et al.*, 2021), we successfully assigned functional annotations to 26,403 of the 37,467 ITAG4-merge genes, considerably improving coverage compared to the previous ITAG4.1 annotation in SolGenomics, which included only 13,142 functionally annotated genes or previous annotations for iTAG4.0 (25,285 genes) (Rivera-Silva *et al.*, 2024).

TF prediction remains a challenging task, as automated approaches usually rely on protein sequence scanning for known DNA-binding domains, which may lead to the inclusion of proteins with DNA-binding capabilities unrelated to TF function (Itzkovitz *et al.*, 2006; Liebold *et al.*, 2024). To avoid such proteins and refine the tomato TF list, we integrated multiple levels of evidence to filter and extract a curated set of 1,840 TFs (representing around 5% of tomato genes). This number

closely aligns with TF numbers reported for tomato in PlantTFDB (1,845 TFs) (Jin *et al.*, 2017) and is slightly higher than those in CisBP (1,773 TFs) (Weirauch *et al.*, 2014) and other annotated TFs from previous studies (1,069 TFs in Kumar et al., 2021). Overall, the percentage of TF coding genes we found is slightly lower than TFs reported for Arabidopsis (approximately 5-10%) (Riechmann and Ratcliffe, 2000) but consistent with estimates reported for other crops, including wheat (5.7%) and rice (6.1%) (Zheng *et al.*, 2016). It is also comparable to values reported for other Solanaceae species, such as eggplant (5.3%) (Wei *et al.*, 2020).

A vast number of transcriptomic studies have been conducted in tomato, covering diverse experimental conditions, organs, and developmental stages. While several efforts have aimed to generate gene expression atlases for tomato gene expression (Ozaki *et al.*, 2010; Fukushima *et al.*, 2012; Gao *et al.*, 2013; Koenig *et al.*, 2013; Arhondakis *et al.*, 2016; Zouine *et al.*, 2017; Bae *et al.*, 2021; Bizouerne *et al.*, 2021; Kumar *et al.*, 2021; Kusano *et al.*, 2022; Li *et al.*, 2024) many studies are limited in scope, often focusing on specific experimental conditions, using older genome assemblies (SL2.4 or SL3.0 with iTAG2.5 or iTAG3.0 annotations), or relying on microarray data. To our knowledge, the gene expression dataset collected in this study represents the most comprehensive to date, compiling over 10,000 RNA-Seq libraries from five major organs and integrating hundreds of Bioprojects performed worldwide. Moreover, the transcriptomes were processed utilizing the latest genome version (SL4.0) and the updated ITAG4-merge annotation, resulting in greater gene coverage. This extensive dataset enabled us to characterize general gene expression patterns at the organ level, facilitating the identification of genes involved in organ-specific functions.

Consistent with previous studies in tomato (Li *et al.*, 2024), we found that most genes meet the threshold for expression across all organs. Similar ubiquitous expression patterns have been reported in other plants, including *Linum usitatissimum* (Qi *et al.*, 2023) and *Zea mays* (Huang *et al.*, 2018), where over 50% of genes are expressed across multiple tissues. Despite this ubiquitous expression, we found differences in expression levels across organs for these genes, showing that organ identity is an important determinant of differential gene

expression, as has been shown in previous studies in Arabidopsis (Aceituno *et al.*, 2008).

Beyond broadly expressed genes, we identified a substantial subset of genes with organ-specific expression that were enriched in biological processes critical for organ function. Our analyses successfully captured known organ-specific genes. Moreover, genes such as *SULTR1;1* and *FER* in roots, *TPD1-l* genes in flowers, *LNG1* and *SlBRC1a* in leaves, stems, and hypocotyls, and ABI TF in seeds were also identified as organ-specific in a recent study in tomato (Li *et al.*, 2024)*.* Furthermore, the TFs also displayed widespread expression across tomato organs, mirroring findings in *Arabidopsis* (Ranjan *et al.*, 2024). Prior studies in tomato indicate that fewer than 20% of expressed TFs are organ-specific (Rohrmann *et al.*, 2012). Nonetheless, despite their broad expression, variations in TF expression levels across organs highlight the dynamic and context-dependent regulation of transcriptional networks governing organ function.

Transcriptomic data has been extensively used to generate biological network models with the purpose of identifying key candidates for functional genomics analyses. Due to the limited availability of TF-target interaction data existing for tomato, the majority of studies have relied on GCNs to infer regulatory relationships and identify co-regulated gene groups (Fukushima *et al.*, 2012; Koenig *et al.*, 2013; Ichihashi *et al.*, 2014; Arhondakis *et al.*, 2016; Yue *et al.*, 2016; Kim *et al.*, 2017; Zouine *et al.*, 2017; Bizouerne *et al.*, 2021; Kusano *et al.*, 2022; Pirona *et al.*, 2023; Wang *et al.*, 2023*a*). Notably, the GCNs lack directionality, making it difficult to establish regulatory interactions. Additionally, many rely on correlations such as Pearson coefficients, which fail to capture non-linear relationships (Escorcia-Rodríguez *et al.*, 2023). To address these limitations, we employed GENIE3, a widely used algorithm for reconstructing directed GRNs in plants *(Chen et al., 2023; De Clercq et al., 2021; Harrington et al., 2020; Huang et al., 2018; Ranjan et al., 2024; Tu et al., 2020)* and other organisms (Huynh-Thu *et al.*, 2010; Huynh-Thu and Geurts, 2019; Cuesta-Astroz *et al.*, 2021; Olivares-Yañez *et al.*, 2021). GENIE3 requires only gene expression data as input, making it particularly suitable for tomato, where TF gene targets remain poorly characterized.

To validate our GRNs, we benchmarked them against available ChIP-Seq data
standard networks, employing the AUPR and AUROC curves. This strategy,
previously used to assess GRN performance in plants (Brooks *et al.*, 2019;
Contreras-López *et al.*, 2022), provides a more centered evaluation of TF-target
interactions than alternative methods that utilize gene co-association to biological
processes or metabolic pathways (Kim *et al.*, 2017; Orduña *et al.*, 2023). Our
analysis determined that a 2% best GENIE3 scores threshold—comprising
660,000–800,000 edges—was a suitable cut-off, aligning with previous GENIE3-
based GRN studies in crops, which typically consider networks containing
around 1 million edges (Harrington et al., 2020; Huang et al., 2018; Ramírez-
González et al., 2018). Notably, the GENIE3-derived GRNs outperformed other
tomato networks from public resources such as PlantRegMap (Tian *et al.*, 2020)
and other genome-scale biological network models (Kim *et al.*, 2017).
Additionally, more than 50% of GENIE3-predicted edges were supported by one
or more independent evidence, including cis-regulatory motif binding predictions,
further reinforcing the biological relevance of these regulatory connections, as it
was proven in other networks (De Clercq *et al.*, 2021; Chen *et al.*, 2023). This
integration of multiple validation strategies enhances the accuracy and functional
significance of inferred GRNs, providing a robust framework for studying
transcriptional regulation in tomato.

Our analysis revealed that while most genes are broadly expressed across
organs, the vast majority of TF-target interactions remain organ-specific. This
pattern has been observed in GRNs from other crops (Huang *et al.*, 2018; Ranjan
et al., 2024), suggesting that gene expression levels play a relevant role in
establishing regulatory interactions underlying organ-specific functions. Notably,
we identified a positive correlation between TF connectivity and target
conservation across organs. Highly connected TFs (hubs) tend to regulate a
similar set of targets in all organs, whereas TFs with fewer connections are more
likely to control organ-specific processes. For example, a highly connected TF
SlHUA (*Solyc12g017410*) primarily regulates genes involved in fundamental
cellular functions, while *SlNGA3* (*Solyc05g004000*) a TF with limited connectivity
exhibits distinct, tissue-specific roles—regulating auxin response in flowers and
defense responses to fungi in seeds, with no enriched function detected in other
organs. Accordingly, *SlNGA3* is homologous to *AtNGA3*, which is associated with

flower development (Salava *et al.*, 2022). An evolutionary constraint may underlie
this phenomenon, as TF-target interactions involving hub genes are more
conserved. Disruptions in these interactions are more likely to be deleterious,
leading to reduced genetic diversity and slower evolutionary rates among hub
TFs. In contrast, tissue-specific TFs, which are less connected, have been
described to evolve faster (Mack *et al.*, 2019).

Biological validation of GRNs involved in ripening

We employed the fruit-specific GRN to identify important TF regulators of fruit
ripening, including RIN and TAGL1 (Karlova *et al.*, 2014). To assess the accuracy
of our networks, we compared their predicted TF-target interactions against
experimentally validated target lists, which included genes confirmed through TF
binding studies (Fujisawa *et al.*, 2013; Zhong *et al.*, 2013; Gao *et al.*, 2019) and
regulatory studies of TF-deficient plants (Li *et al.*, 2018; Zhao *et al.*, 2018; Gao *et
al.*, 2019; Ito *et al.*, 2020). Our results showed a significant enrichment between
the predicted interactions in our GRNs and these validated targets, supporting
the networks' ability to accurately capture *in vivo* regulatory relationships.
Notably, our networks identified RIN targets such as *FUL1*, previously validated
via yeast one-hybrid assays (Fujisawa *et al.*, 2014), as well as essential ripening
genes, including *ACS2, ACS4, E8, EXP1, PSY, NOR,* PSY and *CNR*, which were
confirmed using ChIP-PCR (Martel *et al.*, 2011). Similarly, qPCR validation
supported TAGL1 targets, including *FUL2* (Fujisawa *et al.*, 2014), and ripening-
associated genes like *ACS2, ETR1, ERF2* and *PL* (Itkin et al., 2009). These
findings underscore the predictive strength of our GRNs in identifying key
regulatory interactions governing fruit ripening.
Interestingly, *RIN* and *TAGL1* regulate overlapping sets of ripening-related
genes, but our fruit-specific GRN does not predict a direct regulatory link between
these two TFs. This suggests that their influence on fruit ripening may be
mediated through indirect interactions, such as protein-protein interactions or
epistatic synergistic control of ripening-responsive genes, as proposed in
previous studies (Fujisawa et al., 2014; Jeon et al., 2024). Notably, our GRN
identified *Sl*ARF2A and *Sl*ERF2.E2 as central hubs in the fruit ripening regulatory
network, suggesting that these TFs may play crucial roles in modulating the

ripening process. *Sl*ARF2A has been implicated in the hormonal regulation of
tomato fruit ripening. RNAi-mediated silencing of *Sl*ARF2A results in ripening
defects and ethylene insensitivity, while its overexpression (OX) leads to
accelerated and uneven ripening (Hao *et al.*, 2015; Breitel *et al.*, 2016).
Consistently, our network analysis identified *Sl*ARF2A targets that align with gene
expression changes in OX-ARF2A plants, including *ETR*, *ACS4*, *AP2A*, *ETR3*,
ETR4, *NOR*, and *RIN* (Breitel *et al.*, 2016). In contrast, *Sl*ERF.E2 is associated
with key ripening regulators and ripening-related genes, as its expression is
downregulated in *cnr*, *nor*, and *rin* mutants, yet its precise function remains
unknown (Liu *et al.*, 2016). Given their potential regulatory roles, future studies
should employ TF-binding assays to validate their direct interactions with
ripening-responsive genes and further elucidate their contributions as key hubs
of tomato fruit ripening.

To further explore organ-specific regulatory mechanisms, we examined the
cellular response to abscisic acid (ABA), a biological process that was
ubiquitously enriched in our organ specific gene lists. Our GRN analysis provided
new insights into ABA response cascades, particularly in relation to drought-
response regulation. We identified *Sl*ABF3 and *Sl*ABF5 as important controllers
of drought-responsive genes, as well as ABA-related genes such *as PYL/RCAR*
transporters, *PP2Cs phosphatases* and *SnRK2s protein kinases* (Fujii *et al.*,
2009). We found *Sl*ABF3 targets *Sl*ABF5 and two *PP2C phosphatases*, a direct
interaction previously confirmed in yeast two-hybrid assays (Chen *et al.*, 2016).
Furthermore, *Sl*ABF5 showed a strong connection to drought responses, as
demonstrated by the significant enrichment of its target genes in the drought
responsive transcriptome (Wang et al., 2023). Additionally, its expression levels
vary during drought stress, as reported in earlier studies (Orellana *et al.*, 2010;
Wang *et al.*, 2023*b*). These findings confirm that our leaf-specific GRN
successfully captures ABA-dependent regulatory pathways, reinforcing the role
of *Sl*ABF3/*Sl*ABF5 as regulators of drought-responsive genes expression in
tomato leaves.

SlGBF3 as a novel controller of ABA-related and drought-responsive genes

Our study findings identify the *Sl*GBF3 TF as a novel regulatory hub in
ABA-responsive GRNs, suggesting a key relevance in the drought response and
ABA signaling cascades across different organs, especially in leaves. Prior
research (Bortolami *et al.*, 2024) linked *Sl*GBF3 to drought-responsive co-
expression modules in tomato, but its direct regulatory function remained
unknown. Using DAP-seq analysis, we show direct molecular evidence that
*Sl*GBF3 binds to and regulates multiple ABA-related genes, establishing it as a
key component of ABA-driven transcriptional regulation in tomato. Importantly,
our ortholog and binding motif analyses demonstrated an important conservation
in the DNA-binding preferences between *Sl*GBF3 and its Arabidopsis homolog,
*At*GBF3. In Arabidopsis, *At*GBF3 is documented as a negative regulator of ABA
response, with loss-of-function plants demonstrating an increased sensitivity to
osmotic, drought and pathogen stress, and overexpressor plants showing
reduced ABA sensitivity (Ramegowda *et al.*, 2017; Dixit *et al.*, 2019).
Furthermore, Dixit et al. (2019) found that *At*GBF3 is essential for balancing
drought tolerance and pathogen defense in the context of combined drought
stress and *Pseudomonas syringae* infection.

Our findings suggest that *Sl*GBF3 has conserved its role in ABA-related
responses, with a strong enrichment of genes involved in water deprivation,
abiotic stimulus response and enriched response to ABA target genes. The GRNs
showed that *Sl*GBF3 regulates important ABA-related genes, including *PP2C*
phosphatases, *SnRK2 kinases*, and three ABF TFs (Fujii et al., 2009; Krukowski
et al., 2023; Park et al., 2009). Notably, our study identified targets such as
Abscisic Acid Stress Ripening 1 (ASR1, Solyc04g071610) and *dehydrin-like*
proteins (Solyc02g062390, Solyc02g084850), which are indicators of ABA-
mediated stress responses (Golan *et al.*, 2014).

Exploring tomato GRNs through the TomViz tools

The *Solanum lycopersicum* organ-level GRNs, available through the TomViz
module on the PlantaeViz platform (Santiago et al., 2024), provide a robust and
comprehensive resource for investigating TF-target interactions and organ-
specific regulatory mechanisms. By integrating extensive datasets and
emphasizing regulatory cascades, this tool surpasses traditional GCNs-based

approaches, which are often limited to fruit tissue and a narrower gene set. The

web application offers an accessible yet powerful platform for in-depth regulatory

analysis, enabling researchers to explore tomato gene regulation across diverse

developmental stages, environmental conditions, and genetic backgrounds. Its

utility extends to various research contexts, facilitating novel discoveries in

tomato biology and advancing functional genomics studies.

Supplementary data

Supplementary Figures:
971
Supplementary Fig. S1. Distribution of tomato transcriptomic datasets across
organs and experimental conditions.
Supplementary Fig. S2. Heatmap of Gene Set Enrichment Analysis (GSEA)
results for organ-specific gene expression (FDR-adjusted p-value < 0.05).
Supplementary Fig. S3. Accuracy analysis of organ-specific GRNs.
Supplementary Fig. S4. GSEA results (FDR-adjusted p-value < 0.05) for the five
highest connected TFs shared across organ-level GRNs (conserved targets).
Supplementary Fig. S5. GSEA results (FDR-adjusted p-value < 0.05) for the five
lowest connected TFs shared across organ-level GRNs (specific targets).
Supplementary Fig. S6. Genome-wide identification of *Sl*GBF3 binding targets
using DAP-seq.
Supplementary Fig. S7. Enrichment analysis of *Sl*GBF3 targets in organ-
specific GRNs compared to DAP-seq binding targets**.**
Supplementary Fig. S8. GSEA results (FDR-adjusted p-value < 0.05) of *Sl*GBF3
DAP-seq targets.
Supplementary Fig. S9. Network visualization of *Sl*GBF3-regulated abscisic
acid (ABA)-related genes within Root, Seed, Flower and Fruit GRNs.
989
990 **Supplementary Tables:**
991
Supplementary Table S1. Summary of Tomato Gene Annotations.
Supplementary Table S2. Gene Ontology (GO) annotations for *Solanum*
lycopersicum (ITAG 4.0c).
Supplementary Table S3. Transcription factor (TF) evidence and position weight
matrices (PWMs).
Supplementary Table S4. Transcriptomic libraries used for GRN construction.
Supplementary Table S5. Organ-level expression patterns.
Supplementary Table S6. Enriched GO terms shared across tomato organs.
Supplementary Table S7. Metadata of ChIP-seq libraries.
Supplementary Table S8. Enrichment analysis of GENIE3 predicted TF-target
pairings.
Supplementary Table S9. Comparative enrichment analysis of GENIE3-
predicted TF-target interactions and previously reported networks.
Supplementary Table S10. Metadata of Open Chromatin Sites (OCS) libraries.

Supplementary Table S11. TFs analysis reveals connectivity and target
conservation distribution across tomato organs-level GRNs.
Supplementary Table S12. Ripening-related genes and their classification as
RIN or TAGL1 targets.
Supplementary Table S13. Network analysis of key TFs in the fruit ripening
network.
Supplementary Table S14. Genes annotated with the GO term "Response to
ABA" or ABA-related genes.
Supplementary Table S15. ABA-related genes in tomato regulated by ABF TFs.
Supplementary Table S16. Leaf GRN targets of ABF3 and ABF5 regulating
ABA-related and drought-responsive genes.
Supplementary Table S17. Network analysis of *Sl*GBF3 (Solyc01g095460)
across all organ-specific GRNs.
Supplementary Table S18. DAP-seq analysis results for *Sl*GBF3.
1020

Acknowledgements

This research was supported by the computing infrastructure of the Center for
Genomics and Bioinformatics, Universidad Mayor and the HPC cluster Garnatxa
at the Institute for Integrative Systems Biology (I^{2}SysBio). We thank the staff at
SolGenomics (Dr. Surya Saha) for providing the iTAG4.2 beta annotation file.

Author contributions

JDF, JMA, JTM and EAV: conceptualization; JDF, DN-P, AS: data curation; JDF,
DN-P, AS, J Canan, SC-R, AC, TCM, LM and NRJ: methodology and
investigation; JDF, DN-P, AS, TCM: formal analysis; EAV, JMA and JTM: funding
acquisition; J Canales, JMA, JTM, and EAV: supervision; JDF, EAV: writing -
original draft; JDF, J Canales, JMA, JTM and EAV writing - review and editing.
All authors provided critical feedback and approved the final version of the
manuscript.

Conflict of interest

No conflict of interest declared.

Funding

This work was supported by the Agencia Nacional de Investigación y Desarrollo
(ANID)-Millennium Science Initiative Program, [Millennium Institute for Integrative
Biology iBio ICN17_022 to EAV, JMA and JC, Millennium Nucleus in Data
Science for Plant Resilience NCN2024_047 to EAV and JMA], ANID-Fondo de

Desarrollo Científico y Tecnológico (FONDECYT) [grant 1211130 and 1250631

to EAV, 1230833 and 1211040 to JC and 1210389 and 1250403 to JMA], ANID-

Anillo [grant ACT210007 to EAV], ANID-Vinculación Internacional [grant

FOVI230159 to EAV, JC, JMA and JTM], Ministerio de Ciencia, Innovación y

Universidades (MCIU, Spain) [grant Valinet-PID2021-128865NB-I00 to JTM],

Agencia Estatal de Investigación (AEI, Spain) and Fondo Europeo de Desarrollo

Regional (FEDER, European Union) to JTM, Doctoral grant GVA-

PROMETEO/2021/056-01 to AS and ANID-Beca Doctoral 21230478 to JDF.

These funding agencies were not involved in the design of the study, data

collection, analysis, interpretation of data, or in writing the manuscript.

Data availability

The organ-level GRNs can be extracted and visualized through the GRN apps

found within the TomViz module of the PlantaeViz platform (Santiago *et al.*, 2024)

available at https://plantaeviz.tomsbiolab.com/tomviz. The Supplementary

Tables S4, S7, and S10 include the metadata of the public RNA-seq, ATAC-seq,

DNASE-seq experiments, as well as the ChIP-seq libraries analyzed in this study.

The DAP-seq data underlying this article is available at the NCBI Sequence Read

Archive (SRA) database and can be accessed with accession number

PRJNA1236412. The scripts used for analyzing RNA-seq, ATAC-seq, DNASE-

seq, ChIP-seq, DAP-seq, the GENIE3 network generation pipeline as well of all

the GRNs are available at:

https://github.com/ibioChile/VidalLab/tree/master/Pipelines.

1063

1064   **References**

1065

Abuqamar S, Luo H, Laluk K, Mickelbart M V., Mengiste T. 2009. Crosstalk

between biotic and abiotic stress responses in tomato is mediated by the AIM1

transcription factor. The Plant Journal **58**, 347–360.

Aceituno FF, Moseyko N, Rhee SY, Gutiérrez RA. 2008. The rules of gene

expression in plants: Organ identity and gene body methylation are key factors

for regulation of gene expression in Arabidopsis thaliana. BMC Genomics **9**, 1–

14.

Arhondakis S, Bita CE, Perrakis A, Manioudaki ME, Krokida A, Kaloudas D, Kalaitzis P, Pezzotti M. 2016. In silico transcriptional regulatory networks involved in tomato fruit ripening. Frontiers in Plant Science **7**, 195142.

Aviña-Padilla K, Zambada-Moreno O, Jimenez-Limas MA, Hammond RW, Hernández-Rosales M. 2023. Dynamic co-expression modular network analysis of bHLH transcription factors regulation in the potato spindle tuber viroid-tomato pathosystem. bioRxiv, 2023.11.10.566618.

Babu MM, Luscombe NM, Aravind L, Gerstein M, Teichmann SA. 2004. Structure and evolution of transcriptional regulatory networks. Current Opinion in Structural Biology **14**, 283–291.

Bae SH, Park J, Park SJ, Han J, Oh JH. 2021. Transcriptome data for tissue-specific genes in four reproductive organs at three developmental stages of micro-tom tomato. Data in Brief **34**, 106715.

Van Bel M, Silvestri F, Weitz EM, Kreft L, Botzki A, Coppens F, Vandepoele K. 2022. PLAZA 5.0: extending the scope and power of comparative and functional genomics in plants. Nucleic Acids Research **50**, D1468–D1474.

Bizouerne E, Buitink J, Vu BL, Vu JL, Esteban E, Pasha A, Provart N, Verdier J, Leprince O. 2021. Gene co-expression analysis of tomato seed maturation reveals tissue-specific regulatory networks and hubs associated with the acquisition of desiccation tolerance and seed vigour. BMC Plant Biology **21**, 1–23.

Bortolami G, de Werk TA, Larter M, Thonglim A, Mueller-Roeber B, Balazadeh S, Lens F. 2024. Integrating gene expression analysis and ecophysiological responses to water deficit in leaves of tomato plants. Scientific Reports 2024 14:1 **14**, 1–18.

Breitel DA, Chappell-Maor L, Meir S, *et al.* 2016. AUXIN RESPONSE FACTOR 2 Intersects Hormonal Signals in the Regulation of Tomato Fruit Ripening. PLOS Genetics **12**, e1005903.

Bres C, Petit J, Reynoud N, Brocard L, Marion D, Lahaye M, Bakan B, Rothan C. 2022. The SlSHN2 transcription factor contributes to cuticle formation and epidermal patterning in tomato fruit. Molecular Horticulture **2**, 1–20.

Brooks MD, Cirrone J, Pasquino A V., *et al.* 2019. Network Walking charts transcriptional dynamics of nitrogen signaling by integrating validated and predicted genome-wide interactions. Nature Communications **10**.

Cantalapiedra CP, Herṇandez-Plaza A, Letunic I, Bork P, Huerta-Cepas J.
2021. eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and
Domain Prediction at the Metagenomic Scale. Molecular Biology and Evolution
38, 5825–5829.

Castro-Mondragon JA, Riudavets-Puig R, Rauluseviciute I, *et al.* 2022.
JASPAR 2022: the 9th release of the open-access database of transcription
factor binding profiles. Nucleic Acids Research **50**, D165–D173.

Chai LE, Loh SK, Low ST, Mohamad MS, Deris S, Zakaria Z. 2014. A review
on the computational approaches for gene regulatory network construction.
Computers in Biology and Medicine **48**, 55–65.

Chang HC, Tsai MC, Wu SS, Chang IF. 2019. Regulation of ABI5 expression by
ABF3 during salt stress responses in Arabidopsis thaliana. Botanical Studies **60**,
1–14.

Chen Y, Guo Y, Guan P, *et al.* 2023. A wheat integrative regulatory network from
large-scale complementary functional datasets enables trait-associated gene
discovery for crop improvement. Molecular Plant **16**, 393–414.

Chen P, Sun YF, Kai W Bin, Liang B, Zhang YS, Zhai XW, Jiang L, Du YW,
Leng P. 2016. Interactions of ABA signaling core components (SlPYLs,
SlPP2Cs, and SlSnRK2s) in tomato (Solanum lycopersicon). Journal of Plant
Physiology **205**, 67–74.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics **34**, i884–i890.

Chong L, Xu R, Huang P, Guo P, Zhu M, Du H, Sun X, Ku L, Zhu JK, Zhu Y.
2022. The tomato OST1–VOZ1 module regulates drought-mediated flowering.
The Plant Cell **34**, 2001–2018.

De Clercq I, Van de Velde J, Luo X, Liu L, Storme V, Van Bel M, Pottie R,
Vaneechoutte D, Van Breusegem F, Vandepoele K. 2021. Integrative
inference of transcriptional networks in Arabidopsis yields novel ROS signalling
regulators. Nature Plants 2021 7:4 **7**, 500–513.

Contreras-López O, Vidal EA, Riveras E, *et al.* 2022. Spatiotemporal analysis
identifies ABF2 and ABF3 as key hubs of endodermal response to nitrate.
Proceedings of the National Academy of Sciences of the United States of
America **119**, e2107879119.

Cuesta-Astroz Y, Gischkow Rucatti G, Murgas L, SanMartín CD, Sanhueza M, Martin AJM. 2021. Filtering of Data-Driven Gene Regulatory Networks Using Drosophila melanogaster as a Case Study. Frontiers in Genetics **12**.

Ding X, Zhang D, Gu D, Li Z, Liang H, Zhu H, Jiang Y, Duan X. 2022. The histone H3K27 demethylase SlJMJ4 promotes dark- and ABA-induced leaf senescence in tomato. Horticulture Research **9**.

Dixit SK, Gupta A, Fatima U, Senthil-Kumar M. 2019. AtGBF3 confers tolerance to Arabidopsis thaliana against combined drought and Pseudomonas syringae stress. Environmental and Experimental Botany **168**, 103881.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics **29**, 15–21.

Doidy J, Li Y, Neymotin B, Edwards MB, Varala K, Gresham D, Coruzzi GM. 2016. 'Hit-and-Run' transcription: De novo transcription initiated by a transient bZIP1 'hit' persists after the 'run'. BMC Genomics **17**.

Du M, Zhao J, Tzeng DTW, *et al.* 2017. MYC2 orchestrates a hierarchical transcriptional cascade that regulates jasmonate-mediated plant immunity in tomato. Plant Cell **29, 1883–1906.

Emms DM, Kelly S. 2019. OrthoFinder: Phylogenetic orthology inference for comparative genomics. Genome Biology **20**, 1–14.

Escorcia-Rodríguez JM, Gaytan-Nuñez E, Hernandez-Benitez EM, Zorro-Aranda A, Tello-Palencia MA, Freyre-González JA. 2023. Improving gene regulatory network inference and assessment: The importance of using network structure. Frontiers in Genetics **14**, 1143382.

Ezura K, Ji-Seong K, Mori K, Suzuki Y, Kuhara S, Ariizumi T, Ezura H. 2017. Genome-wide identification of pistil-specific genes expressed during fruit set initiation in tomato (Solanum lycopersicum). PLoS ONE **12**.

Fernandez-Pozo N, Menda N, Edwards JD, *et al.* 2015. The Sol Genomics Network (SGN)—from genotype to phenotype to breeding. Nucleic Acids Research **43, D1036–D1041.

Fujii H, Chinnusamy V, Rodrigues A, Rubio S, Antoni R, Park SY, Cutler SR, Sheen J, Rodriguez PL, Zhu JK. 2009. In vitro reconstitution of an abscisic acid signalling pathway. Nature 2009 462:7273 **462**, 660–664.

Fujisawa M, Nakano T, Ito Y. 2011. Identification of potential target genes for
the tomato fruit-ripening regulator RIN by chromatin immunoprecipitation. BMC
Plant Biology **11**.

Fujisawa M, Nakano T, Shima Y, Ito Y. 2013. A Large-Scale Identification of
Direct Targets of the Tomato MADS Box Transcription Factor RIPENING
INHIBITOR Reveals the Regulation of Fruit Ripening. The Plant Cell **25**, 371–
386.

Fujisawa M, Shima Y, Nakagawa H, Kitagawa M, Kimbara J, Nakano T,
Kasumi T, Ito Y. 2014. Transcriptional Regulation of Fruit Ripening by Tomato
FRUITFULL Homologs and Associated MADS Box Proteins. The Plant Cell **26**,
89–101.

Fukushima A, Nishizawa T, Hayakumo M, Hikosaka S, Saito K, Goto E,
Kusano M. 2012. Exploring Tomato Gene Functions Based on Coexpression
Modules Using Graph Clustering and Differential Coexpression Approaches   .
Plant Physiology **158**, 1487–1502.

Gao C, Ju Z, Li S, Zuo J, Fu D, Tian H, Luo Y, Zhu B. 2013. Deciphering
Ascorbic Acid Regulatory Pathways in Ripening Tomato Fruit Using a Weighted
Gene Correlation Network Analysis Approach. Journal of Integrative Plant
Biology **55**, 1080–1091.

Gao Y, Zhu N, Zhu X, *et al.* 2019. Diversity and redundancy of the ripening
regulatory networks revealed by the fruitENCODE and the new CRISPR/Cas9
CNR and NOR mutants. Horticulture Research **6**.

Gascuel Q, Diretto G, Monforte AJ, Fortes AM, Granell A. 2017. Use of natural
diversity and biotechnology to increase the quality and nutritional content of
tomato and grape. Frontiers Media S.A.

Golan I, Guadalupe Dominguez P, Konrad Z, Shkolnik-Inbar D, Carrari F,
Bar-Zvi D. 2014. Tomato ABSCISIC ACID STRESS RIPENING (ASR) Gene
Family Revisited. PLoS ONE **9**, e107117.

Grant CE, Bailey TL, Noble WS. 2011. FIMO: Scanning for occurrences of a
given motif. Bioinformatics **27**, 1017–1018.

Hao Y, Hu G, Breitel D, Liu M, Mila I, Frasse P, Fu Y, Aharoni A, Bouzayen
M, Zouine M. 2015. Auxin Response Factor SlARF2 Is an Essential Component
of the Regulatory Mechanism Controlling Fruit Ripening in Tomato. PLOS
Genetics **11**, e1005649.

Harrington SA, Backhaus AE, Singh A, Hassani-Pak K, Uauy C. 2020. The Wheat GENIE3 Network Provides Biologically-Relevant Information in Polyploid Wheat. G3 Genes|Genomes|Genetics **10**, 3675–3686.

Hawar A, Xiong S, Yang Z, Sun B. 2022. Histone Acetyltransferase SlGCN5 Regulates Shoot Meristem and Flower Development in Solanum lycopersicum. Frontiers in Plant Science **12**, 805879.

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK. 2010. Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. Molecular Cell **38**, 576–589.

Hendelman A, Zebell S, Rodriguez-Leal D, *et al.* 2021. Conserved pleiotropy of an ancient plant homeobox gene uncovered by cis-regulatory dissection. Cell **184, 1724-1739.e16.

**Hitz BC, Lee J-W, Jolanki O, *et al.* 2023. The ENCODE Uniform Analysis Pipelines. bioRxiv, 2023.04.04.535623.

**Hosmani PS, Flores-Gonzalez M, Van De Geest H, *et al.* 2019. An improved de novo assembly and annotation of the tomato reference genome using single-molecule sequencing, Hi-C proximity ligation and optical maps. bioRxiv, 767764.

Hsieh TH, Li CW, Su RC, Cheng CP, Sanjaya, Tsai YC, Chan MT. 2010. A tomato bZIP transcription factor, SlAREB, is involved in water deficit and salt stress response. Planta **231**, 1459–1473.

Huang Y, An J, Sircar S, *et al.* 2023. HSFA1a modulates plant heat stress responses and alters the 3D chromatin organization of enhancer-promoter interactions. Nature Communications **14, 469.

Huang J, Zheng J, Yuan H, McGinnis K. 2018. Distinct tissue-specific transcriptional regulation revealed by gene regulatory networks in maize. BMC Plant Biology **18**.

Hutin S, Blanc-Mathieu R, Rieu P, Parcy F, Lai X, Zubieta C. 2023. Identification of Plant Transcription Factor DNA-Binding Sites Using seq-DAP-seq. Methods in Molecular Biology **2698**, 119–145.

Huynh-Thu VA, Geurts P. 2019. Unsupervised Gene Network Inference with Decision Trees and Random Forests. 195–215.

Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. 2010. Inferring Regulatory
Networks from Expression Data Using Tree-Based Methods. PLoS ONE **5**,
e12776.

Ichihashi Y, Aguilar-Martínez JA, Farhi M, Chitwood DH, Kumar R, Millon L
V., Peng J, Maloof JN, Sinha NR. 2014. Evolutionary developmental
transcriptomics reveals a gene network module regulating interspecific diversity
in plant leaf shape. Proceedings of the National Academy of Sciences of the
United States of America **111**.

Itkin M, Seybold H, Breitel D, Rogachev I, Meir S, Aharoni A. 2009. TOMATO
AGAMOUS-LIKE 1 is a component of the fruit ripening regulatory network. The
Plant Journal **60**, 1081–1095.

Ito Y, Sekiyama Y, Nakayama H, *et al.* 2020. Allelic Mutations in the Ripening -
Inhibitor Locus Generate Extensive Variation in Tomato Ripening. Plant
Physiology **183**, 80–95.

Itzkovitz S, Tlusty T, Alon U. 2006. Coding limits on the number of transcription
factors. BMC Genomics **7**, 1–15.

Jeon C, Chung MY, Lee JM. 2024. Reassessing the contribution of TOMATO
AGAMOUS-LIKE1 to fruit ripening by CRISPR/Cas9 mutagenesis. Plant Cell
Reports **43**, 1–7.

Jiang G, Li Z, Ding X, Zhou Y, Lai H, Jiang Y, Duan X. 2023. WUSCHEL-
related homeobox transcription factor SlWOX13 regulates tomato fruit ripening.
Plant Physiology doi: 10.1093/PLPHYS/KIAD623.

Jin J, Tian F, Yang D-C, Meng Y-Q, Kong L, Luo J, Gao G. 2017. PlantTFDB
4.0: toward a central hub for transcription factors and regulatory interactions in
plants. Nucleic Acids Research **45**, D1040–D1045.

Jones P, Binns D, Chang HY, *et al.* 2014. InterProScan 5: genome-scale protein
function classification. Bioinformatics (Oxford, England) **30**, 1236–1240.

Kajala K, Gouran M, Shaar-Moshe L, *et al.* 2021. Innovation, conservation, and
repurposing of gene function in root cell type development. Cell **184**, 3333-
3348.e19.

Kang JY, Choi HI, Im MY, Soo YK. 2002. Arabidopsis Basic Leucine Zipper
Proteins That Mediate Stress-Responsive Abscisic Acid Signaling. The Plant Cell
14, 343–357.

Kans J. 2010. Entrez Direct: E-utilities on the Unix Command Line.

Karlova R, Chapman N, David K, Angenent GC, Seymour GB, De Maagd RA.
2014. Transcriptional control of fleshy fruit development and ripening. Journal of
Experimental Botany **65**, 4527–4541.

Kim H, Kim BS, Shim JE, Hwang S, Yang S, Kim E, Iyer-Pascuzzi AS, Lee I.
2017. TomatoNet: A Genome-wide Co-functional Network for Unveiling Complex
Traits of Tomato, a Model Crop for Fleshy Fruits. Molecular Plant **10**, 652–655.

Kimura S, Sinha N. 2008. Tomato (Solanum lycopersicum): A model fruit-
bearing crop. Cold Spring Harbor Protocols **3**.

Koenig D, Jiménez-Gómez JM, Kimura S, *et al.* 2013. Comparative
transcriptomics reveals patterns of selection in domesticated and wild tomato.
Proceedings of the National Academy of Sciences of the United States of
America **110**, E2655–E2662.

Krukowski P, Colanero S, Sutti A, Martignago D, Conti L. 2023. How Changes
in ABA Accumulation and Signaling Influence Tomato Drought Responses and
Reproductive Development. International Journal of Plant Biology **14**, 162–176.

Kumar V, Singh D, Majee A, Singh S, Roohi, Asif MH, Sane AP, Sane VA.
2021. Identification of tomato root growth regulatory genes and transcription
factors through comparative transcriptomic profiling of different tissues.
Physiology and Molecular Biology of Plants **27**, 1173.

Kusano M, Worarad K, Fukushima A, *et al.* 2022. Transcriptomic, Hormonomic
and Metabolomic Analyses Highlighted the Common Modules Related to
Photosynthesis, Sugar Metabolism and Cell Division in Parthenocarpic Tomato
Fruits during Early Fruit Set. Cells **11**, 1420.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2.
Nature Methods **9**, 357–359.

Lee YK, Kim GT, Kim IJ, Park J, Kwak SS, Choi G, Chung W II. 2006.
LONGIFOLIA1 and LONGIFOLIA2, two homologous genes,regulate longitudinal
cell elongation in Arabidopsis. Development **133**, 4305–4314.

Leinonen R, Sugawara H, Shumway M. 2011. The Sequence Read Archive.
Nucleic Acids Research **39**, D19–D21.

Li S, Chen K, Grierson D. 2019. A critical evaluation of the role of ethylene and
MADS transcription factors in the network controlling fleshy fruit ripening. New
Phytologist **221**, 1724–1741.

Li S, Chen K, Grierson D. 2021*a*. Molecular and Hormonal Mechanisms Regulating Fleshy Fruit Ripening. Cells 2021, Vol. 10, Page 1136 **10**, 1136.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics **25**, 2078–2079.

Li C, Hou X, Qi N, Liu H, Li Y, Huang D, Wang C, Liao W. 2021*b*. Insight into ripening-associated transcription factors in tomato: A review. Scientia Horticulturae **288**, 110363.

Li J, Sun B, Xu Q, Jiang L, Wang N. 2024. Transcriptome-level analysis of gene expressions in different tissues of tomato and key gene identifications during seed germination. Scientia Horticulturae **337**, 113565.

Li S, Xu H, Ju Z, Cao D, Zhu H, Fu D, Grierson D, Qin G, Luo Y, Zhu B. 2018. The RIN-MC Fusion of MADS-Box Transcription Factors Has Transcriptional Activity and Modulates Expression of Many Ripening Genes. Plant Physiology **176**, 891–909.

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics **30**, 923–930.

Liebold J, Neuhaus F, Geiser J, Kurtz S, Baumbach J, Newaz K. 2024. Transcription factor prediction using protein 3D secondary structures. (PL Martelli, Ed.). Bioinformatics **41**.

Lira BS, Oliveira MJ, Shiose L, Wu RTA, Rosado D, Lupi ACD, Freschi L, Rossi M. 2020. Light and ripening-regulated BBX protein-encoding genes in Solanum lycopersicum. Scientific Reports **10**, 19235.

Liu M, Gomes BL, Mila I, *et al.* 2016. Comprehensive Profiling of Ethylene Response Factor Expression Identifies Ripening-Associated ERF Genes and Their Link to Key Regulators of Fruit Ripening in Tomato. Plant Physiology **170**, 1732–1744.

Liu Y, Shi Y, Zhu N, Zhong S, Bouzayen M, Li Z. 2020. SlGRAS4 mediates a novel regulatory pathway promoting chilling tolerance in tomato. Plant Biotechnology Journal **18**, 1620–1633.

López-Vidriero I, Godoy M, Grau J, Peñuelas M, Solano R, Franco-Zorrilla JM. 2021. DNA features beyond the transcription factor binding site specify target

recognition by plant MYC2-related bHLH proteins. Plant Communications **2**, 100232.

Lü P, Yu S, Zhu N, *et al.* 2018. Genome encode analyses reveal the basis of convergent evolution of fleshy fruit ripening. Nature Plants **4**, 784–791.

Machanick P, Bailey TL. 2011. MEME-ChIP: motif analysis of large DNA datasets. Bioinformatics **27**, 1696–1697.

Mack KL, Phifer-Rixey M, Harr B, Nachman MW. 2019. Gene Expression Networks Across Multiple Tissues Are Associated with Rates of Molecular Evolution in Wild House Mice. Genes 2019, Vol. 10, Page 225 **10**, 225.

Maere S, Heymans K, Kuiper M. 2005. BiNGO: a Cytoscape plugin to assess overrepresentation of Gene Ontology categories in Biological Networks. Bioinformatics **21**, 3448–3449.

Maher KA, Bajic M, Kajala K, *et al.* 2018. Profiling of accessible chromatin regions across multiple plant species and cell types reveals common gene regulatory principles and new control modules. Plant Cell **30**, 15–36.

Martel C, Vrebalov J, Tafelmeyer P, Giovannoni JJ. 2011. The Tomato MADS-Box Transcription Factor RIPENING INHIBITOR Interacts with Promoters Involved in Numerous Ripening Processes in a COLORLESS NONRIPENING-Dependent Manner. Plant Physiology **157**, 1568–1579.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal **17**, 10.

Martín-Trillo M, Grandío EG, Serra F, Marcel F, Rodríguez-Buey ML, Schmitz G, Theres K, Bendahmane A, Dopazo H, Cubas P. 2011. Role of tomato BRANCHED1-like genes in the control of shoot branching. The Plant Journal **67**, 701–714.

Meuleman W, Muratov A, Rynes E, *et al.* 2020. Index and biological spectrum of human DNase I hypersensitive sites. Nature **584**, 244–251.

Moyano TC, Gutiérrez RA, Alvarez JM. 2021. Genomic Footprinting Analyses from DNase-seq Data to Construct Gene Regulatory Networks. 25–46.

Olivares-Yañez C, Sánchez E, Pérez-Lara G, Seguel A, Camejo PY, Larrondo LF, Vidal EA, Canessa P. 2021. A comprehensive transcription factor and DNA-binding motif resource for the construction of gene regulatory networks in Botrytis cinerea and Trichoderma atroviride. Computational and Structural Biotechnology Journal **19**, 6212–6228.

Orduña L, Santiago A, Navarro-Payá D, Zhang C, Wong DCJ, Matus JT.
2023. Aggregated gene co-expression networks predict transcription factor
regulatory landscapes in grapevine. Journal of Experimental Botany **74**, 6522–
6540.

Orellana S, Yañez M, Espinoza A, Verdugo I, González E, Ruiz-Lara S,
Casaretto JA. 2010. The transcription factor SlAREB1 confers drought, salt
stress tolerance and regulates biotic and abiotic stress-related genes in tomato.
Plant, Cell & Environment **33**, 2191–2208.

Ozaki S, Ogata Y, Suda K, *et al.* 2010. Coexpression Analysis of Tomato Genes
and Experimental Verification of Coordinated Expression of Genes Found in a
Functionally Enriched Coexpression Module. DNA Research **17**, 105–116.

Pan Y, Bradley G, Pyke K, *et al.* 2013. Network Inference Analysis Identifies an
APRR2-Like Gene Linked to Pigment Accumulation in Tomato and Pepper Fruits.
Plant Physiology **161**, 1476–1485.

Pan X, Wang C, Liu Z, Gao R, Feng L, Li A, Yao K, Liao W. 2023. Identification
of ABF/AREB gene family in tomato (Solanum lycopersicum L.) and functional
analysis of ABF/AREB in response to ABA and abiotic stresses. PeerJ **11**,
e15310.

Park P. 2009. ChIP–seq: advantages and challenges of a maturing technology.
Nature Reviews Genetics 2009 10:10 **10**, 669–680.

Park SY, Fung P, Nishimura N, *et al.* 2009. Abscisic acid inhibits type 2C protein
phosphatases via the PYR/PYL family of START proteins. Science **324**, 1068–
1071.

Pirona R, Frugis G, Locatelli F, Mattana M, Genga A, Baldoni E. 2023.
Transcriptomic analysis reveals the gene regulatory networks involved in leaf and
root response to osmotic stress in tomato. Frontiers in Plant Science **14**,
1155797.

Qi F, Wang F, Xiaoyang C, Wang Z, Lin Y, Peng Z, Zhang J, Wang N, Zhang
J. 2023. Gene Expression Analysis of Different Organs and Identification of AP2
Transcription Factors in Flax (Linum usitatissimum L.). Plants **12**, 3260.

Qiu Z, Li R, Zhang S, Wang K, Xu M, Li J, Du Y, Yu H, Cui X. 2016. Identification
of Regulatory DNA Elements Using Genome-wide Mapping of DNase I
Hypersensitive Sites during Tomato Fruit Development. Molecular Plant **9**, 1168–
1182.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics **26**, 841–842.

Ramegowda V, Gill US, Sivalingam PN, *et al.* 2017. GBF3 transcription factor
imparts drought tolerance in Arabidopsis thaliana. Scientific Reports 2017 7:1 **7**,
1–13.

Ramírez-González RH, Borrill P, Lang D, *et al.* 2018. The transcriptional
landscape of polyploid wheat. Science **361**.

Ranjan R, Srijan S, Balekuttira S, *et al.* 2024. Organ-delimited gene regulatory
networks provide high accuracy in candidate transcription factor selection across
diverse processes. Proceedings of the National Academy of Sciences of the
United States of America **121**, e2322751121.

Reynoso MA, Kajala K, Bajic M, *et al.* 2019. Evolutionary flexibility in flooding
response circuitry in angiosperms. Science **365**, 1291–1295.

Ricardi MM, González RM, Zhong S, *et al.* 2014. *Genome-wide data (ChIP-*
seq) enabled identification of cell wall-related and aquaporin genes as targets of
tomato ASR1, a drought stress-responsive transcription factor.

Riechmann JL, Ratcliffe OJ. 2000. A genomic perspective on plant transcription
factors. Current Opinion in Plant Biology **3**, 423–434.

Rivera-Silva R, Chávez Montes RA, Jaimes-Miranda F. 2024. Gene ontology
functional annotation datasets for the ITAG3.2 and ITAG4.0 tomato (Solanum
lycopersicum) genome annotations. Data in Brief **54**, 110401.

Salava H, Thula S, Sánchez AS, Nodzyński T, Maghuly F. 2022. Genome
Wide Identification and Annotation of NGATHA Transcription Factor Family in
Crop Plants. International Journal of Molecular Sciences **23**, 7063.

Salavaty A, Ramialison M, Currie PD. 2020. Integrated Value of Influence: An
Integrative Method for the Identification of the Most Influential Nodes within
Networks. Patterns **1**, 100052.

Santiago A, Orduña L, David Fernández J, Vidal Á, de Martín-Agirre I, Lisón
P, Vidal EA, Navarro-Payá D, Tomás Matus J. 2024. The Plantae Visualization
Platform: a comprehensive web-based tool for the integration, visualization, and
analysis of omic data across plant and related species. bioRxiv,
2024.12.19.629382.

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N,
Schwikowski B, Ideker T. 2003. Cytoscape: A Software Environment for

Integrated Models of Biomolecular Interaction Networks. Genome Research **13**, 2498–2504.

Siloto RMP, Findlay K, Lopez-Villalobos A, Yeung EC, Nykiforuk CL, Moloney MM. 2006. The Accumulation of Oleosins Determines the Size of Seed Oilbodies in Arabidopsis. The Plant Cell **18**, 1961–1974.

Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms. PLOS ONE **6**, e21800.

Swift J, Coruzzi GM. 2017. A matter of time — How transient transcription factor interactions create dynamic gene regulatory networks. Biochimica et Biophysica Acta - Gene Regulatory Mechanisms **1860**, 75–83.

Takahashi H, Watanabe-Takahashi A, Smith FW, Blake-Kalff M, Hawkesford MJ, Saito K. 2000. The roles of three functional sulphate transporters involved in uptake and translocation of sulphate in Arabidopsis thaliana. The Plant Journal **23**, 171–182.

Tian F, Yang DC, Meng YQ, Jin J, Gao G. 2020. PlantRegMap: charting functional regulatory maps in plants. Nucleic Acids Research **48**, D1104–D1113.

Tu X, Mejía-Guerra MK, Valdes Franco JA, *et al.* 2020. Reconstructing the maize leaf regulatory network using ChIP-seq data of 104 transcription factors. Nature Communications **11**, 5089.

Tu X, Ren S, Shen W, *et al.* 2022. Limited conservation in cross-species comparison of GLK transcription factor binding suggested wide-spread cistrome divergence. Nature Communications 2022 13:1 **13**, 1–12.

Uno Y, Furihata T, Abe H, Yoshida R, Shinozaki K, Yamaguchi-Shinozaki K. 2000. Arabidopsis basic leucine zipper transcription factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions. Proceedings of the National Academy of Sciences of the United States of America **97**, 11632–11637.

Vidal EA, Alvarez JM, Araus V, *et al.* 2020. Nitrate in 2020: Thirty Years from Transport to Signaling Networks. The Plant Cell **32**, 2094–2119.

Vishwakarma K, Upadhyay N, Kumar N, *et al.* 2017. Abscisic acid signaling and abiotic stress tolerance in plants: A review on current knowledge and future prospects. Frontiers in Plant Science **8**, 228798.

Wang B, Wang J, Yang T, Wang J, Dai Q, Zhang F, Xi R, Yu Q, Li N. 2023*a*. The transcriptional regulatory network of hormones and genes under salt stress

in tomato plants (Solanum lycopersicum L.). Frontiers in Plant Science **14**, 1115593.

Wang L, Zhou Y, Ding Y, Chen C, Chen X, Su N, Zhang X, Pan Y, Li J. 2023*b*. Novel flavin-containing monooxygenase protein FMO1 interacts with CAT2 to negatively regulate drought tolerance through ROS homeostasis and ABA signaling pathway in tomato. Horticulture Research **10**.

Wei Q, Wang J, Wang W, Hu T, Hu H, Bao C. 2020. A high-quality chromosome-level genome assembly reveals genetics for important traits in eggplant. Horticulture Research 2020 7:1 **7**, 1–15.

Weirauch MT, Yang A, Albu M, *et al.* 2014. Determination and inference of eukaryotic transcription factor sequence specificity. Cell **158**, 1431–1443.

Wolfe CJ, Kohane IS, Butte AJ. 2005. Systematic survey reveals general applicability of 'guilt-by-association' within gene coexpression networks. BMC Bioinformatics **6**, 227.

Xie L, Dong C, Shang Q. 2019. Gene co-expression network analysis reveals pathways associated with graft healing by asymmetric profiling in tomato. BMC Plant Biology **19**, 1–12.

Xu T, Liu X, Wang R, Dong X, Guan X, Wang Y, Jiang Y, Shi Z, Qi M, Li T. 2016. SlARF2a plays a negative role in mediating axillary shoot formation. Scientific Reports 2016 6:1 **6**, 1–13.

Yang D, Liu Y, Ali M, Ye L, Pan C, Li M, Zhao X, Yu F, Zhao X, Lu G. 2022. Phytochrome interacting factor 3 regulates pollen mitotic division through auxin signalling and sugar metabolism pathways in tomato. New Phytologist **234**, 560–577.

Yin W, Mendoza L, Monzon-Sandoval J, Urrutia AO, Gutierrez H. 2021. Emergence of co-expression in gene regulatory networks. PLOS ONE **16**, e0247671.

Yue J, Xu W, Ban R, Huang S, Miao M, Tang X, Liu G, Liu Y. 2016. PTIR: Predicted Tomato Interactome Resource. Scientific Reports **6**, 25047.

Zhang Y, Liu T, Meyer CA, *et al.* 2008. Model-based analysis of ChIP-Seq (MACS). Genome Biology **9**, 1–9.

Zhao X, Yuan X, Chen S, Meng L, Fu D. 2018. Role of the tomato TAGL1 gene in regulating fruit metabolites elucidated using RNA sequence and metabolomics analyses. PLOS ONE **13**, e0199083.

Zheng Y, Jiao C, Sun H, *et al.* 2016. iTAK: A Program for Genome-wide
Prediction and Classification of Plant Transcription Factors, Transcriptional
Regulators, and Protein Kinases. Molecular Plant **9**, 1667–1670.
Zhong S, Fei Z, Chen YR, *et al.* 2013. Single-base resolution methylomes of
tomato fruit development reveal epigenome modifications associated with
ripening. Nature Biotechnology 2013 31:2 **31**, 154–159.
Zhu F, Wen W, Cheng Y, Fernie AR. 2022. The metabolic changes that effect
fruit quality during tomato fruit ripening. Molecular Horticulture 2022 2:1 **2**, 1–19.
**Zhu Y, Zhu G, Xu R, Jiao Z, Yang J, Lin T, Wang Z, Huang S, Chong L, Zhu
JK**. 2023. A natural promoter variation of SlBBX31 confers enhanced cold
tolerance during tomato domestication. Plant Biotechnology Journal **21**, 1033–
1043.
**Zouine M, Maza E, Djari A, Lauvernier M, Frasse P, Smouni A, Pirrello J,
Bouzayen M**. 2017. TomExpress, a unified tomato RNA-Seq platform for
visualization of expression data, clustering and correlation networks. The Plant
Journal **92**, 727–735.

1525
1526

Tables

Table 1. Enrichment metrics for organ-Level GRNs. Summary of enrichment
metrics for organ-specific GRNs, considering the top 2% of interactions identified
by the GENIE3 algorithm. Enrichment results obtained from a Fisher's exact test
that assessed gene set overlap significance, reporting log2 fold change, p-value,
and intersection size to the validation network (ChIp-seq). -Inf represents $\log_{10}$
adjusted p-values < -400.

1534

	Root	Leaf	Flower	Fruit	Seed
Total TFs	1,297	1,216	1,300	1,058	1,241
Total Genes	23,226	22,513	23,988	20,888	23,124
Total edges	797,120	743,902	798,851	665,733	817,504
$\log_2$ Fisher odd ratio	1.89	2.66	2.01	3.01	2.22
$\log_{10}$ adj p-value	-156.72	-inf	-196.33	-305.17	-inf
Genes in Overlap	1714	2967	1912	2226	4204
AUROC	0.72	0.65	0.65	0.72	0.53
AUPR	0.51	0.45	0.46	0.52	0.32

1535

Figure Legends

Fig. 1. Organ-Level transcriptomic landscape in tomato. Organ-Level transcriptomic landscape in tomato. (A) Distribution of expressed genes across organs. (B) Distribution of expressed transcription factors (TFs) across organs. (C) Heatmap of normalized (Z-scored) expression levels of shared genes across organs. (D) Heatmap of normalized (Z-scored) expression levels of shared TFs across organs. (E) Enriched biological process Gene Ontology (GO) terms associated with expressed genes across organs (FDR-adjusted p-value < 0.05).

Fig. 2. Multi-omics data integration for generating organ-level GRNs in tomato. Overview of datasets used to generate organ-specific gene regulatory networks (GRNs) in *Solanum lycopersicum*. Bar plots indicate the number of available datasets per organ for transcriptomics (RNA-seq), chromatin accessibility (ATAC-seq/DNase-seq), and transcription factor binding sites (ChIP-seq). Arrows illustrate data flow into regulatory network generation, including GRNs, co-expression networks (GCNs), TF-binding networks, and validation datasets.

Fig. 3. Comparative analysis of organ-level GRNs reveals regulatory signatures and TF connectivity patterns. (A) Stacked bar plot showing the proportion of TF-target pairs in the GENIE3-inferred GRNs supported by multiple evidence sources. Darker purple shades indicate interactions validated by GENIE3 alone (G3) or by 2–4 sources, including open chromatin site binding (OCS), gene co-expression networks (Coex), and promoter binding (Promo). (B) UpSet plot displaying the overlap of TF-target interactions across organ-specific GRNs, with an inset showing the distribution of shared versus unique interactions (1 to 5 organs). (C) Relationship between TF mean connectivity (average number of target genes across organs) and target conservation (proportion of shared targets across organs). Colored dots represent TFs with highest and lowest values. A black trend line highlights the general pattern in the data.

Fig. 4. Identification of key transcriptional regulators of fruit ripening regulatory cascades in tomato. (A–B) Enrichment and validation of fruit GRN for RIN (A) and TAGL1 (B) using knockout mutant data and ChIP-binding analyses. Box heatmaps display enrichment obtained from a Fisher's exact test

(log2 fold change, p-value, and intersection size) to the validation network (ChIp-
binding). (C–D) Network representation of ripening-associated genes (Li et al.,
2019; Zhu et al., 2022) for RIN (C) and TAGL1 (D) derived from the fruit GRN.
Triangles represent TFs, squares represent target genes. Node colors indicate
function. Edges are colored in red (RIN) or blue (TAGL1) when interactions are
validated by ChIP-binding evidence. (E) Network of key regulators of ripening-
associated genes (Top scored TFs), applying the same node and edge color
scheme as in (C–D). Edge darker shades indicate accumulated regulatory
evidence.

1578

Fig. 5. Tomato GRNs reveal the role of ABF TFs and identify a key regulator
of ABA-related GRNs. (A) Bar plots showing organ-specific expression levels
and the number of ABA-related target genes regulated by ABF TFs. Bars indicate
target counts, while black lines represent $\log_{10}$ TPM expression values. (B)
Enrichment and validation of leaf GRNs for *ABF3* and *ABF5* using drought-
responsive DEGs (Gao et al., 2019). Box heatmaps (left) display enrichment
results from a Fisher's exact test (log2 fold change, p-value, and intersection
size). The networks (right) show distribution of ABA-related and drought
regulated targets of these TFs. (C) Gene Set Enrichment Analysis (GSEA, FDR-
adjusted p-value < 0.05) of *SlGBF3* target genes in organ-specific GRNs. Dot
size represents gene number, while color intensity reflects enrichment values. (D)
Network visualization of *SlGBF3*-regulated ABA-related genes in the leaf GRN.
Triangles represent TFs, squares represent target genes. Node colors indicate
function. Orange-bordered nodes indicate DAP-seq validated targets, while
enclosed nodes are DEGs from drought-stressed leaves (Gao et al., 2019). Edge
darker shades indicate accumulated regulatory evidence, (GENIE3: G3).

1595

Fig. 6. TomViz-GRNs: A web-based platform for exploring tomato organ-
level GRNs. (A) TomViz interface within the PlantaeViz platform (Santiago et al.,
2024), providing access to GRN exploration tools. (B) Regulatory Targets Tab:
Users can query TFs or genes to explore regulatory interactions and validation
layers. The interface includes options to download TF target lists, perform GSEA,
and visualize target distributions on a chromosome map. (C) D3 Subnetwork Tab:
Users can upload gene lists, visualize GRNs, and analyze regulatory pathways

at the organ level. The visualization includes directional edges representing regulatory interactions from TFs to targets, with edge colors indicating the level of supporting evidence. Additional options allow customization of network layout and node separation.

Fig. S1. Distribution of tomato transcriptomic datasets across organs and experimental conditions. (A) Stacked bar plot showing the number of transcriptomic libraries classified by organ of origin (Seed, Fruit, Flower, Leaf, Root). (B) Stacked bar plot showing the distribution of transcriptomic libraries categorized by treatment or experimental condition (Abiotic stress, Biotic interactions, Developmental, Genetic modifications, Undetermined). The legend indicates the total number of studies per category.

Fig. S2. Heatmap of Gene Set Enrichment Analysis (GSEA) results for organ-specific gene expression (FDR-adjusted p-value < 0.05). Color intensity reflects the enrichment values.

Fig. S3. Accuracy analysis of organ-specific GRNs. Receiver Operating Characteristic (ROC) curves (A) and Precision-Recall (PR) curves (B) comparing organ-specific GRNs to ChIP-seq validation networks. The shaded areas indicate variability across multiple iterations. Black and grey lines represent the maximum and minimum quartiles of randomly generated TF-target pairings, respectively.

Fig. S4. GSEA results (FDR-adjusted p-value < 0.05) for the five highest connected TFs shared across organ-level GRNs (conserved targets). Dot size represents gene number, while color intensity reflects enrichment values per TF.

Fig. S5. GSEA results (FDR-adjusted p-value < 0.05) for the five lowest connected TFs shared across organ-level GRNs (specific targets). Dot size represents gene number, while color intensity reflects enrichment values per TF.

Fig. S6. Genome-wide identification of *Sl*GBF3 binding targets using DAP-seq. (A) Venn diagram showing the overlap of DAP-seq peaks associated genes

between two replicates (2). (B) *De novo* binding motif identified from *Sl*GBF3 DAP-seq peaks using the MEME suite (Machanick and Bailey, 2011). (C) Comparison of binding motifs from *At*GBF3 orthologs (MA1351.1, MA1351.2, MA1351.3). (D) Pearson correlation heatmap comparing the enrichment of *Sl*GBF3 binding motif (M_*Sl*GBF3) with the *At*GBF3 orthologs.

Fig. S7. Enrichment analysis of *Sl*GBF3 targets in organ-specific GRNs compared to DAP-seq binding targets. Box heatmap display enrichment results from a Fisher's exact test (log2 fold change, p-value, and intersection size).

Fig. S8. GSEA results (FDR-adjusted p-value < 0.05) of *Sl*GBF3 DAP-seq targets. Dot size represents gene number, while color intensity reflects enrichment values.

Fig. S9. Network visualization of *Sl*GBF3-regulated abscisic acid (ABA)-related genes within Root, Seed, Flower and Fruit GRNs. Triangles represent transcription factors, and squares represent genes. Node colors indicate function. Orange-bordered nodes highlight DAP-seq validated genes. Edge color shades represent accumulated regulatory evidence (G3: GENIE3).

Fig. S5. GSEA results (FDR-adjusted p-value < 0.05) for the five lowest connected TFs shared across organ-level GRNs (specific targets). Dot size represents gene number, while color intensity reflects enrichment values per TF.

Fig. S6. Genome-wide identification of SlGBF3 binding targets using DAP-seq. (A) Venn diagram showing the overlap of DAP-seq peaks associated genes between two replicates (2). (B) De novo binding motif identified from SlGBF3 DAP-seq peaks using the MEME suite (Machanick and Bailey, 2011). (C) Comparison of binding motifs from AtGBF3 orthologs (MA1351.1, MA1351.2, MA1351.3). (D) Pearson correlation heatmap comparing the enrichment of SlGBF3 binding motif (M_SIGBF3) with the AtGBF3 orthologs.

Fig. S7. Enrichment analysis of *Sl*GBF3 targets in organ-specific GRNs compared to DAP-seq binding targets. Box heatmap display enrichment results from a Fisher's exact test (log2 fold change, p-value, and intersection size).

Fig. S8. GSEA results (FDR-adjusted p-value < 0.05) of *Sl*GBF3 validated targets.
Dot size represents gene number, while color intensity reflects enrichment values.

Fig. S9. Network visualization of SlGBF3-regulated abscisic acid (ABA)-related genes within Root, Seed, Flower and Fruit GRNs. Triangles represent transcription factors, and squares represent genes. Node colors indicate function. Orange-bordered nodes highlight DAP-seq validated genes. Edge color shades represent accumulated regulatory evidence (G3: GENIE3).

Fig. 1. Organ-Level transcriptomic landscape in tomato. (A) Distribution of expressed genes across organs. (B) Distribution of expressed transcription factors (TFs) across organs. (C) Heatmap of normalized (Z-scored) expression levels of shared genes across organs. (D) Heatmap of normalized (Z-scored) expression levels of shared TFs across organs. (E) Enriched biological process Gene Ontology (GO) terms associated with expressed genes across organs (FDR-adjusted p-value < 0.05).

Fig. 2. Multi-omics data integration for generating organ-level GRNs in tomato. Overview of datasets used to generate organ-specific gene regulatory networks (GRNs) in *Solanum lycopersicum.* Bar plots indicate the number of available datasets per organ for transcriptomics (RNA-seq), chromatin accessibility (ATAC-seq/DNase-seq), and transcription factor binding sites (ChIP-seq). Arrows illustrate data flow into regulatory network generation, including GRNs, co-expression networks (GCNs), TF-binding networks, and validation datasets.

Fig. 3. Comparative analysis of organ-level GRNs reveals regulatory signatures and TF connectivity patterns. (A) Stacked bar plot showing the proportion of TF-target pairs in the GENIE3-inferred GRNs supported by multiple evidence sources. Darker purple shades indicate interactions validated by GENIE3 alone (G3) or by 2–4 sources, including open chromatin sites binding (OCS), gene co-expression networks (Coex), and promoter binding (Promo). (B) UpSet plot displaying the overlap of TF-target interactions across organ-specific GRNs, with an inset showing the distribution of shared versus unique interactions (1 to 5 organs). (C) Relationship between TF mean connectivity (average number of target genes across organs) and target conservation (proportion of shared targets across organs). Colored dots represent TFs with highest and lowest values. A black trend line highlights the general pattern in the data.

Fig. 4. Identification of key transcriptional regulators of fruit ripening regulatory cascades in tomato. (A–B) Enrichment and validation of fruit GRN for RIN (A) and TAGL1 (B) using knockout mutant data and ChIP-binding analyses. Box heatmaps display enrichment obtained from a Fisher's exact test (log2 fold change, p-value, and intersection size) to the validation network (ChIp-binding). (C–D) Network representation of ripening-associated genes (Li et al., 2019; Zhu et al., 2022) for RIN (C) and TAGL1 (D) derived from the fruit GRN. Triangles represent TFs, squares represent target genes. Node colors indicate function. Edges are colored in red (RIN) or blue (TAGL1) when interactions are validated by ChIP-binding evidence. (E) Network of key regulators of ripening-associated genes (Top scored TFs), applying the same node and edge color scheme as in (C–D). Edge darker shades indicate accumulated regulatory evidence.

Fig. 5. Tomato GRNs reveal the role of ABF TFs and identify a key regulator of ABA-related GRNs. (A) Bar plots showing organ-specific expression levels and the number of ABA-related target genes regulated by ABF TFs. Bars indicate target counts, while black lines represent log10 TPM expression values. (B) Enrichment and validation of leaf GRNs for ABF3 and ABF5 using drought-responsive DEGs (Gao et al., 2019). Box heatmaps (left) display enrichment results from a Fisher's exact test (log2 fold change, p-value, and intersection size). The networks (right) show distribution of ABA-related and drought regulated targets of these TFs. (C) Gene Set Enrichment Analysis (GSEA, FDR-adjusted p-value < 0.05) of *Sl*GBF3 target genes in organ-specific GRNs. Dot size represents gene number, while color intensity reflects enrichment values. (D) Network visualization of *Sl*GBF3-regulated ABA-related genes in the leaf GRN. Triangles represent TFs, squares represent target genes. Node colors indicate function. Orange-bordered nodes indicate DAP-seq validated targets, while enclosed nodes are DEGs from drought-stressed leaves (Gao et al., 2019). Edge darker shades indicate accumulated regulatory evidence, (GENIE3: G3).

Fig. 6. TomViz-GRNs: A web-based platform for exploring tomato organ-level GRNs. (A) TomViz interface within the PlantaeViz platform (Santiago et al., 2024), providing access to GRN exploration tools. (B) Regulatory Targets Tab: Users can query TFs or genes to explore regulatory interactions and validation layers. The interface includes options to download TF target lists, perform GSEA, and visualize target distributions on a chromosome map. (C) D3 Subnetwork Tab: Users can upload gene lists, visualize GRNs, and analyze regulatory pathways at the organ level. The visualization includes directional edges representing regulatory interactions from TFs to targets, with edge colors indicating the level of supporting evidence. Additional options allow customization of network layout and node separation.

Fig. S1. Distribution of tomato transcriptomic datasets across organs and experimental conditions. (A) Stacked bar plot showing the number of transcriptomic libraries classified by organ of origin (Seed, Fruit, Flower, Leaf, Root). (B) Stacked bar plot showing the distribution of transcriptomic libraries categorized by treatment or experimental condition (Abiotic stress, Biotic interactions, Developmental, Genetic modifications, Undetermined). Each category total number of studies is indicated in the legend

Fig. S2. Heatmap of Gene Set Enrichment Analysis (GSEA) results for organ-specific gene expression (FDR-adjusted p-value < 0.05). Color intensity reflects the enrichment values.

Fig. S3. Accuracy analysis of organ-specific GRNs. Receiver Operating Characteristic (ROC) curves (A) and Precision-Recall (PR) curves (B) comparing organ-specific GRNs to ChIP-seq validation networks. The shaded areas indicate variability across multiple iterations. Black and grey lines represent the maximum and minimum quartiles of randomly generated TF-target pairings, respectively.

Fig. S4. GSEA results (FDR-adjusted p-value < 0.05) for the five highest connected TFs shared across organ-level GRNs (conserved targets). Dot size represents gene number, while color intensity reflects enrichment values per TF.