Neural sensitivity to the heartbeat is modulated by fluctuations in affective arousal during spontaneous thought

Abbreviated title: Interoceptive sensitivity and affective arousal

David Braun[1], Lotus Shareef-Trudeau[1], Swetha Rao[1], Christine Chesebrough[2], Julia W. Y. Kam[3], Aaron Kucyi[1]

[1]Department of Psychological & Brain Sciences, Drexel University, 19104

[2]Northwell Health Feinstein Institutes for Medical Research, 11030

[3]Department of Psychology, University of Calgary, T2N 1N4

Correspondence to:

Aaron Kucyi, Ph.D.
Dept. of Psychological & Brain Sciences, Drexel University
3201 Chestnut St.
Philadelphia, PA 19104
United States
Email: aaron.kucyi@drexel.edu

Number of pages: 51
Number of figures: 4 (in text) & 3 (supplementary)
Number of tables: 2 (supplementary)
Number of words in abstract: 249
Number of words in introduction: 632
Number of words in discussion: 1,476

Acknowledgments: This work was supported by the National Institute of Mental Health of the National Institutes of Health under award numbers R21MH129630 and R21MH127384 (to AK).[1]

---

[1] The authors declare no competing financial interests.

Abstract

Spontaneous thoughts, occupying much of one's awake time in daily life, are often colored by

emotional qualities. While spontaneous thoughts have been associated with various neural

correlates, the relationship between subjective qualities of ongoing experiences and the brain's

sensitivity to bodily signals (i.e., interoception) remains largely unexplored. Given the well-

established role of interoception in emotion, clarifying this relationship may elucidate how

processes relevant to mental health, such as arousal and anxiety, are regulated. We used EEG and

ECG to measure the heartbeat evoked potential (HEP), an index of interoceptive processing, while

51 adult participants (34 male, 20 female) visually fixated on a cross image and let their minds

wander freely. At pseudo-random intervals, participants reported their momentary level of arousal.

This measure of affective arousal was highly variable within and between individuals but was

statistically unrelated to several markers of physiological arousal, including heart rate, heart rate

variability, time on task, and EEG alpha power at posterior electrodes. A cluster-based permutation

analysis revealed that the HEP amplitude was increased during low relative to high affective

arousal in a set of frontal electrodes during the $340 - 356$ millisecond window after heartbeat onset.

This HEP effect was more pronounced in individuals who reported high, relative to low, levels of

trait anxiety. Together, our results offer novel evidence that at varying levels of trait anxiety, the

brain differentially modulates sensitivity to bodily signals in coordination with the momentary,

spontaneous experience of affective arousal—a mechanism that may operate independently of

physiological arousal.

Significance Statement

Our findings highlight the relationships between spontaneous fluctuations in affective arousal, brain-body interactions, and anxiety, offering new insights into how interoception fluctuates with changes in internal states. By showing that interoceptive processing is heightened during lower affective arousal and that this effect is amplified in individuals with higher trait anxiety, our study suggests the brain adaptively downregulates interoceptive sensitivity in response to fluctuating internal states. These results have implications for understanding how spontaneous thoughts shape interoception and emotion, particularly in clinical contexts where dysregulated interoception is linked to anxiety and mood disorders. More broadly, our work underscores the need to distinguish between different forms of arousal, advancing understanding of the taxonomy and ways of measuring arousal.

Introduction

The human brain continually processes signals from the body to guide perception and

behavior, a capacity referred to as *interoception* (Engelen et al., 2023). Although early theorists

(e.g., James, 1884/1948; Sherrington, 1906) noted its role in cognition and emotion, interoception

has recently become a critical focus in cognitive neuroscience (Azzalini et al., 2019; Kluger et al.,

2024). A method of measuring interoception—first introduced in the late 20[th] century(Montoya et

al., 1993; Schandry et al., 1986) and now commonly applied with refined approaches (Park &

Blanke, 2019)—is the *heartbeat-evoked potential (HEP)*, which involves averaging neural

electrophysiological signals time-locked to features of the electrocardiogram (*ECG*) representing

the neural processing of heartbeats. Larger HEP amplitude is often interpreted as greater

interoception (Judah et al., 2018; Pang et al., 2019; Verdonk et al., 2024), and the HEP has been

linked to affective arousal (Fourcade et al., 2024), a factor central to the present investigation.

Arousal is historically defined as physiological and psychological activation or intensity

(Russell, 1980; Russell & Barrett, 1999). Recently, researchers have debated whether arousal is

best characterized as unidimensional or multidimensional (Sabat et al., 2025), for example

separating affective and physiological aspects as independent subcomponents (Reid et al., 2024;

Satpute et al., 2019). Differences in the HEP have been found in frontal electrodes in the 200–450

ms interval post-heartbeat across various conditions that broadly targeted arousal, including

exposure to emotional stimuli (Marshall et al., 2017, 2018, 2020; Sel et al., 2017), food deprivation

(Schulz et al., 2015), pain (Shao et al., 2011), and cortisol injections (Schulz et al., 2013). Notably,

stimulus- and task-evoked arousal has previously been associated with both increased (Luft &

Bhattacharya, 2015) and diminished (Fourcade et al., 2024) interoception.

Another way to target arousal is to measure natural, spontaneous fluctuations during

wakeful rest (Chang et al., 2016; Kucyi & Parvizi, 2020), an approach that is also increasingly

used to investigate the neural basis of ongoing mental experience (e.g., "spontaneous thoughts" or

"mind-wandering"; Christoff et al., 2016; Kam et al., 2021; Shareef-Trudeau et al., In press;

Smallwood et al., 2018). For example, HEP amplitudes vary depending on spontaneous

fluctuations in self- or other-referenced thoughts (Babo-Rebelo et al., 2019; Babo-Rebelo, Richter,

et al., 2016; Babo-Rebelo, Wolpert, et al., 2016). However, no study to our knowledge has linked

the HEP to spontaneous fluctuations in affective arousal. Spontaneous mental experiences are

pervasive in daily life (Killingsworth & Gilbert, 2010), and the arousal component of experience

is particularly relevant to mental health conditions involving anxiety.

Anxiety can involve dysregulated interoception, such as heightened or inconsistent

sensitivity to bodily signals (Khalsa & Lapidus, 2016). Individuals scoring higher in anxiety tend

to exhibit larger HEP amplitudes (Judah et al., 2018; Pang et al., 2019; Verdonk et al., 2024). Given

the intimate associations between (i) anxiety and interoception (Janelle, 2002; Noteboom et al.,

2001; Roos et al., 2021) and (ii) anxiety and abnormal spontaneous thinking (Fell et al., 2023),

spontaneous fluctuations in affective arousal and their interoceptive underpinnings may prove

important for understanding anxiety.

Here we examined the relationship between subjective qualities of ongoing experiences

and interoception as well as how this relationship might be influenced by anxiety. Participants

completed anxiety surveys and underwent EEG and ECG recordings during a wakeful resting state,

intermittently rating affective arousal via experience sampling. To better understand our "affective

arousal" construct, we assessed whether self-reported arousal converged with physiological

measures or was linked to other aspects of ongoing experience. We next leveraged HEP analysis

to examine whether affective arousal is associated with higher or lower levels of interoceptive

processing. Finally, we explored whether individual variability in trait anxiety modulated the

relationship between HEP amplitude and affective arousal. Our findings offer important

implications for how arousal is conceptualized and how spontaneous internal fluctuations shape

the link between interoception and emotion.

Materials and Methods

Participants

Fifty-four participants (34 female, 20 male) were recruited to participate in this study.

Participants were between the ages of 18 and 31, and 50.00% identified as Asian, 35.19% identified

as White, 11.11% identified as Black or African American, 1.85% identified as two or more races,

and 1.85% identified as other. Participants were recruited both from undergraduate psychology

courses for optional course credit and from recruitment posters placed around Drexel and Temple

University campuses. Those recruited through the posters received $60 USD of compensation for

participation. All participants were fluent in English, had normal or corrected-to-normal vision,

were right-handed, and reported no history or current diagnosis of any neurological or psychiatric

disorders. All participants provided informed consent for procedures approved by the Drexel

University Institutional Review Board. One participant was excluded due to removal of an

excessive number of noisy epochs in EEG recordings, one participant was excluded due to EEG

recording failure, and one participant was excluded due to behavioral data recording failure,

leaving 51 participants for the final analysis.

EEG Data Acquisition

EEG recordings were obtained using a 32-channel Brain Products system, including the

BrainAmp MRPlus and the BrainAmp ExG (*BrainCap MR*, 2020). We recorded EEG data with

the BrainVision Recorder Vers. 1.25.0201 on a Windows operating system using a Dell Latitude

7430 64-bit operating system with a 12[th] Gen Intel Core i7-1265U and 16GB of RAM. Data were

continuously sampled at 5000 Hz. We fitted participants using the BrainCap MR (based on the

extended 10/20 system) with embedded carbon wire loops. The system includes ground and

reference electrodes, each with 15 kOhms of resistance, 31 scalp electrodes with 10 kOhms of resistance each and an ECG channel with 20 kOhms of resistance. We recorded heartbeat activity via the ECG which was applied to the center of each participant's back as far down as the cord could reach, typically mid-back (Yamashita et al., 2024). Cap setup was concluded when all electrodes' impedance measurements were at levels below 20 kOhms.

Mental Health Surveys

Prior to EEG setup, participants filled out a series of surveys pertaining to their demographic (e.g., age, sex, race) and mental health status. Of these, only the Generalized Anxiety Disorder-7 (GAD-7; Spitzer et al., 2006) and the state scales of the State Trait Anxiety Inventory (STAI-S; Speilberger et al., 1983) were analyzed to capture both trait and state anxiety, respectively, given our focus on the links of anxiety with arousal and interoception (for the complete set of surveys administered in the experiment, see *Table S1*). The state anxiety measure (STAI-S) was a late addition to the experiment after 23 participants' data had been collected. All surveys, except for the STAI-S, were administered online using REDCap electronic data capture tools (Harris et al., 2009, 2019). STAI-S was collected either online or with paper and pencil. Participants were asked to complete all but the state anxiety (i.e., STAI-S) measure prior to arriving for the experiment. Both of the GAD-7 and STAI-S scales have shown high validity and reliability across a wide range of contexts and populations (Gunning et al., 2010; Löwe et al., 2008; Pretorius & Padmanabhanunni, 2023; Vitasari et al., 2011). The GAD-7 and STAI-S are shown to have high internal consistency, both in past research (Cronbach's $\alpha_{GAD} = 0.89$, Löwe et al., 2008; $\alpha_{STAI} = 0.87$, Spielberger et al., 2017) and in our sample ($\alpha_{GAD} = 0.91$, $\alpha_{STAI} = 0.89$).

Experience Sampling

Throughout the EEG/ECG experiment, participants were intermittently prompted to report

on the nature of their ongoing experience using thought probes. Each probe consisted of 13

experience sampling items presented sequentially in a fixed order. For each item, participants

provided a rating on a 0–100 slider scale assessing a different dimension of their experience in the

moments immediately preceding the probe's onset. The primary experience sampling item relevant

to this study measured affective arousal, asking: *"How activated or energized were you feeling?"*

The highest end of the anchor was labeled *"Completely activated"*, while the lowest end was

labeled *"Completely deactivated."* Affective arousal was defined to participants as *"feelings of*

energy linked to emotion, which may result in heightened bodily energy, tension, or

responsiveness", a definition which is consistent with prior research (Barrett, 1998; Bliss-Moreau

et al., 2020; Haj-Ali et al., 2020; Kron et al., 2015; Kuppens et al., 2013; Maher et al., 2019; Nook

et al., 2017; Russell, 1980, 2003; Russell & Barrett, 1999). Other relevant experience sampling

items for present purposes included probes about future thinking ("Were your thoughts oriented

towards the future?" from "Not future oriented" to "Completely future oriented"), deliberate

thinking ("How intentional were your thoughts?" from "Completely unintentional" to "Completely

intentional"), self-related thinking ("Were your thoughts about yourself?" from "Nothing about

you"   to   "Completely   about   you

"), and difficulty to disengage from thinking ("How difficult was it to disengage from your

thoughts?" from "Extremely easy" to "Extremely difficult"). Participants additionally rated their

level of confidence in their responses at the end of each thought probe (from "Completely

confident" to "Completely unconfident"). For the complete set of experience sampling items used

in the experiment, see *Table S2*.

Procedure

Upon arrival, participants provided informed consent and were fitted with an EEG cap

while receiving task instructions. They were introduced to each experience sampling item and

tested for comprehension using a hypothetical scenario in which a student was experiencing stress

while preparing for an exam, and participants were asked to respond to experience sampling items

as if they were the hypothetical student studying for the exam. The experimenter provided

feedback to ensure participant comprehension of instructions before proceeding, and evidence of

comprehension was supported by participants giving generally high ratings of confidence in their

responses throughout the experiment ($M = 17.89$, where 0 is most confident and 100 is least; SD

$= 14.41$).

The main task consisted of experience sampling during wakeful rest (see *Figure 1A*).

During wakeful rest, participants sat at a computer and were instructed to "Let your mind wander

freely" while focusing on a black screen with a central white fixation cross. At a randomized

integer interval every 15–45 s, the fixation cross was replaced by a thought probe with a series of

experience sampling items, which participants answered using arrow keys on a laptop computer

to adjust a slider scale and entered their chosen response using the spacebar. After the last item,

the fixation cross reappeared, and the cycle repeated.

Prior to the wakeful rest and experience sampling task, participants first alternated between

opening and closing their eyes for eight minutes to validate that EEG was recording as expected.

Participants next performed the main phase of the experiment, where each cycle of wakeful rest

and experience sampling repeated 10 times per run, with 5 total runs, yielding 50 total trials (i.e.,

thought probe sets). Prior to the start of each task run, the experimenter exited the room. The

experimenter reentered the room between each run to give the participants a short break and cue

the next task/run. Data collection lasted approximately 60 minutes, with the full session taking about 2 hours.

The first 22 participants were tested in a windowless lab room, which was equipped with a stimulus laptop for task administration and an EEG recording laptop. During the main task, the experimenter visually monitored participants remotely via Zoom (*Zoom Video Communications*, 2024) using a separate monitoring computer. Due to equipment and facility constraints, the remaining 31 participants were tested in a different room that included a glass wall. Participants were seated with their backs to the glass wall and, to minimize external distractions during the task, participants wore earplugs.

Figure 1. Trial sequence and EEG/ECG analysis approach. (A) Participants were instructed to "Let your mind wander freely" while focusing on a black screen with a central, white fixation cross. EEG and ECG were simultaneously recorded throughout. After 15 – 45 s, a series of thought probes appeared, prompting participants for ratings about their experience. This trial sequence was repeated 50 times throughout the experiment. (B) Pre-probe epochs were first defined in the

window 10 s prior to thought probe onset. Within pre-probe epochs, heartbeat epochs were defined in the EEG data from -0.1 – 0.65 s around each R-peak in the ECG data. We then conducted trial-wise regressions, using EEG voltage at each timepoint and channel within the heartbeat-locked epochs to predict affective arousal ratings. This yielded a timepoint-by-channel map of regression slopes for each participant, which we submitted to a cluster-based permutation analysis using one-sample t-tests against zero to identify significant clusters.

EEG Preprocessing

Electrophysiological data preprocessing was performed using custom Python scripts based on the MNE library (Gramfort et al., 2013; Larson et al., 2023). Data were downsampled to 250 Hz, bandpass filtered between 1 and 55 Hz, and referenced to the common average. Initial artifact removal was conducted using Extended Infomax independent components analysis (ICA; Lee et al., 1999), with automatic classification and rejection of non-brain artifacts via ICLabel (Pion-Tonachini et al., 2019). Noisy channels were identified via manual inspection and interpolated on the pre-ICA data. Five participants had one channel interpolated, two participants had two channels interpolated, and two participants had three or four channels interpolated. A second ICA decomposition was then performed, and ICLabel results were manually reviewed for further artifact rejection. We chose not to perform baseline correction for HEP analyses because, unlike in stimulus-evoked designs, heartbeat-evoked designs lack a true baseline. Cardiac activity is cyclical, and electrical activity before a heartbeat can reflect both the preceding heartbeat and preparatory processes for the upcoming one (Azzalini et al., 2019).

For HEP analyses, R-peaks in the ECG signal were detected using MNE's preprocessing library. We defined the 10 s period prior to the onset of the thought probe as the time period of interest, as brain states linked to self-report probes can emerge up to 10 s before probe onset (Christoff et al., 2009; Kucyi et al., 2024). Heartbeats occurring within 10 s prior to each thought probe onset were retained for analysis. Following prior work (Petzschner et al., 2019), heartbeats

within 700 ms of each other were excluded to prevent overlapping epochs. Within this 10 s pre-probe period, epochs were defined on the EEG channels from -100 ms to +650 ms relative to each R-peak. Noisy epochs were automatically identified with the autoreject library (Jas et al., 2017), and all epochs were manually inspected before removal.

The cardiac field artifact (CFA) is noise introduced into the EEG signal due to electrical activity from the heart. While ICA is a powerful method for removing many types of artifacts, including the CFA, from EEG (Debener et al., 2010; Devuyst et al., 2008; Hoffmann & Falkenstein, 2008), no known method can entirely remove the CFA from EEG data. To ensure the CFA was not impacting the EEG signal differently across experimental conditions, we conducted several control analyses. First, we examined the association between ECG metrics—including peak and mean ECG amplitude—and affective arousal (Petzschner et al., 2019). This association was examined by first averaging both ECG metrics in the significant time window revealed by the cluster-based permutation on the EEG data (see below), and then separately regressing both averaged ECG metrics on to affective arousal ratings across trials, using Bayesian estimation to provide Bayes factors in support of the null hypothesis. Second, we applied the same cluster-based permutation analysis used in our EEG analysis (see below) directly to the ECG data. Any statistically significant associations between ECG amplitude and affective arousal would raise concerns about a potential CFA-related confound.

For spectral analyses involving parieto-occipital alpha power, epochs were defined over the full 10 s period prior to thought probe onset. Noisy epochs were again removed by manually inspecting results from the autoreject library. We examined parieto-occipital alpha power in the 10 s pre-probe period, as past studies have shown a connection between parieto-occipital alpha power and various types of arousal (Hofmann et al., 2021; Lozano-Soldevilla, 2018; Schubring &

Schupp, 2021; Tuladhar et al., 2007). To examine parieto-occipital power, the EEG signal was decomposed into time-frequency dimensions via MNE's Morlet wavelet transform algorithm with 50 logarithmically spaced frequencies from 2 to 30 Hz and a variable number of cycles per wavelet (n_{cycles} = freq/2) for occipital and parietal channels only, namely O1, O2, Oz, P3, P4, P7, P8, and Pz (Fourcade et al., 2024). We decomposed the signal from 2-30 Hz to allow for exploratory flexibility in examining power across multiple frequency bands, though our primary focus in this study is on parieto-occipital alpha power.

Experimental Design and Statistical Analysis

Affective Arousal, Physiological Arousal, and Other Experience Sampling Measures

To assess the relationship between physiological and affective arousal, measures of the two constructs were compared. Measures of physiological arousal included heart rate, heart rate variability (HRV), time-on-task, and parieto-occipital alpha power amplitude (Hofmann et al., 2021; Lozano-Soldevilla, 2018; Schubring & Schupp, 2021; Tuladhar et al., 2007), each of which except for time-on-task were summarized over all heartbeats in the 10 s period prior to probe onset (i.e., not removing heartbeats occurring within 700 ms of each other, a procedure only done for the HEP analyses). Time-on-task, calculated as the probe (i.e., trial) order number, was included because of its well established links to self-reported fatigue, impaired task performance, and negative association with physiological arousal measures (Gergelyfi et al., 2015; Herlambang et al., 2019; Karthikeyan et al., 2021, 2022; Matuz et al., 2019; Melo et al., 2017; Qin et al., 2021; Shi et al., 2022; Zhao et al., 2012). HRV was calculated using the root mean square of differences between successive heartbeats (Shaffer & Ginsberg, 2017). Affective arousal was measured by self-reported levels of arousal obtained at each thought probe.

We used hierarchical Bayesian linear regressions to estimate relationships between

physiological and affective measures of arousal. These regressions were implemented via the brms

(Bürkner, 2017) and Stan (Stan Development Team, 2024) packages in R (R Core Team, 2024)

using default, weakly-informative priors. Hierarchical Bayesian estimation is ideal for accurately

estimating trends at both the group and individual level. Separate models were fit for each

physiological arousal measure—heart rate, HRV, time on task, and alpha power—with the

physiological arousal measure predicting affective arousal. Random intercepts and slopes were

included into the model, grouped by participant. To aid estimation, affective arousal was z-scored

within participants and alpha power was log-transformed prior to model sampling. Estimation of

each model used four chains, each drawing 5,000 samples (including 2,500 warm-up).

Convergence was assessed by inspecting $\hat{R}$ values, ensuring all values were ≤ 1.01. All models

showed adequate effective sample sizes and no convergence warnings. We summarized both

group- and subject-level slope estimates using posterior means and 95% credible intervals.

Intervals excluding zero were interpreted as evidence of a nonzero relationship. These posterior

summaries were complemented by significance estimates from traditional (i.e., frequentist)

hierarchical models estimated using *lme4* (Bates et al., 2015) using the Satterthwaite degrees of

freedom approximation from the *lmerTest* package (Kuznetsova et al., 2017) to obtain *p* values.

We used the same modeling methods to assess the degree to which ratings of affective arousal

were related to other self-reported experience sampling measures.

Cardiac Interoceptive Sensitivity and Affective Arousal

Cardiac interoceptive sensitivity was analyzed in relation to affective arousal by first

averaging heartbeat epochs at the trial level within participants. Time points in the epochs less than

250 ms post-R peak were excluded from statistical analysis to avoid overlap with the CFA (Al et

al., 2021; Fourcade et al., 2024; Tanaka et al., 2023). Moreover, much previous research using both scalp and intracranial EEG have found HEP effects prior to 450 ms post heartbeat (Al et al., 2020, 2021; Babo-Rebelo, Wolpert, et al., 2016; Canales-Johnson et al., 2015; Flasbeck et al., 2024; Fukushima et al., 2011; Kern et al., 2013; Park et al., 2018; Yuan et al., 2008), including those studies focused specifically on arousal (Coll et al., 2021; Fourcade et al., 2024). Thus, in line with prior work, we chose to restrict our statistical analysis to the 250 – 450 ms window post heartbeat. For each participant, we performed a regression at every channel and epoch timepoint, using affective arousal ratings to predict EEG voltage across trials. The resulting slope coefficients were compiled across participants, yielding a time (250–450 ms) by channel (31 EEG electrodes) array of slopes per participant. We then submitted these slope arrays to a group-level cluster-based permutation test, with the statistical test being a one-sample t-test against zero, identifying clusters where EEG voltage showed a consistent linear relationship with arousal ratings across participants. The cluster-based permutation test used a $p = .01$ cluster forming threshold, 5,000 simulations, and evaluated two-tailed significance.

Validating Heartbeat Evoked Effects. To validate that effects observed in cluster-based permutation analyses were indeed heartbeat-evoked—and not driven by external events or spontaneous drift unrelated to heartbeats—we conducted a *surrogate heartbeat control analysis* (Azzalini et al., 2019; Babo-Rebelo, Wolpert, et al., 2016; Park et al., 2016; Park & Blanke, 2019; Steinfath et al., 2024). This surrogate analysis compares the observed cluster effect to a null distribution generated by disrupting the time-locking between heartbeats and EEG epochs. The test statistic for the surrogate analysis was derived by performing a participant-specific median-split version of the cluster-based permutation test described above (as the nature of repeated simulations required for the surrogate analysis rendered the regression method computationally

infeasible). Trial-level epochs were further averaged to a within-participants median split on affective arousal ratings, generating a time (250–450 ms) by channel (31 EEG electrodes) array of EEG voltages for each affective arousal condition (i.e., high and low) for each participant. The statistical test was a paired t test, comparing these arrays across conditions and participants. Obtaining the test statistic for the surrogate test involved summing these t values from the significant cluster. To generate the null distribution, we simulated surrogate heartbeats by randomly sampling k timepoints—where k is equal to the number of true heartbeats—within each 10 s pre-probe period, continuing to sample if any surrogate heartbeat times were in the set of true heartbeat times. We constructed EEG epochs (-100 to +650 ms relative to each surrogate timepoint) and applied the autoreject library (Jas et al., 2017) to identify and remove noisy epochs. Autoreject was fit separately for each participant across all epochs, using its automated thresholding and sensor interpolation methods to optimize artifact rejection without manual intervention. After cleaning, the remaining epochs were processed using the same median-split pipeline as in the main analysis, resulting in a time (250–450 ms) by channel (31 electrodes) array for each condition and participant. We then submitted these arrays to a cluster-based permutation analysis identical to the original, summed the t values within each cluster, identified the cluster with the largest absolute value t sum, and recorded the original (signed) sum for that cluster as one observation in the null distribution. This process was repeated 100 times, yielding an empirical null distribution of cluster statistics. The original test statistic was evaluated for two-tailed significance by comparing its percentile within this null distribution (i.e., whether it exceeded the 97.5th or fell below the 2.5th percentile).

Cardiac Interoceptive Sensitivity, Affective Arousal, and Anxiety

To examine whether individual differences in anxiety modulates the link between cardiac interoceptive sensitivity and affective arousal, the magnitude of HEP effects was correlated with anxiety scale scores. To determine HEP effect magnitude for each participant, we averaged the regression slopes in a significant time by channels cluster. This averaged slope was correlated separately with GAD (trait anxiety) and STAI-S (state anxiety) composite scores.

For t tests and non-hierarchical regressions reported in the results, we report Bayes factors characterizing evidence either in favor of or against the null hypothesis using the *BayesFactor* package (Morey et al., 2011; Morey & Rouder, 2011; Rouder et al., 2009) in the *R* programming language (R Core Team, 2024). We use the conventional notation of BF_{01} to denote evidence in favor of the null hypothesis and BF_{10} to denote evidence in favor of the alternative hypothesis. We used default prior settings in all applications, which are a noninformative Jeffrey's prior placed on the variance of the normal population, and a Cauchy prior placed on the standardized effect size. We interpret Bayes factors using canonical thresholds: < 3 as anecdotal evidence, > 3 & < 10 as substantial evidence, > 10 & < 31.6 as strong evidence, > 31.6 & < 100 as very strong evidence, and > 100 as decisive evidence (Jeffreys, 1961/1998). Note that for hierarchical Bayesian models, we do not report Bayes factors, as estimating marginal likelihoods necessary for these Bayes factors is notoriously difficult. Estimation methods such as bridge sampling, while available, are known to be highly sensitive to prior specification and model complexity (Gronau et al., 2017, 2020).

Results

Sufficient trial-level data and no modulation of cardiac field artifact by affective arousal

Given that the HEP is relative to the onset of a heartbeat and not to an external stimulus,

enough time needs to elapse between heartbeats to cleanly measure the HEP (Petzschner et al.,

2019). To prevent overlapping heartbeat events, we dropped detected R peaks that were within 700

ms of each other, resulting in an average dropout rate of 11.48% (SD = 14.28%) per participant.

An average of 9.76 epochs per trial (i.e., 10-second pre-thought probe interval) (SE = 0.20) were

retained per participant, and there was strong evidence that the number of epochs retained per trial

was not associated with affective arousal ratings ($b = 0.001, CIs = [-0.002, 0.002]$, $t(2355) =$

0.43, $p = .666$, $BF_{01} = 19.68$). These results suggest that a sufficient number of epochs were

being averaged per trial.

To control for the CFA, we sought to ensure that peak and mean ECG amplitude were not

associated with affective arousal ratings. Comparing peak ECG amplitude within the significant

time window (revealed by the cluster-based permutation analysis, 340 – 356 ms post R-peak; see

below) also revealed strong evidence in favor of no association between peak ECG amplitude and

affective   arousal   ratings   ($b = -1.12 \times 10^{-8}, 95\ CIs = [-1.08 \times 10^{-7}, 8.50 \times$

$10^{-8}]$, $t(2355) = 0.236$, $p = .819$, $BF_{01} = 21.03$). Additionally, there was strong evidence in

favor of no association between difference in mean ECG amplitude and affective arousal ratings

($b = -5.43 \times 10^{-9}, 95\ CIs = [-9.82 \times 10^{-8}, 8.74 \times 10^{-8}]$, $t(2355) = 0.115$, $p =$

.909, $BF_{01} = 21.44$). Further, a cluster-based permutation analysis of ECG signals, using the

same parameters as the EEG analysis (see below), failed to find any significant clusters. These

results suggest that the CFA was well controlled and not modulated by affective arousal. Finally,

a visual inspection revealed that the averaged ECG waveform was nearly identical across a

participant-specific median split in affective arousal ratings (*Figure S1*).

Affective arousal is distinct from physiological arousal and other experience sampling

measures

We next examined how affective arousal fluctuates within and between participants and

how it relates to other measures collected in the experiment. Affective arousal, rated on a 100-

point scale, varied substantially between participants ($M = 49.1$, $SD = 12.4$), with some participants

exhibiting a bimodal pattern of responding while others exhibited a unimodal, normal-shaped

pattern (see *Figure 2A and Figure S2*). For some participants, affective arousal varied widely over

time, whereas others displayed more stable patterns (see *Figure* 2B).

We next examined the relationship between subjective and physiological arousal measures

(see top row of *Figure 2C*). Group-level 95% credible intervals included zero for all physiological

measures: time-on-task ($\beta = -0.14$, $95\ CIs = [-1.34, 1.13]$, $t(50.21) = 0.24$, $p = .812$), heart

rate ($\beta = -1.02$, $95\ CIs = [-3.43, 1.25]$, $t(52.62) = 0.78$, $p = .441$), alpha power ($\beta =$

-0.02, $95\ CIs = [-0.04,\ 0.01]$, $t(48.09) = 1.21$, $p = .232$),   and   HRV   ($\beta =$

-0.003, $95\ CIs = [-0.01, 4.92 \times 10^{-5}]$, $t(2302) = 1.99$, $p = .047$).   Note   that,   while

frequentist estimation indicated a significant association between HRV and affective arousal ($p =$

$.045$), the Bayesian 95% credible interval included zero. This discrepancy was likely driven by

extremely low between-participant variance in the HRV slope estimate ($\sigma = 0.002$; see Figure

2C), which inflated the estimated degrees of freedom and, in turn, lowered the threshold for

significance in the t distribution. These results converge to support the idea that physiological

measures of arousal were largely unrelated to affective arousal ratings at the group-average level.

Having explored the relationship between subjective and physiological arousal, we next

sought to more fully characterize what affective arousal measures capture by examining its

relationship with other self-reported dimensions of ongoing experience (see bottom row of *Figure*

2C). We report results from hierarchical Bayesian linear regression models for the four experience

sampling items that exhibited the highest average participant-level correlations with affective

arousal. Group-level 95% credible intervals excluded zero for each of these experience sampling

items, indicating significant relationships with affective arousal. Specifically, higher levels of

affective arousal were associated with increased future thinking ($\beta = 5.03$, $95\ CIs =$

$[2.89, 7.16]$, $t(52.05) = 4.86$, $p < .001$), increased deliberate thinking ($\beta = 4.69$, $95\ CIs =$

$[2.54, 6.92]$, $t(52.0) = 4.30$, $p < .001$), increased self-related thinking ($\beta = 4.45$, $95\ CIs =$

$[2.19, 6.61]$, $t(51.49) = 3.97$, $p < .001$), and increased difficulty disengaging from thinking

($\beta = 3.51$, $95\ CIs = [1.44, 5.56]$, $t(48.62) = 3.34$, $p = .002$). While affective arousal was

overall related to these four dimensions of ongoing experience, the moderate strength of the

standardized regression coefficients across participants suggests that affective arousal carries

independent variance from other dimensions of ongoing experience.

Figure 2. Characteristics of affective arousal ratings and relationships with other physiological and
subjective measures. (A) Gaussian kernel density estimates for affective arousal ratings across an
example subset of participants. A distribution reflects the frequency of responses on the affective
arousal scale for a particular participant across the experiment. (B) Timeseries of affective arousal
ratings across all 50 trials for an example set of 3 participants. Each point reflects a particular
affective arousal rating submitted by a participant on one trial. (C) Within-participant posterior
distributions of the slope parameter between affective arousal and measures of both physiological
arousal and other dimensions of ongoing experience. For experience measures, higher values on
the x-axis reflect (i) increased future thinking, (ii) increased deliberate thinking, (iii) increased
self-related thinking, and (iv) increased difficulty to disengage from thinking. Points reflect means
of participant-level posterior distributions; bands around points reflect 95% credible intervals of
the participant-level posterior distribution. Top line of panel text reflects the group-level posterior
mean and 95% credible interval for the slope parameter. Bottom line reports the *p* value estimated
from a frequentist hierarchical model using the Satterthwaite approximation for degrees of
freedom.

Lower affective arousal is associated with greater HEP amplitude

To assess the relationship between affective arousal and cardiac interoceptive sensitivity,

we examined the association between the HEP and affective arousal ratings using a regression-

based cluster permutation analysis. This analysis revealed a significant cluster at frontal electrodes

(F3, F4, FC1, FC2, Fp1, Fp2, Fz; see *Figure 3B*) from 0.340 to 0.356 s post R peak, where

regression coefficients were negative in this cluster (see *Figure 3A*), suggesting that decreased

affective arousal ratings are associated with increased HEP (see *Figure 3C*. The direction of this

effect was largely consistent across participants (see *Figure 3D*). A surrogate cluster control test

revealed that the cluster-wise statistic is significantly greater than what would be expected from a

null effect not time-locked to heartbeats ($p = .02$; see *Figure S3*). Since larger HEP amplitude is

thought to reflect greater interoception (Judah et al., 2018; Pang et al., 2019; Verdonk et al., 2024),

the direction of this effect implies lower affective arousal during states of greater cardiac

interoceptive sensitivity. Interestingly, the topology, time course, and direction of this effect closely

replicate and extend a finding from past work, where affective arousal was driven by an external

event rather than spontaneously fluctuating at rest (Fourcade et al., 2024).

13
14   *Figure 3*. HEP effect across time, participants, and electrodes. (A) Average regression slope across
15   the heartbeat epoch. Timepoint zero reflects the R peak, and colors denote whether EEG voltage
16   was positively (orange) or negatively (blue) associated with affective arousal ratings. Shaded

regions reflect standard error of the mean. The green dashed lines reflect the significant time
window indicated by the cluster-based permutation analysis. The regression slopes are averaged
across significant electrodes. (B) Topographical map of *t*-values for slopes comparing HEP
amplitude to affective arousal. (C) HEP grand average across the heartbeat epoch, visualized across
within-participant median split in affective arousal ratings. Timepoint zero reflects the R-peak, and
colors denote the affective arousal direction. The green shaded region reflects the significant time
window indicated by the cluster-based permutation analysis (0.340 – 0.356 s), while the colored
shaded regions reflect standard errors of the mean. The EEG waveform is averaged across
significant electrodes (i.e., F3, F4, FC1, FC2, Fp1, Fp2, Fz). (D) HEP averaged across a
participant-level median split in affective arousal collapsed across the significant time window and
channels, shown separately for each participant (dotted lines connect averages across conditions
from the same participant).

The relationship between affective arousal and cardiac interoceptive sensitivity depends on

trait anxiety

15   To investigate whether increased cardiac interoceptive sensitivity during states of lower

affective arousal is linked to anxiety, we examined whether the size of the participant-level HEP

effect correlated with individual difference measures in both trait (GAD) and state (STAI-S)

anxiety. While there was a non-significant relationship between the HEP effect and STAI-S (r(29)

= -.25, p = .182), there was a significant positive correlation between the HEP effect and GAD

(r(49) = -.34 , p = .015; see *Figure 4A*). In other words, the extent to which HEP amplitude was

more positive for lower states of arousal was increased for those individuals who reported greater

trait   anxiety   (*Figure*   *4B*).

Figure 4. Association between HEP arousal effect (averaged HEP-arousal slope) and individual
differences in anxiety. (A) Correlations of trait (GAD) and state (STAI-S) anxiety with individual
differences in the HEP effect magnitude (lower on the y-axis reflects larger HEP amplitude for low
vs. high affective arousal). (B) HEP grand average across the heartbeat epoch, broken down by
within-participant median split on affective arousal ratings and a median split on trait anxiety
(GAD). Insets zoom in to the significant time window, indicated by the shaded green region.
Shaded colored regions reflect one standard error from the mean.

Discussion

Our experiment found that cardiac interoceptive sensitivity increased during lower

affective arousal during spontaneous thought, with this effect amplified in individuals with high

trait anxiety. Affective arousal was largely distinct from physiological measures like time-on-task,

parieto-occipital alpha power, and heart rate, though the relationship with heart rate variability was

more equivocal. Affective arousal also differed from other aspects of spontaneous mental

experience and was positively associated with future thinking, deliberate thinking, self-related

thinking, and difficulty disengaging from thoughts. Cardiac interoceptive sensitivity, indexed by

the HEP, showed greater amplitude prior to lower affective arousal ratings, localized at frontal

electrodes (0.340 – 0.356 ms post-heartbeat). This effect was stronger in those with high trait

anxiety. Our findings refine the concept of arousal and have clinical relevance for anxiety and

interoceptive dysregulation (Khalsa & Lapidus, 2016).

Arousal is widely studied but inconsistently defined (Lindquist et al., 2016; Satpute et al.,

2019). Measures such as self-report, skin conductance, heart rate, respiratory frequency, and EEG

alpha power often show weak or inconsistent correlations among each other (Agren, 2023;

Leonidou & Panayiotou, 2022; Wenzler et al., 2017). This inconsistency has led some to

conceptualize arousal as multidimensional (Reid et al., 2024), while others argue that distinct types

of arousal may share underlying cortical pathways, namely the presupplementary motor areas and

dorsal anterior insula (Sabat et al., 2025). These competing perspectives underscore the need to

differentiate aspects of arousal. In our data, spontaneous fluctuations in self-reported affective

arousal showed little association with physiological measures: 95% credible intervals for all

physiological predictors included zero. Although HRV reached nominal significance in a

frequentist model, the effect size was small and exhibited minimal between-subject variability.

These findings add to growing evidence that subjective and physiological arousal may reflect

distinct components of a broader arousal construct.

Our findings lend support to the idea that arousal is a multidimensional construct with

independent subcomponents (Reid et al., 2024). Although arousal has been widely studied, few

investigations have examined spontaneous fluctuations in affective arousal during wakeful rest

(but see Hoemann et al., 2020). The association between affective and physiological arousal may

depend on whether attention is focused on external stimuli (Leonidou & Panayiotou, 2022), raising

the possibility that self-reported arousal reflects deliberate rather than purely physiological

processes. Future research should examine how spontaneous fluctuations in affective arousal relate

to attentional focus and how this interaction influences the link (or lack thereof) between subjective

and physiological arousal.

Affective arousal was moderately associated with other dimensions of ongoing experience,

such as future thinking, deliberate thinking, self-related thinking, and difficulty to disengage from

thoughts. Deliberate thinking, which involves effortful problem solving, has been linked to

increased physiological and affective arousal, particularly when deliberation is framed as task

engagement (Kahneman, 2011; Matthews et al., 2010). Our results extend these ideas by showing

a positive link between affective arousal and deliberate thinking even when deliberate thinking is

not tied to an active task. Difficulty disengaging from thoughts—often linked to repetitive negative

thinking and worrying, key features of generalized anxiety disorder (Ehring & Watkins, 2008) and

state anxiety (Vălenaş et al., 2017)—was also associated with affective arousal. This finding aligns

with prior research showing that self-reported trait anxiety positively correlates with physiological

arousal (Hoehn-Saric & McLeod, 2000; Noteboom et al., 2001; Roos et al., 2021). Finally,

affective arousal was positively associated with both self- and future-thinking. This finding

contrasts somewhat with recent work showing a negative association, at the inter-individual level, between conceptual thinking (as opposed to bodily focus) and physiological arousal (Banellis et al., 2024). Self- and future-thinking tend to co-occur and are associated with improved future mood (Ruby et al., 2013). One possibility is that affective arousal facilitates this relationship, where engaging in self- and future-thinking enhances arousal, which in turn supports mood regulation. However, how affective arousal and self- and future-thinking impact future mood likely depends on individual differences in anxiety. Anxiety has been associated with more vivid and detailed autobiographical future thinking (Du et al., 2022), which may amplify or distort the impact of future thinking on mood.

We investigated how interoceptive processing varies as a function of spontaneous fluctuations in affective arousal by examining the HEP. We found that in the 340–356 ms window post-heartbeat, HEP amplitude was increased at frontal electrodes in the 10 s preceding lower affective arousal ratings. This finding suggests that interoceptive processing is enhanced during states of lower affective arousal. The timing and topography of this effect align with previous studies linking the HEP to arousal (Fourcade et al., 2024; Fukushima et al., 2011; Gentsch et al., 2019; Ito et al., 2019; Luft & Bhattacharya, 2015; MacKinnon et al., 2013; Marshall et al., 2017, 2018, 2020; Park et al., 2016; Schulz et al., 2013, 2015; Sel et al., 2017; Shao et al., 2011), though definitions of arousal vary widely across studies. One study particularly relevant to our work recorded electrophysiological responses while participants experienced virtual reality (VR) roller coaster rides and retrospectively rated fluctuations in affective arousal (Fourcade et al., 2024). Their results closely parallel ours, showing increased HEP for lower affective arousal at fronto-central electrodes in a similar time window. This convergence across methodologies strengthens the link between affective arousal and interoceptive processing indexed by the HEP. Our findings

extend this prior work that showed differences in HEP driven by external events (i.e., roller coaster

videos; Fourcade et al., 2024) by revealing that such differences in HEP can also be driven by

spontaneously fluctuating internal states during wakeful rest. Fourcade et al. (2024) performed

source localization analyses on their EEG results to suggest that their effect, which closely parallels

ours, originates from somatosensory areas. Indeed, past work using intracranial EEG has

implicated a variety of brain regions in HEP activity during rest, including somatosensory cortex,

insula, opercular cortex, inferior frontal gyrus, and amygdala (Babo-Rebelo, Wolpert, et al., 2016;

Canales-Johnson et al., 2015; Kern et al., 2013; Park et al., 2018; Park & Blanke, 2019).

One possible explanation for increased HEP during lower affective arousal is that reduced

arousal may, in some contexts, enhance interoception. This inverse relationship is evident in

practices like meditation, which both lower arousal and improve interoceptive awareness (Hanh,

2015; Kushner & Marnocha, 2008; Nathoo, 2021; Shapiro & Carlson, 2009). In Theravada

Buddhism, bodily calm is cultivated to heighten sensitivity to internal sensations and support

insight into impermanence (Analayo, 2018). Empirical work links meditation to enhanced

interoceptive awareness, potentially through altered neural structure and function, particularly in

the insula (Farb et al., 2013; Hölzel et al., 2008; Luders et al., 2012). An alternative explanation is

that lower HEP during high arousal reflects a compensatory downregulation of interoception—

especially in individuals prone to anxiety, who may suppress bodily awareness during high-arousal

states (Russell, 1980, 2003). Supporting this view, the present study found larger HEP differences

among individuals higher in trait anxiety (see orange lines, Figure 4B). Although clinical anxiety

has been linked to disrupted interoception, findings remain mixed (Abrams et al., 2018; Ardizzi &

Ferri, 2018; Gaebler et al., 2013; Khalsa & Lapidus, 2016; Stewart et al., 2001). These results

underscore the importance of considering how momentary affective arousal shapes the interplay

between anxiety and interoceptive processing.

This study had several limitations that can be addressed in future work. The state anxiety

measure (i.e., STAI-S) was a late addition to the experiment, and as such we have data from only

30 of the 51 total participants. Our sample also contained few individuals scoring high on trait

anxiety (i.e., GAD; see *Figure 4a*). The present experiment was not specifically designed to

examine these correlations and may have been underpowered to detect them; thus, associations

with anxiety measures should be interpreted as preliminary. Future studies should replicate our

findings related to trait anxiety in a larger and more clinically diverse sample. Such a replication

is particularly important given the inconsistent links between HEP and anxiety in the literature (Qi

et al., 2025). A further limitation concerns the degree to which findings in our laboratory setting

translate to real-world contexts. During daily life, many individuals are rarely truly at rest (i.e.,

devoid of prominent external stimuli or task demands). What may prove crucial in the study of

mental health is understanding how spontaneously fluctuating internal states of affective arousal

interact with salient external conditions or environmental cues to influence the regulation of

anxiety.

In conclusion, our findings offer advanced knowledge of the neural basis of affective

arousal and have broad implications for understanding various mental health conditions marked

by emotional and interoceptive dysregulation. The interactions between HEP, affective arousal,

and anxiety that we identified suggest a potential naturally-operating mechanism in which

interoception is spontaneously downregulated to facilitate emotional regulation. Further research

is needed to causally test for such a mechanism, how it may be modulated, and whether evidence

of this mechanism is present in other clinical populations, such as individuals with generalized

anxiety disorder.

Code Accessibility

We will make the custom code used to analyze the electrophysiological data available upon

publication.

References

Abrams, K., Cieslowski, K., Johnson, S., Krimmel, S., La Rosa, G. B.-D., Barton, K., &

Silverman, P. (2018). The effects of alcohol on heartbeat perception: Implications for

anxiety. *Addictive Behaviors*, *79*, 151–158. https://doi.org/10.1016/j.addbeh.2017.12.023

Agren, T. (2023). Physiological and subjective arousal to prospective mental imagery: A

mechanism for behavioral change? *PLOS ONE*, *18*(12), e0294629.

https://doi.org/10.1371/journal.pone.0294629

Al, E., Iliopoulos, F., Forschack, N., Nierhaus, T., Grund, M., Motyka, P., Gaebler, M., Nikulin,

V. V., & Villringer, A. (2020). Heart–brain interactions shape somatosensory perception

and evoked potentials. *Proceedings of the National Academy of Sciences*, *117*(19),

10575–10584. https://doi.org/10.1073/pnas.1915629117

Al, E., Iliopoulos, F., Nikulin, V. V., & Villringer, A. (2021). Heartbeat and somatosensory

perception. *NeuroImage*, *238*, 118247. https://doi.org/10.1016/j.neuroimage.2021.118247

Analayo, B. (2018). *Satipatthana Meditation: A Practice Guide*. Windhorse Publications.

Ardizzi, M., & Ferri, F. (2018). Interoceptive influences on peripersonal space boundary.

Cognition, *177*, 79–86. https://doi.org/10.1016/j.cognition.2018.04.001

Azzalini, D., Rebollo, I., & Tallon-Baudry, C. (2019). Visceral Signals Shape Brain Dynamics

and Cognition. *Trends in Cognitive Sciences*, *23*(6), 488–509.

https://doi.org/10.1016/j.tics.2019.03.007

Babo-Rebelo, M., Buot, A., & Tallon-Baudry, C. (2019). Neural responses to heartbeats

distinguish self from other during imagination. *NeuroImage*, *191*, 10–20.

https://doi.org/10.1016/j.neuroimage.2019.02.012

Babo-Rebelo, M., Richter, C. G., & Tallon-Baudry, C. (2016). Neural Responses to Heartbeats in the Default Network Encode the Self in Spontaneous Thoughts. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *36*(30), 7829–7840. https://doi.org/10.1523/JNEUROSCI.0262-16.2016

Babo-Rebelo, M., Wolpert, N., Adam, C., Hasboun, D., & Tallon-Baudry, C. (2016). Is the cardiac monitoring function related to the self in both the default network and right anterior insula? *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *371*(1708), 20160004. https://doi.org/10.1098/rstb.2016.0004

Banellis, L., Nikolova, N., Brændholdt, M., Vejlø, M., Fardo, F., Smallwood, J., & Allen, M. G. (2024). *Body-wandering reveals an embodied dimension of thought with distinct affective and neural signatures* (p. 2024.10.25.620252). bioRxiv. https://doi.org/10.1101/2024.10.25.620252

Barrett, L. F. (1998). Discrete Emotions or Dimensions? The Role of Valence Focus and Arousal Focus. *Cognition and Emotion*, *12*(4), 579–599. https://doi.org/10.1080/026999398379574

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*, 1–48. https://doi.org/10.18637/jss.v067.i01

Bliss-Moreau, E., Williams, L. A., & Santistevan, A. C. (2020). The immutability of valence and arousal in the foundation of emotion. *Emotion*, *20*(6), 993–1004. https://doi.org/10.1037/emo0000606

BrainCap MR. (2020). [Computer software].

Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software*, *80*(1), 1–28. https://doi.org/10.18637/jss.v080.i01

Canales-Johnson, A., Silva, C., Huepe, D., Rivera-Rei, Á., Noreika, V., Garcia, M. del C., Silva, W., Ciraolo, C., Vaucheret, E., Sedeño, L., Couto, B., Kargieman, L., Baglivo, F., Sigman, M., Chennu, S., Ibáñez, A., Rodríguez, E., & Bekinschtein, T. A. (2015). Auditory Feedback Differentially Modulates Behavioral and Neural Markers of Objective and Subjective Performance When Tapping to Your Heartbeat. *Cerebral Cortex*, *25*(11), 4490–4503. https://doi.org/10.1093/cercor/bhv076

Chang, C., Leopold, D. A., Schölvinck, M. L., Mandelkow, H., Picchioni, D., Liu, X., Ye, F. Q., Turchi, J. N., & Duyn, J. H. (2016). Tracking brain arousal fluctuations with fMRI. *Proceedings of the National Academy of Sciences*, *113*(16), 4518–4523. https://doi.org/10.1073/pnas.1520613113

Christoff, K., Gordon, A. M., Smallwood, J., Smith, R., & Schooler, J. W. (2009). Experience sampling during fMRI reveals default network and executive system contributions to mind wandering. *Proceedings of the National Academy of Sciences*, *106*(21), 8719–8724. https://doi.org/10.1073/pnas.0900234106

Christoff, K., Irving, Z. C., Fox, K. C. R., Spreng, R. N., & Andrews-Hanna, J. R. (2016). Mind-wandering as spontaneous thought: A dynamic framework. *Nature Reviews Neuroscience*, *17*(11), 718–731. https://doi.org/10.1038/nrn.2016.113

Coll, M.-P., Hobson, H., Bird, G., & Murphy, J. (2021). Systematic review and meta-analysis of the relationship between the heartbeat-evoked potential and interoception. *Neuroscience & Biobehavioral Reviews*, *122*, 190–200. https://doi.org/10.1016/j.neubiorev.2020.12.012

Debener, S., Thorne, J., Schneider, T., & Campos Viola, F. (2010). Using ICA for the Analysis of Multi-Channel EEG Data. In M. Ullsperger & S. Debener (Eds.), *Simultaneous EEG and fMRI: Recording, Analysis, and Application*. Oxford University Press, New York.

Devuyst, S., Dutoit, T., Stenuit, P., Kerkhofs, M., & Stanus, E. (2008). Removal of ECG artifacts from EEG using a modified independent component analysis approach. *2008 30th Annual International Conference of the IEEE Engineering in Medicine and Biology Society*, 5204–5207. https://doi.org/10.1109/IEMBS.2008.4650387

Du, J. Y., Hallford, D. J., & Busby Grant, J. (2022). Characteristics of episodic future thinking in anxiety: A systematic review and meta-analysis. *Clinical Psychology Review*, *95*, 102162. https://doi.org/10.1016/j.cpr.2022.102162

Ehring, T., & Watkins, E. R. (2008). Repetitive Negative Thinking as a Transdiagnostic Process. *International Journal of Cognitive Therapy*, *1*(3), 192–205. https://doi.org/10.1521/ijct.2008.1.3.192

Engelen, T., Solcà, M., & Tallon-Baudry, C. (2023). Interoceptive rhythms in the brain. *Nature Neuroscience*, *26*(10), 1670–1684. https://doi.org/10.1038/s41593-023-01425-1

Farb, N. A. S., Segal, Z. V., & Anderson, A. K. (2013). Mindfulness meditation training alters cortical representations of interoceptive attention. *Social Cognitive and Affective Neuroscience*, *8*(1), 15–26. https://doi.org/10.1093/scan/nss066

Fell, J., Chaieb, L., & Hoppe, C. (2023). Mind wandering in anxiety disorders: A status report. *Neuroscience & Biobehavioral Reviews*, *155*, 105432. https://doi.org/10.1016/j.neubiorev.2023.105432

Flasbeck, V., Jungilligens, J., Lemke, I., Beckers, J., Öztürk, H., Wellmer, J., Seliger, C., Juckel, G., & Popkirov, S. (2024). Heartbeat evoked potentials and autonomic arousal during

dissociative seizures: Insights from electrophysiology and neuroimaging. *BMJ Neurology Open*, *6*(1), e000665. https://doi.org/10.1136/bmjno-2024-000665

Fourcade, A., Klotzsche, F., Hofmann, S. M., Mariola, A., Nikulin, V. V., Villringer, A., & Gaebler, M. (2024). Linking brain–heart interactions to emotional arousal in immersive virtual reality. *Psychophysiology*, e14696. https://doi.org/10.1111/psyp.14696

Fukushima, H., Terasawa, Y., & Umeda, S. (2011). Association between interoception and empathy: Evidence from heartbeat-evoked brain potential. *International Journal of Psychophysiology*, *79*(2), 259–265. https://doi.org/10.1016/j.ijpsycho.2010.10.015

Gaebler, M., Daniels, J. K., Lamke, J.-P., Fydrich, T., & Walter, H. (2013). Heart rate variability and its neural correlates during emotional face processing in social anxiety disorder. *Biological Psychology*, *94*(2), 319–330. https://doi.org/10.1016/j.biopsycho.2013.06.009

Gentsch, A., Sel, A., Marshall, A. C., & Schütz-Bosbach, S. (2019). Affective interoceptive inference: Evidence from heart-beat evoked brain potentials. *Human Brain Mapping*, *40*(1), 20–33. https://doi.org/10.1002/hbm.24352

Gergelyfi, M., Jacob, B., Olivier, E., & Zénon, A. (2015). Dissociation between mental fatigue and motivational state during prolonged mental activity. *Frontiers in Behavioral Neuroscience*, *9*. https://doi.org/10.3389/fnbeh.2015.00176

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., Goj, R., Jas, M., Brooks, T., Parkkonen, L., & Hämäläinen, M. (2013). MEG and EEG data analysis with MNE-Python. *Frontiers in Neuroscience*, *7*. https://doi.org/10.3389/fnins.2013.00267

Gronau, Q. F., Sarafoglou, A., Matzke, D., Ly, A., Boehm, U., Marsman, M., Leslie, D. S., Forster, J. J., Wagenmakers, E.-J., & Steingroever, H. (2017). A tutorial on bridge

sampling. *Journal of Mathematical Psychology*, *81*, 80–97.

https://doi.org/10.1016/j.jmp.2017.09.005

Gronau, Q. F., Singmann, H., & Wagenmakers, E.-J. (2020). bridgesampling: An R Package for

Estimating Normalizing Constants. *Journal of Statistical Software*, *92*, 1–29.

https://doi.org/10.18637/jss.v092.i10

Gunning, M. D., Denison, F. C., Stockley, C. J., Ho, S. P., Sandhu, H. K., & Reynolds, R. M.

(2010). Assessing maternal anxiety in pregnancy with the State-Trait Anxiety Inventory

(STAI): Issues of validity, location and participation. *Journal of Reproductive and Infant

Psychology*, *28*(3), 266–273. https://doi.org/10.1080/02646830903487300

Haj-Ali, H., Anderson, A. K., & Kron, A. (2020). Comparing three models of arousal in the

human brain. *Social Cognitive and Affective Neuroscience*, *15*(1), 1–11.

https://doi.org/10.1093/scan/nsaa012

Hanh, T. N. (2015). *How to relax* (Vol. 5). Parallax Press.

Harris, P. A., Taylor, R., Minor, B. L., Elliott, V., Fernandez, M., O'Neal, L., McLeod, L.,

Delacqua, G., Delacqua, F., Kirby, J., & Duda, S. N. (2019). The REDCap consortium:

Building an international community of software platform partners. *Journal of

Biomedical Informatics*, *95*, 103208. https://doi.org/10.1016/j.jbi.2019.103208

Harris, P. A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., & Conde, J. G. (2009). Research

electronic data capture (REDCap)—A metadata-driven methodology and workflow

process for providing translational research informatics support. *Journal of Biomedical

Informatics*, *42*(2), 377–381. https://doi.org/10.1016/j.jbi.2008.08.010

Herlambang, M. B., Taatgen, N. A., & Cnossen, F. (2019). The Role of Motivation as a Factor in

Mental Fatigue. *Human Factors*, *61*(7), 1171–1185.

https://doi.org/10.1177/0018720819828569

Hoehn-Saric, R., & McLeod, D. R. (2000). Anxiety and arousal: Physiological changes and their

perception. *Journal of Affective Disorders*, *61*(3), 217–224.

https://doi.org/10.1016/S0165-0327(00)00339-6

Hoemann, K., Khan, Z., Feldman, M. J., Nielson, C., Devlin, M., Dy, J., Barrett, L. F.,

Wormwood, J. B., & Quigley, K. S. (2020). Context-aware experience sampling reveals

the scale of variation in affective experience. *Scientific Reports*, *10*(1), 12459.

https://doi.org/10.1038/s41598-020-69180-y

Hoffmann, S., & Falkenstein, M. (2008). The Correction of Eye Blink Artefacts in the EEG: A

Comparison of Two Prominent Methods. *PLOS ONE*, *3*(8), e3004.

https://doi.org/10.1371/journal.pone.0003004

Hofmann, S. M., Klotzsche, F., Mariola, A., Nikulin, V., Villringer, A., & Gaebler, M. (2021).

Decoding subjective emotional arousal from EEG during an immersive virtual reality

experience. *eLife*, *10*, e64812. https://doi.org/10.7554/eLife.64812

Hölzel, B. K., Ott, U., Gard, T., Hempel, H., Weygandt, M., Morgen, K., & Vaitl, D. (2008).

Investigation of mindfulness meditation practitioners with voxel-based morphometry.

Social Cognitive and Affective Neuroscience, *3*(1), 55–61.

https://doi.org/10.1093/scan/nsm038

Ito, Y., Shibata, M., Tanaka, Y., Terasawa, Y., & Umeda, S. (2019). Affective and temporal

orientation of thoughts: Electrophysiological evidence. *Brain Research*, *1719*, 148–156.

https://doi.org/10.1016/j.brainres.2019.05.041

James, W. (1948). What is emotion? 1884. In *Readings in the history of psychology* (pp. 290–303). Appleton-Century-Crofts. https://doi.org/10.1037/11304-033

Janelle, C. M. (2002). Anxiety, arousal and visual attention: A mechanistic account of performance variability. *Journal of Sports Sciences*, *20*(3), 237–251. https://doi.org/10.1080/026404102317284790

Jas, M., Engemann, D. A., Bekhti, Y., Raimondo, F., & Gramfort, A. (2017). Autoreject: Automated artifact rejection for MEG and EEG data. *NeuroImage*, *159*, 417–429. https://doi.org/10.1016/j.neuroimage.2017.06.030

Jeffreys, H. (1998). *The Theory of Probability* (3rd ed.). Oxford, England. (Original work published 1961)

Judah, M. R., Shurkova, E. Y., Hager, N. M., White, E. J., Taylor, D. L., & Grant, D. M. (2018). The relationship between social anxiety and heartbeat evoked potential amplitude. *Biological Psychology*, *139*, 1–7. https://doi.org/10.1016/j.biopsycho.2018.09.013

Kahneman, D. (2011). *Thinking, Fast and Slow*. Macmillan.

Kam, J. W. Y., Irving, Z. C., Mills, C., Patel, S., Gopnik, A., & Knight, R. T. (2021). Distinct electrophysiological signatures of task-unrelated and dynamic thoughts. *Proceedings of the National Academy of Sciences*, *118*(4), e2011796118. https://doi.org/10.1073/pnas.2011796118

Karthikeyan, R., Carrizales, J., Johnson, C., & Mehta, R. K. (2022). A Window Into the Tired Brain: Neurophysiological Dynamics of Visuospatial Working Memory Under Fatigue. *Human Factors*, *66*(2), 528–543. https://doi.org/10.1177/00187208221094900

Karthikeyan, R., Smoot, M. R., & Mehta, R. K. (2021). Anodal tDCS augments and preserves working memory beyond time-on-task deficits. *Scientific Reports*, *11*(1), 19134. https://doi.org/10.1038/s41598-021-98636-y

Kern, M., Aertsen, A., Schulze-Bonhage, A., & Ball, T. (2013). Heart cycle-related effects on event-related potentials, spectral power changes, and connectivity patterns in the human ECoG. *NeuroImage*, *81*, 178–190. https://doi.org/10.1016/j.neuroimage.2013.05.042

Khalsa, S. S., & Lapidus, R. C. (2016). Can Interoception Improve the Pragmatic Search for Biomarkers in Psychiatry? *Frontiers in Psychiatry*, *7*. https://doi.org/10.3389/fpsyt.2016.00121

Killingsworth, M. A., & Gilbert, D. T. (2010). A wandering mind is an unhappy mind. *Science*, *330*(6006), 932–932.

Kluger, D. S., Allen, M. G., & Gross, J. (2024). Brain–body states embody complex temporal dynamics. *Trends in Cognitive Sciences*, *28*(8), 695–698. https://doi.org/10.1016/j.tics.2024.05.003

Kron, A., Pilkiw, M., Banaei, J., Goldstein, A., & Anderson, A. K. (2015). Are valence and arousal separable in emotional experience? *Emotion*, *15*(1), 35–44. https://doi.org/10.1037/a0038474

Kucyi, A., Anderson, N., Bounyarith, T., Braun, D., Shareef-Trudeau, L., Treves, I., Braga, R. M., Hsieh, P.-J., & Hung, S.-M. (2024). Individual variability in neural representations of mind-wandering. *Network Neuroscience*, 1–66. https://doi.org/10.1162/netn_a_00387

Kucyi, A., & Parvizi, J. (2020). Pupillary Dynamics Link Spontaneous and Task-Evoked Activations Recorded Directly from Human Insula. *Journal of Neuroscience*, *40*(32), 6207–6218. https://doi.org/10.1523/JNEUROSCI.0435-20.2020

Kuppens, P., Tuerlinckx, F., Russell, J. A., & Barrett, L. F. (2013). The relation between valence and arousal in subjective experience. *Psychological Bulletin*, *139*(4), 917–940. https://doi.org/10.1037/a0030811

Kushner, K., & Marnocha, M. (2008). Meditation and relaxation. In *Evidence-based adjunctive treatments* (pp. 177–205). Elsevier Academic Press. https://doi.org/10.1016/B978-012088520-6.50010-X

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*, 1–26. https://doi.org/10.18637/jss.v082.i13

Larson, E., Gramfort, A., Engemann, D. A., Leppakangas, J., Brodbeck, C., Jas, M., Brooks, T., Sassenhagen, J., Luessi, M., McCloy, D., King, J.-R., Höchenberger, R., Goj, R., Favelier, G., Brunner, C., van Vliet, M., Wronkiewicz, M., Holdgraf, C., Rockhill, A., … luzpaz. (2023). *MNE-Python* (Version v1.5.1) [Computer software]. Zenodo. https://doi.org/10.5281/zenodo.8322569

Lee, T.-W., Girolami, M., & Sejnowski, T. J. (1999). Independent Component Analysis Using an Extended Infomax Algorithm for Mixed Subgaussian and Supergaussian Sources. *Neural Computation*, *11*(2), 417–441. Neural Computation. https://doi.org/10.1162/089976699300016719

Leonidou, C., & Panayiotou, G. (2022). Can we predict experiential avoidance by measuring subjective and physiological emotional arousal? *Current Psychology*, *41*(10), 7215–7227. https://doi.org/10.1007/s12144-020-01317-8

Lindquist, K. A., Satpute, A. B., Wager, T. D., Weber, J., & Barrett, L. F. (2016). The Brain Basis of Positive and Negative Affect: Evidence from a Meta-Analysis of the Human

Neuroimaging Literature. *Cerebral Cortex*, *26*(5), 1910–1922.

https://doi.org/10.1093/cercor/bhv001

Löwe, B., Decker, O., Müller, S., Brähler, E., Schellberg, D., Herzog, W., & Herzberg, P. Y.

(2008). Validation and standardization of the Generalized Anxiety Disorder Screener

(GAD-7) in the general population. *Medical Care*, *46*(3), 266–274.

https://doi.org/10.1097/MLR.0b013e318160d093

Lozano-Soldevilla, D. (2018). On the Physiological Modulation and Potential Mechanisms

Underlying Parieto-Occipital Alpha Oscillations. *Frontiers in Computational*

Neuroscience, *12*, 23. https://doi.org/10.3389/fncom.2018.00023

Luders, E., Kurth, F., Mayer, E. A., Toga, A. W., Narr, K. L., & Gaser, C. (2012). The Unique

Brain Anatomy of Meditation Practitioners: Alterations in Cortical Gyrification. *Frontiers*

in Human Neuroscience, *6*. https://doi.org/10.3389/fnhum.2012.00034

Luft, C. D. B., & Bhattacharya, J. (2015). Aroused with heart: Modulation of heartbeat evoked

potential by arousal induction and its oscillatory correlates. *Scientific Reports*, *5*(1),

15717. https://doi.org/10.1038/srep15717

MacKinnon, S., Gevirtz, R., McCraty, R., & Brown, M. (2013). Utilizing Heartbeat Evoked

Potentials to Identify Cardiac Regulation of Vagal Afferents During Emotion and

Resonant Breathing. *Applied Psychophysiology and Biofeedback*, *38*(4), 241–255.

https://doi.org/10.1007/s10484-013-9226-5

Maher, J. P., Dzubur, E., Nordgren, R., Huh, J., Chou, C.-P., Hedeker, D., & Dunton, G. F.

(2019). Do fluctuations in positive affective and physical feeling states predict physical

activity and sedentary time? *Psychology of Sport and Exercise*, *41*, 153–161.

https://doi.org/10.1016/j.psychsport.2018.01.011

Marshall, A. C., Gentsch, A., Jelinčić, V., & Schütz-Bosbach, S. (2017). Exteroceptive expectations modulate interoceptive processing: Repetition-suppression effects for visual and heartbeat evoked potentials. *Scientific Reports*, *7*(1), 16525. https://doi.org/10.1038/s41598-017-16595-9

Marshall, A. C., Gentsch, A., Schröder, L., & Schütz-Bosbach, S. (2018). Cardiac interoceptive learning is modulated by emotional valence perceived from facial expressions. *Social Cognitive and Affective Neuroscience*, *13*(7), 677–686. https://doi.org/10.1093/scan/nsy042

Marshall, A. C., Gentsch, A., & Schütz-Bosbach, S. (2020). Interoceptive cardiac expectations to emotional stimuli predict visual perception. *Emotion*, *20*(7), 1113–1126. https://doi.org/10.1037/emo0000631

Matthews, G., Warm, J. S., Reinerman, L. E., Langheim, L. K., & Saxby, D. J. (2010). Task Engagement, Attention, and Executive Control. In A. Gruszka, G. Matthews, & B. Szymura (Eds.), *Handbook of Individual Differences in Cognition: Attention, Memory, and Executive Control* (pp. 205–230). Springer. https://doi.org/10.1007/978-1-4419-1210-7_13

Matuz, A., Van der Linden, D., Topa, K., & Csathó, Á. (2019). Cross-Modal Conflict Increases With Time-on-Task in a Temporal Discrimination Task. *Frontiers in Psychology*, *10*. https://doi.org/10.3389/fpsyg.2019.02429

Melo, H. M., Nascimento, L. M., & Takase, E. (2017). Mental fatigue and heart rate variability (HRV): The time-on-task effect. *Psychology & Neuroscience*, *10*(4), 428–436. https://doi.org/10.1037/pne0000110

Montoya, P., Schandry, R., & Müller, A. (1993). Heartbeat evoked potentials (HEP): Topography and influence of cardiac awareness and focus of attention. *Electroencephalography and Clinical Neurophysiology/Evoked Potentials Section*, *88*(3), 163–172. https://doi.org/10.1016/0168-5597(93)90001-6

Morey, R. D., & Rouder, J. N. (2011). Bayes factor approaches for testing interval null hypotheses. *Psychological Methods*, *16*(4), 406–419. https://doi.org/10.1037/a0024377

Morey, R. D., Rouder, J. N., Pratte, M. S., & Speckman, P. L. (2011). Using MCMC chain outputs to efficiently estimate Bayes factors. *Journal of Mathematical Psychology*, *55*(5), 368–378. https://doi.org/10.1016/j.jmp.2011.06.004

Nathoo, A. (2021). Relaxation and Meditation. In M. Farias, D. Brazier, & M. Lalljee (Eds.), *The Oxford Handbook of Meditation* (p. 0). Oxford University Press. https://doi.org/10.1093/oxfordhb/9780198808640.013.22

Nook, E. C., Sasse, S. F., Lambert, H. K., McLaughlin, K. A., & Somerville, L. H. (2017). Increasing verbal knowledge mediates development of multidimensional emotion representations. *Nature Human Behaviour*, *1*(12), 881–889. https://doi.org/10.1038/s41562-017-0238-7

Noteboom, J. T., Barnholt, K. R., & Enoka, R. M. (2001). Activation of the arousal response and impairment of performance increase with anxiety and stressor intensity. *Journal of Applied Physiology*, *91*(5), 2093–2101. https://doi.org/10.1152/jappl.2001.91.5.2093

Pang, J., Tang, X., Li, H., Hu, Q., Cui, H., Zhang, L., Li, W., Zhu, Z., Wang, J., & Li, C. (2019). Altered Interoceptive Processing in Generalized Anxiety Disorder—A Heartbeat-Evoked Potential Research. *Frontiers in Psychiatry*, *10*, 616. https://doi.org/10.3389/fpsyt.2019.00616

Park, H.-D., Bernasconi, F., Bello-Ruiz, J., Pfeiffer, C., Salomon, R., & Blanke, O. (2016). Transient Modulations of Neural Responses to Heartbeats Covary with Bodily Self-Consciousness. *Journal of Neuroscience*, *36*(32), 8453–8460. https://doi.org/10.1523/JNEUROSCI.0311-16.2016

Park, H.-D., Bernasconi, F., Salomon, R., Tallon-Baudry, C., Spinelli, L., Seeck, M., Schaller, K., & Blanke, O. (2018). Neural Sources and Underlying Mechanisms of Neural Responses to Heartbeats, and their Role in Bodily Self-consciousness: An Intracranial EEG Study. *Cerebral Cortex*, *28*(7), 2351–2364. https://doi.org/10.1093/cercor/bhx136

Park, H.-D., & Blanke, O. (2019). Heartbeat-evoked cortical responses: Underlying mechanisms, functional roles, and methodological considerations. *NeuroImage*, *197*, 502–511. https://doi.org/10.1016/j.neuroimage.2019.04.081

Petzschner, F. H., Weber, L. A., Wellstein, K. V., Paolini, G., Do, C. T., & Stephan, K. E. (2019). Focus of attention modulates the heartbeat evoked potential. *NeuroImage*, *186*, 595–606. https://doi.org/10.1016/j.neuroimage.2018.11.037

Pion-Tonachini, L., Kreutz-Delgado, K., & Makeig, S. (2019). ICLabel: An automated electroencephalographic independent component classifier, dataset, and website. *NeuroImage*, *198*, 181–197. https://doi.org/10.1016/j.neuroimage.2019.05.026

Pretorius, T. B., & Padmanabhanunni, A. (2023). Anxiety in Brief: Assessment of the Five-Item Trait Scale of the State-Trait Anxiety Inventory in South Africa. *International Journal of Environmental Research and Public Health*, *20*(9), 5697. https://doi.org/10.3390/ijerph20095697

Qi, M., Shen, X., Zeng, Y., Lin, X., Suliman, M., & Li, P. (2025). Interoception and mental health in middle-aged and elderly adults: A systematic review and meta-analysis.

Neuroscience & Biobehavioral Reviews, *172*, 106104.

https://doi.org/10.1016/j.neubiorev.2025.106104

Qin, H., Zhou, X., Ou, X., Liu, Y., & Xue, C. (2021). Detection of mental fatigue state using

heart rate variability and eye metrics during simulated flight. *Human Factors and*

Ergonomics in Manufacturing & Service Industries, *31*(6), 637–651.

https://doi.org/10.1002/hfm.20927

R Core Team. (2024). *R: A Language and Environment for Statistical Computing*. R Foundation

for Statistical Computing. https://www.R-project.org/

Reid, T., Nielson, C., & Wormwood, J. B. (2024). Measuring Arousal: Promises and Pitfalls.

Affective Science. https://doi.org/10.1007/s42761-024-00288-4

Roos, A.-L., Goetz, T., Voracek, M., Krannich, M., Bieg, M., Jarrell, A., & Pekrun, R. (2021).

Test Anxiety and Physiological Arousal: A Systematic Review and Meta-Analysis.

Educational Psychology Review, *33*(2), 579–618. https://doi.org/10.1007/s10648-020-

09543-z

Rouder, J. N., Speckman, P. L., Sun, D., Morey, R. D., & Iverson, G. (2009). Bayesian t tests for

accepting and rejecting the null hypothesis. *Psychonomic Bulletin & Review*, *16*(2), 225–

237. https://doi.org/10.3758/PBR.16.2.225

Ruby, F. J. M., Smallwood, J., Engen, H., & Singer, T. (2013). How Self-Generated Thought

Shapes Mood—The Relation between Mind-Wandering and Mood Depends on the

Socio-Temporal Content of Thoughts. *PLOS ONE*, *8*(10), e77554.

https://doi.org/10.1371/journal.pone.0077554

Russell, J. A. (1980). A circumplex model of affect. *Journal of Personality and Social*

Psychology, *39*(6), 1161–1178. https://doi.org/10.1037/h0077714

Russell, J. A. (2003). Core affect and the psychological construction of emotion. *Psychological Review*, *110*(1), 145–172. https://doi.org/10.1037/0033-295X.110.1.145

Russell, J. A., & Barrett, L. F. (1999). Core affect, prototypical emotional episodes, and other things called emotion: Dissecting the elephant. *Journal of Personality and Social Psychology*, *76*(5), 805–819. https://doi.org/10.1037/0022-3514.76.5.805

Sabat, M., De Dampierre, C., & Tallon-Baudry, C. (2025). Evidence for domain-general arousal from semantic and neuroimaging meta-analyses reconciles opposing views on arousal. *Proceedings of the National Academy of Sciences*, *122*(6), e2413808122. https://doi.org/10.1073/pnas.2413808122

Satpute, A. B., Kragel, P. A., Barrett, L. F., Wager, T. D., & Bianciardi, M. (2019). Deconstructing arousal into wakeful, autonomic and affective varieties. *Neuroscience Letters*, *693*, 19–28. https://doi.org/10.1016/j.neulet.2018.01.042

Schandry, R., Sparrer, B., & Weitkunat, R. (1986). From the heart to the brain: A study of heartbeat contingent scalp potentials. *International Journal of Neuroscience*, *30*(4), 261–275. https://doi.org/10.3109/00207458608985677

Schubring, D., & Schupp, H. T. (2021). Emotion and Brain Oscillations: High Arousal is Associated with Decreases in Alpha- and Lower Beta-Band Power. *Cerebral Cortex*, *31*(3), 1597–1608. https://doi.org/10.1093/cercor/bhaa312

Schulz, A., Ferreira de Sá, D. S., Dierolf, A. M., Lutz, A., van Dyck, Z., Vögele, C., & Schächinger, H. (2015). Short-term food deprivation increases amplitudes of heartbeat-evoked potentials. *Psychophysiology*, *52*(5), 695–703. https://doi.org/10.1111/psyp.12388

Schulz, A., Strelzyk, F., Ferreira de Sá, D. S., Naumann, E., Vögele, C., & Schächinger, H. (2013). Cortisol rapidly affects amplitudes of heartbeat-evoked brain potentials—

Implications for the contribution of stress to an altered perception of physical sensations? *Psychoneuroendocrinology*, *38*(11), 2686–2693. https://doi.org/10.1016/j.psyneuen.2013.06.027

Sel, A., Azevedo, R. T., & Tsakiris, M. (2017). Heartfelt Self: Cardio-Visual Integration Affects Self-Face Recognition and Interoceptive Cortical Processing. *Cerebral Cortex*, *27*(11), 5144–5155. https://doi.org/10.1093/cercor/bhw296

Shaffer, F., & Ginsberg, J. P. (2017). An Overview of Heart Rate Variability Metrics and Norms. *Frontiers in Public Health*, *5*, 258. https://doi.org/10.3389/fpubh.2017.00258

Shao, S., Shen, K., Wilder-Smith, E. P. V., & Li, X. (2011). Effect of pain perception on the heartbeat evoked potential. *Clinical Neurophysiology*, *122*(9), 1838–1845. https://doi.org/10.1016/j.clinph.2011.02.014

Shapiro, S. L., & Carlson, L. E. (2009). *The art and science of mindfulness: Integrating mindfulness into psychology and the helping professions* (pp. xvii, 194). American Psychological Association. https://doi.org/10.1037/11885-000

Shareef-Trudeau, L., Lydon-Staley, D. M., Medaglia, J., & Kucyi, A. (In press). Brain-based predictive modeling of mind-wandering and involuntary thought: The idiographic approach. *Technology, Mind and Behavior*.

Sherrington, C. S. (1906). *The integrative action of the nervous system* (pp. xvi, 411). Yale University Press. https://doi.org/10.1037/13798-000

Shi, L., Zheng, L., Jin, D., Lin, Z., Zhang, Q., & Zhang, M. (2022). Assessment of Combination of Automated Pupillometry and Heart Rate Variability to Detect Driving Fatigue. *Frontiers in Public Health*, *10*. https://doi.org/10.3389/fpubh.2022.828428

Smallwood, J., Margulies, D., Bernhardt, B. C., & Jefferies, E. (2018). Investigating the Elements of Thought: Toward a Component Process Account of Spontaneous Cognition. In K. Christoff & K. C. R. Fox (Eds.), *The Oxford Handbook of Spontaneous Thought: Mind-Wandering, Creativity, and Dreaming* (p. 0). Oxford University Press. https://doi.org/10.1093/oxfordhb/9780190464745.013.34

Speilberger, C. D., Gorsuch, Rl., Lushene, R., Vagg, P., & Jacobs, G. (1983). Manual for the state-trait anxiety inventory. *Palo Alto, CA: Consulting Psychologists*.

Spielberger, C. D., Gonzalez-Reigosa, F., Martinez-Urrutia, A., Natalicio, L. F. S., & Natalicio, D. S. (2017). The State-Trait Anxiety Inventory. *Revista Interamericana de Psicología/Interamerican Journal of Psychology*, *5*(3 & 4), Article 3 & 4. https://doi.org/10.30849/rip/ijp.v5i3

Spitzer, R. L., Kroenke, K., Williams, J. B. W., & Löwe, B. (2006). A Brief Measure for Assessing Generalized Anxiety Disorder: The GAD-7. *Archives of Internal Medicine*, *166*(10), 1092–1097. https://doi.org/10.1001/archinte.166.10.1092

Stan Development Team. (2024). *Stan Reference Manual*. Stan Development Team. https://mc-stan.org

Steinfath, P., Herzog, N., Fourcade, A., Sander, C., Nikulin, V., & Villringer, A. (2024). *Validating genuine changes in Heartbeat Evoked Potentials using Pseudotrials and Surrogate Procedures* (p. 2024.08.24.609348). bioRxiv. https://doi.org/10.1101/2024.08.24.609348

Stewart, S. H., Buffett-Jerrott, S. E., & Kokaram, R. (2001). Heartbeat awareness and heart rate reactivity in anxiety sensitivity: A further investigation. *Journal of Anxiety Disorders*, *15*(6), 535–553. https://doi.org/10.1016/S0887-6185(01)00080-9

Tuladhar, A. M., ter Huurne, N., Schoffelen, J.-M., Maris, E., Oostenveld, R., & Jensen, O.

(2007). Parieto-occipital sources account for the increase in alpha activity with working

memory load. *Human Brain Mapping*, *28*(8), 785–792.

https://doi.org/10.1002/hbm.20306

Vălenaş, S. P., Szentágotai-Tătar, A., Grafton, B., Notebaert, L., Miu, A. C., & MacLeod, C.

(2017). Prediction of pre-exam state anxiety from ruminative disposition: The mediating

role of impaired attentional disengagement from negative information. *Behaviour*

Research and Therapy, *91*, 102–110. https://doi.org/10.1016/j.brat.2017.01.014

Verdonk, C., Teed, A. R., White, E. J., Ren, X., Stewart, J. L., Paulus, M. P., & Khalsa, S. S.

(2024). Heartbeat-evoked neural response abnormalities in generalized anxiety disorder

during peripheral adrenergic stimulation. *Neuropsychopharmacology*.

https://doi.org/10.1038/s41386-024-01806-5

Vitasari, P., Wahab, M. N. A., Herawan, T., Othman, A., & Sinnadurai, S. K. (2011). Re-test of

State Trait Anxiety Inventory (STAI) among Engineering Students in Malaysia:

Reliability and Validity tests. *Procedia - Social and Behavioral Sciences*, *15*, 3843–3848.

https://doi.org/10.1016/j.sbspro.2011.04.383

Wenzler, S., Hagen, M., Tarvainen, M. P., Hilke, M., Ghirmai, N., Huthmacher, A.-C., Trettin,

M., van Dick, R., Reif, A., & Oertel-Knöchel, V. (2017). Intensified emotion perception

in depression: Differences in physiological arousal and subjective perceptions. *Psychiatry*

Research, *253*, 303–310. https://doi.org/10.1016/j.psychres.2017.03.040

Yamashita, S., Saitoh, T., Iguchi, K., Suwa, K., Ohtani, H., Saotome, M., & Maekawa, Y. (2024).

Electrocardiogram Electrode Positioning on the Back During Echocardiography: An

Exploratory Cross-Sectional Study. *Cureus*, *16*(4), e57967.

https://doi.org/10.7759/cureus.57967

Yuan, H., Yan, H.-M., Xu, X.-G., Han, F., & Yan, Q. (2008). Effect of heartbeat perception on

heartbeat evoked potential waves. *Neuroscience Bulletin*, *23*(6), 357–362.

https://doi.org/10.1007/s12264-007-0053-7

Zhao, C., Zhao, M., Liu, J., & Zheng, C. (2012). Electroencephalogram and electrocardiograph

assessment of mental fatigue in a driving simulator. *Accident Analysis & Prevention*, *45*,

83–90. https://doi.org/10.1016/j.aap.2011.11.019

Zoom Video Communications (Version 5.17.0). (2024). [Computer software]. https://zoom.us