Divergence of *germ cell-less* roles in germ line development across
insect species

Authors: Jonchee A. Kao[1], Ben Ewen-Campen[2,3], Cassandra G. Extavour[1,2,4]

[1]Department of Molecular and Cellular Biology, Harvard University, 16 Divinity Ave,
Cambridge, MA 02138, USA

[2]Department of Organismic and Evolutionary Biology, Harvard University, 16 Divinity Ave,
Cambridge, MA 02138, USA

[3]Current address: Department of Genetics, Blavatnik Institute, Harvard Medical School, Harvard
University, Boston, MA 02115, USA

[4]Howard Hughes Medical Institute, Chevy Chase, MD, USA

Corresponding author: Cassandra G. Extavour

Email: extavour@oeb.harvard.edu

Phone: (617) 496-1935

ORCIDs: JAK 0000-0003-1701-3265; BE-C 0000-0002-4326-3033, CGE 0000-0003-2922-5855

Keywords: *Oncopeltus fasciatus*; *Gryllus bimaculatus*; germ cell specification; *torso*; pole cells

Abstract

During development, sexually reproducing animals must specify and maintain the germ line, the

lineage of cells that gives rise to the next generation of animals. In the fruit fly *Drosophila*

melanogaster, *germ cell-less* (*gcl*) is required for the formation of primordial germ cells in the

form of cells that cellularize at the posterior pole of the embryo, called pole cells. Forming pole

cells is a mechanism of germ cell formation unique to a subset of insects. Even though most

animals do not form pole cells as primordial germ cells, *gcl* is conserved across Metazoa, raising

the question of how this conserved gene acquired its central role in the evolutionarily derived

process of pole cell formation. Here, we examine the functions of *gcl* in two other insects with

different modes of germ cell specification: the milkweed bug *Oncopeltus fasciatus* and the

cricket *Gryllus bimaculatus*. We found that *gcl* is involved in germ cell development, but not

strictly required for germ cell specification, in *O. fasciatus*, although it appears to function

through a mechanism different from that in *D. melanogaster*. In contrast, we could not detect any

impact on the embryonic germ line upon *gcl* knockdown in *G. bimaculatus*. This work serves as

a case study into how the roles of genes in the process of germ line development can change over

evolutionary time across animals.

Keywords: Germ cells; pole cells; *Oncopeltus fasciatus*; *Gryllus bimaculatus*; Hemimetabola;
milkweed bug; cricket; Torso

Highlights

• *germ cell-less* knockout reduces germ cell number in *Oncopeltus fasciatus*.
• *germ cell-less* knockdown does not affect germ cell number in *Gryllus bimaculatus*.
• The genetic mechanisms through which *gcl* affects germ cell number have diverged among insects.

Graphical abstract

Introduction

One of the most crucial processes in animal development is the specification and

maintenance of the germ line, the lineage of cells uniquely capable of giving rise to the next

generation of animals (Extavour and Akam, 2003). Despite the importance of this process,

mechanisms of germ line development vary widely even within closely related taxa, including

across insects (Extavour and Akam, 2003). For example, in the fruit fly *Drosophila*

melanogaster, germ cells first appear at the posterior pole of the embryo very early in

embryogenesis, before cellularization occurs in the rest of the embryo (Huettner, 1923). The

formation of germ cells in embryos of this fly depends on the presence of the germ plasm, a

specialized region of the cytoplasm at the posterior pole containing maternally synthesized

RNAs and proteins that collectively direct the formation and specification of germ cells

(Illmensee and Mahowald, 1974). In contrast, germ cells in the cricket *Gryllus bimaculatus* are

first detectable comparatively later in embryogenesis, after segmentation of the abdomen has

begun (Ewen-Campen et al., 2013a). There is no evidence of germ plasm in cricket embryos, and

germ cells are thought to be specified instead by inductive signals from neighboring cells,

including BMP signaling (Donoughe et al., 2014; Ewen-Campen et al., 2013a). In yet another

mechanism of insect germ cell specification, germ cells of the milkweed bug *Oncopeltus*

fasciatus appear at a stage intermediate between those of *D. melanogaster* and *G. bimaculatus*,

after blastoderm cellularization but before abdominal segmentation (Butt, 1949; Ewen-Campen

et al., 2013b). There is also no evidence of germ plasm in embryos of this bug (Ewen-Campen et

al., 2013b), but specific germ cell specification signals have not been identified.

Despite the variation in modes of germ line development across species, many genes

involved in this process are conserved. These include *vasa* and *piwi*, which are commonly used

as germ cell markers in many different animals (Ewen-Campen et al., 2010). *germ cell-less* (*gcl*)

is also sometimes used as a germ cell marker, but its role in germ line development is less well

understood (Jongens et al., 1992). Embryos laid by *D. melanogaster* females homozygous for a

loss of function mutation in *gcl* have significantly fewer germ cells than controls, if any (Jongens

et al., 1994, 1992). The gene is thought to have a role in suppressing somatic cell fate by

targeting the receptor tyrosine kinase Torso for degradation (Colonnetta et al., 2021; Pae et al.,

2017). *gcl* is also required for germ cell transcriptional quiescence in flies (Leatherman et al.,

2002). However, *gcl* is not required for germ cell specification in the mouse *Mus musculus* nor in

the flour beetle *Tribolium castaneum*, but it is required for proper mouse spermatogenesis

(Ansari et al., 2018; Kimura et al., 2003). Outside the germ line, *gcl* plays a role in establishing

the anterior–posterior axis in *T. castaneum* (Ansari et al., 2018). Despite the wide divergence in

their physiological roles, *gcl* homologs are highly conserved in amino acid sequence, with 45%

similarity between homologs from *D. melanogaster* and *M. musculus*. In addition, these

homologs appear to share at least some biochemical functions, since the mouse *gcl* homolog can

partially restore germ cell formation in a *gcl*-null *D. melanogaster* embryo, though not to wild

type numbers (Leatherman et al., 2000). *gcl* expression has also been observed in the embryonic

germ line in other arthropod species (Ewen-Campen et al., 2013a; Gupta and Extavour, 2013).

However, to our knowledge, functional tests of whether *gcl* is required for insect germ line

development or function remain limited to *D. melanogaster* and *T. castaneum*.

Here, we examine the function of *germ cell-less* in the embryonic and adult germ lines of

O. fasciatus and *G. bimaculatus*. Our data suggest that *gcl* is not required for germ cell

specification nor for adult gametogenesis in either species. However, *gcl* mutant *O. fasciatus*

embryos have significantly fewer germ cells than wild type. Knockdown of *Of-torso* does not

affect germ cell number, suggesting that *gcl* in this species affects germ cell number through a

different mechanism from that in *D. melanogaster*. Overall, these studies into *gcl* function across

species inform our understanding of the evolution of gene function in germ line development.

Materials and Methods

Gene identification

germ cell-less. *Of-germ cell-less* is annotated as OFAS015300 on Scaffold 2113 of the current *O.*

fasciatus genome assembly (Panfilio et al., 2019). *Gb-germ cell-less* is annotated as GBIM03474

on Scaffold 14 of the current *G. bimaculatus* genome assembly (Ylla et al., 2021). We amplified

both genes from mixed stage embryonic cDNA of the respective species to verify the full-length

coding sequences.

Of-torso. A tBLASTn search was performed on the *O. fasciatus* genome (Panfilio et al., 2019)

using *D. melanogaster* Torso as a query, and a reciprocal BLASTp was performed on the top

results with e-value < 1e–40. Of the initial results, OFAS005359 on Scaffold 336 returned *torso*

orthologs from other insects. A reciprocal BLASTp search using the adjacent upstream annotated

gene, OFAS005360, also returned *torso* orthologs from other insects. PCR was performed from

mixed stage embryonic cDNA to confirm that OFAS005359 and OFAS005360 correspond to the

same transcript. To obtain full-length sequence, RACE was performed using the SMARTer

RACE 5'/3' Kit (Takara Bio 634859) following manufacturer's instructions.

The initial tBLASTn search using *D. melanogaster* Torso as a query also returned

OFAS025175 as a result. A reciprocal BLASTp search using the predicted peptide sequence of

this hit returned *fgf receptor* orthologs from other insects. Since the FGF receptor is a family of

receptor tyrosine kinases closely related to Torso (Duncan et al., 2013), we wanted to include

OFAS025175 for phylogenetic analysis. Therefore, we performed RACE using the SMARTer

RACE 5'/3' Kit to obtain the full-length sequence of this gene. Full length sequences for both

Of-torso and *Of-fgfr* have been deposited at GenBank with accession numbers PV361543 and

PV361544, respectively.

Phylogenetic analysis was performed using MEGA 11 (Tamura et al., 2021). Insect

orthologs of Torso were aligned using MUSCLE (Edgar, 2004) along with orthologs of Ret

oncogene and FGF receptor, which are the closest related families of receptor tyrosine kinases

(Duncan et al., 2013). The multiple sequence alignment was used to generate a gene tree using

the Maximum Likelihood method with the JTT substitution model (Jones et al., 1992), a discrete

Gamma-distributed rate model and 1000 bootstrap replicates.

Molecular biology

RNA extraction and cDNA synthesis. Embryos or tissues of interest were homogenized in TRIzol

(Invitrogen 15596026), and RNA was extracted with phenol–chloroform extraction and

isopropanol precipitation. cDNA was synthesized with SuperScript III First-Strand Synthesis

SuperMix for qRT-PCR (Invitrogen 11752250) following manufacturer's instructions.

Cloning and in vitro *transcription*. Fragments of genes of interest were PCR-amplified from

embryonic cDNA and each cloned into the pGEM-T Easy vector (Promega A137A) using

manufacturer's instructions. A fragment of *gfp* was subcloned from an existing plasmid

containing the *gfp* sequence. Insertion was confirmed by Sanger sequencing. Primers used are

listed in Supplemental Table 1.

For *in situ* probe synthesis, gene fragments were PCR-amplified from plasmids using T7

and Sp6 primers. *In vitro* transcription was performed using T7 RNA polymerase (Promega

P2075) or Sp6 RNA polymerase (Roche 10810274001) with DIG RNA labeling mix (Roche

11277073910). 100 ng of template was used for a 20-µl reaction run at 37°C overnight. RNA

was purified with LiCl/ethanol precipitation, then resuspended in 20 µl DEPC-treated water.

Concentration was measured using a NanoDrop Spectrophotometer (ThermoFisher), and the

sample was diluted to 100 ng/µl with hybridization buffer (50% formamide, 1x phosphate-

buffered saline (PBS), 5x saline–sodium citrate (SSC), 0.1% Triton X-100, 0.1% CHAPS, 5 mM

EDTA, 50 µg/mL heparin, 1 µg/mL yeast tRNA). Probes were stored at −20°C.

For double-stranded RNA (dsRNA) synthesis, gene fragments were PCR-amplified from

plasmids using a forward T7 primer and a reverse primer containing both T7 and Sp6 promoter

sequences to generate a template for *in vitro* transcription with the T7 promoter sequence on both

ends. *In vitro* transcription was performed using the MEGAscript T7 Transcription Kit

(Invitrogen AM13345). 100 ng of template was used for an 80-µl reaction run at 37°C overnight.

RNA was purified with LiCl/ethanol precipitation, then resuspended in 100 µl DEPC-treated

water. To anneal the two strands, this solution was heated to 95°C on a heat block for 5 minutes,

then allowed to cool slowly by leaving the tube on the heat block with the power switched off. A

1:10 dilution of annealed RNA was analyzed using a NanoDrop Spectrophotometer

(ThermoFisher) to measure its concentration and purity, then run on a 1% agarose gel to confirm

its size and integrity. For embryonic RNAi, dsRNA was diluted in DEPC-treated water to 2 µg/µl

(*O. fasciatus*) or 5 µg/µl (*G. bimaculatus*), with 0.05% phenol red. For *G. bimaculatus* adult

RNAi, dsRNA was diluted to 7.5 µg/µl in DEPC-treated water.

CRISPR/Cas9-mediated mutagenesis. Alt-R S.p. Cas9 Nuclease V3 (cat. #1081059) and Alt-R

CRISPR-Cas9 tracrRNA (cat. #1072533) were ordered from IDT, along with gene-specific Alt-R

CRISPR-Cas9 crRNA. tracrRNA and crRNA were resuspended to 100 µM in Nuclease-Free

Duplex Buffer (IDT cat. #11-01-03-01). 5 µl of each RNA were combined in a PCR tube, heated

to 95°C for 5 minutes, and allowed to cool to room temperature to form 50 µM gRNA duplex. 3

µl gRNA and 2 µl Cas9 protein (10 µg/µl or ~62 µM) were combined in a PCR tube and

incubated at room temperature for 10 minutes to form Cas9/gRNA ribonucleoprotein complex

(RNP) at roughly 25 µM. Finally, RNP was diluted tenfold in 1x PBS with 0.05% phenol red.

Of-Vasa antibody generation

A rabbit polyclonal antibody was raised against a recombinant _Of_-Vasa protein fragment

(McGill Biology CIAN facility, Canada). The sequence encoding the 565 C-terminal amino acids

of _Of_-Vasa, the stop codon and the first 96 base pairs of 3' UTR (GenBank accession KC261571

(Ewen-Campen et al., 2013b)) was cloned into the pET151/dTOPO expression vector (Invitrogen

K151-01), thus introducing an N-terminal 6x-polyhistidine tag. Protein expression was induced

in _E. coli_ BL21 (DE3) by addition of IPTG to a final concentration of 0.25 mM and shaking at

30°C for four hours. The overexpressed proteins were highly insoluble, so the cell pellets were

disrupted by sonication in non-denaturing buffer, and the soluble fraction was separated by

centrifugation and discarded. The pellet was resuspended in 8 M urea (crude prep). Crude preps

were further purified by SDS gel-purification. Protein bands were excised from the gel after

reverse staining with zinc chloride and the proteins were collected by electroelution. Acetone

precipitation was performed, and the precipitated protein was washed and resuspended in 8 M

urea. Protein concentration was adjusted to 1–2 mg/mL for injection. Rabbits were injected using

a standard 80-day protocol with three boosts after the initial injection; the final bleed was

performed after 87 days. The serum was processed by addition of sodium azide to 0.02% (w/v).

The antibody was affinity-purified against bacterially expressed *Of*-Vasa protein.

Embryo microinjection

Needles for injection were made by pulling glass capillaries (World Precision Instruments

1B100-4) with a Sutter P-97 Flaming/Brown micropipette puller. Needle tips were manually

opened with fine forceps just prior to injection. Eggs were injected with a Narishige IM300

microinjector.

O. fasciatus were allowed to lay eggs in dry cotton for four hours, after which eggs were

picked out of the cotton for injection. Two glass microscope slides were taped together with

double-sided tape to create a ledge with exposed double-sided tape against which embryos were

lined up. Eggs were then submerged in tap water for at least 20 minutes to soften the eggshell

before injection. Slides with injected eggs were kept at 28°C to continue development until the

desired age.

G. bimaculatus were allowed to lay eggs in wet cotton for four hours, after which eggs

were washed out of the cotton for injection. Eggs were aligned in grooves (570 µm wide × 2930

µm long × 650 µm deep) made in 2% agarose in Milli-Q water using a custom mold as

previously described (Donoughe et al., 2018). Eggs were injected while submerged in 1x PBS.

After injection, the solution was replaced with fresh 1x PBS. Injected eggs were kept submerged

in 1x PBS at 28°C overnight. The following day, dead embryos, if any, were removed, and the

remaining eggs were transferred onto damp Kimwipes in a covered Petri dish kept at 28°C to

continue development. Three days after injection, eggs were submerged in tap water again and

monitored every day until they reached the desired stage for dissection.

Embryo dissection and fixation

O. fasciatus eggs were stuck to the bottom of a 60-mm Petri dish with double-sided tape

and a slit was created in the eggshell with a microscalpel. 1x PBS with 4% paraformaldehyde

(PFA) was then poured into the dish to fix the embryos for 30 minutes at room temperature,

followed by at least three washes with PBT (1x PBS with 0.1% Triton X-100). Embryos were

then dissected free of their yolk in PBT with fine forceps. Embryos were washed at least three

times with PBT at room temperature before continuing with staining or dehydration in methanol.

G. bimaculatus embryos were dissected free of their yolk in 1x PBS with fine forceps and

fixed overnight in 1x PBS with 4% PFA at 4°C. The following morning, they were washed at

least three times with PBT at room temperature before continuing with staining or dehydration in

methanol.

G. bimaculatus adult RNA interference (RNAi)

2 µl dsRNA was injected into the abdomen of an adult cricket with a Hamilton syringe

through the base of a metathoracic leg. Gonads were dissected seven days after injection for

analysis.

Colorimetric in situ hybridization

Dissected embryos were stored in methanol at −20°C overnight or longer before *in situ*

hybridization. After rehydration in PBT and washing in hybridization buffer, embryos were

hybridized overnight in a 70°C water bath with 1 ng/µl probe in hybridization buffer. The

following day, the following series of 20-minute washes were performed at 70°C: once in

hybridization buffer, three times in wash 1 (2x SSC, 0.1% CHAPS) and three times in wash 2

(0.2x SSC, 0.1% CHAPS). Embryos were then washed three times for ten minutes each at room

temperature in KTBT (1x Tris-buffered saline with KCl and 0.1% Triton X-100) followed by an

hour-long incubation at room temperature in *in situ* blocking solution (1x PBS, 0.1% Tween-20,

5% NGS, 2 mg/mL bovine serum albumin, 1% dimethyl sulfoxide). Embryos were incubated for

two hours at room temperature with sheep anti-digoxigenin–AP (Roche 11093274910), diluted

1:2000 in *in situ* blocking solution, then washed at least four times over two hours with KTBT

and incubated in KTBT overnight at 4°C. The following day, embryos were washed with NTMT

(100 mM Tris pH 9.5, 100 mM NaCl, 50 mM $MgCl_2$, 0.1% Tween-20), then developed in the

dark at 37°C with NBT/BCIP in NTMT with 5% polyvinyl alcohol. Development was monitored

at least every hour under a dissection microscope. To stop development of the enzymatic

reaction, embryos were washed in PBT, incubated with 5 ng/µl Hoechst 33342 in PBT at room

temperature for an hour, post-fixed in 4% PFA in PBS, then incubated in 70% glycerol before

mounting in 100% glycerol.

Immunohistochemistry

Embryos were incubated in blocking solution (PBT with 5% normal goat serum (Jackson

ImmunoResearch 005-000-121)) for one hour at room temperature, then incubated overnight at

4°C with primary antibodies diluted in blocking solution. The following day, embryos were

washed for at least four times over four hours with PBT, incubated for an hour in blocking

solution, then incubated overnight at 4°C with secondary antibodies diluted in blocking solution.

Finally, embryos were washed in PBT, post-fixed in 4% PFA in PBS, then incubated in 70%

glycerol before mounting in glycerol. Antibody staining was also performed with a no-primary

control.

Primary antibodies used were: rabbit anti-*Gb*-Piwi (Ewen-Campen et al., 2013a) at a final

concentration of 1:300; rabbit anti-*Of*-Vasa at 1:100; mouse anti-α-tubulin (DM1A; Sigma

T9026) at 1:100; rabbit anti-β-catenin (Sigma C2206) at 1:500; and mouse anti-RNA polymerase

II (H5; Covance MMS-129R) (Bregman et al., 1995; Warren et al., 1992) at 1:100. Secondary

antibodies used were goat anti-rabbit conjugated to Alexa Fluor 647 (Invitrogen A-21246), goat

anti-rabbit Alexa Fluor 488 (Invitrogen A-11008), goat anti-rabbit Alexa Fluor 568 (Invitrogen

A-11011), donkey anti-mouse Alexa Fluor 488 (Invitrogen A-21202), and goat anti-mouse Alexa

Fluor 647 (Invitrogen A-21236). All secondary antibodies were used at a final concentration of

1:400 along with Hoechst 33342 (Sigma B2261) at 5 ng/µl.

Embryos were imaged on a Nikon CSU-W1 spinning disk inverted confocal microscope

with 10x air or 20x air objectives or a Zeiss AxioObserver inverted compound microscope with

10x air or 20x air objectives. Images were analyzed in Fiji.

Quantitative RT-PCR

Quantitative RT-PCR was performed on an AriaMx Real-time PCR system (Agilent)

using PerfeCTa SYBR Green SuperMix, Low ROX (Quantabio 95056). Fold-change differences

were calculated using the $\Delta\Delta C_t$ method for RNAi experiments. Expression levels were

normalized against *Of-gapdh* or *Gb-β-tubulin*. All primers used are listed in Supplemental Table

1.

Results.

germ cell-less expression across insect species.

During oogenesis in *D. melanogaster*, Gcl protein is detectable at the nuclear membrane

of nurse cells and oocytes (Moore et al., 2009). In addition, *gcl* transcripts are transcribed by

nurse cells, transported into the oocyte and deposited at the oocyte posterior, where they remain

as part of the germ plasm into early embryogenesis (Jongens et al., 1992). These transcripts are

translated into protein starting at the cleavage stages of embryogenesis (Jongens et al., 1992).

Gcl protein is localized to the nuclear membranes of germ cells after they form and remains

detectable in germ cells until mid-gastrulation, when germ cells enter the primordial midgut

(Jongens et al., 1992). We asked whether *gcl* is also expressed in other insect embryos at the time

and place of germ cell specification.

We (Ewen-Campen 2013) and others (Butt 1949) have qualitatively described the first

appearance of primordial germ cells in blastoderm stage *O. fasciatus* embryos, and their

proliferation and migration during embryogenesis. However, to our knowledge there has been no

embryonic developmental staging system established for this insect. We therefore we defined

stages based on major embryological, morphogenetic and morphological milestones, for

describing embryonic development from egg-laying through to katatrepsis, with special

emphasis on the development of germ cells (Supplemental Figure 1; Supplemental Table 2). We

assessed *gcl* transcript levels using quantitative RT-PCR on *O. fasciatus* embryos collected in 12-

hour windows spanning the first four days of embryogenesis. *Of-gcl* transcripts were detected at

the highest levels 24–36 hours after egg-lay at 28°C (Figure 1A). This window corresponds to

the late blastoderm stage (stage 3), when germ cells are first distinguishable (Ewen-Campen et

al., 2013b), and the beginning of germ band elongation (stages 4–5). We speculate that this peak

in expression is likely due to zygotic rather than maternal expression, since it is higher than

expression in freshly laid eggs. However, we were unable to detect *Of-gcl* transcripts by *in situ*

hybridization above background levels in embryos at any stage of development, nor in adult

ovaries or testes (Supplemental Figure 2A–B).

We similarly assessed *gcl* transcript levels using qRT-PCR on *G. bimaculatus* embryos

collected in 8-hour windows spanning the first two days of embryogenesis and 24-hour windows

spanning the following eight days of embryogenesis. *Gb-gcl* transcripts were detectable at the

highest levels in the first eight hours after egg-lay (Figure 1B), suggesting maternal deposition of

Gb-gcl transcript. Consistent with this hypothesis, *Gb-gcl* transcript was detectable by *in situ*

hybridization in ovaries but not during embryogenesis (Supplemental Figure 2C–D).

Although we were unable to detect localized *gcl* expression in either *O. fasciatus* or *G.*

bimaculatus embryos, we still sought to determine whether the gene's documented roles in germ

cell specification (*D. melanogaster*) or adult spermatogenesis (*M. musculus*) might be conserved

(Jongens et al., 1992; Kimura et al., 2003). To this end, we analyzed *gcl* loss of function

conditions in embryos and adult gonads of both species. *gcl* is also required for transcriptional

quiescence in germ cells in *D. melanogaster* (Leatherman et al., 2002). However, we could not

detect any reduction in transcriptional activity in germ cells relative to somatic cells of either *O.*

fasciatus or *G. bimaculatus* (Supplemental Figure 3), so we did not assess this phenotype in our

functional studies of *gcl*.

Of-gcl mutation decreases germ cell number in milkweed bug embryos.

To determine whether *gcl* is required for germ cell specification in *O. fasciatus*, we generated an *Of-gcl* putative protein-null allele with CRISPR/Cas9-mediated mutagenesis (Figure 2A–B). This allele contains a 2-bp insertion at amino acid residue 14. The resulting frameshift leads to a premature stop codon, resulting in a predicted peptide of 30 amino acid residues, compared with the 478-residue wild type protein. Surprisingly, individuals homozygous for this mutant allele were viable and fertile, and we have maintained a stable mutant line for at least twelve generations at the time of writing. There were no detectable morphological defects in adult ovaries or testes from this mutant line, nor any evidence of abnormal gametogenesis in either males or females (Supplemental Figures 4–5).

In *O. fasciatus*, germ cells are first detectable in stage 3 on the interior of the cellular blastoderm, forming part of an invaginating structure called the posterior pit (Ewen-Campen et al., 2013b). The germ cells remain at the posterior end of the germ band as it forms by invaginating into the egg (Ewen-Campen et al., 2013b) (Figure 2B; Supplemental Figure 1). At stage 3, *gcl* mutant embryos had on average 35% fewer germ cells than wild type embryos ($p < 0.001$; $n = 14$ wild type, 8 *gcl* embryos; Figure 2C–D; Supplemental Table 3). Germ cell number remained significantly lower in *gcl* mutant embryos relative to wild type embryos throughout all embryonic stages examined (Figure 2D). We detected a marginal decrease in germ cell proliferation in *gcl* mutant embryos relative to wild type only at stages 6 and 9, as measured by the proportion of germ cells with detectable phospho-histone 3 (Figure 2E; Supplemental Table 3). However, there was no detectable difference in germ cell proliferation between the two genotypes at any other embryonic stage examined (Figure 2E; Supplemental Table 3). We were

also unable to detect cleaved caspase 3-dependent cell death in stages 3–4 of either genotype ($n =$

27 wild type, 23 *gcl* embryos).

It is formally possible that *gcl* loss of function affects the expression of Vasa protein,

which we use as an experimental proxy for identifying germ cells. However, we noted that the

Vasa-positive cells in *gcl* mutant embryos expressed Vasa at comparable levels to wild type

embryos. Thus, while we cannot rule out the possibility that germ cells that do not express

detectable Vasa may be present in our *gcl* loss of function embryos, we believe it unlikely that

lack of *gcl* would eliminate Vasa expression in some germ cells without significantly affecting

expression in others. Therefore, we favor the interpretation that the reduction in Vasa-positive

cells in *gcl* mutant embryos reflects a reduction in germ cell number. Given that the discrepancy

in germ cell number did not appear to be the result of differences in cell proliferation or cell

death, these data suggest that *Of-gcl* is involved in, but not strictly necessary for, one or more of

germ cell specification, formation or maintenance.

In *D. melanogaster*, Gcl promotes germ cell fate by targeting the receptor Torso for

degradation, and *torso* RNAi in *gcl* mutant flies restores germ cell numbers to wild type levels

(Pae et al., 2017). To determine whether this mechanism of germ cell number determination is

conserved in *O. fasciatus*, we asked whether *torso* loss of function also affects germ cell number

in this species. Despite previous reports that the *O. fasciatus* genome lacks a *torso* ortholog

(Skelly et al., 2019; Weisbrod et al., 2013), we identified a high-confidence homolog in the

published genome assembly (Panfilio et al., 2019) (Figure 3A). Knockdown of *Of-torso* by

embryonic RNAi did not affect germ cell number in stages 3–4 in either wild type or *gcl* mutant

backgrounds (Figure 3B–C). qPCR analysis of injected embryos confirmed that RNAi was

effective at reducing *torso* transcript levels to 50% relative to controls (Supplemental Figure 6A).

This result suggests that the mechanism through which *gcl* controls germ cell number is not

conserved between *O. fasciatus* and *D. melanogaster*.

Gb-gcl knockdown does not affect germ cell number in cricket embryos.

To determine whether *gcl* is required for germ cell specification in *G. bimaculatus*, we

knocked down *gcl* expression using embryonic RNAi. We found no significant difference in total

germ cell number in stage 8 embryos under *gcl* knockdown relative to *gfp* controls, nor did we

detect any significant difference in germ cell number in any segment (Figure 4). qPCR analysis

of injected embryos confirmed that RNAi was effective at reducing *gcl* transcript levels to 34%

relative to controls (Supplemental Figure 6B). It is possible that knockdown efficiency is not

high enough with RNAi to affect germ cell specification. However, we note that we have

previously detected significant changes in germ cell number with comparable or lower

knockdown efficiencies (Barnett et al., 2019; Donoughe et al., 2014; Nakamura and Extavour,

2016). Therefore, we speculate that the role of *Gb-gcl* in germ cell specification or maintenance,

if any, must be subtle.

Next, we asked whether *gcl* has any role in adult gametogenesis in *G. bimaculatus* by

performing RNAi in adult crickets. We could not detect any morphological or gametogenesis

defects in either ovaries or testes (Supplemental Figures 4-5). These data suggest that *gcl* is not

required for adult gametogenesis in this cricket.

Discussion.

Our analysis of *gcl* loss of function in embryos of two different hemimetabolous insects

suggests that *gcl* is not required for germ cell specification in these species, in contrast to the

gene's importance in germ cell formation in *D. melanogaster*. However, *Of-gcl* knockout

embryos have significantly fewer germ cells than wild type embryos, suggesting that the gene

may be involved in germ cell specification or maintenance in this bug, even if it is not strictly

required.

Gcl protein suppresses the somatic cell fate in *D. melanogaster* by targeting the receptor

tyrosine kinase Torso for degradation (Colonnetta et al., 2021; Pae et al., 2017). Germ cell

number can thus be modulated in flies by up- or downregulating Torso activity (Pae et al., 2017).

However, RNAi knockdown of *torso* in *O. fasciatus* does not affect germ cell number in either

wild type or *gcl* mutant embryos (Figure 3). Therefore, we hypothesize that germ cell

specification in this species does not require downregulation of Torso by Gcl, and that *Of-gcl*

regulates germ cell specification through some other unknown pathway. Torso's role in germ cell

specification may have been co-opted from its ancestral functions in neuroendocrine signaling,

just as it is thought to have been co-opted into its role in terminal patterning in flies (Duncan et

al., 2013; Schoppmeier and Schröder, 2005).

In *D. melanogaster*, the interaction between Gcl and Torso proteins depends on sequence

motifs present on both proteins (Pae et al., 2017). In Gcl protein, this motif is the GCL domain,

which is conserved across insect homologs (Supplemental Figure 7) (Pae et al., 2017). Torso

interacts with Gcl through a motif in the linker domain between its split kinase domains. This

linker region is significantly longer in *Dm*-Torso compared with other insect homologs

(Supplemental Figure 8), and the specific motif identified by Pae and colleagues (Pae et al.,

2017) is not present in other homologs. The conservation of the GCL domain despite the lack of

conservation in the Torso interaction motif may suggest that Gcl has some other unknown

interaction partner, or that the Gcl–Torso interaction may not be as sequence-specific as

originally thought. AlphaFold3 (Abramson et al., 2024) predicts potential interaction between

Gcl and Torso homologs in both *O. fasciatus* and *G. bimaculatus* through the split kinase linker

domain of Torso (Supplemental Figure 9). Further experiments will be needed to test whether

these proteins interact *in vivo*, and to identify additional conserved interaction partners of Gcl.

In mice, *gcl* knockout results in abnormal nuclear morphology in the liver, pancreas and

testis, though the only functional defect identified was impaired spermatogenesis (Kimura et al.,

2003). Similarly, *gcl* is expressed at low levels in somatic tissues in flies including the gut,

muscle and nervous system, but no defects have been reported in those tissues in *gcl*-null flies

(Jongens et al., 1992; Robertson et al., 1999). We were also unable to detect any defects in

spermatogenesis under either *Of-gcl* knockout or *Gb-gcl* knockdown conditions (Supplemental

Figure 4). Given that the mouse *gcl* homolog can rescue germ cell formation in *gcl*-null flies

(Leatherman et al., 2000), its biochemical functions are likely to be conserved across these

species. We speculate that *gcl* has a ubiquitous but subtle cell biological role in maintaining

nuclear morphology that only yields a phenotype in certain tissue contexts, such as in mouse

spermatogenesis. Under this hypothesis, *gcl* then gained an additional role in germ cell

specification in certain lineages of insects. In addition, the mechanism underlying this germ cell

specification has diverged between insect lineages, perhaps through the evolution of new

interaction partners such as Torso. This model reflects the complexity and variation present in the

mechanisms of germ cell specification and their evolution, including the co-option of new genes

with the shifting mechanisms of this process across animals.

Author contributions

JAK: Conceptualization, Methodology, Formal Analysis, Investigation, Writing – Original Draft, Writing – Review & Editing, Visualization; BE-C: Resources; CGE: Conceptualization, Writing – Review & Editing, Supervision, Funding Acquisition

Acknowledgments

The authors thank members of the Extavour lab for valuable discussions. This research was supported by National Science Foundation (NSF) Awards IOS-2220747 and IOS-0817678 to CGE, and an NSF Graduate Research Fellowship to BE-C. CGE is a Howard Hughes Medical Institute Investigator.

Figure 1. *gcl* expression is detectable by qPCR in *O. fasciatus* (A) and *G. bimaculatus* (B)
embryos. C_q values were normalized against *Of-gapdh* or *Gb-tubulin* expression and averaged
over four biological replicates for each time point. Error bars represent standard deviations. The
width of each bar represents the length of time for each embryo collection.

439
440
441
442

444   **(single column width)**

A

Of-Gcl

73   173 181   259   348 387

BTB   BACK   GCL   478

mutant ——— 30

B

allele	DNA sequence	translation
wild type	...CTGCCC<u>AG--A GCGTGTTGCTGCAACTCTC</u>...	...YCPERVAATL...
mutant	**...CTGCCCAGCCA** **GCGTGTTGCTGCAACTCTC...**	**...YCPASVLLQL...**

C

wild type | *gcl*

Of-Vasa

D

(total germ cells vs stage)

E

(PH3+ germ cells vs stage)

● wild type   ● *gcl* mutant

Figure 2 (previous page). *Of-gcl* **mutation decreases germ cell number in embryos.**

(**A**) Predicted domain architecture of *Of*-Gcl and predicted mutant protein. Wild type *Of*-Gcl has two predicted folded domains (BTB and BACK domains) as well as a conserved motif called the GCL domain. The predicted mutant protein contains none of these domains and is truncated after residue 30.

(**B**) *Of-gcl* mutant allele contains a two base pair insertion, creating a frameshift and leading to a predicted truncated protein. Guide sequence is underlined; arrowhead marks predicted Cas9 cut site. Inserted bases and predicted mutated residues are indicated in magenta.

(**C**) Representative antibody stains against *Of*-Vasa to mark germ cells in stage 3 wild type and *gcl* mutant embryos. Mutant embryos still have germ cells at the expected location, indicated by arrows. Scale bar = 100 µm.

(**D**) Germ cell counts across embryogenesis in wild type and *gcl* mutant embryos. Mutant embryos have significantly fewer germ cells per embryo compared to wild type.

(**E**) Proportion of PH3$^+$ germ cells across embryogenesis in wild type and *gcl* mutant embryos. There is no significant difference in the proportion of proliferating germ cells between wild type and *gcl* mutant embryos.

All statistical tests are non-parametric bootstrap comparisons. Embryonic stages are defined in Supplemental Figure 1 and Supplemental Table 3. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

(1.5-column width)

A

73 — fgfr|Schistocerca gregaria|isoform X1|XP 049864650.1	***FGF receptor***
76 — fgfr|Camponotus floridanus|isoform X1|XP 011254079.1	
60 — fgfr|Tribolium castaneum|isoform X1|XP 008190747.1	
50 — fgfr|Oncopeltus fasciatus|OFAS025175 OFAS016138	
98 — fgfr|Halyomorpha halys|XP 014289518.1	
100 — heartless|Drosophila melanogaster|isoform A|NP 732286.1	
100 — breathless|Drosophila melanogaster|isoform A|NP 729956.1	
fgfr|Bombyx mori|BAE94421.1	

Ret

Ret|Niloparvata lugens|XP 039288327.1
100 — Ret|Drosophila melanogaster|isoform A|NP 477044.1
68 — Ret|Tribolium castaneum|XP 015839320.2
61 — Ret|Camponotus floridanus|isoform X1|XP 025270917.1
85 — Ret|Cryptotermes secundus|XP 023720364.1
99 — Ret|Gryllus bimaculatus|GLH10125.1

torso

torso|Drosophila melanogaster|isoform A|NP 476762.1
torso|Bombyx mori|ACZ81657.1
100 — torso|Halyomorpha halys|XP 014279301.1
33 — torso|Oncopeltus fasciatus|OFAS005359 OFAS005360
92 — torso|Tribolium castaneum|EFA07525.1
55 — torso|Athalia rosae|isoform X1|XP 048509519.1
47 — torso|Gryllus bimaculatus|GLH08284.1
88 — torso|Cryptotermes secundus|isoform X1|XP 033610372.1

1.00

B

C

Figure 3 (previous page). *Of-torso* **knockdown does not affect embryonic germ cell number.**

(**A**) Phylogenetic analysis of *torso* and related receptor tyrosine kinases (*Ret oncogene* and *FGF receptor*) across insects. Each clade is individually highlighted for clarity. The putative *torso* homolog from *O. fasciatus* (OFAS005359/OFAS0005360) falls within the *torso* clade.

(**B**) Germ cell counts in stage 3–4 *Of-torso* RNAi embryos, performed in wild type and *gcl* mutant backgrounds. *torso* knockdown does not affect germ cell number in either background.

(**C**) Proportion of PH3$^+$ germ cells in stage 3–4 *Of-torso* RNAi embryos, performed in wild type and *gcl* mutant backgrounds. *torso* knockdown marginally reduces germ cell proliferation in *gcl* mutant embryos but not in wild type embryos.

All statistical tests are non-parametric bootstrap comparisons.
* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

(full page width)

A

Gb-Gcl
75   175   183   261   349   388
            BTB   BACK   GCL   498
dsRNA

B

| control | *Gb-gcl* RNAi |

Gb-Piwi

C

germ cells per embryo

n.s.

ctrl   *gcl*

n =   46   68

D

germ cells per hemisegment

n.s.   *n.s.*   *n.s.*

A2   A3   A4

n =   92   136   92   136   92   136

Figure 4 (previous page). *Gb-gcl* **knockdown does not affect embryonic germ cell number.**

(**A**) Predicted domain architecture of *Gb*-Gcl. *Gb*-Gcl has two predicted folded domains (BTB and BACK domains) and a conserved motif called the GCL domain. The fragment region used for dsRNA synthesis is marked.

(**B**) Representative antibody stains against *Gb*-Piwi to mark germ cells in stage 8 control and *gcl* knockdown embryos. Embryos in which *Gb-gcl* has been knocked down still have germ cells in the expected positions. Scale bar = 100 µm.

(**C**) Germ cell counts in stage 8 *Gb-gcl* RNAi embryos. Germ cell numbers are not significantly different in *Gb-gcl* knockdown embryos relative to controls. Each point represents one embryo.

(**D**) Germ cell counts in stage 8 *Gb-gcl* RNAi embryos, divided into hemisegments (*i.e.*, the left or right half of one segment). *Gb-gcl* knockdown does not affect the number of germ cells in any individual segment. Each point represents a hemisegment.

All statistical tests are non-parametric bootstrap comparisons.

(1.5 column width)

References

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A.J., Bambrick, J., Bodenstein, S.W., Evans, D.A., Hung, C.-C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A., Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A., Cherepanov, A., Congreve, M., Cowen-Rivers, A.I., Cowie, A., Figurnov, M., Fuchs, F.B., Gladman, H., Jain, R., Khan, Y.A., Low, C.M.R., Perlin, K., Potapenko, A., Savy, P., Singh, S., Stecula, A., Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E.D., Zielinski, M., Bapst, V., Kohli, P., Jaderberg, M., Hassabis, D., Jumper, J.M., 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. Nature 630, 493–500. https://doi.org/10.1038/s41586-024-07487-w

Ansari, S., Troelenberg, N., Dao, V.A., Richter, T., Bucher, G., Klingler, M., 2018. Double abdomen in a short-germ insect: Zygotic control of axis formation revealed in the beetle,Tribolium castaneum. Proc. Natl. Acad. Sci. 115, 1819–1824. https://doi.org/10.1073/pnas.1716512115

Barnett, A.A., Nakamura, T., Extavour, C.G., 2019. Hox genes limit germ cell formation in the short germ insect,Gryllus bimaculatus. Proc. Natl. Acad. Sci. 116, 16430–16435. https://doi.org/10.1073/pnas.1816024116

Bregman, D.B., Du, L., Zee, S. van der, Warren, S.L., 1995. Transcription-dependent redistribution of the large subunit of RNA polymerase II to discrete nuclear domains. J. cell Biol. 129, 287–298. https://doi.org/10.1083/jcb.129.2.287

Butt, F.H., 1949. Embryology of the Milkweed Bug, Oncopeltus fasciatus (Hemiptera). Cornell Experiment Station Memoir 3–43.

Colonnetta, M.M., Lym, L.R., Wilkins, L., Kappes, G., Castro, E.A., Ryder, P.V., Schedl, P., Lerit, D.A., Deshpande, G., 2021. Antagonism between,germ cell-less,and Torso receptor regulates transcriptional quiescence underlying germline/soma distinction. eLife 10, e54346. https://doi.org/10.7554/elife.54346

Donoughe, S., Kim, C., Extavour, C.G., 2018. High-throughput live-imaging of embryos in microwell arrays using a modular specimen mounting system. Biol. Open 7, bio031260. https://doi.org/10.1242/bio.031260

Donoughe, S., Nakamura, T., Ewen-Campen, B., Green, D.A., Henderson, L., Extavour, C.G., 2014. BMP signaling is required for the generation of primordial germ cells in an insect. Proc. Natl. Acad. Sci. 111, 4133–4138. https://doi.org/10.1073/pnas.1400525111

Duncan, E.J., Benton, M.A., Dearden, P.K., 2013. Canonical terminal patterning is an evolutionary novelty. Dev. Biol. 377, 245–261. https://doi.org/10.1016/j.ydbio.2013.02.010

Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res. 32, 1792–1797. https://doi.org/10.1093/nar/gkh340

Ewen-Campen, B., Donoughe, S., Clarke, D.N., Extavour, C.G., 2013a. Germ cell specification requires zygotic mechanisms rather than germ plasm in a basally branching insect. Curr. Biol. 23, 835–842. https://doi.org/10.1016/j.cub.2013.03.063

Ewen-Campen, B., Jones, T.E.M., Extavour, C.G., 2013b. Evidence against a germ plasm in the milkweed bug,Oncopeltus fasciatus, a hemimetabolous insect. Biol. Open 2, 556–568. https://doi.org/10.1242/bio.20134390

Ewen-Campen, B., Schwager, E.E., Extavour, C.G., 2010. The molecular machinery of germ line specification. Mol. Reprod. Dev. 77, 3–18. https://doi.org/10.1002/mrd.21091

Extavour, C.G., Akam, M., 2003. Mechanisms of germ cell specification across the metazoans: epigenesis and preformation. Development 130, 5869–5884. https://doi.org/10.1242/dev.00804

Gupta, T., Extavour, C.G., 2013. Identification of a putative germ plasm in the amphipod,Parhyale hawaiensis. EvoDevo 4, 34. https://doi.org/10.1186/2041-9139-4-34

Huettner, A.F., 1923. The origin of the germ cells in,Drosophila melanogaster. J. Morphol. 37, 385–423. https://doi.org/10.1002/jmor.1050370204

Illmensee, K., Mahowald, A.P., 1974. Transplantation of posterior polar plasm in,Drosophila. Induction of germ cells at the anterior pole of the egg. Proc. Natl. Acad. Sci. 71, 1016–1020. https://doi.org/10.1073/pnas.71.4.1016

Jones, D.T., Taylor, W.R., Thornton, J.M., 1992. The rapid generation of mutation data matrices from protein sequences. Bioinformatics 8, 275–282. https://doi.org/10.1093/bioinformatics/8.3.275

Jongens, T.A., Ackerman, L.D., Swedlow, J.R., Jan, L.Y., Jan, Y.N., 1994. Germ cell-less encodes a cell type-specific nuclear pore-associated protein and functions early in the germ-cell specification pathway of Drosophila. Genes Dev. 8, 2123–2136. https://doi.org/10.1101/gad.8.18.2123

Jongens, T.A., Hay, B., Jan, L.Y., Jan, Y.N., 1992. The germ cell-less,gene product: A posteriorly localized component necessary for germ cell development in,Drosophila. Cell 70, 569–584. https://doi.org/10.1016/0092-8674(92)90427-e

Kimura, T., Ito, C., Watanabe, S., Takahashi, T., Ikawa, M., Yomogida, K., Fujita, Y., Ikeuchi, M., Asada, N., Matsumiya, K., Okuyama, A., Okabe, M., Toshimori, K., Nakano, T., 2003. Mouse Germ cell-less as an essential component for nuclear integrity. Mol. Cell. Biol. 23, 1304–1315. https://doi.org/10.1128/mcb.23.4.1304-1315.2003

Leatherman, J.L., Kaestner, K.H., Jongens, T.A., 2000. Identification of a mouse,germ cell-less,homologue with conserved activity in,Drosophila. Mech. Dev. 92, 145–153. https://doi.org/10.1016/s0925-4773(99)00335-4

Leatherman, J.L., Levin, L., Boero, J., Jongens, T.A., 2002. germ cell-less,acts to repress transcription during the establishment of the Drosophila germ cell lineage. Curr. Biol. 12, 1681–1685. https://doi.org/10.1016/s0960-9822(02)01182-x

Moore, J., Han, H., Lasko, P., 2009. Bruno negatively regulates,germ cell-less,expression in a BRE-independent manner. Mech. Dev. 126, 503–516. https://doi.org/10.1016/j.mod.2009.04.002

Nakamura, T., Extavour, C.G., 2016. The transcriptional repressor Blimp-1 acts downstream of BMP signaling to generate primordial germ cells in the cricket,Gryllus bimaculatus. Development 143, 255–263. https://doi.org/10.1242/dev.127563

Pae, J., Cinalli, R.M., Marzio, A., Pagano, M., Lehmann, R., 2017. GCL and CUL3 Control the Switch between Cell Lineages by Mediating Localized Degradation of an RTK. Dev. Cell 42, 130-142.e7. https://doi.org/10.1016/j.devcel.2017.06.022

Panfilio, K.A., Jentzsch, I.M.V., Benoit, J.B., Erezyilmaz, D., Suzuki, Y., Colella, S., Robertson, H.M., Poelchau, M.F., Waterhouse, R.M., Ioannidis, P., Weirauch, M.T., Hughes, D.S.T., Murali, S.C., Werren, J.H., Jacobs, C.G.C., Duncan, E.J., Armisén, D., Vreede, B.M.I., Baa-Puyoulet, P., Berger, C.S., Chang, C., Chao, H., Chen, M.-J.M., Chen, Y.-T., Childers, C.P., Chipman, A.D., Cridge, A.G., Crumière, A.J.J., Dearden, P.K., Didion, E.M., Dinh, H., Doddapaneni, H.V., Dolan, A., Dugan, S., Extavour, C.G., Febvay, G., Friedrich, M., Ginzburg, N., Han, Y., Heger, P., Holmes, C.J., Horn, T., Hsiao, Y., Jennings, E.C., Johnston, J.S., Jones, T.E., Jones, J.W., Khila, A., Koelzer, S., Kovacova, V., Leask, M., Lee, S.L., Lee, C.-Y., Lovegrove, M.R., Lu, H., Lu, Y., Moore, P.J., Munoz-Torres, M.C., Muzny, D.M., Palli, S.R., Parisot, N., Pick, L., Porter, M.L., Qu, J., Refki, P.N., Richter, R., Rivera-Pomar, R., Rosendale, A.J., Roth, S., Sachs, L., Santos, M.E., Seibert, J., Sghaier, E., Shukla, J.N., Stancliffe, R.J., Tidswell, O., Traverso, L., Zee, M. van der, Viala, S., Worley, K.C., Zdobnov, E.M., Gibbs, R.A., Richards, S., 2019. Molecular evolutionary trends and feeding ecology diversification in the Hemiptera, anchored by the milkweed bug genome. Genome Biol. 20, 64. https://doi.org/10.1186/s13059-019-1660-0

Robertson, S.E., Dockendorff, T.C., Leatherman, J.L., Faulkner, D.L., Jongens, T.A., 1999. germ cell-less,is required only during the establishment of the germ cell lineage of,Drosophila,and has activities which are dependent and independent of its localization to the nuclar envelope. Dev. Biol. 215, 288–297. https://doi.org/10.1006/dbio.1999.9453

Schoppmeier, M., Schröder, R., 2005. Maternal Torso Signaling Controls Body Axis Elongation in a Short Germ Insect. Curr. Biol. 15, 2131–2136. https://doi.org/10.1016/j.cub.2005.10.036

Skelly, J., Pushparajan, C., Duncan, E.J., Dearden, P.K., 2019. Evolution of the Torso activation cassette, a pathway required for terminal patterning and moulting. Insect Mol. Biol. 28, 392–408. https://doi.org/10.1111/imb.12560

Tamura, K., Stecher, G., Kumar, S., 2021. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Mol. Biol. Evol. 38, 3022–3027. https://doi.org/10.1093/molbev/msab120

Warren, S.L., Landolfi, A.S., Curtis, C., Morrow, J.S., 1992. Cytostellin: a novel, highly conserved protein that undergoes continuous redistribution during the cell cycle. J. Cell Sci. 103, 381–388. https://doi.org/10.1242/jcs.103.2.381

Weisbrod, A., Cohen, M., Chipman, A.D., 2013. Evolution of the insect terminal patterning system—Insights from the milkweed bug,Oncopeltus fasciatus. Dev. Biol. 380, 125–131. https://doi.org/10.1016/j.ydbio.2013.04.030

Ylla, G., Nakamura, T., Itoh, T., Kajitani, R., Toyoda, A., Tomonari, S., Bando, T., Ishimaru, Y., Watanabe, T., Fuketa, M., Matsuoka, Y., Barnett, A.A., Noji, S., Mito, T., Extavour, C.G., 2021. Insights into the genomic evolution of insects from cricket genomes. Commun. Biol. 4, 733. https://doi.org/10.1038/s42003-021-02197-9